Focusing attention in human working memory and long-term memory:

benefits through dissociable processes

Dongyu Gong[1-4], Dejan Draschkow[3,4], Anna C. Nobre[1-4]

[1]Department of Psychology, Yale University, New Haven, USA

[2]Wu Tsai Institute, Yale University, New Haven, USA

[3]Department of Experimental Psychology, University of Oxford, Oxford, UK

[4]Oxford Centre for Human Brain Activity, Wellcome Centre for Integrative Neuroimaging, Department of Psychiatry, University of Oxford, Oxford, UK

Author notes

Correspondence concerning this article should be addressed to Dongyu Gong (dongyu.gong@yale.edu), Dejan Draschkow (dejan.draschkow@psy.ox.ac.uk), or Anna C. Nobre (kia.nobre@yale.edu). Address: 100 College Street, New Haven, CT 06510, United States.

Abstract

We developed a new experimental approach to compare how attentional orienting facilitates
retrieval from working memory (WM) and long-term memory (LTM), and how selective attention
within these two memory types impacts incoming sensory information processing. In three
experiments with healthy young adults, retrospective attention cues prioritized an item represented
in WM or LTM. Participants then retrieved a memory item or performed a perceptual task. The
retrocue was informative for the retrieval task but not for the perceptual task. Attentional orienting
benefited performance for both WM and LTM, with stronger effects for WM. Eye-tracking
revealed significant gaze shifts and microsaccades correlated with attention in WM but not LTM.
Visual discrimination of unrelated visual stimuli was consistently improved for items matching
attended WM locations. Similar effects occurred at LTM locations but less consistently. The
findings suggest potent and at least partly dissociable attention-orienting processes for different
memory types.

Introduction

Selective attention is fundamental to human cognition, delivering information that is relevant to

ongoing behaviour among boundless competing signals[1–4]. While standard models emphasise

external sensory inputs as a primary target of attentional control[1,5], the arrow of attention can also

point inward[6,7] to flexibly prioritise and select items within working memory (WM). Retrospective

cues presented during WM delay period that predict goal-relevant items have proved highly useful

for investigating the modulatory mechanisms of internal selective attention[8–10].

Interestingly, like during attentional selection in the external world[11–13], orienting attention within

visual WM is accompanied by subtle gaze shifts[14–16]. Studies have also shown that selecting an

item in WM leads to the capture of attention by sensory stimuli with matching locations or

features[17–19]. These capture effects appear obligatory, operating even when they are counter-

productive for the perceptual task[20], thus underscoring the strong role of WM in guiding external

attention.

While there is a solid appreciation for the strong interrelation between attention and WM, a similar

symbiotic relationship has been proposed for attention and long-term memory (LTM). LTM has

long been considered to guide sensory processing, as exemplified by Helmholtz's view that

perception results from unconscious inference based on prior memories[21]. Experimental tasks

manipulating prior experience with stimulus contexts have since confirmed the role LTM plays in

guiding attention[22–28].

In turn, researchers have posited that selective attention also operates within LTM, guiding

retrieval. Corroborating evidence comes from the observation that brain areas controlling external

attentional orienting are activated in tasks of episodic memory retrieval[29–32] (but see ref. [33]).

Studies showing similar modulation of alpha-band neural activity during visual memory retrieval

as during external visual attention further support a role for attentional selection in guiding LTM

retrieval[34,35]. However, less is understood about *how* selective attention prioritizes different

contents within LTM. One major open question is whether the ability to focus attention on specific

contents within LTM, to guide selective retrieval, relies on the same internal attention processes

that operate in WM. Most standard models of memory suggest that LTM retrieval is mediated by

WM[36,37]. If representations move to WM, then the same internal attention mechanisms should

select relevant mnemonic content for behaviour. However, in principle, focusing on LTM content

to guide retrieval may also occur directly and independently.

To test how selective attention modulates LTM retrieval, we need experimental tasks that

manipulate the goal relevance of items in LTM. Such tasks should provide direct evidence that

orienting attention to a specific item in LTM yields performance benefits relative to retrieving

other items associated with the same context. With evidence suggesting the ability to orient

attention in LTM flexibly and selectively, it becomes possible to probe how similar the processes

and consequences are compared to orienting in WM. Would orienting in LTM also be accompanied

by gaze biases? Would the selective attention in LTM spill over to modulate sensory processing?

Answering these questions would inform enduring debates regarding the relationship between WM

and LTM[36,38–41].

Here, we introduce a new experimental framework to test the ability to orient attention selectively

among competing LTM contents and to compare directly selective attention in WM and LTM. We

borrow from retrospective cue (referred to as "retrocue" hereafter) designs to direct attention to a

specific item within a pre-learned array of items in LTM or to a specific item of an encoded WM

array presented earlier in the trial. Across three experiments, we measured the consequences of

attentional orienting in WM vs. LTM for memory retrieval and additionally tested for spill-over

effects on sensory processing. At the end of each trial, participants either retrieved a memory item

or performed a perceptual discrimination task. The retrocue was informative for the retrieval task

but not for the perceptual task. This allowed us to investigate how orienting selective attention in

WM and LTM enhanced the retrieval of relevant memoranda and biased perceptual processing in

an unrelated task in similar ways.

The findings clearly vindicate the strong effects of internal attention in WM and LTM and,

interestingly, show that dissociable mechanisms are at play. Orienting attention in LTM

consistently speeds selective retrieval and only sometimes improves retrieval accuracy. Benefits

of internal attention in WM are stronger, with larger gains for response speed and consistent

accuracy improvements. Eye-tracking reveal striking differences in engagement of oculomotor

mechanisms during shifts of attention within WM vs. LTM domains. Perceptual discrimination of

unrelated visual stimuli is consistently superior at locations associated with LTM items and is

modulated by internal attention in both WM and LTM domains.

Results

Across three experiments, we demonstrate benefits conferred by orienting attention in WM and
LTM, which differed qualitatively. In Experiment 1, participants acquired two colour-location
bindings in LTM during a training session. In a subsequent testing session, they encoded two
additional colour-location bindings into WM on every trial. Half of the trials tested for benefits of
orienting attention in WM or LTM. Participants indicated the location of an item based on
informative or non-informative colour retrocues. The remaining half tested the impact of
prioritizing an item location in WM or LTM on sensory processing. Participants discriminated the
direction of one of four arrows briefly flashed at WM and LTM locations. Experiments 2 and 3
followed the same training-testing approach, with testing trials measuring benefits of internal
attention on memory performance or sensory processing. Experiment 2 extended the findings to
colour-shape bindings. A non-spatial task probed for benefits of internal attention. Participants
reproduced item shapes using a randomly oriented wheel after informative or non-informative
colour retrocues. This shape reproduction method controlled for the possible contributions of
motor preparation in memory retrieval performance. The perceptual discrimination trials were the
same as in Experiment 1. Experiment 3 was based on Experiment 2, with training and testing
sessions separated across two successive days to ensure that LTM effects did not reflect lingering
priming or shorter-term memory traces. The perceptual discrimination task was replaced with a
simpler task. Only one arrow appeared, and participants identified its direction.

Training LTM associations

For each experiment, we first confirmed that our LTM training was effective. In Experiment 1 (**Fig.**
1), a linear-contrast analysis showed that participants reproduced the LTM colours with smaller
errors on late as compared to early trial bins ($F (1, 26) = 16.156$, $p < .001$, partial $\eta^2 = 0.157$). For
the location reproduction task, participants remained near ceiling-level accuracy across the four
bins (first bin: 0.990 ± 0.005 ($M \pm SEM$), last bin: 0.997 ± 0.002, $F (1, 26) = 0.480$, $p = .490$). In
Experiment 2 (**Fig. 3**), shape reproduction improved significantly across bins (linear contrast: F

$(1, 40) = 17.939$, $p < .001$, partial $\eta^2 = 0.122$). Although improvement in colour reproduction was

not significant ($F(1, 40) = 2.304$, $p = .132$), the errors remained relatively low with a numerically

decreasing trend across bins (first bin: $23.107° ± 2.295$, last bin:$19.182° ± 1.612$). In Experiment

3 (**Fig. 6b**), linear contrasts showed that learning performance improved significantly for both

colour reproduction ($F (1, 115) = 9.015$, $p < .001$, partial $\eta^2 = 0.054$) and shape reproduction (F

$(1, 115) = 38.470$, $p < .001$, partial $\eta^2 = 0.196$).

Attentional orienting exerts stronger effects in WM than LTM

Across three experiments, attentional orienting exerted stronger effects in WM than LTM,

featuring higher benefits in both retrieval speed and accuracy.

In Experiment 1 (**Fig. 2b**), both retrocueing (neutral vs. informative) and memory conditions (WM

vs. LTM) significantly impacted retrieval speed: RTs in the memory retrieval task were faster when

retrocues were informative ($F (1, 29) = 83.205$, $p < .001$, partial $\eta^2 = 0.742$) and when WM items

were probed ($F (1, 29) = 8.239$, $p = .008$, partial $\eta^2 = 0.221$). Crucially, retrocueing and memory

conditions also interacted ($F (1, 29) = 18.627$, $p < .001$, partial $\eta^2 = 0.391$). Both WM and LTM

retrocues conferred a significant benefit (WM: $t (29) = 9.217$, $p_{Bonferroni} < .001$, $d = 1.521$; LTM: t

$(29) = 6.575$, $p_{Bonferroni} < .001$, $d = 0.833$), but retrocue benefits were stronger for the speed of

retrieving WM items ($t (29) = 4.316$, $p < .001$, $d = 0.748$). Similar analyses on retrieval quality

showed significant main effects of retrocueing ($F (1, 29) = 4.340$ $p = .046$, partial $\eta^2 = 0.130$) and

memory conditions ($F (1, 29) = 4.667$, $p = .039$, partial $\eta^2 = 0.139$), as well as a significant

interaction ($F (1, 29) = 12.115$, $p = .002$, partial $\eta^2 = 0.295$). Post-hoc comparisons revealed a

significant improvement in retrieval accuracy by retrocues for WM items ($t (29) = 2.981$, $p_{Bonferroni}$

$= .012$, $d = 0.465$) but no significant effect of retrocues for LTM items ($t (29) = 1.530$, $p_{Bonferroni}$

$= .274$). This could possibly be explained by the overall very high retrieval accuracy for LTM

$(0.977 ± 0.005)$, leaving little room for improvement. A shortcoming of Experiment 1 was the

spatial nature of the memory-retrieval report. Upon presentation of an informative retrocue,

participants could immediately prepare their response, thus confounding the quality and speed of

item selection with response preparation. The non-spatial nature of Experiments 2 and 3, combined

with using a randomly oriented response wheel, overcame this limitation.

In Experiment 2 (**Fig. 4b**), retrocueing significantly shortened RTs ($F (1, 43) = 179.198$, $p < .001$,

partial $\eta^2 = 0.807$). Although RTs were equivalent for both memory conditions ($F (1, 43) = 0.436$,

$p = .513$), there was a significant interaction between the two factors ($F (1, 43) = 24.568$, $p < .001$,

partial $\eta^2 = 0.364$). RT benefits were present for both WM ($t (43) = 11.657$, $p_{Bonferroni} < .001$, $d = $

0.594) and LTM ($t (43) = 9.016$, $p_{Bonferroni} < .001$, $d = 0.273$) but were stronger for WM ($t (43) = $

4.957, $p < .001$, $d = 0.929$). For retrieval accuracy, both retrocueing ($F (1, 43) = 8.971$, $p = .005$,

partial $\eta^2 = 0.173$) and memory conditions ($F (1, 43) = 32.995$, $p < .001$, partial $\eta^2 = 0.434$) exerted

significant main effects. The two factors also interacted ($F (1, 43) = 7.441$, $p = .006$, partial $\eta^2 = $

0.163). Post-hoc comparisons revealed that WM retrocues significantly reduced shape

reproduction errors ($t (43) = 4.322$, $p_{Bonferroni} < .001$, $d = 0.429$) but LTM retrocues did not ($t (43)$

$= 1.250$, $p_{Bonferroni} = .436$). Shape reproduction was very accurate, and errors were smaller when

LTM shapes were reproduced ($16.610° \pm 1.519$) as compared to WM shapes ($27.603° \pm 2.417$),

possibly reflecting ceiling effects.

In Experiment 3 (**Fig. 6c**), the one-day interval between training and testing brought LTM

performance below ceiling, allowing us to test for LTM accuracy benefits by internal attention.

Overall, the next-day LTM representations in Experiment 3 were weaker than the same-day

representations in Experiment 2 (reproduction errors in Experiment 3 vs. Experiment 2: $51.547 \pm$

2.880 vs. 17.378 ± 1.819, Welch's $t = 10.031$, $p < .001$). Informative retrocues significantly

improved RTs ($F (1, 118) = 113.287$, $p < .001$, partial $\eta^2 = 0.490$). Response times did not differ

between memory condition ($F (1, 29) = 0.271$, $p = .603$) but the two factors interacted ($F (1,118)$

$= 12.643$, $p < .001$, partial $\eta^2 = 0.097$). Post-hoc t-tests showed that RT benefits were present for

both WM ($t (118) = 10.080$, $p_{Bonferroni} < .001$, $d = 0.318$) and LTM ($t (118) = 9.381$, $p_{Bonferroni} < .001$,

$d = 0.253$) but were stronger for WM ($t (118) = 3.556$, $p < .001$, $d = 0.267$). For retrieval accuracy,

retrocueing ($F (1, 118) = 55.519$, $p < .001$, partial $\eta^2 = 0.320$) and memory condition ($F (1, 118) = $

6.947, $p = .010$, partial $\eta^2 = 0.056$) both exerted significant main effects and interacted ($F (1, 118)$

$= 15.216$, $p < .001$, partial $\eta^2 = 0.114$). Post-hoc comparisons revealed that informative retrocues

significantly reduced shape reproduction errors for both WM items ($t (118) = 9.581$, $p_{Bonferroni}$

$< .001$, $d = 0.540$) and LTM items ($t (118) = 2.792$, $p_{Bonferroni} = .018$, $d = 0.143$). Error reduction

was stronger for WM items ($t (118) = 3.901$, $p < .001$, $d = 0.474$). The results from Experiment 3

thus showed that attentional orienting can improve LTM retrieval accuracy when overall LTM

performance is below ceiling.

Sensory spill-over effects of prioritizing WM and LTM items

The current study also yielded insights on how prioritizing items in WM or LTM impacts sensory
information processing at the item location.

In Experiment 1 (**Fig. 2c**), selective prioritization of WM and LTM memoranda by retrocues
impacted perceptual discrimination of items occurring at the matching location. Perceptual
discrimination accuracy showed a main effect of retrocue matching ($F(1, 29) = 31.541$, $p < .001$,
partial $\eta^2 = 0.521$), with superior accuracy for discriminating stimuli at retrocue-matching
locations than for stimuli on neutral-retrocue trials. This accuracy benefit was significantly larger
than zero for both memory conditions (WM: 0.094 ± 0.020, $t(29) = 4.804$, $p < .001$, $d = 0.877$;
LTM: 0.121 ± 0.028, $t(29) = 4.362$, $p < .001$, $d = 0.797$) and did not differ in size ($t(29) = 0.962$,
$p = .344$). We also observed a main effect of memory condition ($F(1, 29) = 4.887$, $p = .035$, partial
$\eta^2 = 0.144$). Accuracy was higher when LTM locations were probed (0.655 ± 0.026) compared to
WM locations (0.601 ± 0.032). A separate analysis confirmed there was no effect of congruence
between item location and arrow-pointing direction ($t(29) = 0.894$, $p = 0.379$).

Improvement of perceptual discrimination in Experiment 1 could have resulted from response
preparation related to reporting item location rather than prioritization of an item in WM or LTM.
In Experiment 2 (**Fig. 4c**), there was no direct link between spatial location and responses in the
memory retrieval task, so that any effect on perceptual discrimination could be linked to
prioritization of items in WM or LTM. Under these conditions, perceptual discrimination still
benefitted at locations matching the retrocued items ($F(1, 43) = 9.504$, $p = .004$, partial $\eta^2 = 0.181$).
Perceptual benefits for items matching retrocued WM and LTM locations were both significantly
larger than zero (WM: 0.037 ± 0.003, $t(43) = 2.191$, $p = .034$, $d = 0.330$; LTM: 0.032 ± 0.002, t
$(43) = 2.176$, $p = .035$, $d = 0.328$) and did not differ in size ($t(43) = 0.259$, $p = .797$). As in
Experiment 1, discrimination performance was better at LTM than WM locations overall ($F(1, 43)$
$= 55.213$, $p < .001$, partial $\eta^2 = 0.562$). A separate analysis confirmed there was no effect of
congruence between item location and arrow-pointing direction ($t(43) = 0.354$, p $= 0.725$).

In Experiment 3 (**Fig. 6d**), the main effect of retrocue matching did not reach significance ($F(1,$
$118) = 3.473$, $p = .065$). Memory condition still exerted a main effect, with perceptual
discrimination better at LTM than WM locations ($F(1, 118) = 32.213$, $p < .001$, partial $\eta^2 = 0.214$).

Retrocueing and memory condition interacted (F (1,118) = 10.328, p = .002, partial η^2 = 0.080).

Perceptual benefits of WM retrocues were significantly larger than zero (t (118) = 3.308, p = .001,

d = 0.214), while benefits from LTM retrocues were not (t (118) = 0.896, p = .372).

Attentional orienting in WM but not LTM elicited directional gaze biases

In Experiment 2, eye tracking investigated the engagement of the oculomotor system during

attentional orienting. Replicating previous findings[14–16], WM retrocues elicited significant gaze

biases in the direction of attentional orienting. In contrast, no evidence of similar gaze biases

occurred when orienting attention in LTM (**Fig. 5a-5d**). Compared to the baseline pattern of gaze

shifts following neutral cues, systematic changes were observed for trials with WM retrocues

depending on the direction of the retrocued item, both horizontally and vertically. No significant

changes occurred for trials with LTM retrocues, on which the baseline pattern of gaze shifts was

observed regardless of the retrocued item direction.

The differences between gaze biases following WM and LTM retrocues were quantified using the

associated towardness time courses. Cluster-based permutation testing confirmed significant

differences following WM and LTM retrocues in both horizontal and vertical directions (horizontal:

approximately 450-1000 ms after cue onset, p = .002; vertical: approximately 420-870 ms after

cue onset, p = .003). No significant clusters occurred following LTM retrocues.

A complementary analysis of directional microsaccades (see ref. [42]) showed a similar pattern (**Fig.**

5e-5f). After WM retrocues, microsaccades occurred more often in the direction of the prioritized

item than the other item (approximately 380-490 ms after cue onset, p < .001). No systematic

directional changes in microsaccade rates occurred after LTM retrocues.

Discussion

We demonstrate that internal attention-directing cues benefit both WM and LTM recall in humans,

with qualitatively different consequences, and through non-coextensive mechanisms. Our findings

reinforce the increasing recognition of the strong and multidirectional relationship between

attention and memories of different durations[43,44] as well as the plurality of mechanisms supporting

their interactions[7].

We introduced a new approach for investigating and comparing selective attention operating in
LTM vs. WM using equivalent stimulus parameters and response requirements within the same
tasks. In our tasks, orienting attention to contents in LTM significantly improved retrieval speed
in a similar, though less pronounced way as for WM. Accuracy benefits were more consistent for
WM than LTM. No improvements in LTM accuracy occurred when retrieval performance was at
very high levels overall (Experiments 1 and 2) but improvements emerged when there was greater
room for improvement in retrieval accuracy (Experiment 3). The current pattern of results suggests
that attention can, at least, consistently enhance the accessibility or "readiness" to act on LTM
memory representations. The modulations observed for WM compared to LTM retrieval were
stronger, shortening response times to a greater degree and consistently affecting retrieval accuracy.
The findings suggest differences in the mechanisms of internal attention within these different
memory domains. However, on their own, these behavioural results could result from variations
in the strength of the memory representations themselves and the resulting retrieval demands.

Our eye-tracking data yielded the most suggestive evidence for a functional dissociation between
the mechanisms for orienting attention in WM and LTM. By measuring directional biases in eye
gaze and microsaccades, we revealed a striking difference in oculomotor involvement in trials with
WM vs. LTM retrocues. Consistent with previous work[14–16], we observed that gaze was biased
toward retrocued items in WM. Critically, no significant gaze biases were detected following LTM
retrocues.

Our eye-tracking results thus clearly demonstrate that internal attention in LTM is not co-extensive
with internal attention in WM. Orienting attention in WM engages a control network of frontal,
parietal, and subcortical areas modulating activity in task-relevant sensory and motor areas[45–47].
The close relationship between the internal attention control regions and regions involved in
oculomotor control results in correlated gaze shifts and microsaccades. The network and
mechanisms for orienting attention in LTM have not yet been fully characterized. They may be
qualitatively different than the network for attention in WM. For example, posterior parietal areas
have been implicated in selective LTM retrieval, with parallels drawn to external attention
mechanisms (e.g.,[29]). However, the degree of anatomical overlap has been questioned[33].

Engagement of attention-related frontal areas has been less conspicuous. In addition, established

plasticity patterns related to the longer-term associations may confer alternative or additional

mechanisms for prioritizing feature values of LTM items. Informative LTM retrocues may interact

with latent functional states that have been intrinsically reinforced by associative plasticity.

We also measured the spill-over effects of orienting in WM and LTM on sensory processing. While

previous reports of this phenomenon focused on the influence of WM[48,49], our results showed that

comparable effects can result from prioritizing LTM representations. In the perceptual

discrimination task, prioritising items within the spatial layout of either WM or LTM boosted the

discrimination accuracy of location-matching stimuli in two of the three experiments (Experiments

1 and 2). This occurred despite there being no strategic benefit for using the retrocue information

in the perceptual task since the location of the retrocued item only matched the probed location on

25% of the trials. Consequences of memory prioritization for sensory processing were therefore

likely incidental. These perceptual improvements at the matching locations of attended items

reflect the engagement of spatial attention. It is worth noting that spatial representations are not

strictly required in the Testing phase of Experiment 2; cueing the item colour required retrieval of

the corresponding shape. The automatic recruitment of spatial attention, even when not required,

has been commonly observed in WM tasks[14,50]. The spatial layout of encoded items has been

proposed to provide an important scaffolding for WM representations[51–53]. The findings suggest

that the spatial layout may also be a useful or intrinsic preserved property of encoded LTM arrays.

Interestingly, in Experiment 2, significant perceptual benefits at LTM-matching locations were

observed despite no significant differences in gaze biases or microsaccades. Previous studies have

shown that oculomotor markers are strongly correlated with orienting attention within the spatial

layout of WM but are not obligatory[42]. The current findings reinforce the non-obligatory nature of

the relationship between internal spatial attention and eye movements. Focusing within LTM can

facilitate sensory discrimination in a spatially specific way without accompanying oculomotor

readouts.

Experiment 3 introduced a simpler perceptual identification task to corroborate further the findings

on perceptual benefits from the previous two experiments. Retrocueing WM items still conferred

perceptual benefits, whereas retrocueing LTM items no longer did. The lack of perceptual benefits

linked to prioritizing items in LTM could reflect the weaker nature of the next-day LTM
representations in Experiment 3 compared to Experiments 1 and 2. Alternatively, spillover effects
may have been overridden by capture from the single arrow cue or dampened by reduced
competition in the perceptual discrimination task used in Experiment 3 (see ref. [54]). Greater
variability of online performance may also have contributed. It will be interesting to explore further
the boundary conditions for perceptual spillover effects of internal attention in LTM.

An additional, unexpected, and interesting finding was the superior performance in the perceptual
discrimination at locations associated with LTM compared to WM items across all three
experiments. Involvement of longer-term plasticity mechanisms for the LTM items may have
contributed to better sensory performance for items appearing in LTM locations. Cortical plasticity
mechanisms akin to operating in perceptual[55] or statistical[56] learning could render the functional
states associated with LTM locations and features more accessible to re-activation. Another
possible explanation for the disparity in perceptual sensitivity at the locations of LTM vs. WM
items is the degree of active sensory recruitment supporting the two types of memory(see ref. [45]).
Stronger activation of feature-specific sensory signals during attention to WM could interfere with
processing other incoming visual stimuli. However, this explanation would not account for better
performance for discriminating stimuli at attended compared to unattended WM locations. The
mechanisms behind the intriguing findings warrant further study.

Some of the findings in the current study inevitably depend on specific task parameters and
demands. The LTM representations investigated in this study were simple visual associations of
object features and locations (e.g., [57,58]). These representations are similar to those often studied in
visual WM tasks but are less often considered in LTM studies. Given the unimodal and simple
nature of these associations, they may not require the involvement of multisensory or contextual
association hubs, such as the medial temporal lobe[59]. Using these simple visual associative LTMs,
we observed that selective feature retrieval did not engage oculomotor markers of spatial attention.
However, under some conditions, focusing within LTM may rely on spatial codes. For example,
systematic eye-movement changes accompany attention to remembered items within scenes or
contexts[26,60,61].

Nevertheless, some of the observations in the study carry generalizable lessons. The dissociation

between the degree of reliance on spatial codes when orienting attention in WM vs. LTM in our

task shows that focusing on contents within LTM is not necessarily mediated by placing

representations within WM and using standard internal attention mechanisms for WM. It will be

interesting and important to build on our laboratory-based study to test for the generalizability of

our observations within natural, immersive contexts, where psychological mechanisms can

sometimes deviate from those predicted by experiments in more controlled, artificial settings (see

ref. [62]).

The nature of the spatial frameworks used when orienting attention in WM is open to further

investigation. In contrast to the extensive literature considering how various representational

frameworks contribute to human performance based on LTM (e.g., [63–65]), the topic is

underexplored in WM tasks (with notable exceptions, e.g., [66–68]). Recent virtual-reality tasks have

shown that orienting attention to items within WM in immersive settings engages both an

egocentric spatiotopic framework and an allocentric framework based on the layout of the initially

encoded items[16]. Further investigation will likely reveal a much richer array of interacting spatial

frameworks in WM tasks.

In summary, we compared the effects of focusing attention within human WM vs. LTM and

discovered that both bring significant benefits to retrieval and subsequent sensory processing, but

through dissociable mechanisms. The strikingly distinct oculomotor signatures of covert spatial

attention in WM and LTM corroborate a plurality of functional properties when memories of

different durations guide adaptive behaviour and open novel opportunities for furthering our

understanding of the relationship between WM and LTM.

Methods

Experiment 1

Participants. Thirty individuals (23 females, 7 males, $M = 25.13$ years, $SD = 4.15$) with reported

normal or corrected visual acuity volunteered and received monetary compensation for

participation. The experiment was approved by the Central University Research Ethics Committee

of the University of Oxford, and all participants provided informed consent before any

experimental procedure began.

Apparatus and stimuli. Participants sat in front of a 27″ monitor (1920 × 1080 pixels, 100 Hz)

and rested their chin on a chinrest placed 95 cm away from the monitor. The experiment was

programmed in MATLAB (MathWorks, Natick, MA, U.S.A.) using the Psychophysics Toolbox[69].

Stimuli appeared overlaid on a grey background. Throughout the experiment, four squares (2.5°

in diameter) were always presented as placeholders at the four quadrants, at 5° horizontally and

vertically from the central fixation to the centre of each square. The stimuli consisted of four

equiluminant colours (brown [183.5, 113, 19], green [65, 143, 110.5], lilac [138, 117.5, 190],

magenta [245.5, 37, 112.5]) drawn from a circle in CIELAB colour space.

Procedure and design. The experiment included a learning session and a testing session,

separated by a 5-minute break. During the learning session, participants were trained to encode

two colours and their corresponding locations into LTM (**Fig. 1**). These two colours were randomly

selected from the four colours defined above, and they were always located along one of the two

pairs of diagonal locations (i.e., top left and bottom right, or top right and bottom left,

counterbalanced across participants). We refer to this pair of locations as LTM locations. Each

learning trial began with a fixation display lasting randomly between 800 and 1000 ms, after which

the LTM display was presented for 150 ms. Following a delay of 850 ms, participants were probed

to reproduce either the colour at one location or the location of one colour.

On colour reproduction trials, a colour wheel (containing 360 colours) was presented at the centre,

and participants responded by rotating the dial and selecting a colour along the wheel. The colour

wheel was presented in a random orientation on every trial. Immediately after the response,

feedback was presented for 1000 ms in the form of an integer ranging from 0 to 100, with 100

indicating a perfect reproduction of the probed colour and 0 indicating the exact opposite on the

wheel. On location reproduction trials, one of the two colours was presented at the centre, and

participants responded by pressing one of four keys mapped to the four locations (Q for top left,

W for top right, A for bottom left, and S for bottom right). Performance feedback was presented

for 500 ms indicating whether the chosen location was correct or wrong. Each to-be-learned

attribute (two colours and two locations) was probed on 20 trials, resulting in a total of 80 learning

trials presented in random order.

During the testing session, participants performed either a memory retrieval task or a perceptual
discrimination task on each trial (**Fig. 2a**). Each testing trial began with a fixation display (800-
1000 ms). A WM display followed, where the two colours unused in the learning session were
presented for 150 ms at the unused pair of diagonal locations. We refer to this pair of locations as
WM locations. To make sure the contents in WM were not fixed across trials, each WM colour
was randomly assigned to one of the WM locations on every trial. Following a delay of 850 ms, a
retrocue that was either neutral or informative was presented for 200 ms. The retrocue was neutral
on one-third of the trials; the fixation display changed to white, providing no information about
the item to be probed. The retrocue was informative on two-thirds of the trials; the fixation display
changed to one of the four colours, matching either a WM or LTM item with equal probability.
Informative retrocues indicated the item to be probed in the memory retrieval task with 100%
validity. Following a second delay of 800 ms after the retrocue, the memory retrieval task and the
perceptual discrimination tasks were equally likely to be presented.

In the memory retrieval task, participants were required to retrieve the location of the probed item.
On trials containing an informative retrocue, participants reported the location of the WM or LTM
item indicated by the retrocue. On trials containing a neutral cue, the probed item was indicated
by a centrally presented colour, chosen randomly from the WM or LTM colours. Responses were
delivered by pressing one of four keys mapped to the four locations (the same keys as used in the
location reproduction task during the learning session). Feedback was then presented for 500 ms
indicating whether the response was correct or wrong.

In the perceptual discrimination task, four arrows (length: 1.25°, width of the tail: 0.625°, RGB
value: [128, 128, 128]) were presented in the placeholders for 100 ms, after which randomly
generated Gaussian noise masks were applied to the four locations for 100 ms. Following the mask,
one location was probed, and participants pressed one of the arrow keys to report the arrow
direction at that location. Each of the four locations had an equal possibility of being probed. The
arrow direction at each location was independently drawn from four possible directions ($\uparrow$, $\leftarrow$, $\downarrow$,
$\rightarrow$), resulting in $4^4 = 256$ combinations. The choice of presenting arrow stimuli at all four locations
and using a post-cue to elicit a response was intended to avoid the sensory capture by stimuli with
different attributes.

Participants were informed that the memorized items and the retrocue bore no predictive relation

concerning the location of the perceptual item to be discriminated. When retrocues were

informative, however, participants always needed to maintain the retrocued item in mind for

potential future use because of the randomization of the memory retrieval and perceptual

discrimination trial order, which allowed us to examine any spill-over benefits elicited by the

retrocue on the perceptual discrimination task. The relationship between the perceptual

discrimination task and the retrocue was totally incidental because all locations were equally likely

to be probed no matter which location the retrocue would preferentially bias attention to. As a

result, only 25% of informative-retrocue trials in the perceptual discrimination task were

"matching" trials on which the probed sensory location coincided with the location of the retrocued

item.

The testing session consisted of 480 trials divided into 10 blocks (each including 48 trials). To

become familiarized with the task, participants performed an additional 48 practice trials before

testing.

Behavioural Analysis. Behavioural data were analysed in MATLAB. For the learning session, we

examined the average colour reproduction errors and location reproduction accuracy respectively,

by sorting the learning trials of each type into 4 bins (each containing 10 trials). Colour

reproduction errors (in units of degrees) were calculated by taking the absolute difference between

the angle of the target colour and the reproduced colour on the colour wheel. One-way repeated-

measures ANOVAs with linear contrast weights ([-3, -1, 1, 3]) across four bins tested for the

efficacy of training. For the testing session, data from the memory retrieval and perceptual

discrimination tasks were analysed separately. During pre-processing, we excluded trials on which

RTs were 3 SD above the individual mean across all conditions in either task. After this exclusion

step, an average of 98.35% (SD = 0.44%) trials were retained in the analyses.

To test for benefits of internal selective attention on WM and LTM retrieval, we analysed the

average RTs and accuracy for the memory retrieval task as a function of retrocueing (neutral vs.

informative) and the memory timescale of probed items (WM vs. LTM). To examine whether

orienting attention to a memory item benefited subsequent perceptual processing at matching

locations, we compared perceptual discrimination accuracy on retrocue matching trials vs. neutral

trials (matching vs. neutral) when locations associated with WM or LTM items were probed (WM

vs. LTM). To gauge the quality of perception, accuracy was the dependent variable of interest, but

RTs were also evaluated for completeness.

When comparing behavioural performance between conditions, we applied a repeated-measures

ANOVA and reported partial η^2 as a measure of effect size. For post hoc t-tests, we reported

Bonferroni-corrected p values that we denoted as "$p_{Bonferroni}$". We reported Cohen's d as a measure

of effect size for all the t-tests. Where relevant, the within-subject standard error of the mean (*SEM*)

was calculated from normalized data[41]. When evaluating potential perceptual benefits elicited by

WM and LTM retrocues, we applied one-sample t-tests against 0.

Experiment 2

Participants. A total of 44 volunteers (27 females, 17 males, $M = 25.93$ years, $SD = 4.51$) with

reported normal or corrected-to-normal visual acuity were recruited. The sample size was

calculated using G*Power[70] to achieve 90% power for the one-sample t-tests performed to test the

significance of perceptual benefits following WM and LTM retrocues. The effect sizes for these

comparisons in Experiment 1 were 0.877 and 0.797. We assumed a conservative approach and

aimed to power for the detection of a medium effect size (0.5) because we expected that the

manipulation in Experiment 2 would lead to a smaller effect due to the incidental nature of the

spatial attributes in the task. Most of the experimental setup was identical to Experiment 1, with

the following modifications.

Apparatus and stimuli. Eye movements were recorded with the EyeLink 1000 Desktop Mount

(SR Research, Ottawa, ON, Canada) at 1000 Hz. Eye-tracker calibration used the built-in

calibration and validation protocols from the EyeLink software. When possible, horizontal and

vertical gaze positions were continuously recorded for both eyes. For some participants ($N = 13$),

only one eye was tracked due to a lack of good-quality binocular tracking (mostly because of

wearing glasses). Stimuli appeared on a white background. Four shapes were equidistantly

sampled from the Validated Circular Shape (VCS) space[71], and then randomly assigned to each

WM and LTM item, adding a new feature dimension to the existing configurations in Experiment

1. These same four shapes were used across all participants, but the shapes assigned to WM and

LTM items were randomized across participants.

Procedure and design. During the learning session, participants were trained to memorize the
colours and shapes of the two LTM items (**Fig. 3**). On every trial, they were probed to reproduce
either the colour or the shape of one item. At the response stage, either a colour wheel or a shape
wheel was presented at the centre, indicating the feature dimension to be reproduced in this trial.
The spatial location of the item was used to probe the colour or shape reproduction, but participants
were never asked to report the item location. Both the colour and shape wheels were presented in
a random orientation each time. The shape wheel consisted of 360 shapes from the VCS space. To
avoid clustering, eight shapes sampled from equidistant positions on the wheel were displayed
along the cardinal axes (i.e., every 45 degrees). These eight shapes served as visual anchors, which
were also randomly chosen every time based on the orientation of the shape wheel. Participants
responded using a computer mouse that controlled the dial on the wheel. Participants had unlimited
time to retrieve the item from memory and to decide what to reproduce. However, once they started
moving the dial, they had only 2500 ms to complete their reproduction. This was intended to
encourage participants to retrieve the exact colour or shape before moving the dial. The position
of the dial when participants clicked the left mouse button or when the time limit was reached was
taken as the response. Immediately after their response, participants received feedback for 500 ms.
Each colour and shape were probed on 20 trials, resulting in a total of 80 learning trials presented
in random order.

During the testing session, each WM shape was randomly combined with one of the WM colours
on every trial (**Fig. 4a**). As in Experiment 1, participants performed either a memory retrieval task
or a perceptual discrimination task on each trial equiprobably. In the memory retrieval task,
participants reproduced the shape of the item matching the retrocued colour or a randomly probed
colour when the retrocue was neutral. The shape wheel was identical to that used during the
learning session and was randomly rotated across trials. Uninformative, grey neutral retrocues
appeared on one-fifth of the trials. The remaining trials contained informative coloured retrocues
matching each of the LTM or WM colours with equal probability. In contrast to Experiment 1, the
memory retrieval task in Experiment 2 was non-spatial. Retrieving the item shape based on the
cued colour does not require using any spatial association. Although spatial locations were used
for training the colour-shape associations, participants were never cued or asked to report the item
location in the memory retrieval task.

The perceptual discrimination task was the same as that in Experiment 1.

The testing session consisted of 600 trials divided into 10 blocks (each including 60 trials). To
become familiarized with the task, participants performed an additional 30 practice trials before
testing.

Behavioural Analysis. The analyses of interest were basically the same as in Experiment 1, with
location reproduction in the learning and testing sessions replaced by shape reproduction. Shape
reproduction errors (in units of degrees) were calculated by taking the absolute difference between
the angle of the target shape and the reproduced shape on the shape wheel. The RTs in the memory
retrieval task were calculated as the time from probe onset to when the response was recorded,
either when participants clicked the left mouse button or when the time limit was reached. After
excluding memory retrieval and perceptual discrimination trials on which RTs were 3 SD above
the individual mean across all conditions, an average of 98.66% (SD = 0.47%) trials were retained
in the analyses.

Eye-tracking Analysis. Data were first converted from edf to asc format and subsequently read
into RStudio. For binocularly tracked participants, data from the left and right eyes were averaged
to obtain a single horizontal and a single vertical gaze position channel. Blinks were marked by
detecting NaN clusters in the eye-tracking data and then interpolated using a linear interpolation
procedure. Data were epoched from 250 ms before to 1000 ms after cue onset. To make our
analyses more robust to drift of the eyes during fixation, we obtained the average gaze position
within the 250 ms window before cue onset for every trial and participant and subtracted it from
every corresponding time course. We performed analyses on horizontal and vertical channels
separately. For both channels, we only included trials on which gaze position remained within ±50%
from fixation (with 100% denoting the centres of the original item locations at a ± 5° visual angle)
throughout the trial, as previous work showed that the gaze bias phenomenon is constituted by a
bias in gaze around fixation[14–16]. For the horizontal channel, data for all 44 participants were used,
with an average of 9.6% ± 1.7% (*M ± SEM*) trials excluded per participant. For the vertical channel,
six participants had to be removed due to a high number of excluded trials (> 50%). For the 38
participants retained, an average of 14.6% ± 2.1% trials were excluded. Gaze time courses were
smoothed using a 25-ms average moving window.

For the horizontal channel, we compared trial-averaged gaze-position time courses between

conditions in which the retrocued item occupied the left side (top/bottom left location) or the right

side (top/bottom right location) during encoding, separately for trials with WM retrocues and trials

with LTM retrocues. For the vertical channel, we did the same between conditions in which the

retrocued item occupied the bottom side (bottom left/right location) or the top side (top left/right

location) during encoding, also separately for trials with WM and LTM retrocues. For both

channels, trial-averaged gaze-position time courses were also obtained for trials with neutral

retrocues. To increase sensitivity, we also constructed a measure of towardness separately for trials

with WM and LTM retrocues, which expressed the gaze bias toward the side of the retrocued item

in a single value[14–16]. We did this for both horizontal and vertical channels to obtain towardness in

horizontal and vertical directions, respectively.

Analysis of microsaccade rates toward vs. away from prioritized items in WM or LTM followed

previously described methods[42]. Saccades were detected using a velocity-based method and only

those with a magnitude smaller than 1 degree of visual angle were considered microsaccades[42].

Furthermore, for direct comparisons with previous work[42,72], we focused exclusively on

microsaccades along the horizontal axis. Depending on the side of the retrocued item, we labelled

microsaccades as "toward" or "away" based on whether they were moving toward or away from

the retrocued item. The resulting time courses of "toward" and "away" microsaccade rates were

smoothed using a moving average with a 50-ms sliding window.

Statistical evaluation of the towardness time courses used a cluster-based permutation approach[73]

implemented in the *permuco* package, which is ideally suited to evaluate physiological effects

across multiple time points while retaining high sensitivity.

Experiment 3

Participants. Given the assumption that LTM would decay on the second day, we aimed to power

for detecting an effect size of 0.3 for perceptual benefits, which led G*power to yield a sample

size of 119 participants. Participants (86 males, 33 females; mean age = 28.69 years, *SD* = 6.04

years; 106 right-handed, 12 left-handed, 1 ambidextrous) were recruited on Prolific

(https://www.prolific.co/). They were pre-screened on demographic criteria (age range 18 to 40,

fluent in English), general health (normal or corrected-to-normal vision, no previous or ongoing

mental illnesses), and participation history on Prolific Academic (participated in at least 10 studies,

with a study approval rate above 90%). All participants provided informed consent before

participating and were paid $12 per hour.

Stimuli and procedure. Participants performed 80 learning trials on the first day, which was

identical to the learning session in Experiment 2. Participants were invited to complete the testing

session around 24 hours after training. The differences in the testing session compared to

Experiment 2 are as follows.

Experiment 3 used a simpler perceptual discrimination task, which avoided post-cues. The

motivation was to tap into perceptual sensitivity more directly, without requiring post-encoding

transfer and selection before behavioural reports. In the perceptual array, only one placeholder

contained an arrow drawn from four directions (↑, ←, ↓, →). The other three placeholders

contained randomly selected horizontal or vertical bars as control stimuli (**Fig. 6a**).

Correspondingly, at the probe stage, participants were asked to report the direction of the single

presented arrow by pressing an arrow key. Considering the greater variability of stimulus timing

in online studies[74], the duration of the perceptual array was lengthened to 200 ms. The testing

session consisted of 480 trials divided into 12 blocks (each including 40 trials). Participants

performed an additional block of 40 practice trials before the 480 testing trials. During pre-

processing, we excluded memory retrieval and perceptual discrimination trials with RTs exceeding

3 SD above the individual mean across all conditions. After this procedure, an average of 98.65%

(SD = 0.61%) trials were retained in the analyses.

Code availability

Analysis code is available at https://github.com/Daniel-Gong/Orienting-in-WM-and-LTM/.

Data availability

All data are publicly available through the Open Science Framework at https://osf.io/n629s/.

References

1. Desimone, R. & Duncan, J. S. Neural Mechanisms of Selective Visual Attention. *Annu. Rev. Neurosci.* **18**, 193–222 (1995).

2. Fawcett, J., Risko, E. & Kingstone, A. *The Handbook of Attention*. (MIT Press, 2015).

3. Nobre, A. C. Attention. in *Stevens' Handbook of Experimental Psychology and Cognitive Neuroscience* (ed. Wixted, J. T.) 1–75 (Wiley, 2018).

4. Nobre, K. & Kastner, S. *The Oxford Handbook of Attention*. (Oxford University Press, 2014).

5. Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the brain. *Nat. Rev. Neurosci.* **3**, 201–215 (2002).

6. Chun, M. M., Golomb, J. D. & Turk-Browne, N. B. A Taxonomy of External and Internal Attention. *Annu. Rev. Psychol.* **62**, 73–101 (2011).

7. Van Ede, F. & Nobre, A. C. Turning Attention Inside Out: How Working Memory Serves Behavior. *Annu. Rev. Psychol.* **74**, 137–165 (2023).

8. Griffin, I. C. & Nobre, A. C. Orienting Attention to Locations in Internal Representations. *J. Cogn. Neurosci.* **15**, 1176–1194 (2003).

9. Landman, R., Spekreijse, H. & Lamme, V. A. F. Large capacity storage of integrated objects before change blindness. *Vision Res.* **43**, 149–164 (2003).

10. Souza, A. S. & Oberauer, K. In search of the focus of attention in working memory: 13 years of the retro-cue effect. *Atten. Percept. Psychophys.* **78**, 1839–1860 (2016).

11. Engbert, R. & Kliegl, R. Microsaccades uncover the orientation of covert attention. *Vision Res.* **43**, 1035–1045 (2003).

12. Hafed, Z. M. & Clark, J. J. Microsaccades as an overt measure of covert attention shifts. *Vision*

619   *Res.* **42**, 2533–2545 (2002).

620   13. Corneil, B. D. & Munoz, D. P. Overt Responses during Covert Orienting. *Neuron* **82**, 1230–

621   1243 (2014).

622   14. van Ede, F., Chekroud, S. R. & Nobre, A. C. Human gaze tracks attentional focusing in

623   memorized visual space. *Nat. Hum. Behav.* **3**, 462–470 (2019).

624   15. van Ede, F., Board, A. & Nobre, A. Goal-directed and stimulus-driven selection of internal

625   representations. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 24590–24598 (2020).

626   16. Draschkow, D., Nobre, A. C. & van Ede, F. Multiple spatial frames for immersive working

627   memory. *Nat. Hum. Behav.* (2022).

628   17. Awh, E. & Jonides, J. Overlapping mechanisms of attention and spatial working memory.

629   *Trends Cogn. Sci.* **5**, 119–126 (2001).

630   18. Downing, P. E. Interactions between visual working memory and selective attention. *Psychol.*

631   *Sci.* **11**, 467–473 (2000).

632   19. Soto, D., Humphreys, G. W. & Rotshtein, P. Dissociating the neural mechanisms of memory-

633   based guidance of visual selection. *Proc. Natl. Acad. Sci.* **104**, 17186–17191 (2007).

634   20. Soto, D. & Humphreys, G. W. Working Memory Biases in Human Vision. in *The Oxford*

635   *Handbook of Attention* (eds. Nobre, A. C. (Kia) & Kastner, S.) 0 (Oxford University Press,

636   2014).

637   21. Von Helmholtz, H. *Handbuch Der Physiologischen Optik*. vol. 9 (Voss, 1867).

638   22. Brockmole, J. R., Castelhano, M. S. & Henderson, J. M. Contextual cueing in naturalistic

639   scenes: Global and local contexts. *J. Exp. Psychol. Learn. Mem. Cogn.* **32**, 699–706 (2006).

640   23. Chun, M. M. & Jiang, Y. Contextual cueing: Implicit learning and memory of visual context

641   guides spatial attention. *Cognit. Psychol.* **36**, 28–71 (1998).

24. Fan, J. E. & Turk-Browne, N. B. Incidental biasing of attention from visual long-term memory.

J. Exp. Psychol. Learn. Mem. Cogn. **42**, 970–977 (2016).

25. Stokes, M. G., Atherton, K. E., Patai, E. Z. & Nobre, A. C. Long-term memory prepares neural

activity for perception. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 1838–1839 (2012).

26. Summerfield, J. J., Lepsien, J., Gitelman, D. R., Mesulam, M.-M. & Nobre, A. C. Orienting

attention based on long-term memory experience. *Neuron* **49**, 905–916 (2006).

27. Theeuwes, J., Bogaerts, L. & van Moorselaar, D. What to expect where and when: how

statistical learning drives visual selection. *Trends Cogn. Sci.* **26**, 860–872 (2022).

28. Võ, M. L.-H., Boettcher, S. E. & Draschkow, D. Reading scenes: how scene grammar guides

attention and aids perception in real-world environments. *Curr. Opin. Psychol.* **29**, 205–210

(2019).

29. Cabeza, R., Ciaramelli, E., Olson, I. R. & Moscovitch, M. The parietal cortex and episodic

memory: an attentional account. *Nat. Rev. Neurosci.* **9**, 613–625 (2008).

30. Chun, M. M. & Johnson, M. K. Memory: Enduring Traces of Perceptual and Reflective

Attention. *Neuron* **72**, 520–535 (2011).

31. Ciaramelli, E., Grady, C. L. & Moscovitch, M. Top-down and bottom-up attention to memory:

a hypothesis (AtoM) on the role of the posterior parietal cortex in memory retrieval.

Neuropsychologia **46**, 1828–1851 (2008).

32. Sestieri, C., Shulman, G. L. & Corbetta, M. The contribution of the human posterior parietal

cortex to episodic memory. *Nat. Rev. Neurosci.* **18**, 183–192 (2017).

33. Hutchinson, J. B., Uncapher, M. R. & Wagner, A. D. Posterior parietal cortex and episodic

retrieval: convergent and divergent effects of attention and memory. *Learn. Mem. Cold Spring

Harb. N* **16**, 343–356 (2009).

34. Sutterer, D. W., Foster, J. J., Serences, J. T., Vogel, E. K. & Awh, E. Alpha-band oscillations

track the retrieval of precise spatial representations from long-term memory. *J. Neurophysiol.*

122, 539–551 (2019).

35. Sabo, M. & Schneider, D. Pattern reinstatement and attentional control overlap during episodic

long-term memory retrieval. *Sci. Rep.* **12**, 10739 (2022).

36. Atkinson, R. C. & Shiffrin, R. M. Human Memory: A Proposed System and its Control

Processes. in *Psychology of Learning and Motivation* (eds. Spence, K. W. & Spence, J. T.) vol.

2 89–195 (Academic Press, 1968).

37. Baddeley, A. D., Hitch, G. & Allen, R. A multicomponent model of working memory. *Work.*

Mem. State Sci. 10–43 (2021).

38. Cowan, N. What are the differences between long-term, short-term, and working memory?

Prog. Brain Res. **169**, 323–338 (2008).

39. Fukuda, K. & Woodman, G. F. Visual working memory buffers information retrieved from

visual long-term memory. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5306–5311 (2017).

40. Hirschstein, Z. & Aly, M. Long-term memory and working memory compete and cooperate to

guide attention. *Atten. Percept. Psychophys.* **85**, 1517–1549 (2023).

41. Logie, R. H. *Visuo-Spatial Working Memory*. (Lawrence Erlbaum Associates Ltd., Hove, UK,

1995).

42. Liu, B., Nobre, A. C. & van Ede, F. Functional but not obligatory link between microsaccades

and neural modulation by covert spatial attention. *Nat. Commun.* **13**, 3503 (2022).

43. Nobre, A. C. & Stokes, M. G. Premembering Experience: A Hierarchy of Time-Scales for

Proactive Attention. *Neuron* **104**, 132–146 (2019).

44. Nobre, A. & Stokes, M. Memory and attention: the back and forth. *Cogn. Neurosci.* 291–300

(2020).

45. D'Esposito, M. & Postle, B. R. The Cognitive Neuroscience of Working Memory. *Annu. Rev.*

Psychol. **66**, 115–142 (2015).

46. Nobre, A. C. *et al.* Orienting Attention to Locations in Perceptual Versus Mental

Representations. *J. Cogn. Neurosci.* **16**, 363–373 (2004).

47. Wallis, G., Stokes, M., Cousijn, H., Woolrich, M. & Nobre, A. C. Frontoparietal and Cingulo-

opercular Networks Play Dissociable Roles in Control of Working Memory. *J. Cogn. Neurosci.*

27, 2019–2034 (2015).

48. Olivers, C. N. L., Peters, J., Houtkamp, R. & Roelfsema, P. R. Different states in visual

working memory: when it guides attention and when it does not. *Trends Cogn. Sci.* **15**, 327–

334 (2011).

49. Soto, D., Hodsoll, J., Rotshtein, P. & Humphreys, G. W. Automatic guidance of attention from

working memory. *Trends Cogn. Sci.* **12**, 342–348 (2008).

50. Kuo, B.-C., Rao, A., Lepsien, J. & Nobre, A. C. Searching for Targets within the Spatial Layout

of Visual Short-Term Memory. *J. Neurosci.* **29**, 8032–8038 (2009).

51. Pertzov, Y. & Husain, M. The privileged role of location in visual working memory. *Atten.*

Percept. Psychophys. **76**, 1914–1924 (2014).

52. Schneegans, S. & Bays, P. M. Neural Architecture for Feature Binding in Visual Working

Memory. *J. Neurosci.* **37**, 3913–3925 (2017).

53. Treisman, A. & Zhang, W. Location and Binding in Visual Working Memory. *Mem. Cognit.*

34, 1704–1719 (2006).

54. Sawetsuttipan, P. *et al.* Perceptual Difficulty Regulates Attentional Gain Modulations in

Human Visual Cortex. *J. Neurosci.* **43**, 3312–3330 (2023).

55. Watanabe, T. & Sasaki, Y. Perceptual learning: Toward a comprehensive theory. *Annu. Rev.*

Psychol. **66**, 197–221 (2015).

56. Conway, C. M. How does the brain learn environmental structure? Ten core principles for

understanding the neurocognitive mechanisms of statistical learning. *Neurosci. Biobehav. Rev.*

112, 279–299 (2020).

57. Balaban, H., Assaf, D., Arad Meir, M. & Luria, R. Different features of real-world objects are

represented in a dependent manner in long-term memory. *J. Exp. Psychol. Gen.* **149**, 1275

(2020).

58. Shimi, A. & Logie, R. H. Feature binding in short-term memory and long-term learning. *Q. J.*

Exp. Psychol. **72**, 1387–1400 (2019).

59. Sanders, D. M. W. & Cowell, R. A. The locus of recognition memory signals in human cortex

depends on the complexity of the memory representations. *Cereb. Cortex* **33**, 9835–9849

(2023).

60. Hayhoe, M. M., Bensinger, D. G. & Ballard, D. H. Task constraints in visual working memory.

Vision Res. **38**, 125–137 (1998).

61. Henderson, J. M. & Hollingworth, A. Eye movements and visual memory: Detecting changes

to saccade targets in scenes. *Percept. Psychophys.* **65**, 58–71 (2003).

62. Draschkow, D., Kallmayer, M. & Nobre, A. C. When Natural Behavior Engages Working

Memory. *Curr. Biol.* **31**, 869-874.e5 (2021).

63. Committeri, G. *et al.* Reference frames for spatial cognition: different brain areas are involved

in viewer-, object-, and landmark-centered judgments about object location. *J. Cogn. Neurosci.*

16, 1517–1535 (2004).

64. Moraresku, S. & Vlcek, K. The use of egocentric and allocentric reference frames in static and

dynamic conditions in humans. *Physiol. Res.* **69**, 787–801 (2020).

65. Kerkhoff, G. Spatial hemineglect in humans. *Prog. Neurobiol.* **63**, 1–27 (2001).

66. Aagten-Murphy, D. & Bays, P. M. Independent working memory resources for egocentric and

allocentric spatial information. *PLOS Comput. Biol.* **15**, e1006563 (2019).

67. Golomb, J. D. & Kanwisher, N. Higher Level Visual Cortex Represents Retinotopic, Not

Spatiotopic, Object Location. *Cereb. Cortex* **22**, 2794–2810 (2012).

68. Shafer-Skelton, A. & Golomb, J. D. Memory for retinotopic locations is more accurate than

memory for spatiotopic locations, even for visually guided reaching. *Psychon. Bull. Rev.* **25**,

1388–1398 (2018).

69. Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

70. Faul, F., Erdfelder, E., Lang, A.-G. & Buchner, A. G*Power 3: a flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. *Behav. Res. Methods* **39**,

175–191 (2007).

71. Li, A. Y., Liang, J. C., Lee, A. C. H. & Barense, M. D. The validated circular shape space:

Quantifying the visual similarity of shape. *J. Exp. Psychol. Gen.* **149**, 949–966 (2020).

72. de Vries, E., Fejer, G. & van Ede, F. No obligatory trade-off between the use of space and time

for working memory. *Commun. Psychol.* **1**, 1–10 (2023).

73. Maris, E. & Oostenveld, R. Nonparametric statistical testing of EEG- and MEG-data. *J.*

Neurosci. Methods **164**, 177–190 (2007).

74. Bridges, D., Pitiot, A., MacAskill, M. R. & Peirce, J. W. The timing mega-study: comparing a

range of experiment generators, both lab-based and online. *PeerJ* **8**, e9414 (2020).

Acknowledgements

This research was funded by a Clarendon Scholarship, a Medical Research Council Studentship,

and a New College-Yeotown Scholarship to D.G.; a Wellcome Trust Senior Investigator Award

(104571/Z/14/Z) and a James S. McDonnell Foundation Understanding Human Cognition

Collaborative Award (220020448) to A.C.N.; and by the NIHR Oxford Health Biomedical

Research Centre. The Wellcome Centre for Integrative Neuroimaging is supported by core funding

from the Wellcome Trust (203139/Z/16/Z). For the purpose of open access, the author has applied

a CC BY public copyright licence to any Author Accepted Manuscript version arising from this

submission.

Author contributions

767   D.G., D.D. and A.C.N. conceived and designed the experiments. D.G. acquired and analysed the

768   data. D.G., D.D. and A.C.N. interpreted the data. D.G., D.D. and A.C.N. drafted and revised the

769   manuscript.

Competing Interests

772   The authors declare no competing interests.

Fig. 1. LTM training in Experiment 1. On every learning trial, participants memorized the same
colours and locations of two squares at diagonal locations. When prompted, they reproduced the
colour at the probed location or the location of the probed colour. Colour reproduction trials and
location reproduction trials were presented in random order. Learning performance was aggregated
into 4 bins (of 10 trials each) and shown in the form of errors for colour reproduction and accuracy
for location reproduction, respectively. Error bars represent ±1 *SEM* (*n* = 30).

Fig. 2. Behavioural results in Experiment 1. a, task schematic of the testing trials in Experiment 1. Participants first encoded two items into WM, with different colours and locations from those in LTM. After the first delay, neutral retrocues were uninformative, whereas informative retrocues (WM or LTM) indicated which location would be relevant in the memory retrieval task. After a second delay, the memory retrieval task and the perceptual discrimination task were equally likely to be presented. In the retrieval task, participants retrieved the location of a randomly chosen colour on neutral-retrocue trials and the location of the retrocued colour on informative-retrocue trials. In the perceptual task, participants discriminated the direction of one of the four arrows that briefly appeared at the four quadrants before being masked. The location of the probed arrow was indicated by a post-cue. **b**, memory retrieval mean RTs and accuracy for WM and LTM items, grouped by neutral-retrocue and informative-retrocue trials. **c**, perceptual discrimination accuracy at WM and LTM locations, grouped by retrocue neutral and matching trials. **d**, means and distributions of perceptual accuracy benefits for WM and LTM locations matching retrocues. Error bars in **b-d** represent ± 1 *SEM* ($n = 30$).

Fig. 3. LTM training in Experiment 2. On every learning trial, participants memorized two
consistent items at diagonal locations defined by unique colours and shapes. When prompted, they
reproduced the colour or shape at one location. Colour reproduction trials and shape reproduction
trials appeared in random order. Learning performance was sorted into 4 bins (with 10 trials each)
and shown in the form of errors for both colour and shape reproduction. Error bars represent ±1
SEM (*n* = 44).

Fig. 4. Behavioural results in Experiment 2. a, task schematic of the testing trials in Experiment
2. Participants first encoded two items in WM. Their colours, shapes, and locations differed from
those in LTM. After the first delay, neutral retrocues were uninformative, whereas informative
retrocues (WM or LTM) indicated which item shape would be relevant in the memory retrieval
task. After a second delay, the memory retrieval task and the perceptual discrimination task were
equally likely to be presented. In the retrieval task, participants retrieved the shape of the item
matching a randomly chosen colour on neutral-retrocue trials and the shape of the item matching
the retrocued colour on informative-retrocue trials. In the perceptual task, participants
discriminated the direction of one of the four arrows that briefly appeared at the four quadrants
before being masked. The location of the probed arrow was indicated by a post-cue. **b**, memory
retrieval mean RTs and accuracy for WM and LTM items, grouped by neutral-retrocue and
informative-retrocue trials. **c**, perceptual discrimination accuracy at WM and LTM locations,
grouped by retrocue neutral and matching trials. **d**, means and distributions of perceptual accuracy
benefits for WM and LTM locations matching retrocues. Error bars in **b-d** represent ± 1 *SEM* (*n* =
44).

Fig. 5. Eye-tracking results in Experiment 2. a, time courses of horizontal gaze biases following
neutral, WM, and LTM retrocues. For WM and LTM retrocues, time courses are shown separately
for conditions in which the retrocued item occupied the left side (top/bottom left location) or the
right side (top/bottom right location) during encoding. **b**, time courses of horizontal gaze bias
towardness on trials with WM and LTM retrocues, corresponding to the data in **a**. **c**, time courses
of vertical gaze biases following neutral, WM, and LTM retrocues. For WM and LTM retrocues,
time courses were shown separately for conditions in which the retrocued item occupied the top
side (top left/right location) or the bottom side (bottom left/right location) during encoding. **d**, time

courses of vertical gaze bias towardness on trials with WM and LTM retrocues, corresponding to
the data in **c**. **e-f**, time courses of toward vs. away microsaccade rate in the horizontal channel,
following WM and LTM retrocues respectively. Horizontal lines below the time courses in **b**, **d**,
and **e** indicate significant temporal clusters (green lines: WM compared to zero; black lines in **b**
and **d**: WM vs. LTM difference; black line in **e**: toward vs. away difference).

Fig. 6. Behavioural results in Experiment 3. a, task schematic of the testing trials in Experiment
3, where only the perceptual task was different from Experiment 2. In this new task, participants
identified the direction of the only arrow appearing at one of the four quadrants. **b**, LTM learning
curves during the first day of training. Colour reproduction (left panel) and shape reproduction

(right panel) errors were sorted into 4 bins (with 10 trials each). **c**, memory retrieval mean RTs and
accuracy for WM and LTM items, grouped by neutral-retrocue and informative-retrocue trials. **d**,
perceptual identification accuracy at WM and LTM locations, grouped by retrocue neutral and
matching trials (left panel); means and distributions of perceptual accuracy benefits for WM and
LTM locations matching retrocues (right panel). Error bars in **b-d** represent ± 1 *SEM* (*n* = 119).

850