Title: Components of Isolated Skeletal Muscle Differentiated Through Antibody Validation

Authors and Affiliations: Dominique C. Stephens[1,2*], Margaret Mungai[3*], Amber Crabtree[1*,] Heather K. Beasley[1], Edgar Garza-Lopez[3], Larry Vang[1], Kit Neikirk[1], Zer Vue[1], Neng Vue[1], Andrea G. Marshall[1], Kyrin Turner[2], Jian-qiang Shao[4], Bishnu Sarker[5], Sandra Murray[6], Jennifer A. Gaddy[7,8], Jamaine Davis[9], Steven M. Damo[2#], Antentor O. Hinton Jr[1#]

1. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA

2. Department of Life and Physical Sciences, Fisk University, Nashville, TN, 37232, USA

3. Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA

4. Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA

5. School of Applied Computational Sciences, Meharry Medical College, Nashville, TN, 37232, USA

6. Department of Cell Biology, College of Medicine, University of Pittsburgh, Pittsburgh, TN, 15260, USA

7. Division of Infectious Diseases, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

8. Tennessee Valley Healthcare Systems, U.S. Department of Veterans Affairs, Nashville, Tennessee, USA

9. Department of Biochemistry and Cancer Biology. Meharry Medical College, Nashville, TN, USA

*Co-first Authors

[#]Corresponding Authors:

Steven Damo, PhD
Fisk University
sdamo@fisk.edu

Antentor O. Hinton, Jr, PhD
Vanderbilt School of Medicine Basic Sciences
antentor.o.hinton.jr@Vanderbilt.edu

Abstract

Isolation of skeletal muscles allows for the exploration of many complex diseases. Fibroblasts
and myoblast play important roles in skeletal muscle morphology and function. However,
skeletal muscles are complex and made up of many cellular populations and validation of these
populations is highly important. Therefore, in this article, we discuss a comprehensive method to
isolate mice skeletal muscle, create satellite cells for tissue culture, and use immunofluorescence
to validate our approach.

Novel imaging technology is increasing our ability to visualize and analyze cellular organelles
and compartments.

Tweetable Abstract

44 Proper antibody validation of cellular populations within isolated skeletal muscle can lead to
45 better elucidation of skeletal muscle structure and function, and their roles in complex diseases.

Graphical Abstract

Method Summary

56 Myoblasts are isolated from mouse limb muscles. They are plated for immunofluorescence-based
57 validation using confocal microscopy. This method demonstrates the need for reliable antibodies
58 to correctly determine and differentiate between cellular populations within isolated skeletal
59 muscles.

60 **Keywords:** Skeletal muscles, antibody validation, myoblast, fibroblast, immunofluorescence

Introduction

Skeletal muscles allow for animals and humans to be mobile [1]. Defects in skeletal muscle (SkM) mass can cause atrophy and other pathological diseases [2]. Since the first description of skeletal muscle diseases [3], there have been numerous discoveries describing their pathology and the next step in studying these pathologies is characterizing the different cellular populations residing within them. Isolating cells from these muscles allows for models to develop more complex studies to understand how these pathological mechanisms work. In addition to muscle diseases, skeletal muscles are also used to study immunological, neuronal, and other chronic diseases [4]. Specifically, skeletal muscle cells are essential for studies on exercise and insulin stimulation. They are also useful experimental model to answer more complex questions, such as the effects of insulin stimulation [5] on organelle morphology and the efficacy of new microscopy methods like Focused Ion Beam Scanning Electron Microscopy (FIB-SEM)[6].

Here we offer two aims, firstly to show how to develop isolate myoblasts, or differentiated myotubes, from murine skeletal muscle (**Figure 1**). Secondly, developing antibody-based approaches for validating SkM cells has been a challenge. Here we also offer a technique for myoblast validation.

Antibodies are useful for validating different populations of skeletal muscle cells. Antibodies allow researchers to study the diversity of muscle fibers and cells while providing important insights into cellular processes and disease development. Here, we listed common antibodies used to study different cell populations in SkM tissue (**Table 1**).

SkM tissue is composed of various cell types with different functions, including myoblasts and fibroblasts [2]. Skeletal myoblasts drive muscle regeneration after injury, while fibroblasts create extracellular matrix components and secrete growth factors [7] (**Figure 2**). Morphologically, fibroblasts are larger than myoblasts and contain more vesicles [8]. Given that these populations have morphological differences [9], validating that myoblast or myotube differentiation is successful is of critical importance, especially for experiments that seek to study homogenous populations and fine ultrastructural changes. Antibodies and fluorescence light microscopy can be used to validate different cell populations in skeletal muscle tissue (**Figure 3A-D**). Together, here we propose a standardized approach to isolate and identify different skeletal muscle cell populations.

Before you Begin:

Initial PBS Wash mixture:

Name	Volume
PBS	25 mL
Fungizone	75 µL
Penicillin-Streptomycin	250 µL

Initial DMEM-F12 incubation mixture:

Name	Volume
DMEM-F12	250 mL DMEM (+ 4.5 g/L D-glucose, + L-Glut, -

Name	Volume
	Sodium Pyruvate) + 250 mL F-12 (+L-Glut)
Collagenase II	1300 mg
Penicillin-Streptomycin	6.4 mL
Fungizone	2.0 mL

100   Secondary DMEM-F12 incubation mixture:

Name	Volume
DMEM-F12	250 mL DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate) + 250 mL F-12 (+L-Glut)
Collagenase II	650 mg
Penicillin-Streptomycin	6.4 mL
Fungizone	2.0 mL
Dipase	325 mg

102   DMEM-F12 Growth Media:

103   Mix the following. Use a sterile filter with Millipore brand 0.22 uM filter units. Store at 4C for
104   no longer than 2 months. Add bFGF (10ng/mL) to the aliquot just before adding it to plate.

Name	Volume	Example Catalog
DMEM-F12	250 mL DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate) + 250 mL F-12 (+L-Glut)	Gibco 11965-092 Gibco 11765-054
FBS Note: Do not heat inactivate FBS. Just thaw, swirl to mix, and go.	129 mL	Atlanta Biologicals S11550
Penicillin-Streptomycin	6.4 mL	Gibco 15140
Fungizone	2.0 mL	Gibco 15290-018
MEM Non-essential Amino Acids	6.4 mL	Gibco 11140

Beta-Mercaptoethanol	6.4 µL	Gibco 21985-023

107 Permeabilization Buffer:

Name	Volume
PBS	495.5 mL
Triton X-100	0.5 mL

110 Differentiation medium:

Name	Volume
DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate)	250 mL
F-12 (+L-Glut)	250 mL
FBS Note: Do not heat inactivate FBS. Just thaw, swirl to mix, and go.	10.5 mL
Insulin-transferrin-selenium-X (100x)	5.3 mL

Reconstitute Human FGF-basic (FGF-2/bFGF) Recombinant Protein (here we use ThermoFisher
13256-029). Briefly, to prepare a stock solution of bFGF at a concentration of 0.1 mg/mL,
reconstitute it in 100 µL of 10 mM Tris (pH 7.6). Dilute in buffer containing 0.1% BSA and
store in polypropylene vials for up to six months at -20°C. Avoid freezing and rethawing.

Guide:

Myoblast Isolation

1. Collect muscle tissue from the gastrocnemius, quadriceps, and hamstring muscles at 4-8
weeks of age from mice.
2. Wash Isolated tissue 2-3 times with Initial PBS Wash Mixture.
122
123 **NOTE**: The PBS solution is prepared right before dissecting the tissue.

124   3. Incubate muscle tissue in the initial DMEM-F12 incubation mixture.

CRITICAL: Avoid filtering the DMEM-F12 media containing collagenase, 1% pen/strep, and 3
μL/mL Fungizone. This solution must be added cold to reduce temperature shock.

4. Maintain the muscle solution in a 37 °C water bath for 10-15 mins.
5. Shake at 220 rpm, for an overall time of 1.5 hrs.
6. After incubation, wash the tissue 3-4 times with PBS.
7. Incubate in warm the initial DMEM-F12 incubation mixture while the tissue is shaken for
    30 mins in a 37 °C water bath.

NOTE: Solution has to be pre-warmed to 37°C to ensure efficient mixing of dispase and as the
muscles were at 37°C after incubation

8. After shaking, ground tissue with mortar and pestle in the presence of liquid nitrogen.
9. Pass through a 100 μm, then 70μm, cell strainer.
10. Centrifuge the solution at 1000 rpm for five mins to pellet the cells.
11. Transfer the to a plate and resuspended using DMEM-F12 growth media supplemented
    with 40 ng/mL bFGF.
12. Pre-plate the cells for 1-3 hours on UNCOATED dishes to reduce the number of
    fibroblasts.

CRITICAL: Fibroblasts can dilute satellite cells. Recommended for dystrophic or injured
muscle. Pre-plating on an uncoated plate causes fibroblasts to stick and be isolated. Fibroblasts
can separately be used to isolate and for other experiments.

13. Dilute cells 1:15 in PBS, then plate in a Matrigel-coated dish.

NOTE: To create Matrigel-coated dishes, dilute stock concentration (while keeping on ice) to
1:15 in sterile PBS in the hood. Put Matrigel solution on flask/plate, shake/tilt to coat the bottom,
incubate at room temperature in hood for 30 mins, and remove Matrigel solution back into its
original tube. Matrigel solution may be reused up to 5 times total.

14. Wait for activation, which takes 24-48 hrs, after which myoblasts will grow rapidly.
    a. To maintain healthy myoblast cells, use the growth media supplemented with
        bFGF (10 ng/mL).

CRITICAL: Use Differentiation Medium to go from myoblasts to myocytes and then to
myotubes (Figure 2).

15. Plate primary myoblast at ~.8 X 106 cells per well and to differentiate the cells, add
    differentiation media, supplemented with 1:10,000 bFGF.

NOTE: This will depend on # of cell passages and type of treatment, adjust accordingly.

164   16. Incubate for 4 to 5 days.
165

166   **NOTE**: Switch out with fresh differentiation media every 2 days, supplemented with 1:10,000
167   bFGF.

168   17. Cells are split using accutase.
169
170   **Note: DO NOT** use trypsin to split the cells. Accutase is less harsh to the extracellular matrix,
171   surface proteins, and cytoskeleton of skeletal cells than trypsin, so it is highly preferred [10].

172   18. Cells are maintained in a hypoxic environment (5% O_2) at 37°C. If growing myotubes, A
173   confluency of 70-85% has to be reached prior to adding growth media.
174

Myoblast Validation

Immunofluorescence staining is effective for examining differences in skeletal muscles
simultaneously. Refer to Table 1 for a list of validated primary antibodies for skeletal muscles.
Select secondary antibodies that are compatible with the epifluorescence or confocal microscope
available to you.

CRITICAL: All steps are performed at room temperature unless otherwise indicated.

NOTE: This protocol for Immunofluorescence staining and antibody validation of isolated
skeletal muscle cells is an adaptation of Esper et al., skeletal muscle tissue immunofluorescence
labeling protocol [11].

1. To prepare the cells for fluorescence microscopy, the cells fix them by incubating them in
4% PFA for five mins.
2. Wash three times for five mins using PBS.
187

188   **NOTE:** Ice-cold 100% methanol or acetone is an effective fixative for cryosections and more
189   suited for some antigens. Acetone is less harsh than methanol.

190   3. Incubate cells in permeabilization buffer for 10 mins.
191   4. Incubate cells in blocking solution for 1 hr at room temperature or overnight at 4 °C.
192

NOTE: When using permeabilization buffer, keep the solution away from the hydrophobic
barrier to avoid loss of hydrophobicity. If this happens, wash the slide well with PBS. Include
Mouse on Mouse (MOM) blocking reagent at a 1:40 dilution when staining mouse tissue with
antibodies raised in the mouse.

5. To begin immunostaining, dilute the primary and secondary antibodies in a blocking
solution according to the manufacturer's suggested ratio.
199
NOTE: It is acceptable to dilute antibodies in hybridoma supernatant when targeting multiple
antigens.

6. Aspirate the blocking buffer and cover the slide with the primary antibody solution.
7. Incubate the slides overnight at 4 °C.
8. On the following day, wash three times for five mins with PBS.
9. After washing, cover cells with secondary antibodies diluted in blocking buffer for 1 hr at
room temperature in the dark.
207
NOTE: Keep slides in the dark for the remainder of the protocol.

10. After incubation, wash the slides three times for five mins with PBS.
11. Incubate the cells with 1 μg/mL DAPI diluted in PBS for five mins.
12. Wash once with PBS for five min.
13. Aspirate the PBS and place 1–2 drops of mounting media on to the cells
14. Carefully place a coverslip on the slide, while avoiding air bubbles.
15. Let the slides dry in the dark for 1–2 hr before sealing the slides with clear nail polish.
16. Store the slides at 4 °C and image within 2 weeks.
216
Expected Outcomes:

Upon isolation of myoblast and myotube, we validated their structure in light microscopy
(Figure 3A). Furthermore, we viewed multinucleated myotubes through TEM to validate that
ultrastructure was as expected (Figure 3B). Transfection further showed myotubes demonstrated
fluorescence as expected (Figure 3C). From there, we performed staining for myosin and desmin,
muscle-specific proteins that play crucial roles in muscle cell structure and function [12], to
confirm that filaments were present as expected (Figure 3 D-D'').

Once myoblasts and myotubes are validated, they can be used for a variety of studies
including to measure mitochondrial efficiency with oxygen consumption rate, western blot
analysis to look for expression of specific proteins in knockout studies, or a variety of electron
microscopy techniques such as serial block-face scanning-electron microscopy to perform 3D
reconstruction of organelles [13] (Figure 4). As an example, to validate this method, we sought to
understand how insulin treatment (10 nM/L) in 2-hour increments may alter myoblast and
myotube function through the usage of a Seahorse XF96 analyzer, per past protocols [14].

To begin with, time points 1-3 measure basal respiration, or baseline rate of oxygen
consumption by cells in culture without any treatment. Oligomycin (1 μg/ml) was added to
inhibit ATP synthase, which reduces mitochondrial respiration and leads to an increase in proton
gradient, to measure the amount of oxygen consumed by the myoblasts and myotubes to
maintain the proton gradient in time points 4-6. carbonyl cyanide 4-
(trifluoromethoxy)phenylhydrazone (FCCP; 1 μM) was then added in time points 7-9 which
allows electrons to flow freely through the chain for reserve capacity and maximum oxygen
consumption to be measured. Finally, rotenone (1 μM) and antimycin A (10 μM) were added in
time points 10-12 which inhibit electron transfer from NADH to ubiquinone and ubiquinol to
cytochrome c, respectively, to measure non-mitochondrial respiration [15].

We found that for myoblasts, there is a significantly increased basal, maximum, and non-
mitochondrial OCR after 2 hours of insulin treatment, while this difference is retained or

exacerbated after 4 hours of insulin treatment (Figure 5A-B). After 6 hours of insulin treatment, OCR conversely showed significant decreases in all of these parameters (Figure 5C). In myotubes, after 2 and 4 hours of insulin treatment, we similarly noted a significant increase in mitochondrial OCR (Figure 5D-E). Notably, the increase in basal, ATP-linked, maximum, and non-mitochondrial OCR is much higher in 4 hours than 2 hours. Unlike myoblasts, 6 hours of insulin treatment myotubes did not differ significantly from untreated cells (Figure 5F). Importantly, there may be a differential response to insulin treatment in myoblasts and myotubes, highlighting the importance of studying both models. This validated that the function of myoblasts and myotubes are intact following this isolation.

From there we sought to elucidate if organelle proteins are affected following insulin treatment and we targeted Optic atrophy protein 1 (OPA-1), which is a mitochondrial inner membrane (IMM) fusion protein that mediates the fusion of the IMM between two mitochondria while also serving roles in mitochondrial bioenergetics and cristae architecture [17]. OPA-1 is just one of several proteins which modulate mitochondrial structure. For example, contrastingly, Dynamin-related protein 1 (DRP-1) is a protein which initiates the fission process through constriction of the mitochondria which divides the mitochondria into two separate organelles [18]. However, given that OCR increased following insulin treatment, it is possible this is due to increased mitochondrial area caused by upregulated mitochondrial fusion. To see if OPA-1 may be changed in expression, we performed western blotting. When looking at OPA-1, we noticed a significant continuous increase in protein levels in myoblasts across 2 and 4 hours of insulin stimulation when normalized (Figure 6A-B). We further differentiated primary myotubes and carried out these experiments again to see if any differences existed (Figure 6C-D). We noticed significant increases in OPA-1 levels after 4 hours of insulin stimulation (Figure 6C-D). Together, this suggests that insulin stimulation causes increased expression of OPA-1 in a short time frame which is exacerbated in myotubes compared with myoblasts. These results are suggestive that mitochondrial fusion may be a compensatory of insulin stimulation which occurs through OPA-1-mediated mechanisms in myoblasts and myotubes. Together these data validate this isolation and validation technique allows for the application of experimental models to elucidate cellular processes.

Quantification and Statistical Analysis:

After differentiation, quantification can be done for many experimental designs. Here, we performed seahorse analysis per prior methods [14] with GraphPad Prism version 8.4.0 (GraphPad Software, La Jolla, CA) was used to perform students' T-tests to measure statistical significance.

Limitations: This protocol has been optimized for mice gastrocnemius, quadriceps, and hamstring muscles and may not be applicable to other model organisms or tissue types. Compared with other protocols, ours takes a similar period of time [19], but this can still be a slow process that must be carried out across multiple days. While C2C12 myoblasts are ideal for this protocol, increasingly human skeletal myoblasts are important to study and past protocols

282 indicate that differences in the procedure must be made, such as antisense miR-133a addition, to
283 promote the fast differentiation of human skeletal myoblasts [20].

Trouble Shooting:

Problem: Ultrastructure or Gross morphology of Myoblasts are Degraded

Potential Solution: This may be due to too much damage incurred to myoblasts during
preparation. Here, we found that tissue should first be digested with type II collagenase and
dispase, then ground by being put in liquid nitrogen with a mortar with a pestle, and finally
passed through cell strainers optimizes this procedure. However, reducing the time grounded or
reducing the amount of digestion can avoid potential damage to the myoblasts if it is occurring.

Problem: Contamination with Fibroblasts

Potential Solution: It is important to plate first on an uncoated plate. However, if fibroblasts are
still observed, pre-plating can be done twice. Antibody-based selection of fibroblast may cause
certain issues but can also be explored as an option to remove fibroblasts. If this remains an
issue, other methods have shown that using flowing cytometry can be used to identify and
remove fibroblasts [21].

Problem: Low Cell Yield or Viability

Potential Solution: If myoblast or myotube viability is low increasing the concentration of
growth factors and assuming a sterile environment is attained is important. Reducing time with
accutase can also ensure cells are not treated too harshly.

Resource Availability:

Lead contact
Further information and requests for resources and reagents should be directed to and will be
fulfilled by the lead contact, Antentor Hinton (antentor.o.hinton.jr@Vanderbilt.Edu).

Materials availability
All generated materials, if applicable, are created in methods highlighted in the text above.

Data and code availability
Full data utilized and requests for data and code availability should be directed to and will be
fulfilled by the lead contact, Antentor Hinton (antentor.o.hinton.jr@Vanderbilt.Edu).

Author Contributions

Acknowledgements

All antibodies were obtained from the Iowa Developmental Studies Hybridoma Bank (DSHB).

Financial & Competing Interests' Disclosure

All authors have no competing interests.

This project was funded by the UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, BWF Career Awards at the Scientific Interface Award, BWF Ad-hoc Award, NIH Small Research Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program, DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation (to A.H.J.). NSF EES2112556, NSF EES1817282, and CZI Science Diversity Leadership grant number 2022-253614 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation (to S.M. D.) and National Institutes of Health grant HD090061 and the Department of Veterans Affairs Office of Research award I01 BX005352 (to J.G.). Additional support was provided by the Vanderbilt Institute for Clinical and Translational Research program supported by the National Center for Research Resources, Grant UL1 RR024975–01, and the National Center for Advancing Translational Sciences, Grant 2 UL1 TR000445–06 and the Cell Imaging Shared Resource.

Data Sharing and Open Access

All data is available upon request to the corresponding author.

Figures and Legends

344 **Figure 1: The process of myoblast isolation from gastrocnemius muscle.**

Figure 2: The process of myotube differentiation from myoblasts and utilization for serum

Figure 3: Myotubes as characterized by A. Light microscopy, B. Transmission electron microscopy, and C. Transfection with adenovirus containing the green fluorescent protein gene (Ad-GFP). D. Straining with BA-D5-s, D' SC-71-s, and D'' D3-s to show myosin and desmin.

Figure 4: Examples of experiments that may be performed following myotube differentiation and isolation.

Figure 5: Oxygen consumption rate (OCR) altered in myoblasts and myotubes upon altered insulin stimulation which shows changes in mitochondrial efficiency. (A) Seahorse plot for primary myoblasts following 2 hours of insulin stimulation (B) 4 hours of insulin stimulation and (C) 6 hours of insulin stimulation. (D) Oxygen consumption rate was measured after several inhibitors to measure respiration in primary myotubes after 2 hours, (E) 4 hours and (F) 6 hours of insulin stimulation. (A'-F') Basal OCR which represents respiration under normal, unstressed conditions. (A''-F'') ATP-linked OCR, which is respiration associated with ATP synthesis during oxidative phosphorylation which is marked by a reduction in OCR due to oligomycin. (A'''-F''') Maximum OCR which is the maximal capacity at which mitochondria may utilize oxygen. (A''''-F'''') Non-mitochondrial respiration which can be attributed to factors such as glycolysis or ROS and not due to mitochondrial respiration. These values were compared to the control (blue) in all of these examples. N = 6 per treatment, and * indicates p-value < .05.

Figure 6: Comparison of mitochondrial fusion proteins following insulin stimulation in primary
myoblasts and myotubes. (A) Western blotting for mitochondrial fusion protein OPA-1
following 2 hours, 4 hours, and 6 hours of insulin stimulation. (B) OPA-1 levels normalized to
Alpha tubulin following insulin stimulation. (C) This was replicated in primary myotubes, as
western blotting for mitochondrial fusion OPA1. (D) OPA-1 levels, normalized to Alpha tubulin,
in primary myotubes following insulin treatment. N = 6 per treatment, and * indicates p-value <
.05.

Antibody Name	Antigen	IF/IHC Rec
BF-F3-s	Myosin heavy chain Type IIB	2-5 ug/ml
SC-71-s	Myosin heavy chain Type IIA	2-5 ug/ml
BA-D5-s	Myosin heavy chain Type I	2-5 ug/ml
F5D-s	Myogenin	2-5 ug/ml
PAX7-s	Pax7	2.5 ug/ml
D3-s	Desmin, intermediate filament	2.5 ug/ml
9.1 ITGA7-s	Integrin alpha-7, extracellular domain	2-5 ug/ml
JLT12-s	troponin T, fast skeletal muscle specific	2.5 ug/ml

Table 1: A list of antibodies and their respective antigens for skeletal muscle validation after isolation. The list was obtained from https://dshb.biology.uiowa.edu/.

References

1. Figueiredo, P.A., Mota, M.P., Appell, H.J., and Duarte, J.A. (2008). The role of mitochondria in aging of skeletal muscle. Biogerontology *9*, 67–84. 10.1007/s10522-007-9121-7.

2. Dave, H.D., Shook, M., and Varacallo, M. (2022). Anatomy, Skeletal Muscle. In StatPearls [Internet] (StatPearls Publishing).

3. Walusinski, O. (2022). François-Amilcar Aran (1817-1861) and the recognition of spinal muscular atrophy. Rev Neurol (Paris) *178*, 756–765. 10.1016/j.neurol.2022.01.011.

4. Morgan, J., and Partridge, T. (2020). Skeletal muscle in health and disease. Disease Models
& Mechanisms *13*, dmm042192. 10.1242/dmm.042192.

5. Alonge, K.M., Meares, G.P., and Hillgartner, F.B. (2017). Glucagon and Insulin
Cooperatively Stimulate Fibroblast Growth Factor 21 Gene Transcription by Increasing the
Expression of Activating Transcription Factor 4 *. Journal of Biological Chemistry *292*,
5239–5252. 10.1074/jbc.M116.762922.

6. Marshall, A.G., Damo, S.A., and Hinton, A. (2023). Revisiting focused ion beam scanning
electron microcopy. Trends Biochem Sci, S0968-0004(23)00056-7.
10.1016/j.tibs.2023.02.005.

7. Thummarati, P., and Kino-Oka, M. (2020). Effect of Co-culturing Fibroblasts in Human
Skeletal Muscle Cell Sheet on Angiogenic Cytokine Balance and Angiogenesis. Front
Bioeng Biotechnol *8*, 578140. 10.3389/fbioe.2020.578140.

8. Yablonka-Reuveni, Z., Anderson, S.K., Bowen-Pope, D.F., and Nameroff, M. (1988).
Biochemical and morphological differences between fibroblasts and myoblasts from
embryonic chicken skeletal muscle. Cell Tissue Res *252*, 339–348. 10.1007/BF00214376.

9. Mingueitti, G., and Mair, W.G. (1980). The developing human muscle: ultrastructural
differences between myoblasts and fibroblasts. Rev Bras Pesqui Med Biol *13*, 1–8.

10. Nowak-Terpiłowska, A., Śledziński, P., and Zeyland, J. (2021). Impact of cell harvesting
methods on detection of cell surface proteins and apoptotic markers. Braz J Med Biol Res
54, e10197. 10.1590/1414-431X202010197.

11. Esper, M.E., Kodippili, K., and Rudnicki, M.A. (2023). Immunofluorescence Labeling of
Skeletal Muscle in Development, Regeneration, and Disease. Methods Mol Biol *2566*, 113–
132. 10.1007/978-1-0716-2675-7_9.

12. Agnetti, G., Herrmann, H., and Cohen, S. (2022). New roles for desmin in the maintenance
of muscle homeostasis. The FEBS Journal *289*, 2755–2770. 10.1111/febs.15864.

13. Garza-Lopez, E., Vue, Z., Katti, P., Neikirk, K., Biete, M., Lam, J., Beasley, H.K., Marshall,
A.G., Rodman, T.A., Christensen, T.A., et al. (2022). Protocols for Generating Surfaces and
Measuring 3D Organelle Morphology Using Amira. Cells *11*, 65. 10.3390/cells11010065.

14. Pereira, R.O., Marti, A., Olvera, A.C., Tadinada, S.M., Bjorkman, S.H., Weatherford, E.T.,
Morgan, D.A., Westphal, M., Patel, P.H., and Kirby, A.K. (2021). OPA1 deletion in brown
adipose tissue improves thermoregulation and systemic metabolism via FGF21. Elife *10*,
e66519.

15. Rose, S., Frye, R., Slattery, J., Wynne, R., Tippett, M., Pavliv, O., Melnyk, S., and James, S.
(2014). Oxidative Stress Induces Mitochondrial Dysfunction in a Subset of Autism
Lymphoblastoid Cell Lines in a Well-Matched Case Control Cohort. PloS one *9*, e85436.
10.1371/journal.pone.0085436.

16. Chen, H., Detmer, S.A., Ewald, A.J., Griffin, E.E., Fraser, S.E., and Chan, D.C. (2003).
Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for
embryonic development. J Cell Biol *160*, 189–200. 10.1083/jcb.200211046.

17. Barrera, M., Koob, S., Dikov, D., Vogel, F., and Reichert, A.S. (2016). OPA1 functionally
interacts with MIC60 but is dispensable for crista junction formation. FEBS Letters *590*,
3309–3322. 10.1002/1873-3468.12384.

18. Peng, L., Men, X., Zhang, W., Wang, H., Xu, S., Xu, M., Xu, Y., Yang, W., and Lou, J.
(2011). Dynamin-related protein 1 is implicated in endoplasmic reticulum stress-induced
pancreatic β-cell apoptosis. Int J Mol Med *28*, 161–169. 10.3892/ijmm.2011.684.

19. Shahini, A., Vydiam, K., Choudhury, D., Rajabian, N., Nguyen, T., Lei, P., and Andreadis,
S.T. (2018). Efficient and high yield isolation of myoblasts from skeletal muscle. Stem Cell
Research *30*, 122–129. 10.1016/j.scr.2018.05.017.

20. Cheng, C.S., El-Abd, Y., Bui, K., Hyun, Y.-E., Hughes, R.H., Kraus, W.E., and Truskey,
G.A. (2014). Conditions that promote primary human skeletal myoblast culture and muscle
differentiation in vitro. Am J Physiol Cell Physiol *306*, C385–C395.
10.1152/ajpcell.00179.2013.

21. Stellato, M., Czepiel, M., Distler, O., Błyszczuk, P., and Kania, G. (2019). Identification and
Isolation of Cardiac Fibroblasts From the Adult Mouse Heart Using Two-Color Flow
Cytometry. Frontiers in Cardiovascular Medicine *6*.

472