Title: Components of Isolated Skeletal Muscle Differentiated Through Antibody Validation

Authors and Affiliations: Dominique C. Stephens[1,2*], Margaret Mungai[3*], Amber Crabtree[1*,]
Heather K. Beasley[1], Edgar Garza-Lopez[3], Larry Vang[1], Kit Neikirk[1], Zer Vue[1], Neng Vue[1],
Andrea G. Marshall[1], Kyrin Turner[2], Jian-qiang Shao[4], Bishnu Sarker[5], Sandra Murray[6], Jennifer
A. Gaddy[7,8], Jamaine Davis[9], Steven M. Damo[2#], Antentor O. Hinton Jr[1#]
1. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN,
37232, USA
2. Department of Life and Physical Sciences, Fisk University, Nashville, TN, 37232, USA
3. Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA
4. Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA
5. School of Applied Computational Sciences, Meharry Medical College, Nashville, TN, 37232,
USA
6. Department of Cell Biology, College of Medicine, University of Pittsburgh, Pittsburgh, TN,
15260, USA
7. Division of Infectious Diseases, Vanderbilt University School of Medicine, Nashville,
Tennessee, USA
8. Tennessee Valley Healthcare Systems, U.S. Department of Veterans Affairs, Nashville,
Tennessee, USA
9. Department of Biochemistry and Cancer Biology. Meharry Medical College, Nashville, TN,
USA
21
22 *Co-first Authors
23 #Corresponding Authors:
24 Steven Damo, PhD
25 Fisk University
26 sdamo@fisk.edu
27
28 Antentor O. Hinton, Jr, PhD
29 Vanderbilt School of Medicine Basic Sciences
30 antentor.o.hinton.jr@Vanderbilt.edu
31
32
33

Summary

Isolation of skeletal muscles allows for the exploration of many complex diseases. Fibroblasts
and myoblast play important roles in skeletal muscle morphology and function. However,
skeletal muscles are complex and made up of many cellular populations and validation of these
populations is highly important. Therefore, in this article, we discuss a comprehensive method to
isolate mice skeletal muscle, create satellite cells for tissue culture, and use immunofluorescence
to validate our approach.

Graphical Abstract
42
43
44
45

Keywords: Skeletal muscles, antibody validation, myoblast, fibroblast, immunofluorescence

Before you Begin:

Skeletal muscles (SkM) allow for animals and humans to be mobile [1], serving many important roles and constituting nearly half of the total mass of the adult human body [2]. Defects in skeletal muscle mass can cause atrophy and other pathological diseases [3]. Beyond only mediating glucose uptake in an insulin-dependent manner, skeletal muscle also plays important roles in the metabolism and development of diabetes [4]. Since the first description of skeletal muscle diseases [5], there have been numerous discoveries describing their pathology and the next step in studying these pathologies is characterizing the different cellular populations residing within them. Isolating cells from these muscles allows for models to develop more complex studies to understand how these pathological mechanisms work. In addition to muscle diseases, skeletal muscles are also used to study immunological, neuronal, and other chronic diseases [6]. While past studies have used immortalized myogenic cells, myoblasts offer unique advantages to understanding the process of myogenesis, which is an avenue for the repair of injured myofibers [7]. Specifically, skeletal muscle cells are essential for studies on exercise and insulin stimulation. They are also useful experimental models to answer more complex questions, such as the effects of insulin stimulation [8] on organelle morphology and the efficacy of new microscopy methods like Focused Ion Beam Scanning Electron Microscopy (FIB-SEM)[9]. Yet, protocols that allow for the differentiation and isolation of myotubes and myoblasts remain limited.

Here we offer two aims, firstly to show how to develop isolate myoblasts, or differentiated myotubes, from murine skeletal muscle (**Figure 1**). Secondly, developing antibody-based approaches for validating SkM cells has been a challenge. Here we also offer a technique for myoblast validation. Antibodies are useful for validating different populations of skeletal muscle cells. Antibodies allow researchers to study the diversity of muscle fibers and cells while providing important insights into cellular processes and disease development. Here, we listed common antibodies used to study different cell populations in SkM tissue (**Table 1**).

SkM tissue is composed of various cell types with different functions, including myoblasts and fibroblasts [3]. Skeletal myoblasts drive muscle regeneration after injury, while fibroblasts create extracellular matrix components and secrete growth factors [10] (**Figure 2**). Morphologically, fibroblasts are larger than myoblasts and contain more vesicles [11]. Beyond this, while mononuclear cells replicate, as they form sheets of multinucleated myotubes, proliferation is impaired and myogenin is elevated [12]. Myoblasts' process of differentiation mimics that of *in vivo* myogenesis, with the structure of myoblasts affecting that of differentiated myotubes [13]. Given that these populations have morphological differences [14], validating the myoblast or myotube stage is of critical importance, especially for experiments that seek to study homogenous populations and fine ultrastructural changes. Here, we also present how antibodies and fluorescence light microscopy can be used to validate different cell populations in skeletal muscle tissue. However, before you begin care should be taken in experimental design and selection for which populations you want to obtain. Here, we propose a standardized approach to isolate and identify different skeletal muscle cell populations. Using these methods, we looked at the effects on insulin stimulation on oxygen consumption rate (OCR) and have verified past studies which have implicated changes in respiration following insulin stimulation in a potential optic atrophy protein-1 (OPA-1) mediated manner [15].

Methods and Materials:

<u>Animal Studies:</u>
All mice utilized had a C57Bl/6J background. Mice studies followed previous studies with [16–18]
with weaning at 3 weeks of age and maintained on standard chow (2920X Harlan Teklad,
Indianapolis, IN, USA), and at 22°C with a 12 hr light, 12 hr dark cycle with free access to water
and standard chow. All mouse experiments were conducted in alignment with the animal
research guidelines from NIH and were approved by the University of Iowa IACUC.
104
105 <u>Light Microscopy:</u>
106
Staining of myotubes was performed on Olympus IX-81. Myotubes were plated on 35 mm
dishes with a glass bottom and imaged for light microscopy. Transmission electron microscopy
performed with Joel 1400, operating at 80 kV, as we have done previously [17]. 3 days after
differentiation, myotubes were infected with GFP-expressing adenovirus per previous procedures
[16]. All imaging experiments were performed at Central Microscopy Research Facility, University
of Iowa, while adenoviruses were obtained from University of Iowa Viral Vector Core facility.
113
<u>Seahorse Analyzer:</u>
Mice were anesthetized using a mixture of 5% isoflurane/oxygen. Oxygen consumption
rate was measured for using an XF24 bioanalyzer (Seahorse Bioscience: North Billerica, MA,
USA), as previously described [16,19]. Myotubes and myoblasts were plated at a density of 20×10^3
per well and differentiated for 3 days. Isolated myotubes and myoblasts were either treated with
10 nmol/L insulin for the specified time [15]. Media was replaced with XF☐DMEM (supplemented
with 1 g/L D-Glucose, 0.11 g/L sodium pyruvate, and 4 mM L-Glutamine) and cells were
deprived of CO_2 for 60 minutes. Oligomycin (1 μg/ml), carbonyl cyanide 4-
(trifluoromethoxy)phenylhydrazone (FCCP; 1 μM), rotenone (1 μM), and antimycin A (10 μM)
treatment occurred.
For quantifications, time points 1-3 measure basal respiration, or baseline rate of oxygen
consumption by cells in culture without any treatment. Oligomycin (1 μg/ml) was added to
inhibit ATP synthase, which reduces mitochondrial respiration and leads to an increase in proton
gradient, to measure the amount of oxygen consumed by the myoblasts and myotubes to
maintain the proton gradient in time points 4-6. carbonyl cyanide 4-
(trifluoromethoxy)phenylhydrazone (FCCP; 1 μM) was then added in time points 7-9 which
allows electrons to flow freely through the electron transport chain allowing for measurements of
reserve capacity and maximum oxygen consumption. Finally, rotenone (1 μM) and antimycin A
(10 μM) were added in time points 10-12 which inhibit electron transfer from NADH to
ubiquinone and ubiquinol to cytochrome c, respectively, to measure non-mitochondrial
respiration [20].
For normalization of proteins, after measurement 20 μl of 10 mM Tris with 0.1% Triton
X-100 was added at pH 7.4 to lyse cells per prior protocols [19], and media was replaced with 480
μl of Bradford reagent.
138
<u>Transmission Electron Microscopy:</u>
Myoblasts and myotubes were isolated according to the step-by-step below and placed in six-
well poly-D-lysine-coated plates for TEM processing per established protocols [1,2]. Briefly, cells
were fixed by incubating at 37 °C with 2.5% glutaraldehyde in 0.1 m sodium cacodylate buffer
for one hour, then rinsed twice with 0.1 m sodium cacodylate buffer, prior to fixation at room

temperature for 30 min to 1 h using 1% osmium tetroxide and 1.5% potassium ferrocyanide in
0.1 m sodium cacodylate buffer. Samples were washed for 5 min with 0.1 m sodium cacodylate
buffer (7.3 pH), then diH2O (2 X 5 minutes). Samples were incubated with 2.5% uranyl acetate,
diluted with H2O, at 4 °C overnight. Samples were dehydrated and ethanol was replaced with
Eponate 12 mixed in 100% ethanol in a 1:1 solution for 30 min at RT. This was repeated three
times for 1 h using 100% Eponate 12, the media was replaced and plates were cured in an oven
at 70 °C overnight.
After cracking and submerging the plate in liquid nitrogen, an 80 nm thickness jeweler's
saw was used to cut the block to fit in a Leica UC6 ultramicrotome sample holder. From there,
the section was placed on formvar-coated copper grids. These grids were counterstained in 2%
uranyl acetate for 2 min and Reynold's lead citrate for 2 min. Images were acquired by TEM on
either a JEOL JEM-1230, operating at 120 kV, or a JEOL 1400, operating at 80 kV. Analysis
performed via protocols by Lam et al. (2021) [1].
157
158
159 **Key Resource Table:**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
4% paraformaldehyde	ThermoFisher	J61899-AK
Beta-Mercaptoethanol	Gibco	21985-023
Collagenase II	Gibco	17101015
DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate)	Gibco	11965-092
F-12 (+L-Glut)	Gibco	11765-054
FBS	Atlanta Biologicals	S11550
Fungizone	Gibco	15290-018
Human FGF-basic (FGF-2/bFGF) Recombinant Protein	ThermoFisher	13256-029
Insulin-transferrin-selenium-X (100x)	Gibco	41400045
MEM Non-essential Amino Acids	Gibco	11140
PBS, pH 7.4	Gibco	10010023
Penicillin-Streptomycin	Gibco	15140
StemPro™ Accutase™	ThermoFisher	A1110501
Triton X-100 (1%)	Gibco	HFH10
Quantification and Software		

GraphPad	GraphPad Software, San Diego, California USA	www.graphpad.com
Image J	Schindelin et al.5	https://imagej.net/

160
161
162
163
164 Prior to the protocol, make the following reagents and solutions and sterilize all equipment in an
165 autoclave.
166
167 ⏱*Timing: 1 hr*
168
169 Initial PBS Wash mixture:

Name	Volume
PBS	25 mL
Fungizone	75 µL
Penicillin-Streptomycin	250 µL

170
171 Initial DMEM-F12 incubation mixture:

Name	Volume
DMEM-F12	250 mL DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate) + 250 mL F-12 (+L-Glut)
Collagenase II	1300 mg
Penicillin-Streptomycin	6.4 mL
Fungizone	2.0 mL

172
173
174 Secondary DMEM-F12 incubation mixture:

Name	Volume
DMEM-F12	250 mL DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate) + 250 mL F-12 (+L-Glut)
Collagenase II	650 mg
Penicillin-Streptomycin	6.4 mL
Fungizone	2.0 mL

Dispase	325 mg

175
176 DMEM-F12 Growth Media:
177 Mix the following. Use a sterile filter with Millipore brand 0.22 uM filter units. Store at 4C for
178 no longer than 2 months. Add bFGF (10ng/mL) to the aliquot just before adding it to plate.

Name	Volume
DMEM-F12	250 mL DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate) + 250 mL F-12 (+L-Glut)
FBS Note: Do not heat inactivate FBS. Just thaw, swirl to mix, and go.	129 mL
Penicillin-Streptomycin	6.4 mL
Fungizone	2.0 mL
MEM Non-essential Amino Acids	6.4 mL
Beta-Mercaptoethanol	6.4 µL

179
180
181 Permeabilization Buffer:

Name	Volume
PBS	495.5 mL
Triton X-100	0.5 mL

182
183
184 Differentiation medium:

Name	Volume
DMEM (+ 4.5 g/L D-glucose, + L-Glut, - Sodium Pyruvate)	250 mL
F-12 (+L-Glut)	250 mL
FBS Note: Do not heat inactivate FBS. Just thaw, swirl to mix, and go.	10.5 mL
Insulin-transferrin-selenium-X (100x)	5.3 mL

185
186 Reconstitute Human FGF-basic (FGF-2/bFGF) Recombinant Protein (here we use ThermoFisher
187 13256-029). Briefly, to prepare a stock solution of bFGF at a concentration of 0.1 mg/mL,
188 reconstitute it in 100 µL of 10 mM Tris (pH 7.6). Dilute in buffer containing 0.1% BSA and
189 store in polypropylene vials for up to six months at -20°C. Make aliquots to avoid repeated
190 freezing and rethawing.
191

192 **Step-by-Step:**
193 *Myoblast Isolation: Removal of Muscle*
194 The below offers a basic text for the isolation of myoblast.
195
196 🕐*Timing: 30 minutes.*
197

1. Following IACUC guidelines, euthanize mice.
2. Collect muscle tissue from the gastrocnemius, quadriceps, soleus, and hamstring muscles in both legs at 4-8 weeks of age from 4-6 mice.

NOTE: Less or more mice can be utilized but we generally found batches of 4-6 to be a good quantity.

CRITICAL: This protocol works to make tissue-type specific cell lines. If the muscle types will not be combined, double the number of mice, to 8-12 mice, to ensure an adequate sample count.

3. Transfer the collected tissue to an Eppendorf tube.

NOTE: From this point on, ensure a sterile environment is utilized.

Myoblast Isolation: Preparation of muscle, shaking, and grinding of tissue
🕐*Timing: 2.5 hrs.*

4. Wash isolated tissue 2-3 times with the initial PBS wash mixture.

NOTE: The PBS solution is prepared right before dissecting the tissue.

5. Incubate muscle tissue in the initial DMEM-F12 incubation mixture.

CRITICAL: Avoid filtering the DMEM-F12 media containing collagenase, 1% pen/strep, and 3 μL/mL Fungizone. This initial DMEM-F12 incubation mixture must be chilled (4 °C) when added to reduce temperature shock.

6. Maintain the muscle solution in a 37 °C water bath for 10-15 mins.
7. Shake at 220 rpm, for an overall time of 1.5 hrs.
8. After incubation, wash the tissue 3-4 times with PBS.
9. Incubate in warmed secondary DMEM-F12 incubation mixture while the tissue is shaken for 30 mins in a 37 °C water bath.

NOTE: Secondary DMEM-F12 incubation mixture has to be pre-warmed to 37°C to ensure efficient mixing of dispase and since the muscles were at 37°C after incubation.

10. After shaking, grind tissue with a mortar and pestle in the presence of liquid nitrogen.
11. Pass through a 100 μm, then 70 μm, cell strainer.
12. Centrifuge the solution at 1000 rpm for five mins to pellet the cells.

236 *Myoblast Isolation: Plating*
237 ⏱*Timing: 1-3 hrs.*
238
239   13. Transfer the to a plate and resuspended using DMEM-F12 growth media supplemented
240   with 40 ng/mL bFGF.
241   14. Pre-plate the cells for 1-3 hours on UNCOATED dishes to reduce the number of
242   fibroblasts.
243
244 **CRITICAL**: Fibroblasts can dilute satellite cells. Recommended for dystrophic or injured
245 muscle. Pre-plating on an uncoated plate causes fibroblasts to stick and be isolated. Fibroblasts
246 can separately be used to isolate and for other experiments.
247
15. Dilute cells 1:15 in PBS, then plate in a Matrigel-coated dish.

NOTE: To create Matrigel-coated dishes, dilute stock concentration (while keeping on ice) to
1:15 in sterile PBS in the hood. Put Matrigel solution on flask/plate, shake/tilt to coat the bottom,
incubate at room temperature in a chemical hood for 30 mins, and remove Matrigel solution back
into its original tube. Matrigel solution may be reused up to 5 times in total.

Myoblast Isolation: Differentiation
254
255 The below protocol offers subsequent differentiation to myotubes, if desired.
256
257 ⏱*Timing: 1-2 weeks.*
258
259   1. Wait for activation, which takes 24-48 hrs, after which myoblasts will grow rapidly.
260   a. To maintain healthy myoblast cells, use the growth media supplemented with
261   bFGF (10 ng/mL).
262
263 **CRITICAL**: Use Differentiation Medium to go from myoblasts to myocytes and then to
264 myotubes (Figure 2).
265   2. Plate primary myoblast at ~.8 X 10^6 cells per well and to differentiate the cells, add
266   differentiation media, supplemented with 1:10,000 bFGF.
267
268 **NOTE**: This will depend on # of cell passages and type of treatment, adjust accordingly.
269
270   3. Incubate for 4 to 7 days for differentiation to myotubes.
271
272 **NOTE**: Switch out with fresh differentiation media every 2 days, supplemented with 1:10,000
273 bFGF.
274
275   4. Cells are split using 2-5 mL of accutase for 5-15 minutes, dependent on cell count.
276
277 **Note: DO NOT** use trypsin to split the cells. Accutase is less harsh to the extracellular matrix,
278 surface proteins, and cytoskeleton of skeletal cells than trypsin, so it is highly preferred [21].
279 Accutate total incubation time and volume will differ depending on cell yield.

280
281   5. Cells are maintained in (5% CO_2) at 37°C. If growing myotubes, A confluency of 70-
282   85% has to be reached prior to adding growth media.
283
284   *Validation Day 1: Permeabilization and Blocking*
285   Immunofluorescence staining is effective for examining differences in skeletal muscles
286   simultaneously. Refer to Table 1 for a list of validated primary antibodies for skeletal muscles.
287   Select secondary antibodies that are compatible with the epifluorescence or confocal microscope
288   available to you.
289
290   ⏱*Timing: 1.5 hours.*
291
292   **CRITICAL**: All steps are performed at room temperature unless otherwise indicated.
293   **NOTE:** This protocol for Immunofluorescence staining and antibody validation of isolated
294   skeletal muscle cells is an adaptation of Esper et al., skeletal muscle tissue immunofluorescence
295   labeling protocol [22].
296
297   1. Fix cells by incubating them in 4% paraformaldehyde (PFA) for five mins.
298   2. Wash three times for five mins using phosphate-buffered saline (PBS).
299
300   **NOTE:** Ice-cold 100% methanol or acetone is an effective fixative for cryosections and more
301   suited for some antigens. Acetone is less harsh than methanol.
302
303   3. Incubate cells in permeabilization buffer for 10 mins.
304   4. Incubate cells in blocking solution for 1 hr at room temperature or overnight at 4 °C.
305
306   **NOTE:** When using permeabilization buffer, keep the solution away from the hydrophobic
307   barrier to avoid loss of hydrophobicity. If this happens, wash the slide well with PBS. Include
308   Mouse on Mouse (MOM) blocking reagent at a 1:40 dilution when staining mouse tissue with
309   antibodies raised in the mouse.
310
311   *Validation Day 2: Antibodies*
312   ⏱*Timing: 30 minutes.*
313
314
315   5. To begin immunostaining, dilute the primary and secondary antibodies in a blocking
316   solution according to the manufacturer's suggested ratio.
317
318   **NOTE**: It is acceptable to dilute antibodies in hybridoma supernatant when targeting multiple
319   antigens.
320
321   6. Aspirate the blocking buffer and cover the slide with the primary antibody solution.
322   7. Incubate the slides overnight at 4 °C.
323
324   *Validation Day 3: Preparing Slides for Imaging*
325   ⏱*Timing: 2-3 hours.*

326
327   8. On the following day, wash three times for five mins with PBS.
328   9. After washing, cover cells with secondary antibodies diluted in blocking buffer for 1 hr at
329   room temperature in the dark.
330
331 **NOTE**: Keep slides in the dark for the remainder of the protocol.
332
10. After incubation, wash the slides three times for five mins with PBS.
11. Incubate the cells with 1 μg/mL DAPI diluted in PBS for five mins.
12. Wash once with PBS for five min.
13. Aspirate the PBS and place 1–2 drops of mounting media onto the cells
14. Carefully place a coverslip on the slide, while avoiding air bubbles.
15. Let the slides dry in the dark for 1–2 hr before sealing the slides with clear nail polish.
16. Store the slides at 4 °C and image within 2 weeks.
340

Expected Outcomes:

Upon isolation of myoblast and differentiation into myotubes, we validated their structure in light microscopy (**Figure 3A**). Furthermore, we viewed multinucleated myotubes through TEM to validate the ultrastructure (**Figure 3B**). Transfection further showed myotubes demonstrated fluorescence as expected (**Figure 3C**). From there, we performed staining for myosin and desmin, muscle-specific proteins that play crucial roles in muscle cell structure and function [23], to confirm that filaments were present (**Figure 3 D-D''**).

Once myoblasts and myotubes are validated, they can be used for a variety of studies including to measure mitochondrial efficiency with oxygen consumption rate, western blot analysis to look for expression of specific proteins in knockout studies, or a variety of electron microscopy techniques such as serial block-face scanning electron microscopy to perform 3D reconstruction of organelles (**Figure 4**). In the past, following this isolation and differentiation protocol, we have successfully used the protocol by Garza-Lopez et al. (2022) [24] for 3D reconstruction and quantification of mitochondria and endoplasmic reticulum, alongside the protocol by Neikirk et al. (2023) [17] for 3D reconstruction and quantification of lipid droplets, lysosomes, and autophagosomes. However, from our experience, any validated scanning electron microscopy protocol should be effective following isolation and differentiation.

As an example, to validate this method, we sought to understand how insulin treatment (10 nM/L) in 2-hour increments may alter myoblast and myotube function through the usage of a Seahorse XF96 analyzer, per past protocols [25]. We found that for myoblasts, there is a significantly increased basal, maximum, and non-mitochondrial OCR after 2 hours of insulin treatment, while this difference is retained or exacerbated after 4 hours of insulin treatment (**Figure 5A-B**). After 6 hours of insulin treatment, OCR conversely showed significant decreases in all of these parameters (**Figure 5C**). In myotubes, after 2 and 4 hours of insulin treatment, we similarly noted a significant increase in mitochondrial OCR (**Figure 5D-E**). Notably, the increase in basal, ATP-linked, maximum, and non-mitochondrial OCR is much higher in 4 hours than 2 hours. Unlike myoblasts, 6 hours of insulin treatment myotubes did not differ significantly from untreated cells (**Figure 5F**). Importantly, there may be a differential response to insulin treatment in myoblasts and myotubes, highlighting the importance of studying both models. This validated that the function of myoblasts and myotubes are intact following this isolation.

From there we sought to elucidate if organelle proteins are affected following insulin treatment and we targeted Optic atrophy protein 1 (OPA-1), which is a mitochondrial inner membrane (IMM) fusion protein that mediates the fusion of the IMM between two mitochondria while also serving roles in mitochondrial bioenergetics and cristae architecture [26]. OPA-1 is just one of several proteins which modulate mitochondrial structure. For example, contrastingly, Dynamin-related protein 1 (DRP-1) is a protein that initiates the fission process through constriction of the mitochondria which divides the mitochondria into two separate organelles [27]. However, given that OCR increased following insulin treatment, it is possible this is due to the increased mitochondrial area caused by upregulated mitochondrial fusion. To see if OPA-1 may be changed in expression, we performed western blotting. When looking at OPA-1, we noticed a significant continuous increase in protein levels in myoblasts across 2 and 4 hours of insulin stimulation when normalized (**Figure 6A-B**). We further differentiated primary myotubes and carried out these experiments again to see if any differences existed (**Figure 6C-D**). We noticed significant increases in OPA-1 levels after 4 hours of insulin stimulation (**Figure 6C-D**). Together, this suggests that insulin stimulation causes increased expression of OPA-1 in a short time frame which is exacerbated in myotubes compared with myoblasts.

Since we saw changes in OPA-1 expression, which is known to trigger fusion, we also validated this technique of myoblasts and myotubes isolation through the quantification of mitochondria and cristae[1] following insulin treatment. Using transmission electron microscopy, we compared mitochondrial morphology without (**Figure 7A**) and with 2-hour insulin treatment in myoblasts (**Figure 7A'**). When quantified we saw that mitochondria reduce in number (**Figure 7B**), while becoming less spherical and larger in area (**Figure 7C-D**). Together, this suggests an uptick in fusion following insulin treatment in myoblasts. We also considered how cristae morphology may be affected following insulin treatment (**Figure 7E-E'**) and we saw that although the cristae score, a measurement of relative cristae quality, did not change, cristae number and area increased suggesting a greater capacity for oxidative function (**Figure 7F-H**). From there, we similarly sought to see if mitochondria in myotubes had changed following insulin stimulation (**Figure 8A-A'**). Similar to myoblasts, we saw while mitochondria decrease, their size increases (**Figure 8B-D**), again indicating increased mitochondrial fusion. When evaluating the cristae structure, we noticed that while similar to myoblasts cristae score was unchanged, the cristae number had a more significant increase (**Figure 8E-G**). Similar to myoblasts, the cristae area also increased following 2 hours of insulin stimulation (**Figure 8F).** Together these quantifications show that changes in OCR may be due to OPA-1-mediated changes in mitochondrial and cristae architecture following insulin stimulation.

Together these data validate this isolation and validation technique allows for the application of experimental models to elucidate cellular processes. This demonstrates the viability of the protocol outlined here for skeletal muscle. After differentiation, quantification can be done for many experimental designs. Here, we performed seahorse analysis per prior methods [25] with GraphPad to perform students' T-tests to measure statistical significance.

Past results have demonstrated that myoblasts can be assayed through staining with Pax7 and MyoD, while multinucleated myotubes are visualized with phase contrast microscopy or after staining for myosin heavy chain [7]. Past protocols are also available which allow for incredibly dense populations of myoblasts (1.5×10^7, 200 times cell expansion) to be obtained through an intelligent culture system with suppression of myotube formation [12]. Another technique has also allowed for $1 \times 10^7 - 2 \times 10^7$ myoblasts to be isolated from murine hindlimbs from a single organism without the need for cell straining or sorting [28]. Tangentially, techniques

allow for induced pluripotent stem cells to be differentiated into myotubes which allow for the
study of insulin-resistance [29]. Our technique is advantageous in offering the ability to obtain
either subpopulation, as well as fibroblasts, allowing for a wide range of experiments to perform.
Further, the study of specific genes in myoblasts and myotubes can then be affected through
verified techniques such as adenovirus, or herpes simplex virus type 1 amplicon vectors,
depending on the specific gene [30].
423
Quantification and Statistical Analysis:
For all analyses, GraphPad Prism software package was used (La Jolla, CA, USA), with black
bars representing the standard error of meanwhile dots represent individual data points shown. If
only two groups were used for comparison, an unpaired t-test was the statistical test, while more
than two groups were compared with a one-way ANOVA and Tukey *post hoc* tests for multiple
comparisons, or their non-parametric equivalent if applicable. A minimum threshold of $p < 0.05$
indicated a significant difference.
431
Limitations: This protocol has been optimized for mice gastrocnemius, quadriceps, and
hamstring muscles and may not be applicable to other model organisms or tissue types.
Compared with other protocols, ours takes a similar period of time [28], but this can still be a slow
process that must be carried out across multiple days. While C2C12 myoblasts are ideal for this
protocol, increasingly human skeletal myoblasts are important to study, and past protocols
indicate that differences in the procedure must be made, such as antisense miR-133a addition, to
promote the fast differentiation of human skeletal myoblasts [31].
439
Trouble Shooting:
Problem: Ultrastructure or Gross morphology of Myoblasts is Degraded
Potential Solution: This may be due to too much damage incurred to myoblasts during
preparation. We found that optimizing the process by first digesting tissue with type II
collagenase and dispase, followed by grinding the tissue in liquid nitrogen with a mortar with a
pestle, and passing it through cell strainers resulted in an improved procedure. However,
reducing the time grounded or reducing the amount of digestion can avoid potential damage to
the myoblasts if it is occurring.
Problem: Contamination with Fibroblasts
Potential Solution: It is important to plate first on an uncoated plate. However, if fibroblasts are
still observed, pre-plating can be done twice. Antibody-based selection of fibroblasts may cause
certain issues but can also be explored as an option to remove fibroblasts. If this remains an
issue, other methods have shown that using flowing cytometry can be used to identify and
remove fibroblasts [32].
Problem: Low Cell Yield or Viability
Potential Solution: If myoblast or myotube viability is low, it is important to increase the
concentration of growth factors and ensure a sterile environment is maintained. Reducing time
with accutase can also ensure cells are not treated too harshly.
458
459 **Resource Availability:**
460 *Lead contact*
461 Further information and requests for resources and reagents should be directed to and will be
462 fulfilled by the lead contact, Antentor Hinton (antentor.o.hinton.jr@Vanderbilt.Edu).

Materials availability

464 All generated materials, if applicable, are created in methods highlighted in the text above.

Data and code availability

468 Full data utilized and requests for data and code availability should be directed to and will be
469 fulfilled by the lead contact, Antentor Hinton (antentor.o.hinton.jr@Vanderbilt.Edu).

470
471
Author Contributions

473
Acknowledgments
475 All antibodies were obtained from the Iowa Developmental Studies Hybridoma Bank (DSHB).

Financial & Competing Interests' Disclosure
All authors have no competing interests.
This project was funded by the UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, BWF
Career Awards at the Scientific Interface Award, BWF Ad-hoc Award, NIH Small Research
Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program,
DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease
Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from
the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation
(to A.H.J.). NSF EES2112556, NSF EES1817282, and CZI Science Diversity Leadership grant
number 2022-253614 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon
Valley Community Foundation (to S.M. D.) and National Institutes of Health grant HD090061
and the Department of Veterans Affairs Office of Research award I01 BX005352 (to J.G.).
Additional support was provided by the Vanderbilt Institute for Clinical and Translational
Research program supported by the National Center for Research Resources, Grant UL1
RR024975–01, and the National Center for Advancing Translational Sciences, Grant 2 UL1
TR000445–06 and the Cell Imaging Shared Resource.

493
494 BioRender was used for the creation of Figures.

495
Data Sharing and Open Access
497 All data is available upon request to the corresponding author.

498
499
Figures and Legends
501

502
503　**Figure 1: The process of myoblast isolation from gastrocnemius muscle.**

504
505 **Figure 2: The process of myotube differentiation from myoblasts and utilization for serum**
506
507

508
509
510 **Figure 3:** Myotubes as characterized by A. Light microscopy, B. Transmission electron
511 microscopy, and C. Transfection with adenovirus containing the green fluorescent protein gene
512 (Ad-GFP). D. Straining with BA-D5-s, D' SC-71-s, and D'' D3-s to show myosin and desmin.
513

514
515 **Figure 4:** Examples of experiments that may be performed following myotube differentiation
516 and isolation.

517
Figure 5: Oxygen consumption rate (OCR) altered in myoblasts and myotubes upon altered
insulin stimulation which shows changes in mitochondrial efficiency. (A) Seahorse plot for
primary myoblasts following 2 hours of insulin stimulation (B) 4 hours of insulin stimulation and
(C) 6 hours of insulin stimulation. (D) Oxygen consumption rate was measured after several
inhibitors to measure respiration in primary myotubes after 2 hours, (E) 4 hours, and (F) 6 hours
of insulin stimulation. (A'-F') Basal OCR, which represents respiration under normal, unstressed
conditions. (A''-F'') ATP-linked OCR, which is respiration associated with ATP synthesis
during oxidative phosphorylation, which is marked by a reduction in OCR due to oligomycin.
(A'''-F''') Maximum OCR, which is the maximal capacity at which mitochondria may utilize
oxygen. (A''''-F'''') Non-mitochondrial respiration, which can be attributed to factors such as
glycolysis or ROS and not due to mitochondrial respiration. These values were compared to the
control (blue) in all of these examples. N = 6 per treatment, and * indicates p-value < .05.

530
Figure 6: Comparison of mitochondrial fusion proteins following insulin stimulation in primary
myoblasts and myotubes. (A) Western blotting for mitochondrial fusion protein OPA-1
following 2 hours, 4 hours, and 6 hours of insulin stimulation. (B) OPA-1 levels normalized to
Alpha tubulin following insulin stimulation. (C) This was replicated in primary myotubes, as
western blotting for mitochondrial fusion OPA1. (D) OPA-1 levels, normalized to Alpha tubulin,
in primary myotubes following insulin treatment. N = 6 per treatment, and * indicates p-value <
.05.

539
540
Figure 7. TEM Quantification of Myoblasts. (A) Representative transmission electron
micrographs from control and (A') insulin-treated cells, with red arrows showing fused
mitochondria. (B) Quantifications of number of mitochondria, (C) circularity of mitochondria,
(D) and area of mitochondria. (E) Representative transmission electron micrographs from control
and (E') insulin-treated cells, with red arrows showing cristae. (F) Quantifications from cristae
score, (G) cristae quantity, and (H) cristae area comparing non-insulin and insulin-treated
myoblasts. Dots show the number of samples. ** and **** indicates $p < 0.01$ and $p < 0.0001$,
respectively.
549

Figure 8. TEM Quantification of Myotubes. (A) Representative transmission electron micrographs from control and (A') insulin-treated cells, with red arrows showing fused mitochondria. (B) Quantifications of number of mitochondria, (C) circularity of mitochondria, (D) and area of mitochondria. (E) Representative transmission electron micrographs from control and (E') insulin-treated cells, with red arrows showing cristae. (F) Quantifications from cristae score, (G) cristae quantity, and (H) cristae area comparing non-insulin and insulin-treated myotubes. Dots show the number of samples. ** and **** indicates p < 0.01 and p < 0.0001, respectively.

Antibody Name	Antigen	IF/IHC Rec		Isotype	Antigen Species	Host Species	Positive Tested Species Reactivity	Info
BF-F3-s	Myosin heavy chain Type IIB	2-5 ug/ml	Monoclonal	MIgM	Bovine	mouse	Bovine, Mouse, Porcine, Rat, Sheep	
SC-71-s	Myosin heavy chain Type IIA	2-5 ug/ml	Monoclonal	MIgG1	Bovine	mouse	Bovine, Canine, Goat, Horse, Human, Mole, Mouse, Porcine, Rabbit, Rat, elephant seal, guinea pig, llama and goat muscle. Sheep	This antibody recognizes the fast twitch isoform MyHC IIA specifically in horses and rodents.
BA-D5-s	Myosin heavy chain Type I	2-5 ug/ml	Monoclonal	MIgG2b	Bovine	mouse	Bovine, Canine, Fish, Goat, Guinea Pig, Horse, Human, Lamb, llama, Mouse, Porcine, Rabbit, Rat, Zebrafish	Muscle contraction
F5D-s	Myogenin	2-5 ug/ml	Monoclonal	MIgG1, kappa light chain	Rat	mouse	canine, Feline, Human, Mammal, Mouse, Porcine, Rat	This antibody immunoprecipitates deletion mutants containing a.a. 138-158, region immediately carboxy-terminal to the bHLH domain.
PAX7-s	Pax7	2-5 ug/ml	Monoclonal	MIgG1, kappa light chain	Chicken	mouse	Amphibian, Avian, Bovine, Canine, Fish, Goat, Human, Mouse, Ovine, Porcine, Quail, Rat, Turtle, Xenopus, Zebrafish	nuclei in adult skeletal muscle satellite cells
D3-s	Desmin, intermediate filament	2-5 ug/ml	Monoclonal	MIgA, kappa light chain	Chicken	mouse	Chicken, Hamster, Mouse, Rat	D3 also stained primary cultures of embryonic cardiac myocytes
9.1 ITGA7-s	Integrin alpha-7, extracellular domain	2-5 ug/ml	Monoclonal	MIgG2c	Human	mouse	Human	Alpha 7/beta-1 integrin is the primary laminin-1 receptor. This receptor is expressed in skeletal and cardiac muscle and certain tumor cells. The 9.1 antibody recognizes the extracellular domain of alpha 7 integrin in both native and denatured conformations.
JLT12-s	troponin T, fast skeletal muscle specific	2-5 ug/ml	Monoclonal	MIgG1	Rabbit	mouse	Axolotl, Bovine, Broad species, Chicken, Human, Rabbit, Rat	Paraformaldehyde fixation for immunostaining is recommended. JLT12 recognizes all Troponin T, fast skeletal muscle isoforms and doesn't recognize slow isoforms. This antibody was initially characterized in Lin et al. [Lin, J.J. C., Feramisco, J.R., Blose, S.H., and Matsumura, F. (1984). Monoclonal antibodies to cytoskeletal proteins. In Monoclonal Antibodies and Hybridomas: Progress and Applications. (eds. R.H. Kennett, T.J. McKearn, and K.B. Bechtol) pp. 119-151, Plenum Press, New York].
MANEX46B(7G1)-s	dystrophin	2-5 ug/ml	Monoclonal	MIgG1	human	mouse	Human, Mouse	Stains muscle membranes (no staining of Duchenne muscle membrane) Frozen, unfixed sections. May not work on formalin fixed tissue. Dilution: 1/4

Table 1: A list of antibodies and their respective antigens for skeletal muscle validation after isolation. The Table was obtained from https://dshb.biology.uiowa.edu/.

References

Automatic citation updates are disabled. To see the bibliography, click Refresh in the Zotero tab.