Title: Ablation of *Sam50* is associated with fragmentation and alterations in metabolism in murine and human myotubes

Authors and Affiliations: Bryanna Shao[1*], Mason Killion[1*], Ashton Oliver[1*], Chia Vang[1*], Faben Zeleke[1], Kit Neikirk[1], Zer Vue[1], Edgar Garza-Lopez[2], Jian-qiang Shao[3], Margaret Mungai[2], Jacob Lam[2], Qiana Williams[1], Christopher T. Altamura[1], Aaron Whiteside[1,4], Kinuthia Kabugi[1], Jessica McKenzie[1], Alice Koh[1], Estevão Scudese[1], Larry Vang[1], Andrea G. Marshall[1], Amber Crabtree[1], Janelle I. Tanghal[5], Dominique Stephens[1], Ho-Jin Koh[6], Brenita C. Jenkins[7,8], Sandra A. Murray[9], Anthonya T. Cooper[9], Clintoria Williams[4], Steven M. Damo[10], Melanie R. McReynolds[7,8], Jennifer A. Gaddy[11,12], Celestine N. Wanjalla[11], Heather K. Beasley[1#], Antentor Hinton, Jr[1#]

Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA

Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA

Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA.

Department of Neuroscience, Cell Biology and Physiology, Wright State University, Dayton, OH 45435 USA

Morehouse School of Medicine, Atlanta, GA 30310-1495

Department of Biological Sciences, Tennessee State University, Nashville, TN 37209

Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA

Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA

Department of Cell Biology, School of Medicine, University of Pittsburgh, Pittsburgh, PA, 15260, USA

Department of Life and Physical Sciences, Fisk University, Nashville, TN, 37208, USA

Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, 37232, USA

Tennessee Valley Healthcare Systems, U.S. Department of Veterans Affairs, Nashville, TN, 37212, USA

*-Indicates Co-First Authorship
#-Indicates Co-Senior Authorship
Corresponding Authors:
Antentor O. Hinton, Jr, PhD
Vanderbilt School of Medicine Basic Sciences
antentor.o.hinton.jr@Vanderbilt.edu

Keywords: 3D reconstruction, mitochondria, SAM complex, Sam50, MICOS

Abstract: The Sorting and Assembly Machinery (SAM) Complex is responsible for assembling β-barrel proteins in the mitochondrial membrane. Comprising three subunits, Sam35, Sam37, and Sam50, the SAM complex connects the inner and outer mitochondrial membranes by interacting with the mitochondrial contact site and cristae organizing system (MICOS) complex. Sam50, in particular, stabilizes the mitochondrial intermembrane space bridging (MIB) complex, which is crucial for protein transport, respiratory chain complex assembly, and regulation of cristae integrity. While the role of Sam50 in mitochondrial structure and metabolism in skeletal muscle remains unclear, this study aims to investigate its impact. Serial block-face-scanning electron microscopy (SBF-SEM) and computer-assisted 3D renderings were employed to compare mitochondrial structure and networking in *Sam50*-deficient myotubes from mice and humans with wild-type (WT) myotubes. Furthermore, autophagosome 3D structure was assessed in human myotubes. Mitochondrial metabolic phenotypes were assessed using Gas Chromatography-Mass Spectrometry-based metabolomics to explore differential changes in WT and *Sam50*-deficient myotubes. The results revealed increased mitochondrial fragmentation and autophagosome formation in *Sam50*-deficient myotubes compared to controls. Metabolomic analysis indicated elevated metabolism of propanoate and several amino acids, including ß-Alanine, phenylalanine, and tyrosine, along with increased amino acid and fatty acid metabolism in *Sam50*-deficient myotubes. Furthermore, impairment of oxidative capacity was observed upon *Sam50* ablation in both murine and human myotubes, as measured with the XF24 Seahorse Analyzer. Collectively, these findings support the critical role of Sam50 in establishing and maintaining mitochondrial integrity, cristae structure, and mitochondrial metabolism. By elucidating the impact of *Sam50*-deficiency, this study enhances our understanding of mitochondrial function in skeletal muscle.

Introduction:

Mitochondria are generally associated with their bioenergetic roles in the cell, as they are the hubs of oxidative phosphorylation (1,2). Aside from energy production, mitochondria are also integral in other processes such as maintaining calcium homeostasis and apoptosis (2,3). Being such a functionally important organelle, mitochondria are also implicated in the pathophysiology of numerous diseases when they become dysfunctional. Understanding the processes involved in mitochondrial dysfunction may provide novel therapeutic treatment options for many common diseases. For example, Diabetes mellitus is considered a mitochondrial disease (4) and increasing the understanding of mitochondrial function and morphology in normal and pathophysiological functions may be key in developing novel protocols to treat and even prevent disease. This is exemplified when considering reactive oxidative species (ROS) signaling which may alter various mitochondrial functions including ATP production, which may be linked to the development of metabolic diseases and structural changes (5). Mutations or changes in the function of the mitochondrial DNA (mtDNA) can result in altered mitochondrial function; this can have consequences on insulin production, lead to hypertension, and ultimately be related to the progression of Diabetes Mellitus (6,7). Thus, the

prospect of understanding how changes in mitochondrial structure may affect metabolism and
remains pertinent for discovering new therapeutic treatments for diseases with high morbidity
and mortality (8).
Critical to the functioning of mitochondria are the structural folds of the inner membrane
known as cristae (9). Cristae provide a means for mitochondria to maximize the amount of
oxidative phosphorylation machinery that can be present relative to the internal volume (9). A
key determining factor in mitochondrial ultrastructure is mitochondrial dynamics which consist
of continuous cycles of fusion and fission. Mitochondrial dynamics give rise to various
mitochondrial phenotypes which range from the typical spherical to stress-states including
megamitochondria and donut-shaped (10,11). GTPases facilitate the dynamic events with
Mitofusin-1 (Mfn1) and Mitofusin-2 (Mfn2) fusing the outer membrane and Optic atrophy 1
(Opa1) fusing the inner membrane (9,12,13). Opa1 has been shown to be linked to mitochondrial
fragmentation while also affecting cristae dimensions, shapes, and size (14). Specifically, Opa1
exists in two principal isoforms, the short and long forms, which believed to be responsible for
prompting fission and the fusion of the inner membrane, respectively (15). The maintenance of
the proper long vs short isoforms are crucial for mitochondrial health and proper cell signaling
(16). Thus, understanding the mechanisms of how modulators of cristae affect overall
mitochondrial structure remains important and is still not completely understood.
Mitochondria can be separated into several distinct compartments. The inner
mitochondrial membrane that invaginates to form crista and at the cristae junctions, the
mitochondrial contact site and cristae organizing system (MICOS) complex resides (14,17,18).
The MICOS complex bonds to the sorting and assembly machinery (SAM) complex (17). The
MICOS complex is currently known to be made up of Mic10, Mic13, Mic19, Mic25, Mic26,
Mic27, and Mic60 (17). Together, the interaction between the MICOS and SAM complexes
connect the inner and outer mitochondrial membranes to form the mitochondrial intermembrane
space bridging (MIB) complex (17). These proteins and their interactions are important for
organizing and structuring the mitochondria to allow for maximum and efficient respiration and
ATP generation (19). The MICOS complex, SAM complex, and OPA1 can all affect the
dynamics of the cristae in varying ways (14,17,20,21). OPA1 is vital in promoting the formation
of cristae and as it is responsible for determining the width of the cristae (14). Additionally, the
MICOS subunits interact with the SAM complex to assemble β-barrel proteins and maintain
contact between the outer and inner membranes (14,21,22). These all are necessary to maintain
the overall shape and distances in mitochondrial networks. Overall, cristae membranes are
crucial as they host the electron transport chain machinery and the ATP synthase dimers, which
are necessary for the production of ATP through oxidative phosphorylation (9,20). However,
given the critical relationship observed between mitochondrial morphology and function,
determining the effects of SAM-associated gene knockout is necessary to understand the
functional implications concomitant with changes in the mitochondrial ultrastructure and
morphology.

While the functions of the SAM complex may easily be considered as a subset of the
MIB complex, the SAM complex remains an interesting target distinct from the MICOS
complex. The SAM complex is made up of three subunits, Sam35, Sam37, and Sam50 (23).
While both Sam35 and Sam37 are peripheral membrane proteins that are not required for
survival, the central component, Sam50, a 7–8 nm diameter β-barrel channel, interacts with the
MICOS complex to modulate protein transport and mitochondrial morphology (17,21,22,24).
Specifically, Sam50 stabilizes the MIB complex for protein transport, respiratory chain complex
assembly, and cristae integrity regulation (25). In this way, it is understood that to structurally
form and sustain the cristae, proteins of the MICOS complex assemble at the cristae junction and
bind directly to Sam50 (19,21,26). Within the entire MIB complex, Sam50 is considered to be
among the most important proteins, alongside members of the MICOS complex Mitofilin and
Chchd3, for proper mitochondria structure and function (27). The importance of Sam50 is
relevant, as loss of *Sam50* can result in diminished cristae count, reduced cristae dynamic events,
abnormal cristae, and a lack of cristae junctions, the sites at which cristae are tethered to the
mitochondrial inner boundary membrane (14). However, the implications of the loss of *Sam50*
on mitochondrial dynamics in 3D is unknown.
2D microscopy techniques allow for the viewing of typical tubular mitochondria (28,29),
however, 3D phenotypes of mitochondria differ between tissue types and disease states
(10,30,31). Mitochondrial swelling can occur as a response to dysregulation of ion homeostasis
(32), while toroid mitochondria can arise in a calcium-dependent manner in stress states (33).
Through fusion and fission dynamics, it is understood that mitochondria can respond to cellular
stress to balance factors including volume, which typically allows for greater ATP generation,
and surface area, which typically allows for organelle-to-organelle contacts which are crucial for
extraneous processes such as calcium homeostasis (10,34,35). Therefore, 3D microscopy
techniques (34,36–38) are required for an understanding of many of the diverse phenotypes
which mitochondria might display.
This study specifically examined the metabolomics of the knockdown of *Sam50* in
murine and human myotubes. For the knockout of genes, myotubes, derived from satellite cell
fusion, have emerged as robust cellular models which mimic the properties found in skeletal
muscle (39). Importantly, human and murine myotubes remain distinct, with separate phenotypes
and metabolic progression (40). Therefore, human myotubes were *ex vivo* isolated from skeletal
muscle biopsies and differentiated in multinucleated myotubes in culture (41). To measure
structure, we used a mixed-method approach utilizing principally serial block-face-scanning
electron microscopy to allow for manual tracings to be performed on z-direction slices, known as
orthos, which were then aligned in analysis software Amira (34), and reconstructed into 3D
structures. Serial block-face-scanning electron microscopy (SBF-SEM) importantly allows for
the ability to see mitochondrial complexity, as it offers high x- and y- resolution and range. We
also sought to compare differences in mitochondrial structure and function based on human
myotubes to better elucidate if *Sam50* may have a differential role in these models. Beyond
structural dynamics, we sought to combine how changes in mitochondrial morphology upon

Sam50-deficinecy alters mitochondrial function and general biochemical pathways, which may
offer further implications for therapeutical targets involved in *Sam50*.
161
162 **Materials and Methods:**
163
Mice Care:
All procedures for the care of mice were in accordance with humane and ethical protocols
approved by the University of Iowa Animal Care and Use Committee (IACUC) which follows
the National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals, as
described previously (42). Experiments utilized WT male C57Bl/6J mice housed at 22 °C on a
12-hour light/12-hour dark cycle with free access to water and standard chow. Mice were
anesthetized with 5% isoflurane/95% oxygen.
171
Murine and Human Myotube Isolation:
Satellite cell isolation was performed as previously described (42). Satellite cells from C57Bl/6J
mice were plated on BD Matrigel-coated dishes and activated to differentiate into myoblasts and
myotube. Specifically, for myoblast differentiation, mouse-derived satellite cells with
Dulbecco's modified Eagle medium (DMEM)-F12 containing 20% fetal bovine serum (FBS),
40 ng/ml basic fibroblast growth factor, 1× non-essential amino acids, 0.14 mM β-
mercaptoethanol, 1× penicillin/streptomycin, and Fungizone. then maintained with 10 ng/ml
basic fibroblast growth factor. When myoblasts were 90% confluent, they were differentiated to
myotubes in DMEM-F12 containing 2% FBS and 1× insulin–transferrin–selenium. Three days
after differentiation, myotubes were infected with 1 µg CRISPR/Cas9 (Santa Cruz Sam50
CRISPR Plasmid; sc-427129) to achieve *Sam50* deletion, which was validated by qPCR.
Experiments were performed 3-7 days after infection.
184
RNA Extraction and RT-qPCR
The RNA was isolated with an RNeasy kit (Qiagen Inc) and quantified by measuring absorbance
at 260 nm and 280 nm with a NanoDrop 1000 spectrophotometer (NanoDrop products,
Wilmington, DE, USA). Isolated RNA (~1µg) was reverse transcribed with a High Capacity
cDNA Reverse Transcription Kit (Applied Biosciences, Carlsbad CA) and amplified by real-time
quantitative PCR (qPCR) with SYBR Green (Life Technologies, Carlsbad, CA) (43). Three for
each qPCR, t samples (~50 ng DNA each) were placed in a 384-well plate and underwent
thermal cycling in an ABI Prism 7900HT instrument (Applied Biosystems). Thermal cycling
conditions were set as follows: 1 cycle at 95°C for 10 min; 40 cycles of 95°C for 15 s, 59°C for
15 s, 72°C for 30 s, and 78°C for 10 s; 1 cycle of 95°C for 15 s; 1 cycle of 60°C for 15 s; and one
cycle of 95°C for 15 s. Results were normalized to glyceraldehyde-3-phosphate dehydrogenase
(*GAPDH*) and presented as relative fold changes.

Measurement of OCR Using Seahorse

Oxygen consumption rate was measured for *Sam50* KD fibroblasts using an XF24 bioanalyzer (Seahorse Bioscience: North Billerica, MA, USA) as previously described (42,44). Briefly, cells were plated at a density of 20×10^3 per well and differentiated for 3 days. For *Sam50* KO models, three days after differentiation, myotubes were transfected with CRISPR/Cas9 to achieve KO as described above. Cells were incubated in XF☐DMEM (supplemented with 1 g/L D-Glucose, 0.11 g/L sodium pyruvate, and 4 mM L-Glutamine), without CO_2 for 60 minutes. Cells were subsequently treated with mitochondrial stress modulators oligomycin (1 μg/ml), carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP; 1 μM), rotenone (1 μM), and antimycin A (10 μM), in that order, while remaining in the XF-DMEM media and assessed using an XF24 Seahorse Analyzer. After analysis, cells were lysed in 20μl lysis buffer (10mM Tris, 0.1% TX-100, pH 7.4) (44),and 480 μl of Bradford reagent. was added to each well as described previously (40). Total protein concentration was measured at an absorbance at 595 nm and used for normalization. For each sample, three independent experiments were performed for each condition with representative data from the replicates being shown.

Segmentation and Quantification of 3D SBF-SEM Images Using Amira

SBF-SEM ortho slices were 3D reconstructed using contour tracing (manual segmentation) in Amira to perform 3D reconstruction, as described previously (34,45). 300–400 slices ortho slices were transferred to Amira, and 50-100 serial sections were chosen, stacked, aligned, and visualized. A blinded individual familiar with organelle morphology traced structural features manually on sequential slices of micrograph blocks. Dots represent the total number of mitochondria surveyed, which varies.

Gas Chromatography-Mass Spectrometry (GC-MS) and Analyzing Metabolomic Data

Samples were extracted for metabolites and prepared as previously designed (46) and performed at University of Iowa High Resolution Mass Spectrometry Facility according to standard procedures. TraceFinder 4.1 was used preloaded with standard verified peaks and retention times to compare metabolite peaks in each sample against an in-house library of standards. to correct for drift over time by using QC samples, we used previously described protocols (47). All data were normalized to an internal standard to control for extraction, derivatization, and/or loading effects. Metabolomic analysis was performed as described previously (46) using the web service MetaboAnalyst 5.0 (https://www.metaboanalyst.ca/MetaboAnalyst/ModuleView.xhtml, last accessed on 1 September 2022). All tests performed were using built-in comparison, including one-way ANOVA and Fisher's LSD multiple comparison test. The fold enrichment number was calculated by the observed hits by the expected hits, which were determined by MetaboAnalyst 5.0.

Data Analysis

In graphs, black bars represent the standard error, and dots represent individual data points. An unpaired t-test was used for data with only two groups while ANOVA with Tukeys' *post-hoc* analysis was used for data with two or more groups. For both types of analyses, GraphPad Prism software package was used (La Jolla, CA, USA). A minimum threshold of $p < 0.05$ indicated a significant difference. Higher degrees of statistical significance (**, ***, ****) were defined as $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively.

Results:

<u>Loss of Sam50 causes Loss of Mitochondrial Size and Alterations in Morphology in Murine and Human Myotubes</u>

Given that mitochondrial size is linked with respiratory efficiency (31), we first sought to determine if mitochondrial size changed as a result of *Sam50* ablation. Following CRISPR/Cas9 knockout of Sam50 (Figure 1A), SBF-SEM micrographs were obtained (Figure 1B), and we performed manual contour segmentation for 3D reconstruction (Figure 1C-D) and subsequent quantification (Figure 1E). For each experimental condition, 10 myotubes were used with sectioning performed on 10 µm by 10 µm slices with a z-directional depth of 300 µm (Figure 1B). Fifty ortho slices (Figure 2A-B) were surveyed and manual contour tracing was performed on each mitochondrion to perform 3D reconstructions (Figure 2A'-B'). This allowed for observation of the complete mitochondrial 3D structure (Figure 2A''-B'') in murine myotubes as well as human myotubes (Figure 2A'''-B''''') (48). In each experimental condition, 3 regions of interests were considered, within which approximately 250 mitochondria were quantified for a total of 750 mitochondria (Figure S1).

We observed that mitochondrial fragmentation was increased in *Sam50*-deficient murine myotubes compared to control myotubes (Figure 2C). Similarly, the area in 3D was also decreased upon *Sam50* loss (Figure 2D). This mimicked a loss in perimeter (Figure 2E), reducing the total mitochondrion surface area. In considering *Sam50*-deficient human myotubes (Figure 2C'-E'), we noticed a similar phenotype with loss of mitochondrial volume, area, and perimeter when compared to human myotubes (Figure 2C'-E'). From there, we sought to understand how beyond only size, mitochondria may differentially change in shape in murine and human myotubes following loss of *Sam50*.

To begin with, we showed mitochondria, from the transverse plane, in control (Figure 3A-A') and *Sam50*-deficient murine myotubes (Figure 3B-B'), as well as human myotubes (Figure 3A''-B'''). Viewing mitochondria from both of these planes showed that at baseline murine myotubes are much more complex than human myotubes. While both murine and human myotubes showed loss in mitochondrial structure following loss of *Sam50*, this was a more significant loss in murine myotubes. Next, we sought to determine whether mitochondrial sphericity was altered. *Sam50*-deficiency was associated with an increase in mitochondrial

sphericity which is indicative of reduced complexity in murine (Figure 3C) and human myotubes
(Figure 3C'). To further understand mitochondrial complexity, we measured the mitochondrial
complexity index which is a measure of the surface area-to-volume ratio. Mitochondria become
significantly less complex upon loss of *Sam50* (35) in murine (Figure 3D) and human myotubes
(Figure 3D'). This was further validated with mito-otyping, an approach that involves organizing
mitochondria by volume in a similar manner to the arrangement of chromosomes during
karyotyping. With mito-otyping, we observed that while control mitochondria were highly
elongated *Sam50*-deficient mitochondria generally become less elongated, resulting in a decrease
in overall volume in both murine (Figure 3F) and human (Figure 3F') myotubes. Overall, loss of
Sam50 has been shown to be associated with a dysregulation of mitochondrial structure.
Considering the dramatic decrease in mitochondrial size observed in *Sam50*-deficient myotubes,
we sought to also investigate whether these morphological effects influenced the efficiency of
mitochondrial functions.
288
289   Increased Autophagosomes Occurs Upon Loss of *Sam50* in Human Myotubes
290   Most abundantly, macroautophagy is the routine cellular mechanism for organelles,
including mitochondria, through principally autophagosomes (49). Targeting of mitochondria in
autophagy, known as mitophagy, can serve as a critical cellular response to reactive oxygen
species or other forms of dysfunctional mitochondria; as such, mitophagy has been proposed as a
key target in limiting the pathological progression of neurodegenerative diseases (49). *Sam50*
may play a crucial role in the biochemical triggering of these mitophagy events. Of note, *Sam50*
interacts with autophagy receptor p62 (50). Beyond this, past research has also found that *Sam50*
deficiency results in autophagic flux, with mtDNA protecting from complete mitochondrial
degradation (51). However, autophagy remains a dynamic process that may be more effectively
studied using 3D reconstruction (52), and the effects of *Sam50*-deficiency on the structure of
autophagic organelle machinery are poorly elucidated.
301   Since humans and murine myotubes reflected mitochondrial structural arrangements with
loss of *Sam50*, we focused on human myotubes. Here, again using 3D reconstruction from 10
µm by 10 µm slices obtained from 10 human myotubes, with a z-directional depth of 300 µm,
we surveyed 50 ortho slices (Figure 4A). Following manual segmentation of these 50 ortho
slices, 3D renderings of autophagosomes were able to be created for control and *Sam50*-deficient
human myotubes (Figure 4B-B'). We found that autophagosomes, when normalized to the same
area regions of interests (ROI), were much higher in count upon loss of *Sam50* in human
myotubes (Figure 4C). Beyond only count, the average volume of each autophagosome increases
(Figure 4D). These increases in overall size can be illustrated when autophagosomes are
organized on the basis of their volume (Figure 4E). This suggests larger autophagosomes that
may be more active or have increased cargo loads. Together, these results show that
autophagosome formation was increased in *Sam50*-deficient myotubes compared to control
myotubes, suggesting an uptick in mitophagy.
314

<u>Alterations of Mitochondrial Efficiency Occurs in *Sam50*-deficient Murine and Human</u>
<u>Myotubes</u>
To investigate whether observed structural changes mitochondria influenced
mitochondrial efficiency, we assessed the overall oxidative capacity of control and *Sam50*-
deficient human and murine myotubes. For these experiments, we used an XF24 Seahorse
Analyzer which measures oxygen consumption rate (OCR), an indicator of mitochondrial
oxidative photophosphorylation, across various drug applications. Prior to mitochondrial
efficiency studies, knockout of Sam50 was confirmed in both murine and human myotubes
(Figure 5A-4A'). Overall, *Sam50*-deficient murine and human myotubes showed reduced OCR
at all time intervals (Figure 4B-B'). To determine basal OCR, cells were assessed under normal
oxidative conditions. Basal OCR was impaired in cells lacking Sam50, compared to control cells
(Figure 5C-C'). A series of oxidation modulators were applied to further investigate the effects
of *Sam50* on oxidative processes. First, oligomycin was applied to inhibit ATPase and induce
hyperpolarization of the membrane, which allowed us to examine the role of *Sam50* in ATP-
linked respiration (Figure 5D-D'). Carbonyl cyanide-4 (trifluoromethoxy) phenylhydrazone
(FCCP), a mitochondrial oxidative phosphorylation uncoupling agent, was then applied. With
FCCP treatment, we show that under conditions of maximal respiration, OCR is reduced with the
loss of *Sam50* (Figure 5E-E'). compared to control Finally, myotubes were treated antimycin-a
and rotenone inhibitors electron transport, which allowed for the glycolytic reserve to be
determined as a function between maximum OCR and baseline OCR (Figure 5F-F'). Overall,
with our mitochondrial functional assay, we demonstrate that oxidative capacity is impaired
upon ablation of *Sam50* in murine and human myotubes.
337
<u>*Sam50*-deficiency in Human Myotubes Changes Metabolomic Pathways</u>
Since human and murine myotubes had similar structural changes with the loss of *Sam50*,
we specifically looked at human myotubes to investigate the specific biochemical changes
resulting from the loss of *Sam50*. To investigate the metabolic effects of *Sam50* loss, we
employed Gas Chromatography-Mass Spectrometry-based (GC-MS) metabolomics. Analysis
using a volcano plot revealed significant downregulation of ß-alanine, gamma aminobutyric acid
(GABA), and hypotaurine upon *Sam50* loss (Figure 6A). Heat map analysis further showed
downregulation of various metabolites, including amino acids such as glutamine, phenylalanine,
glutamate, threonine, and proline, suggesting a decrease in amino acid products due to increased
amino acid metabolism. On the other hand, ribose, ß-Hydroxy, indolepropionic, inotisol, and
dihydrophenylalanine were upregulated (Figure 6B). Enrichment analysis confirmed these
findings, with significant enrichment observed in the metabolism and degradation of amino acids
and fatty acids. Additionally, the metabolomic analysis revealed an increase in amino acid
metabolism and fatty acid metabolism (Figure 6C). Collectively, these findings suggest that
altered biochemical pathways may arise in response to, or as a mechanism of, the disrupted 3D
structure of mitochondria upon *Sam50* loss.
354

Discussion:

       Together, these data suggest Sam50 is critical for the establishment and maintenance of mitochondria, mitochondrial cristae structure, and mitochondrial metabolism. To our knowledge, we are the first group to look at the 3D structure of mitochondria in murine and human myotubes following the genetic knockdown of *Sam50*. Previously, transmission electron microscopy has been utilized to observe that Sam50 can be used as a therapeutic target, with overexpression rescuing mitochondrial morphology upon injury-induced loss of morphology (53). Interestingly, our results show that *Sam50* may be functionally required for many of the diverse phenotypes previously observed in other tissue types (10). Loss of *Sam50*, importantly, showed a loss of mitochondrial morphology diversity, which may have functional impacts on functions including calcium homeostasis and mitochondria-organelle contacts.

       Previously, studies have shown that Sam50 levels may rise in response to mitochondrial dysfunction, suggesting it acts as a sort of recovery molecule (53). This highlights the importance of understanding how Sam50 interacts with the MICOS complex to influence overall mitochondrial structure. While it is well understood that the MICOS structure modulates cristae changes, here we also show a structure loss in mitochondria structure upon loss of the SAM complex. It is possible this occurs due to the interaction of the MICOS complex, which may also affect overall mitochondrial dynamics, potentially through the modulation of fusion and fission dynamics such as DRP1 and OPA1, which also affects cristae structure (13). Future experiments may look at cristae to understand how cristae structural rearrangement alters 3D mitochondrial general phenotype in the loss of the MIB complex. Therefore, future experiments should consider further utilizing focused ion beam-scanning electron microscopy, which allows for machine-learning outputs of 3D cristae morphology to be constructed (34,36,54).

       In this study, we use 3D microscopy to dissect changes in mitochondrial morphology and function in myotubes with altered *Sam50* expression. From images captured with this technique, we demonstrated mitochondrial volume concomitantly occurs with a loss of complexity in both murine and human myotubes. By studying both human and murine myotubes we were able to *show* how murine myotubes mitochondria are relatively larger than human myotubes, but loss of Sam50 cause mitochondria to undergo significant reductions in size and morphology in both models. It has been suggested microscopic imaging can be targeted for the development of treatments to correct cellular defects (53). Specifically, overexpression of Sam50 was shown to rescue ischemia/reperfusion injury-induced changes by Yin and colleagues in a neural model (53), highlighting Sam50 may be an important therapeutic in a mitochondrial-dependent manner.

       Importantly, our results show the importance of understanding the functional impact of mitochondria decreasing in elongation, as past mechanistic studies remain conflicting. Previous studies have shown that mitochondrial elongation may be indicative of reduced fission dynamics, such as due to the loss of Drp1 (55). Mitochondria dysfunction may also arise as a result of mtDNA mutations (7). Interestingly, in a Drp1-dependent manner, mtDNA mutations may cause mitochondrial morphological alterations in Parkinson's disease; however, knockdown of Drp1 can rescue mitochondria through elongation (55). Elongated mitochondria can also serve a

distinct function to distribute membrane potential across a wider area (10). Past studies have
shown that loss of the mitochondrial membrane potential alters mitochondria volume (56).
Membrane potential is also regulated in cristae dynamics (57), suggesting alteration in the MIB
complex may affect membrane potential. However, past research has shown that *Sam50*
downregulation does not drastically alter membrane potential (25). Beyond this, it has also been
suggested that elongated mitochondria offer increased ATP production and protect against
mitophagy, which may aid in explaining the uptick in autophagy observed following *Sam50* KO
(58).
Outside of elongated mitochondria, a key mitochondrial characteristic observed
exclusively in wild-type myotubes is a diversity of phenotypes. Interestingly, there is greater
complexity in murine myotubes but reduced elongation when compared to murine myotubes. For
both myotubes, following *Sam50* KO, many phenotypes have much higher complexity (Figure
3D), as well as higher surface area. This is epitomized by toroid-shaped mitochondria, which
offer mitochondria an increased surface area, at the expense of volume for cristae (10). Although
calcium-dependent donut mitochondria have been observed in states of ROS stress and disease
(10,33), they may also serve beneficial tissue-dependent roles (59). Their absence in *Sam50-*
deficient models may signify a potential inhibition of calcium. Beyond this, their increased
surface area may allow wild-type myotubes to have increased contact sites with the endoplasmic
reticulum, lipid droplets, and other organelles. This remains relevant as mitochondria
endoplasmic reticulum sites (MERCs) serve a multitude of functions including calcium
homeostasis (60,61) and regulating autophagy as autophagosomes form at ER sites (52,62).
Future studies may consider further exploring this link by understanding how mitochondrial
contact sites may change upon loss of *Sam50*.
Also offering valuable clues about what may be causing mitochondria function to be
affected is metabolic changes we observed using GC-MS. Many of the results from the
metabolic analysis show that loss of *Sam50* may result in metabolic changes that parallel disease
states. For example, elevations in levels of 3,4-dihydroxyphenylalanine, which we observed
increased in *Sam50*-deficient myotubes (Figure 5B), are understood to occur in association with
oxidative stress during disease states (63,64). Past results have also shown that *Sam50*-deficiency
is associated with liver disease through increased lipid accumulation due to increased fatty acid
oxidation (65). We noted a similar effect as beta-oxidation of fatty acids increasingly occurred in
Sam50-deficiency. In the human heart, energetic demands are often carried out through the beta-
oxidation of fatty acids (66). Therefore, these results suggest that it is possible that fatty acid
oxidation is increased upon loss of *Sam50* to allow for energy production in states when
mitochondria lose optimal functioning; however, future studies may further explore if in
mitochondrial disease states, fatty acid oxidation changes in a *Sam50*-dependent manner.
Of note, we saw largely significant reductions in levels of GABA, hypotaurine, and ß-
alanine in *Sam50*-deficient myotubes (Figure 5A), which may have several potential
implications. Beyond functioning in embryonic neurogenesis and as a neurotransmitter, GABA
acts in a potential mTOR-dependent manner with accumulation resulting in mitophagy and

abnormal morphology (67). Typically GABA is metabolized by a pathway known as the GABA
shunt, found in the mitochondrial matrix; decreased frequency of GABA may be linked to
upregulated upstream targets in the GABA shunt (68). It is well understood that the primary role
of hypotaurine is as a precursor for taurine when reacted with hydrogen peroxide (69). Critically,
taurine serves numerous functions beyond antioxidant roles, protecting against mitochondrial
disease, potentially through posttranslational modifications of mitochondrial tRNAs (70,71).
Together, these studies suggest a potential antioxidant role of hypotaurine in mitochondria (72).
Similar to hypotaurine, ß-alanine is also a precursor to taurine production. Studies have
suggested that ß-alanine supplement increases skeletal muscle carnosine content, which affects
calcium homeostasis and offers antioxidant abilities, which can increase muscle endurance, yet
these results remain debated (73,74). Interestingly, however, increased ß-alanine has also been
shown to result in mitochondrial fragmentation and oxidative stress (75). Overall, results across
prior studies show varied results suggesting a tissue-dependent role of ß-alanine which requires
further elucidation.
These reductions in ß-alanine parallel a larger trend of amino acid metabolism,
evidenced by decreased metabolites of many amino acids (Figure 5B). This can have numerous
implications on the dependency of *Sam50* for overall mitochondrial health during disease states.
For example, the catabolism of fatty acids and amino acids is commonly observed as a method to
bolster cancer growth (76). These changes may also be affecting the mTORC pathway (77).
Recently it was found that the mTORC pathway interacts in an OPA-1 mediated manner and
may be associated with the overall mitochondrial structure in stress states (78). Indeed, it has
been found that the ATF4-mediated integrated stress response (ISR) is responsible for mediating
cellular responses to oxidative stress and regulating amino acid metabolism (79). This pathway
may be activated by ER or mitochondrial stress-dependent eIF2α phosphorylation (80). Future
studies may aim to understand if metabolite pathways are changing in response to the structural-
related declines in respiratory efficiency, or if the inverse is happening.
Alongside metabolic changes, mitochondrial structure may be related to alterations in
autophagy that we observed. Upregulation of autophagy may be targeting ROS in mitochondria,
which is often the cause of mitophagy activation (81,82). Interestingly, past research has found
that cytotoxic ROS-dependent molecules enter the mitochondria in a Sam50-mediated manner to
cause ROS build-up or apoptosis (83). Therefore, loss of Sam50 may serve a cytoprotective
effect by preventing apoptosis and increasing proliferation in cancer cells (84). Beyond this,
Sam50 loss also contributes to an excess of ROS species (24,84), which contributes to many
pathologies. Therefore, restoring *Sam50* may act as a mechanism to prevent ROS overproduction
(84) and restore mitochondrial and cristae structure (53).
Together, our results additionally show that *Sam50* loss may further contribute to
pathology through impairing mitochondrial structures, such as toroid shape and elongated
mitochondria, reducing OCR, and altering metabolic pathways. Therefore, future research
evaluating the targeting of *Sam50* to restore mitochondria 3D structure, function, and metabolism
during *in vivo* disease states may prove a valuable future avenue.

Financial & Competing Interests Disclosure

All authors have no competing interests.

This project was funded by the National Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; National Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to A.C.; Integrated Training in Engineering and Diabetes", Grant Number T32 DK101003; Burroughs Wellcome Fund Postdoctoral Enrichment Program #1022355 to D.S.; The UNCF/ Bristol-Myers Squibb (UNCF/BMS)- E.E. Just Postgraduate Fellowship in Life sciences Fellowship and Burroughs Wellcome Fund/ PDEP #1022376 to H.K.B.; NSF MCB #2011577I to S.A.M.; NSF EES2112556, NSF EES1817282, NSF MCB1955975, and CZI Science Diversity Leadership grant number 2022-253614 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation to S.D.; The UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, Career Award at the Scientific Interface (CASI Award) from Burroughs Welcome Fund (BWF) ID # 1021868.01, BWF Ad-hoc Award, NIH Small Research Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program, DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation to A.H.J.; and National Institutes of Health grant HD090061 and the Department of Veterans Affairs Office of Research award I01 BX005352 (to J.G.); Howard Hughes Medical Institute Hanna H. Gray Fellows Program Faculty Phase (Grant# GT15655 awarded to M.R.M); and Burroughs Wellcome Fund PDEP Transition to Faculty (Grant# 1022604 awarded to M.R.M). National Institutes of Health Grants: R21DK119879 (to C.R.W.) and R01DK-133698 (to C.R.W.), American Heart Association Grant 16SDG27080009 (to C.R.W.) and by an American Society of Nephrology KidneyCure Transition to Independence Grant (to C.R.W.). Additional support was provided by the Vanderbilt Institute for Clinical and Translational Research program supported by the National Center for Research Resources, Grant UL1 RR024975–01, and the National Center for Advancing Translational Sciences, Grant 2 UL1 TR000445–06 and the Cell Imaging Shared Resource. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of the NIH. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Contributions:

B.S., M.K., A.O., and C.V. performed the experimental procedures, data analysis, interpretation of results, and writing of the manuscript. They share equal authorship.

F.Z., K.N., Z.V., and E.G.L. contributed to the experimental design and data analysis.

A.K, Z.V., and E.G.L. J.Q.S., M.M., J.L., and Q.W. contributed to the data collection and analysis.

K.N., C.T.A., A.W., J.B., A.K., A.G.M., E.S., L.V., H.J.K., M.R.M., J.AG., S.D., B.C.J., A.T.C.,

S.M., D.S., and A.C. provided critical feedback and contributed to the writing and revision of the

manuscript.

H.K.B and A.H. conceptualized the study, supervised the research, provided overall guidance,

and revised the manuscript.

All authors reviewed and approved the final version of the manuscript.

Data Sharing and Open Access

All data is available upon request to the corresponding author.

Figure Legend:

Figure 1: Workflow of SBF-SEM manual contour reconstruction to create 3D

mitochondrial structure upon loss of *Sam50*. (A) Workflow depicting isolation of *Sam50*-

deficient human myotubes. SBF-SEM allows for (B) ortho slice alignment. (C) Manual

segmentation of ortho slices was performed to yield (D) 3D reconstructions of mitochondria. (E)

Once 3D reconstruction was performed, quantifications were performed for mitochondria.

Figure 2: SBF-SEM shows changes in human and murine myotubes mitochondrial 3D

structure upon loss of *Sam50*. (A) Control and (B) *Sam50* knockout murine myotubes

orthoslice, displayed. Approximately 50 of these orthoslices are segmented to perform 3D

reconstruction. (A') Control and (B') *Sam50*-deficient myotubes 3D renderings displayed

overlaid on SBF-SEM orthoslice. (A'') Isolated 3D mitochondria from control and (B'') *Sam50*-

deficient myotubes. (C) Upon loss of Sam50, on average mitochondria have reduced volume, (D)

surface area, (E) and perimeter. (A''') Control and (B''') *Sam50* knockout human myotubes

orthoslice, displayed. Approximately 50 of these orthoslices are segmented to perform 3D

reconstruction. (A'''') Control and (B'''') *Sam50*-deficient myotubes 3D renderings displayed

overlaid on SBF-SEM orthoslice. (A''''') Isolated 3D mitochondria from control and (B''''')

Sam50-deficient myotubes. (C') Upon loss of *Sam50* in human myotubes, on average

mitochondria have reduced volume, (D') surface area, (E') and perimeter.

Figure 3: **Changes in human and murine myotubes mitochondrial branching and**

networking upon loss of Sam50 revealed in SBF-SEM. (A) 3D Reconstructions showing

control and (B) *Sam50*-deficient murine and human (A''-B'') myotubes from a transverse-point

of view. Murine (A') and human (A''') 3D Reconstructions showing control and (B') *Sam50*-

deficient murine and human (B''') myotubes from a longitudinal point of view. (C) Sphericity of

mitochondria in control and *Sam50*-deficient murine and (C') murine myotubes. (D)

Mitochondrial complexity index (MCI), which is analogous to sphericity, comparing control and

Sam50-deficient murine and (D') human myotubes. (E) Mito-otyping, to display the diversity of

mitochondrial phenotypes as ordered by volume, in control and *Sam50*-deficient murine and (E')
human myotubes.

Figure 4: Changes in macroautophagic autophagosomes count and volume upon loss of
***Sam50* revealed in human myotubes through SBF-SEM.** (A) 3D Reconstruction overlaid on
z-directional orthoslice of autophagosomes in control myotubes and (B) *Sam50* knockout (KO)
myotubes. (A') Isolated 3D reconstruction of autophagosomes in control and (B') Sam50 KO
myotubes. Quantifications of (C) the number of autophagosomes and (D) the average volume of
autophagosomes upon loss of *Sam50* in human myotube. (E) Autophagosomes organized on the
basis of their volume going from left to right to illustrate the diversity of recycling machinery.

Figure 5: XF24 Seahorse Analyzer measured respiration rate in murine and human
myotubes. Validation through quantitative polymerase chain reaction of knockout of *Sam50*. (A)
Decreased mRNA transcripts of *Sam50* in murine myotubes and (A') human myotubes following
CRISPR/Cas9 transient knockout. (B) *Sam50*-deficient murine and (B') human myotubes
showing reduced oxygen consumption rate (OCR) in comparison to control. (C) The first 4 time
points measure basal respiration, or mitochondrial OCR at baseline in murine and (C') human
myotubes. (D) The next three time points following application of oligomycin (1 µg/mL) shows
decoupling of respiration dependent on ATP in murine and (D') human myotubes. (E) The next
three time points measured following the addition of carbonyl cyanide 4-
(trifluoromethoxy)phenylhydrazone (FCCP; 1 µM) shows OCR under maximum respiration in
murine and (E') human myotubes. (F) The final three time points show OCR following rotenone
(1 µM) and antimycin A (10 µM) show minimum OCR that is not dependent on mitochondria in
murine and (F') human myotubes.

Figure 6: Metabolomic Comparison of *Sam50*-deficient human myotubes and control
myotubes. (A) Volcano plot of metabolites that were differentially regulated in *Sam50*-deficient
myotubes compared to WT samples. (B) Relative abundance of ions in *Sam50*-deficient versus
control myotubes visualized on a heatmap. (C) Enrichment analysis displaying enriched
metabolites in Sam50-*deficient* myotubes.

References:
1. Alston CL, Rocha MC, Lax NZ, Turnbull DM, Taylor RW. The genetics and pathology of
   mitochondrial disease. *J Pathol* (2017) 241:236–250. doi: 10.1002/path.4809

2. Duchen MR, Szabadkai G. Roles of mitochondria in human disease. *Essays Biochem*
   (2010) 47:115–137. doi: 10.1042/bse0470115

3. Bustos G, Cruz P, Lovy A, Cárdenas C. Endoplasmic reticulum–mitochondria calcium
communication and the regulation of mitochondrial metabolism in cancer: a novel potential
target. *Front Oncol* (2017) 7:199.

4. Yee ML, Wong R, Datta M, Fazio TN, Mohammad Ebrahim M, Mcnamara EC, De Jong G,
Gilfillan C. Mitochondrial disease: an uncommon but important cause of diabetes mellitus.
Endocrinol Diabetes Metab Case Rep (2018) 2018:18–0091. doi: 10.1530/EDM-18-0091

5. Rolo AP, Palmeira CM. Diabetes and mitochondrial function: Role of hyperglycemia and
oxidative stress. *Toxicol Appl Pharmacol* (2006) 212:167–178. doi:
10.1016/j.taap.2006.01.003

6. Li L, Wu C-S, Hou G-M, Dong M-Z, Wang Z-B, Hou Y, Schatten H, Zhang G-R, Sun Q-
Y. Type 2 diabetes increases oocyte mtDNA mutations which are eliminated in the
offspring by bottleneck effect. *Reprod Biol Endocrinol* (2018) 16:110. doi:
10.1186/s12958-018-0423-1

7. Dabravolski SA, Orekhova VA, Baig MS, Bezsonov EE, Starodubova AV, Popkova TV,
Orekhov AN. The Role of Mitochondrial Mutations and Chronic Inflammation in Diabetes.
Int J Mol Sci (2021) 22:6733. doi: 10.3390/ijms22136733

8. Bratic A, Larsson N-G. The role of mitochondria in aging. *J Clin Invest* (2013) 123:951–
957.

9. Cogliati S, Enriquez JA, Scorrano L. Mitochondrial cristae: where beauty meets
functionality. *Trends Biochem Sci* (2016) 41:261–273.

10. Glancy B, Kim Y, Katti P, Willingham TB. The Functional Impact of Mitochondrial
Structure Across Subcellular Scales. *Front Physiol* (2020) 11:
https://www.frontiersin.org/articles/10.3389/fphys.2020.541040 [Accessed December 16,
2022]

11. Long Q, Zhao D, Fan W, Yang L, Zhou Y, Qi J, Wang X, Liu X. Modeling of
Mitochondrial Donut Formation. *Biophys J* (2015) 109:892–9. doi:
10.1016/j.bpj.2015.07.039

12. Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE, Chan DC. Mitofusins Mfn1 and
Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic
development. *J Cell Biol* (2003) 160:189–200. doi: 10.1083/jcb.200211046

13. Frezza C, Cipolat S, De Brito OM, Micaroni M, Beznoussenko GV, Rudka T, Bartoli D,
Polishuck RS, Danial NN, De Strooper B. OPA1 controls apoptotic cristae remodeling
independently from mitochondrial fusion. *Cell* (2006) 126:177–189.

14. Hu C, Shu L, Huang X, Yu J, Li L, Gong L, Yang M, Wu Z, Gao Z, Zhao Y, et al. OPA1
and MICOS Regulate mitochondrial crista dynamics and formation. *Cell Death Dis* (2020)
11:1–17. doi: 10.1038/s41419-020-03152-y

15. Lee H, Smith SB, Yoon Y. The short variant of the mitochondrial dynamin OPA1 maintains
mitochondrial energetics and cristae structure. *J Biol Chem* (2017) 292:7115–7130. doi:
10.1074/jbc.M116.762567

16. Kühlbrandt W. Structure and function of mitochondrial membrane protein complexes. *BMC*
Biol (2015) 13:89. doi: 10.1186/s12915-015-0201-x

17. Kozjak-Pavlovic V. The MICOS complex of human mitochondria. *Cell Tissue Res* (2017)
367:83–93.

18. Friedman JR, Mourier A, Yamada J, McCaffery JM, Nunnari J. MICOS coordinates with
respiratory complexes and lipids to establish mitochondrial inner membrane architecture.
Elife (2015) 4:e07739.

19. Rampelt H, Zerbes RM, van der Laan M, Pfanner N. Role of the mitochondrial contact site
and cristae organizing system in membrane architecture and dynamics. *Biochim Biophys*
Acta BBA-Mol Cell Res (2017) 1864:737–746.

20. Anand R, Reichert AS, Kondadi AK. Emerging Roles of the MICOS Complex in Cristae
Dynamics and Biogenesis. *Biology* (2021) 10:600. doi: 10.3390/biology10070600

21. Ding C, Wu Z, Huang L, Wang Y, Xue J, Chen S, Deng Z, Wang L, Song Z, Chen S.
Mitofilin and CHCHD6 physically interact with Sam50 to sustain cristae structure. *Sci Rep*
(2015) 5:1–11.

22. Darshi M, Taylor SS. Mitochondrial ChChD3 acts as a Scaffold for Mitofilin, Sam50 and
PKA. (2008)

23. Kozjak V, Wiedemann N, Milenkovic D, Lohaus C, Meyer HE, Guiard B, Meisinger C,
Pfanner N. An Essential Role of Sam50 in the Protein Sorting and Assembly Machinery of
the Mitochondrial Outer Membrane *. *J Biol Chem* (2003) 278:48520–48523. doi:
10.1074/jbc.C300442200

24. Lionello S, Marzaro G, Martinvalet D. SAM50, a side door to the mitochondria: The case
of cytotoxic proteases. *Pharmacol Res* (2020) 160:105196. doi:
10.1016/j.phrs.2020.105196

25. Ott C, Ross K, Straub S, Thiede B, Götz M, Goosmann C, Krischke M, Mueller MJ,
Krohne G, Rudel T, et al. Sam50 Functions in Mitochondrial Intermembrane Space
Bridging and Biogenesis of Respiratory Complexes. *Mol Cell Biol* (2012) 32:1173–1188.
doi: 10.1128/MCB.06388-11

26. Körner C, Barrera M, Dukanovic J, Eydt K, Harner M, Rabl R, Vogel F, Rapaport D,
Neupert W, Reichert AS. The C-terminal domain of Fcj1 is required for formation of crista
junctions and interacts with the TOB/SAM complex in mitochondria. *Mol Biol Cell* (2012)
23:2143–2155. doi: 10.1091/mbc.E11-10-0831

27. Ott C, Dorsch E, Fraunholz M, Straub S, Kozjak-Pavlovic V. Detailed analysis of the
human mitochondrial contact site complex indicate a hierarchy of subunits. *PloS One*
(2015) 10:e0120213.

28. Lam J, Katti P, Biete M, Mungai M, AshShareef S, Neikirk K, Garza Lopez E, Vue Z,
Christensen TA, Beasley HK, et al. A Universal Approach to Analyzing Transmission
Electron Microscopy with ImageJ. *Cells* (2021) 10:2177. doi: 10.3390/cells10092177

29. Neikirk K, Lopez E-G, Marshall AG, Alghanem A, Krystofiak E, Kula B, Smith N, Shao J,
Katti P, Hinton AO. Call to Action to Properly Utilize Electron Microscopy to Measure
Organelles to Monitor Disease. *Eur J Cell Biol* (2023)151365. doi:
10.1016/j.ejcb.2023.151365

30. Vue Z, Garza-Lopez E, Neikirk K, Vang L, Beasley H, Marshall AG, Murray SA, Hinton Jr
A. Mouse Skeletal Muscle Decrease in the MICOS Complex and Altered Mitochondrial
Networks with age. *bioRxiv* (2022)2022–03.

31. Vue Z, Neikirk K, Vang L, Garza-Lopez E, Christensen TA, Shao J, Lam J, Beasley HK,
Marshall AG, Crabtree A, et al. Three-Dimensional Mitochondria Reconstructions of
Murine Cardiac Muscle Changes in Size Across Aging. *Am J Physiol-Heart Circ Physiol*
(2023) doi: 10.1152/ajpheart.00202.2023

32. Javadov S, Chapa-Dubocq X, Makarov V. Different approaches to modeling analysis of
mitochondrial swelling. *Mitochondrion* (2018) 38:58–70. doi: 10.1016/j.mito.2017.08.004

33. Ahmad T, Aggarwal K, Pattnaik B, Mukherjee S, Sethi T, Tiwari B, Kumar M, Micheal A,
Mabalirajan U, Ghosh B, et al. Computational classification of mitochondrial shapes
reflects stress and redox state. *Cell Death Dis* (2013) 4:e461. doi: 10.1038/cddis.2012.213

34. Garza-Lopez E, Vue Z, Katti P, Neikirk K, Biete M, Lam J, Beasley HK, Marshall AG,
Rodman TA, Christensen TA, et al. Protocols for Generating Surfaces and Measuring 3D
Organelle Morphology Using Amira. *Cells* (2022) 11:65. doi: 10.3390/cells11010065

35. Vincent AE, White K, Davey T, Philips J, Ogden RT, Lawless C, Warren C, Hall MG, Ng
YS, Falkous G, et al. Quantitative 3D Mapping of the Human Skeletal Muscle
Mitochondrial Network. *Cell Rep* (2019) 26:996-1009.e4. doi:
10.1016/j.celrep.2019.01.010

36. Marshall AG, Damo SM, Hinton A. Revisiting focused ion beam scanning electron
microcopy. *Trends Biochem Sci* (2023) 0: doi: 10.1016/j.tibs.2023.02.005

37. Marshall AG, Krystofiak E, Damo SM, Hinton A. Correlative light-electron microscopy:
integrating dynamics to structure. *Trends Biochem Sci* (2023)S0968-0004(23)00127-5. doi:
10.1016/j.tibs.2023.05.003

38. Marshall AG, Neikirk K, Stephens DC, Vang L, Vue Z, Beasley HK, Crabtree A, Scudese
E, Lopez EG, Shao B, et al. Serial Block Face-Scanning Electron Microscopy as a
Burgeoning Technology. *Adv Biol* (2023)e2300139. doi: 10.1002/adbi.202300139

39. Smolina N, Kostareva A, Bruton J, Karpushev A, Sjoberg G, Sejersen T. Primary Murine
Myotubes as a Model for Investigating Muscular Dystrophy. *BioMed Res Int* (2015)
2015:e594751. doi: 10.1155/2015/594751

40. Boldrin L, Muntoni F, Morgan JE. Are Human and Mouse Satellite Cells Really the Same?
J Histochem Cytochem (2010) 58:941–955. doi: 10.1369/jhc.2010.956201

41. Aas V, Bakke SS, Feng YZ, Kase ET, Jensen J, Bajpeyi S, Thoresen GH, Rustan AC. Are
cultured human myotubes far from home? *Cell Tissue Res* (2013) 354:671–682. doi:
10.1007/s00441-013-1655-1

42. Pereira RO, Tadinada SM, Zasadny FM, Oliveira KJ, Pires KMP, Olvera A, Jeffers J,
Souvenir R, Mcglauflin R, Seei A. OPA 1 deficiency promotes secretion of FGF 21 from
muscle that prevents obesity and insulin resistance. *EMBO J* (2017) 36:2126–2145.

43. Boudina S, Sena S, Theobald H, Sheng X, Wright JJ, Hu XX, Aziz S, Johnson JI, Bugger
H, Zaha VG, et al. Mitochondrial energetics in the heart in obesity-related diabetes: direct
evidence for increased uncoupled respiration and activation of uncoupling proteins.
Diabetes (2007) 56:2457–2466. doi: 10.2337/db07-0481

44. Dranka BP, Benavides GA, Diers AR, Giordano S, Zelickson BR, Reily C, Zou L, Chatham
JC, Hill BG, Zhang J, et al. Assessing bioenergetic function in response to oxidative stress
by metabolic profiling. *Free Radic Biol Med* (2011) 51:1621–1635. doi:
10.1016/j.freeradbiomed.2011.08.005

45. Hinton A, Katti P, Christensen TA, Mungai M, Shao J, Zhang L, Trushin S, Alghanem A,
Jaspersen A, Geroux RE. A comprehensive approach for artifact-free sample preparation
and assessment of mitochondrial morphology in tissue and cultured cells. *bioRxiv* (2021)

46. Phillips MA, Arnold KR, Vue Z, Beasley HK, Garza-Lopez E, Marshall AG, Morton DJ,
McReynolds MR, Barter TT, Hinton A. Combining Metabolomics and Experimental
Evolution Reveals Key Mechanisms Underlying Longevity Differences in Laboratory
Evolved Drosophila melanogaster Populations. *Int J Mol Sci* (2022) 23:1067. doi:
10.3390/ijms23031067

47. Li B, Tang J, Yang Q, Li S, Cui X, Li Y, Chen Y, Xue W, Li X, Zhu F. NOREVA:
normalization and evaluation of MS-based metabolomics data. *Nucleic Acids Res* (2017)
45:W162–W170.

48. Marshall AG, Garza-Lopez E, Vue Z, Vang L, Hinton A Jr. Sam50 is Associated with
Fragmentation and Alterations in Metabolism in Human and Murine Myotubes. *Microsc*
Microanal (2023) 29:1189–1191. doi: 10.1093/micmic/ozad067.613

49. Kamat PK, Kalani A, Kyles P, Tyagi SC, Tyagi N. Autophagy of Mitochondria: A
Promising Therapeutic Target for Neurodegenerative Disease. *Cell Biochem Biophys*
(2014) 70:707–719. doi: 10.1007/s12013-014-0006-5

50.   Abudu YP, Shrestha BK, Zhang W, Palara A, Brenne HB, Larsen KB, Wolfson DL,
Dumitriu G, Øie CI, Ahluwalia BS, et al. SAMM50 acts with p62 in piecemeal basal- and
OXPHOS-induced mitophagy of SAM and MICOS components. *J Cell Biol* (2021)
220:e202009092. doi: 10.1083/jcb.202009092

51.   Jian F, Chen D, Chen L, Yan C, Lu B, Zhu Y, Chen S, Shi A, Chan DC, Song Z. Sam50
Regulates PINK1-Parkin-Mediated Mitophagy by Controlling PINK1 Stability and
Mitochondrial Morphology. *Cell Rep* (2018) 23:2989–3005. doi:
10.1016/j.celrep.2018.05.015

52.   Neikirk K, Vue Z, Katti P, Rodriguez BI, Omer SA, Shao J, Christensen T, Lopez EG,
Marshall A, Palavicino-Maggio CB, et al. Systematic Transmission Electron Microscopy-
Based Identification and 3D Reconstruction of Cellular Degradation Machinery.
(2022)2021.09.26.461841. doi: 10.1101/2021.09.26.461841

53.   Yin X, Wang J, Yang S, Li H, Shen H, Wang H, Li X, Chen G. Sam50 exerts
neuroprotection by maintaining the mitochondrial structure during experimental cerebral
ischemia/reperfusion injury in rats. *CNS Neurosci Ther* (2022) 28:2230–2244. doi:
10.1111/cns.13967

54.   Vidavsky N, Akiva A, Kaplan-Ashiri I, Rechav K, Addadi L, Weiner S, Schertel A. Cryo-
FIB-SEM serial milling and block face imaging: Large volume structural analysis of
biological tissues preserved close to their native state. *J Struct Biol* (2016) 196:487–495.
doi: 10.1016/j.jsb.2016.09.016

55.   Santos D, Esteves AR, Silva DF, Januário C, Cardoso SM. The Impact of Mitochondrial
Fusion and Fission Modulation in Sporadic Parkinson's Disease. *Mol Neurobiol* (2015)
52:573–586. doi: 10.1007/s12035-014-8893-4

56.   Safiulina D, Veksler V, Zharkovsky A, Kaasik A. Loss of mitochondrial membrane
potential is associated with increase in mitochondrial volume: physiological role in
neurones. *J Cell Physiol* (2006) 206:347–353. doi: 10.1002/jcp.20476

57.   Ghochani M, Nulton JD, Salamon P, Frey TG, Rabinovitch A, Baljon ARC. Tensile forces
and shape entropy explain observed crista structure in mitochondria. *Biophys J* (2010)
99:3244–3254. doi: 10.1016/j.bpj.2010.09.038

58.   Putti R, Sica R, Migliaccio V, Lionetti L. Diet impact on mitochondrial bioenergetics and
dynamics. *Front Physiol* (2015) 6:
https://www.frontiersin.org/articles/10.3389/fphys.2015.00109 [Accessed February 7,
2023]

59.   Bleck CK, Kim Y, Willingham TB, Glancy B. Subcellular connectomic analyses of energy
networks in striated muscle. *Nat Commun* (2018) 9:1–11.

60.   Moltedo O, Remondelli P, Amodio G. The mitochondria–endoplasmic reticulum contacts
and their critical role in aging and age-associated diseases. *Front Cell Dev Biol* (2019)
7:172.

61.   Giacomello M, Pellegrini L. The coming of age of the mitochondria–ER contact: a matter
775   of thickness. *Cell Death Differ* (2016) 23:1417–1427.

776   62.   Axe EL, Walker SA, Manifava M, Chandra P, Roderick HL, Habermann A, Griffiths G,
777   Ktistakis NT. Autophagosome formation from membrane compartments enriched in
778   phosphatidylinositol 3-phosphate and dynamically connected to the endoplasmic reticulum.
779   *J Cell Biol* (2008) 182:685–701.

780   63.   Zhang X, Monroe ME, Chen B, Chin MH, Heibeck TH, Schepmoes AA, Yang F, Petritis
781   BO, Camp DG, Pounds JG, et al. Endogenous 3,4-Dihydroxyphenylalanine and
782   Dopaquinone Modifications on Protein Tyrosine: Links to Mitochondrially Derived
783   Oxidative Stress via Hydroxyl Radical *. *Mol Cell Proteomics* (2010) 9:1199–1208. doi:
784   10.1074/mcp.M900321-MCP200

785   64.   Schoneich C, Hewarathna A, Pal R, Jiang L, Michaelis E. P 236 - Oxidative Stress Markers
786   of Alzheimer's Disease in Peripheral Cell Mitochondria. *Free Radic Biol Med* (2017)
787   108:S100. doi: 10.1016/j.freeradbiomed.2017.04.321

788   65.   Li Z, Shen W, Wu G, Qin C, Zhang Y, Wang Y, Song G, Xiao C, Zhang X, Deng G, et al.
789   The role of SAMM50 in non-alcoholic fatty liver disease: from genetics to mechanisms.
790   *FEBS Open Bio* (2021) 11:1893–1906. doi: 10.1002/2211-5463.13146

791   66.   Parra V, Verdejo HE, Iglewski M, del Campo A, Troncoso R, Jones D, Zhu Y, Kuzmicic J,
792   Pennanen C, Lopez☐Crisosto C, et al. Insulin Stimulates Mitochondrial Fusion and
793   Function in Cardiomyocytes via the Akt-mTOR-NFκB-Opa-1 Signaling Pathway. *Diabetes*
794   (2013) 63:75–88. doi: 10.2337/db13-0340

795   67.   Lakhani R, Vogel KR, Till A, Liu J, Burnett SF, Gibson KM, Subramani S. Defects in
796   GABA metabolism affect selective autophagy pathways and are alleviated by mTOR
797   inhibition. *EMBO Mol Med* (2014) 6:551–566. doi: 10.1002/emmm.201303356

798   68.   Ravasz D, Kacso G, Fodor V, Horvath K, Adam-Vizi V, Chinopoulos C. Catabolism of
799   GABA, succinic semialdehyde or gamma-hydroxybutyrate through the GABA shunt impair
800   mitochondrial substrate-level phosphorylation. *Neurochem Int* (2017) 109:41–53. doi:
801   10.1016/j.neuint.2017.03.008

802   69.   Grove RQ, Karpowicz SJ. Reaction of hypotaurine or taurine with superoxide produces the
803   organic peroxysulfonic acid peroxytaurine. *Free Radic Biol Med* (2017) 108:575–584. doi:
804   10.1016/j.freeradbiomed.2017.04.342

805   70.   Jong CJ, Sandal P, Schaffer SW. The Role of Taurine in Mitochondria Health: More Than
806   Just an Antioxidant. *Molecules* (2021) 26:4913. doi: 10.3390/molecules26164913

807   71.   Suzuki BH. Revisiting the Model Minority Stereotype: Implications for Student Affairs
808   Practice and Higher Education. *New Dir Stud Serv* (2002) 2002:21–32. doi: 10.1002/ss.36

809   72.   Ubuka T, Okada A, Nakamura H. Production of hypotaurine from l-cysteinesulfinate by rat
810   liver mitochondria. *Amino Acids* (2008) 35:53–58. doi: 10.1007/s00726-007-0633-x

73.   Saunders B, DE Salles Painelli V, DE Oliveira LF, DA Eira Silva V, DA Silva RP, Riani L,
Franchi M, Gonçalves LDS, Harris RC, Roschel H, et al. Twenty-four Weeks of β-Alanine
Supplementation on Carnosine Content, Related Genes, and Exercise. *Med Sci Sports Exerc*
(2017) 49:896–906. doi: 10.1249/MSS.0000000000001173

74.   Bhattacharya TK, Pence BD, Ossyra JM, Gibbons TE, Perez S, McCusker RH, Kelley KW,
Johnson RW, Woods JA, Rhodes JS. Exercise but not (–)-epigallocatechin-3-gallate or β-
alanine enhances physical fitness, brain plasticity, and behavioral performance in mice.
Physiol Behav (2015) 145:29–37. doi: 10.1016/j.physbeh.2015.03.023

75.   Shetewy A, Shimada-Takaura K, Warner D, Jong CJ, Mehdi A-BA, Alexeyev M,
Takahashi K, Schaffer SW. Mitochondrial defects associated with β-alanine toxicity:
relevance to hyper-beta-alaninemia. *Mol Cell Biochem* (2016) 416:11–22. doi:
10.1007/s11010-016-2688-z

76.   Li Z, Zhang H. Reprogramming of glucose, fatty acid and amino acid metabolism for
cancer progression. *Cell Mol Life Sci* (2016) 73:377–392. doi: 10.1007/s00018-015-2070-4

77.   Goberdhan DCI, Wilson C, Harris AL. Amino Acid Sensing by mTORC1: Intracellular
Transporters Mark the Spot. *Cell Metab* (2016) 23:580–589. doi:
10.1016/j.cmet.2016.03.013

78.   Hinton A, Katti P, Mungai M, Hall D, Koval O, Shao J, Vue Z, Christensen TA, Lopez EG,
Rostami R, et al. OPA1 Downregulation in Skeletal Muscle Induces MERC formation in an
ATF4-Dependent Manner. (2022)2022.09.12.507669. doi: 10.1101/2022.09.12.507669

79.   Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C, Popko B,
Paules R, et al. An Integrated Stress Response Regulates Amino Acid Metabolism and
Resistance to Oxidative Stress. *Mol Cell* (2003) 11:619–633. doi: 10.1016/S1097-
2765(03)00105-9

80.   Tian X, Zhang S, Zhou L, Seyhan AA, Hernandez Borrero L, Zhang Y, El-Deiry WS.
Targeting the Integrated Stress Response in Cancer Therapy. *Front Pharmacol* (2021) 12:
https://www.frontiersin.org/articles/10.3389/fphar.2021.747837 [Accessed February 7,
2023]

81.   Basit F, van Oppen LM, Schöckel L, Bossenbroek HM, van Emst-de Vries SE, Hermeling
JC, Grefte S, Kopitz C, Heroult M, Hgm Willems P, et al. Mitochondrial complex I
inhibition triggers a mitophagy-dependent ROS increase leading to necroptosis and
ferroptosis in melanoma cells. *Cell Death Dis* (2017) 8:e2716. doi: 10.1038/cddis.2017.133

82.   Chen G, Kroemer G, Kepp O. Mitophagy: An Emerging Role in Aging and Age-Associated
Diseases. *Front Cell Dev Biol* (2020) 8:
https://www.frontiersin.org/articles/10.3389/fcell.2020.00200 [Accessed December 21,
2022]

83.   Chiusolo V, Jacquemin G, Yonca Bassoy E, Vinet L, Liguori L, Walch M, Kozjak-Pavlovic
V, Martinvalet D. Granzyme B enters the mitochondria in a Sam50-, Tim22- and mtHsp70-

849    dependent manner to induce apoptosis. *Cell Death Differ* (2017) 24:747–758. doi:
850    10.1038/cdd.2017.3

851  84. Capala ME, Pruis M, Vellenga E, Schuringa JJ. Depletion of SAM50 Specifically Targets
852    BCR-ABL-Expressing Leukemic Stem and Progenitor Cells by Interfering with
853    Mitochondrial Functions. *Stem Cells Dev* (2016) 25:427–437. doi: 10.1089/scd.2015.0151

A

Extraction of
skeletal muscle

CRISPR/Cas9
knockout of *Sam50*

Differentiation to
Myotubes

B

SBF-SEM
3D imaging

10μm
10μm
50μm

2μm

C

1	2	3	4
5	6	7	8
9	10	11	12
13	14	15	16

0.5μm

D

90°

0.5μm

E

Surface Area = Σ pixels
outside of object x

3D perimeter =
Boundary Pixel Count

Volume = Total
Displacement/
Internal Pixels
of Mitochondria

Sphericity =

$$\frac{\pi^{1/3}(6*Volume)^{2/3}}{Surface\ Area}$$

Mitochondria
Complexity
Index =

$$\frac{Surface\ Area^3}{16\pi^2 Volume^2}$$

Murine

Human

WT — *Sam50* KO

A — Transverse
A' — Longitudinal
B — Transverse
B' — Longitudinal

C — Sphericity ****
D — Complexity Index ****

A'' — Transverse
A''' — Longitudinal
B'' — Transverse
B''' — Longitudinal

C' — Sphericity ****
D' — Complexity Index ****

E — Mito-otype
WT
Sam50 KO

E' — Mito-otype
WT
Sam50 KO

1µm

0.5µm