A method for extracting an approximated connectome from libraries of single anatomically co-registered cell reconstructions

Manjunatha K.K.H.[2,4#], Bruzzone M.[1,3,#], Nicoletti G.[2,3], Suweis S.[2,3,@] and dal Maschio M. [1,3,#,@]

[1,] Department of Biomedical Sciences, Università degli Studi di Padova, Padua, IT
[2,] Department of Physics and Astronomy, Università degli Studi di Padova, Padua, IT
[3,] Padova Neuroscience Center, Università degli Studi di Padova, Padua, IT
[4,] Modelling and Engineering Risk and Complexity, Scuola Superiore Meridionale, Napoli, IT

[#,] equal contribution
[@,] corresponding author

Lead contact: marco.dalmaschio@unipd.it

Abstract

Mapping functional brain information onto a circuit wiring diagram supporting possible brain circuit working mechanisms remains challenging. When available, the analysis of electron microscopy based high-resolution connectomes, leveraging synapse annotation, allows for unraveling portions of the nervous system's wiring diagram or specific circuit motifs. However, obtaining a complete brain reconstruction with annotation of the synapses and extracting the general organization principles of the neuronal networks across the brain remains a demanding effort. To extend the available methods, we present here an approach to obtain an approximate brain connectome starting from libraries of single cell reconstructions co-registered in a reference anatomical space. Our approach relies on both the Strahler numbering of the nodes characterizing the cell morphological reconstructions and a proximity range criterion, to infer the general connectivity structures between the different cells, in the absence of synapse annotation. First, we confronted this approach on a ground truth derived from a fully annotated EM dataset of *Drosophila melanogaster*. The connectivity modules identified with our approach show generally a similar organization compared to those resulting from the EM annotated dataset, in terms of Normalized Mutual Information and Adjusted Rand Index, always with high significance with respect to random null models. Building on this, we then applied this approach to extract an approximate connectome of the zebrafish larvae brain starting from a light microscopy-based dataset of about 3000 co-registered neuronal skeletonizations. Modularity analysis of the retrieved

directed and undirected connectomes revealed a hierarchical organization of the resulting graph. The

identified architecture encompasses a series of high-order brain-wide neuronal modules composed of

a number of lower-order submodules. Importantly, we show that this approximated representation is

sufficient for capturing precise and topographically organized connection patterns as well as known

functional circuit architectures. In conclusion, we present a scalable, from-circuit-to-brain range,

approach to reveal neuronal architectures supporting brain mechanisms, potentially suitable for

hypothesis drawing and as for guiding the exploration of and integration with EM connectomes.

Introduction

The understanding of the brain mechanisms takes advantage of the continuous refinement of

molecular, hardware, and software approaches. These methods allow reconstructing, in suitable model

organisms, the brain dynamics from synapse level to brain-wide scales[1–5]. Current optical and

electrophysiological methods represent invaluable tools for studying the neuronal dynamics associated

with sensory information, the fine-tuning of motor programs, and spatiotemporal patterns

characterizing brain activity[6–8]. The obtained dataset becomes fundamental building blocks for the

formulation, and the possible falsification, of circuit models describing the functional working

mechanisms[9–11]. However, even under optimal conditions, one faces a typical scenario where different

models, despite their intrinsic architectural difference, number of parameters, and complexity, perform

similarly well in capturing the same circuit dynamics[12,13]. With the aim of better constraining this kind

of problem, one possible strategy is offered by the availability of the anatomical information

characterizing the circuit wiring diagram supporting a certain functional mechanism. Then, it is

convenient to try to refine a circuit working model combining functional aspects with the wiring

properties of the underlying circuit. For this purpose, one can take advantage of dense anatomical

reconstructions of large brain parcels based on Electron Microscopy (EM)[14–18], which achieves sufficient

resolution levels to characterize the synaptic contacts and possibly the neurotransmitters involved

between neuronal partners, while covering a large field of view. Retrieving the connectome across the

whole brain, along with an accessible organism, requires a relatively long acquisition time, supervised

or partially-supervised cell tracing, a cell segmentation step, annotation of the synaptic contacts, and

lastly a thorough control of the precision of the reconstructions. Today, even with the support of AI-

based algorithms, this is still a time-demanding process that limits its applicability to a few exemplars

and requires computational strategies to minimize human workload[15,19]. Once the information on the

neuronal connectivity schemes i.e. the organization of the input-output contacts characterizing the cells

involved- has been retrieved, it becomes possible to reconstruct an adjacency matrix representing a

connectome and identify the circuit wiring diagrams. Because of the amount and the complexity of the

data one has to face, this type of anatomical reconstruction frequently focuses on targeted circuit

motifs, specific neuronal subpopulations, or particular brain regions. On the other hand, unraveling the

anatomical organization of brain-wide networks requires suitable datasets, with a sufficient sampling

of the cellular morphologies, coverage across different regions of the brain, and, ideally, based on the

evaluation of a large number of exemplars. Currently, even if there are several reported whole-brain

EM reconstructions[15,16,18,20–22], partial or whole-brain connectomes that provide skeletonization of

neurons along with detailed annotations of their synaptic contacts are extremely limited. To extend

and complement the methods currently available, here, we present a computational approach to

generate an approximated connectome starting from an anatomical dataset with neuronal cell

skeletonizations, belonging to or co-registered in the same reference space, where the actual synaptic

information is not accessible or not yet available. This is the typical setting of mesoscale atlases based

on optical microscopy acquisitions, as well as electron microscopy datasets not yet fully annotated at

the detail level corresponding to the neuronal input-output features[23–28]. For this type of dataset, here,

we devised an approach to infer brain network organization by assigning cell-to-cell connections, which

are subsequently translated into adjacency matrices, representing the brain wide cell resolution

connectivity scheme. Following its evaluation on EM-based ground truth, we applied this approach to

a zebrafish single-cell dataset, which comprises more than three thousand neuronal reconstructions.

Results

1. Building an approximated connectome based on Strahler numbering and spatial proximity

For the cases where actual synaptic information is unavailable, we evaluated an approach to infer an

approximated brain network structure by assigning cell-to-cell connections on the basis of a proximity

rule between the nodes along the reconstructed cell skeleton (see the Methods section). For this

purpose, we devised to start from the information of the cell anatomy and calculated for each point of

the cell skeleton (or neurite) the corresponding Strahler number. This is a parameter that represents

the relative level of the cell skeleton points in the cell branching hierarchy[28] (Figure 1A,B). Terminal

points (endpoints) along the neurite structure, characterized by a unitary value of the Strahler number,

are considered putative points of connection between different cells if they fall within a defined

proximity range one from the other. For a typical light microscopy dataset, considering the intrinsic

resolution of the acquisition process, the typical errors in the anatomical reconstruction, and co-
registration procedures, we adopted a proximity range between 0 and 5 μm (Figure 1D).

As reconstructing the connectome relies on the determination of the adjacency matrix A_{ij}, we
envisaged two different possible scenarios. In the first one, we assume no subcellular segregations for
the input and output points of a neuron, resulting in a symmetric adjacency matrix A_{ij}^{U} corresponding
to an undirected connectome (Figure 1E). In the second one, as most of the neuronal reconstructions
available are characterized by a bimodal spatial distribution of the neuronal endpoints[29]
(Supplementary Figure 1), we assign for each cell a separation in two putative components: input
dendritic points in the regions proximal to the soma, and output axonal points in more distal areas. This
allows us to constrain the connectome according to an axon-to-dendrite synaptic scheme. In this latter
case, we obtain an asymmetric adjacency matrix A_{ij}^{D} representing a directed connectome (Figure 1E
panels at the bottom).

2. Evaluating the connectome reconstruction strategy against an EM-based ground truth

To assess the actual applicability of the approach and its capability to capture brain circuit organization
patterns, we took advantage from the synapse-annotated connectome of the Drosophila melanogaster
hemibrain, composed by a library of 21739 connected neurons[30]. Starting from the available
information on the input-output contacts obtained from the EM analysis, we identified using a
modularity analysis algorithm a hierarchical organization with eight main modules (module 1-8), each
accounting on average for about three thousand cells distributed in a variable number of submodules,
from four to nine, associated with a same main module (Figure 2A). We then considered the same
dataset and calculated the adjacency matrix and the modularity organization, but this time without
considering the available synapse annotation, but relying solely on the Strahler numbering of the cell
skeleton and on the proximity range definition to build an undirected graph (see Methods section). The
pairs of adjacency matrices obtained for each main module, one based on the synapse annotation (EM-
based) and one relying on the Strahler numbering (Strahler-based) , present Pearson correlation
coefficients ranging from 0.05 to 0.215 always with p-value smaller than 0.001 (see Methods), with

minimal variations considering the 1 μm and 5 μm proximity range (0.146 ± 0.053 and 0.14 ± 0.052,

average ± SD, respectively). In order to evaluate the capability of the approximated approach to actually

capture the expected wiring, we calculated for each of the main modules identified in the EM-based

analysis the Adjusted Rand Index (ARI)[31–33] based on the cell grouping associated with the organization

of the corresponding submodules. This metric is frequently used for comparing different partitions or

clustering patterns and returns a value between -1 and 1 to quantify the level of overlap in the

contingency matrix. We obtained for the 1 μm proximity range an average ARI of 0.36 ± 0.17 (average

± SD, covering a range between 0.04 and 0.6), pointing to a general overlap of the Strahler-based

approach with the ground truth sub-communities. A similar trend resulted for the 5 μm proximity

range, 0.38 ± 0.17 (average ± SD, Figure 2B, covering a range between 0.04 and 0.6). To compare the

community detection accuracy between the two datasets, we also computed the Normalized Mutual

Information (NMI)[34,35]. This metric returns a parameter between 0 and 1, depending on the level of

overlap between the communities using adjacency matrices and modularity analysis obtained with the

two approaches. NMI ranged between 0.1 and 0.55 (0.38 ± 0.17 average ± SD) and between 0.1 and

0.56 (0.38 ± 0.17 average ± SD) for 1 μm and 5 μm, respectively (Figure 2B). Then, to corroborate this

outcome, we evaluated the significance of the calculated ARI and NMI scores, with a comparison against

two types of random null models. We calculated the Z-score for NMI and ARI between the Strahler-

based connectome modules and the EM-based connectome modules, focusing here on two modules in

particular, module 6 and module 8, those characterized by the lowest and the highest, ARI and NMI

values. We constructed 100 random networks for the two Strahler-based connectome modules either

by randomizing the links or the weights of the links between the neurons (see Methods). Then, we

quantified the p-value by counting the number of random networks that had its own z-scores (for ARI

and NMI) higher than that one of the corresponding Strahler-based modules[36]. For each module we

found $p < 0.05$ highlighting that the Strahler-based model can capture connectivity patterns in the

networks better than models with the same number of nodes, but different links or link strength

organizations.

*Figure 2: Comparison with EM-based ground truth and null models. **A)** Decomposition of the higher order modules in submodules obtained with the second iteration of the algorithm. Colors and numbers are assigned according to the submodules. Length of the lines represents the number of submodules for each module. **B)** ARI and NMI computed between the EM ("ground truth") and Strahler reconstructed network at different proximity ranges. **C, D)** Z-scores computed over the ARI (panel **C**) and NMI (panel **D**) between Module 6 (or 8) of EM and Strahler reconstructed network (red point) and z-scores computed over the same mentioned similarity measures between EM and 100 random null models of Strahler reconstructed network (green points). The larger the gaps between the red point and the green points, the farther is the Strahler based networks from random behavior and closer is to the ground truth (EM based networks). *: p<0.05.*

3. Building an approximated connectome of the zebrafish larval brain

Aiming to build an approximated connectome for the zebrafish larval brain, we took advantage of an available dataset with anatomical reconstructions of individual neurons obtained by means of optical microscopy z-stack acquisitions co-registered on the same reference brain[27] (Figure 1A). Applying the proposed method, we retrieved two connectome scenarios showing a rather articulated scheme of neuronal interactions between the different regions of the brain (Figure 3B, Figure 3F). From the obtained adjacency matrices, A_{ij}^U, and A_{ij}^D (Figure 3A and Figure 3B), we extracted a set of graph parameters. In general, being the non-zero entries of the directed matrix a subset of the undirected

ones, A_{ij}^D resulted sparser than A_{ij}^U. The analysis of the undirected graph, A_{ij}^U, revealed the existence of

23 connected subgraphs, with a Giant Component (GC) accounting for 99% of the edges of the original

graph and a set of small, poorly populated disconnected subgraphs (accounting for 32 out of total 3163

cells), that have been neglected from further analyses. The GC presents a network density of 0.0397,

pointing to a rather limited connectivity, a small weighted clustering coefficient of 0.002 and an average

shortest path length of 3.792. The undirected graph shows a heterogeneous degree distribution, with

most of the nodes characterized by a relatively small number of connections ($<10^3$) and fewer ones –

hubs-like - with high degrees ($>10^4$) (Figure 3C). Interestingly, for low-degree nodes, the distribution

shows a flat trend, suggesting deviations from the power-law behavior. An assortativity analysis gives

a correlation coefficient β =0.345, meaning that the graph is assortative and that nodes tend to connect

with those of a similar degree – i.e., small-degree nodes to small-degree nodes and hubs to hubs. We

mapped hubs to the corresponding anatomical regions normalizing for the relative number of cells; it

appears that the majority belongs to Medulla Oblungata (Figure 3D - Pink), followed by the

Telencephalon (Figure3D - Cyan) and the Tectum Opticum (Figure 3D - Brown). On the other hand,

regarding the weighted directed graph, A_{ij}^D (Figure 3E), we found a low clustering coefficient (0.0005)

and a path length of 6.10. While the in- and out-degree distributions resemble the general

heterogeneity of the undirected scenario, the low-degree region did not show a corresponding

saturation (Figure 3G).

Figure 3. Topological properties of undirected and directed networks. A, E) Binarized adjacency matrixes showing connections in the undirected (A) and directed (E) networks with neurons sorted by brain regions in which the soma is located. The matrix is binarized for visualization purposes. B, F) Chord plots representing inter-regions connections for the undirected (B) network and incoming – outgoing inter-regions connection for the directed network (F). Labels for the colors are reported in A and E. C, G) Weighted degree distributions for the undirected (C) and directed (G) network. Dashed lines depict the threshold above which neurons are defined as hubs. D, H) Bar plots representing the distribution of hubs in the brain regions for the undirected (D) and directed (H) networks. Labels for the colors are reported in A and E.

4. Analysis of brain-wide modules from the approximated zebrafish larvae connectome

One of the most appealing uses of a brain wide connectome is unravelling possible network architecture and circuit organization motives that characterize an organism brain. We considered to use an established network community discovery strategy based on the modularity optimization, i.e. the Leiden algorithm[37], to analyze the network organization of the zebrafish connectome. Running the

algorithm 10,000 times on the undirected GC typically resulted in the stable and robust identification

of 12 neuronal modules, with small deviations in the assignment of a few nodes from trial to trial

(<0.5%). However, only eight of the identified modules are prominent and well populated, the

remaining containing an extremely limited number of nodes (4 out of 3116 and 14 out of 3131 in total

for directed and undirected GC graphs) (Figure 4A). To characterize the resulting networks, we first

mapped all identified undirected modules (U) on the brain regions containing the body of neurons

belonging to each corresponding module (Figure 4B). In general, all modules present rather distributed

spatial patterns, covering a number of brain regions in a non-exclusive manner. In addition, most of the

brain regions in this analysis, although to different extents, participate in more than one module (Figure

4C). Similar to the undirected connectome, the analysis of the directed connectome revealed the

presence of seven main modules (D) (Figure 4E-G), showing a substantial overlap, beside two directed

modules. D2 includes two undirected modules, U4 and U6, and D1 combines U1 with a fraction of U3

(Figure 4D, H). This substantial overlap between the two modularity analyses, directed and undirected

cases, points to the possibility that the two corresponding connectomes generated in this way rely to a

large extent on a common backbone represented by a directed graph and axo-dendrite communication

as reported for other datasets with synapse annotation[21].

*Figure 4. Distribution of the modules in the different regions for the undirected and directed networks. **A,E)** Adjacency matrices recapitulating neuronal connections in the undirected (A) and directed (E) networks sorted by the identified modules. The matrix is binarized for visualization purposes only. **B,F)** Distribution of the modules across the brain regions for the undirected (B) and directed (F) network. Number refers to neurons belonging to a certain module and with the soma located in the regions. Matrix cells are color-coded based on their proportion within the module. U: undirected module, D: directed module. **C,G)** Proportion of the distribution of the modules across the brain regions for the undirected (C) and the directed (G) networks. Labels for the colors are reported in B and F. **D)** Proportion of neurons of the undirected network modules present in the directed network modules. **H)** Number of neurons of the undirected network modules present in the directed network modules. Labels for the colors are reported in D.*

5. Hierarchical anatomical organization of the modules reflects circuit functional role

To better capture the 3D anatomical organization of the 8 identified modules in the undirected graph, we represented the wiring diagrams corresponding to each module in the anatomical reference space (Figure 5). In some of the cases, the identified modules correspond to previously described brain-wide circuits underlying information processing or motor control functions, like the visual network module[19] (Figure 5D and Figure 5F) and the cerebellar network module (Figure 5C). To unravel finer structures in the identified modules, we re-iterated the Leiden algorithm in each main module. This approach, where each module is considered an independent subnetwork, led to the finding of a variable number of subcomponents, typically ranging between five and eight (Figure 5 and Supplementary Figures 2 - 7).

237   These submodule patterns, even with the limitations deriving from the neuronal contact approximation

238   employed here, captured an additional level of the circuit organization, involving features of spatial

239   organization of the neuronal populations across the brain.

Figure 5. Modules of the zebrafish brain. *A-H) Anatomical representation of the eight high-order modules with the corresponding submodules found in the zebrafish dataset overlaid on the brain silhouette. Colors are assigned to the cells according to the soma position in the different brain regions. Scale bar = 100 µm. I) Tree diagram reporting the structure of the zebrafish network with the organization of the submodules for the different modules. L) Distribution of connections between the different modules.*

242   First, we considered the modules including the organization of the Retinal Ganglion Cells, (RGCs),

243   projections to the tectum (modules 4 and 6). The analysis of the relationship of the RGC body position

in the retina and the connected PeriVentricular Neurons (PVNs) in the contralateral tectum showed a retinotopic organization in submodules 4.2 and 4.6 (right tectum, Figure 5A), and in the sub modules 6.1 and 6.2 (left tectum) (Figure 5B). Indeed, these submodules show projections from RGC cells with soma located in the caudal part of the retina to connect with PVNs in the rostral part of the contralateral periventricular region at the level of the tectum and, conversely, for the RGC connections involving the rostral part of the retina and the caudal part of the contralateral retino- recipient area. Similarly, our analysis revealed an articulated spatial pattern of connections involving different brain areas associated with the module 3, representing a pretectal-cerebellar network (Figure 6C). The connections formed between neurons located in the medulla oblongata, cerebellum, and tegmentum (Figure 6C1, submodules 3.3 and 3.4) resemble those associated with vestibulo-ocular reflex (VOR)[32]. In this case, the ascending and ascending/descending tangential neurons are located in the medial octavolateral nucleus (MON) of the medulla (Figure 6C1, submodules 3.3 and 3.4), the recipient of utricular inputs. These neurons project to the oculomotor nucleus (nIII) where extraocular motor neurons mediate gaze stabilization[38]. Furthermore, connections between neurons located in the cerebellum have been shown to be involved in VOR[39,40]. The cerebellum displays a spatially segregated pattern of distinct cells, with lateral Purkinje cells projecting to the MON (Figure 6C2, submodule 3.2) and eurydendroid cells located in the middle with less specific projections[27,41] (Figure 6C2, submodules 3.1 and 3.5). Even though our approach highlighted the existence of principal networks supported by modules of neurons characterized by a relatively higher connection degree, it is clear that the different identified subnetworks show mutual intermodule connection schemes. In order to quantify this aspect, we evaluated the links formed across different modules (see Supplementary Figure 8). This analysis showed a rather heterogeneous scenario, with some modules characterized by a relevant number of intermodular connections, while others with a very limited number of intermodules links.

A Module 4

4.1 4.2 4.3 4.4 4.5 4.6 4.7 4.8

Antero-posterior
Dorso-ventral
Lateral

A1 4.2 4.6

Tectum and tori Retina Hypothalamus
Pretectum Ventral thalamus Eminentia talami
Medulla oblongata Dorsal thalamus Tegmentum
Cerebellum

A2

Antero-posterior axis (um)

4.2
4.6

Retina Optic tectum

B Module 6

6.1 6.2 6.3 6.4 6.5

B1 6.1 6.2

B2

Antero-posterior axis (um)

6.1
6.2

Retina Optic tectum

C Module 3

3.1 3.2 3.3 3.4 3.5

C1 3.3 3.4

C2 3.1 3.2 3.5

Figure 6. Functional description of the modules. A) Anatomical representation of the high-order module U4. Colors are assigned according
to the soma distribution in the brain regions. Scale bar = 100 μm. A1) Sub-modules associated with the right Tectum opticum involving
Peri Ventricular Neurons and Retinal Ganglion Cells, highlighting two different sub-populations of neurons (4.2 caudal tectum, 4.6 medial
tectum). A2) Median of the distributions along the antero-posterior axis of retina and optic tectum neurons of modules 4.2 (magenta) and
4.6 (teal). B) Anatomical representation of the high-order module U6. Colors are assigned according to the soma distribution in the brain
regions. B1) Sub-modules associated with the left Tectum opticum involving PeriVentricular Neurons and Retinal Ganglion Cells,
highlighting two different sub-populations of neurons (6.1 caudal tectum, 6.2 medial tectum). B2) Median of the distributions along the
antero-posterior axis of retina and optic tectum neurons of modules 6.1 (magenta) and 6.2 (teal). C) Anatomical representation of the
high-order module U3 as composed by the five lower-order modules (3.1-3.5). Colors are assigned according to the lower-order modules.
MON: medial octavolateral nucleus, nIII: oculomotor nucleus. C1-C2) Anatomical representation of the individual lower-order modules.
Colors are assigned according to the location of the soma. Scale bar = 100 μm.

CONCLUSIONS

We have devised a method to generate approximated connectomes from datasets of single cell

reconstruction, covering neuronal circuits composed of different neuronal components represented in

terms of single cell morphologies and extending across the complete brain. The proposed approach,

inferring an approximated cell-to-cell contact structure on the basis of a spatial proximity rule, can be

applied to all datasets with single cell reconstructions where synapse direct annotation is not

accessible/available or is not yet complete. This typically is the case of anatomical reconstructions of

sparse cells based on light microscopy[23–27]. Within the characteristic resolution limits dictated by the

optical constraints, using laser scanning or light-sheet microscopy in combination with restricted or

sparse fluorescence labelling, z-stacks of large portions of the brain or even of the complete brain are

acquired to map the neuronal cell interregional connectivity or projection patterns[24,42]. In such

scenarios, even considering a synapse tagging strategy in combination with pre- and post-synaptic

partner labelling[43,44], the synaptic information retrieved could be partial or biased, so overlooking the

involvement of other potential players. On the other hand, considering high-resolution, so-called

"dense" tissue reconstructions based on serial block electron microscopy, the amount of data to

process, even with artificial intelligence approaches, becomes easily relevant even for small brain

sections[30]. While automatically reconstructing cell anatomies and their manual validation could

become possible for small datasets, a complete synapse annotation may still represent a prohibitive

challenge. In this case, one must either focus on a specific circuit or brain region or use the lower-

hanging information available about the cell skeletons to extract approximated information about

brain-wide circuit organization. The two approaches are not mutually exclusive, but rather answer two

different types of questions. The first aims at characterizing the microcircuit structure or the circuit

working motifs[45]. The second, instead, seeks to understand the general organization of brain

networks[46].

We have validated the applicability of our approach by leveraging the synapse-annotated connectome

of the Drosophila melanogaster hemibrain, containing 21,739 connected neurons. Through modularity

analysis, we uncovered a hierarchical organization comprising eight main modules, each with several

submodules. When comparing the adjacency matrices generated from synapse annotations (EM-based)

and those derived using Strahler numbering and proximity range definitions (Strahler-based), we

observed significant correlations with respect two null models where links or weights have been re-

shuffled. Importantly, Adjusted Rand Index (ARI) and Normalized Mutual Information (NMI) scores

confirmed a general overlap between the Strahler-based and EM-based community structures, with

significant p-values supporting the robustness of the approach.

We then have applied our approach to a mesoscale atlas of about 3200 single cell reconstructions

obtained from a vertebrate organism, *Danio rerio,* where the input-output relationships were

estimated based on a proximity rule, and we have analyzed the graph properties of the inferred

connectome. From the corresponding adjacency matrices, we extracted the network communities by

modularity maximization. In this way, we were able to identify, according to a top-down process, those

neuronal modules characterized by a relatively higher degree of connections. With this approach and

its reiterations, we have obtained a decomposition of the connectomes of two phylogenetically distant

organisms on a set of basic components that form a sort of neuronal backbone of the brain wiring

diagram. For most of the identified modules, it is possible to associate a functional role, either based

on known anatomical characterizations of the circuits involved, or on the evidence provided by

functional analyses[27,28,38,41]. In conclusion, these results highlight the capability of our Strahler-based

method to approximate neuronal circuit architectures, demonstrating its potential as a scalable tool for

capturing connectivity patterns in large-scale brain networks. Moreover, the organization of the

modules obtained from the zebrafish larva light microscopy brain dataset shows that the application of

such top-down approach of community detection is able to capture relevant circuit motifs and may be

applied to several datasets at both the circuit and whole-brain scale.

Methods

Dataset and data extraction

We used a whole-brain atlas of the zebrafish larva[27]. The atlas comprises more than 3000 neurons,

approximately 3% of the total neurons in the zebrafish larvae brain, covering almost the entire brain,

anatomically reconstructed by random and sparse Green Fluorescent Protein (GFP) labelling using

BGUG[47] or with other labelling techniques. With a typical resolution of 300x300x1000 nm (X, Y, Z), the

size of the skeletonized dataset was approximately 60 MB. Skeletonized neurons have been

downloaded in the .swc format[48] from the mapzebrain website (https://mapzebrain.org/home).

Computing the Strahler ordering for the neuronal anatomy reconstructions

For a given neuronal skeleton i from the original dataset of 3163 reconstructions, we imported the

.swc file and used the "networkx" library (https://networkx.org/)[49] to convert the corresponding

collection of ordinated endpoints to a tree-like directed graph representing its anatomical structure.

The soma is considered as a root and the connected endpoints as children. In order to build an

approximated connectome from this dataset, we first classified each skeleton endpoint within its

branching hierarchy using Strahler ordering[28]. In this representation, terminal endpoints, those without

branching children, obtain a value one. For the other endpoints in the .swc file, the integer value of the

Strahler progressively increases with the number of upstream branches along the anatomical structure

to reach the maximum at the level of the soma. For the calculation of the putative synaptic contacts

and hence for the calculation of the connectome, we considered only endpoints with unitary Strahler

numbers (below referred to as endpoint unless otherwise specified), which we denoted as E_{μ}^{i}

where the italic superscript refers to i-th neuron within the dataset and the Greek subscript stands for

the μ-th endpoint of unitary Strahler number in the .swc file of i-th neuron.

Comparison with the ground truth in Drosophila EM

We considered the EM annotated connectome of the Drosophila hemibrain as ground-truth and

compared how close the reconstruction results when we applied on the same dataset our simplified

method based on Strahler numbering and proximity range (SP method). For that, we identified eight

modules using a community discovery approach on the EM connectome and considered the resulting

partitions in submodules, $P^{EM} = \{P_i^{EM} | i \in (1,..,8)\}$ where $P_i^{EM} = \{P_{i,j}^{EM} | j \in (1,..,..)\}$ represents the

i-module and $P_{i,j}^{EM}$ the j-sub-module associated with the module. Each P_i^{EM} was considered for

evaluating the sub-module partitions obtained based on the EM annotation ($P_{i,j}^{EM}$) or retrieved with the

approximated approach, $P_i^{EM} = \{P_{i,q}^{SP}|q \in (1,..,..)\}$, where $P_{i,q}^{SP}$ is the q-sub-module obtained with

the SP method in association with the reference module P_i^{EM}. For each module P_i^{EM}, we calculated the

corresponding ARI_i according to the following equation, as reported[31,50].

$$ARI_i\left(P_{i,j}^{EM}, P_{i,q}^{SP}\right) = \frac{\sum_{j,q}\binom{n_{jq}}{2} - [\sum_j\binom{a_j}{2}\sum_q\binom{b_q}{2}]/\binom{n}{2}}{\frac{1}{2}[\sum_j\binom{a_j}{2} + \sum_q\binom{b_q}{2}] - [\sum_j\binom{a_j}{2}\sum_q\binom{b_q}{2}]/\binom{n}{2}}$$

364 where n_{jq} represents the entries of the contingency matrix associated with the i-module, i.e. the

365 number of neurons that are belonging to $P_{i,j}^{EM} \cap P_{i,q}^{SP}$, a_j and b_q are the sum of n_{jq} along q and j indices

366 respectively, and n is the total number of partition elements. As for the NMI, we started by calculating

367 the probability $\pi_{jq} = \frac{n_{jq}}{n}$, as the probability to belong simultaneously to $P_{i,j}^{EM}$ and $P_{i,q}^{SP}$, and the marginal

368 distributions computed as $\pi_{j*} = \sum_q \pi_{jq}$ and $\pi_{*q} = \sum_j \pi_{jq}$. We could then calculate the NMI as:

$$NMI_i\left(P_{i,j}^{EM}, P_{i,q}^{SP}\right) = \frac{-2\sum_{j,q}\pi_{jq}\log(\frac{\pi_{jq}}{\pi_{j*}\pi_{*q}})}{\sum_j \pi_{j*}\log(\pi_{j*}) + \sum_q \pi_{*q}\log(\pi_{*q})}$$

Validation of ARI and NMI scores using Random null models

For validating the NMI and ARI parameters, we constructed 100 random networks for each of the

mentioned 8 networks obtained with the SP method using two different approaches. In a first approach,

called "Link randomization or directed configuration", we used directed_edge_swap function from the

Python NetworkX library to swap the edges of the graph while keeping the node degree fixed. In a

second approach, called Weight Randomization, we kept the original links between the nodes

unchanged but changed the weight of the connection by randomly sampling from the original weight

distribution. The z-score was calculated for ARI and NMI considering two different proximity range, 1

and 5 μm, and the two types of random models as the difference of the between the actual value with

respect to the average value obtained from the 100 null-model networks, divided by the corresponding

standard deviation.

Zebrafish undirected network reconstruction

We assume that all endpoints between two neurons form connections with each other. We introduce

a proximity range u that represents the maximal effective distance at which two endpoints can form a

connection. We set $u = 5\,\mu m$ based on the typical reconstruction error in optical microscopy[27,51,52].

Then, a link between two neurons is present if at least two of their endpoints are at a distance smaller

than u. That is, given two neurons i, j and their endpoints $\{E_\mu^{(i)}\}$ and $\{E_\nu^{(j)}\}$, we compute

the Euclidean distance matrix $d_{\mu\nu}^{(ij)} = d(E_\mu^{(i)}, E_\nu^{(j)})$, for all $\mu = 1, \dots, N_i$

and $\nu = 1, \dots, N_j$. The elements of the resulting adjacency matrix A_{ij}^U are the number of

entries of $d_{\mu\nu}^{(ij)}$ such that $d_{\mu\nu}^{(ij)} \leq u$. Since $d_{\mu\nu}^{(ij)}$ is

symmetric in the i, j indexes, A_{ij}^U is symmetric as well.

Zebrafish directed network reconstruction

The morphological data we considered do not contain information between the dendritic and axonal

endpoints. Therefore, given the endpoints $\{E_\mu^{(i)}\}$ of a neuron, we computed the radial

distance $r_\mu^{(i)}$ between each endpoint $E_\mu^{(i)}$ and the soma. The associated

probability distribution $p_i(r)$ resulted often bimodal, pointing to the fact that a large number of the

endpoints are concentrated either close to the soma or far from it. We introduce a threshold T which

approximately corresponds to the antimode of $p_i(r)$, i.e., the local minimum between the two modes.

If $r_\mu^{(i)} < T$, we assume that the corresponding synaptic endpoint is a dendritic endpoint for the neuron

i, whereas $E_\mu^{(i)}$ is an axonal endpoint if $r_\mu^{(i)} \geq T$.

Then, the connections from the i-th neuron to the j-th one are obtained from the distances $d_{\mu\nu}^{(ij)} < u$,

similarly to the undirected reconstruction. However, we now impose that a connection is counted only

for those elements of $d_{\mu\nu}^{(ij)}$ such that μ is an axonal endpoint for neuron i, and ν is a dendritic endpoint

for neuron j. Clearly, the resulting adjacency matrix A_{ij}^D is not symmetric, since an axonal endpoint

connecting to a dendritic endpoint does not imply that the reverse connection is present.

Network properties

The analysis of network parameters[53] was performed using dedicated Python libraries. The following

functions from the 'networkx'[49] (https://networkx.org/) library have been used:

'connected_components' to identify the network connected components,

'average_shortest_path_length' to compute the average shortest path length, 'density' to compute the

density of the network, 'average_clustering' to compute the average clustering coefficient and

'degree_assortativity_coefficient' to compute assortativity correlation coefficient.

Leiden algorithm for neuronal module detection

We adopted Leiden algorithm[41] to extract neuronal modules. The analysis of network parameters was

performed in Python using the 'leidenalg' package[54]

(https://leidenalg.readthedocs.io/en/stable/index.html) .

The Leiden community detection algorithm iteratively refines an initial partition of nodes in a network

to optimize a quality function called modularity Q. It has three main steps, which are fast local

movement of nodes to communities, refinement of partition and lastly aggregation.

1. **Initialization**. The algorithm starts with a singleton partition which means each node is
    considered as its own community. The quality function Q is computed using the formulas below.
    We have chosen the well-known quality function for the undirected case[55]

    $$Q_{undir} = \frac{1}{2m} \sum_{ij} \left(A_{ij} - \gamma \frac{k_i k_j}{2m} \right) \delta(\sigma_i, \sigma_j)$$

    and for the directed case[56]

    $$Q_{dir} = \frac{1}{m} \sum_{ij} \left(A_{ij} - \gamma \frac{k_i^{out} k_j^{in}}{m} \right) \delta(\sigma_i, \sigma_j)$$

    , where m is total number of edges or total edge weight, A is the corresponding adjacency
    matrix for directed or undirected graph, $\gamma > 0$ is the linear resolution parameter[57] and is
    assumed equal to 1 in our analysis. Higher value of this parameter leads to more communities
    k_i and k_j are the weighted node degree of node i node j for the undirected case while for
    the directed it is mentioned in superscript to specify whether it is weighted out-degree
    $k_i^{out} = \sum_j A_{ij}$ or weighted in-degree $k_j^{in} = \sum_i A_{ij}$ and finally σ_i
    denotes the community of node i with δ representing the Kronecker delta.

2. **Fast local movement of nodes.** A set of all nodes is created. The following steps are repeated until there are no further nodes in a list of nodes: (i) randomly select and remove a node from this set; (ii) for each community, calculate the gain in modularity as if the node were to move to that community; (iii) consider the move that results in the largest modularity gain, or no move if the modularity cannot be improved; (iv) update the partition by moving the node to the selected community; (v) all the neighboring nodes of the selected node which are not in the new community are added to the set of nodes. The steps are repeated until there is no node in the set of nodes.

3. **Refinement phase.** The partition P obtained from the fast local moving phase is refined as follows. Let the refined partition P* be initialized to a singleton partition, where each node represents a community. Randomly select one of the singleton communities in P*, and merge it to another community C* from P* with a probability that depends on the increase of the quality function. The higher the increase, the higher the probability of the merging. Repeat till all singleton community of P* are merged. Iterate these steps over all the communities of C belonging to P and obtain a new refined partition P*.

4. **Aggregation phase.** In this phase, each community in P* is considered as a node, while an edge between such communities is present if any of their nodes are connected. In this aggregated graph, communities are formed by merging all the aggregated nodes which are also subsets of the same community in the original partition.

5. **Convergence.** Return to step 2 of fast local moving of nodes and then terminate if there is no further change in the quality function.

Code and data availability

The scripts and data used to produce the results of the manuscript are available at https://github.com/dalMaschio-lab/Connectome_analysis.

Funding

The funding was provided by Horizon 2020 Framework Programme (734862) to M.D.M.

Contributions

K.K.H.M. : conceptualization and data analysis.

M.B.: conceptualization, data analysis, anatomical analysis, supervision.

G.N. : conceptualization, data analysis, supervision.

S.S.: conceptualization and supervision.

M.D.M: conceptualization, supervision and funding acquisition.

All the authors participated in the writing, review and editing of the manuscript.

Acknowledgments

We thank Delfina Iriarte, Physics department of University of Padua and Nora Nikoloska, Mathematics
Department of University of Padua for their support in preliminary analyses; we thank Prof. Tomaso
Erseghe from the Information Engineering Department of University of Padua for useful discussion.

Supplementary Figure 1. Distribution of endpoints of neurons. A) Typically, the distance between the endpoints of a neuron and the soma is distributed accordingly to two main distribution classes, following the Galtung classification[40]. **B**) Bimodal class S. The antimode of the distribution is used to set a threshold (dashed line) to distinguish between dendrites and axon. C) Bimodal class U. The antimode of the distribution is used to set a threshold (dashed line) to distinguish between dendrites and axon.

Supplementary Figure 2. The zebrafish module 1. A) Anatomical representation of the high-order module U1. Colors are assigned according to the soma distribution in the brain regions. Scale bar = 100 μm. **B)** Anatomical representation of the individual lower-order modules. Colors are assigned according to the location of the soma. Scale bar = 100 μm.

Supplementary Figure 3. The zebrafish module 2. A) Anatomical representation of the high-order module U2. Colors are assigned according to the soma distribution in the brain regions. Scale bar = 100 µm. **B)** Anatomical representation of the individual lower-order modules. Colors are assigned according to the location of the soma. Scale bar = 100 µm.

Supplementary Figure 4. The zebrafish module 4. A) Anatomical representation of the high-order module U4. Colors are assigned according to the soma distribution in the brain regions. Scale bar = 100 μm. **B)** Anatomical representation of the individual lower-order modules. Colors are assigned according to the location of the soma. Scale bar = 100 μm.

Supplementary Figure 5. The zebrafish module 5. A) Anatomical representation of the high-order module U5. Colors are assigned according to the soma distribution in the brain regions. Scale bar = 100 μm. **B)** Anatomical representation of the individual lower-order modules. Colors are assigned according to the location of the soma. Scale bar = 100 μm.

Supplementary Figure 6. The zebrafish module 6. A) Anatomical representation of the high-order module U6.
Colors are assigned according to the soma distribution in the brain regions. Scale bar = 100 μm. **B)** Anatomical
representation of the individual lower-order modules. Colors are assigned according to the location of the
soma. Scale bar = 100 μm.

Supplementary Figure 7. The zebrafish module 7. A) Anatomical representation of the high-order module U7.
Colors are assigned according to the soma distribution in the brain regions. Scale bar = 100 μm. **B)** Anatomical
representation of the individual lower-order modules. Colors are assigned according to the location of the
soma. Scale bar = 100 μm.

Supplementary Figure 8. Relationships between zebrafish modules. Distribution of connections between the
modules of the directed network. D: directed modules.

Supplementary Information

Validation of ARI and NMI scores using Random null models, details

526 We considered the 8 networks from the SP method and let's represent them as $\{M_i^{SP} : 1 \le i \le 8\}$.
527 Before proceeding to compute the z-scores, we constructed 100 random networks for each of the
528 mentioned 8 networks of SP method using two unique random null models:

1. **Link randomization or directed configuration**[58]**:** In this method, we use *directed_edge_swap()* function
of NetworkX library of python to Swap three edges in a directed graph while keeping the node degrees
fixed. This method considers randomly chosen three connected edges such that $a \to b \to c \to d$ and
after the swap is applied it becomes $a \to c \to b \to d$. In this way, we reach all possible states with the
same in- and out-degree distribution in a directed graph and finally giving us directed configuration
random null model.

2. **Weight randomization:** In this method, while keeping the links between the nodes unchanged or
unswapped, we assign weight to each edge by sampling from weight distribution of all existing edge
weights.

We computed the z-score for ARI and NMI for the 8 networks reconstructed with 1um proximity
range and 5 μm proximity range using SP method. The mathematical formulation is as follows:

$$\text{z-score}(ARI, M_i, PR, \text{randomization type}) = \frac{ARI(M_i, PR) - \mu_{100}(M_i, PR, \text{randomization type})}{\sigma_{100}(M_i, PR, \text{randomization type})} \qquad [1]$$

where the notations refer to:

• M_i - refers to the module(set of nodes belonging to community i) for which we compute the similarity
measures $ARI\left(P_i^{EM}, P_i^{SP}\right)$ or $NMI\left(P_i^{EM}, P_i^{SP}\right)$.

• PR - refers to proximity range which is either 1 µm or 5 µm used for reconstructing the network M_i^{SP}
using SP method.

• randomization type - refers to the type of random null model used to randomize the network M_i^{SP} i.e.
directed configuration model or weight randomization.

• μ_{100} and σ_{100} are the mean and standard deviations over ARI or NMI computed on 100 random
networks. Specifically, it is $ARI\left(P_i^{EM}, P_i^{SP,rand}\right)$ or $NMI\left(P_i^{EM}, P_i^{SP}\right)$

The above formula similarly can be written by replacing it with NMI measure.

p-values computation

We have computed the p-values for the z scores(Eq. [1]) obtained for ARI or NMI in two ways:

$$\text{p-value}(M_i) = 1 - \Phi(\text{z-score}) \qquad [2]$$

where $\Phi(\ )$ refers to cumulative distribution function of standard normal distribution.

$$\text{p-value}(M_i) = \frac{\sum_j 1_{\text{z-score}(rand_j) > \text{z-score}(M_i)} : 1 \leq j \leq 100}{\text{No. of realizations of random networks(i.e. 100)}} \qquad [3]$$

where $1 \leq i \leq 8$.

REFERENCES

1.   Brunner, C. *et al.* Whole-brain functional ultrasound imaging in awake head-fixed mice. *Nat Protoc* **16**,

560   3547–3571 (2021).

561   2.   Ahrens, M. B., Orger, M. B., Robson, D. N., Li, J. M. & Keller, P. J. Whole-brain functional imaging at cellular

562   resolution using light-sheet microscopy. *Nat Methods* **10**, 413–420 (2013).

563   3.   Bruzzone, M. *et al.* Whole brain functional recordings at cellular resolution in zebrafish larvae with 3D

564   scanning multiphoton microscopy. *Sci Rep* **11**, 11048 (2021).

565   4.   Schrödel, T., Prevedel, R., Aumayr, K., Zimmer, M. & Vaziri, A. Brain-wide 3D imaging of neuronal activity in

566   Caenorhabditis elegans with sculpted light. *Nat Methods* **10**, 1013–1020 (2013).

567   5.   Brondi, M., Bruzzone, M., Lodovichi, C. & Dal Maschio, M. Optogenetic Methods to Investigate Brain

568   Alterations in Preclinical Models. *Cells* **11**, 1848 (2022).

569   6.   Ahrens, M. B. *et al.* Brain-wide neuronal dynamics during motor adaptation in zebrafish. *Nature* **485**, 471–

570   477 (2012).

571   7.   Chen, X. *et al.* Brain-wide Organization of Neuronal Activity and Convergent Sensorimotor Transformations

572   in Larval Zebrafish. *Neuron* **100**, 876-890.e5 (2018).

573   8.   Ahrens, M. B. & Engert, F. Large-scale imaging in small brains. *Current Opinion in Neurobiology* **32**, 78–86

574   (2015).

575   9.   Naumann, E. A. *et al.* From Whole-Brain Data to Functional Circuit Models: The Zebrafish Optomotor

576   Response. *Cell* **167**, 947-960.e20 (2016).

577   10.   Mariani, B. *et al.* Disentangling the critical signatures of neural activity. *Sci Rep* **12**, 10770 (2022).

578   11.   Barzon, G., Nicoletti, G., Mariani, B., Formentin, M. & Suweis, S. Criticality and network structure drive

579   emergent oscillations in a stochastic whole-brain model. *J. Phys. Complex.* **3**, 025010 (2022).

580   12.   Panzeri, S., Harvey, C. D., Piasini, E., Latham, P. E. & Fellin, T. Cracking the Neural Code for Sensory

581   Perception by Combining Statistics, Intervention, and Behavior. *Neuron* **93**, 491–507 (2017).

13. Schaeffer, R., Khona, M. & Fiete, I. R. *No Free Lunch from Deep Learning in Neuroscience: A Case Study*

through Models of the Entorhinal-Hippocampal Circuit.

http://biorxiv.org/lookup/doi/10.1101/2022.08.07.503109 (2022) doi:10.1101/2022.08.07.503109.

14. Wanner, A. A., Genoud, C., Masudi, T., Siksou, L. & Friedrich, R. W. Dense EM-based reconstruction of the

interglomerular projectome in the zebrafish olfactory bulb. *Nat Neurosci* **19**, 816–825 (2016).

15. Svara, F. *et al.* Automated synapse-level reconstruction of neural circuits in the larval zebrafish brain. *Nat*

Methods **19**, 1357–1366 (2022).

16. White, J. G., Southgate, E., Thomson, J. N. & Brenner, S. The structure of the nervous system of the

nematode Caenorhabditis elegans. *Philos Trans R Soc Lond B Biol Sci* **314**, 1–340 (1986).

17. Takemura, S. *et al.* The comprehensive connectome of a neural substrate for 'ON' motion detection in

Drosophila. *eLife* **6**, e24394 (2017).

18. Zheng, Z. *et al.* A Complete Electron Microscopy Volume of the Brain of Adult Drosophila melanogaster.

Cell **174**, 730-743.e22 (2018).

19. Schubert, P. J. *et al.* SyConn2: dense synaptic connectivity inference for volume electron microscopy. *Nat*

Methods **19**, 1367–1370 (2022).

20. Ryan, K., Lu, Z. & Meinertzhagen, I. A. The CNS connectome of a tadpole larva of Ciona intestinalis (L.)

highlights sidedness in the brain of a chordate sibling. *eLife* **5**, e16962 (2016).

21. Winding, M. *et al.* The connectome of an insect brain. *Science* **379**, eadd9330 (2023).

22. Dorkenwald, S. *et al.* Neuronal wiring diagram of an adult brain. *Nature* **634**, 124–138 (2024).

23. Arganda-Carreras, I. *et al.* A Statistically Representative Atlas for Mapping Neuronal Circuits in the

Drosophila Adult Brain. *Front Neuroinform* **12**, 13 (2018).

24. Oh, S. W. *et al.* A mesoscale connectome of the mouse brain. *Nature* **508**, 207–214 (2014).

25. Jenett, A. *et al.* A GAL4-driver line resource for Drosophila neurobiology. *Cell Rep* **2**, 991–1001 (2012).

26. Bogovic, J. A. *et al.* An unbiased template of the Drosophila brain and ventral nerve cord. *PLOS ONE* **15**,

e0236495 (2020).

27. Kunst, M. *et al.* A Cellular-Resolution Atlas of the Larval Zebrafish Brain. *Neuron* **103**, 21-38.e5 (2019).

28. Helmbrecht, T. O., dal Maschio, M., Donovan, J. C., Koutsouli, S. & Baier, H. Topography of a Visuomotor

Transformation. *Neuron* **100**, 1429-1445.e4 (2018).

29. Galtung, J. *Theory and Methods of Social Research*. (Oslo, Universitetsforlaget; New York, Columbia

University Press [etc.], 1967).

30. Scheffer, L. K. *et al.* A connectome and analysis of the adult Drosophila central brain. *Elife* **9**, e57443

(2020).

31. Hubert, L. & Arabie, P. Comparing partitions. *Journal of Classification* **2**, 193–218 (1985).

32. Rand, W. M. Objective Criteria for the Evaluation of Clustering Methods. *Journal of the American Statistical*

Association **66**, 846–850 (1971).

33. Vinh, N. X., Epps, J. & Bailey, J. Information Theoretic Measures for Clusterings Comparison: Variants,

Properties, Normalization and Correction for Chance. *Journal of Machine Learning Research* **11**, 2837–2854

(2010).

34. McDaid, A. F., Greene, D. & Hurley, N. Normalized Mutual Information to evaluate overlapping community

finding algorithms. Preprint at https://doi.org/10.48550/arXiv.1110.2515 (2013).

35. Lancichinetti, A., Fortunato, S. & Kertész, J. Detecting the overlapping and hierarchical community

structure in complex networks. *New J. Phys.* **11**, 033015 (2009).

36. Suárez, L. E. *et al.* Connectome-based reservoir computing with the conn2res toolbox. *Nat Commun* **15**,

656 (2024).

37. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected

communities. *Sci Rep* **9**, 5233 (2019).

38. Bianco, I. H. *et al.* The Tangential Nucleus Controls a Gravito-inertial Vestibulo-ocular Reflex. *Current*

Biology **22**, 1285–1295 (2012).

39. Matsui, H., Namikawa, K., Babaryka, A. & Köster, R. W. Functional regionalization of the teleost cerebellum

analyzed in vivo. *Proceedings of the National Academy of Sciences* **111**, 11846–11851 (2014).

632 40. Favre-Bulle, I. A., Vanwalleghem, G., Taylor, M. A., Rubinsztein-Dunlop, H. & Scott, E. K. Cellular-Resolution

633   Imaging of Vestibular Processing across the Larval Zebrafish Brain. *Current Biology* **28**, 3711-3722.e3

634   (2018).

635 41. Bae, Y.-K. *et al.* Anatomy of zebrafish cerebellum and screen for mutations affecting its development. *Dev*

636   *Biol* **330**, 406–426 (2009).

637 42. Yao, S. *et al.* A whole-brain monosynaptic input connectome to neuron classes in mouse visual cortex. *Nat*

638   *Neurosci* **26**, 350–364 (2023).

639 43. Kim, J. *et al.* mGRASP enables mapping mammalian synaptic connectivity with light microscopy. *Nat*

640   *Methods* **9**, 96–102 (2011).

641 44. Shearin, H. K., Quinn, C. D., Mackin, R. D., Macdonald, I. S. & Stowers, R. S. t-GRASP, a targeted GRASP for

642   assessing neuronal connectivity. *J Neurosci Methods* **306**, 94–102 (2018).

643 45. Eschbach, C. *et al.* Multilevel feedback architecture for adaptive regulation of learning in the insect brain.

644   649731 Preprint at https://doi.org/10.1101/649731 (2019).

645 46. van der Plas, T. L. *et al.* Neural assemblies uncovered by generative modeling explain whole-brain activity

646   statistics and reflect structural connectivity. *Elife* **12**, e83139 (2023).

647 47. Scott, E. K. *et al.* Targeting neural circuitry in zebrafish using GAL4 enhancer trapping. *Nat Methods* **4**, 323–

648   326 (2007).

649 48. Cannon, R. C., Turner, D. A., Pyapali, G. K. & Wheal, H. V. An on-line archive of reconstructed hippocampal

650   neurons. *Journal of Neuroscience Methods* **84**, 49–54 (1998).

651 49. Hagberg, A. A., Schult, D. A. & Swart, P. J. Exploring Network Structure, Dynamics, and Function using

652   NetworkX. (2008).

653 50. Labatut, V. Generalized Measures for the Evaluation of Community Detection Methods. Preprint at

654   https://doi.org/10.48550/arXiv.1303.5441 (2016).

655 51. Vishwanathan, A. *et al.* Predicting modular functions and neural coding of behavior from a synaptic wiring

656   diagram. 2020.10.28.359620 Preprint at https://doi.org/10.1101/2020.10.28.359620 (2022).

52. Stepanyants, A. & Chklovskii, D. B. Neurogeometry and potential synaptic connectivity. *Trends Neurosci* **28**,

387–394 (2005).

53. Preface. in *Networks: An Introduction* (ed. Newman, M.) 0 (Oxford University Press, 2010).

doi:10.1093/acprof:oso/9780199206650.002.0005.

54. Traag, V. *et al.* vtraag/leidenalg: 0.10.0. Zenodo https://doi.org/10.5281/zenodo.8147844 (2023).

55. Newman, M. E. J. & Girvan, M. Finding and evaluating community structure in networks. *Phys. Rev. E* **69**,

026113 (2004).

56. Leicht, E. A. & Newman, M. E. J. Community Structure in Directed Networks. *Phys. Rev. Lett.* **100**, 118703

(2008).

57. Reichardt, J. & Bornholdt, S. Statistical mechanics of community detection. *Phys. Rev. E* **74**, 016110 (2006).

58. Erdős, P. L., Miklós, I. & Toroczkai, Z. A simple Havel-Hakimi type algorithm to realize graphical degree

sequences of directed graphs. Preprint at https://doi.org/10.48550/arXiv.0905.4913 (2010).

669