Frequency-and circuit-specific effects of septohippocampal deep brain stimulation in mice

as measured by functional ultrasound imaging.

Authors

Lindsey M. Crown, PhD[*2], Kofi Agyeman, MS[*5,7], Wooseong Choi, BS[*1], Nancy Zepeda, BS[1],

Ege Iseri MS [5], Pooyan Pahlavan MS[5], Steven J. Siegel, MD, PhD[2], Charles Liu MD, PhD[§1,3,4,6],

Vasileios Christopoulos, PhD[§3,5,7], Darrin J. Lee, MD, PhD[§1,2,3,4,6]

Affiliations

[1]Department of Neurological Surgery, Keck School of Medicine, University of Southern

California, Los Angeles, CA, USA

[2]Department of Psychiatry and Behavioral Sciences, Keck School of Medicine, University of

Southern California, Los Angeles, CA, 90033, USA

[3]Neurorestoration Center, Keck School of Medicine, University of Southern California, Los

Angeles, CA, USA

[4]Viterbi School of Engineering, University of Southern California, Los Angeles, CA, USA

[5]Department of Bioengineering, University of California Riverside, Riverside, CA, USA

[6]Rancho Los Amigos National Rehabilitation Center, Downey, CA, USA

[7]Neuroscience Graduate Program, University of California Riverside, Riverside, CA, USA

[*] These authors contributed equally

[§] These authors jointly supervised this work (Correspondence)

Please send correspondence to: Dr. Darrin J. Lee, MD, PhD (Darrin.Lee@med.usc.edu)

Abstract

Background

Deep brain stimulation (DBS) has shown remarkable success in treating neurological and

psychiatric disorders such as Parkinson's disease, dystonia, epilepsy, and obsessive-compulsive

disorder. Despite this success, the underlying mechanism of action remains unknown. DBS is

now being explored to improve functional outcomes in other psychiatric conditions, such as

those characterized by reduced N-methyl-D-aspartate (NMDA) function (i.e. schizophrenia).

While DBS for movement disorders requires high-frequency continuous stimulation, there is

evidence that intermittent low-frequency stimulation in neuropsychiatric conditions may have

persisting cognitive benefits, necessitating a broader exploration of how DBS alters brain

networks.

Objective

We characterize the effects of pharmacologic NMDA antagonism on the septohippocampal

network and the impact of high- and low-frequency MSN DBS on cerebral blood volume (CBV)

in brain structures within and outside of the septohippocampal network.

Methods

In this study, we utilize a novel technology, functional ultrasound imaging (fUSI), to characterize

the cerebrovascular impact of medial septal nucleus (MSN) DBS under conditions of NMDA

antagonism (pharmacologically using Dizocilpine [MK-801]) in anesthetized male mice.

Results

Imaging from a sagittal plane across a variety of brain regions, we find that MSN theta-

frequency (7.7Hz) DBS has a larger effect on hippocampal CBV after stimulation offset. This is

observed following an intraperitoneal (i.p.) injection of either saline vehicle or MK-801 (1

mg/kg). This effect is not present using standard high-frequency DBS stimulation parameters

(i.e. gamma [100Hz]).

<u>Conclusion</u>

These results indicate the MSN DBS increases circuit-specific hippocampal neurovascular

activity in a frequency-dependent manner that continues beyond the period of electrical

stimulation.

Introduction

There is growing interest in utilizing neuromodulation to treat cognitive impairment associated with neurological and psychiatric disorders. Many of these disorders involve aberrant electrophysiology and cerebral blood perfusion [1,2]. Recent evidence demonstrates that electrical neuromodulation can improve functional outcomes [3,4]. Particularly, deep brain stimulation (DBS) has shown remarkable success in treating neurological diseases such as movement disorders and epilepsy, and there is increasing evidence for its efficacy in cognitive and psychiatric conditions.

Despite this growing utility, the underlying mechanism of DBS for cognitive outcomes remains largely unknown. Pre-clinical studies have been hindered by technological limitations, such as the inability to record electrical brain activity during stimulation and the low spatial resolution of electrographic measures. Functional ultrasound imaging (fUSI) is a relatively new technology that enables large-scale estimates of neural activity through measures of cerebral blood volume (CBV). fUSI provides a unique combination of high spatiotemporal resolution (~100 μm^3, up to 10 ms) and high sensitivity to slow blood flow (~1 mm/s velocity) across a large field of view. In fact, fUSI has already been proven to be an effective tool for imaging large-scale brain activity and pharmacodynamics [5–8]. As such, it is well-positioned to improve our understanding of the impact of DBS on large-scale brain networks during and immediately following stimulation.

In many disorders, cognitive dysfunction is accompanied by altered septohippocampal network activity. For instance, neural oscillatory patterns within the septohippocampal network, such as gamma- and theta-band activity, are characteristically altered in disorders such as schizophrenia and Alzheimer's disease and are often correlated with impaired memory [1,9–11].

Alterations in specific neurotransmitter signal transduction pathways also play an important role

in modulating neural oscillatory activity. In particular, the glutamatergic *N*-methyl-D-aspartate

(NMDA) receptor is crucial to regulating hippocampal theta and gamma oscillations and is the

predominant molecular control for synaptic plasticity and memory function [12–14] As such,

pharmacologic NMDA receptor antagonism (i.e. via MK-801) results in characteristic changes to

neural oscillatory patterns and memory dysfunction [15–19].

The medial septal nucleus (MSN) is a key structure in the septohippocampal network that

modulates sensory-motor processing and acts as a "pacemaker" for hippocampal theta

oscillations via dense glutamatergic, cholinergic, and GABAergic projections to the

hippocampus [20,21]. This makes the MSN a promising target for DBS in cognitive disorders

involving memory impairments [22]. Our recent work as well as that of others suggest that

modulating the septohippocampal network via MSN theta frequency-specific (7.7 Hz) DBS can

restore cognitive impairment and memory dysfunction in preclinical models of epilepsy,

traumatic brain injury, Alzheimer's disease, and schizophrenia [22–27].

In the current study, we utilize fUSI to characterize the effects of MK-801 on CBV in the

septohippocampal network (Fig. 1A) including the hippocampus, and medial prefrontal cortex

(mPFC). Within the same sagittal plane, we also image CBV changes (ΔCBV) to regions of

interest (ROIs) outside this network including the striatum, thalamus, hypothalamus, and

pallidum. Note that other regions that are connected with the MSN, such as amygdala, habenula,

or raphe nucleus were not recorded, since they were not accessible from the selected 2D image

plane. Additionally, we determine the effect of direct MSN stimulation on blood flow in these

areas using two distinct frequencies, theta (7.7Hz) and gamma (100Hz). Finally, we test the

hypothesis that MSN theta-frequency-specific stimulation can improve blood flow under

conditions of reduced NMDA function.

<u>Materials and Methods</u>

Animals and surgical procedures

82 male 8–12-week-old C57BL/6 mice (Charles River Laboratories; Hollister, CA) were

used in this study. fUSI data from two animals were excluded due to extreme values (Grubbs

test for outliers, 98[th] percentile of all maximum change in pD intensities) that did not appear

physiological [28,29]. The animals were divided into two main groups: saline vehicle control

(n=46) and MK-801 drug-administered (n=34). Each group was then sub-divided into three

categories: no stimulation (saline: n=14; MK-801: n=10), theta stimulation (saline: n=16, MK-

801: n=12), and gamma stimulation (saline: n=14, MK-801: n=14).

Mice were anaesthetized with 5% isoflurane in O_2/N_2O (1:2) carrier gas and then

maintained at a constant rate (1.5- 2%) through surgery and data acquisition. Body temperature

was kept constant throughout recordings by placing animals on an electric heating pad. Hair was

removed from the mouse's head for fUSI using a commercially available depilatory cream (Nair,

Pharmapacks).

To implant DBS electrodes, mice were head-fixed in a stereotaxic frame (David Kopf

instruments, Tujunga, CA) and a midline incision of the scalp was made to expose the skull. A

2mm burr hole was then drilled to implant bipolar stimulating electrodes (E363T/2/SPC ELEC

.008"/.2MM, Plastics One Inc., Roanoke, VA) targeting the midline MSN (AP: +0.7mm, ML: -

0.9mm, from bregma. Z: -4.39mm at 11.8 degrees) from the left hemisphere. Prior to

implantation, the electrodes were connected to an electronic interface board (Neuralynx Inc.,

Bozeman, MT) and bent at 4.5mm from the tip to maximize the proximity of the fUSI probe to

the skull (Fig. 1B). Recordings took place over one hour, animals were injected after 5 minutes

with saline vehicle or MK-801 (1mg/kg) and stimulation or sham stimulation was delivered for 5

minutes 45 minutes into the recording period (Fig. 1C). All procedures were approved by the

University of Southern California, Institutional Animal Care and Use Committee (IACUC

#21006).

Figure 1. Experimental setup and fUSI recording protocol. A) *Schematic illustration of connectivity between the MSN and ROIs. Arrowheads represent axonal projections to and/or from MSN.* ***B)*** *Experimental set-up showing the anesthetized mouse in a stereotaxic frame under the Iconeous One motorized probe mount. DBS stimulating electrodes were implanted on the left hemisphere and a sagittal plane of the right hemisphere was imaged.* ***C)*** *Diagram of the protocol for 60 minutes of continuous fUSI acquisition consisting of saline or 1.0 mg/kg MK-*

801 drug injection (at the 5-minute mark) and 5 minutes of gamma- or theta-frequency DBS (at

the 45-minute mark).

Figure 2. Histological mapping of electrode placement in the MSN. *A) Representative Nissl*

*stain of the mouse brain with a blood mark indicating the electrode placement **B**) Annotation*

of the MSN from the Allen Reference Atlas – Mouse Brain in the same slice position as A.

Histology

Mice were euthanized immediately after the fUSI recording by isoflurane overdose followed by

transcardial perfusion using 50 mL of 0.1M sodium phosphate buffer saline (PBS) and 50 mL of

4% paraformaldehyde. Brains were harvested and stored in phosphate buffered saline at 4°C. To

confirm electrode positioning within the MSN, coronal sections were cut at 100µm thickness

with a vibratome (Leica VT 1200; Leica Biosystems, Buffalo Grove, IL) and then Nissl stained

with Cresyl Violet (Fig. 2A,B).

Computational Modeling of the Induced Voltages: Volume of Tissue Activation

Our group has developed a computational analysis method for applications in electrophysiology

and bioelectromagnetic interactions, namely the Admittance Method (AM)[32]. AM discretizes a

bulk tissue model into cuboid voxels, where each voxel is represented by an equivalent circuit of

lumped passive elements. The admittance value at each voxel is calculated using the material

properties such as conductivity and permittivity. A set of linear equations using iterative methods

are used to calculate the induced voltage at each node of the voxelized network. Here, a 3-D bulk

tissue model with the desired electrode configuration was constructed, a current-controlled

stimulation pulse was applied through the electrodes and the induced voltages at each voxel were

calculated. A mouse head model (Fig. 3A) was used for this study where three types of tissue were

considered (grey matter, white matter (WM) and cerebrospinal fluid (CSF). Electrodes were set as

platinum and the medium surrounding the mouse head was set as air. The locations and geometries

of the WM and CSF with respect to the septal medial nucleus were approximated using an

interactive online mouse brain atlas (http://labs.gaidi.ca/mouse-brain-atlas/). The output from AM

was then processed using a MATLAB script to produce an illustration of the voltage mapping in

the mouse head (Fig. 3B). A one-dimensional voltage profile in the proximity of the electrodes

was plotted to estimate the volume of tissue that was electrically affected/activated (Fig. 3C).

Figure 3. Volume of Tissue Activation A) 3-D illustration of the discretized mouse head model. The bulk tissue model is divided into voxels at a resolution of 31.25 μm. Each voxel represents an equivalent electrical circuit according to its material type. The brain model is mostly uniform as grey matter. The proximity of the medial septal nucleus, where the electrodes are placed, also contains areas of white matter and CSF modeled as basic geometries. B) The voltage mapping near the medial septal nucleus when stimulated with a bipolar electrode configuration. Top row: two sides of the sagittal slice, looking into both the positively and negatively polarized electrodes. Bottom row: the horizontal and coronal slices capturing the voltage distribution around both electrodes. The absolute voltage decays to zero rapidly moving away from the electrodes, demonstrating that the volume of activation is confined within a diameter of about 1 millimeter. C) The voltage amplitude plotted along a single axis. The voltage decay profile moving away from the electrode is very similar along all three axes and the volume of activation

can be modeled as a sphere. The "equivalent diameter" for the volume affected by stimulation is

approximated by choosing points where the slope is approaching zero (green circles), which

determines the current density generated at that location. Thresholding can also be used to

visualize the points where the target voltage amplitude is maintained. The red circles label the

coordinates at which an amplitude of 1 Volt or more is observed. C+ and C- labels mark the

centers of the positive and negative electrodes, respectively.

Data analysis

Data pre-processing

The Iconeus One acquisition system generates power Doppler (pD) images pre-processed

with built-in phase-correlation based sub-pixel motion registration and singular-value-

decomposition (SVD) based clutter filtering algorithms [33]. These algorithms were used to

separate tissue signal from blood signal to obtain pD images (Fig. 4A/B). To correct potential

physiological and motion artifacts, we adopted rigid motion correction techniques that have

successfully been used in fUSI and other neuroimaging studies [34–36]. These were combined

with high frequency filtering algorithms to eliminate noise artifacts. We employed a low-pass

filter with normalized passband frequency of 0.02 Hz, with a stopband attenuation of 60 dB that

compensates for delay introduced by the filter, to remove high-frequency fluctuations in the pD

signals.

Effects of MK-801 on CBV

We investigated the temporal effects of intraperitoneal MK-801 administration on the

septohippocampal circuit (hippocampus, mPFC) and surrounding regions (striatum, pallidum,

thalamus, hypothalamus). To do so, we generated event-related average (ERA) time course

curves of the CBV changes (ΔCBV) as a percentage change of the pD signal from baseline

activity for the selected ROIs. The average pD signal from 2 minutes prior to the saline or drug

injection was used as the baseline. We utilized a repeated measures analysis of variance

(rmANOVA) to assess the effects and interactions between drug (saline and MK801) and ROI

factors over time. We fitted a repeated measures '*within-design*' model to the CBV percentage

change signals over 42-min interval (including 2 minutes just before the drug injection and the

40 minutes after injection) for each mouse and ROI for the rmANOVA analysis. To further

quantify the relative differences in ΔCBV between saline-vehicle and MK-801-treated mice in

various ROIs, we used the last 2 minutes of recordings to compute the mean effects-size

differences in ΔCBVs and the 95% confidence interval of the effect size (if 95% confidence

interval contains zero, then the effect is not significant at the 0.05 level) in each ROI. We also

computed the Cohen's *d* value in each ROI as a measure of the drug effect size that describes the

standardized difference between the means of ΔCBVs in the two groups of animals [37]. A

Cohen's d value of 0.2 represents a small effect size, 0.5 represents a moderate effect size, 0.8

represents a large size and greater than 0.8 represents a very large size.

Effects of MSN DBS during stimulation on brain hemodynamics

We computed the ΔCBV for each ROI to investigate effects of MSN theta- and gamma-

frequency stimulation on the cerebral hemodynamics following saline control or MK-801

injection. We observed ERA time-series of the ΔCBV of each selected ROI relative to the

average pD signal acquired 2 minutes prior to stimulation onset to visualize the temporal

dynamics of DBS effects to CBVs of the ROIs (Fig. 1C).   We utilized a three-way rmANOVA

to assess the effects and interactions between drug (saline and MK801), stimulation (gamma,

theta, no-stimulation) and ROI factors across time during the stimulation process. Here, we fit a

repeated measures '*within-design*' model to the ΔCBV signals over the 7-min period (including 2

minutes prior to stimulation, 5 minutes during stimulation) for each animal and ROI.

Subsequently, we computed the mean effects-size differences in ΔCBVs, the 95% confidence

interval of the effect size, and the Cohen's *d* value in each ROI as a measure of the stimulation

effect size. We measured the mean ΔCBVs during stimulation in theta [saline: n=16, MK-801:

n=12], gamma [saline: n=14, MK-801: n=14], and no-stimulation [saline: n=14; MK-801: n=10]

animal groups. The mean ΔCBVs were calculated utilizing the last 2 minutes of pD signal during

stimulation across animals in each of the three stimulation categories.

Effects of MSN DBS after stimulation offset on brain hemodynamics

We are also interested in assessing the effects of MSN DBS after stimulation offset. To do

so, we repeated the same analysis described above, but we used the 10 minutes of pD signal after

stimulation offset, plus 2 minutes prior to stimulation onset as a baseline. The mean ΔCBVs were

calculated utilizing the last 2 minutes of pD signal after stimulation offset across animals in each

of the three stimulation categories. Our goal is to identify whether MSN DBS causes persistent

and delayed changes in CBVs within and/or outside the septohippocampal network.

Statistical analysis of drug and stimulation effects on ΔCBVs

All the analysis was performed using Matlab Version 9.13.0.2193358 (R2022b). We

assessed the drug and stimulation effects and interactions utilizing the Matlab functions '*fitrm*'

and '*ranova*' for the repeated measures model fitting and the rmANOVA respectively. We

utilized the Greenhouse-Geisser approximation to correct for the possibility of non-compound

symmetry (same variance in means and shared common correlation in paired responses) in the

ROI-time series assessed. The mean effect-size differences and the *Cohen's d* values for the

mean difference in ΔCBVs between saline and MK801, as well as the between no-stimulation-

gamma, no-stimulation-theta, and gamma-theta were computed with the *'meanEffectSize'* Matlab

function.

Figure 4. Functional ultrasound imaging (fUSI) of the mouse brain. A) 3D mouse brain model with fUSI probe positioning (white bar) and ROIs – hippocampus (HPF), medial

prefrontal cortex (mPFC), hypothalamus (HYT), thalamus (TLM), pallidum (PDM), and striatum (STM), superimposed onto a mean grayscale fUSI vascular map of the sagittal mouse brain. **B)** *pD image of cerebral blood volume (CBV) in a sagittal plane (max-min normalized relative scale).*

Results

NMDA antagonist MK-801 reduced blood perfusion

We analyzed 40 minutes of pD signal from the septohippocampal circuit (hippocampus,

mPFC) as well as surrounding structures (hypothalamus, thalamus, pallidum, and striatum) to

assess the effects of MK-801 on cerebral hemodynamics (Fig. 4A). We quantified changes in

CBV (ΔCBV) as a percent change relative to baseline activity (average of 2 minutes pD signal

acquired prior to drug injection). A repeated measures ANOVA (factors treatment $\times$ ROI; where

treatment is saline vs. MK-801 and ROI is the 6 recorded brain areas) revealed that there was a

statistically significant effect over time ($F_{(2519, 56406)} = 8.76$, $p = 6.3$ e-5) after Greenhouse-

Geiser approximation correction. To quantify the effect of MK-801 on CBV, we computed the

percentage change (i.e., ΔCBV) relative to baseline using the last two minutes before stimulation

onset (38-40 minutes after drug injection) and found a reduction of ΔCBV (mean $\pm$ SEM) in the

hippocampus (-3.6 $\pm$ 1.1 %), mPFC (-4.1 $\pm$ 0.61 %), hypothalamus (-1.0 $\pm$ 0.2 %), pallidum (-1.5

$\pm$ 0.4 %), striatum (-1.7 $\pm$ 0.3 %), and thalamus (-3.0 $\pm$ 0.5 %). For saline-treated animals these

values were: hippocampus (-0.7 $\pm$ 0.6 %), mPFC (-0.1 $\pm$ 0.2 %), hypothalamus (-0.7 $\pm$ 0.6 %),

pallidum (0.9 $\pm$ 0.4 %), striatum (-0.3 $\pm$ 0.4 %), and thalamus (0.2 $\pm$ 0.4 %) (Fig. 5A-F, radar

chart insert). The mean effect-size difference in ΔCBVs and Cohen's d analysis comparing mean

ΔCBV induced by saline and MK-801 over a 2-minute interval (38 – 40 minutes post-drug

injection), revealed that MK-801 induces greater decreases in CBV than saline control in all

ROIs investigated (Fig. 5A-F, radar chart insert) - i.e., mPFC (ΔCBV mean difference between

saline and MK-801 animals ± confidence, Cohen's d; 3.96 ± 0.38 %, d= 0.42), thalamus ($3.17 \pm$

0.23 %, d= 0.55), hippocampus (2.82 ± 0.42 %, d= 0.27), pallidum (2.42 ± 0.20 %, d= 0.49),

hypothalamus (1.72 ± 0.12 %, d= 0.59) and striatum (1.33 ± 0.18 %, d= 0.29). Together, these

results indicate that MK-801 reduces CBV both within and outside of the septohippocampal

network.

Figure 5. Event related average (ERA) temporal course curves prior to stimulation onset and mean ΔCBVs during the last 2 minutes in saline and MK-801 treated animals. A – F) Temporal course (42 minutes) of ΔCBV relative to baseline (2-minute average pD signal before saline or MK-801 drug injection) in the A) hippocampus, B) hypothalamus, C) mPFC, D) pallidum, E) striatum, and F) thalamus after saline [blue] and 1.0 mg/kg MK-801 [1] injection. The radar chart insert shows MK-801-induced decreases in CBV in all ROIs compared to saline over the last 2 minutes interval from 38-40 minutes post injection.

<u>MSN stimulation increases CBV in saline control animals in a frequency- and region-dependent</u>

<u>manner.</u>

We assessed whether theta- and gamma-frequency MSN stimulation has disparate

impacts on CBV measures in saline-treated control mice. ERAs of ΔCBV reflect the temporal

responses to theta, gamma, and no stimulation during and post stimulation time periods (Fig. 6,

7). A three-way repeated measures ANOVA (factors, treatment × ROI × DBS; where treatment

is saline vs. MK-801, ROI is the 6 recorded brain areas, and DBS is theta frequency vs. gamma

frequency vs no DBS) was utilized to examine the effects and interactions of drug, stimulation,

and ROIs during the 5 minutes period after onset of MSN stimulation. The two-minute baseline

period pre-stimulation was included in the analysis. We found significant effects of drug over

time ($F(419, 186036) = 7.35$, $p = 1.85$ e-8), stimulation over time ($F(823, 186036) = 2.71$, $p =$

7.20 e-4), as well as, interaction of drug and stimulation over time ($F(838, 186036) = 1.96$, $p =$

1.95 e-2) during the 5 minutes stimulation interval, after Greenhouse-Geisser approximation

correction. To further quantify the effects of DBS, we computed the mean ΔCBV in the last 2

minutes during stimulation across animals in each stimulation category and compared the mean-

effect size differences in ΔCBVs between no stimulation and stimulation in each ROI. We found

that MSN theta stimulation increased CBV compared to no-stimulation only in the mPFC (mean

ΔCBV difference between theta- and no-stimulation ± confidence, Cohen's d; 0.82 ± 0.12 %, d=

0.45) and hippocampus (0.39 ± 0.12 %, d= 0.21). For the rest on ROIs the effect size magnitude

was either very small (i.e., Cohen's d <0.08) or theta-frequency stimulation caused further

reduction in CBVs compared to no-stimulation. On the other hand, MSN gamma stimulation

caused increases in CBV compared to no-stimulation in the mPFC (0.60 ± 0.11 %, d= 0.36),

pallidum (0.38 ± 0.09 %, d= 0.30) and striatum (0.22 ± 0.06 %, d= 0.26). For the rest of the ROIs

the effect size magnitude was either very small (i.e., Cohen's d < 0.095) or gamma-frequency

stimulation resulted in further reduction in CBVs compared to no-stimulation. When comparing

the ΔCBV induced by the theta and gamma stimulation in mPFC – the only ROI that exhibited

moderate effect on both types of stimulations – we found very small effect size on ΔCBV

between the two types of stimulations (mean ΔCBV differences between theta- and gamma-

stimulation ± confidence, Cohen's d; 0.22 ± 0.10, d= 0.14).

The next step was to assess the effects of DBS after stimulation offset (i.e., post-

stimulation). To do so, we conducted a three-way repeated measures ANOVA (factors, treatment

× ROI × DBS; where treatment is saline vs. MK-801, ROI is the 6 recorded brain areas, and DBS

is theta frequency vs. gamma frequency vs no DBS) over the 10 minutes period after the offset of

MSN stimulation. We found significant effects of drug over time $(F(1019, 452436) = 5.28$, p =

1.27 e-4), stimulation over time $(F(2038, 452436) = 3.67$, p = 1.17 e-4), as well as, interaction of

drug and stimulation over time $(F(2038, 186036) = 3.09$, p = 9.40 e-4), after Greenhouse-Geisser

approximation correction, across the 10 minutes post-stimulation interval. We further quantified

the post-effects of DBS by computing the mean ΔCBV in the last 2 minutes of the acquisition –

i.e., 8-10 minutes post-stimulation and comparing the mean-effect size differences on ΔCBV

between no stimulation and stimulation in each ROI. The results showed that MSN theta

stimulation causes increases in CBV compared to no-stimulation in the hippocampus (mean

ΔCBV differences between theta- and no-stimulation ± confidence, Cohen's d; 1.30 ± 0.21, d=

0.42), mPFC (1.20 ± 0.22 %, d= 0.37) and thalamus (0.97 ± 0.11 %, d= 0.58) (Fig. 7A, F – radar

chart). On the other hand, MSN gamma stimulation resulted in an increase in CBV compared to

no stimulation in the mPFC (2.01 ± 0.22 %, d= 0.60), striatum (0.77 ± 0.14 %, d= 0.37) and

pallidum (0.61 ± 0.16 %, d= 0.25). When comparing the differences in ΔCBVs induced by theta

and gamma stimulation in the mPFC – the only ROI that exhibited medium to moderate effect on

both types of stimulations – we found that gamma induces higher ΔCBV than theta stimulation

with medium effect size difference (mean ΔCBV differences between gamma- and theta-

stimulation $\pm$ confidence, Cohen's d; 0.81 ± 0.17, d= 0.31).

<u>Theta-frequency stimulation elicits stronger CBV increases than gamma-frequency stimulation</u>

<u>in MK-801 treated animals.</u>

Recently, our group showed that theta, but not gamma frequency DBS of the MSN

improves spatial memory in MK-801 treated rats [27]. Therefore, we sought to determine if

MSN theta and gamma frequency stimulation had differing impacts on neurovascular activity

measures within memory-associated regions including the mPFC and hippocampus as well as

neighboring regions outside the septohippocampal network (striatum, pallidum, thalamus,

hypothalamus) following MK-801 drug-administration. Again, we assessed the mean effect-size

differences in ΔCBV between MSN theta-, gamma-, and no-stimulation in each of the selected

ROIs, relative to 2 minutes of pD signal recordings just prior to stimulation onset. The analysis

was performed after repeated measures ANOVA over the stimulation and post-stimulation time

intervals to examine the effects and interactions of drug, stimulation, and ROI (results presented

in previous section). Fig. 8 and 9 display the ERA curves for ROIs in response to theta, gamma,

and no DBS during and post stimulation periods in the MK-801 group. We found that MSN

theta stimulation in the MK-801 treated group caused increased ΔCBV with respect to no-

stimulation group with medium to large effect size in all ROIs except mPFC – i.e., hippocampus

(mean ΔCBV differences between theta- and no-stimulation ± confidence, Cohen's d; 2.01 ±

0.20 %, d= 0.78), thalamus (0.49 ± 0.07 %, d= 0.54), pallidum (0.36 ± 0.07 %, d= 0.43), striatum

(0.18 ± 0.04 %, d= 0.31) and hypothalamus (0.14 ± 0.04 %, d= 0.30)   (Fig. 8). Intriguingly, we

only found a medium effect-size increase in ΔCBV for the pallidum (0.60 ± 0.08 %, d= 0.53) and

striatum (0.23 ± 0.05 %, d= 0.38) between MSN gamma- stimulation and no-stimulation groups,

during the stimulation period. For the rest of the ROIs, the effect-size was very small.

Importantly, we found that ΔCBV increased in theta- relative to no-stimulation animals

after stimulation offset with medium to larger effects in the pallidum (1.26 ± 0.15 %, d=0.66),

hippocampus (2.8 ± 0.37 %, d=0.60) and thalamus (1.24 ± 0.17 %, d= 0.57) and small to

medium effects in the striatum (0.50 ± 0.13 %, d= 0.31), hypothalamus (0.21 ± 0.06 %, d= 0.28)

and mPFC (0.64 ± 0.22 %, d= 0.22) (Fig. 9). Importantly, we found only small to medium effects

of gamma MSN stimulation on ΔCBV in the pallidum, relative to stimulation animals, and after

stimulation offset (mean ΔCBV differences between gamna- and no-stimulation ± confidence,

Cohen's d; 0.66 ± 0.13 %, d=0.39) (Fig. 9). For the rest of the ROIs, the effect-size was either

very small (i.e., Cohen's d < 0.1 in hippocampus, striatum and thalamus) or gamma-frequency

stimulation resulted in further CBV reduction (i.e., in hypothalamus and mPFC) compared to no-

stimulation.

Additionally, our results showed medium or large mean effect-size differences in

ΔCBVs between the theta and gamma stimulated MK801 groups after stimulation offset in the

hippocampus (2.67 ± 0.28 %, d= 0.70), mPFC (1.83 ± 0.23 %, d= 0.59), and thalamus (1.15 ±

0.14 %, d= 0.56) (Fig. 9). Together these results demonstrate that theta-frequency stimulation

elicits the strongest CBV response in MK-801-treated mice in the hippocampus and pallidum,

while gamma stimulation had almost no effect on hippocampal CBV (Cohen's d = 0.03) and

decreased CBV in the mPFC compared to no-stimulation mice.

Figure 6. DBS ERA curves after stimulation onset and mean ΔCBVs in saline mice during the last 2 minutes of stimulation period. A – F) Temporal course (theta [orange], gamma [blue], no-stimulation [black]) of mean ΔCBV relative to baseline (2 minutes average pD signal prior to DBS) for the A) hippocampus, B) hypothalamus, C) mPFC, D) pallidum, E) striatum, and F) thalamus regions in the saline-treated animals. Radar chart insert gives the

mean percentage ΔCBVs during stimulation for theta [orange], gamma [blue], and no-stimulation [dark gray] animals in the ROIs investigated. Means were calculated utilizing the last 2 minutes of pD signals acquired during stimulation (3rd – 5th minute after stimulation onset) across animals in each stimulation category.

Figure 7. DBS ERA curves after stimulation onset and mean ΔCBVs in saline mice during the last 2 minutes of recordings in the post-stimulation period in 6 ROIs. *Similar to Figure 6, but the radar chart gives the mean percentage ΔCBVs during post-stimulation for theta*

*[orange], gamma [blue], and no-stimulation [dark gray] for the **A**) hippocampus, **B**) hypothalamus, **C**) mPFC, **D**) pallidum, **E**) striatum, and **F**) thalamus regions in the saline-treated animals. Means were calculated utilizing the last 2 minutes of pD signals acquired post stimulation (8th – 10th minute after stimulation offset) across animals in each stimulation category.*

***Figure 8. DBS ERA curves after stimulation onset and mean ΔCBVs in MK-801 treated mice during the last 2 minutes of stimulation period A – F)** Temporal course (theta [1],*

*gamma [blue], no-stimulation [black]) of mean ΔCBV relative to baseline (2 minutes average pD signal prior to DBS) for **A**) hippocampus, **B**) hypothalamus, **C**) mPFC, **D**) pallidum, **E**) striatum, and **F**) thalamus regions in the MK-801 drug injected mice. Radar chart insert gives the mean percentage ΔCBVs during stimulation for theta [1], gamma [blue], and no-stimulation [dark gray] animals in the ROIs investigated. Means were calculated utilizing the last 2 minutes of pD signals acquired during stimulation ($3^{rd} – 5^{th}$ minute after stimulation onset) across animals in each stimulation category.*

***Figure 9. DBS ERA curves after stimulation onset and mean ΔCBVs in MK-801 treated
mice during the last 2 minutes of recordings in the post-stimulation period.*** *Similar to Figure
8, but the radar chart gives the mean percentage ΔCBVs in post-stimulation period for theta
[1], gamma [blue], and no-stimulation [dark gray] for the **A)** hippocampus, **B)** hypothalamus,
C) mPFC, **D)** pallidum, **E)** striatum, and **F)** thalamus regions in the MK-801 treated animals.
Means were calculated utilizing the last 2 minutes of pD signals acquired post stimulation (8th
– 10th minute after stimulation offset) across animals in each stimulation category.*

Discussion

The present study utilized the high spatiotemporal resolution and sensitivity of fUSI to

demonstrate that acute administration of MK-801 causes a significant reduction in CBV across

all ROIs. Furthermore, we demonstrated that theta frequency MSN DBS alters regions within the

septohippocampal network, with the strongest effect on the hippocampus. Intriguingly, the

observed increase in hippocampal CBV remain even after cessation of DBS. On the other hand,

structures outside the septohippocampal network, such as the hypothalamus and striatum, show

less of a response to DBS. These effects were less pronounced with gamma frequency

stimulation with very small effects on the hippocampus. These findings suggest that MSN theta

frequency DBS precisely modulates neurovascular activity in cognitive networks [27].

MK-801 reduced CBV in all ROIs

MK-801 and other NMDA antagonists have been widely used in preclinical models to

mimic the behavioral and electrophysiological deficits associated with schizophrenia

[16,18,38,39]. However, the regionally specific effects of MK-801 on CBV in such models is

not well known. We observed that MK-801 reduced CBV across all ROIs. Importantly,

previous fMRI studies have observed reduced BOLD signals in hippocampal and prefrontal areas

in schizophrenia patients [40–42]. In this context, our findings support the use of MK-801 as a

neurovascular model of schizophrenia. Furthermore, our study demonstrates the feasibility of

using fUSI to identify network-specific hemodynamic changes as an additional modality for

studying neurocognitive disorders.

MSN theta stimulation was relatively specific to cognitive networks.

We observed that theta frequency MSN DBS resulted in an increase to CBV in the

hippocampus in both the saline- and MK-801-treated animals. Importantly, this effect was

greatest in the hippocampus, which receives direct projections from the MSN, and is a primary

target for neuromodulatory interventions to treat memory dysfunction [23–25,27]. In saline

control animals, significant increases to CBV were also observed in the mPFC (during and after

stimulation) and thalamus (after stimulation), both of which play important roles in memory

function and receive direct projections from the MSN (Fig. 2A) [22]. Interestingly, gamma

stimulation did not alter hippocampal CBV, but did increase regions of the brain that were

anatomically closer to the MSN. Specifically, mPFC CBV was increased in saline- treated

animals, while the pallidum and striatum were increased to a lesser degree in MK-801-treated

animals. This suggests that MSN gamma stimulation may have a local response to stimulation

but is less specific to the neural circuitry being stimulated.

MSN theta stimulation increased hippocampal CBV during and after stimulation despite NMDA

antagonism.

A leading hypothesis is that reduced N-methyl-D-aspartate (NMDA) receptor-mediated

glutamatergic transmission underlies psychiatric conditions involving cognitive and memory

dysfunction [16,18,38,39,43,44]. Reductions to NMDA activity either pharmacologically or

through genetic manipulation have been shown to decrease theta activity, increase gamma

activity and lead to deficits in spatial navigation and memory [16,17,27,45,46]. Research by our

group has found that acute (<5 minutes) theta frequency (7.7 Hz), but not gamma frequency (100

Hz) stimulation of the MSN during the Barnes maze task improves spatial memory in rodents

following pharmacological NMDA antagonism [27]. Further, we have also demonstrated that

MSN theta stimulation prior to the task can also improve spatial memory [23]. Interestingly,

hippocampal theta oscillations return to baseline after cessation of MSN DBS. Therefore, the

question remains open as to how MSN DBS mediates sustained improvements. In this current

study, we demonstrate that hippocampal CBV remains elevated after cessation of MSN

stimulation.

<u>MSN theta stimulation may drive high frequency or spiking activity via hippocampal</u>

<u>interneurons.</u>

A number of studies utilizing various modalities in combination with fUSI have found a

strong relationship between pD signal and neuronal activity [47–49]. This relationship was also

true for high frequency oscillatory activity (~100Hz) but was much weaker for lower frequency

oscillations [49]. This suggests that the theta-induced increases to hippocampal CBV may reflect

increased hippocampal gamma or spiking activity, rather than increases to theta oscillatory

activity itself. However, given that gamma band activity is often correlated with spiking activity,

differentiating between changes in oscillatory dynamics and spiking activity is not possible in the

present study [50].

GABAergic interneurons in the hippocampus play an important role in synchronizing

hippocampal oscillatory activity. Indeed, inhibitory neurons are hypothesized to be a primary

source of dysfunction in pathologies involving NMDA dysfunction (for review see [51]) and are

inhibited by NMDA-antagonists [52,53]. One possibility is that theta frequency MSN

stimulation may, by briefly stimulating afferent populations at theta-frequency, act as a 'reset'

and allow synchronous innervation of hippocampal interneurons that regulate the activity of

glutamatergic pyramidal cells. Indeed, it has been previously shown that optogenetic stimulation

of GABAergic neurons decreases spontaneous neural activity and leads to an increase in local

blood flow [54]. This hypothesis is further supported by a recent study by Nunez-Elizadle and

colleagues who observed that the relationship between the fUSI signal and firing rates were

greatest for putative interneurons [49]. Future studies combining single unit activity and fUSI

could test this hypothesis.

MSN gamma stimulation did not affect hippocampal CBV.

Our previous work suggests that MSN theta, but not gamma stimulation can improve

spatial memory in MK-801-treated rodents [27]. While MSN theta stimulation increased

hippocampal CBV during and after stimulation in MK-801-treated animals, this was not true of

MSN gamma stimulation. Gamma stimulation resulted in delayed increases to mPFC CBV in

saline-treated animals and had no effect on MK-801-treated animals in any of the ROIs. These

results suggest that MSN gamma stimulation is not sufficient to engage hippocampal activity and

highlights the importance of frequency parameters in DBS paradigms for spatial memory. This is

further supported by our previous study demonstrating no improvement to spatial memory in

MK-801 treated animals with MSN gamma stimulation [27].

Implications for neuromodulation

We observed that theta-frequency stimulation of the MSN increased blood perfusion in

the hippocampus following cessation of the stimulus. These effects were not observed using

gamma stimulation and were still present even under conditions of pharmacologic NMDA

antagonism. It is worth noting that arguably the most effective form of neuromodulation is still

electroconvulsive therapy (ECT), which is performed under anesthesia and also results in

increases to cerebral blood flow [55,56]. However, ECT is very non-specific, this lack of

specificity may well contribute to its detrimental effect on memory [57]. Alternatively,

transcranial magnetic stimulation (TMS) is most effective when applied focally to awake, alert

patients [58,59]. Because DBS can combine a relatively high degree of modulation in deep

structures with greater spatial and temporal specificity than ECT or TMS, it is plausible that DBS

may have more benefits beyond that of ECT or TMS in treating disorders of cognitive function.

<u>Limitations and future directions</u>

While the current study was performed in anesthetized animals, futures studies will

investigate the effects of reduced NMDA function and MSN DBS in awake, behaving animals

during memory-associated behavioral tasks (e.g., novel object recognition and Barnes Maze).

The goal will be to determine if the observed ΔCBV within the septo-hippocampal network

following theta- frequency MSN DBS is also associated with improved memory function,

linking the present study with our previous study demonstrating improved memory following

MSN DBS in MK-801 treated animals [27]. Another limitation of our study is that fUSI

recordings performed using the conventional 1-dimensional linear ultrasound transducer array

necessarily generates 2-dimensional pD vascular maps of the animals' CBV. As a result, other

regions that are connected with the MSN besides the hippocampus and mPFC (e.g., amygdala,

habenula, raphe nucleus) were not accessible from the selected sagittal 2-dimensional image

plane. Recent studies are tackling this challenge using whole-brain 3-dimensional fUSI with

either moving linear arrays (similar to the array used in our study), matrix arrays or raw column

arrays (RCAs) [60–62]. Future studies can use these probes to cover volumes rather slices of the

mouse brain providing access to all areas of the septohippocampal network. Overall and

regardless of these limitations, our findings demonstrate the feasibility of using fUSI to

characterize network-specific neurovascular changes in disease models as well understand what

parameters in neuromodulatory techniques most impact cerebral perfusion dynamics.

References:

[1]   Ranasinghe KG, Petersen C, Kudo K, Mizuiri D, Rankin KP, Rabinovici GD, et al. Reduced synchrony in alpha oscillations during life predicts post mortem neurofibrillary tangle density in early-onset and atypical Alzheimer's disease. Alzheimer's & Dementia 2021;17:2009–19. https://doi.org/10.1002/alz.12349.

[2]   Tu M-C, Chung H-W, Hsu Y-H, Yang J-J, Wu W-C. Stage-Dependent Cerebral Blood Flow and Leukoaraiosis Couplings in Subcortical Ischemic Vascular Disease and Alzheimer's Disease. J Alzheimers Dis 2022;86:729–39. https://doi.org/10.3233/JAD-215405.

[3]   Dougherty DD, Chou T, Corse AK, Arulpragasam AR, Widge AS, Cusin C, et al. Acute deep brain stimulation changes in regional cerebral blood flow in obsessive-compulsive disorder. J Neurosurg 2016;125:1087–93. https://doi.org/10.3171/2015.9.JNS151387.

[4]   Laxton AW, Tang-Wai DF, McAndrews MP, Zumsteg D, Wennberg R, Keren R, et al. A phase I trial of deep brain stimulation of memory circuits in Alzheimer's disease. Ann Neurol 2010;68:521–34. https://doi.org/10.1002/ana.22089.

[5]   Macé E, Montaldo G, Cohen I, Baulac M, Fink M, Tanter M. Functional ultrasound imaging of the brain. Nat Methods 2011;8:662–4. https://doi.org/10.1038/nmeth.1641.

[6]   Mace E, Montaldo G, Osmanski B-F, Cohen I, Fink M, Tanter M. Functional ultrasound imaging of the brain: theory and basic principles. IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control 2013;60:492–506. https://doi.org/10.1109/TUFFC.2013.2592.

[7]   Rabut C, Ferrier J, Bertolo A, Osmanski B, Mousset X, Pezet S, et al. Pharmaco-fUS: Quantification of pharmacologically-induced dynamic changes in brain perfusion and connectivity by functional ultrasound imaging in awake mice. NeuroImage 2020;222:117231. https://doi.org/10.1016/j.neuroimage.2020.117231.

[8]   Norman SL, Maresca D, Christopoulos VN, Griggs WS, Demene C, Tanter M, et al. Single-trial decoding of movement intentions using functional ultrasound neuroimaging. Neuron 2021;109:1554-1566.e4. https://doi.org/10.1016/j.neuron.2021.03.003.

[9]   Uhlhaas PJ, Singer W. Abnormal neural oscillations and synchrony in schizophrenia. Nature Reviews Neuroscience 2010;11:100–13. https://doi.org/10.1038/nrn2774.

[10]   Barr MS, Rajji TK, Zomorrodi R, Radhu N, George TP, Blumberger DM, et al. Impaired theta-gamma coupling during working memory performance in schizophrenia. Schizophrenia Research 2017;189:104–10. https://doi.org/10.1016/j.schres.2017.01.044.

[11]   Jafari Z, Kolb BE, Mohajerani MH. Neural oscillations and brain stimulation in Alzheimer's disease. Prog Neurobiol 2020;194:101878. https://doi.org/10.1016/j.pneurobio.2020.101878.

[12]   Gonzalez-Burgos G, Lewis DA. NMDA Receptor Hypofunction, Parvalbumin-Positive Neurons, and Cortical Gamma Oscillations in Schizophrenia. Schizophrenia Bulletin 2012;38:950–7. https://doi.org/10.1093/schbul/sbs010.

[13]   Gu Z, Alexander GM, Dudek SM, Yakel JL. Hippocampus and Entorhinal Cortex Recruit Cholinergic and NMDA Receptors Separately to Generate Hippocampal Theta Oscillations. Cell Reports 2017;21:3585–95. https://doi.org/10.1016/j.celrep.2017.11.080.

[14]   Collingridge G. The role of NMDA receptors in learning and memory. Nature 1987;330:604–5. https://doi.org/10.1038/330604a0.

[15] Spangler EL, Bresnahan EL, Garofalo P, Muth NJ, Heller B, Ingram DK. NMDA receptor
channel antagonism by dizocilpine (MK-801) impairs performance of rats in aversively
motivated complex maze tasks. Pharmacol Biochem Behav 1991;40:949–58.
https://doi.org/10.1016/0091-3057(91)90111-e.
[16] Newcomer JW, Farber NB, Jevtovic-Todorovic V, Selke G, Melson AK, Hershey T, et al.
Ketamine-Induced NMDA Receptor Hypofunction as a Model of Memory Impairment and
Psychosis. Neuropsychopharmacology 1999;20:106–18. https://doi.org/10.1016/S0893-
133X(98)00067-0.
[17] Korotkova T, Fuchs EC, Ponomarenko A, von Engelhardt J, Monyer H. NMDA receptor
ablation on parvalbumin-positive interneurons impairs hippocampal synchrony, spatial
representations, and working memory. Neuron 2010;68:557–69.
https://doi.org/10.1016/j.neuron.2010.09.017.
[18] Saunders JA, Gandal MJ, Siegel SJ. NMDA antagonists recreate signal-to-noise ratio and
timing perturbations present in schizophrenia. Neurobiology of Disease 2012;46:93–100.
https://doi.org/10.1016/J.NBD.2011.12.049.
[19] Billingslea EN, Tatard-Leitman VM, Anguiano J, Jutzeler CR, Suh J, Saunders JA, et al.
Parvalbumin Cell Ablation of NMDA-R1 Causes Increased Resting Network Excitability
with Associated Social and Self-Care Deficits. Neuropsychopharmacol 2014;39:1603–13.
https://doi.org/10.1038/npp.2014.7.
[20] Fuhrmann F, Justus D, Sosulina L, Kaneko H, Beutel T, Friedrichs D, et al. Locomotion,
Theta Oscillations, and the Speed-Correlated Firing of Hippocampal Neurons Are
Controlled by a Medial Septal Glutamatergic Circuit. Neuron 2015;86:1253–64.
https://doi.org/10.1016/j.neuron.2015.05.001.
[21] Leão RN, Targino ZH, Colom LV, Fisahn A. Interconnection and synchronization of
neuronal populations in the mouse medial septum/diagonal band of Broca. Journal of
Neurophysiology 2015;113:971–80. https://doi.org/10.1152/jn.00367.2014.
[22] Takeuchi Y, Nagy AJ, Barcsai L, Li Q, Ohsawa M, Mizuseki K, et al. The Medial Septum
as a Potential Target for Treating Brain Disorders Associated With Oscillopathies. Frontiers
in Neural Circuits 2021;15.
[23] Lee DJ, Gurkoff GG, Izadi A, Berman RF, Ekstrom AD, Muizelaar JP, et al. Medial Septal
Nucleus Theta Frequency Deep Brain Stimulation Improves Spatial Working Memory after
Traumatic Brain Injury. Journal of Neurotrauma 2013;30:131–9.
https://doi.org/10.1089/neu.2012.2646.
[24] Lee DJ, Gurkoff GG, Izadi A, Seidl SE, Echeverri A, Melnik M, et al. Septohippocampal
Neuromodulation Improves Cognition after Traumatic Brain Injury. Journal of
Neurotrauma 2015;32:1822–32. https://doi.org/10.1089/neu.2014.3744.
[25] Lee DJ, Izadi A, Melnik M, Seidl S, Echeverri A, Shahlaie K, et al. Stimulation of the
medial septum improves performance in spatial learning following pilocarpine-induced
status epilepticus. Epilepsy Research 2017;130:53–63.
https://doi.org/10.1016/j.eplepsyres.2017.01.005.
[26] Cole ER, Grogan DP, Laxpati NG, Fernandez AM, Skelton HM, Isbaine F, et al. Evidence
supporting deep brain stimulation of the medial septum in the treatment of temporal lobe
epilepsy. Epilepsia 2022;63:2192–213. https://doi.org/10.1111/epi.17326.
[27] Zepeda NC, Crown LM, Medvidovic S, Choi W, Sheth M, Bergosh M, et al. Frequency-
specific medial septal nucleus deep brain stimulation improves spatial memory in MK-801-

treated male rats. Neurobiology of Disease 2022;170:105756.
https://doi.org/10.1016/j.nbd.2022.105756.

[28] Grubbs FE. Procedures for Detecting Outlying Observations in Samples. Technometrics
1969;11:1–21. https://doi.org/10.1080/00401706.1969.10490657.

[29] Stefansky W. Rejecting Outliers in Factorial Designs. Technometrics 1972;14:469–79.
https://doi.org/10.2307/1267436.

[30] Wegener N, Nagel J, Gross R, Chambon C, Greco S, Pietraszek M, et al. Evaluation of
brain pharmacokinetics of (+)MK-801 in relation to behaviour. Neurosci Lett 2011;503:68–
72. https://doi.org/10.1016/j.neulet.2011.08.012.

[31] Wang Q, Ding S-L, Li Y, Royall J, Feng D, Lesnar P, et al. The Allen Mouse Brain
Common Coordinate Framework: A 3D Reference Atlas. Cell 2020;181:936-953.e20.
https://doi.org/10.1016/j.cell.2020.04.007.

[32] Paknahad J, Kosta P, Iseri E, Farzad S, Bouteiller J-MC, Humayun MS, et al. Modeling ON
Cone Bipolar Cells for Electrical Stimulation. Annu Int Conf IEEE Eng Med Biol Soc
2021;2021:6547–50. https://doi.org/10.1109/EMBC46164.2021.9629884.

[33] Ledoux LA, Brands PJ, Hoeks AP. Reduction of the clutter component in Doppler
ultrasound signals based on singular value decomposition: a simulation study. Ultrason
Imaging 1997;19:1–18. https://doi.org/10.1177/016173469701900101.

[34] Stringer C, Pachitariu M. Computational processing of neural recordings from calcium
imaging data. Current Opinion in Neurobiology 2019;55:22–31.
https://doi.org/10.1016/j.conb.2018.11.005.

[35] Friedrich J, Giovannucci A, Pnevmatikakis EA. Online analysis of microendoscopic 1-
photon calcium imaging data streams. PLOS Computational Biology 2021;17:e1008565.
https://doi.org/10.1371/journal.pcbi.1008565.

[36] Pnevmatikakis EA, Giovannucci A. NoRMCorre: An online algorithm for piecewise rigid
motion correction of calcium imaging data. Journal of Neuroscience Methods 2017;291:83–
94. https://doi.org/10.1016/j.jneumeth.2017.07.031.

[37] Liu XS. Statistical Power Analysis for the Social and Behavioral Sciences: Basic and
Advanced Techniques. Routledge; 2013.

[38] Balu DT. The NMDA Receptor and Schizophrenia: From Pathophysiology to Treatment.
Adv Pharmacol 2016;76:351–82. https://doi.org/10.1016/bs.apha.2016.01.006.

[39] Farber NB. The NMDA Receptor Hypofunction Model of Psychosis. Annals of the New
York Academy of Sciences 2003;1003:119–30. https://doi.org/10.1196/annals.1300.008.

[40] MacDonald SWS, Nyberg L, Bäckman L. Intra-individual variability in behavior: links to
brain structure, neurotransmission and neuronal activity. Trends in Neurosciences
2006;29:474–80. https://doi.org/10.1016/J.TINS.2006.06.011.

[41] Gur RE, Gur RC. Functional magnetic resonance imaging in schizophrenia. Dialogues in
Clinical Neuroscience 2010;12:333–43. https://doi.org/10.31887/DCNS.2010.12.3/rgur.

[42] Bedford NJ, Surguladze S, Giampietro V, Brammer MJ, David AS. Self-evaluation in
schizophrenia: an fMRI study with implications for the understanding of insight. BMC
Psychiatry 2012;12:106. https://doi.org/10.1186/1471-244X-12-106.

[43] Moghaddam B, Javitt D. From Revolution to Evolution: The Glutamate Hypothesis of
Schizophrenia and its Implication for Treatment. Neuropsychopharmacol 2012;37:4–15.
https://doi.org/10.1038/npp.2011.181.

[44] Lin C-H, Huang Y-J, Lin C-J, Lane H-Y, E. Tsai G. NMDA Neurotransmission
Dysfunction in Mild Cognitive Impairment and Alzheimer's Disease. Current
Pharmaceutical Design 2014;20:5169–79.

[45] Stark E, Eichler R, Roux L, Fujisawa S, Rotstein HG, Buzsáki G. Inhibition-induced theta
resonance in cortical circuits. Neuron 2013;80.
https://doi.org/10.1016/j.neuron.2013.09.033.

[46] Ward KR, Featherstone RE, Naschek MJ, Melnychenko O, Banerjee A, Yi J, et al. Src
deficient mice demonstrate behavioral and electrophysiological alterations relevant to
psychiatric and developmental disease. Progress in Neuro-Psychopharmacology and
Biological Psychiatry 2019;93:84–92. https://doi.org/10.1016/j.pnpbp.2019.02.017.

[47] Boido D, Rungta RL, Osmanski B-F, Roche M, Tsurugizawa T, Le Bihan D, et al.
Mesoscopic and microscopic imaging of sensory responses in the same animal. Nat
Commun 2019;10:1110. https://doi.org/10.1038/s41467-019-09082-4.

[48] Aydin A-K, Haselden WD, Goulam Houssen Y, Pouzat C, Rungta RL, Demené C, et al.
Transfer functions linking neural calcium to single voxel functional ultrasound signal. Nat
Commun 2020;11:2954. https://doi.org/10.1038/s41467-020-16774-9.

[49] Nunez-Elizalde AO, Krumin M, Reddy CB, Montaldo G, Urban A, Harris KD, et al. Neural
correlates of blood flow measured by ultrasound. Neuron 2022;110:1631-1640.e4.
https://doi.org/10.1016/j.neuron.2022.02.012.

[50] Buzsáki G, Schomburg EW. What does gamma coherence tell us about inter-regional
neural communication? Nat Neurosci 2015;18:484–9. https://doi.org/10.1038/nn.3952.

[51] Marín O. Interneuron dysfunction in psychiatric disorders. Nat Rev Neurosci 2012;13:107–
20. https://doi.org/10.1038/nrn3155.

[52] Homayoun H, Moghaddam B. NMDA receptor hypofunction produces opposite effects on
prefrontal cortex interneurons and pyramidal neurons. J Neurosci 2007;27:11496–500.
https://doi.org/10.1523/JNEUROSCI.2213-07.2007.

[53] Cohen SM, Tsien RW, Goff DC, Halassa MM. The impact of NMDA Receptor
hypofunction on GABAergic interneurons in the pathophysiology of schizophrenia.
Schizophr Res 2015;167:98–107. https://doi.org/10.1016/j.schres.2014.12.026.

[54] Anenberg E, Chan AW, Xie Y, LeDue JM, Murphy TH. Optogenetic Stimulation of GABA
Neurons can Decrease Local Neuronal Activity While Increasing Cortical Blood Flow. J
Cereb Blood Flow Metab 2015;35:1579–86. https://doi.org/10.1038/jcbfm.2015.140.

[55] Espinoza RT, Kellner CH. Electroconvulsive Therapy. New England Journal of Medicine
2022;386:667–72. https://doi.org/10.1056/NEJMra2034954.

[56] Singh A, Kar SK. How Electroconvulsive Therapy Works?: Understanding the
Neurobiological Mechanisms. Clinical Psychopharmacology and Neuroscience
2017;15:210–21. https://doi.org/10.9758/cpn.2017.15.3.210.

[57] Andrade C, Arumugham SS, Thirthalli J. Adverse Effects of Electroconvulsive Therapy.
Psychiatric Clinics 2016;39:513–30. https://doi.org/10.1016/j.psc.2016.04.004.

[58] Gersner R, Kravetz E, Feil J, Pell G, Zangen A. Long-Term Effects of Repetitive
Transcranial Magnetic Stimulation on Markers for Neuroplasticity: Differential Outcomes
in Anesthetized and Awake Animals. J Neurosci 2011;31:7521–6.
https://doi.org/10.1523/JNEUROSCI.6751-10.2011.

[59] Xu T, Sturgeon D, Ramirez JSB, Froudist-Walsh S, Margulies DS, Schroeder CE, et al.
Interindividual Variability of Functional Connectivity in Awake and Anesthetized Rhesus

Macaque Monkeys. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging
2019;4:543–53. https://doi.org/10.1016/j.bpsc.2019.02.005.

[60] Bertolo A, Nouhoum M, Cazzanelli S, Ferrier J, Mariani J-C, Kliewer A, et al. Whole-
Brain 3D Activation and Functional Connectivity Mapping in Mice using Transcranial
Functional Ultrasound Imaging. JoVE (Journal of Visualized Experiments) 2021:e62267.
https://doi.org/10.3791/62267.

[61] Rabut C, Correia M, Finel V, Pezet S, Pernot M, Deffieux T, et al. 4D functional ultrasound
imaging of whole-brain activity in rodents. Nat Methods 2019;16:994–7.
https://doi.org/10.1038/s41592-019-0572-y.

[62] Sauvage J, Porée J, Rabut C, Férin G, Flesch M, Rosinski B, et al. 4D Functional Imaging
of the Rat Brain Using a Large Aperture Row-Column Array. IEEE Transactions on
Medical Imaging 2020;39:1884–93. https://doi.org/10.1109/TMI.2019.2959833.