SNAP25 disease mutations change the energy landscape for synaptic exocytosis

due to aberrant SNARE interactions

Anna Kádková[1,3], Jacqueline Murach[2,3], Maiken Ø. Pedersen[1,3], Andrea Malsam[2], Jörg Malsam[2],

Thomas H. Söllner[2], and Jakob B. Sørensen[1]

[1] Department of Neuroscience, University of Copenhagen, Blegdamsvej 3B, 2200 Copenhagen N,

Denmark.

[2] Heidelberg University Biochemistry Center, 69120 Heidelberg, Germany.

[3] These authors contributed equally.

Running title: SNAP25 disease mutations changing priming and energy barriers

*Correspondence to:

Jakob B. Sørensen
Department of Neuroscience
University of Copenhagen
Blegdamsvej 3B
2200 Copenhagen N
Denmark.
Email: jakobbs@sund.ku.dk.

Abstract

SNAP25 is one of three neuronal SNAREs driving synaptic vesicle exocytosis. We studied three mutations in SNAP25 that cause epileptic encephalopathy: V48F, and D166Y in the Synaptotagmin-1 (Syt1) binding interface, and I67N, which destabilizes the SNARE-complex. All three mutations reduced Syt1-dependent vesicle docking to SNARE-carrying liposomes and Ca^{2+}-stimulated membrane fusion *in vitro* and in neurons. The V48F and D166Y mutants (with potency D166Y > V48F) led to reduced Readily Releasable Pool (RRP) size, due to increased spontaneous (mEPSC) release and decreased priming rates. These mutations lowered the energy barrier for fusion and increased the release probability, which are gain-of-function features not found in *Syt1* knockout (KO) neurons; normalized mEPSC release rates were higher (potency D166Y>V48F) than in the *Syt1* KO. These mutations (potency D166Y > V48F) increased spontaneous association to partner SNAREs, resulting in unregulated membrane fusion. In contrast, the I67N mutant decreased mEPSC frequency and EPSC amplitudes due to an increase in the apparent height of the energy barrier for fusion, whereas the RRP size was unaffected. This could be partly compensated by positive charges lowering the energy barrier. Overall, pathogenic mutations in SNAP25 cause complex changes in the energy landscape for priming and fusion.

Introduction

The fusion machinery responsible for chemical synaptic transmission is well known: it consists of

the SNARE-complex, a ternary complex formed by the proteins VAMP2, syntaxin-1 and SNAP25

(Sutton et al., 1998). This complex is under tight control by upstream partner protein Munc18-1,

which acts as a template for SNARE-complex formation, and Munc13s, which assist in the transitions

required along the pathway of assembly (Rizo, 2022). SNARE-complexes assemble in a zipper-like

manner; partially assembled SNAREpins bind synaptotagmin-1 (Syt1) and complexin within two

separate interaction sites: the primary interface formed between a synaptotagmin 1 molecule and

syntaxin-1 and SNAP25 (Zhou et al., 2015), and the tripartite interface formed by syntaxin-1 and

VAMP2 with another Syt1 molecule and complexin (Zhou et al., 2017). Upon arrival of an action

potential, Ca^{2+} binds to the two C2-domains of Syt1, which results in rapid vesicle-plasma membrane

fusion and release of neurotransmitter within a fraction of a millisecond (Sudhof, 2013).

The strong functional integration and specialization of the neuronal SNARE for speed has rendered

the release machinery exquisitely susceptible to insults. *De novo* mutations in SNAREs and

associated proteins lead to complex neurological disease, characterized by drug-resistant epilepsy,

intellectual disability, movement disorders and often autism; a syndrome which has been denoted

"SNAREopathy" (Verhage and Sorensen, 2020). Although rare, these are devastating conditions for

the patients and their families. Consequently, there is considerable interest in revealing the

molecular/cellular mechanisms for these conditions, which is seen as key to the development of

treatment. Disease-causing mutations in the SNARE-machinery fall into distinct categories according

to the nature of the defect (Verhage and Sorensen, 2020). These include different forms of haplo-

insufficiency, where the mutated protein is either lost altogether or has lost its functionality, and

dominant-negative or recessive variants, as well as variants with new or changed protein

interactions, referred to as neomorphs.

Most disease mutations have been described in *STXBP1* (Abramov et al., 2021a; Verhage and

Sorensen, 2020; Xian et al., 2022), the gene encoding Munc18-1. These mutations are generally

found to cause Munc18-1 hypo-expression and thus the mutations belong in the haploinsufficiency

category. In some cases, associated synaptic phenotypes have been identified, often a decrease in

evoked release (Kovacevic et al., 2018); however, for other mutants no electrophysiological

correlate was found (Kovacevic et al., 2018). The reason for haploinsufficiency/hypoexpression of

most *STXBP1* disease mutations has been suggested to be protein instability, leading to degradation

(Guiberson et al., 2018; Kovacevic et al., 2018; Martin et al., 2014; Saitsu et al., 2008). Consequently,

attempts to rescue the disease phenotypes have focused on mechanisms for increasing general

expression levels, for instance chemical chaperones that might prevent Munc18-1 misfolding and

degradation (Abramov et al., 2021b; Guiberson et al., 2018).

The situation is different for disease mutations in Syt1 and SNAP25. In these cases, mutation

generally does not cause protein instability, but instead changes function in different ways. For Syt1,

disease mutations were found to cluster in the C2B-domain around the top loops that coordinate

Ca^{2+} (Baker et al., 2018). Expression of these mutants in neurons demonstrated reduced evoked

release, in some cases an increase in spontaneous release, and a dominant-negative phenotype

when co-expressed with wildtype protein (Bradberry et al., 2020). The molecular mechanism was

identified as a decrease in Ca^{2+}-dependent lipid binding (Bradberry et al., 2020). In SNAP25, disease

causing mutations are found within the SNARE-domains (Hamdan et al., 2017; Klockner et al., 2021;

Rohena et al., 2013; Shen et al., 2014). Alten et al. studied a selection of SNAP25 mutants, and found

no changes in expression levels, but changes in both spontaneous and evoked release (Alten et al.,

2021). Specifically, mutations in the primary Syt1:SNARE interface (Zhou et al., 2015) caused an

increase in spontaneous release rates, and a decrease in evoked release amplitudes. Conversely, C-

terminal mutations in the so-called 'layer residues', whose side chains point to the center of the

SNARE complex and are involved in SNAREpin zippering (Sutton et al., 1998), led to a decrease in

both evoked and spontaneous release (Alten et al., 2021). This is expected because the C-terminal

end of the SNARE-complex is required for both types of release (Weber et al., 2010).

Alten et al. (2021) described striking phenotypes for most SNAP25 mutations tested, but the

molecular reason for these phenotypes remains incompletely understood, and a few findings were

surprising. For instance, Alten et al. reported that the V48F and D166Y mutants supported an

unchanged Readily Releasable Pool (RRP) of vesicles. The RRP is the pool of vesicles available for

immediate release upon invasion of the presynapse by the action potential, and it is often assessed

by applying a hyperosmotic solution (often 0.5 M sucrose), which causes primed vesicles to fuse

(Rosenmund and Stevens, 1996). Previously, we concluded that the primary Syt1:SNARE-interface

is involved in vesicle priming (Schupp et al., 2016), which agrees with the suggestion that Syt1 binds

to the SNAREs before Ca^{2+} arrival (Zhou et al., 2015). Conversely, Alten et al. reported a smaller RRP

for the I67N mutant, but the I67 residue is present in the internal of layer +4, which we expected to

affect final SNARE complex zippering causing membrane fusion rather than priming (Sorensen et al.,

2006; Weber et al., 2010). Another open question is whether the phenotype of the mutants in the

primary interface (V48F, D166Y) are explainable solely by the loss of Syt1 binding, or whether other

features of these mutations add to, or detract from, the phenotype.

Here, we reexamined three SNAP25 disease mutations: I67N, V48F and D166Y. We confirmed the

increase in spontaneous release rate and decrease in evoked release by V48F and D166Y reported

previously (Alten et al., 2021). Through *in vitro* analysis we find that both mutations lower Syt1

association to SNARE-protein liposomes. Additional experiments both *in vitro* and in cells

demonstrate that V48F and especially D166Y represent partial gain-of-function mutations that

increase association to partner SNAREs and lower the energy barrier for fusion, bypassing Syt1-

dependent control. Thus, these mutants do not phenocopy the loss of Syt1, but combine loss of Syt1

binding with a gain-of-function phenotype. At the same time, the mutants act as loss-of-function in

upstream reactions, through effects on priming. The I67N is a classical dominant-negative mutation,

which increases the energy barrier for fusion, but does not change the size of the RRP if probed by

a sufficiently high concentration of sucrose, nor does the I67N change the electrostatics of triggering

itself. Thus, for V48F and D166Y, loss-of-function and gain-of-function features combine to change

the energy landscape for vesicle priming and fusion in a complex way. These findings have

consequences for our understanding of the simultaneous role of the primary SNARE:Syt1 interface

in vesicle priming and release clamping. It further demonstrates the challenge faced by finding

mechanism-based treatments of these disorders in the presence of multiple effects caused by single

point mutations.

Results

We first investigated two SNAP25 disease-causing mutations within the primary Syt1:SNARE-interface (V48F, D166Y; Fig. 1A-C). The mutations occurred *de novo* and were identified in single heterozygous subjects. The V48F mutant was identified in a 15-year-old female with encephalopathy, intellectual disability and generalized epilepsy with seizures started at 5 months of age; MRI was normal except for delayed myelination (Rohena et al., 2013). The D166Y mutants was found in a 23-year-old male with global developmental delay, nocturnal tonic-clonic seizures, and moderate intellectual disability; the MRI showed mild diffuse cortical atrophy (Hamdan et al., 2017). We aimed to achieve a detailed understanding of the reason for synaptic dysfunction.

The V48F and D166Y mutants disinhibit spontaneous release and desynchronize evoked release

We constructed lentiviral vectors, which expressed wildtype (WT) or mutant SNAP25b fused N-terminally to EGFP as an expression marker (Delgado-Martinez et al., 2007). In the absence of SNAP25, neuronal viability is compromised (Delgado-Martinez et al., 2007; Peng et al., 2013; Santos et al., 2017; Weber et al., 2010) with low-density autaptic glutamatergic neurons dying within a few days, whereas *Snap25* KO neurons growing at higher densities, or in the intact brain, can survive longer, but also eventually degenerate (Bronk et al., 2007; Hoerder-Suabedissen et al., 2019). We therefore first examined the morphology and survival of autaptic hippocampal neurons from *Snap25* KO after expressing mutated or WT EGFP-SNAP25b (henceforth denoted 'WT' in rescue experiments; not to be confused with '*Syt1* WT', which refers to wildtype littermates of *Syt1* KO mice).

As expected, *Snap25* KO autaptic neurons did not survive in the absence of expression of exogeneous SNAP25 (Fig. 2E), whereas expression of WT EGFP-SNAP25b restored survival (Delgado-Martinez et al., 2007; Ruiter et al., 2019; Weber et al., 2010). Both mutations (V48F and D166Y) caused rescue of survival; however, the number of neurons per islet was lower than in WT-expressing neurons prepared in parallel. This difference was statistically significant for D166Y, but not for V48F (Fig. 2E). Since patients harbor one mutated and one WT allele, we coexpressed WT EGFP-SNAP25b with mutated EGFP-SNAP25b, by combining infection with separate viruses encoding WT and mutant protein in a 1:1 ratio, as done previously by others for Syt1 (Bradberry et al., 2020). As a prerequisite for this, we demonstrated by Western blot analysis that all mutant

viruses expressed similar amounts of protein as WT viruses (Fig. 2A). In co-expressing neurons, we added half the volume of WT and mutant virus as compared to the WT condition, thus keeping the total amount of virus constant. We preferred to combine two single-cistronic viruses, rather than to construct bicistronic viruses, since the larger insert would be expected to lower the titer of viruses, which might compromise survival rescue of *Snap25* KO neurons. Using co-expression of WT and mutant SNAP25b, neuronal survival was mildly and significantly reduced in the V48F + WT condition from WT (Fig. 2E). Overall, mutations seemed to mildly reduce survivability of neurons. Staining of neurons against VGlut1 (a marker for glutamatergic synapses) and MAP2 (a dendritic marker) revealed no significant difference in the dendritic length or the number of synapses in neurons expressing V48F, and D166Y alone (Fig. 2B-D).

Patch-clamp was performed on days 10-14 on autaptic neurons expressing either the WT EGFP-SNAP25b, the mutants, or both the mutant and WT in a 1:1 ratio. Spontaneous miniature events were recorded at a holding voltage of -70 mV. Both primary interface mutations (V48F, D166Y) strongly increased the frequency of mEPSCs (Fig. 3A-B, D-E); the mEPSC amplitude was significantly increased in the V48F, and insignificantly (p=0.14) increased for the D166Y (Fig. 3C, F). The mEPSC frequency was at least as high as in *Syt1* KO neurons measured in separate experiments (Fig. 3G-I). V48F and D166Y co-expressed with WT resulted in mEPSC frequencies close to the arithmetic mean between frequencies in WT and the mutant alone (Fig. 3B, E), indicating that the mutations are incompletely dominant, or co-dominant.

Brief depolarization elicits an unclamped action potential in the axon, which makes it possible to study evoked release, which is essentially absent in *Snap25* KO neurons (Bronk et al., 2007; Delgado-Martinez et al., 2007). The V48F and D166Y mutants both supported evoked release, but the evoked EPSC (eEPSC) amplitude was significantly reduced in the mutant condition (Fig. 4A-B, E-F), whereas when co-expressed with the WT, only the D166Y mutant had a significantly reduced EPSC (Fig. 4B,F). Integration of individual eEPSCs allowed determination of the total charge and the assessment of synchronous and asynchronous release components (Fig. 4 – Figure supplement 1). The eEPSC charge was significantly reduced in the D166Y, but not in the V48F (Fig. 4C,G), whereas in both cases the kinetics was significantly shifted in the direction of more asynchronous release, and the fast time constant was prolonged (Fig. 4D,H; Fig. 4 - Figure supplement 2). We compared the data obtained from V48F and D166Y SNAP25 to *Syt1* KO neurons recorded in separate experiments. *Syt1*

WT neurons in this set of experiments had larger EPSC amplitudes, and a larger charge than WT-
rescued *Snap25* KO neurons (Fig. 4I,J,K), which might be caused by differences between cell cultures,
or animal lines (*SNAP25* vs *Syt1* lines). Note that our experiments using WT, mutant and WT +
mutant SNAP25 was always carried out in neurons prepared and recorded in parallel from the same
SNAP25 KO embryos (Materials and Methods). In the *Syt1* KO, the kinetic change was similar to the
V48F and D166Y (Fig. 4L), but the total charge was also strongly reduced (Fig. 4K), unlike the two
mutations that displayed at most a mild reduction (Fig. 4C,G).

Overall, D166Y and V48F caused a strong disinhibition of spontaneous release and a
desynchronization of evoked release, as demonstrated before (Alten et al., 2021), and consistent
with previous mutational studies of the primary interface, both in SNAP25 (Schupp et al., 2016) and
Syt1 (Zhou et al., 2015). Accordingly, these phenotypes are similar to *Syt1* KO, but the effect of V48F
and D166Y on total evoked charge was milder than in the *Syt1* KO. The preserved overall charge of
evoked release in V48F and mild reduction in D166Y might point to a compensatory gain-of-function
aspect to these mutations, in addition to the impaired Syt1 interaction.

V48F and D166Y mutations lower the energy barrier for release and reduce the readily releasable
pool size

The role of Syt1 and Syt1:SNARE interactions in vesicle priming has been controversially discussed.
The RRP is often assessed by applying a pulse of hypertonic solution to the neurons, usually 0.5 M
sucrose (Rosenmund and Stevens, 1996; Schotten et al., 2015). However, in some experiments this
did not lead to a change in RRP size in the absence of Syt1 (Bacaj et al., 2015; Xu et al., 2009),
whereas experiments both from our laboratories and others showed a decrease in RRP in the
absence of Syt1 (Bouazza-Arostegui et al., 2022; Chang et al., 2018; Courtney et al., 2019; Huson et
al., 2020; Liu et al., 2009; Ruiter et al., 2019). The reason for this discrepancy is likely partly technical
and has to do with how fast sucrose can be applied to the dendritic tree (see also Discussion). Using
neurons growing on small autaptic islands makes it possible to apply sucrose to the entire dendritic
tree (which is confined to the 30-50 µm island) within tens of milliseconds, which is fast enough to
dissect the RRP with a short (few s) sucrose application. We set out to understand whether the V48F
and D166Y mutants changed the size of the RRP.

Application of 0.5M sucrose to neurons with the V48F or D166Y mutation resulted in estimates of

the RRP (denoted $RRP_{0.5}$) that were significantly reduced compared to the WT condition (Fig. 5

A,C,F,H). Also the co-expressed condition displayed significantly reduced $RRP_{0.5}$. Application of two

different sucrose concentrations is used to probe the size of the energy barrier for release (Basu et

al., 2007; Schotten et al., 2015), because the use of a lower sucrose concentration (typically 0.25 M)

will only release a fraction of the RRP, and this fraction depends sensitively on the energy barrier

for release; the size of the RRP at 0.5M sucrose ($RRP_{0.5}$) is used for normalization. Application of 0.25

M sucrose to V48F and D166Y expressing neurons strikingly led to unchanged pool release ($RRP_{0.25}$)

(Fig. 5B,G). Consequently, the ratio of pools ($RRP_{0.25}/RRP_{0.5}$) was significantly increased for both the

V48F and the D166Y mutations (Fig. 5D,I), indicating that these two mutations decrease the

apparent energy barrier for fusion (Schotten et al., 2015). The co-expressed conditions were in-

between WT and mutant and did not reach statistical significance. Consistent with these results,

both mutations on average increased the release probability, calculated as the ratio between the

eEPSC charge and the $RRP_{0.5}$ charge measured in the same cell. The increase was statistically

significant for the D166Y mutation (Fig. 5J), but not for V48F (Fig. 5E). This is different from the

situation in the *Syt1* KO, where the $RRP_{0.25}/RRP_{0.5}$ and the release probability were both significantly

decreased (Fig. 5K-O), although in this data set the reducint in $RRP_{0.5}$ size by by removing Syt1 did

not reach statistical significance (P=0.0548, unpaired t-test). Thus, the increased mEPSC release rate

from the *Syt1* KO does not correlate with a reduced energy barrier as assayed by sucrose, which was

reported before (Huson et al., 2020), and the V48F and D166Y have specific gain-of-function

features that lower the energy barrier for vesicle fusion.

To investigate the reasons for the change in RRP-size by D166Y and V48F, we considered a one-pool

model for the RRP (Fig. 6A), where the RRP is filled by priming (k_1) from an upstream pool and

depleted by de-priming (k_{-1}) or fusion (k_f). Dividing the miniature release rates (r_{mini}) with the size of

the RRP yields the spontaneous fusion rate k_f. The current plateau during 0.5 M sucrose application

(Fig. 6B) essentially reports on the priming rate (k_1), providing that the fusion rate is sufficiently

increased by sucrose (Materials and Methods). However, high sucrose concentrations can change

the baseline current level, which will cause an error in estimation of k_1. We therefore corrected the

plateau level using a plot of the variance versus mean during the sucrose application (Fig. 6C); this

plot is linear for the type of noise generated by synaptic transmission (shot noise). Back-

extrapolation of a regression line allows a determination of the baseline current level in the

presence of sucrose (Fig. 6B-C; see Materials and Methods). Combining RRP size with the estimates

of k_1 and k_f allows determining the depriming rate, k_{-1}.

These calculations showed that k_f, the spontaneous fusion rate was strongly increased in the V48F

and D166Y, and this increase was much larger for the mutations than for the *Syt1* KO (Fig. 6D)

fulfilling D166Y>V48F>*Syt1* KO. Further analysis showed that in both mutations, the forward priming

rate, k_1, and the depriming rate, k_{-1}, were both decreased – the latter effect was only significant for

D166Y (Table 1). Summing up the effects of changes in the three parameters on the RRP-size for the

V48F, D166Y and *Syt1* KO (Fig. 6E), we can conclude that for the V48F and even more for the D166Y,

spontaneous release contributes to the reduction in RRP size, whereas this effect is minimal in the

Syt1 KO. However, the major reason for the smaller RRP size is a reduction in priming rate, k_1, which

is partly counteracted by the decrease in k_{-1} (which would increase RRP size). Overall, changes in

priming, depriming and spontaneous fusion rates combine to change RRP size.

Repetitive stimulation to determine the RRPs often results in lower estimates than sucrose

application, because action potentials draw on a sub-pool of the RRP (Moulder and Mennerick,

2005), whereas sucrose releases the entire RRP (Rosenmund and Stevens, 1996). To determine the

RRP sub-pool that evoked release draws on (denoted RRP_{ev}) we applied repetitive stimulation (50

APs @ 40 Hz) and used back-extrapolation to determine the RRP (Neher, 2015). Performing this in

our standard 2 mM Ca^{2+}-containing extracellular solution resulted in overall smaller estimates for

RRP_{ev} for V48F and D166Y compared to WT (Fig. 7 - Figure supplement 1); however, the differences

were not statistically significant. The back-extrapolation method works best with high release

probabilities (Neher, 2015); therefore, we repeated these experiments in the presence of 4 mM

extracellular Ca^{2+}. Under these conditions, the RRP_{ev} was significantly reduced for both the V48F

and D166Y mutation (Fig. 7C,G; Fig. 7 - Figure supplement 2). Strikingly, the release probabilities

(calculated as the charge of the first EPSC of the train divided by the RRP_{ev}) were decreased for both

mutations (Fig. 7D,H), which correlated with an increased paired-pulse ratio (Fig. 7A,E, *inserts)*

(Zucker and Regehr, 2002). Thus, although the release probability calculated by normalizing evoked

charge to the sucrose-determined RRP ($RRP_{0.5}$) was increased (see above), the release probability

when normalizing to RRP_{ev} was decreased. This points to a difference in the organization of the

RRP_{ev} and $RRP_{0.5}$ (see Discussion). The forward priming rate is determined as the slope of the linear

fit used for the back extrapolation; this parameter was reduced in both mutations (Fig. 7B, F; Fig. 7

- Figure supplement 2). Thus, the difference in priming rate extends to the RRP_{ev}.

Overall, these data demonstrate a rather complex phenotype of the D166Y and V48F mutations,

which combine a lowering of the energy barrier – a gain-of-function feature – with a loss of vesicle

priming – a loss-of-function feature. The D166Y and V48F phenotypes can be summarized as 1)

desynchronized EPSCs with at most mildly reduced total charge, 2) lowered energy barrier for

fusion, 3) increased release probability when normalized to the sucrose pools, 4) decreased RRP-

size due to unclamped spontaneous release and lowered forward priming rates, 5) short-term

facilitation. These phenotypes are distinctive from the Syt1 KO, which does not have a preserved

charge of the EPSC, or lowered energy barrier when probed by sucrose, or an increased release

probability. Thus, the D166Y and V48F cannot be understood solely in terms of a lack of Syt1

coupling; instead, gain-of-function features are present in the mutants which are absent upon

deletion of Syt1.

V48F and D166Y mutants show increased partner SNARE interactions and cause unregulated fusion

in vitro

We next tried to identify the biochemical properties of V48F and D166Y, which could support a gain-

of-function phenotype during exocytosis. These data are displayed in Fig. 8-9 - data on the I67N

mutant were obtained in parallel and will be presented later. To test to which degree the mutants

may change the stability of SNARE complexes, full-length t-SNAREs (syntaxin-1 and SNAP25) were

incubated with the cytosolic domain of VAMP2 (VAMP2cd) overnight. Cis-SNARE complex stability

was tested in the presence of SDS at the indicated temperatures (Fig. 8A), and the release of

syntaxin-1 as a single protein band was used as a measure of the complex dissociation. The wildtype

(WT), V48F and D166Y v-/t-SNARE complexes showed a similar stability with half-maximal

dissociation occurring at approximately 71°C.

Next, we asked whether the V48F and D166Y mutants would change interaction with Syt1. To this

end, Atto647 labelled Giant Unilamellar Vesicles (GUVs) filled with isosmotic sucrose and containing

preassembled t-SNARE complexes were preincubated with Atto488/Atto550 labeled Small

Unilamellar Vesicles (SUVs) containing Syt1 as well as VAMP2 for 10 minutes on ice, followed by

centrifugation to reisolate GUVs with attached SUVs. Fusion was blocked by performing the assay

on ice. Attachment of SUVs was determined by measuring the Atto550 fluorescence of SUVs. In the

absence of PI(4,5)P$_2$, vesicle attachment occurs by Syt1:SNARE interactions (Kim et al., 2012;

Parisotto et al., 2012), probably involving the primary interface (Zhou et al., 2015). In the cell,

PI(4,5)P$_2$-binding by Syt1 appears to happen first (Honigmann et al., 2013), whereas subsequent

Syt1:SNARE-binding leads to a tightly docked state (Chen et al., 2021). Under our conditions, both

SNAP25 mutants (V48F and D166Y) showed significantly impaired attachment of Syt1/VAMP2 SUVs

to t-SNARE GUVs (Fig. 8B). Both V48 and D166 directly interact with Syt1 (Fig. 1B-C, (Zhou et al.,

2015)), so that changing these two amino acids to more bulky or hydrophobic amino acids reduced

the docking from 42.7 ± 2.6 % wildtype (WT) docking efficiency to 29.6 ± 4.1 % and 20.9 ± 3.7 %,

respectively. In the presence of 1% PI(4,5)P$_2$, the vesicle attachment increased from 42.7 ± 2.6 % to

66.0 ± 0.5 % for WT t-SNARE. Although the Syt1:PI(4,5)P$_2$ interaction predominated SUV-GUV

docking, both mutants still showed significantly reduced vesicle attachment by approximately 6%

compared to WT (Fig. 8C) (V48F: 59.1 ± 1.4 %, and D166Y: 60.0 ± 2.0 %).

To understand how the mutations affect fusion in a well-defined reconstituted membrane fusion

system, we performed *in vitro* lipid mixing assays using GUVs containing both t-SNAREs (syntaxin-1

and SNAP25) with 1% PI(4,5)P$_2$, and 0.5% Atto488/0.5% Atto550 labeled SUVs containing Syt1 and

VAMP2 in the presence or absence of complexin-II (6 µM) (Kedar et al., 2015; Malsam et al., 2012).

Fusion was measured at 37°C by Atto488 fluorescence dequenching, which occurs upon lipid mixing

with GUVs. Calcium (100 µM free Ca^{2+} final) was added after 2 minutes to the t-SNARE GUV assay.

Measurements were continued for another 5 minutes. SUVs treated with botulinum toxin D, which

cleaves VAMP2 and abolishes membrane fusion, served as negative control and the corresponding

background fluorescence was subtracted. Measurements were normalized to total fluorescence

after detergent lysis.

Complexin-II clamps spontaneous Ca^{2+}-independent membrane fusion in the reconstituted assay

(i.e. fusion before addition of Ca^{2+}), via laterally binding the membrane-proximal C-terminal ends of

SNAP25 and VAMP2 (Malsam et al., 2020). Remarkably, V48F and D166Y showed impaired clamping

by complexin, as apparent by increased fusion before Ca^{2+} addition (Fig. 8D). The decreased

clamping is likely caused by the reduced interaction of V48F and D166Y with Syt1. As a note, the

clamping function of Syt1 becomes only obvious in the presence of complexin, whereas the

clamping function of complexin depends on the presence of Syt1 (Malsam et al., 2012); thus, the

clamping function of the two proteins cannot be separated. After Ca^{2+}-triggering, WT SNAP25

supported the largest amount of fusion, followed by V48F, whereas D166Y Ca^{2+}-dependent fusion

was clearly reduced (Fig. 8D). Notably, this is the same sequence as found for evoked release in

synapses, considering either the amplitude or the charge of the EPSC (Fig. 4B-C, F-G). Thus, the *in*

vitro assay reproduces both the increased spontaneous release and the reduced Ca^{2+}-dependent

release found in neurons, indicating that these features are present within the minimal set of fusion

proteins included in this assay (i.e. the SNAREs, Syt1 and complexin).

There is evidence that SNAP25 might enter the SNARE-complex last, after syntaxin-1 and VAMP2

are joined by Munc18-1 (Baker et al., 2015; Jiao et al., 2018; Sitarska et al., 2017), although another

view is that a syntaxin/SNAP25 dimer bound to Munc18-1 acts as an intermediary (Jakhanwal et al.,

2017). In the former case, mutations changing association of SNAP25 to the SNAREs might change

exocytosis efficiency. To test this, we used a syntaxin-1 GUV assay, where the incorporation of

soluble SNAP25 into the SNARE complex becomes a rate-limiting step. In this assay, fusion kinetics

are much slower when compared with fusion reactions containing GUVs with pre-assembled t-

SNAREs. Accordingly, we allowed 30 min for pre-stimulation fusion to take place, and after addition

of Ca^{2+} (100 μM free Ca^{2+}), fusion was followed for another 30 min (Fig. 9A). The assay was

performed in the presence and absence of complexin-II. D166Y and to a lesser degree V48F revealed

enhanced stimulation of fusion in comparison to WT before calcium was added, regardless of the

presence or absence of complexin (Fig. 9B). Comparing data with and without complexin established

that complexin barely suppressed spontaneous fusion in reactions containing the V48F mutant. The

presence of complexin partially reduced Ca^{2+}-independent fusion, but D166Y still stimulated pre-

Ca^{2+} fusion compared to wildtype. Overall, these data demonstrate that D166Y and V48F are gain-

of-function mutants under conditions where SNAP25 association to the other SNAREs is rate

limiting.

To test directly whether V48F and especially D166Y enhance SNARE interactions, co-flotation assays

were performed. SUVs containing either syntaxin-1 (Stx-1), or VAMP2, or Syt1, or the Syt1/VAMP2

combination were incubated with SNAP25 and re-isolated by flotation using a Nycodenz density

gradient. An additional reaction, reflecting the fusion assay, contained SNAP25 in combination with

both Syt1/VAMP2 SUVs and Syntaxin-1 SUVs (Fig. 9D). SNAP25 recruitment for each condition was

determined by SDS PAGE followed by Coomassie blue and silver staining. Silver stained SNAP25

bands were quantified, and the mutants were plotted relatively to the wildtype SNAP25 (Fig. 9D and

Fig. 9 – Figure supplement 1).

SNAP25 WT did not show any binding to Syt1 SUVs (Fig. 9 – Figure supplement 1) , although direct

interactions with Syt1 would be predicted based on the primary interface, indicating that such

interactions are not stable under the employed conditions, which is expected because SNAP25 is

unstructured until it binds its SNARE partners (Fasshauer et al., 1997). D166Y showed profoundly

increased interactions with SUVs containing either syntaxin-1, or VAMP2, or Syt1/VAMP2, and the

combination used in the fusion assay (Fig. 9D; Fig. 9 – Figure supplement 1). V48F displayed mildly

increased binding to syntaxin-1 and Syt1/VAMP1 SUVs, and a tendency to increased association to

VAMP2 SUVs (p=0.0629, one-sample t-test). These data show that loss of Syt1 interaction upon

mutation in the primary interface can be accompanied by a gain-of-function phenotype stimulating

interactions with the other SNARE-partners. This association of SNAP25 to the other SNAREs might

happen as one of the last steps towards fusion; consequently, when D166Y and V48F join the

complex prematurely, it will bypass layers of control and result in uncontrolled fusion.

The I67N mutation supports an intact RRP, but an increased energy barrier for fusion

We next addressed the phenotype of the I67N disease mutation, which was found in an 11-year-old

female, who suffered from myasthenia, cortical hyperexcitability, ataxia and intellectual disability,

but with normal brain MRI (Shen et al., 2014). The I67 is found within the interaction layer +4

(Fasshauer et al., 1998), which helps in assembly of the C-terminal of the SNARE-complex (Gao et

al., 2012), and it might therefore have a different synaptic phenotype than V48F and D166Y.

In *in vitro* experiments, SNARE-complexes formed with the I67N mutant displayed a lower stability,

with the melting temperature reduced from 71°C to approximately 56°C (Fig. 8A), as expected for a

mutation that destabilizes the SNARE-complex. I67N also showed a strong decrease in SUV docking

(Fig. 8B-C), which is likely caused by the destabilization of the t-SNARE complex, which indirectly

perturbs the Syt1 binding interface(s). In lipid mixing assays, I67N strongly reduced both Ca^{2+}-

independent and Ca^{2+}-dependent fusion, whether the t-SNAREs were preassembled (Fig. 8D-E) or

not (Fig. 9B-C). The co-flotation assay did not display any binding of I67N to t-SNAREs or Syt1 (Fig.

9D). The binding to Syt1/VAMP2 SUVs was reduced compared to SNAP25 WT, but since binding of

the WT protein is already very low, the biological significance of this result is unclear. Overall, these

data indicate that I67N is inferior in membrane fusion.

Lentiviruses encoding the I67N mutant N-terminally fused to EGFP expressed similar amounts as WT

EGFP-SNAP25 (Fig. 10A). Expression in SNAP25 KO neurons resulted in reduced rescue of survival in

neurons expressing I67N alone, whereas neurons co-expressing WT and I67N had intermediate

survival, not significantly different from WT (Fig. 10C). Staining against MAP2 (dendritic marker) and

VGlut1 (synapse marker) showed that the number of synapses on average was reduced in the I67N

(Fig. 10B,D), and the dendritic length was on average reduced, but the changes did not reach

statistical significance (Fig. 10E). Patch-clamp measurements demonstrated strongly reduced

spontaneous release frequencies (Fig. 10F-H) and evoked release amplitude (Fig 10I-L) with the I67N

mutation, see also (Alten et al., 2021). mEPSCs were absent in most I67N expressing neurons,

whereas WT and I67N co-expressing neurons had a very low mEPSC rate, much closer to the I67N

than the WT phenotype; similar for EPSC amplitudes (Fig. 10I-L). The I67N therefore is dominant-

negative for both types of release, in contrast to the incompletely dominant phenotypes of the V48F

and I166Y mutants (see above). The fraction of synchronous release was unchanged in WT and I67N

coexpressing neuron (Fig. 10L); this number could not be estimated for I67N-expressing neurons

due to the low amount of release.

Reduced spontaneous and evoked release could result from a decrease in priming, or fusion, or

both. To distinguish between these possibilities, we turned to sucrose applications. Application of

0.25M sucrose did not lead to any measurable release in I67N expressing cells, and only minimal

release in cells co-expressing WT and I67N (Fig. 11A-D). This indicates that the energy barrier is

increased in amplitude, and therefore the RRP might be underestimated when probed by 0.5M

sucrose (Schotten et al., 2015). Indeed, 0.5M sucrose displayed reduced release in the I67N and

WT+I67N condition (Fig. 11C), but this could be due to defects in priming or fusion. To investigate

this, we applied a stronger stimulus, 0.75M sucrose, to these cells (Schotten et al., 2015). Strikingly,

the RRP as assessed by 0.75M sucrose ($RRP_{0.75}$) was unchanged between WT, I67N and WT+I67N co-

expressed cells (Fig. 11E,G). Application of 0.375M sucrose led to small amounts of release in the

I67N, but more in the WT+I67N co-expressed situation (Fig. 11E,F). Forming the ratio

$RRP_{0.375}/RRP_{0.75}$ revealed a statistically significant reduction in I67N compared to WT (Fig. 11H).

Therefore, the RRP *per se* appears intact in the I67N (when probed by sufficiently high

concentrations of sucrose), but the vesicles face a higher apparent fusion barrier, which makes the

RRP appear smaller if assessed by 0.5M sucrose. The higher apparent fusion barrier explains the

lower frequency of mEPSC in the I67N mutation, the lower degree of spontaneous fusion in *in vitro*

assays, as well as the lower amount of Ca^{2+}-dependent release in vitro and in the cell.

The I67N mutation profoundly affected trains in 2 mM extracellular Ca^{2+} (Fig. 11I) leading to strong

facilitation, which is expected due to the strong phenotype of this mutation, which radically lowers

release probability. Even when co-expressed with WT protein, the train facilitated over the first

several stimulations, attesting to the strong dominant-negative feature of the I67N mutation (Fig.

11I-J). Consequently, the paired-pulse ratio was increased in the I67N and intermediate in the

WT+I67N condition (Fig. 11K). Back-extrapolation of these trains was not reliable, because the low

release probability in the I67N mutation made it impossible to achieve sufficient depletion of the

RRP.

The energy barrier for fusion is exquisitely sensitive to the charges on the surface of the SNARE-

complex, with positive charges decreasing and negative charges increasing the fusion barrier

amplitude (Ruiter et al., 2019). To investigate whether the same electrostatic mechanism applies to

the I67N, we combined the I67N mutation with a mutation of four amino acids ("4K"=SNAP25

E183K/S187K/T190K/E194K) in the second SNARE-domain of SNAP25, constructing the quintuple

mutation ("I67N/4K"=SNAP25 I67N/E183K/S187K/T190K/E194K). The 4K mutation lowers the

energy barrier for fusion by increasing the charge of the SNARE-complex surface by +6 via charge

introduction and charge reversal (Ruiter et al., 2019). The 4K-mutation increased the mEPSC release

rate compared to WT (Ruiter et al., 2019), whereas in the combined I67N/4K mutation the

spontaneous release rate was indistinguishable from WT (Fig. 12A-C), showing that increased

positive charges rescued the defect of spontaneous release in the I67N mutant. Evoked release was

also increased in the 4K mutation compared to WT (Ruiter et al., 2019), but in the I67N/4K mutation,

evoked release was still strongly depressed compared to WT (Fig. 12D-E). Nevertheless, evoked

release was noticeable in the combined mutation, whereas it was almost absent in I67N-expressing

cells (comp. Fig. 10I-L), indicating a positive effect of the 4K mutation, which amounted to an

increase by a factor ~5 (eEPSC amplitude, I67N: 0.0475 ± 0.0087; I67N/4K: 0.2668 ± 0.12 nA; Mann-

Whitney test, P=0.035).

We previously created a simple mathematical model that links the release rate to the number of

charges added to the SNARE-complex (Ruiter et al., 2019). This model includes both spontaneous

and evoked release, which are separated by the addition of 35 positive charges in the latter case

(Fig. 12F, black points are WT spontaneous and evoked release rates; blue line is the model fitted to

WT data). Placing the spontaneous release rates for I67N and I67N/4K on this curve (by dividing the

spontaneous release rate with RRP size and finding a corresponding charge-value using the model)

resulted in two points (red) separated by 5.6 charges, which is close to the nominal 6 charges added

by the 4K mutation (Fig. 12F). Similarly, evoked release in the I67N and the I67N/4K were separated

by 5.9 charges (Fig. 12F). The fact that these numbers are close to the nominal 6 charges introduced

shows that even for the I67N mutation, the same basal electrostatic model still applies, but the

deleterious effect of the I67N on evoked release rates is larger than on spontaneous rates and

therefore the rescue of evoked release rate by positive charges is insufficient to reach WT values.

Moreover, because of the saturating form of the curve (i.e. the model) adding positive charges is an

effective way of rescuing spontaneous, but not evoked release.

Overall, the I67N disease mutation increases the amplitude of the energy barrier for fusion, and it

does so more for evoked than for spontaneous release, but the electrostatic mechanism, which we

assume is part of release triggering (Ruiter et al., 2019), appears to be intact.

Discussion

We have shown that two SNAP25 mutations (V48F, D166Y) that compromise interaction with Syt1

lead to complex phenotypes characterized by a combination of loss-of-function and gain-of-function

features. Thereby, the mutations fall into the 'neomorph' category, where the mutated protein has

novel or changed interactions or functions (Verhage and Sorensen, 2020). In contrast, the I67N

substitution within the SNARE-bundle is a dominant-negative mutation.

SNAP25 disease mutations change protein-protein interactions and the energy landscape of fusion

Both the V48F and the D166Y resulted in a decrease in the amplitude of the apparent energy barrier

for fusion, whereas the I67N increased the amplitude of the apparent fusion barrier. A vesicle's

release willingness can only be assessed by fusing it; therefore, it is not possible to distinguish

between effects on the fusion barrier *per se*, and effects on the fusion machinery. In recognition of

this fact, we here refer to the "apparent energy barrier". Using Arrhenius' equation to convert

fusion, priming and depriming rates to their respective energy barrier heights (see Materials and

Methods for the assumptions behind this procedure), we can derive the energy landscape for fusion

of the three mutants (Fig. 13A-D). This shows the multiple changes in the V48F and D166Y, which

affect at least two different barriers (priming and fusion, Fig 13A-B), leading to a complex

phenotype, whereas for I67N the fusion barrier is primarily (or solely) affected (Fig. 13D).

When combining the I67N with the 4K-mutation, which introduces 6 extra positive fixed charges,

we could place our data within the framework of our model for electrostatic triggering (Ruiter et al.,

2019) and show that the effect of charge *per se* is approximately the same in the I67N mutant as in

the WT. Note that there are endogenous positively charged amino acids towards the C-terminal end

of SNAP25 that are important for release rates (Fang et al., 2015). Rescue of spontaneous release

was completed by adding 6 positive charges, which is consistent with the idea that the assembly of

the C-terminal end of the SNARE-complex, which is compromised by I67N (Rebane et al., 2018),

works against the electrostatic energy barrier, which is affected by the SNARE surface charge (Ruiter

et al., 2019). In contrast, rescue of evoked release by charges were incomplete, due to the larger

effect of I67N on evoked release, combined with the shallow effect of charges on evoked release

(Ruiter et al., 2019). The larger susceptibility of evoked release to C-terminal mutation of the

SNAREs might be partly due to the higher number of SNARE-complexes involved in evoked than in sustained/spontaneous release (Mohrmann et al., 2010).

The effect of the I67N mutation in the energy domain at rest can be calculated from the spontaneous release rate, which was reduced by a factor 22.4 (from 1.31 ± 0.36 to 0.0583 ± 0.0274 Hz). This corresponds to an effect in the energy domain of 3.1 k_BT (where k_B is Boltzmann's constant; assuming unchanged RRP size, pool normalization is not required). Work with single-molecular optical tweezers showed that the I67N mutation destabilizes the overall SNARE C-terminal and linker domain, which are supposed to deliver the power stroke for membrane fusion, by 14 k_BT (Rebane et al., 2018), which is substantially more. The reduction in spontaneous release rate is more comparable to the reduction in the transition rate of folding by the C-terminal and linker domain by a factor of ~10 (Rebane et al., 2018). Since at least three SNARE-complexes, possibly more, contribute to vesicle fusion (Bao et al., 2018; Manca et al., 2019; Mohrmann et al., 2010; Shi et al., 2012), folding kinetics correlates better to spontaneous fusion rates than overall SNARE-complex stability.

The D166Y and V48F mutations lead to increases in spontaneous release, and more asynchronous EPSCs, consistent with their localization in the primary SNARE:Syt1 interface (Schupp et al., 2016; Zhou et al., 2015; Zhou et al., 2017), and the demonstrated impaired Syt1 binding (Fig. 8B-C), see also (Alten et al., 2021). These phenotypes are at first glance similar to Syt1 knockout/knockdown (Bouazza-Arostegui et al., 2022; Chang et al., 2018; Huson et al., 2020; Ruiter et al., 2019). However, when normalized to the RRP size, mEPSC frequencies were much higher in the V48F and the D166Y than in the *Syt1* KO (Fig. 6D), and dual sucrose applications indicated a decrease in the amplitude of the sucrose-probed apparent energy barrier and increased release probability, features not found in the *Syt1* KO (Bouazza-Arostegui et al., 2022; Huson et al., 2020). This indicates a gain-of-function feature of these mutations, which fulfills D166Y>V48F; such a feature was identified as an increased interaction of V48F and D166Y with VAMP2- and syntaxin-1-containing SUVs (Fig. 9D); the interaction was stronger for D166Y than for V48F. AlphaFold prediction of the SNAP25 mutants did not reveal major structural changes in the SNAP25 backbone. However, the replacement of D166 with Y shows stronger interactions with the neighboring H162 and V48F alters the interactions with the neighbouring E52 (Fig. 9 - Figure supplement 2; (Jumper et al., 2021; Mirdita et al., 2022)). This may result in stabilization of a structure consistent with SNARE-complex formation. However, note

that the AlphaFold prediction is identical to the structure of SNAP25 in the assembled SNARE-complex (Zhou et al., 2015), whereas unassembled SNAP25 is likely less structured or unstructured (Fasshauer et al., 1997). Overall, these disease mutants do not only fail in their interaction with Syt1, they bypass fusion control, resulting in premature SNARE-complex assembly. Although the increased spontaneous release rate might cause patient symptoms, Alten et al. (2021) suggested that the resulting postsynaptic depolarization might compensate for the smaller EPSC amplitude to normalize the overall firing rate.

In a SUV:GUV fusion assay, where folding of SNAP25 onto syntaxin-1/VAMP2 is rate-limiting, D166Y and V48F caused an increase in pre-stimulation fusion rates. This is consistent with recent data showing that folding of SNAP25 onto a template formed by VAMP2 and syntaxin-1 held in place by Munc18-1 might be a late, rate-limiting, step in exocytosis (Jiao et al., 2018). This process is regulated and sped up by Munc13-1 (Kalyana Sundaram et al., 2021; Shu et al., 2020; Wang et al., 2019). The fact that a similar effect on spontaneous fusion was seen in the assay with preformed t-SNAREs in the presence of complexin indicates that there is an assembly step, even in that assay, which can be sped up by the mutations. This aligns with the demonstration by single molecule FRET of a further assembled ('tighter') state of the trans-SNARE-complex induced by Ca^{2+}-unbound Syt1, which becomes committed for fusion once Ca^{2+} binds (Das et al., 2020).

V48F and D166Y change the size of the RRP via effects on priming, depriming and fusion

We found that V48F and D166Y cause a decrease in RRP size, whether measured by sucrose application or by train stimulation. The reduced RRP is at variance with the publication by Alten et al., who used sucrose application to larger mixed cultures (Alten et al., 2021). There might be two reasons for this discrepancy. First, Alten et al. used longer duration of sucrose application (~1 min) and applied sucrose to a large mixed culture, which might result in a variable delay such that all synapses are not stimulated simultaneously. This will result in temporally overlapping release of RRP and upstream vesicle pools, which then cannot be distinguished from each other. Indeed, the RRP size reported by Alten et al. is ~10-fold larger than RRPs measured from autaptic neurons. When grown on 50-μm micro-islands, sucrose can be applied acutely (within ~0.05 s) to the entire dendritic tree and all synapses using a local perfusion system, which allows distinguishing the RRP from upstream pools. Second, it was recently shown that sufficient neuronal maturation is

necessary to detect the decrease in RRP upon Syt1 elimination (Bouazza-Arostegui et al., 2022),

which is likely modulated by the presence of other neurons (Chang et al., 2018; Liu et al., 2009;

Wierda and Sorensen, 2014). This adds an additional layer of complexity, since neuronal maturation

likely varies between laboratories, or even over time within the same laboratory.

Dissection of the three rates that determine RRP size (priming, depriming and fusion rates) showed

that all three are changed in the V48F and D166Y mutations. Especially for D166Y, but also for V48F,

spontaneous release contributed to RRP depletion by triggering premature fusion. Significant RRP

depletion has not been expected for moderate increases in mEPSC frequency (Rhee et al., 2005;

Ruiter et al., 2019), based on the argument that RRP refilling should be fast enough to counteract

depletion. However, to properly make this argument, priming, depriming and spontaneous release

rates must be compared in the steady-state situation. Similarly, we recently showed by cryo-

electron tomography that the 4K-mutation, which increased the mEPSC frequency to ~30 Hz and

had a reduced RRP (Ruiter et al., 2019) caused a loss of synaptic vesicles tethered to the membrane

with three tethers (Radecke et al., 2023), which is the structural correlate of the RRP (Fernandez-

Busnadiego et al., 2010).

The major effect on RRP size is caused by a reduction in forward priming rate by D166Y and V48F.

Comparison to the Syt1 KO showed qualitatively similar changes, but of a smaller magnitude, with

spontaneous release playing a negligible role for RRP size. Notably, in all three cases a reduction in

depriming rate partly counteracted the lowered priming rate (Table 1) – this was only significant for

D166Y. Indeed, part of the role of Syt1:SNARE interaction might be catalytic, lowering the energy

level of a transition state along the path to priming, which will affect both rates (Walter et al., 2013).

This might happen because transient binding to Syt1 might structure SNAP25 and assist in formation

of the SNARE-complex, whereas SNAP25 mutants might prestructure the protein, bypassing the

need for Syt1. In an energy diagram (Fig. 13) it becomes clear that vesicle priming and regulation of

spontaneous release are interdependent. Stabilization of the RRP state will both increase RRP size

at rest and reduce spontaneous release, since lowering the energy level of (i.e. stabilizing) RRP

vesicles will increase the size of the energy barrier that the RRP vesicles face (Fig. 13). Consistently,

downregulation (clamping) of spontaneous release and upregulation of evoked release are often

interdependent under conditions where Syt1 expression level, or Syt1 interaction with the SNAREs,

are up or down regulated (Courtney et al., 2021; Courtney et al., 2019; Vevea and Chapman, 2020;

Zhou et al., 2015; Zhou et al., 2017). This does not rule out the existence of mutations that can affect

one mode of release more than the other (e.g. the I67N/4K mutation). By inference, assembly of

the primary Syt1:SNARE interface (Zhou et al., 2017) is most likely involved in both clamping release

and setting up a RRP. In further support of this, the minimal *in vitro* assay with preassembled t-

SNARE dimers displayed a qualitatively similar reduction in calcium-dependent release, with D166Y

being more impaired than V48F (Fig. 8D). Overall, the energetic contribution of Syt1:SNARE

interaction to Ca^{2+}-triggered release is a stabilization of upstream steps, and the increase of the

fusion barrier downstream of the RRP (Fig. 13). The electrostatic nature of the fusion barrier (Ruiter

et al., 2019) ensures its rapid dissolution by Ca^{2+}, possibly by unbinding of Syt1 from the SNARE

complex (Voleti et al., 2020).

For the I67N-mutation, the sucrose-RRP was also reduced in size when using 0.5 M sucrose, which

is consistent with previous observations (Alten et al., 2021). However, when the energy barrier for

release is increased, 0.5 M sucrose is insufficient to deplete the RRP (Schotten et al., 2015).

Accordingly, 0.75 M sucrose released an RRP of similar size in WT, I67N and I67N+WT co-expressing

cells, showing that the vesicle priming reaction is intact, but the vesicles face a larger fusion barrier

(Fig. 11). In this mutant, train stimulations in the I67N or co-expressed WT + I67N case resulted in a

phenotype quite distinct from the WT, with strong facilitation throughout the train. Thus, the I67N

is a strongly dominant-negative mutant, whether considering mEPSC frequency, eEPSC amplitude

or train stimulations, whereas V48F and D166Y are incompletely dominant. This can be explained

by the different function of SNAP25 domains during fusion, where V48 and D166 help set up an

arrested primed vesicle state by interacting with Syt1 (Schupp et al., 2016; Zhou et al., 2015; Zhou

et al., 2017), whereas I67N participates in the final conformational change, the 'stroke' that leads

to assembly of the C-terminal end of the complex and the linker domain. Due to their defect in

priming, V48F and D166Y might not enter the super-complex (the complex of SNARE-complexes

driving fusion) as often as WT protein. In contrast, I67N will readily enter the super-complex and

compromise its function, which will lead to a dominant-negative phenotype due to the multiple

SNARE-complexes involved in fusion (Bao et al., 2018; Mohrmann et al., 2010; Shi et al., 2012).

Back-extrapolation of 40 Hz trains (@4 mM Ca^{2+}) also led to the conclusion that V48F and D166Y

have a reduced RRP_{ev}, the RRP-subpool that action potentials draw on, and a reduced forward

priming rate. Interestingly, the release probability when normalized to the sucrose pool ($RRP_{0.5}$) was

increased non-significantly for the V48F and significantly for the D166Y mutation, but when

considering 40Hz trains, the release probabilities of both mutations were decreased, consistent with

a shift towards facilitation. The latter finding is likely caused by the defective interaction with Syt1,

which leads to suboptimal priming and/or defective super-priming, which is an additional priming

step after entry of the vesicle into the RRP (Lee et al., 2013; Taschenberger et al., 2016). This will

lead to reduction of the first EPSC of a train and thereby a lowered nominal release probability.

However, when comparing the EPSC charge to the $RRP_{0.5}$, the strong reduction in the $RRP_{0.5}$ pool

(especially in the D166Y mutation) accounts for the increase in the overall release probability. This

can be explained if spontaneous release causes a disproportional depletion of vesicles, which are in

the $RRP_{0.5}$, but not in the RRP_{ev}.

Conclusion

SNAP25 disease missense mutations change the function of the protein without compromising its

expression, leading to dominant negative or neomorphic mutations. Missense mutations in the

primary SNARE:Syt1 interface (V48F, D166Y) result in a complex phenotype characterized by loss-

of-function in the priming step and gain-of-function in the fusion step. Missense mutation in the

SNARE-bundle (I67N) leads to an increased amplitude of the energy barrier for fusion. In addition,

disease mutations display inefficient rescue of neuronal survival. Overall, SNAP25 encephalopathy

caused by single missense point mutations presents with interdependent functional deficits, which

must be overcome for successful treatment.

Materials and Methods

Animals

SNAP25 KO C57/Bl6-mice: Heterozygous animals were routinely backcrossed to Bl6 to generate new heterozygotes. The strain was kept in the heterozygous condition and timed pregnancies were used to recover knockout embryos by caesarean section at embryonic day 18 (E18). Pregnant females were killed by cervical dislocation; E18 embryos of either sex were collected and killed by decapitation. Permission to keep and breed *Snap25* and *Syt1* mice was obtained from the Danish Animal Experiments Inspectorate (2018-15-0202-00157) and followed institutional guidelines as overseen by the Institutional Animal Care and Use Committee (IACUC). CD1 outbred mice were used to create astrocytic cultures and mass cultures for Western blotting. Newborns (P0-P2) of either sex were used. Pups were killed by decapitation.

Cell lines

HEK293-FT cells for production of lentiviruses were obtained from the Max-Planck-Institute for biophysical chemistry. The cells were passaged once a week, and they were used between passage 11 and 25 for generation of lentiviral particles. The cells were kept in DMEM + Glutamax (Gibco, cat. 31966047) supplemented with Fetal Bovine Serum (Gibco, cat. 10500064), Pen/Strep (Gibco, cat. 15140122) and Geneticin G418 (Gibco, cat. 11811064) at 37°C in 5% CO_2.

Preparation of neuronal culture

Self-innervating ("autaptic") hippocampal cultures were used (Bekkers and Stevens, 1991). Astrocytes were isolated from CD1 outbred mice (P0-P2). The cortices were isolated from the brains in HBSS-HEPES medium (HBSS supplemented with 1 M HEPES) and the meninges were removed. The cortical tissue was chopped into smaller fragments, transferred to 0.25% trypsin dissolved in DMEM solution (450 ml Dulbecco's MEM with 10% Foetal calf serum, 20000 IU Penicillin, 20 mg Streptomycin, 1% MEM non-essential Amino Acids) and incubated for 15 min at 37 °C. Subsequently, inactivation medium (12,5 mg Albumin + 12,5 mg Trypsin-Inhibitor in 5 ml 10% DMEM) was added, the tissue was washed with HBSS-HEPES, triturated and the cells were plated in 75 cm^2 flasks with pre-warmed DMEM solution (one hemisphere per flask) and stored at 37 °C with 5% CO_2. Glial cells were used after 10 days.

Glass coverslips were washed overnight in 1 M HCl; for an hour in 1 M NaOH and washed with water

before storage in 96% ethanol. Coverslips were first coated by 0.15 % agarose and islands were

made by stamping the coating mixture (3 parts acetic acid (17 mM), 1 part collagen (4 mg/ml) and

1 part poly-D-lysine (0.5 mg/ml)) onto the glass coverslips using a custom rubber stamp. Glial cells

were washed with pre-warmed HBSS-HEPES. Trypsin was added and the flasks were incubated at

37 °C for 10 min. Cells were triturated and counted in a Bürker chamber before plating onto the

glass coverslip with DMEM solution. After 2-5 days, neurons were plated on the islands.

Neurons for autaptic culture were isolated from E18 *Snap25* KO mice. Pups were selected based on

the absence of motion after tactile stimulation and bloated neck (Washbourne et al., 2002); the

genotype was confirmed by PCR in all cases. The cortices were isolated from the brains in the HBSS-

HEPES medium. The meninges were removed and the hippocampi were cut from the cortices before

being transferred to 0.25% trypsin dissolved in HBSS-HEPES solution. The hippocampi were

incubated for 20 min at 37 °C, subsequently washed with HBSS-HEPES, triturated and the cell count

was determined with a Bürker chamber before plating on the islands (7000-8000 neurons per well).

Cells were incubated in the NB medium (Neurobasal with 2% B-27, 1 M HEPES, 0,26% Glutamax,

14,3 mM β-mercaptoethanol, 20000 IU Penicillin, 20 mg Streptomycin) and used for measurements

between DIV10-14.

Hippocampal neurons for high-density cell culture for Western blotting were obtained from P0-P1

CD1 outbred mice. The dissection, tissue digestion and cell counting were performed the same way

as the neurons for the autaptic culture, the high-density culture (600,000 neurons per well) was

then kept in NB-A medium (Neurobasal-A with 2% B-27, 1 M HEPES, 0,26% Glutamax, 20000 IU

Penicillin, 20 mg Streptomycin) and half the volume of the media was replaced every 2-3 days before

harvesting the cells at DIV14.

Constructs for rescue experiments

SNAP25B was N-terminally fused to GFP and cloned into a pLenti construct with a CMV promoter

(Delgado-Martinez et al., 2007). Mutations were made using the QuikChange II XL kit (Agilent).

Primers were ordered from TAGC Copenhagen. All mutations were verified by sequencing before

virus production. Viruses were prepared as previously described using transfection of HEK293FT-

cells (Naldini et al., 1996). Neurons were infected with lentiviruses on DIV 0-1. 30 µl total virus per

well (WT or mutant; 15 µl + 15 µl WT + mutant virus for co-expressed condition).

Immunostaining and confocal microscopy

Autaptic hippocampal neurons were fixed at DIV14 in 2% PFA in culture medium for 10 min and

subsequently in 4% PFA for 10 min. Cells were then washed with PBS, permeabilized by 0.5% Triton

X-100 for 5 min and blocked with 4% normal goat serum in 0.1% Triton X-100 (blocking solution) for

30 min. Cells were incubated with primary antibodies diluted in blocking solution (anti-MAP2, 1:500,

chicken, ab5392, Abcam; and anti-vGlut1 1:1000, guinea pig, AB5905, Merck Millipore) for 2 h at RT.

After washing with PBS, the cells incubated with secondary antibodies in blocking solution for 1h at

RT in the dark (anti-chicken Alexa 568, 1:1000, A11041, Thermo Fisher Scientific; and anti-guinea

pig Alexa 647, 1:1000, A-21450, Thermo Fisher Scientific) and washed again. Coverslips were

mounted with FluorSave and imaged on Zeiss CellObserver spinning disc confocal microscope (40x

water immersion objective; NA 1.2) with Zeiss Zen Blue 2012 software. Images were acquired as Z-

stack and 9 images per plane to capture the whole island in the field of view. The images were post-

processed with Zeiss Zen Black software and neuronal morphology was analyzed using SynD

automated image analysis (Schmitz et al., 2011).

Western Blotting

Harvesting the high-density hippocampal culture, BCA assay and transferring the protein samples

on a PVDF membrane were performed as described before (Ruiter et al., 2019). Incubation in

primary antibodies (a-SNAP25: mouse, 1:10000, SYSY 111011, Synaptic Systems; a-VCP: mouse,

1:2000, ab11433, Abcam) was performed overnight with 70 rpm shaking at 4 °C, followed by

washing in TBST (0.1%) and a 1 h incubation in secondary antibody (goat a-mouse-HRP: 1:10000,

P0447, Dako). After washing, Pierce ECL Western blotting substrate was added and

chemiluminescence was visualized with FluorChem E (Proteinsimple).

Electrophysiology

Autaptic cultures were used from DIV10 until DIV14. The intracellular pipette medium contained:

KCl 136 mM, HEPES 17.8 mM, Creatine Phosphate 15 mM, Na-ATP 4 mM, Creatine Phosphokinase

50 U, $MgCl_2$ 4.6 mM, EGTA 1 mM (pH 7.4, osmolarity ~300 mOsm). The standard extracellular

recording medium contained: NaCl 140 mM, KCl 2.4 mM, HEPES 10 mM, Glucose 14 mM, $CaCl_2$ 2

mM, $MgCl_2$ 2 mM. The extracellular recording medium for de-priming experiments contained: NaCl

140 mM, KCl 2.4 mM, HEPES 10 mM, Glucose 14 mM, $CaCl_2$ 4 mM, $MgCl_2$ 4 mM (pH 7.4, osmolarity

~300 mOsm). An Axio Observer A1 inverted microscope (Zeiss) was used to visualize the cells. The

recordings were performed at room temperature. An EPC10 amplifier (HEKA) was used with the

program Patchmaster v2.73 (HEKA). Traces were filtered with a 3kHz Bessel low-pass filter and data

were acquired at 20 kHz. The series resistance was compensated to 70%. Glass pipettes were freshly

pulled on a P1000 pipette puller (Sutter Instruments) from borosilicate glass capillaries. Pipets

ranging from 2.5 to 5 MΩ were selected for recordings. Cells with starting access resistance above

10 MΩ were rejected. Recordings were performed in voltage clamp, with the holding potential kept

at -70 mV. Evoked excitatory postsynaptic currents (EPSC) were induced by raising the holding

voltage to 0 mV for 2 ms. Sucrose was dissolved into the standard extracellular recording medium.

Application of the extracellular media was done using a custom-made barrel system, controlled by

SF-77B perfusion fast step (Warner Instruments) controlled via digital output switches from the

EPC10. Electrophysiological data was analyzed in IGOR Pro (v6.21 and v6.37, WaveMetrics) using a

custom-written script. mEPSCs were analyzed with MiniAnalysis (v6.0.7, Synaptosoft).

Electrophysiological data: Calculations

Deconvolution was calculated and the electrostatic model for triggering was fitted to 4K and

I67N/4K data as described before (Ruiter et al., 2019). Since the evoked release for I67N was so

small that deconvolution became unreliable, we downscaled the peak release rate of the I67N/4K

mutation with a factor of 5.616, corresponding to the reduction in eEPSC amplitude in the I67N

compared to I67N/4K.

In order to determine the reasons for the reduced RRP size for V48F and D166Y, we considered a

single pool model for the RRP, with priming rate k_1, depriming rate k_{-1}, and fusion rate k_f (Fig. 6A).

The equation describing the evolution in RRP size is:

(Eq. 1) $\qquad \dfrac{d\mathrm{RPP}(t)}{dt} = k_1 - \left(k_{-1} + k_f\right) \cdot \mathrm{RRP}(t)$

where k_1 has the unit vesicles/s, whereas k_{-1}, and k_f have the unit 1/s. The solution of this differential

equation is

(Eq. 2) $\mathrm{RPP}(t) = \frac{k_1}{(k_{-1}+k_f)}\left(1 - e^{-(k_{-1}+k_f)t}\right)$

The steady-state RPP size is

(Eq. 3) $RRP = \frac{k_1}{(k_{-1}+k_f)}$

The miniature release rate at steady state is

(Eq. 4) $r_{mini} = \frac{k_f k_1}{(k_{-1}+k_f)}$

When stimulated by sucrose, k_f increases, and if $k_f \gg k_{-1}$ (for justification, see below) the current

plateau will report on the forward priming rate alone:

(Eq. 5a) $r_{prime} \approx k_1$

Thus, the current plateau during sucrose application can be used for estimating the forward priming

rate. However, application of 0.5 M sucrose causes cell shrinkage and changes in solution viscosity,

which in turn can change the leak current. This might cause the plateau current to change, which

might be invisible during the experiment due to the synaptic events. To correct for this, we

implemented variance-mean analysis to identify the true baseline current (the current

corresponding to the lack of synaptic release). Synaptic release is essentially a source of shot noise,

for which the variance is proportional to the mean. We therefore calculated the variance (after

subtraction of a running average) and mean of the current in 50 ms intervals during the sucrose

application, and performed linear regression in a variance-mean plot. The corrected baseline was

identified by backextrapolation to the variance level found in the absence of synaptic activity (using

a stretch of current before sucrose application), as illustrated in Fig. 6C.

If sucrose does not sufficiently increase k_f, the equation 5a above would be replaced by

(Eq. 5b) $r_{mini} = \frac{k_f k_1 N_{suc}}{(k_{-1} + k_f N_{suc})}$

where N_{suc} is the fold-increase in k_f induced by sucrose application. Plotting the solutions to Eqs 3,

4, and 5b at different values of N_{suc}, the dependency of the estimated k_{-1} and k_1 upon N_{suc} can be

investigated (Figure 6 – Figure supplement 1). Notably, in wildtype cells $N_{suc} > 5,000$ (Schotten et al.,

2015), indicating that the estimated values of k_{-1} and k_1 using Eq. 5a (Table 1) is quite accurate for

the WT case, and the estimation in the case of the D166Y and V48F is even less dependent on N_{suc},

because the value of k_f is higher at rest (Figure 6 – Figure supplement 1). Importantly, for no realistic

value of N_{suc} would the conclusion of decreased k_1 and k_{-1} in the two mutants be in jeopardy.

For calculating the energy profiles of WT and mutants, we used Arrhenius' Equation:

(Eq. 6) $k = Ae^{\left(\frac{-E_A}{RT}\right)}$

where E_A is the activation energy, R and T are the gas constants and the absolute temperature,

respectively, and A is an empirical constant that depends on collision rates (Schotten et al., 2015).

Solving for the activation energy, we get:

(Eq. 7) $E_a = RT(\ln(A) - \ln(k))$

Since A is unknown, we cannot use this equation to calculate the absolute values of the transition

energies; however, when we compare a mutation to the WT condition, and under the assumption

that A is unchanged by mutation, we can calculate the difference in energy level of the transition

states:

(Eq. 8) $\Delta E_a^{Mut-WT} = RT(\ln(k^{WT}) - \ln(k^{Mut}))$

Using this equation sequentially, for the priming rate, the depriming rate and the fusion rate, we

can derive the entire energy diagram, under the additional assumption that the energy in the pre-

primed state is identical between WT and mutation, and using that at room temperature, RT=2.479

kJ/mol.

The assumption that the empiric factor, *A*, is unchanged by mutation is likely to hold for the fusion

reaction, which depends on conformational changes in a preformed complex. In contrast, collision

rates might be involved in priming; in that case, the effect of the V48F and D166Y mutations on

priming, which we here attribute to an increase in the energy level of the priming transition state

might reflect a lower collision rate between vesicles and plasma membrane fusion machinery,

and/or a lower energy level of the pre-primed state.

Constructs for in vitro protein expression

The following constructs were used: Glutathione S-Transferase (GST) - full length VAMP2 is encoded

by plasmid pSK28 (Kedar et al., 2015), GST-cytosolic domain VAMP2 (amino acids 1-94) (pSK74,

(Ruiter et al., 2019)), synaptotagmin 1-His6 lacking the luminal domain (amino acids 57-421) (pLM6,

(Mahal et al., 2002), His6-complexin II (CpxII) (pMDL80, (Malsam et al., 2012)), His6-syntaxin-1A

(pSK270, (Schollmeier et al., 2011)), His6-SNAP25B (pFP247, (Parlati et al., 1999)), t-SNARE

consisting of syntaxin-1A and His6-SNAP25B (pTW34, (Parlati et al., 1999)). Point mutants in soluble

His6-SNAP25B and in the t-SNARE complex were generated by using the DNA templates pFP247 and

pTW34, respectively, and the Quikchange DNA mutagenesis kit (Qiagen) (Ruiter et al., 2019).

Thereby, the following SNAP25 constructs were established: His6-SNAP25B mutant I67N (pUG1),

V48F (pUG2), D166Y (pUG3), and t-SNAREs containing the corresponding SNAP25B mutants I67N

(pUG7), V48F (pUG8), and D166Y (pUG9). The identity of all constructs was validated by DNA

sequencing.

Protein expression and purification

In general, the expression vectors, encoding the desired protein constructs were transfected into

Escheria coli BL21 (DE3) (Stratagene). At an OD_{660} of 0.8, protein expression was induced by the

addition of 0.3 mM IPTG. Alternatively, proteins were expressed by autoinduction using buffered

media containing lactose (Studier, 2005). Cells were harvested by centrifugation (3500 rpm, 15

minutes, H-12000 rotor, Sorvall) and lysed using the high-pressure pneumatic processor 110L

(Microfluidics). Cell fragments were removed by centrifugation at 60.000 rpm (70Ti rotor, Beckman)

for 1 hour and the clarified supernatant was snap-frozen in liquid nitrogen.

The purification of full length VAMP2 was performed as described previously (Kedar et al., 2015)

with the following modifications. Cells were grown in ZYM media (Studier, 2005) and protein

expression was induced with IPTG for 3 hours at 25°C. The purification and expression of the GST-

tagged cytosolic domain of VAMP2 was described previously (Ruiter et al., 2019). Synaptotagmin 1-

His6 lacking the luminal domain was purified as described previously (Malsam et al., 2012) with the

following modification. After dilution to 50 mM salt, the protein was further purified on a MonoS

Sepharose column (GE healthcare) applying a gradient of 50 to 500 mM KCl in 25 mM Hepes-KOH

(pH 7.4).

His6-syntaxin-1A purification was performed as outlined by (Schollmeier et al., 2011) with the

following modifications. Briefly, cells were grown in ZYM media (Studier, 2005) and autoinduction

was used for the protein expression at 22°C overnight. Syntaxin-1A was eluted from Ni-NTA-beads

(Qiagen) by over-night cleavage with Prescission protease (GE healthcare) at 4°C, removing the His$_6$

tag. After dilution to 80 mM salt, the protein was further purified on a MonoQ Sepharose column

(GE healthcare) applying a gradient of 50 to 500 mM KCl in 25 mM HEPES-KOH (pH 7.4).

His6-SNAP25 was expressed as depicted for syntaxin-1A and purified via Ni-NTA beads, followed by

MonoQ Sepharose column chromatography (Ruiter et al., 2019). Preassembled full-length t-SNARE

complexes were expressed and purified as described previously (Weber et al., 1998). His$_6$-Complexin

II expression and purification was performed according to (Malsam et al., 2012) with the following

modifications. His6-CpxII was expressed in BL21-DE3 codon+ bacteria for 2 hours at 27°C.

The concentrations of purified proteins were determined by SDS-PAGE and Coomassie Blue staining

using BSA as a standard and the Fiji software for quantification. Furthermore, mutant and wildtype

protein concentrations were directly compared on a single gel.

Protein reconstitution into liposomes

All lipids were from Avanti Polar Lipids, except of Atto488-DPPE and Atto550-DPPE, which were

purchased from ATTO-TEC. For VAMP2 and Syt1 reconstitution into small unilamellar vesicles

(SUVs), lipid mixes (3 µmol total lipid) with the following composition were prepared: 25 mol% POPE

(1-hexadecanoyl-2-octadecenoyl-SN-glycero-3-phosphoethanolamine), 15 mol% DOPS (1,2-

dioleoyl-SN-glycero-3-phosphoserine), 29 mol% POPC (1-palmitoyl-2-oleoyl-SN-glycero-3-

phosphocholine), 25 mol% cholesterol (from ovine wool), 5 mol% PI (L-α-phosphatidylinositol), 0.5

mol% Atto488-DPPE (1,2-dipalmitoyl-SN-glycero-3-phosphoethanolamine) and 0.5 mol% Atto550-

DPPE. For docking assays, the t-SNARE liposome lipid mix (5 µmol total lipid) had the following

composition: 35 mol% POPC, 15 mol% DOPS, 20 mol% POPE, 25 mol% cholesterol, 4 mol% PI, 0,05

mol% Atto647-DPPE and 0,5 mol% tocopherol. For the preparation of PI(4,5)P2-containing t-SNARE

and syntaxin-1A liposomes, the t-SNARE liposome lipid mix was used, but 1 mol% PI(4,5)P2 (L-α-

phosphatidylinositol-4,5-bisphosphate) was added and the POPC concentration lowered by 1%

accordingly.

Proteins were reconstituted as described previously (Malsam et al., 2012). For the docking and

fusion assays, t-SNARE wildtype and mutants were reconstituted at a protein to lipid ratio of 1:900.

For the syntaxin-1A membrane fusion assay, syntaxin-1A was reconstituted at a protein to lipid ratio

of 1:1000. Briefly, 5 µmol dried lipids were dissolved in 0.7 ml reconstitution buffer (25 mM HEPES-

KOH, pH 7.4, 550 mM KCl, 1 mM EDTA-NaOH, 1 mM DTT) containing final 1.4 weight% octyl-ß-D-

glucopyranoside (ß-OG) and either 6.5 nmol (390 µg) t-SNARE complex or 5 nmol (165 µg) syntaxin-

1A. SUVs containing either t-SNARE or syntaxin-1A were formed by rapid ß-OG dilution below the

critical micelle concentration by adding 1.4 ml reconstitution buffer. For the quantification of lipid

recovery, a 20 µl aliquot (GUV input) was removed and stored at -20°C. The liposome suspension

was desalted using a PD10 column (GE Healthcare) equilibrated with desalting buffer 1 (1 mM

HEPES-KOH, pH 7.5, 1 w/v% glycerol, 10 µM EGTA-KOH, 1 mM DTT) and snap frozen in four aliquots

in liquid nitrogen and stored in -80°C. These syntaxin-1A and t-SNARE SUVs were later used to

prepare giant unilamellar vesicles (GUVs). The final protein to lipid ratios were determined by SDS-

PAGE and Coomassie Blue staining of the proteins and Atto647 fluorescence intensity

measurements of the lipids.

VAMP2/Synaptotagmin 1-SUVs were prepared as described previously (Malsam et al., 2012; Weber

et al., 1998) with the following modification: VAMP2 and Synaptotagmin 1 were reconstituted at a

protein to lipid ratio of 1:350 and 1:800, respectively. SUVs, which were used for the SNAP25

recruitment assay, were not dialyzed twice, but directly harvested after the density gradient

flotation and lipid recovery and protein-to-lipid-ratio were determined (Stx1-SUVs: 1:1900; Syt1/V2-

SUVs: Syt1 1:700, VAMP2 1:300; V2-SUVs: VAMP2 1:200; Syt1-SUVs: Syt1 1:700).

GUV preparation

GUVs were prepared as described previously (Kedar et al., 2015). Briefly, t-SNARE or syntaxin-1A

liposomes (1.25 µmol lipid) were loaded onto a midi column (GE Healthcare) equilibrated with

desalting buffer 2 containing trehalose (1 mM HEPES-KOH, pH 7.5, 0.5 w/v% glycerol, 10 µM EGTA-

KOH, pH 7.4, 50 µM MgCl2, 1 mM DTT, 10 mM trehalose). 1.4 ml eluate were collected and

liposomes sedimented in a TLA-55 rotor (Beckman) at 35.000 rpm for 2 h at 4°C. The pellet was

resuspended by rigorous vortexing and, while vortexing, was diluted with 10 µl of pellet

resuspension buffer (1 mM HEPES-KOH, pH 7.4, 10 µM EGTA-KOH, 50 µM MgCl$_2$) to lower the

osmotic strength. The total volume (20-25 µl) was spread as a uniform layer (14 mm diameter) on

the surface of a platinum foil (Alfa Aesar; 25 x 25 mm, 0.025 mm thick) attached to a glass slide as

support. After drying the liposome suspension for 50 min at 50 mbar, the incubation chamber was

assembled using an O-ring (2 mm x 18 mm), filled with 620 µl of swelling buffer (1mM EPPS-KOH,

pH 8.0, 240 mM sucrose (Ca^{2+} free), 1 mM DTT) and closed using a second platinum plate.

Conductive copper tape (3M) was attached to the platinum foil to connect the assembly with a

function generator (Voltcraft 8202). GUVs were generated by electro-formation at 10 Hz and 1 V at

0°C overnight.

Lipid mixing assay

The membrane fusion assay was performed as described previously (Malsam et al., 2012), except

that in the fusion buffer HEPES was replaced with 20 mM MOPS pH 7.4. Briefly, t-SNARE-GUVs (14

nmol lipid, 15 ± 0,7 pmol t-SNARE) were preincubated with or without 6 µM (0,6 nmol) CpxII for 5

min on ice in fusion buffer containing 0,1 mM EGTA-KOH and 0.5 mM MgCl$_2$. When using syntaxin-

1A-GUVs (14 nmol lipid, 14 pmol syntaxin-1A), these preincubations contained 2 µM (0.2 nmol) of

soluble SNAP25 in addition. Subsequently, VAMP2/Syt1-SUVs (2.5 nmol lipid, 4.5 pmol VAMP2, 2

pmol Syt1) were added to the GUV reaction mix resulting in 104 µl sample volume. After 10 minutes

on ice, 100 µl of the GUV-SUV mixes were transferred into a prewarmed 96-well plate (37°C) and

fluorescence emitted by Atto488 (λ_{ex} = 485 nm, λ_{em} = 538 nm) was measured in a Synergy 4 plate

reader (BioTek Instruments GmbH) at intervals of 10 seconds. Ca^{2+} was added to a final free

concentration of 100 µM

(https://somapp.ucdmc.ucdavis.edu/pharmacology/bers/maxchelator/CaEGTA-TS.htm) after 2

minutes or 30 minutes for t-SNARE-GUVs or syntaxin-1A-GUVs, respectively. The fusion reactions

were stopped after 4 minutes for t-SNARE-GUVs or after 1 h for syntaxin-1A-GUVs by the addition

of 0.7% SDS and 0.7% n-Dodecyl-ß-D-Maltosid (DDM). The resulting "maximum" fluorescent signal

was used to normalize the fusion-dependent fluorescence. As a negative control, SUVs were treated

with Botulinum NeuroToxin type D (BoNT-D) and their fluorescence signals were subtracted from

individual measurement sets. Three independent fusion experiments were performed for each

mutant.

SUV-GUV binding assay

All SUV-GUV binding studies were performed in an ice bath and all pipetting steps were carried out

in the cold room to avoid membrane fusion (Parisotto et al, 2012; Malsam et al, 2012; Weber et al,

1998). Before starting the incubation, potential SUV aggregates were removed by centrifugation. T-

SNARE-GUVs (28 nmol lipid, 30 ± 1,4 pmol t-SNARE) were preincubated with VAMP2/Syt1-SUVs (5

nmol lipid, 9 pmol VAMP2, 4 pmol Syt1) on ice in 100 µL fusion buffer (20 mM MOPS-KOH, pH 7.4,

135 mM KCl, 1mM DTT) with 0.1 mM EGTA and 1 mM $MgCl_2$. After 10 min incubation, the reactions

were underlaid with 20 µL of a sucrose cushion (1 mM MOPS-KOH, pH 7.4, 60 mM sucrose, 1 mM

DTT) and the GUVs with attached SUVs were re-isolated by centrifugation for 10 min. After removing

the supernatant, the pellets (in 10 µL remaining volume) were resuspended, transferred into new

tubes, treated with 100 µL of 1% SDS/1% DDM and the SUV recovery was determined by measuring

the Atto488 fluorescence.

To determine the respective inputs, 28 nmol GUV lipids or 5 nmol SUV lipids were treated with 1%

SDS/1% DDM (final) and the corresponding Atto647 and Atto488 fluorescence was measured at λ_{ex}

= 620/40 nm, λ_{em} = 680/30 nm and λ_{ex} = 485/20 nm, λ_{em} = 528/20 nm, respectively. A sample lacking

SUVs was used to determine the GUV recovery (usually 80-95%). GUV recovery of each sample was

used to normalize the respective SUV docking. A sample without GUVs was used to determine the

absolute background (usually <15%), which was subtracted from all samples.

SNAP25 recruitment assay / SUV flotation assay

Small unilamellar liposomes (SUVs) containing 35 pmol syntaxin-1A or 210 pmol VAMP2 and/or 100

pmol Synaptotagmin-1 (66 nmol lipid for Stx1-, V2/Syt1- or Syt1-SUVs, 40 nmol lipid for V2-SUVs)

were mixed with 180 pmol SNAP25 in a final assay volume of 50 µl in fusion buffer (20 mM MOPS-

KOH, pH 7.4, 135 mM KCl, 1 mM DTT). After two hours on ice, allowing the SNARE complex

formation, the samples were diluted five times with fusion buffer and mixed with the equivalent

volume of 80% nycodenz solution. After overlaying the sample with 100 µl 35% nycodenz, 25 µl 20%

nycodenz solution and finally with 5 µl fusion buffer in an ultra-clear tube (5 x 41 mm), the liposomes

were isolated by centrifugation for 3 h 40 mins at 55,000 rpm at 4°C in a SW 60 rotor (SW 60 Ti,

Beckman). 20 µl were harvested from the top of the gradient and mixed with 8 µl of 4x Laemmli

buffer (final 62.5 mM Tris-HCl, pH 6.8, 10% glycerol, 2% SDS, 50 mM ß-mercaptoethanol, 0.1%

bromphenol blue). 18 µl of this mixture were used to quantify the amount of recruited SNAP25 by

SDS PAGE followed by Coomassie Staining and Silver Staining. Using the Fiji software (Image J based)

the Coomassie stained band intensity of syntaxin-1A or VAMP2 or Syt1, respectively, were

determined and normalized to the respective mean. Subsequently, Silver Staining was used to

quantify the band intensities of SNAP25, and these values were normalized to the intensities of the

relative protein (e.g. Syntaxin-1A) based on the Coomassie Staining. From this, the ratios between

the SNAP25 mutants and wild type were determined.

Temperature-dependent dissociation of v-/t-SNARE complexes in SDS

SNARE complex stability was determined as described previously (Schupp et al., 2016). Briefly,

preassembled full-length t-SNARE complexes (WT and mutants, 10 µM) were incubated with the

cytoplasmic domain of VAMP2 (30 µM) in 25 mM MOPS (3-(Nmorpholino) propanesulfonic acid)-

KOH, pH 7.4, 135 mM KCl, 1% Octyl ß-D-glucopyranoside, 1 mM Dithiothreitol, 10 mM TECEP (Tris(2-

carboxyethyl)phosphine hydrochloride)-KOH, pH 7.4, 1 mM EDTA-NaOH, pH 7.4 overnight at 0°C

and subsequently for 1 h at 25°C. Subsequently, 37 µl of reaction mixture (36 µg of total protein)

was diluted with 213 µl of 1x Laemmli buffer. 7.5 µl aliquots were incubated at the indicated

temperatures for 5 min. Samples were analyzed by SDS-PAGE (15% gels) and proteins were

visualized by Coomassie brilliant blue staining. Temperature-dependent dissociation of the SNARE

complex was quantified by the appearance of free syntaxin-1A (35 kDa protein band released from

the high MW SNARE complex) using the Fiji (Image J based) software. Data were normalized to the

maximum value of a measurement set.

Silver Staining

Coomassie prestained gels from SDS PAGE were destained overnight in destain solution (30 %

methanol, 10 % acetic acid). Gels were gently washed 30 minutes with 10% ethanol and one minute

with 0.02% sodium thiosulfate. After short washing with deionized water, gels were stained (0.03%

paraformaldehyde, 0.002% silver nitrate) for 15 minutes and again quickly washed with water.

Developing solution (0.06% sodium carbonate, 0.018% paraformaldehyde, 0.0002% sodium

thiosulfate) was applied to the gels until protein band intensities were satisfactorily stained. The

reactions were stopped by replacing the staining solution with 0.07% acetic acid. Gels were scanned

and quantified by using the Fiji (Image J based) software.

Statistical analysis

Graphs (bar and line) display mean ± SEM with all points displayed, except when otherwise noted;

for electrophysiological experiments *n* denotes the number of cells recorded and is given in the

legends. For *in vitro* experiments the number of biological replicates was 3 unless stated otherwise

in the legends. Statistics were performed using GraphPad Prism 9. Unless otherwise noted,

statistical differences between several groups were determined by one-way ANOVA; post-test was

Dunnett's test comparing to the WT condition as a reference. Equal variance of groups was tested

by the Brown-Forsythe test; in case of a significant test, the Brown-Forsythe ANOVA test, which

does not assume equal variances, was used instead. Kruskal-Wallis test was used in cases, where

the data structure contained many identical values (zeros). Pairwise testing was carried out using

unpaired t-test, or Welch's t-test in case of significantly different variances as determined by an F-

test. The test is mentioned in the Figure legend; if no test is mentioned, the difference was not

significantly different. Significance was assumed when $p<0.05$ and the level of significance is

indicated by asterisk: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Acknowledgements

978 We would like to thank Dorte Lauritsen and Ursula Göbel for excellent technical assistance. This

979 investigation was supported by the Novo Nordic Foundation (NNF19OC0058298, JBS), the

980 Independent Research Fund Denmark (8020-00228A, JBS), the Lundbeck Foundation (R277-2018-

981 802, JBS), and the German Reseach Foundation DFG (grant 278001972- TRR 186, THS, JM).

983 **Table 1.**

Mean±SEM	WT	V48F	WT	D166Y	*Syt1* WT	*Syt1* KO
k_1 [vesicles/s]	385.6 ±52.2	79.87 *** ±12.5	457.4 ±77.4	37.68 **** ±7.33	1227 ±189	646 * ±114
k_{-1} [1/s]	0.0903 ±0.0091	0.0605 # ±0.011	0.1114 ±0.012	0.0294 **** ±0.0122	0.140 ±0.016	0.114 ¤ ±0.013
k_f [1/s]	0.000844 ±0.000178	0.0164**** ±0.00230	0.000398 ±0.000077	0.03522**** ±0.00352	0.000235 ±0.000043	0.00286*** ±0.00062

985 **Estimated parameters affecting the size of the RRP.** Displayed is mean ±SEM. Two-sample t-test or
986 Welch's t-test comparing mutant to WT: *$p<0.05$; ***$p<0.001$; ****$p<0.0001$, # non-significant
987 (p=0.125), ¤ non-significant (p=0.210).

Literature

Abramov, D., N.G.L. Guiberson, and J. Burre. 2021a. STXBP1 encephalopathies: Clinical spectrum, disease mechanisms, and therapeutic strategies. *J Neurochem*. 157:165-178.

Abramov, D., N.G.L. Guiberson, A. Daab, Y. Na, G.A. Petsko, M. Sharma, and J. Burre. 2021b. Targeted stabilization of Munc18-1 function via pharmacological chaperones. *EMBO Mol Med*. 13:e12354.

Alten, B., Q. Zhou, O.H. Shin, L. Esquivies, P.Y. Lin, K.I. White, R. Sun, W.K. Chung, L.M. Monteggia, A.T. Brunger, and E.T. Kavalali. 2021. Role of Aberrant Spontaneous Neurotransmission in SNAP25-Associated Encephalopathies. *Neuron*. 109:59-72 e55.

Bacaj, T., D. Wu, J. Burre, R.C. Malenka, X. Liu, and T.C. Sudhof. 2015. Synaptotagmin-1 and -7 Are Redundantly Essential for Maintaining the Capacity of the Readily-Releasable Pool of Synaptic Vesicles. *PLoS Biol*. 13:e1002267.

Baker, K., S.L. Gordon, H. Melland, F. Bumbak, D.J. Scott, T.J. Jiang, D. Owen, B.J. Turner, S.G. Boyd, M. Rossi, M. Al-Raqad, O. Elpeleg, D. Peck, G.M.S. Mancini, M. Wilke, M. Zollino, G. Marangi, H. Weigand, I. Borggraefe, T. Haack, Z. Stark, S. Sadedin, G. Broad Center for Mendelian, T.Y. Tan, Y. Jiang, R.A. Gibbs, S. Ellingwood, M. Amaral, W. Kelley, M.A. Kurian, M.A. Cousin, and F.L. Raymond. 2018. SYT1-associated neurodevelopmental disorder: a case series. *Brain*. 141:2576-2591.

Baker, R.W., P.D. Jeffrey, M. Zick, B.P. Phillips, W.T. Wickner, and F.M. Hughson. 2015. A direct role for the Sec1/Munc18-family protein Vps33 as a template for SNARE assembly. *Science*. 349:1111-1114.

Bao, H., D. Das, N.A. Courtney, Y. Jiang, J.S. Briguglio, X. Lou, D. Roston, Q. Cui, B. Chanda, and E.R. Chapman. 2018. Dynamics and number of trans-SNARE complexes determine nascent fusion pore properties. *Nature*. 554:260-263.

Basu, J., A. Betz, N. Brose, and C. Rosenmund. 2007. Munc13-1 C1 domain activation lowers the energy barrier for synaptic vesicle fusion. *J Neurosci*. 27:1200-1210.

Bekkers, J.M., and C.F. Stevens. 1991. Excitatory and inhibitory autaptic currents in isolated hippocampal neurons maintained in cell culture. *Proc Natl Acad Sci U S A*. 88:7834-7838.

Bouazza-Arostegui, B., M. Camacho, M.M. Brockmann, S. Zobel, and C. Rosenmund. 2022. Deconstructing Synaptotagmin-1's Distinct Roles in Synaptic Vesicle Priming and Neurotransmitter Release. *J Neurosci*. 42:2856-2871.

Bradberry, M.M., N.A. Courtney, M.J. Dominguez, S.M. Lofquist, A.T. Knox, R.B. Sutton, and E.R. Chapman. 2020. Molecular Basis for Synaptotagmin-1-Associated Neurodevelopmental Disorder. *Neuron*. 107:52-64 e57.

Bronk, P., F. Deak, M.C. Wilson, X. Liu, T.C. Sudhof, and E.T. Kavalali. 2007. Differential effects of SNAP-25 deletion on Ca2+ -dependent and Ca2+ -independent neurotransmission. *J Neurophysiol*. 98:794-806.

Chang, S., T. Trimbuch, and C. Rosenmund. 2018. Synaptotagmin-1 drives synchronous Ca(2+)-triggered fusion by C2B-domain-mediated synaptic-vesicle-membrane attachment. *Nat Neurosci*. 21:33-40.

Chen, Y., Y.H. Wang, Y. Zheng, M. Li, B. Wang, Q.W. Wang, C.L. Fu, Y.N. Liu, X. Li, and J. Yao. 2021. Synaptotagmin-1 interacts with PI(4,5)P2 to initiate synaptic vesicle docking in hippocampal neurons. *Cell Rep*. 34:108842.

Courtney, K.C., J.D. Vevea, Y. Li, Z. Wu, Z. Zhang, and E.R. Chapman. 2021. Synaptotagmin 1 oligomerization via the juxtamembrane linker regulates spontaneous and evoked neurotransmitter release. *Proc Natl Acad Sci U S A*. 118.

Courtney, N.A., H. Bao, J.S. Briguglio, and E.R. Chapman. 2019. Synaptotagmin 1 clamps synaptic vesicle fusion in mammalian neurons independent of complexin. *Nat Commun*. 10:4076.

Das, D., H. Bao, K.C. Courtney, L. Wu, and E.R. Chapman. 2020. Resolving kinetic intermediates during the regulated assembly and disassembly of fusion pores. *Nat Commun*. 11:231.

Delgado-Martinez, I., R.B. Nehring, and J.B. Sorensen. 2007. Differential abilities of SNAP-25 homologs to support neuronal function. *J Neurosci*. 27:9380-9391.

Fang, Q., Y. Zhao, A.D. Herbst, B.N. Kim, and M. Lindau. 2015. Positively charged amino acids at the SNAP-25 C terminus determine fusion rates, fusion pore properties, and energetics of tight SNARE complex zippering. *J Neurosci*. 35:3230-3239.

Fasshauer, D., D. Bruns, B. Shen, R. Jahn, and A.T. Brunger. 1997. A structural change occurs upon binding of syntaxin to SNAP-25. *J Biol Chem*. 272:4582-4590.

Fasshauer, D., R.B. Sutton, A.T. Brunger, and R. Jahn. 1998. Conserved structural features of the synaptic fusion complex: SNARE proteins reclassified as Q- and R-SNAREs. *Proc Natl Acad Sci U S A*. 95:15781-15786.

Fernandez-Busnadiego, R., B. Zuber, U.E. Maurer, M. Cyrklaff, W. Baumeister, and V. Lucic. 2010. Quantitative analysis of the native presynaptic cytomatrix by cryoelectron tomography. *J Cell Biol*. 188:145-156.

Gao, Y., S. Zorman, G. Gundersen, Z. Xi, L. Ma, G. Sirinakis, J.E. Rothman, and Y. Zhang. 2012. Single reconstituted neuronal SNARE complexes zipper in three distinct stages. *Science*. 337:1340-1343.

Guiberson, N.G.L., A. Pineda, D. Abramov, P. Kharel, K.E. Carnazza, R.T. Wragg, J.S. Dittman, and J. Burre. 2018. Mechanism-based rescue of Munc18-1 dysfunction in varied encephalopathies by chemical chaperones. *Nat Commun*. 9:3986.

Hamdan, F.F., C.T. Myers, P. Cossette, P. Lemay, D. Spiegelman, A.D. Laporte, C. Nassif, O. Diallo, J. Monlong, M. Cadieux-Dion, S. Dobrzeniecka, C. Meloche, K. Retterer, M.T. Cho, J.A. Rosenfeld, W. Bi, C. Massicotte, M. Miguet, L. Brunga, B.M. Regan, K. Mo, C. Tam, A. Schneider, G. Hollingsworth, S. Deciphering Developmental Disorders, D.R. FitzPatrick, A. Donaldson, N. Canham, E. Blair, B. Kerr, A.E. Fry, R.H. Thomas, J. Shelagh, J.A. Hurst, H. Brittain, M. Blyth, R.R. Lebel, E.H. Gerkes, L. Davis-Keppen, Q. Stein, W.K. Chung, S.J. Dorison, P.J. Benke, E. Fassi, N. Corsten-Janssen, E.J. Kamsteeg, F.T. Mau-Them, A.L. Bruel, A. Verloes, K. Ounap, M.H. Wojcik, D.V.F. Albert, S. Venkateswaran, T. Ware, D. Jones, Y.C. Liu, S.S. Mohammad, P. Bizargity, C.A. Bacino, V. Leuzzi, S. Martinelli, B. Dallapiccola, M. Tartaglia, L. Blumkin, K.J. Wierenga, G. Purcarin, J.J. O'Byrne, S. Stockler, A. Lehman, B. Keren, M.C. Nougues, C. Mignot, S. Auvin, C. Nava, S.M. Hiatt, M. Bebin, Y. Shao, F. Scaglia, S.R. Lalani, R.E. Frye, I.T. Jarjour, S. Jacques, R.M. Boucher, E. Riou, M. Srour, L. Carmant, A. Lortie, P. Major, P. Diadori, F. Dubeau, G. D'Anjou, G. Bourque, S.F. Berkovic, L.G. Sadleir, P.M. Campeau, Z. Kibar, R.G. Lafreniere, S.L. Girard, S. Mercimek-Mahmutoglu, C. Boelman, et al. 2017. High Rate of Recurrent De Novo Mutations in Developmental and Epileptic Encephalopathies. *Am J Hum Genet*. 101:664-685.

Hoerder-Suabedissen, A., K.V. Korrell, S. Hayashi, A. Jeans, D.M.O. Ramirez, E. Grant, H.C. Christian, E.T. Kavalali, M.C. Wilson, and Z. Molnar. 2019. Cell-Specific Loss of SNAP25 from Cortical Projection Neurons Allows Normal Development but Causes Subsequent Neurodegeneration. *Cereb Cortex*. 29:2148-2159.

Honigmann, A., G. van den Bogaart, E. Iraheta, H.J. Risselada, D. Milovanovic, V. Mueller, S. Mullar, U. Diederichsen, D. Fasshauer, H. Grubmuller, S.W. Hell, C. Eggeling, K. Kuhnel, and R. Jahn. 2013. Phosphatidylinositol 4,5-bisphosphate clusters act as molecular beacons for vesicle recruitment. *Nat Struct Mol Biol*. 20:679-686.

Huson, V., M. Meijer, R. Dekker, M. Ter Veer, M. Ruiter, J.R. van Weering, M. Verhage, and L.N. Cornelisse. 2020. Post-tetanic potentiation lowers the energy barrier for synaptic vesicle fusion independently of Synaptotagmin-1. *Elife*. 9.

Jakhanwal, S., C.T. Lee, H. Urlaub, and R. Jahn. 2017. An activated Q-SNARE/SM protein complex as a possible intermediate in SNARE assembly. *EMBO J*. 36:1788-1802.

Jiao, J., M. He, S.A. Port, R.W. Baker, Y. Xu, H. Qu, Y. Xiong, Y. Wang, H. Jin, T.J. Eisemann, F.M. Hughson, and Y. Zhang. 2018. Munc18-1 catalyzes neuronal SNARE assembly by templating SNARE association. *Elife*. 7.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S.A.A. Kohl, A.J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A.W. Senior, K. Kavukcuoglu, P. Kohli,

and D. Hassabis. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature*. 596:583-589.

Kalyana Sundaram, R.V., H. Jin, F. Li, T. Shu, J. Coleman, J. Yang, F. Pincet, Y. Zhang, J.E. Rothman, and S.S. Krishnakumar. 2021. Munc13 binds and recruits SNAP25 to chaperone SNARE complex assembly. *FEBS Lett*. 595:297-309.

Kedar, G.H., A.S. Munch, J.R. van Weering, J. Malsam, A. Scheutzow, H. de Wit, S. Houy, B. Tawfik, T.H. Sollner, J.B. Sorensen, and M. Verhage. 2015. A Post-Docking Role of Synaptotagmin 1-C2B Domain Bottom Residues R398/399 in Mouse Chromaffin Cells. *J Neurosci*. 35:14172-14182.

Kim, J.Y., B.K. Choi, M.G. Choi, S.A. Kim, Y. Lai, Y.K. Shin, and N.K. Lee. 2012. Solution single-vesicle assay reveals PIP2-mediated sequential actions of synaptotagmin-1 on SNAREs. *EMBO J*. 31:2144-2155.

Klockner, C., H. Sticht, P. Zacher, B. Popp, H.E. Babcock, D.P. Bakker, K. Barwick, M.V. Bonfert, C.G. Bonnemann, E.H. Brilstra, C. Care4Rare Canada, W.K. Chung, A.J. Clarke, P. Devine, S. Donkervoort, J.L. Fraser, J. Friedman, A. Gates, J. Ghoumid, E. Hobson, G. Horvath, J. Keller-Ramey, B. Keren, M.A. Kurian, V. Lee, K.A. Leppig, J. Lundgren, M.T. McDonald, H.M. McLaughlin, A. McTague, H.C. Mefford, C. Mignot, M.A. Mikati, C. Nava, F.L. Raymond, J.R. Sampson, A. Sanchis-Juan, V. Shashi, J.T.C. Shieh, M. Shinawi, A. Slavotinek, T. Stodberg, N. Stong, J.A. Sullivan, A.C. Taylor, T.L. Toler, M.J. van den Boogaard, S.N. van der Crabben, K.L.I. van Gassen, R.H. van Jaarsveld, J. Van Ziffle, A.F. Wadley, M. Wagner, K. Wigby, S.B. Wortmann, Y.A. Zarate, R.S. Moller, J.R. Lemke, and K. Platzer. 2021. De novo variants in SNAP25 cause an early-onset developmental and epileptic encephalopathy. *Genet Med*. 23:653-660.

Kovacevic, J., G. Maroteaux, D. Schut, M. Loos, M. Dubey, J. Pitsch, E. Remmelink, B. Koopmans, J. Crowley, L.N. Cornelisse, P.F. Sullivan, S. Schoch, R.F. Toonen, O. Stiedl, and M. Verhage. 2018. Protein instability, haploinsufficiency, and cortical hyper-excitability underlie STXBP1 encephalopathy. *Brain*. 141:1350-1374.

Lee, J.S., W.K. Ho, E. Neher, and S.H. Lee. 2013. Superpriming of synaptic vesicles after their recruitment to the readily releasable pool. *Proc Natl Acad Sci U S A*. 110:15079-15084.

Liu, H., C. Dean, C.P. Arthur, M. Dong, and E.R. Chapman. 2009. Autapses and networks of hippocampal neurons exhibit distinct synaptic transmission phenotypes in the absence of synaptotagmin I. *J Neurosci*. 29:7395-7403.

Mahal, L.K., S.M. Sequeira, J.M. Gureasko, and T.H. Sollner. 2002. Calcium-independent stimulation of membrane fusion and SNAREpin formation by synaptotagmin I. *J Cell Biol*. 158:273-282.

Malsam, J., S. Barfuss, T. Trimbuch, F. Zarebidaki, A.F. Sonnen, K. Wild, A. Scheutzow, L. Rohland, M.P. Mayer, I. Sinning, J.A.G. Briggs, C. Rosenmund, and T.H. Sollner. 2020. Complexin Suppresses Spontaneous Exocytosis by Capturing the Membrane-Proximal Regions of VAMP2 and SNAP25. *Cell Rep*. 32:107926.

Malsam, J., D. Parisotto, T.A. Bharat, A. Scheutzow, J.M. Krause, J.A. Briggs, and T.H. Sollner. 2012. Complexin arrests a pool of docked vesicles for fast Ca2+-dependent release. *EMBO J*. 31:3270-3281.

Manca, F., F. Pincet, L. Truskinovsky, J.E. Rothman, L. Foret, and M. Caruel. 2019. SNARE machinery is optimized for ultrafast fusion. *Proc Natl Acad Sci U S A*. 116:2435-2442.

Martin, S., A. Papadopulos, V.M. Tomatis, E. Sierecki, N.T. Malintan, R.S. Gormal, N. Giles, W.A. Johnston, K. Alexandrov, Y. Gambin, B.M. Collins, and F.A. Meunier. 2014. Increased polyubiquitination and proteasomal degradation of a Munc18-1 disease-linked mutant causes temperature-sensitive defect in exocytosis. *Cell Rep*. 9:206-218.

Mirdita, M., K. Schutze, Y. Moriwaki, L. Heo, S. Ovchinnikov, and M. Steinegger. 2022. ColabFold: making protein folding accessible to all. *Nat Methods*. 19:679-682.

Mohrmann, R., H. de Wit, M. Verhage, E. Neher, and J.B. Sorensen. 2010. Fast vesicle fusion in living cells requires at least three SNARE complexes. *Science*. 330:502-505.

Moulder, K.L., and S. Mennerick. 2005. Reluctant vesicles contribute to the total readily releasable pool in glutamatergic hippocampal neurons. *J Neurosci*. 25:3842-3850.

Naldini, L., U. Blomer, P. Gallay, D. Ory, R. Mulligan, F.H. Gage, I.M. Verma, and D. Trono. 1996. In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science*. 272:263-267.

Neher, E. 2015. Merits and Limitations of Vesicle Pool Models in View of Heterogeneous Populations of Synaptic Vesicles. *Neuron*. 87:1131-1142.

Parisotto, D., J. Malsam, A. Scheutzow, J.M. Krause, and T.H. Sollner. 2012. SNAREpin assembly by Munc18-1 requires previous vesicle docking by synaptotagmin 1. *J Biol Chem*. 287:31041-31049.

Parlati, F., T. Weber, J.A. McNew, B. Westermann, T.H. Sollner, and J.E. Rothman. 1999. Rapid and efficient fusion of phospholipid vesicles by the alpha-helical core of a SNARE complex in the absence of an N-terminal regulatory domain. *Proc Natl Acad Sci U S A*. 96:12565-12570.

Peng, L., H. Liu, H. Ruan, W.H. Tepp, W.H. Stoothoff, R.H. Brown, E.A. Johnson, W.D. Yao, S.C. Zhang, and M. Dong. 2013. Cytotoxicity of botulinum neurotoxins reveals a direct role of syntaxin 1 and SNAP-25 in neuron survival. *Nat Commun*. 4:1472.

Radecke, J., R. Seeger, A. Kadkova, U. Laugks, A. Khosrozadeh, K.N. Goldie, V. Lucic, J.B. Sorensen, and B. Zuber. 2023. Morphofunctional changes at the active zone during synaptic vesicle exocytosis. *EMBO Rep*:e55719.

Rebane, A.A., B. Wang, L. Ma, H. Qu, J. Coleman, S. Krishnakumar, J.E. Rothman, and Y. Zhang. 2018. Two Disease-Causing SNAP-25B Mutations Selectively Impair SNARE C-terminal Assembly. *J Mol Biol*. 430:479-490.

Rhee, J.S., L.Y. Li, O.H. Shin, J.C. Rah, J. Rizo, T.C. Sudhof, and C. Rosenmund. 2005. Augmenting neurotransmitter release by enhancing the apparent Ca2+ affinity of synaptotagmin 1. *Proc Natl Acad Sci U S A*. 102:18664-18669.

Rizo, J. 2022. Molecular Mechanisms Underlying Neurotransmitter Release. *Annu Rev Biophys*. 51:377-408.

Rohena, L., J. Neidich, M. Truitt Cho, K.D. Gonzalez, S. Tang, O. Devinsky, and W.K. Chung. 2013. Mutation in SNAP25 as a novel genetic cause of epilepsy and intellectual disability. *Rare Dis*. 1:e26314.

Rosenmund, C., and C.F. Stevens. 1996. Definition of the readily releasable pool of vesicles at hippocampal synapses. *Neuron*. 16:1197-1207.

Ruiter, M., A. Kadkova, A. Scheutzow, J. Malsam, T.H. Sollner, and J.B. Sorensen. 2019. An Electrostatic Energy Barrier for SNARE-Dependent Spontaneous and Evoked Synaptic Transmission. *Cell Rep*. 26:2340-2352 e2345.

Saitsu, H., M. Kato, T. Mizuguchi, K. Hamada, H. Osaka, J. Tohyama, K. Uruno, S. Kumada, K. Nishiyama, A. Nishimura, I. Okada, Y. Yoshimura, S. Hirai, T. Kumada, K. Hayasaka, A. Fukuda, K. Ogata, and N. Matsumoto. 2008. De novo mutations in the gene encoding STXBP1 (MUNC18-1) cause early infantile epileptic encephalopathy. *Nat Genet*. 40:782-788.

Santos, T.C., K. Wierda, J.H. Broeke, R.F. Toonen, and M. Verhage. 2017. Early Golgi Abnormalities and Neurodegeneration upon Loss of Presynaptic Proteins Munc18-1, Syntaxin-1, or SNAP-25. *J Neurosci*. 37:4525-4539.

Schmitz, S.K., J.J. Hjorth, R.M. Joemai, R. Wijntjes, S. Eijgenraam, P. de Bruijn, C. Georgiou, A.P. de Jong, A. van Ooyen, M. Verhage, L.N. Cornelisse, R.F. Toonen, and W.J. Veldkamp. 2011. Automated analysis of neuronal morphology, synapse number and synaptic recruitment. *J Neurosci Methods*. 195:185-193.

Schollmeier, Y., J.M. Krause, S. Kreye, J. Malsam, and T.H. Sollner. 2011. Resolving the function of distinct Munc18-1/SNARE protein interaction modes in a reconstituted membrane fusion assay. *J Biol Chem*. 286:30582-30590.

Schotten, S., M. Meijer, A.M. Walter, V. Huson, L. Mamer, L. Kalogreades, M. ter Veer, M. Ruiter, N. Brose, C. Rosenmund, J.B. Sorensen, M. Verhage, and L.N. Cornelisse. 2015. Additive effects on the energy barrier for synaptic vesicle fusion cause supralinear effects on the vesicle fusion rate. *Elife*. 4:e05531.

Schupp, M., J. Malsam, M. Ruiter, A. Scheutzow, K.D. Wierda, T.H. Sollner, and J.B. Sorensen. 2016. Interactions Between SNAP-25 and Synaptotagmin-1 Are Involved in Vesicle Priming, Clamping Spontaneous and Stimulating Evoked Neurotransmission. *J Neurosci*. 36:11865-11880.

1184 Shen, X.M., D. Selcen, J. Brengman, and A.G. Engel. 2014. Mutant SNAP25B causes myasthenia, cortical
1185   hyperexcitability, ataxia, and intellectual disability. *Neurology*. 83:2247-2255.
1186 Shi, L., Q.T. Shen, A. Kiel, J. Wang, H.W. Wang, T.J. Melia, J.E. Rothman, and F. Pincet. 2012. SNARE proteins:
1187   one to fuse and three to keep the nascent fusion pore open. *Science*. 335:1355-1359.
1188 Shu, T., H. Jin, J.E. Rothman, and Y. Zhang. 2020. Munc13-1 MUN domain and Munc18-1 cooperatively
1189   chaperone SNARE assembly through a tetrameric complex. *Proc Natl Acad Sci U S A*. 117:1036-1041.
1190 Sitarska, E., J. Xu, S. Park, X. Liu, B. Quade, K. Stepien, K. Sugita, C.A. Brautigam, S. Sugita, and J. Rizo. 2017.
1191   Autoinhibition of Munc18-1 modulates synaptobrevin binding and helps to enable Munc13-
1192   dependent regulation of membrane fusion. *Elife*. 6.
1193 Sorensen, J.B., K. Wiederhold, E.M. Muller, I. Milosevic, G. Nagy, B.L. de Groot, H. Grubmuller, and D.
1194   Fasshauer. 2006. Sequential N- to C-terminal SNARE complex assembly drives priming and fusion of
1195   secretory vesicles. *EMBO J*. 25:955-966.
1196 Studier, F.W. 2005. Protein production by auto-induction in high density shaking cultures. *Protein Expr Purif*.
1197   41:207-234.
1198 Sudhof, T.C. 2013. Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. *Neuron*.
1199   80:675-690.
1200 Sutton, R.B., D. Fasshauer, R. Jahn, and A.T. Brunger. 1998. Crystal structure of a SNARE complex involved in
1201   synaptic exocytosis at 2.4 A resolution. *Nature*. 395:347-353.
1202 Taschenberger, H., A. Woehler, and E. Neher. 2016. Superpriming of synaptic vesicles as a common basis for
1203   intersynapse variability and modulation of synaptic strength. *Proc Natl Acad Sci U S A*. 113:E4548-
1204   4557.
1205 Verhage, M., and J.B. Sorensen. 2020. SNAREopathies: Diversity in Mechanisms and Symptoms. *Neuron*.
1206   107:22-37.
1207 Vevea, J.D., and E.R. Chapman. 2020. Acute disruption of the synaptic vesicle membrane protein
1208   synaptotagmin 1 using knockoff in mouse hippocampal neurons. *Elife*. 9.
1209 Voleti, R., K. Jaczynska, and J. Rizo. 2020. Ca(2+)-dependent release of synaptotagmin-1 from the SNARE
1210   complex on phosphatidylinositol 4,5-bisphosphate-containing membranes. *Elife*. 9.
1211 Walter, A.M., P.S. Pinheiro, M. Verhage, and J.B. Sorensen. 2013. A sequential vesicle pool model with a single
1212   release sensor and a Ca(2+)-dependent priming catalyst effectively explains Ca(2+)-dependent
1213   properties of neurosecretion. *PLoS Comput Biol*. 9:e1003362.
1214 Wang, S., Y. Li, J. Gong, S. Ye, X. Yang, R. Zhang, and C. Ma. 2019. Munc18 and Munc13 serve as a functional
1215   template to orchestrate neuronal SNARE complex assembly. *Nat Commun*. 10:69.
1216 Washbourne, P., P.M. Thompson, M. Carta, E.T. Costa, J.R. Mathews, G. Lopez-Bendito, Z. Molnar, M.W.
1217   Becher, C.F. Valenzuela, L.D. Partridge, and M.C. Wilson. 2002. Genetic ablation of the t-SNARE SNAP-
1218   25 distinguishes mechanisms of neuroexocytosis. *Nat Neurosci*. 5:19-26.
1219 Weber, J.P., K. Reim, and J.B. Sorensen. 2010. Opposing functions of two sub-domains of the SNARE-complex
1220   in neurotransmission. *EMBO J*. 29:2477-2490.
1221 Weber, T., B.V. Zemelman, J.A. McNew, B. Westermann, M. Gmachl, F. Parlati, T.H. Söllner, and J.E. Rothman.
1222   1998. SNAREpins: minimal machinery for membrane fusion. *Cell*. 92:759-772.
1223 Wierda, K.D., and J.B. Sorensen. 2014. Innervation by a GABAergic neuron depresses spontaneous release in
1224   glutamatergic neurons and unveils the clamping phenotype of synaptotagmin-1. *J Neurosci*. 34:2100-
1225   2110.
1226 Xian, J., S. Parthasarathy, S.M. Ruggiero, G. Balagura, E. Fitch, K. Helbig, J. Gan, S. Ganesan, M.C. Kaufman,
1227   C.A. Ellis, D. Lewis-Smith, P. Galer, K. Cunningham, M. O'Brien, M. Cosico, K. Baker, A. Darling, F. Veiga
1228   de Goes, C.M. El Achkar, J.H. Doering, F. Furia, A. Garcia-Cazorla, E. Gardella, L. Geertjens, C. Klein, A.
1229   Kolesnik-Taylor, H. Lammertse, J. Lee, A. Mackie, M. Misra-Isrie, H. Olson, E. Sexton, B. Sheidley, L.
1230   Smith, L. Sotero, H. Stamberger, S. Syrbe, K.M. Thalwitzer, A. van Berkel, M. van Haelst, C. Yuskaitis,
1231   S. Weckhuysen, B. Prosser, C. Son Rigby, S. Demarest, S. Pierce, Y. Zhang, R.S. Moller, H. Bruining, A.
1232   Poduri, F. Zara, M. Verhage, P. Striano, and I. Helbig. 2022. Assessing the landscape of STXBP1-related
1233   disorders in 534 individuals. *Brain*. 145:1668-1683.

1234   Xu, J., Z.P. Pang, O.H. Shin, and T.C. Sudhof. 2009. Synaptotagmin-1 functions as a Ca2+ sensor for
1235   spontaneous release. *Nat Neurosci*. 12:759-766.
1236   Zhou, Q., Y. Lai, T. Bacaj, M. Zhao, A.Y. Lyubimov, M. Uervirojnangkoorn, O.B. Zeldin, A.S. Brewster, N.K.
1237   Sauter, A.E. Cohen, S.M. Soltis, R. Alonso-Mori, M. Chollet, H.T. Lemke, R.A. Pfuetzner, U.B. Choi, W.I.
1238   Weis, J. Diao, T.C. Sudhof, and A.T. Brunger. 2015. Architecture of the synaptotagmin-SNARE
1239   machinery for neuronal exocytosis. *Nature*. 525:62-67.
1240   Zhou, Q., P. Zhou, A.L. Wang, D. Wu, M. Zhao, T.C. Sudhof, and A.T. Brunger. 2017. The primed SNARE-
1241   complexin-synaptotagmin complex for neuronal exocytosis. *Nature*. 548:420-425.
1242   Zucker, R.S., and W.G. Regehr. 2002. Short-term synaptic plasticity. *Annu Rev Physiol*. 64:355-405.

1243

1244

Figure Legends

Figure 1 – Localization of three pathogenic mutations in SNAP25

A	Schematic of the neuronal SNARE complex interacting with C2B domain of synaptotagmin-1 (Syt1; not to scale) via the primary interface. Position of the I67N mutation in the first SNARE domain of SNAP25 is depicted by an asterisk.

B	Interaction site of the C2B domain of Syt1 and SNAP25. Syt1 interacts with SNAP25 both electrostatically (region I and II) and within the hydrophobic patch (HP patch)(Zhou et al., 2015).

C	Position of the disease-linked mutations V48F (hydrophobic patch) and D166Y (region I) in the SNARE complex.

Figure 2 – **Pathogenic SNAP25 mutations compromise neuronal viability, but not synaptogenesis**

A	SNAP25 V48F and D166Y mutations are similarly expressed as the WT SNAP25 protein. EGFP-SNAP25 was overexpressed in neurons from CD1 (wildtype) mice; both endogenous and overexpressed SNAP25 are shown. Valosin-containing protein (VCP) was used as loading control.

B	Representative images of control (WT) and mutant (V48F, D166Y) hippocampal neurons stained by dendritic (MAP2) and synaptic (vGlut1) markers. Displayed is MAP2 staining, representing the cell morphology, in inserts MAP2 staining is depicted in red and vGlut staining in cyan. The scale bar represents 50 μm.

C	Number of synapses per neuron in WT and mutant cells.

D	Total dendritic length of WT and mutant neurons.

E	Cell viability represented as the number of neurons per glia island. ****p <0.0001, **p <0.01, *p<0.05, Brown-Forsythe ANOVA test with Dunnett's multiple comparisons test.

Figure 3 –V48F and D166Y mutations increase miniature EPSC frequency

A,D,G	Example traces of mEPSC release for WT, mutant and 1:1 co-expression of WT and mutant SNAP25, or (**G**) Syt1 WT and KO.

B, E	The mEPSC frequencies were increased in both V48F and D166Y mutants and co-expressed conditions (V48F: n = 49, 47, 48 for WT, coexpressed and mutant conditions, respectively; D166Y: n = 54, 43, 50). ****p <0.0001, ***p <0.001, Brown-Forsythe ANOVA test with Dunnett's multiple comparisons test.

C, F	mEPSC amplitudes were on average increased by the V48F and D166Y mutations; this was significant for the V48F. *p <0.05, ANOVA with Dunnett's multiple comparison test.

H, I	Syt1 WT and KO data (Syt1: n = 28, 26 for the WT and KO condition). The mEPSC frequencies and amplitudes were increased and decreased in the KO, respectively. ****p <0.0001, Welch's t-test, *p<0.05, unpaired t-test.

Figure 4 – V48F and D166Y mutations reduce the amplitude of the EPSC

A, E, I	Example evoked excitatory post-synaptic currents (eEPSC) for WT, SNAP25 mutants and co-expressed WT/mutants, or (**I**) Syt1 WT and KO.

B, F, J	EPSC amplitude was decreased by both SNAP25 mutations (V48F: n = 50, 50, 45 for WT, co-expressed and mutant conditions, respectively; D166Y: n = 56, 35, 44) and by Syt1 KO (Syt1: n = 19,

26 for the WT and KO condition). SNAP25 mutations: ****p <0.0001, **p <0.01, Brown-Forsythe
ANOVA test with Dunnett's multiple comparisons test; Syt1: ****p<0.0001, Welch's t-test.
C, G, K Overall evoked charge after a single depolarization (V48F: n = 50, 45, 50 for WT, mutant
and co-expressed conditions, respectively; D166Y: 56, 44, 35; Syt1: 19, 20 for WT and KO). SNAP25:
*p <0.05, Brown-Forsythe ANOVA with Dunnett's multiple comparison test; Syt1: ****p<0.0001,
Welch's t-test.
D, H, L Fractional contribution of the synchronous release component to the overall charge (V48F:
n = 50, 50, 45 for WT, co-expressed and mutant conditions, respectively; D166Y: 56, 35, 44; Syt1:
19, 20 for WT and KO). SNAP25: ****p <0.0001, ***p <0.001, Brown-Forsythe ANOVA (V48F) or
standard ANOVA (D166Y) with Dunnett's multiple comparisons test; Syt1: ****p<0.0001, Welch's t-
test.
1297
Figure 5 – The apparent energy barrier for vesicle fusion is lowered by V48F and D166Y, but not
by removing Syt1.
A, F, K Example traces for the WT, mutant and co-expressed condition. Each cell was stimulated
by 0.25 M (in grey) and 0.5 M sucrose (in black or color).
B, G, L The charge released by 0.25 M sucrose (V48F: n = 28, 30, 29 for WT, co-expressed and
mutant conditions, respectively; D166Y: n = 33, 30, 35; Syt1: n = 23, 18 for WT and KO). Syt1: p<0.05,
Welch's t-test.
C, H, M The charge released by 0.5 M sucrose (V48F: n = 28, 30, 29 for WT, co-expressed and
mutant conditions, respectively; D166Y: n = 33, 30, 35; Syt1: n = 23, 26 for WT and KO). SNAP25:
****p <0.0001, **p <0.01, *p<0.05, Brown-Forsythe ANOVA with Dunnett's multiple comparisons
test; Syt1: p=0.0548, unpaired t-test.
D, I, N The ratio of 0.25 and 0.5 M sucrose pool (V48F: n = 28, 30, 29 for WT, co-expressed and
mutant conditions, respectively; D166Y: n = 33, 30, 35; Syt1: n = 23, 18 for WT and KO). SNAP25:
****p <0.0001, **p <0.01, ANOVA with Dunnett's multiple comparisons test; Syt1: *p < 0.05,
unpaired t-test.
E, J, O Release probability calculated by dividing the charge of an eEPSC with the 0.5 M sucrose
pool (V48F: n = 24, 25, 22 for WT, co-expressed and mutant conditions, respectively; D166Y: n = 33,
24, 30; Syt1, n = 16, 21 for WT and KO). SNAP25: ***p <0.001, ANOVA with Dunnett's multiple
comparisons test; Syt1: ****p<0.0001, unpaired t-test.
1317
Figure 6 – Dissection of the RRP reduction in V48F and D166Y mutations.
A One-pool model of the Readily Releasable Pool (RRP). k_1 is the rate of priming (units
vesicles/s), k_{-1} is the rate of depriming (s^{-1}), k_f is the rate of fusion (s^{-1}).
B Estimation of the three parameters from the response to 0.5 M sucrose and a
measurement of the spontaneous release rate.
C Variance-mean analysis in 50 ms intervals during the sucrose application allows
determination of the corrected baseline by back-extrapolation of a regression line to the variance
of the baseline.
D Normalized mEPSC frequency (k_f) for V48F, D166Y and Syt1 KO. (V48F: n = 23, 24 for WT
and mutant conditions, respectively; D166Y: n = 19, 19; Syt1: n = 23, 26). Brown-Forsythe ANOVA

test with Dunnett's multiple comparison test, testing the three mutant conditions against each
other. ****p <0.0001, ***p <0.001.
E Normalized RRP size for WT and mutant conditions, with indications of the effect of the
mutant-induced changes in k_1, k_{-1}, and k_f on the RRP size.

1332

Figure 7 – SNAP25 V48F and D166Y mutations change short-term plasticity towards facilitation.
A, E EPSCs in response to 50 APs at 40 Hz recorded in 4 mM extracellular Ca^{2+} (V48F: 27, 17, 15
for WT, co-expressed and mutant conditions, respectively; D166Y: 27, 18, 16). Inserts: Normalized
EPSC amplitudes demonstrating facilitation of mutant conditions. ****p<0.0001; **p<0.01, Brown-
Forsythe ANOVA with Dunnett's multiple comparison test.
B, F Priming rate calculated as the slope of a linear fit to the cumulative evoked charges during
the last part of stimulation (V48F: 27, 17, 15 for WT, co-expressed and mutant conditions,
respectively; D166Y: 27, 18, 16). *p <0.05, ANOVA (V48F) or Brown-Forsythe ANOVA (D166Y) with
Dunnett's multiple comparisons test.
C, G RRP calculated by back-extrapolation of a linear fit to the cumulative evoked charges during
the last part of stimulation (V48F: 27, 17, 15 for WT, co-expressed and mutant conditions,
respectively; D166Y: 27, 18, 16). **p <0.01, *p <0.05, ANOVA (V48F) or Brown-Forsythe ANOVA
(D166Y) with Dunnett's multiple comparisons test.
D, H Release probability calculated as the charge of the first evoked response divided by the RRP
obtained by back-extrapolation (V48F: 27, 17, 15 for WT, co-expressed and mutant conditions,
respectively; D166Y: 27, 18, 16). ***p<0.001; **p <0.01, ANOVA (V48F) or Brown-Forsythe ANOVA
(D166Y) with Dunnett's multiple comparisons test.

1350

Figure 8 – Pathogenic SNAP25 mutations affect synaptotagmin-1 interaction and fusion rates *in*
***vitro*.**
A In the presence of SDS, SNAP25 I67N containing v-/t-SNARE complexes were more sensitive
to temperature-dependent dissociation. Shown are mean ± SEM (n = 3) for SNARE-complexes
including SNAP25 WT and the I67N, V48F and D166Y mutations.
B-C *In vitro* Syt1/VAMP2 SUVs docking to t-SNARE GUVs was significantly reduced by SNAP25
V48F, I67N and D166Y mutants either in absence (**B**) or presence (**C**) of PI(4,5)P$_2$. Fusion was blocked
by performing the assay on ice.
D-E *In vitro* lipid mixing assays of VAMP/Syt1 SUVs with t-SNARE GUVs containing SNAP25 V48F,
I67N, or D166Y mutants showed impaired membrane fusion in the absence (left) or presence (right)
of complexin-II. Fusion clamping in the presence of complexin was selectively reduced by V48F and
D166Y. Bar diagrams show lipid mixing just before (pre) and after (post) Ca^{2+} addition and at the end
of the reaction. Shown is mean ± SEM (n = 3).

1364

Figure 9 – The D166Y mutation increases binding to its SNARE partners.
A-C *In vitro* lipid mixing assays of VAMP/Syt1 SUVs with syntaxin-1A GUVs in the presence of
soluble SNAP25. V48F, and D166Y mutants showed impaired fusion clamping in the absence (left)
or presence (right) of complexin-II; I67N (red) showed impaired Ca^{2+}-independent and Ca^{2+}-

triggered fusion. Bar diagrams show lipid mixing just before (pre) and after (post) Ca^{2+} addition and
at the end of the reaction. Mean $\pm$ SEM (n = 3).

D SNAP25 D166Y showed enhanced interactions with SUVS carrying reconstituted syntaxin-
1A (Stx-1), VAMP2, Syt1/VAMP2 or a SUV-mixture containing Syntaxin-1A and VAMP2/Syt1 in co-
flotation assays, whereas V48F displayed weaker increases in interactions with SUVs containing
syntaxin-1A, or Syt1/VAMP2. Shown is mean $\pm$ SEM on a logarithmic scale. *** p<0.001, ** p<0.01,
* p<0.05, two-tailed one-sample t-test comparing to 1.

1376

Figure 10 – The I67N mutation inhibits spontaneous and evoked release.

A SNAP25 I67N is similarly expressed as the WT SNAP25 protein. EGFP-SNAP25 was
overexpressed in neurons from CD1 (wildtype) mice; both endogenous and overexpressed SNAP25
are shown. Valosin-containing protein (VCP) was used as the loading control.

B Representative image of mutant (I67N) hippocampal neurons stained for the dendritic
marker MAP2 and the synaptic markers vGlut1. Displayed is MAP2 staining, representing the cell
morphology, in inserts MAP2 staining is depicted in red and vGlut staining in cyan. The scale bar
represents 50 µm.

C Cell viability represented as the number of neurons per glial islet. ****p <0.0001, Brown-
Forsythe ANOVA test with Dunnett's multiple comparisons test.

D Number of synapses per neuron in WT and mutant cells. The WT data are the same as in
Fig. 2B because these experiments were carried out in parallel. The difference was tested using
ANOVA between all conditions, which was non-significant.

E Total dendritic length of WT and mutant neurons.

F Example traces of mEPSC release for WT, mutant (I67N) and 1:1 co-expression of WT and
SNAP25 mutant.

G The mini frequency was decreased in both I67N mutant and the WT+I67N combination
(I67N: n = 39, 36, 30 for WT, coexpressed and mutant). ****p <0.0001, ***p <0.001, Kruskal-Wallis
with Dunn's multiple comparisons.

H mEPSC amplitudes were unchanged by the I67N mutation.

I Example evoked excitatory post-synaptic currents (eEPSC) for WT, mutant (I67N) and co-
expressed WT and mutant.

J EPSC amplitude was decreased by the I67N mutations (I67N: n = 39, 37, 30 for WT, co-
expressed and mutant conditions, respectively). SNAP25 mutations: ****p <0.0001, **p <0.01,
Brown-Forsythe ANOVA test with Dunnett's multiple comparisons test.

K Overall evoked charge after a single depolarization (I67N: 24, 10, 0 for WT, co-expressed
and mutant conditions, respectively).

L Fractional contribution of the synchronous release component to the overall charge (I67N:
24, 10, 0 for WT, co-expressed and mutant conditions, respectively).

1406

1407 **Figure 11 – The I67N mutation has normal RRP size, but increased energy barrier for fusion.**

1408 **A, E** Example traces for the WT, mutant and co-expressed condition. Each cell was stimulated
1409 by 0.25 M (**A**, in grey) and 0.5 M sucrose (**A**, in color) or 0.375 M sucrose (**E**, in grey) and 0.75 M (**E**,
1410 in color).

B, F The charge released by 0.25 M sucrose (**B**, I67N: n = 21, 15, 8 for WT, co-expressed and mutant conditions, respectively) or 0.375 M sucrose (**F**, I67N: n = 12, 16, 18; a few cells were stimulated with 0.35 M sucrose – shown with open symbols). **B**, **p<0.01; *p<0.05, Kruskal-Wallis test with Dunn's multiple comparison test; **F**, p=0.0339 Brown Forsythe ANOVA test; Dunnett's multiple comparison test, p=0.0571.

C, G The charge released by 0.5 M sucrose (**C**, I67N: n = 21, 15, 8 for WT, co-expressed and mutant conditions, respectively), or 0.75 M sucrose (**G**, I67N: n = 13, 16, 18). **C**, **p<0.01, *p<0.05, Brown-Forsythe ANOVA with Dunnett's multiple comparisons test.

D, H The ratio of the 0.25 and 0.5 M sucrose pool (**D**, I67N: n = 21, 15, 8 for WT, coexpressed and mutant conditions, respectively), or the ratio of 0.375 and 0.75 M sucrose pool (**H**, n = 13, 16, 18). **D**, * p<0.05, Kruskal-Wallis test with Dunn's multiple comparisons test. **H**, **** p<0.0001; *** p<0.001, ANOVA with Dunnett's multiple comparisons test.

I EPSCs in response to 50 APs at 40 Hz recorded in 2 mM extracellular Ca^{2+} (I67N: n = 23, 16, 20 for WT, co-expressed and mutant conditions, respectively). Inserts: Normalized EPSC amplitudes demonstrating facilitation of mutant conditions.

J Normalized EPSC amplitudes in response to 50 APs at 40 Hz recorded in 2 mM extracellular Ca^{2+}.

K Paired pulse ratio at interstimulus interval 25 ms (I67N: n = 24, 14, 17 for WT, co-expressed and mutant conditions, respectively). *p<0.05, ANOVA with Dunnett's multiple comparison test.

Figure 12 - Adding positive surface charges to the SNARE complex partly compensate for the I67N mutation.

A Example traces of mEPSC release for WT, I67N/E183K/S187K/T190K/E194K ('I67N/4K') and E183K/S187K/T190K/E194K ('4K') SNAP25. Data from the 4K mutation was obtained in a separate experiment and is shown for comparison, but statistical tests with 4K mutation data were not carried out.

B The mini frequencies for the WT and I67N/4K are not significantly different; data from the 4K mutation is shown for comparison (n = 19, 25, 13 for WT, I67N/4K and 4K, respectively).

C Mini amplitudes remain unaffected by I67N/4K mutation.

D-E eEPSC examples (**D**) and amplitudes (**E**) for WT and I67N/4K; 4K is shown for comparison (n = 19, 25, 13 for WT, I67N/4K and 4K, respectively). ****p<0.0001 Mann-Whitney test.

F Electrostatic triggering model (blue line; Ruiter et al., 2019) refitted to WT spontaneous and evoked data points (black points). Fitted parameters: rate 0.00029 s^{-1} at zero (0) charge (Z); fraction f=0.030; the maximum rate was fixed at 6000 s^{-1}. WT (black points), I67N, I67N/4K (red points): means of log-transformed data. The charge-values (Z, horizontal axis) for I67N and I67N/4K were found by interpolation in the model; the two spontaneous points (I67N, I67N/4K) are separated by 5.6 charges. For evoked release, rates were found by deconvolution and normalizing to $RRP_{0.5}$ (Ruiter et al., 2019). The Z-values for evoked release were found by interpolation in the model; the two mutants (I67N, I67N/4K) are separated by 5.9 charges.

Figure 13 – Energy landscapes.
The energy landscapes of WT and mutants were calculated as explained in Materials and Methods and displayed to scale. Energy landscapes for D166Y (**A**), V48F (**B**) and *Syt1* KO (**C**) are shown at rest and are characterized by a higher priming barrier ("loss-of-function" phenotype), a destabilized RRP

1455 and a lower fusion barrier ("gain-of-function" phenotype). The I67N (**D**) is characterized by a higher
1456 fusion barrier ("loss-of-function" phenotype). The relative increase in the fusion barrier by the I67N
1457 mutation is higher during stimulation than at rest. Dotted lines represent energy levels for which
1458 less is known.
1459
Figure 4 – Figure Supplement 1. Kinetic parameters of evoked EPSCs.
A   eEPSC (black trace), and integrated eEPSC (after multiplication with -1, red trace) with
double exponential fit (green trace).
B   Zoom-in of eEPSC (black trace), and integrated EPSC (after multiplication with -1, red trace)
with double exponential fit (green trace). Equations for integration and double exponential function
used for fit is given.
1466
Figure 4 – Figure Supplement 2. Kinetic parameters of evoked EPSCs.
A, E, I   Synchronous release components (**A**, V48F: n = 50, 50, 45 for WT, co-expressed and mutant
conditions, respectively; **E**, D166Y: n = 56, 35, 44; **I**, *Syt1*: 19, 20 for WT and KO). **A**: * $p < 0.05$, Welch's
ANOVA with Dunnett's multiple comparison test, **E**: *** $p < 0.001$, Brown-Forsythe ANOVA with
Dunnett's multiple comparison test, **I**: **** $p < 0.0001$, Welch's unpaired t-test.
B, F, J   Asynchronous release components. **B**: ** $p < 0.01$; * $p < 0.05$, Brown-Forsythe ANOVA with
Dunnett's multiple comparison test.
C, G, K   Fast time constants. **C**: **** $p < 0.0001$; ** $p < 0.01$, Brown-Forsythe ANOVA with Dunnett's
multiple comparison test, **G**: **** $p < 0.0001$, Brown-Forsythe ANOVA with Dunnett's multiple
comparison test, **K**: **** $p < 0.0001$, Welch's unpaired t-test.
D, H, L   Slow time constants.
1478
Figure 6 – Figure Supplement 1. Effect of sucrose stimulation on estimates of k_1 and k_{-1}.
The figure shows the effect of the fold-increase in fusion rate (*N*) induced by sucrose (abscissa) on
the estimates of k_1 (**A**, **C**) or k_{-1} (**B**, **D**) using Eqs. 3, 4 and 5b and the values estimated for D166Y (**A**,
B), V48F (**C**, **D**) and WT (Table 1). Previous data showed that 0.5 M sucrose increases the fusion rate
by a factor ~5000 (Schotten et al., 2015). The plots show that the estimates in Table 1 are not strongly
affected by small changes in the effect of sucrose upon k_f.
1485
Figure 7 – Figure Supplement 1. Train stimulations of V48F and D166Y in 2 mM Ca^{2+}.
A, D   EPSCs in response to 50 APs at 40 Hz recorded in 2 mM extracellular Ca^{2+} (V48F: 25, 18, 24
for WT, co-expressed and mutant conditions, respectively; D166Y: 23, 15, 15). Inserts: Normalized
EPSC amplitudes of first five stimulations. * $p < 0.05$, *** $p < 0.001$, one-way ANOVA with Dunnett's
multiple comparison test.
B, E   Priming rate calculated by as the slope of a linear fit to the cumulative evoked charges
during the last part of stimulation (V48F: 24, 18, 24 for WT, co-expressed and mutant conditions,
respectively; D166Y: 23, 15, 15).
C, F   RRP calculated by back-extrapolation of a linear fit to the cumulative evoked charges during
the last part of stimulation (V48F: 24, 18, 24 for WT, co-expressed and mutant conditions,
respectively; D166Y: 23, 15, 15).

1497

Figure 7 – Figure Supplement 2. Cumulative charges of V48F and D166Y trains in 4 mM Ca^{2+}.

A, B Cumulative charges obtained by integrating EPSCs during 40 Hz trains. The slope of the
linear part of the curve reports on the priming rate, which is reduced by the mutations. The back
extrapolation of the linear fit to zero time reports on the RRP$_{ev}$, the part of the RRP which APs draw
on, which is also reduced by mutation (V48F: n= 27, 17, 15 for WT, co-expressed and mutant
conditions, respectively; D166Y: 27, 18, 16).

Figure 9 – Figure Supplement 1. Floatation assay.
Example Coomassie and silver stained gels demonstrating binding of SNAP25 WT and mutants to
different populations of SUVs: Syntaxin-1 (Stx-1) and VAMP2/Syt1, Syntaxin-1 (Stx-1), VAMP2/Syt1,
VAMP2, or Syt1 SUVs. Note increased binding of D166Y and V48F to most SUV populations, strongest
for D166Y (quantified data in Fig. 9D).
1509

1510 **Figure 9 – Figure Supplement 2. AlphaFold prediction of mutants.**
1511 The figure shows an overlay of the WT protein and the three mutants I67N, V48F and D166Y as
1512 predicted by AlphaFold (Jumper et al., 2021; Mirdita et al., 2022). No major structural changes were
1513 detected, but V48F is expected to alter the interaction with neighboring E52, whereas D166Y will
1514 induce a stronger interaction with H162.

Kádková et al., Fig. 1

Kádková et al., Fig. 2

Kádková et al., Fig. 3

Kádková et al., Fig. 4

Kádková et al., Fig. 5

Kádková et al., Fig. 6

Kádková et al., Fig. 7

Kádková et al., Fig. 8

Kádková et al., Fig. 9

Kádková et al., Fig. 10

Kádková et al., Fig. 11

Kádková et al., Fig. 12

Kádkova et al., Fig. 13

Kádková et al., Fig. 4 - Figure supplement 1

Kádková et al., Fig. 4 - Figure supplement 2

Kádková et al., Fig. 6 - Figure supplement 1

Kádková et al., Fig. 7 - Figure supplement 1

Kádková et al., Fig. 7 - Figure supplement 2

Kádková et al., Fig. 9 - Figure supplement 1

WT
I67N
V48F
D166Y

F48
V48
E52

V48

D166
H162
D166
Y166

Kádková et al., Fig. 9 - Figure supplement 2