Dependent variable selection in phylogenetic generalized least squares

regression analysis under Pagel's lambda model

Zheng-Lin Chen[1,2], Hong-Ji Guo[1,2], and Deng-Ke Niu[1,2]

[1]MOE Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life

Sciences, Beijing Normal University, Beijing, 100875, China

[2]Beijing Key Laboratory of Gene Resource and Molecular Development, College of Life

Sciences, Beijing Normal University, Beijing, 100875, China

Correspondence

Deng-Ke Niu

Email: dkniu@bnu.edu.cn, dengkeniu@hotmail.com.

Running title: Dependent variable selection in PGLS analysis

Abstract

Phylogenetic generalized least squares (PGLS) regression is widely used to examine evolutionary associations while accounting for phylogenetic non-independence, but it requires designating one trait as the dependent variable and the other as the independent variable. When causal relationships between traits are unclear, choosing which trait serves as the dependent variable becomes a practical concern. While studying the evolutionary relationship between bacterial growth rate and CRISPR-Cas content, we noticed that switching the roles of dependent and independent variables in PGLS analyses could lead to inconsistent conclusions in a substantial proportion of cases. To confirm this observation, we conducted 16,000 simulations of trait evolution along binary trees with 100 terminal nodes under different evolutionary models. PGLS regressions using Pagel's λ model were applied to each simulation, and the results consistently showed that swapping the dependent and independent variables can lead to inconsistent outcomes. We evaluated seven potential criteria for selecting the dependent variable, including log-likelihood, AIC, R², p-value, Pagel's λ, Blomberg's K, and the estimated λ in the Pagel's λ model. Among these, Pagel's λ, Blomberg's K, and the estimated λ performed equally well and outperformed the others in selecting the dependent variable, providing a reliable basis for PGLS analyses when causal direction between traits is unclear.

Keywords

Evolutionary correlation, dependent variable, phylogenetic comparative methods, phylogenetic generalized least squares (PGLS), phylogenetic signal.

1 INTRODUCTION

Through mutations, genetic drift, and natural selection, biological traits often co-evolve along the phylogeny of species (Felsenstein 1985; Revell & Collar 2009; Goswami *et al.,* 2014; Revell *et al.,* 2022). Studying correlations among such traits can provide insights into how changes in one characteristic may accompany evolutionary changes in another, offering support for or challenges to specific biological hypotheses (Pearman *et al.,* 2014). Understanding these correlations contributes to reconstructing evolutionary histories and uncovering underlying biological processes (Bartomeus *et al.,* 2018; Bawa *et al.,* 2019; Suarez-Castro *et al.,* 2020).

A common approach for assessing trait associations is through correlation analysis,

using Pearson, Spearman, or Kendall coefficients. However, conventional statistical methods assume independence among observations—a condition violated in comparative biological data, where species share evolutionary history. Ignoring phylogenetic non-independence can lead to inflated or distorted correlation estimates (Revell 2010; Whitney & Garland 2010). To address this, phylogenetic comparative methods (PCMs) have been developed, explicitly accounting for the phylogenetic structure of the data (Felsenstein 1985; Grafen 1989; Lynch 1991; Garamszegi 2014; Cooper *et al.,* 2016).

Among PCMs, phylogenetic generalized least squares (PGLS) has become widely used to study associations between traits. Originally proposed by Grafen (1989) and refined by Martins and Hansen (1997), Pagel (1997; 1999), and Rohlf (2001), PGLS extends ordinary least squares (OLS) regression by incorporating the expected covariance among species due to shared ancestry. Despite being fundamentally a regression method, PGLS is frequently used to examine trait correlations, likely due to the lack of simple and flexible tools for phylogenetically controlled correlation analysis. While other approaches such as phylogenetic independent contrasts (PIC) can also estimate correlations (Felsenstein 1985), PIC is limited to traits evolving under a Brownian motion (BM) model and cannot accommodate traits with more complex evolutionary dynamics.

Regression-based methods inherit OLS assumptions, including that one variable is the independent variable (predictor) and the other is the dependent variable (response), implying a causal direction (Waugh 1943). In practice, however, the causal relationships among many biological traits are often uncertain or debated for decades. For example, the relationship between GC content and environmental temperature has been discussed in multiple ways: one hypothesis suggests GC content affects DNA double-strand stability, influencing thermal tolerance, while another emphasizes that temperature-induced DNA damage may not produce symmetric changes in GC-AT content(Travnicek *et al.,* 2019; Hu *et al.,* 2022). Each perspective can support an alternative causal direction, and from a biological standpoint, there is often no basis to prioritize one over the other.

When applying PGLS to study the correlation between two traits with unclear causal relationships, the assignment of dependent and independent variables may not matter if swapping them does not change the outcome—i.e., the sign (positive or negative) of the regression coefficient or the significance test results ($p <$ or ≥ 0.05). In practice, however, we found that using PGLS with Pagel's λ model (Freckleton et al., 2002) on simulated data, swapping dependent and independent variables could substantially affect the results: correlations that were significant in one configuration sometimes became non-significant in

the other, and the sign of the regression coefficient could also change. This finding underscores that, in practical applications, careful consideration of variable assignment is necessary for robust inference and motivates further investigation into criteria for selecting dependent and independent variables in PGLS.

However, in a more comprehensive study of the correlation between bacterial growth rate and CRISPR spacer content in genomes (Liu *et al.*, 2023) using Pagel's λ model (Freckleton *et al.*, 2002), we found that switching the dependent and independent variables could substantially affect the results: correlations that were significant ($p < 0.05$) in one configuration sometimes became non-significant ($p \geq 0.05$) in the other. These observations underscore that, in practical applications, careful consideration of variable assignment is necessary for robust inference and motivate further investigation into criteria for selecting dependent and independent variables in PGLS.

Using simulated data, we further assessed the frequency of inconsistent outcomes caused by switching independent and dependent variables in the PGLS analysis. In addition, we compared seven criteria to determine the appropriate designation of independent and dependent variables in the PGLS analysis using Pagel's λ model (Freckleton *et al.*, 2002).

2 MATERIALS AND METHODS

2.1 Empirical datasets

A dataset including the empirical minimal doubling times, CRISPR spacer numbers, optimal growth temperature, and prophage numbers of 262 bacteria was extracted from the Supplemental Material Table S1 of Liu et al. (2023). The phylogenetic tree of these 262 bacteria was retrieved from the Genome Taxonomy Database (GTDB; accessed April 8, 2022) (Parks *et al.*, 2022).

2.2 Simulation data

First, we generated a binary tree containing 100 terminal nodes using the R package ape (Paradis & Schliep 2019). The trait X_1 evolved according to a Brownian motion model (using the R package ape) with a variance rate $\sigma^2{}_{BM} = 4$ along the tree, and then the trait X_2 was simulated according to

$$X_2 = X_1 + \varepsilon, \tag{1}$$

where ε was a normally distributed random noise term with a mean of 0 and a variance of 10^{-4}, 0.01, 1, 4, 16, 64, 256 to 1024. The term ε introduced noise to the dependent variable

X_2 and a gradient variance of this noise term (from 10^{-4} to 1024) changed the correlation

between X_1 and X_2, from strong to weak. We named this case "BM & BM + Norm."

Then, we simulated the trait X_1 according to a normal distribution with a mean of 0

and a variance of 4. The trait X_2 was simulated according to equation (1), where ε was

simulated according to a Brownian motion model with $\sigma^2{}_{BM}$ in a range of 10^{-4}, 0.01, 1, 4,

16, 64, 256, 1024. We named this case "Norm & Norm + BM."

We ran 1000 simulations for each parameter condition, and a total of 16,000 simulations

were performed.

2.3 Software and packages

All PGLS regressions in this study were performed using the R (version 4.0.2) package

phylolm (version 2.6.2) (Ho & Ane 2014). The estimated λ ($\hat{\lambda}$), log-likelihood (LLK)

(Freckleton *et al.,* 2002), Akaike information criterion (AIC), R^2 of the models, and the *p*-

value of the regression coefficients were calculated in the PGLS regression analyses.

The phylogenetic signals (λ and K) were estimated using the R (version 4.0.2) package

phytools (version 1.0-3) (Revell 2012).

3 RESULTS

3.1 Switching dependent and independent variables in PGLS analyses of empirical data

sometimes gave conflicting outcomes

Using a dataset comprising a range of traits from 262 bacteria in a recent publication

(Liu *et al.,* 2023), we analyzed the correlations of empirical minimal doubling time and

optimal growth temperature with genomic characteristics. Thirty-eight pairs of traits were re-

examined using Pagel's λ model in the PGLS approach. However, switching dependent and

independent variables in 10 pairs resulted in conflicting outcomes, leading to different

conclusions (Table 1). For instance, when considering the average prophage number as the

dependent variable and the optimal growth temperature as the independent variable, the

PGLS analysis showed a significant negative correlation between the two traits ($p = 9 \times 10^{-4}$,

Table 1). In contrast, switching dependent and independent variables showed no statistically

significant correlation ($p = 0.242$, Table 1). Identifying accurate findings rather than

conflicting results is necessary. However, the question is how one can determine which one is

correct.

3.2 Prevalence of conflicts resulting from switching dependent and independent variables

In the above survey on correlations within empirical data, we found that switching dependent and independent variables resulted in conflicting outcomes in 26.3% of cases. To evaluate the prevalence of conflicting results from switching dependent and independent variables, we simulated the progression of two traits along a binary tree of 100 species. Different distributions of the traits were simulated. In the case of "BM & BM + Norm", we constructed the trait X_1 under the Brownian motion model ("BM") and the trait X_2, as "BM + Norm", equals to X_1 plus a noise term "Norm." In the case of "Norm & Norm + BM," the noise term is "BM." To account for varying levels of correlation, we set a gradient of variance in the noise term, with the variance $\sigma^2{}_{Norm}$ varying from 10^{-4}, 0.01, 1, 4, 16, 64, 256, to 1024. For each parameter condition, we simulated 1000 times.

For the data from each simulation, we performed two rounds of PGLS analysis, $X_1 \sim X_2$ and $X_2 \sim X_1$ (representing regressions of X_1 on X_2 and X_2 on X_1, respectively). The results are presented in Table S1 and summarized in Table 2. In 3781 simulations, neither $X_1 \sim X_2$ nor $X_2 \sim X_1$.gave significant correlations ($p \geq 0.05$ for all of them). In 10161 simulations, both $X_1 \sim X_2$ and $X_2 \sim X_1$.gave significant correlations ($p < 0.05$). In each of these 10161 cases, the regression coefficients of $X_1 \sim X_2$ and $X_2 \sim X_1$ have the same sign. In the other 2058 simulations, only one of $X_1 \sim X_2$ and $X_2 \sim X_1$.gave significant correlations ($p < 0.05$ for one and $p \geq 0.05$ for the other). The frequency of conflicting results caused by switching dependent and independent variables in our simulated dataset is 12.9%.

Table 2 shows a relationship between the variance in simulations and the frequency of conflicting results caused by switching dependent and independent variables. When the variance of the noise term is slight (like 10^{-4}, 0.01, 1, and 4), i.e., where there is a strong correlation between X_1 and X_2, switching independent and dependent variables gives almost the same results. As the variance in the term noise increases (i.e., where the correlation between X_1 and X_2 becomes weak), switching independent and dependent variables leads to many conflicting results. In cases where the variance is 16 in "BM & BM + Norm" and 64 in "Norm & Norm + BM," there are even close to 50% of cases with conflicting results. However, as the variance becomes much more pronounced and the correlation between X_1 and X_2 becomes much weaker, the frequency of conflicting results caused by switching dependent and independent variables decreases (Table 2).

3.3 Estimated lambda may change according to the choice of the dependent variable

To control the phylogenetic dependence in the data, PGLS assumes that the model

residual error is distributed according to the variance-covariance matrix representing the

residual variance and the evolutionary relationships among species. In its simplest form,

PGLS assumes a Brownian motion model of evolution at a single rate. By parameter

estimation and altering the properties of the variance-covariance matrix, PGLS can be

extended to various evolutionary models, like Pagel's λ model (Freckleton *et al.,* 2002),

Ornstein–Uhlenbeck model (Hansen 1997), and the early burst model (Harmon *et al.,* 2010).

We suspect that switching independent and dependent variables in the PGLS analysis may

result in inconsistent estimates of the parameters (like Pagel's λ) and so inconsistent

correction for the phylogenetic non-independence of the trait values.

For the data of each simulation, we calculated the phylogenetic signal λ of the two

traits (X_1 and X_2) and the $\hat{\lambda}$ of $X_1 \sim X_2$ and $X_2 \sim X_1$ using Pagel's λ model.

In the "BM & BM + Norm" case, when the variance of the noise term is slight, both

X_1 and X_2 exhibit strong phylogenetic signals with their λ values close to 1, and both

$X_1 \sim X_2$ and $X_2 \sim X_1$ give small $\hat{\lambda}$ values (Figure 1). However, with the increase of the

variance of the noise term, the λ values of X_2 gradually decrease while those of X_1. remain

at their high levels. Together, the $\hat{\lambda}$ values of the $X_1 \sim X_2$ models gradually increase to 1

while those of the $X_2 \sim X_1$ models remain stable at around 0.03 (Figure 1).

In the "Norm & Norm + BM" case, we found a pattern previously reported by Revell

(2010), PGLS regression is possible to give a high $\hat{\lambda}$ value even if both traits have very

weak phylogenetic signals. When the variance of the BM noise term is slight, both X_1 and

X_2 have small λ values but both $X_1 \sim X_2$ and $X_2 \sim X_1$ models yielded high $\hat{\lambda}$ values of

about 1 (Figure 1). With the increase of the variance of the noise term, the phylogenetic

signals of X_2 gradually become stronger and finally approach one, but that of X_1 remains

weak. Meanwhile, the $\hat{\lambda}$ values of $X_1 \sim X_2$ gradually decrease to about 0.03, while those of

$X_2 \sim X_1$ remain close to 1 (Figure 1).

We could see that the two models, $X_1 \sim X_2$ and $X_2 \sim X_1$, generally give similar $\hat{\lambda}$

values when the two variables, X_1 and X_2, have similar λ values (Figure 1). That is,

switching dependent and independent variables would not dramatically affect the parameter

estimation of the PGLS regression analyses when the two analyzed traits have similar levels

of phylogenetic signals. However, when the two variables differ strikingly in their

phylogenetic signals, the two models give quite different $\hat{\lambda}$ values (Figure 1). In this case,

switching dependent and independent variables would dramatically affect the parameter

estimation and potentially give conflicting results. In addition, we noticed that the $\hat{\lambda}$ values

tend to be close to the phylogenetic signal of the dependent variable.

3.4 A golden standard to evaluate the accuracy of PGLS results

When performing PGLS analysis on both empirical and simulated datasets, switching

dependent and independent variables leads to a significant frequency of conflicting results.

Avoiding arbitrary selection of the dependent variable is essential in PGLS analysis using

Pagel's λ model. In order to select a better dependent variable, we will use simulated data to

establish a selection criterion.

In empirical phylogenetic data, we only have access to trait values at terminal nodes,

and potential correlations between traits can be estimated using methods such as PGLS. In

contrast, we can access trait values at internal nodes in simulated phylogenetic data. Since

changes along different phylogenetic branches are independent, we can measure the

evolutionary correlation between two traits by analyzing their changes along phylogenetic

branches using standard statistical methods. First of all, we calculated the changes in the

traits X_1 and X_2 along evolutionary branches per unit time, $\Delta X_1/L$ and $\Delta X_2/L$, where L

is the length of the branch. Then, we did Shapiro-Wilk test on $\Delta X_1/L$ and $\Delta X_2/L$ to see if

they had normal distributions. If both of them satisfy normality, we used Pearson correlation

to detect the correlation between $\Delta X_1/L$ and $\Delta X_2/L$. Otherwise, we used Spearman's rank

correlation. These analyses revealed significant correlations between the two traits, X_1 and

X_2, in 11911 simulations (10988 positives and 923 negatives, $p < 0.05$ for all these cases) but

not in the other 4089 simulations ($p \geq 0.05$ for all these cases) (Table S1). These results

provide "golden standards" for judging whether PGLS analyses of trait values on terminal

nodes (X_1 and X_2) give correct results.

PGLS analyses of $X_1 \sim X_2$ show that there are significant positive correlations in

10326 simulations ($p < 0.05$ for all these cases), significant negative correlations in 29

simulations ($p < 0.05$ for all these cases), and no significant correlations in 5645 simulations

($p \geq 0.05$ for all these cases) (Table S1). The same analyses of $X_2 \sim X_1$ show that there are

significant positive correlations in 12005 simulations ($p < 0.05$ for all these cases), significant

negative correlations in 20 simulations ($p < 0.05$ for all these cases), and no significant

correlations in 3975 simulations ($p \geq 0.05$ for all these cases) (Table S1). By comparing with

the golden standards, we found that PGLS analyses gave correct results for both $X_1 \sim X_2$ and

$X_2 \sim X_1$ in 11474 simulations (71.71%) and incorrect results for both $X_1 \sim X_2$ and $X_2 \sim X_1$ in

2468 simulations (15.43%). In 2058 (12.86%) simulations, only one of the two competing

models, $X_1 \sim X_2$ or $X_2 \sim X_1$, yielded correct results. Therefore, limited by the performance of

PGLS regression analysis, we could at most obtain an accuracy of 84.57% in analyzing the

data of our 16000 simulations by PGLS regression. However, if we arbitrarily select one trait

(X_1 or X_2) as the independent variable with the worst luck, we could get an accuracy of only

71.71%. In the following attempt to find an accurate criterion for the selection of dependent

variables, we hope to perceive more correct cases from the 2058 simulations, where $X_1 \sim X_2$

and $X_2 \sim X_1$ gave conflicting results.

3.5 Looking for an accurate criterion for dependent variable selection

Referring to the golden standards, we evaluated the potential performance of seven

criteria for selecting a better model from $X_1 \sim X_2$ and $X_2 \sim X_1$.

In statistics, the suitability of two competing statistical models is often assessed by

calculating each model's log-likelihood (LLK) value. First, we examined whether a higher (or

lower) LLK could give an accurate prediction of the better model between $X_1 \sim X_2$ and $X_2 \sim X_1$.

By calculating the LLKs of the two models for the 2058 simulations (Table S1), we found that

the models selected from $X_1 \sim X_2$ and $X_2 \sim X_1$ by a lower LLK (denoted as $Model_{LLK,lower}$)

have more correct outcomes than the alternative model (denoted as $Model_{LLK,higher}$), 1076 *vs.*

982. A χ^2 test showed a statistically significant difference ($p = 3.7\times 10^{-3}$, Table 3).

Akaike information criterion (AIC) is a widely used estimator of the quality of

statistical models for a given dataset (Akaike 1974). It balances the excellent fit of the model

with the complexity of the model. Calculating the AIC values of the two competing models

for the 2058 simulations (Table S1), we found that the models selected with a higher AIC

value (denoted as $Model_{AIC,higher}$) had significantly more correct outcomes than the

alternative model (denoted as $Model_{AIC,lower}$), 1076 *vs.* 982, $p = 3.7\times 10^{-3}$ (Table 3).

R^2 describes the proportion of total variation in the dependent variable explained by

independent variables in the regression model, and the *p*-value of the regression coefficient is

the probability of observing the test statistic value under the assumption that the null

hypothesis is true, where the regression coefficient equals 0. These two parameters are widely

used indicators of the suitability of regression models. The models selected by a higher R^2

(denoted as $Model_{R^2,higher}$) and a lower *p*-value (denoted as $Model_{p,lower}$) have

significantly more correct outcomes than the alternative models ($p = 2.7\times 10^{-4}$ for both cases,

Table 3).

The phylogenetic signal measures how much phylogenetic structure influences species trait values. Pagel's λ and Blomberg et al.'s K are the most commonly used indicators of the phylogenetic signal (Pagel 1999; Blomberg *et al.,* 2003). The estimated λ ($\hat{\lambda}$) is a parameter in Pagel's λ model (Pagel 1997) that measures the relatedness of the regression residuals with the phylogenetic structure. We found that the models that use the trait with a higher λ value or a higher K value as the dependent variable (denoted as $Model_{\lambda y,higher}$ and $Model_{K y,higher}$, respectively) have significantly more correct outcomes than the alternate models (denoted as $Model_{\lambda y,lower}$ and $Model_{K y,lower}$, respectively) ($p < 2.2 \times 10^{-16}$ for both case Table 3). And the model chosen by a higher $\hat{\lambda}$ (denoted as $Model_{\hat{\lambda},higher}$) also had significantly more correct results than the alternative model (denoted as $Model_{\hat{\lambda},lower}$) ($p < 2.2 \times 10^{-16}$, Table 3).

In addition, we defined a virtual criterion where one model was randomly selected from the two competing models, $X_1 \sim X_2$ and $X_2 \sim X_1$. The results of the models selected by this virtual criterion ($Model_{rc}$) were compared with the better models selected by the above seven criteria. Although the better models selected by the above seven criteria consistently have more correct results than $Model_{rc}$ (Table 4), none of $Model_{LLK,lower}$, $Model_{AIC,higher}$, $Model_{R^2,higher}$, and $Model_{p,lower}$ is significantly different from $Model_{rc}$ ($p > 0.100$ for all four cases), but the differences of $Model_{\lambda y,higher}$, $Model_{K y,higher}$, and $Model_{\hat{\lambda},higher}$ with $Model_{rc}$ are highly significant ($p < 2.2 \times 10^{-16}$ for all the three cases).

From Tables 3-4 and S2, we can see three distinct groups of criteria: LLK and AIC, R^2 and p-value, and Pagel's λ, Blomberg et al.'s K, and $\hat{\lambda}$. Within each group, the criteria gave the same results. Among the three groups, Pagel's λ, Blomberg et al.'s K, and $\hat{\lambda}$ seem to be the best criterion for dependent variable selection. For a quantitative evaluation of these impressions, we performed the χ^2 tests to compare Pagel's λ with Blomberg et al.'s K, $\hat{\lambda}$, LLK, AIC, R^2, and p-value using the 2058 simulations where $X_1 \sim X_2$ and $X_2 \sim X_1$ gave conflicting results. As shown in Table 5, the equivalency among Pagel's λ, Blomberg et al.'s K, and $\hat{\lambda}$ and the superiority of λ to LLK, AIC, R^2, and p-value have been statistically confirmed.

In summary, Pagel's λ, Blomberg et al.'s K, and $\hat{\lambda}$ are the best criteria for dependent variable selection. Of the 2058 simulations that $X_1 \sim X_2$ and $X_2 \sim X_1$ gave conflicting results, these criteria led to correct results in 1734 simulations. Combined with the 11474 simulations

where both $X_1 \sim X_2$ and $X_2 \sim X_1$ gave correct results, PGLS analysis was able to achieve

82.55% accuracy when the trait with a stronger phylogenetic signal was selected as the

dependent variable.

4 DISCUSSION

In the PGLS correlation analysis of an empirical dataset (Liu *et al.,* 2023), we found

that switching dependent and independent variables could lead to a significant incidence of

conflicting results. To investigate this further, we simulated the evolution of two traits (X_1

and X_2) along a binary tree with 100 terminal nodes, using different models and variances

for 16000 iterations. Our PGLS analyses of these simulated data showed that the frequency of

conflicting results caused by switching dependent and independent variables depends on the

strength of the relationship between the analyzed traits. Furthermore, we showed that

switching dependent and independent variables leads to different parameter estimations in

PGLS regression analysis of two traits that have quite different phylogenetic signals. Using

simulation, we established a golden standard for whether the two traits, X_1 and X_2, are

correlated in each simulation by conventional correlation analysis of the changes of the two

traits along the branches of the phylogenetic tree. With this golden standard, we can tell

which model, $X_1 \sim X_2$ or $X_2 \sim X_1$, is correct. Seven potential criteria for dependent variable

selection, LLK, AIC, R^2, p-value, Pagel's λ, Blomberg et al.'s K, and $\hat{\lambda}$, have been compared.

The last three criteria are equivalent in selecting dependent variables and have shown their

superiority over the other four criteria. Pagel's λ and Blomberg et al.'s K values are generally

calculated at the beginning of the phylogenetic comparative analysis, so they are already

known before the PGLS analysis. If we can choose the trait with a higher λ or K value as the

dependent variable, two rounds of PGLS analysis, like the $X_1 \sim X_2$ and $X_2 \sim X_1$, are not

required. Considering the practical convenience, Pagel's λ and Blomberg et al.'s K are

superior to $\hat{\lambda}$.

As suggested by some previous publications, like Cooper et al. (2016), Paterno et al.

(2018), and Uyeda et al. (2018), PGLS regression is not a perfect method for phylogenetic

correlation analysis. In our 16000 simulations, PGLS regression analyses yielded incorrect

results for 2468 cases, regardless of which trait is designated as the dependent variable. In

2058 simulations, the two competing models, $X_1 \sim X_2$ or $X_2 \sim X_1$, gave conflicting results:

one is right and the other is wrong. That is, PGLS regression analyses of 16000 simulations

with arbitrarily selected dependent variables could yield correct results in the range of

71.79% to 84.57%. By selecting the trait with a strong phylogenetic signal as the dependent

variable, we could obtain an accuracy of 82.55%. As this accuracy is still 2.02% below the

upper limit of PGLS regression using Pagel's λ model, a better criterion may be found in the

future. PGLS regression analysis itself should also be improved.

It is important to emphasize that the terms "independent" and "dependent" variables

can be misleading when using PGLS to study the correlation between traits whose causal

relationships are unclear, and should not be interpreted literally. In such cases, the

independent variable does not necessarily represent the cause of changes in the dependent

variable. PGLS is thus primarily designed to account for phylogenetic non-independence and

effectively replaces conventional correlation methods, such as Pearson and Spearman's rank

tests, in comparative studies. When selecting a dependent variable for PGLS, the focus

should be on obtaining robust and interpretable estimates of the relationship rather than

assigning causal roles. Specifically, even if a trait appears to drive changes in another

biologically, it should be designated as the dependent variable if it exhibits a stronger

phylogenetic signal, as this choice generally produces more consistent results.

This study specifically addresses the issue of dependent variable selection in PGLS

analyses using Pagel's λ model (Freckleton *et al.,* 2002). To examine the theoretical

consistency, we performed PGLS analyses on 16,000 simulated datasets under a Brownian

motion model, alternating X_1 and X_2 as the dependent variable (Table S1). For all simulations,

the results of $X_1 \sim X_2$ and $X_2 \sim X_1$ were consistent in terms of significance (*p* value) and the sign

of the regression coefficient, indicating that arbitrary choice of the dependent variable does

not affect correlation estimates in models without parameter estimation. In contrast, for

models involving parameter estimation, such as the Ornstein–Uhlenbeck model (Hansen

1997) or the early burst model (Harmon *et al.,* 2010), swapping dependent and independent

variables can produce different parameter estimates and potentially conflicting conclusions.

Further research is needed to establish clear criteria for selecting dependent variables in these

models.

Separately, in a more comprehensive empirical study of the correlation between

bacterial growth rate and CRISPR spacer content in genomes (Liu et al., 2023), we observed

that switching dependent and independent variables could change the significance of the

correlation. This observation highlights that in practical applications, especially with real

biological data, careful consideration of variable assignment—favoring traits with stronger

phylogenetic signal—is critical for robust and reproducible inference in PGLS.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (Grant number 31671321).

REFERENCES

Akaike, H. (1974) A new look at the statistical model identification. *IEEE Transactions on Automatic Control,* **19,** 716-723. Https://doi.org/10.1109/TAC.1974.1100705

Bartomeus, I., Cariveau, D.P., Harrison, T. & Winfree, R. (2018) On the inconsistency of pollinator species traits for predicting either response to land-use change or functional contribution. *Oikos,* **127,** 306-315. Https://doi.org/10.1111/oik.04507

Bawa, K.S., Ingty, T., Revell, L.J. & Shivaprakash, K.N. (2019) Correlated evolution of flower size and seed number in flowering plants (monocotyledons). *Annals of Botany,* **123,** 181-190. Https://doi.org/10.1093/aob/mcy154

Blomberg, S.P., Garland, T. & Ives, A.R. (2003) Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. *Evolution,* **57,** 717-745. Https://doi.org/10.1111/j.0014-3820.2003.tb00285.x

Cooper, N., Thomas, G.H. & FitzJohn, R.G. (2016) Shedding light on the 'dark side' of phylogenetic comparative methods. *Methods in Ecology and Evolution,* **7,** 693-699. Https://doi.org/10.1111/2041-210X.12533

Felsenstein, J. (1985) Phylogenies and the comparative method. *The American Naturalist,* **125,** 1-15. Https://doi.org/10.1086/284325

Freckleton, R.P., Harvey, P.H. & Pagel, M. (2002) Phylogenetic analysis and comparative data: A test and review of evidence. *American Naturalist,* **160,** 712-726. Https://doi.org/10.1086/343873

Garamszegi, L.Z. (2014) Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology: Concepts and Practice. Springer, Berlin. Https://doi.org/10.1007/978-3-662-43550-2

Goswami, A., Smaers, J.B., Soligo, C. & Polly, P.D. (2014) The macroevolutionary consequences of phenotypic integration: from development to deep time. *Philosophical Transactions of the Royal Society B: Biological Sciences,* **369,** 20130254. Https://doi.org/10.1098/rstb.2013.0254

Grafen, A. (1989) The phylogenetic regression. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,* **326,** 119-157. Https://doi.org/10.1098/rstb.1989.0106

Hansen, T.F. (1997) Stabilizing selection and the comparative analysis of adaptation. *Evolution,* **51,** 1341-1351. Https://doi.org/10.1111/j.1558-5646.1997.tb01457.x

Harmon, L.J., Losos, J.B., Jonathan Davies, T., Gillespie, R.G., Gittleman, J.L., Bryan Jennings, W., Kozak, K.H., McPeek, M.A., Moreno-Roark, F., Near, T.J., Purvis, A., Ricklefs, R.E., Schluter, D., Schulte Ii, J.A., Seehausen, O., Sidlauskas, B.L., Torres-Carvajal, O., Weir, J.T. & Mooers, A.Ø. (2010) Early bursts of body size

and shape evolution are rare in comparative data. *Evolution, 64,* 2385-2396. Https://doi.org/10.1111/j.1558-5646.2010.01025.x

Ho, L.S.T. & Ane, C. (2014) A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Systematic Biology, 63,* 397-408. Https://doi.org/10.1093/sysbio/syu005

Hu, E.-Z., Lan, X.-R., Liu, Z.-L., Gao, J. & Niu, D.-K. (2022) A positive correlation between GC content and growth temperature in prokaryotes. *BMC Genomics, 23,* 110. Https://doi.org/10.1186/s12864-022-08353-7

Liu, Z.-L., Hu, E.-Z. & Niu, D.-K. (2023) Investigating the relationship between CRISPR-Cas content and growth rate in bacteria. *Microbiology Spectrum, 11,* e03409-03422. Https://doi.org/10.1128/spectrum.03409-22

Lynch, M. (1991) Methods for the analysis of comparative data in evolutionary biology. *Evolution, 45,* 1065-1080. Https://doi.org/10.1111/j.1558-5646.1991.tb04375.x

Martins, E.P. & Hansen, T.F. (1997) Phylogenies and the comparative method: A general approach to incorporating phylogenetic information into the analysis of interspecific data. *American Naturalist, 149,* 646-667. Https://doi.org/10.1086/286013

Pagel, M. (1997) Inferring evolutionary processes from phylogenies. *Zoologica Scripta, 26,* 331-348. Https://doi.org/10.1111/j.1463-6409.1997.tb00423.x

Pagel, M. (1999) Inferring the historical patterns of biological evolution. *Nature, 401,* 877-884. Https://doi.org/10.1038/44766

Paradis, E. & Schliep, K. (2019) ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics, 35,* 526-528. Https://doi.org/10.1093/bioinformatics/bty633

Parks, D.H., Chuvochina, M., Rinke, C., Mussig, A.J., Chaumeil, P.-A. & Hugenholtz, P. (2022) GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Research, 50,* D785-D794. Https://doi.org/10.1093/nar/gkab776

Paterno, G.B., Penone, C. & Werner, G.D.A. (2018) sensiPhy: An r-package for sensitivity analysis in phylogenetic comparative methods. *Methods in Ecology and Evolution, 9,* 1461-1467. Https://doi.org/https://doi.org/10.1111/2041-210X.12990

Pearman, P.B., Lavergne, S., Roquet, C., Wüest, R., Zimmermann, N.E. & Thuiller, W. (2014) Phylogenetic patterns of climatic, habitat and trophic niches in a European avian assemblage. *Global Ecology and Biogeography, 23,* 414-424. Https://doi.org/10.1111/geb.12127

Revell, L.J. (2010) Phylogenetic signal and linear regression on species data. *Methods in Ecology and Evolution, 1,* 319-329. Https://doi.org/10.1111/j.2041-210X.2010.00044.x

Revell, L.J. (2012) phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution,* **3,** 217-223. Https://doi.org/10.1111/j.2041-210X.2011.00169.x

Revell, L.J. & Collar, D.C. (2009) Phylogenetic analysis of the evolutionary correlation using likelihood. *Evolution,* **63,** 1090-1100. Https://doi.org/10.1111/j.1558-5646.2009.00616.x

Revell, L.J., Toyama, K.S. & Mahler, D.L. (2022) A simple hierarchical model for heterogeneity in the evolutionary correlation on a phylogenetic tree. *PeerJ,* **10,** e13910. Https://doi.org/10.7717/peerj.13910

Rohlf, F.J. (2001) Comparative methods for the analysis of continuous variables: Geometric interpretations. *Evolution,* **55,** 2143-2160. Https://doi.org/10.1111/j.0014-3820.2001.tb00731.x

Suarez-Castro, A.F., Mayfield, M.M., Mitchell, M.G.E., Cattarino, L., Maron, M. & Rhodes, J.R. (2020) Correlations and variance among species traits explain contrasting impacts of fragmentation and habitat loss on functional diversity. *Landscape Ecology,* **35,** 2239-2253. Https://doi.org/10.1007/s10980-020-01098-2

Travnicek, P., Certner, M., Ponert, J., Chumova, Z., Jersakova, J. & Suda, J. (2019) Diversity in genome size and GC content shows adaptive potential in orchids and is closely linked to partial endoreplication, plant life-history traits and climatic conditions. *New Phytologist,* **224,** 1642-1656. Https://doi.org/10.1111/nph.15996

Uyeda, J.C., Zenil-Ferguson, R. & Pennell, M.W. (2018) Rethinking phylogenetic comparative methods. *Systematic Biology,* **67,** 1091-1109. Https://doi.org/10.1093/sysbio/syy031

Waugh, F.V. (1943) Choice of the dependent variable in regression analysis. *Journal of the American Statistical Association,* **38,** 210-214. Https://doi.org/10.1080/01621459.1943.10501796

Whitney, K.D. & Garland, T., Jr. (2010) Did genetic drift drive increases in genome complexity? *PLoS Genetics,* **6,** e1001080. Https://doi.org/10.1371/journal.pgen.1001080

DATA AVAILABILITY STATEMENT

Supplementary Tables S1-S2 are submitted as Supporting Information for Online Publication.

FUNDING

This study was supported in part by the National Natural Science Foundation of China (grant number 31671321), with leftover resources used for occasional research expenses.

AUTHOR CONTRIBUTIONS

Deng-Ke Niu and Zheng-Lin Chen conceived the idea and designed the general methodology; Zheng-Lin Chen designed the simulations; Zheng-Lin Chen and Hong-Ji Guo

499 implemented the simulations and data analysis; Deng-Ke Niu led the writing of the

500 manuscript. All authors contributed critically to the drafts and gave final approval for

501 publication.

503 Table 1. Switching dependent and independent variables in PGLS may yield conflicting

504 results.[†]

Dependent Variable ~ Independent Variable	Slope	p	R^2
EMDT ~ min_CRISPR_spacer	0.010	0.049	0.015
min_CRISPR_spacer ~ EMDT	1.376	0.053	0.014
EMDT ~ mean_prophage_number	0.565	0.013	0.024
mean_prophage_number ~ EMDT	0.008	0.558	0.001
EMDT ~ min_prophage_number	0.622	0.007	0.028
min_prophage_number ~ EMDT	0.011	0.427	0.002
EMDT ~ max_prophage_number	0.448	0.037	0.017
max_prophage_number ~ EMDT	0.005	0.722	5×10^{-4}
OPT ~ min_CRISPR_array	0.308	0.061	0.014
min_CRISPR_array ~ OPT	0.053	8×10^{-5}	0.058
OPT ~ mean_prophage_number	−0.197	0.242	0.005
mean_prophage_number ~ OPT	−0.031	9×10^{-4}	0.042
OPT ~ min_prophage_number	−0.154	0.374	0.003
min_prophage_number ~ OPT	−0.026	0.004	0.031
OPT ~ max_prophage_number	−0.183	0.214	0.006
max_prophage_number ~ OPT	−0.036	4×10^{-4}	0.047
OPT ~ prophage_existence	−1.046	0.182	0.007
prophage_existence ~ OPT	−0.009	8×10^{-4}	0.043
OPT ~ Average_Cas_gene_number	0.197	0.610	0.002
Average_Cas_gene_number ~ OPT	0.023	0.012	0.043

[†]Pagel's λ model was used in these analyses. The dataset containing 262 bacteria was

extracted from the Supplemental Material Table S1 of Liu et al. (2023), and the phylogenetic

tree used in PGLS analysis was retrieved from the Genome Taxonomy Database (Parks *et al.,*

2022). EMDT, empirical_minimal_doubling_times; OPT,

mean_optimal_growth_temperature. The variables before and after "~" are dependent and

independent.

 Table 2. Frequency of conflicting results caused by switching dependent and independent variables in PGLS analyses using Pagel's λ model

	BM								Norm							
X_1[†]																
X_2[‡]	BM + Norm								Norm + BM							
Noise Term Variance	10^{-4}	0.01	1	4	16	64	256	1024	10^{-4}	0.01	1	4	16	64	256	1024
Simulation times	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3	10^3
$X_1 \sim X_2$ gave $p < 0.05$[§]	10^3	10^3	10^3	982	543	188	68	53	10^3	10^3	10^3	10^3	926	387	134	74
$X_2 \sim X_1$ gave $p < 0.05$	10^3	10^3	10^3	10^3	954	528	171	81	10^3	10^3	10^3	10^3	10^3	844	318	129
Both $X_1 \sim X_2$ and $X_2 \sim X_1$ gave $p < 0.05$	10^3	10^3	10^3	982	534	169	34	14	10^3	10^3	10^3	10^3	926	375	88	39
Both $X_1 \sim X_2$ and $X_2 \sim X_1$ gave $p \geq 0.05$	0	0	0	0	37	453	795	880	0	0	0	0	0	144	636	836
$X_1 \sim X_2$ and $X_2 \sim X_1$ gave conflicting results[¶]	0	0	0	18	429	378	171	106	0	0	0	0	74	481	276	125

[†]The trait X_1 evolved according to a Brownian motion (BM) model in the first 8000 simulations, and evolved according to a normal distribution in the other

8000 simulations. For details, see the MATERIALS AND METHODS section.

[‡]The trait X_2 was simulated by adding a noise term to X_1. The term noise was normally distributed in the first 8000 simulations and generated according to a

Brownian motion model in the other 8000 simulations. For details, see the MATERIALS AND METHODS section.

[§]The variables before and after "~" are dependent and independent, respectively.

[¶]When one of $X_1 \sim X_2$ and $X_2 \sim X_1$ gives a $p < 0.05$ result and the other gives a $p \geq 0.05$ result, we define these results as conflicting.

Table 3. Differences in the performance of models selected by seven criteria.

Criteria	Models	Correct	Incorrect	χ^2	$p^{\dagger}$
LLK	$Model_{LLK,higher}$	982	1076	8.405	3.7×10^{-3}
	$Model_{LLK,lower}$	1076	982		
AIC	$Model_{AIC,higher}$	1076	982	8.405	3.7×10^{-3}
	$Model_{AIC,lower}$	982	1076		
R^2	$Model_{R^2,higher}$	1088	970	13.30	2.7×10^{-4}
	$Model_{R^2,lower}$	970	1088		
p	$Model_{p,higher}$	970	1088	13.30	2.7×10^{-4}
	$Model_{p,lower}$	1088	970		
λ_y	$Model_{\lambda_y,higher}$	1734	324	1929	$< 2.2 \times 10^{-16}$
	$Model_{\lambda_y,lower}$	324	1734		
K_y	$Model_{K_y,higher}$	1734	324	1929	$< 2.2 \times 10^{-16}$
	$Model_{K_y,lower}$	324	1734		
$\hat{\lambda}$	$Model_{\hat{\lambda},higher}$	1734	324	1929	$< 2.2 \times 10^{-16}$
	$Model_{\hat{\lambda},lower}$	324	1734		

519 †The minimal significance value that the R environment (version 4.1.0) gives is 2.2×10^{-16}.

520 Llk, log-likelihood; AIC, Akaike information criterion. When the χ^2 tests were extended to all

521 16000 simulations, λ_y, K_y, and $\hat{\lambda}$ are also the best criteria (Table S2).

523 Table 4. Comparing the performance of seven criteria with random selection.

Criteria	Models	Correct	Incorrect	χ^2	$p^{\dagger}$
$\hat{\lambda}$	$Model_{\hat{\lambda},higher}$	1734	324	532.3	$< 2.2 \times 10^{-16}$
	$Model_{rc}$	1039	1019		
K_y	$Model_{K_y,higher}$	1734	324	532.3	$< 2.2 \times 10^{-16}$
	$Model_{rc}$	1039	1019		
λ_y	$Model_{\lambda_y,higher}$	1734	324	532.3	$< 2.2 \times 10^{-16}$
	$Model_{rc}$	1039	1019		
LLK	$Model_{LLK,lower}$	1076	982	1.260	0.262
	$Model_{rc}$	1039	1019		
AIC	$Model_{AIC,higher}$	1076	982	1.260	0.262
	$Model_{rc}$	1039	1019		
R^2	$Model_{R^2,higher}$	1088	970	2.242	0.134
	$Model_{rc}$	1039	1019		
p	$Model_{p,lower}$	1088	970	2.242	0.134
	$Model_{rc}$	1039	1019		

524 †The minimal significance value that the R environment (version 4.1.0) gives is 2.2×10^{-16}.

525 LLK, log-likelihood; AIC, Akaike information criterion.

527 Table 5. Comparing the performance of λ_y with other criteria for dependent variable

528 selection

Criteria	Models	Correct	Incorrect	χ^2	$p^\dagger$
λ_y vs. $\hat{\lambda}$	$Model_{\lambda_y,higher}$	1734	324	0	1
	$Model_{\hat{\lambda},higher}$	1734	324		
λ_y vs. K_y	$Model_{\lambda_y,higher}$	1734	324	0	1
	$Model_{K_y,higher}$	1734	324		
λ_y vs. LLK	$Model_{\lambda_y,higher}$	1734	324	484.1	$< 2.2 \times 10^{-16}$
	$Model_{LLK,lower}$	1076	982		
λ_y vs. AIC	$Model_{\lambda_y,higher}$	1734	324	484.1	$< 2.2 \times 10^{-16}$
	$Model_{AIC,higher}$	1076	982		
λ_y vs. R^2	$Model_{\lambda_y,higher}$	1734	324	468.9	$< 2.2 \times 10^{-16}$
	$Model_{R^2,higher}$	1088	970		
λ_y vs. p	$Model_{\lambda_y,higher}$	1734	324	468.9	$< 2.2 \times 10^{-16}$
	$Model_{p,lower}$	1088	970		

529 †The minimal significance value that the R environment (version 4.1.0) gives is 2.2×10^{-16}.

530 LLK, log-likelihood; AIC, Akaike information criterion.

Figure 1 Phylogenetic signals of the variables and the estimated λ of $X_1 \sim X_2$ and $X_2 \sim X_1$

models under different noise term variances. For each variance gradient, 1000 simulations

were conducted. The data points on the graph represent the average value of phylogenetic

diagnostics (phylogenetic signals or model estimated λ) of the noise term under a specific

variance gradient based on the 1000 simulations. (A) In the "BM & BM + Norm" case, X_1

was simulated according to a Brownian motion (BM) model, while X_2 was simulated based

on X_1 plus a normally distributed random noise term. (B) In the "Norm & Norm + BM"

case, X_1 was simulated according to a normal distribution, and X_2 was simulated according

to X_1 plus a noise term, where the noise term followed a BM model. $X_1 \sim X_2$ and $X_2 \sim X_1$

represent PGLS regressions using X_1 and X_2 as dependent variables, respectively.