Decryption of sequence, structure, and functional features of SINE repeat elements in SINEUP non-
coding RNA–mediated post-transcriptional gene regulation

Harshita Sharma[1], Matthew N Z Valentine[1], Naoko Toki[1,], Hiromi Sueki Nishiyori[1], Stefano Gustincich[2],
Hazuki Takahashi[1]* and Piero Carninci[1,3]*

1 Laboratory for Transcriptome Technology, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa,
230-0045, Japan
2 Department of Neuroscience and Brain Technologies, Istituto Italiano di Tecnologia, Genova, Italy
3 Human Technopole, Milan, 20157, Italy
* Corresponding authors
E-mails: carninci@riken.jp; piero.carninci@fht.org; hazuki.takahashi@riken.jp

Abstract

RNA structure folding largely influences RNA regulation by providing flexibility and functional diversity. In
silico and in vitro analyses are limited in their ability to capture the intricate relationships between
dynamic RNA structure and RNA functional diversity present in the cell. Here, we investigate sequence,
structure and functional features of mouse and human SINE-transcribed retrotransposons embedded in
SINEUPs long non-coding RNAs, which positively regulate target gene expression post-transcriptionally.
In-cell secondary structure probing reveals that functional SINEs-derived RNAs contain conserved short
structure motifs essential for SINEUP-induced translation enhancement. We show that SINE RNA structure
dynamically changes between the nucleus and cytoplasm and is associated with compartment-specific
binding to RBP and related functions. Moreover, RNA–RNA interaction analysis shows that the SINE-
derived RNAs interact directly with ribosomal RNAs, suggesting a mechanism of translation regulation.
We further predict the architecture of 18 SINE RNAs in three dimensions guided by experimental
secondary structure data. Overall, we demonstrate that the conservation of short key features involved
in interactions with RBPs and ribosomal RNA drives the convergent function of evolutionarily distant SINE-
transcribed RNAs.

Introduction

In long non-coding RNAs (lncRNAs), evolutionary selection takes place not only at the level of primary

sequence and expression control from syntenic loci, but also at the level of structure and function[1]. This

leads to conservation of function despite divergent sequence and secondary structure among lncRNA

orthologs[1] and highlights the need to look beyond the sequence to understand the relationship between

sequence, structure, and function in lncRNAs. The presence of modular secondary structures in several

functional lncRNAs[2-4] also supports the notion that lncRNA evolution favors the conservation of

functionally relevant short stretches of sequence and/or structure that form distinct interacting domains

for RNA–DNA, RNA–RNA, and RNA–protein regulatory complexes[5,6]. The combination of these different

domains provides the lncRNAs with their unique functional specificity and diversity[5]. Such "modular RNA

code" was validated in recent studies where short structure motifs were found to control the lncRNA

stability and function, such as 3' end triple helix in MALAT1 and NEAT1[7,8], 5' asymmetric G-rich internal

loop (AGIL) motif in mouse Braveheart (Bvht)[4], pseudoknots in human MEG3[9], and a terminal hairpin

stem-loop (SL1) in mouse lncRNA antisense to *Uchl1* mRNA[10-12]. In the antisense to *Uchl1*, the crucial SL1

region is within an embedded short interspersed nuclear element (SINE) from the B2 family, which forms

the effector domain (ED) that is essential for targeted mRNA translation upregulation[10,11].

SINEs are on average 100–300 bp in length and are non-autonomous retrotransposons that are

evolutionarily derived from RNA polymerase III transcripts, mainly 5S ribosomal RNA (rRNA), 7SL RNA, and

transfer RNA (tRNA)[13,14]. These elements, previously considered "junk DNA", are abundant in gene-rich

regions and their role in gene regulatory networks has been established[15-18]. Many SINEs are transcribed

as enhancer RNAs and form the regulatory domains of functional lncRNAs[19-22] as well as participating in

the nuclear retention of lncRNAs[23], STAU1-mediated mRNA decay[24], and promotion of sense mRNA

translation by antisense lncRNA[10], thus supporting the Repeat Insertion Domains of LncRNAs (RIDL)

hypothesis[6]. Yet the majority of SINE regulatory functions and structural features remain poorly

understood. In mouse genome, the B2 family of SINE constitutes approximately 2.4% of the genome with

presence of ~3.5 x 10^5 copies[25]. The SINEB2 family is broadly classified into B3A, B3, B2_Mm2, B2_mm1t,

and B2_mm1a sub-families in descending order of evolutionary age according to the Repbase repeat

elements database[26-28]. Synthetic antisense long non-coding RNAs SINEUPs which comprise a SINEB2

retrotransposon positively regulate the translation of the sense mRNA that they overlap[29,30]. Not only

several mouse SINEB2s but also human free right Alu monomer (FRAM) repeat element and human MIRb

are found in natural antisense lncRNAs and display post-transcriptional SINEUP-induced gene

regulation[10,31], despite their lack of similarity in terms of repeat length, sequence composition, and

predicted secondary structure.

Here, to comprehend more clearly the extent of the SINEUP family, we used a synthetic SINEUP system

to functionally screen 15 SINEB2 elements originally embedded in mouse natural antisense transcripts.

The proven importance of SINE repeats in functional lncRNA domains prompted us to study the primary

sequence and secondary structure of these SINE-derived RNAs. Although we have previously determined

the folding of antisense *Uchl1* SINEB2 in vitro from its chemical footprinting and NMR data[11,12], structural

studies in solution cannot fully grasp the structural dynamics and interaction with cellular machinery.

Moreover, to describe the relationship between sequence, structure, and function in SINE-derived RNAs,

we need to study the RNA structure of multiple SINE-derived RNAs in the cellular environment and its

specific compartments, because RNA can acquire different conformations depending on subcellular

location, functional state, and biomolecular interactions[32] . Additionally, we need to understand RNA

modifications in functional SINE-derived RNAs, specifically pseudouridylation, which is reported to be

crucial for the structural stability, flexibility, and function of RNAs involved in translation[33-35].

LncRNA structures are known to act as a molecular scaffold for proteins[36,37], and in fact, the binding of

heterogeneous nuclear ribonucleoprotein K (HNRNPK) with SINEUP and target mRNA complex is vital for

SINEUP-mediated upregulation of target mRNA translation[38]. Previously, we found that upon HNRNPK

knock-down, the co-localization of endogenous *Uchl1* mRNA and SINEUP RNA in the cytoplasm was

decreased which resulted in loss of translation upregulation by SINEUP-UCHL1[38]. This negative effect of

low HNRNPK expression on SINEUP function was also observed in SINEUP-GFP[38]. Furthermore, RNA

immunoprecipitation analysis showed that HNRNPK binds with SINEUP-EGFP mRNA complex in both

nucleus and cytoplasm and co-sediments with SINEUP and EGFP mRNA in free 40S, monosome and light

polysome fractions in polysome profiling[38]. In other words, HNRNPK not only aids in the export of SINEUP

and target mRNA complexes to the cytoplasm but also participates in SINEUP RNA-translation complex[38],

thus, exhibiting two discrete functions in the nucleus and the cytoplasm. This knowledge makes it

imperative to examine how specific SINE RNA structures in different cellular compartments interact with

RNA binding proteins (RBPs) and support their distinct functions. In addition to intramolecular structure,

intermolecular interactions with other RNAs play a critical role in gene regulation by non-coding RNA[40,41],

notably the binding to rRNAs by linear and circular internal ribosome entry sites (IRESs) for promotion of

translation[42,43]. In fact, our previous molecular dynamics simulation study on NMR solution structure of

SINEB2 SL1 identified motifs in three-dimensions (3D) similar to those in rRNAs[44]. This data and the

association of SINEUPs with ribosomal subunits in polysome profiles[38] made us ponder the possibility that

direct interaction between SINE-derived RNAs and rRNAs may be a mechanism by which SINEUPs enhance

mRNA translation.

To solve the puzzle of SINE-transcribed RNA structure, we utilized a method called in vivo click selective

2'-hydroxyl acylation and profiling experiment (icSHAPE), which chemically probes flexible RNA bases

inside living cells and thus provides an advantage over other chemical or enzymatic methods[45]. We

present two-dimensional (2D) secondary structure models for 17 mouse SINEB2s-derived RNAs and a

human SINE FRAM RNA, providing for the first time experimentally verified in-cell 2D-structure models of

evolutionarily close and distant SINE-transcribed RNAs at this scale. We found that structural deletion

leading to the gain of an apical stem-loop (known as SL1) restores the activity in a SINEUP that is

functionally dormant (i.e., lacking the ability to significantly enhance the target mRNA translation). Based

on the 2D models, we discovered seven short conserved-structure motifs among the SINE RNAs tested,

three of which harbor modified RNA bases. The icSHAPE-guided structures for the nucleus and cytoplasm

showed that SINE RNAs in both mouse (SINEB2, derived from tRNA) and human (FRAM, derived from 7SL

RNA) rearrange into a cloverleaf-like structure in the cytoplasm and have partially conserved structure

motifs. We also observed that compartment-specific structures of SINE RNA are associated with

distinctive binding patterns of HNRNPK in the nucleus and the cytoplasm by using a single-end enhanced

crosslinking and immunoprecipitation assay (seCLIP)[46]. Additionally, when motifs were compared in silico

with those in the Protein Data Bank (PDB), we detected a remarkable similarity of SINEUP 2D structure

motifs with rRNA structure motifs. We also checked RNA–RNA pairs by the psoralen analysis of RNA

interactions and structures (PARIS2) method[41] and identified interactions of mouse and human SINE-

derived RNAs with human 18S and 28S rRNAs in the conserved structure motif regions, which shed some

light on SINEUP's mechanism of action. Finally, we used our experimental 2D structure data to guide the

prediction of structure models for 18 SINE-transcribed RNAs in 3D which can provide a base for future

experimental 3D structure studies. In brief, this study demonstrates that despite sequence variability

among SINE-transcribed RNAs, short stretches of conserved sequence and structure preserve the

regulatory functions by providing binding sites for RBPs and rRNA. These short motifs are likely to be

stabilized by modifications and non-canonical base interactions and are evolutionarily favored to maintain

the SINEUP functionality even in distantly related SINE-derived RNAs.

Results

The SINEUP effect is a prevalent functional feature of mouse SINEB2s-derived RNAs

To experimentally verify the translation-regulatory function of various mouse SINE-transcribed RNAs in

the SINEUP effector domain (ED), we used a previously established synthetic SINEUP-GFP model that

enhances EGFP mRNA translation (**Fig. 1a**). We swapped the SINEB2 from antisense *Uchl1* in SINEUP-GFP

with those from other mouse antisense to genes: *Txnip*, *Gadd45α*, and *Uxt* (**Supplementary Table 1a**).

RepeatMasker revealed the presence of an inverted SINEB2 repeat spanning from 1389 to 1495

nucleotide positions (107 bp long) in antisense *Txnip*, and both antisense *Gadd45α* and antisense *Uxt*

lncRNAs contain two inverted SINEB2 repeats in the region not complementary to the sense mRNA. Both

antisense *Uxt* SINEB2 sequences are from the B3 sub-family; in the FANTOM3 cDNA clone, one sequence

covered positions 160 to 290 (131 bp) and the other 774 to 960 (187 bp) (**Supplementary Table 1a**). Based

on their natural order of occurrence from the 5' end of the transcript, the sequences were named

antisense *Uxt* SINEB2-a (131 bp, proximal to the 5' end) and antisense *Uxt* SINEB2-b (187 bp, distal to the

5' end) (**Supplementary Table 1a**, **Fig. 1b**). In contrast, both SINEB2 repeats of antisense *Gadd45α* belong

to the B2_Mm2 sub-family (according to RepeatMasker nomenclature). We named these two sequences

antisense *Gadd45α* SINEB2-a (131 bp, proximal to the 5' end) and antisense *Gadd45α* SINEB2-b (114 bp,

distal to the 5' end) (**Supplementary Table 1a**). To understand whether both SINEB2 sequences are

essential for SINEUP function and how their order affects SINEUP activity, we designed antisense *Gadd45α*

and antisense *Uxt* constructs with a or b sequences and also with both sequences in the natural (ab) or

the inverted (ba) order (**Fig. 1b**). The shortest SINEB2 of those tested, antisense *Txnip* SINEB2 (107 bp),

enhanced the GFP protein level (**Fig. 1c** and **d**, **Supplementary Table 1**). None of the four constructs with

antisense *Gadd45α* SINEB2s could exert the SINEUP effect, regardless of the order of SINEB2 a and b (**Fig.**

1c and **d**). Indeed, natural antisense *Gadd45α* was found to inhibit translation of sense *Gadd45α*

(**Extended Data Fig. 1**), which implies that the SINEUP activity of antisense *Gadd45α* SINEB2 RNA is

dormant, possibly due to the lack of functional regions. Interestingly, of the antisense *Uxt* SINEB2 repeats,

only SINEB2-b upregulated the GFP protein level significantly, and when the position of the two SINEB2

sequences was switched from the natural order (ab switched to ba), their SINEUP effect was obliterated

(**Fig. 1c** and **d**). As negative controls, in addition to empty vector, we tested direct antisense *Uchl1* SINEB2,

where the orientation of SINEB2 was flipped from the one in the functionally active construct, and the

deletion mutant of antisense *Uchl1* SINEB2 (ΔAS *Uchl1* SINEB2) in which the deletion of SINEB2 shifts

downstream synthetic and non-SINEB2 annotated sequences in its place in the ED of SINEUP[10]. Consistent

with our previous work[10], none of these negative controls could upregulate GFP translation which

demonstrates that SINE RNA region is essential for SINEUP function (Fig. **1c** and **d**). Upholding the post-

transcriptional regulatory nature of SINEUPs, none of the SINEUPs changed the EGFP mRNA expression

level significantly (**Extended Data Fig. 2a**).

To further expand analysis of the SINEB2-derived RNAs present in mouse natural antisense transcripts

(**Supplementary Table 1**), we used the shortest functional model of SINEUP-GFP, named miniSINEUP-

GFP[29]. SINEB2 sequences from antisense *Uchl1* and antisense *Uxt* enhanced GFP protein expression by

around 1.8–2.2-fold (**Fig. 1e**, Lanes 2 and 3 in the left graph). We additionally found that combining the

SINEB2 from antisense *Uchl1* with antisense *Uxt* SINEB2-b upregulated GFP protein production; however,

the protein fold change was lower than the effect of either SINEB2 independently (**Fig. 1e**, Lane 4 in the

left graph). We additionally tested three repeats of antisense *Uchl1* SINEB2 but the SINEUP function

remained similar to that with only one repeat, implying that these repeats do not function in a synergistic

or additive manner (**Fig. 1e**, Lane 5 in the left graph).

Since the SINEB2 sub-family B3 acted as a functionally effective SINEUP ED, we decided to check the

SINEUP potential of consensus subfamily B3 sequence from the Repbase database[27,47]. Indeed, the

sequence is functionally active against GFP (~1.6-fold change) (**Fig. 1e**, Lane 6 in the left graph), but not

to the level of antisense *Uchl1* SINEB2 RNA, which is a similar result to tests on other SINEB2 sequences

in the B3 sub-family from antisense lncRNAs to genes: *Nars2*, *Abhd11*, *Epb4.9*, *Wfdc5*, *Pgbd1*, and *Gsk3b*

(**Fig. 1e**, Lanes 7–10 in the left graph and Lanes 3–4 in the right graph).

The failure of SINEUPs with antisense *Gadd45α* SINEB2 from the B2_Mm2 sub-family prompted us to test

whether the SINEUP effect is limited to the B3 sub-family only. To confirm this, we investigated two

B2_Mm2 sub-family sequences (from antisense *Rhod* and antisense *E4f1*), one B2_Mm1t (from antisense

E4f1) and one B3A (from antisense *Rhod*) (**Fig. 1e**, Lanes 5–8 in the right graph) and found that all the

SINEB2 sequences, regardless of their sub-family type, length, or parent natural antisense transcripts,

upregulated GFP protein expression to more or less the same extent, which reveals the vastness of the

SINEUP family. Similar to that of long SINEUPs, this miniSINEUP effect was also manifested at the post-

transcription level (**Extended Data Fig. 2b**). Although the SINEUP RNA level for SINEB2s from antisense

Wfdc5, *Pgbd1*, and *Gsk3b* was unusually high (3–10-fold higher than other miniSINEUPs), the GFP mRNA

expression level in all miniSINEUPs remained unaltered.

Functional SINEs form double-stranded structures in the cell and consist of conserved short structure

motifs

To check any common sequence features between different SINEB2-transcribed RNAs showing the

SINEUP effect and to identify any functionally important regions, we performed multiple sequence

alignment using ClustalW (**Extended Data Fig. 3a** and **b**). This revealed conserved regions of RNA pol III

internal promoters known as A box and B box[48] (**Extended Data Fig. 3a**) along with sequence diversity

between SINEB2 elements tested in this study. Moreover, in the phylogenetic tree based on the multiple

sequence alignment, the functionally active SINEB2s that significantly upregulated GFP translation were

not clustered separately from the inactive SINEB2s that failed to do so (**Extended Data Fig. 3b**). In other

words, on the basis of sequence, the functional SINEB2 RNAs do not possess any unique features that

confer a functional advantage over the inactive SINEB2 RNAs. This lack of conserved sequence domains

led us to examine the in-cell secondary structure of SINEB2s by using the icSHAPE method to map the RNA

secondary structure inside living cells[45,49].

We compared the icSHAPE antisense *Uchl1* SINEB2 RNA secondary structure with our previous in vitro

chemically probed structure[11], which had identified four internal-loops (IL1–IL4) and three stem-loops

(SL1–SL3) (**Fig. 2a**). Both methods gave a notably similar double-stranded structure for antisense *Uchl1*

SINEB2 RNA (**Fig. 2a**). The icSHAPE-based structure comprises three internal loops similar to the IL2, IL3,

and IL4 found in the chemical footprinting-based structure, with slight differences in the sequence

composition. The in-cell structure lacks SL3, which indicates the influence of intracellular factors and

dynamic structural changes compared to the less complex conditions in vitro. Interestingly, SL1 and SL2

are conserved between the structures derived by these two different methods, which suggests that they

represent the most dominant and stable structure components in the ensemble. The structural

conservation is also understandable knowing that SL1 is crucial for SINEUP function[11]. In a similar way,

secondary structure modelling was performed for other SINEB2s-derived RNAs (**Fig. 2b** and **c**, **Extended**

Data Fig. 4) and revealed that they all retained double-stranded features with a terminal GC-rich stem-

loop similar to SL1 of antisense *Uchl1* SINEB2, even though all SINEB2 RNAs tested here had diverse global

structures with variable sequence length and composition.

The importance of short structure motifs like SL1 in the SINEUP function prompted us to look for other

local motifs that are shared between different SINEB2-transcribed RNAs. For this purpose, we utilized the

ExpaRNA tool[50] and, using the antisense *Uchl1* SINEB2 RNA structure as a base, searched for its exact

sequence and structure patterns in other icSHAPE-derived SINEB2 RNA structures. From this search, we

discovered seven GC-rich structure motifs partially conserved among the SINEB2 RNA structures tested in

this study (**Fig. 2a–c**, **Extended Data Fig. 4**). Motifs 2 and 3 are comprised of 'AGG' trinucleotide(s), while

motifs 4, 5, 6, and 7 are 'UGG' enriched. Motif 6 lies within the SL1 region making it a potentially functional

motif. Furthermore, we wondered if the motifs are conserved in SINE-transcribed RNAs that are

evolutionarily distant from mouse SINEB2 elements, such as human SINE FRAM, which has been proven

to be effective as a SINEUP against both endogenous and exogenous mRNA targets[31]. Indeed, the

structure motifs in the whole-cell icSHAPE structure of FRAM RNA provided five matching regions (**Fig.**

2d). This hints that conservation of such short motifs may contribute in functional preservation. We

further examined the PDB database for 3D RNA structure motifs that were akin to our SINEUP structure

motifs. For this purpose, we employed a sequence- and structure-specific search for each of the seven

SINEUP motifs in the RNA Characterization of Secondary Structure Motifs (CoSSMos) database[51]. This

resulted in hits that matched 16S rRNA (PDB ID: 5JB3) and 18S rRNA (PDB ID: 5A2Q) to 2D structures for

SINEUP motifs 2 and 6, respectively (**Extended Data Fig. 5a**). Interestingly, the matching motifs are very

similar to in-cell derived SINEUP motifs in terms of sequence and structure. On closer inspection, we

observed that the matching region to 18S rRNA is within helix 34, which forms the head of the mRNA

entry channel[52] (**Extended Data Fig. 5b**).

Acquisition of SL1-like structure turns a dormant SINEUP active

Considering the conservation of short local structure motifs in functional SINE-transcribed RNAs, we

theorized that dormant SINEUPs, here the SINEB2s that failed to significantly upregulate GFP protein

expression in **Fig. 1c**, might include embedded inhibitory regions that prevent their proper folding into

functional domains. To test this, we chose a functionally inactive SINEB2 sequence, antisense *GADD45α*

SINEB2-a, and checked the effect of deletion of different sequence regions on its structure. In total, we

created four deletion mutants at positions 27–36, 40–50, 66–76, and 92–101 from the 5' end of SINEB2

including two tRNA-derived conserved internal RNA polymerase III promoter regions, A-box and B-box

(**Extended Data Fig. 3a**). We observed structural changes caused by these deletions by in-silico RNA

structure prediction which we later confirmed in the cell by icSHAPE. Analysis of in-cell structures of

deletion mutants in comparison to that of wild-type antisense *GADD45α* SINEB2-a and antisense *Uchl1*

SINEB2 (**Fig. 2a**, **e**, **Extended Data Fig. 6**) showed that out of all the mutants, Δ27–36 shares the most

sequence and structural similarity in SL1 structure and motif 6 with functional antisense *Uchl1* SINEB2

(**Fig. 2a**, **e**). The wild-type antisense *Gadd45α* SINEB2-a carries a deformed SL1 where the stem is

interrupted by multiple internal loops (**Fig. 2e**). Upon deletion of 27–36 nucleotide positions, this stem

becomes less structured and transforms into a properly folded SL1 similar to antisense *Uchl1* SINEB2. The

Δ40–50 mutant, though shares similarity in overall 2D structure in terms of position of motifs 1, 2, 4, 5,

and 7 with Δ27–36 mutant, lacks a proper SL1 and no motif 6 (**Extended Data Fig. 6**. On the other hand,

Δ66–76 and Δ92–101 mutants contain partial motif 6 but have truncated SL1 (**Extended Data Fig. 6**). This

disparity in structure is reflected in their SINEUP function. When we co-transfected the mutants with

sense EGFP into HEK293T/17 cells, as done in the previous SINEUP-GFP functional analysis experiment we

observed that out of the five mutants Δ27–36 could significantly upregulate GFP protein level compared

to full-length antisense *GADD45α* SINEB2-a (**Fig. 2f**). Deletion mutant Δ40–50 and Δ66–76 could not

upregulate GFP translation while Δ92–101 mutant in some experiments showed around 1.5-fold increase

in GFP protein but had inconsistent performance in others (**Fig. 2f**). This suggested that some structural

features in the wild-type antisense *GADD45α* SINEB2-a are detrimental to function, whereas the

acquisition of short local structure motifs leads to functionally favorable RNA folding, which can restore

SINEUP function. Especially, properly folded SL1 and motif 6 are essential not only for exerting the SINEUP

effect but also for ensuring functional stability.

SINEUP structure dynamics across nucleus and cytoplasm

Our SINEUP icSHAPE data from whole cell lysate showed a set of common motifs among different SINEB2s,

yet it was an average of whole-cell structure ensemble and did not help to infer structure dynamics within

the cell. RNA structure is closely related to cellular processes and can acquire distinct conformations

depending on the cellular compartment[32]. Understanding the different RNA structures becomes more

crucial in the case of SINEUPs because they are involved in post-transcriptional translation regulation, a

function specific to cytoplasm. Therefore, it is logical to examine the structure dynamics of functional

SINEB2s-derived RNAs both in the nucleus and the cytoplasm, to better understand the contribution of

structural conformations to the biological processes specific to these subcellular regions. To achieve this,

we made icSHAPE libraries from nucleoplasmic and cytoplasmic fractions of cells transfected with SINEUP-

GFP carrying either mouse antisense *Uchl1* SINEB2 or human SINE RNA FRAM. The 2D structure models

based on this icSHAPE data revealed that the base of the SINEB2 RNA structure is mostly stable between

the nucleus and the cytoplasm, with only a slight variation observed around the region of motif 2, being

slightly more open in the nucleus than in the cytoplasm (**Fig. 3a**, **b**), and a similar structure difference in

human FRAM-transcribed nuclear RNA (**Fig. 3c**, **d**). In addition, the internal loop between motifs 5, 6, and

7 in the nuclear mouse SINEB2 RNA goes through a structural rearrangement in the cytoplasm and

transforms into a cloverleaf-like structure (**Fig. 3a**, **b**). In the case of human FRAM RNA, both nuclear and

cytoplasmic structures are quite similar and form a cloverleaf-like structure, though the cytoplasmic

structure is slightly more structured (**Fig. 3c**, **d**). This could be possibly due to the transient nature of any

alternative structures, as icSHAPE structure represent only the average of the structure ensemble.

The nucleus-specific internal loop structure might play a role in the export of SINEUP RNA to the

cytoplasm, whereas the cytoplasmic cloverleaf-like structure with SL1 might be crucial for interaction with

translation machinery. Interaction with HNRNPK is crucial in both of these SINEUP processes[38]. To check

how the different nuclear and cytoplasmic structures of SINEB2 RNA are associated with HNRNPK, we

prepared seCLIP[46] libraries from the nuclear and cytoplasmic fractions of SINEUP-GFP co-transfected cells

(**Fig. 3e**). The results showed that in the nucleus HNRNPK interacts with short stretches of SINEB2

coinciding in motifs 1, 3, and 4 (**Fig. 3b**), but in the cytoplasm HNRNPK occupied a long double-stranded

region covering the 5' end of the stem in all the motifs except motif 7. Such disparate interactions could

possibly arise from distinct functions of HNRNPK in SINEUP RNA transport and in translation that are

specific to the cellular compartment.

SINEUP structure motifs interact with rRNA

We wanted to further investigate the role of SINEB2 RNA structure in translation because it has features

reminiscent of tRNA; it forms a cloverleaf-like structure in the cytoplasm and possesses two pseudouridine

(Ψ) sites at 16 bp and 112 bp (**Extended Data Fig. 7**). CircRNA IRESs[43] and many other IRESs are known to

interact with 18S rRNA[42,53] to induce cap-independent translation, however the region by which SINEUP-

RNA interacts with ribosomal subunits and polysomes is not known.

Considering the similarities in sequence and structure, we wondered whether such motifs in SINEB2 RNA

can interact with rRNA. To get a better understanding of SINEUP–rRNA interactions, we employed PARIS2,

which has proved useful in capturing direct mRNA–rRNA short interactions[41,54]. We prepared PARIS2

libraries from HEK293T/17 cells transiently expressing SINEUP-GFP (antisense *Uchl1* SINEB2) or

miniSINEUP-GFP (FRAM) along with their target sense EGFP mRNA and sequenced them on a next-

generation sequencing platform. We could capture few incidents of SINEUP-rRNA pairs (**Fig. 4**, **Fig. 5**,

Extended Data Fig. 8), which is understandable given the technical limitation of psoralen crosslinking and

inefficient detection of scarcely expressed RNA duplexes and highly dynamic interactions in PARIS2[41,55].

Nevertheless, it was intriguing to see interactions of mouse SINEB2 element with human rRNA in both

small and large subunits of ribosome (**Fig. 4a**, **b**, and **c**). We discovered that the 3' side of motifs 1 and 2

of SINEB2 RNA interacted with a region in helices 14 and 15 of 18S rRNA (Int 1 in **Fig. 4a** and **c**). On the

other hand, the region between motifs 5 and 6 formed contacts with domain III helices 55, 56, and 59 of

28S rRNA. Motif 5 and the flanking region also interacted with domain I, helix 25, expansion segments

ES7c and ES7d of 28S rRNA (Int 2 and 3 in **Fig. 4b** and **c**).

Interestingly, functionally active human SINE-derived RNA FRAM displayed conservation of this rRNA

interaction trait. With better expression of FRAM RNA, we were able to detect more instances of FRAM-

rRNA interaction than for SINEB2 RNA. Similar to SINEB2, the 3' side of motifs 1 and 2 and of motifs 5 and

6 of FRAM RNA formed contacts with helix 15 of 18S rRNA (Int 2 in **Fig. 5a**, **b**, **c**). In addition, the FRAM

RNA stem loop analogous to SL1 (containing motifs 2/3, 5, and 6) interacted with 18S rRNA helix 21, es6b

and es6c (Int 1 in **Fig. 5a**, **b**, **c**), and helices 37 and 38 paired with the 3' end region of FRAM RNA (Int 3 in

Fig. 5a, **b**, **c**).

Moreover, the 5' side of FRAM RNA flanking motif 1 interacted with 28S rRNA domain III helix 48 and 49a

(Int1 in **Fig. 5d**, **e**, **f**), and helix 50 and 52 (Int5 **Fig. 5d**, **e**, **f**), whereas FRAM SL1 region interacted with 28S

rRNA domain IV, helix 63, ES27b and with domain I, helix 25, ES7c (Int 2 and 3 in **Fig. 5d**, **e**, **f**). The short

stem-loop at the 3' end of FRAM structure was involved in interaction with 28S rRNA domain V, helix 81

(Int4 in **Fig. 5d**, **e**, **f**).

The presence of SINEUP-rRNA pairs in PARIS2 data in addition to our knowledge of SINEUPs' fate in the

polysomes[38], that over 85% of the cytoplasmic SINEUP-GFP co-sediments with EGFP mRNA and HNRNPK

in free/40S ribosomal subunit and monosome fractions, supports the notion of SINEUP-rRNA

interaction[38].

SINE RNA 3D structure models reveal non-canonical intramolecular interactions

The SINE 2D structure models helped us to understand the conservation and functional importance of

local short motifs in maintaining the SINEUP effect by interacting with RBP and rRNA. However, they

lacked information about non-canonical intramolecular interactions that stabilize the RNA structure and

affect the availability of interaction sites in 3D. This results in canonically unpaired regions in the 2D

structures that are actually non-canonically paired in 3D[56]. Therefore, we took advantage of in vivo SINE

RNA 2D structure information and incorporated it into the machine translation of 3D structure models.

Thus, we created 3D models for 17 mouse and one human SINE-derived RNAs (**Fig. 6** and **Extended Data**

Fig. 9). The 3D models revealed that the SINEUP motifs are stabilized by base-stacking interactions along

the double-helical structure and that bases in the internal loop and stem-loop regions form non-canonical

interactions adding more complexity to the structure (**Fig. 6 a**,**c**,**e**,**g** and **Supplementary Tables 2** and **3**).

The molecular surface view of the nuclear SINEB2 RNA shows an elongated helical structure where SL1

takes the apical position (**Fig. 6b**). Interestingly, human nuclear FRAM RNA, which has a cloverleaf-like

conformation in 2D (**Fig. 3d**), folds similarly to nuclear SINEB2 RNA in 3D, with the three arms positioned

in such a way that all the motifs are in the central arm, SL1 is at the apex, and the other two arms are

close to the central one (**Fig. 2d** and **Fig. 6 e**,**f**). In the cytoplasm, both SINEB2 and FRAM RNAs fold into a

cloverleaf-like conformation where SL1 moves from the center to the left position and in FRAM RNA the other two arms open up (**Fig. 6 c**,**d**,**g**,**h**). We also analyzed the 3D conformation of functionally inactive antisense *Gadd45α* SINEB2-a and its deletion mutants in comparison to antisense *Uchl1* SINEB2 (**Extended Data Fig. 10**). The 3D models showed that among all the mutants, the functional Δ27–36 mutant adopts a similar global structure as well as SL1 and motif 6 sub-structures with that of antisense *Uchl1* SINEB2 in 3D space (**Extended Data Fig. 10 a,c**) consistent with the 2D structure data. The wild-type antisense *Gadd45α* SINEB2-a and functionally inactive mutants Δ40–50, Δ66–76 and Δ92–101 which have deformed SL1, vary in RNA 3D architecture from the functional mutant (**Extended Data Fig. 10 a,b,d–f**).

Discussion

There are about 175 families of SINE scattered across different domains of life[57]. Although generally considered as parasite elements of the genome, a growing number of studies point to their gene-regulatory activity, such as enhancer-like function, polyadenylation signal, transcription factor binding sites, mRNA nuclear retention, and translation upregulation[10,15-18,23]. However, there are no extensive studies on the sequence and structural features of functional SINE-transcribed RNA elements. Here, we examined the SINEUP activity of 15 as-yet-uncharacterized SINEB2-derived RNAs from mouse natural antisense transcripts and assayed their function in GFP protein upregulation. Most (12 out of 15) of the SINEB2 RNAs showed SINEUP activity, but the remaining 3 (antisense *Gadd45α*-SINEB2-a and -b, and antisense *Uxt* SINEB2-a) failed to upregulate GFP protein expression significantly under the current experimental settings. The shortest SINEB2 RNA to enhance GFP protein expression, antisense *Txnip* SINEB2, was found to be 107 nucleotides long, which is half the length of the longest functional natural SINEB2-derived RNAs tested here (antisense *Nars2* SINEB2 and antisense *Epb4.9* SINEB2). This suggests that the SINEUP class of RNAs in nature is vast and diverse and is likely to be a much larger class of antisense RNAs than yet explored. Functional observation of the consensus SINEB2 sub-family B3 sequence also supported this, although it enhanced GFP translation to a lesser extent than the highly functional antisense *Uchl1* SINEB2 RNA. This might be because consensus sequences represent the most frequent occurrence of a particular nucleotide in a multiple sequence alignment of related sequences, which may not necessarily be functionally optimal. We observed that in most cases only one repeat of SINEB2 RNA as the effector domain is sufficient to exert the SINEUP effect, possibly because two SINEB2-derived RNAs (which are generally GC-rich) may share some short complementary regions that can interact and hinder the proper RNA folding into key functional features or may compete for binding

proteins and RNA partners. Another interpretation of this non-synergistic nature of these functional
SINEB2 RNAs is that SINEUP-induced protein upregulation is used to fine-tune translation in the cell,
keeping the protein within a physiological range of up to 1.5–3-fold change. As previously observed, this
fold change is sufficient to rescue haploinsufficient gene dosage and disease phenotype in vivo by
upregulation of protein expression of the subunit 7B of cytochrome c oxidase (*cox*7b) in a medaka fish
model of human microphthalmia with linear skin defects (MLS) syndrome[58], and rescue of endogenous
defective Frataxin (FXN) expression in a cellular model of Friedreich's ataxia[59] as well as rescue of
neurodegeneration and motor phenotype by enhancing endogenous Glial cell-derived neurotrophic
factor (GDNF) protein expression in a mice Parkinson's disease model[60]. Incidentally, this characteristic
makes SINEUPs ideal for therapeutic studies where proteins need to be regulated within a physiological
range. The sequence diversity within the same B3 sub-family or in comparison with other sub-families
such as B3A, B2-Mm2, and B2_Mm1t (as per Repbase nomenclature)[27] does not prevent the SINEUP
function. This observation confirms the universality of the SINEUP function in the SINEB2 repeat family
and the flexibility of its members to maintain key functional features despite varying length and
composition of RNA sequences. It would be interesting to explore other families of SINE elements in other
species and across a wide variety of mammalian and eukaryotic genomes in the future.

Noting the sequence diversity of functional SINEB2-derived RNAs and the importance of RNA structure in
non-coding RNA functions, we looked at the role of the SINE RNA secondary structure in the SINEUP
phenomena in living cells by using icSHAPE experiments. Consequently, we noticed a high correlation
between the in-cell 2D structure of antisense *Uchl1* SINEB2 RNA and the previously reported structure
from chemical footprinting[11]. Moreover, we found that the key functional SL1 structure was conserved
among various SINEB2s RNAs. When the functionally dormant antisense *Gadd45α* SINEB2-a RNA that lacks
an SL1 region was mutated to change its structure, one of the mutants (Δ27–36) acquired a properly
folded SL1 structure and a 2D structure similar to the icSHAPE structure of antisense *Uchl1* SINEB2 RNA,
resulting in SINEUP activity gain. This further emphasizes the functional significance of this short structure
in SINEUP-induced gene regulation.

We discovered 7 conserved structure motifs among the icSHAPE data-driven 2D structure models of all
the SINEB2-derived RNAs tested here, though only partial matches in sequence were observed, given the
sequence variation. Since we observed partial matches to these structure motifs in the functionally
inactive SINEs also, we hypothesize that the SINEUP function is influenced not just by the mere presence
of these structure motifs but also by their relative positions on the structure. Some structures might be

sterically more exposed to interaction with molecular machinery in the 3D space. Additional structural
regions might help in proper folding of these motifs and play an accessory role in maintaining the SINEUP
function. Thus, any deletion might inadvertently disrupt functional structures and interaction with RBPs.

We recently reported that deletion of certain regions in antisense *Uchl1* SINEB2 RNA is detrimental to
SINEUP function[12]. Several of these deleted regions correspond to the SINEUP structure motifs identified
in this study: Δ100-111 disrupts motif 7, Δ130-141 is the deletion of the 3' end side of motifs 3 and 4, and
Δ5-14 disrupts motifs 1 and 2[12]. Interestingly, their Δ65–74 mutant[12], which deletes part of motif 6, does
not show a significant decrease in the SINEUP function, whereas another study demonstrated that
deletion of SINEB2 RNA SL1 consisting of the entire motif 6 abolishes SINEUP activity[11]. The in-cell
structure of human SINE-derived FRAM RNA also showed a partial match for SINEUP motifs and the
presence of a structural feature analogous to SL1. Furthermore, we found similarity between the structure
of the motif 6 sequence in 2D and that of a motif in 18S rRNA in 3D, and the structure of motif 2 was found
to be similar to a region in 16S rRNA in 3D. The matching region in 18S rRNA belongs to helix 34, which
forms the roof of the mRNA entry channel. The similarity of the SL1 region of SINEB2 RNA with rRNA was
previously noticed in a molecular dynamics simulation based on NMR data[44], and here in-cell analysis
again showed that the conserved and functionally crucial SINEUP motifs were similar to rRNA structure
motifs.

Mouse SINEB2s originate from tRNA element[57,61], and pseudouridylation of tRNA bases is reported to be
crucial for its stability and function[35]. Not only tRNAs, but all the RNAs involved in translation contain
several Ψ-modified sites[34,35]. This was also found to be the case for SINEUPs in this study. We identified
two Ψ modifications at positions 16 and 112 of antisense *Uchl1* SINEB2 RNA. The Ψ at position 112 is
inside motif 5, and that at position 16 is just at the motif 2 boundary. This indicates that pseudouridylation
of these SINEB2 RNA positions might help to stabilize the structure of these motifs by forming extra
hydrogen bonds by N1-H position of Ψ or by base-stacking, which is an intrinsic property of Ψ[34,62].

We further analyzed the structural dynamics of both SINEB2 and FRAM RNAs in the nucleus and cytoplasm
and identified two regions that differ between their nuclear and cytoplasmic structures, with motif 2 and
the region between motifs 5 and 6 of SINEB2 RNA slightly more open in the nucleus than in the cytoplasm.
Less structured regions were also observed in the nuclear structure of FRAM RNA. We wondered whether
these nucleus-specific structures can provide binding sites for proteins involved in nuclear transport. We
knew that binding to HNRNPK is crucial for cytoplasmic co-localization of SINEUP-GFP and sense mRNA
complex and their association with polysomes[38]. Furthermore, HNRNPK is reported to interact with c-myc

IRES and stimulate its cap-independent translation[63]. Since HNRNPK participates in both nuclear transport

and translation regulation, we checked if structurally dynamic regions have any effect on its interaction in

subcellular compartments. Indeed, we observed distinctive binding patterns of HNRNPK in the nucleus

and cytoplasm; in the nucleus, it interacts with short regions in motifs 1, 3, and 4 of SINEB2 RNA, but in

the cytoplasm, it interacts with a much longer double-stranded region covering all but one motif. These

particular binding patterns could be connected to different functions of HNRNPK in the nucleus and

cytoplasm.

In the cytoplasm, the loop region between motifs 5, 6, and 7 of mouse SINEB2 RNA assumes a cloverleaf-

like structure, formed from a multiloop where four stems converge. One stem is SL1, which harbors motif

6, another stem consists of motif 7, and a disrupted motif 5 makes the top of the central stem where the

Ψ at position 112 is situated at the multiloop branching point. Interestingly, though the nuclear and

cytoplasmic structures of FRAM RNA are highly similar, we discerned a similar cloverleaf-like structure.

icSHAPE structures are generally enriched for the most stable and dominant structural conformations in

the cell, so any alternative short-lived structures might not have been captured. It is fascinating to see

that mouse SINEB2 RNA that originated from tRNA and human FRAM RNA that originated from 7SL RNA[61]

share common characteristics: short conserved structural motifs, a cloverleaf-like structure in the

cytoplasm, and positive regulation of translation. Previously, we created a minimal functional SINEUP by

deleting nucleotide positions 1-30 and 120-167 from SINEB2 RNA, thus keeping only nucleotide positions

31-119 which is the region corresponding to motifs 5, 6, and 7 as well as SL1 and cytoplasmic cloverleaf-

like structure[12]. This mutant upregulated GFP mRNA translation and retained about 80% of full length

SINEB2 functional activity supporting the significance of this structural region in SINEUP function[12].

In addition to the intramolecular structure, we were interested to better understand the intermolecular

interactions of SINE-derived RNAs with translation machinery. Interaction with 18S rRNA has been

reported to be crucial for both linear and circular IRESs[42,43,53]. Knowing that SINEUP-GFP RNA co-

sediments with free small subunits of ribosomes and monosomes, we checked direct interaction of SINE

RNA with rRNAs by PARIS2 and observed such RNA pairs. The sparsity of SINEUP–rRNA interactions may

be due to a limitation of psoralen cross-linking, which has bias for staggered uridines and may not access

RNA structures positioned inside complex ribonucleoprotein regions[55]. Alternatively, the efficiency of

PARIS2 to capture RNA–RNA interactions may be limited by the level of expression of RNAs as well as by

the brief half-lives of such interactions[41,55]. Nevertheless, our PARIS2 data identifies for the first time a

direct interaction of SINEUP-GFP (SINEB2) RNA and miniSINEUP-GFP (FRAM) RNA with both 18S and 28S

rRNA, which is consistent with previous polysome profiling data[38]. Even though we capture long RNA-RNA

interactions, in terms of sequence complementarity we unambiguously identify only short discontinuous

stretches of base-pairing between SINE-derived RNA and rRNA. However, this may still be sufficient to

influence translation, given that circular IRESs have 7-mer–long complementarity to 18S rRNA[43] and HCV

IRES needs only 3 nucleotide-long Watson-Crick base-pairing with 18S rRNA to induce translation[53].

Moreover, RNA–RNA interactions in 3D depend heavily on non-Watson-Crick base pairing and non-

covalent interactions, such as base-stacking and tertiary contacts by purine to minor-groove and tetraloop

to tetraloop receptor[64], which are not easily predicted. Therefore, we decided to take a closer look at the

SINE-rRNA interactions.

Notably, we discovered an interaction between a region comprising helix 14 and 15 of 18S rRNA (Int 1)

and motif 1 and 2 of SINEB2 RNA. FRAM RNA motifs 1 and 2 along with motifs 5 and 6 also interacted with

helix 15 of 18S rRNA. Helix 14 of 18S rRNA is known to interact with ribosomal protein rpL23[65]. In addition,

a protein ABCE1 (ATP Binding Cassette Subfamily E Member 1), which is involved in translation initiation

and termination, interacts with helix 14 during translation initiation[66]. Moreover, helices 14 and 15 bind

with eukaryotic initiation factor 5b (eIF5b), a ribosome-dependent GTPase that mediates joining of 40S

and 60S subunits of ribosomes[67]. This indicates the possibility that SINEUP motifs 1 and 2 are involved in

the joining of ribosomal subunits during translation initiation. Furthermore, we observed contact of the

region between motif 5 and 6 of SINEB2 RNA with domain III helices 55, 56, and 59 of 28S rRNA. Helix 59

is a part of the ribosomal tunnel exit site, from where the nascent peptide emerges out of the ribosome[68].

It suggests that SINEB2 RNA not only takes part in initiating target mRNA translation but also in elongation.

In addition, SINEB2 RNA motif 5 and the surrounding region was found to bind with domain I, helix 25,

expansion segments ES7c and ES7d of 28S rRNA. The connection with ES7c was also found to be conserved

in FRAM RNA, in which the SL1 region (containing a partial match to motif 5) participated in the

interaction. ES7 is the largest expansion segment positioned at the back of the 60S ribosomal subunit. This

highlights the importance of SINEUP structure motifs in their interaction with the large subunit of the

ribosome. We also observed interaction of FRAM RNA SL1 with 18S rRNA helices 21, 37, and 38. Because

the helix 38–42 region is crucial for mRNA binding and placement of anticodon arms for tRNAs[69], this

interaction is notable for the proper positioning of SINEUP target mRNA and translation initiation. FRAM

RNA also interacted with domains I, III, IV, and V of 28S rRNA.

These contacts at multiple rRNA sites indicate the dynamic changes in SINE RNA–rRNA interaction that

occur as translation progresses and highlight the role of SINEUPs in regulating both translation initiation

and elongation. Moreover, SINEUP RNAs were found to interact with other protein-coding RNAs and

lncRNAs (**Supplementary Table 4**). However, among these RNA partners only *Ank3* (Ankyrin 3) was

observed to interact with the SINEB2 RNA region. ANK3 protein links membrane proteins to spectrin-actin

cytoskeleton and is involved in neuronal development[70], but role of *Ank3* mRNA in translation is not

known. It is to be noted that although we captured the dynamics of SINE RNA structure and interaction

with rRNA and RBP, many of these processes are very transient in nature and it is unlikely to obtain a

complete view of all intermediate RNA structures and interaction kinetics in possibly very short time

windows by existing in vivo methods. Considering this, further 3D structure analysis of the tertiary

connections of SINEUP RNA with rRNA and translation complexes by using cryogenic electron microscopy

or other approaches is needed in the future to construct a full 3D model of larger molecular complexes.

In this study, the experimental 2D structure data-based prediction of 3D structure models for 18 SINE-

derived RNAs identified stabilizing stacking interactions within structure motifs and demonstrated

strikingly similar dynamics of mouse and human SINE-derived RNAs in 3D space in the subcellular

compartments. We list all the base-stacking and non-canonical interactions determined from these

models, which will provide a foundation for future experimental 3D structure analysis. In addition, the 3D

structure models of functionally inactive antisense *Gadd45α* SINEB2-a and related structure mutants

confirmed the similarity in global structure and SL1 of functional Δ27-36 mutant with antisense *Uchl1*

SINEB2 which was missing in the wild-type and other non-functional mutants. This further supports the

idea that such short structure motifs are crucial for non-coding RNA function.

In summary, this study offers a repertoire of the sequences, structures, and functional features of several

SINE-derived RNA elements for the first time and demonstrates that evolutionarily distant SINE RNAs have

a convergent translation upregulatory function driven by common short structure motifs. We show that

mouse and human functional SINE-derived RNAs follow a similar pattern of structural dynamics and SINE

RNA–RBP interactions are compartment-specific. Our study reveals direct interaction of mouse and

human SINE-derived transcripts with human small and large subunit of rRNAs, suggesting a mechanism

for SINEUP-mediated post-transcriptional gene regulation. This is the first study to provide a large number

of RNA 3D structure models of functional SINE-derived RNAs. Altogether the data presented here is a key

step towards filling the void in the structural and functional analysis of SINE-transcribed RNA and will

facilitate future studies that aim to resolve the SINEUP mechanism and biomolecular complexes involved.

Data availability

The icSHAPE sequencing data were deposited in Gene Expression Omnibus (GEO) with accession numbers

GSE146407 (for whole cell icSHAPE libraries of SINEUP-GFP) and GSE224534 (for whole cell icSHAPE

libraries of miniSINEUP-GFP and nuclear-cytoplasmic fractionated icSHAPE libraries), and GSE243220 (for

whole cell icSHAPE libraries of antisense *Gadd45α* SINEB2-a deletion mutants). The PARIS2 sequencing

data and SINEUP RNA modification data are accessible through GEO accession numbers GSE224533 and

GSE224018 respectively. Nuclear-cytoplasmic fractionated seCLIP sequencing data is available through

the GEO accession number GSE227250.

Acknowledgements-

We are grateful to the late Dr. Silvia Zucchelli (Department of Health Sciences, Center for Autoimmune

and Allergic Diseases (CAAD) and Interdisciplinary Research Center of Autoimmune Diseases (IRCAD),

University of Piemonte Orientale, Novara, Italy) for her valuable suggestions and scientific discussion. This

work is dedicated to her memory. We are thankful to Dr. Howard Y. Chang from Standford University for

gifting us icSHAPE-probing reagent. We thank Dr. Byron Lee from Chang lab at Stanford University for

helpful suggestions for successful icSHAPE library preparation. We thank Souta Nagayama from P.

Carninci's lab for his help in deletion mutant analysis.

Funding

This work was supported by RIKEN Centre for Integrative Medical Sciences and RIKEN Centre for Life
Science Technology from the Japanese Ministry of Education, Culture, Sports, Science and Technology
(MEXT); and Basic Science and Platform Technology Program for Innovative Biological Medicine grant
from the Japan Agency for Medical Research and Development (AMED grant no.: 18am0301014 to PC).

Conflict of interest statement. S.G. and P.C. are inventors on patent US9353370B2 and related applications
in Europe, USA and Japan, and H.S., H.T., S.G., and P.C. are inventors on patent application
102018000002411 and the related application PCT/IB2019/050914 held by SISSA in Italy and TranSINE
Therapeutics in the UK, of which S.G. and P.C. are founders and H.T. owns shares.

H.S., N.T., M.V., H.T., and P.C. are inventors on the E.U. patent application EP3992289A1
(WO2022090733A1) held by TranSINE Therapeutics in the UK. S.G., N.T., H.T., and P.C. are inventors on
patent application WO2022064221A1 held by TranSINE Therapeutics in the UK. These conflicts of interest
were not associated with the funding of this work.

Author Contributions

H.S. H.T. and P.C wrote, reviewed, and edited the manuscript with comments from all the authors and interpreted the data. H.S. designed and performed the SINEUP functional analysis, whole cell and nuclear-cytoplasmic fractionated icSHAPE, icSHAPE for deletion mutants and analyzed respective data. H.S. performed 2D and 3D structure modelling and analyzed PARIS data. M.N.Z.V. analyzed seCLIP data and RNA modification data from Oxford Nanopore direct-RNA sequencing. N.T. performed seCLIP, RNA modification experiments and prepared direct-RNA sequencing libraries. H.S.N. prepared PARIS libraries and icSHAPE libraries for deletion mutants. H.T., P.C, and S.G. conceptualized and provided scientific discussion. P.C. provided resources and secured funding for the project.

Materials and Methods

Plasmids and cloning

SINEUP-GFP (pcDNA3.1-based) synthetic SINEUP containing antisense *Uchl1* SINEB2[10] was used to construct long SINEUP effector domain–swapping mutants. To swap the antisense *Uchl1* inverted SINEB2 of SINEUP-GFP with other SINEB2 sequences of interest, two restriction sites for SacII and ClaI restriction enzymes were introduced at the start and end of the SINEB2. For this purpose, a QuikChange II Site-Directed Mutagenesis Kit (Agilent, catalog #200523) was used. All the mutagenesis primers were designed using QuikChange Primer Design Program (Agilent) and are listed in **Supplementary Table 5a**. The SINEB2 regions were PCR-amplified from RIKEN mouse cDNA clones AK143784 (antisense to *Txnip*) and AK054076 (antisense to *Gadd45α*), then digested by the SacII and ClaI restriction enzymes and sub-cloned in SINEUP-GFP to create swapping mutants of antisense *Txnip* and antisense *Gadd45α* SINEB2-a and -b respectively. Antisense *Gadd45α* SINEB2-ab and ba, and all the full constructs of antisense *Uxt* SINEB2–swapping mutants were obtained from Genewiz. The highly active antisense *Uchl1*_Δ5'-32nt mutant[30] was included in screening as a positive control. Direct and deleted antisense (ΔAS) *Uchl1* SINEB2 mutants were created by a conventional restriction digestion, PCR, and cloning method, and the SINEUP mutant with deleted binding domain (ΔBD) was purchased from TransSINE Technologies.

Previously described miniSINEUP-GFP[29] plasmid consisting of antisense *Uchl1* SINEB2 as the effector domain and −40/+4 binding domain design with respect to the start codon of GFP mRNA was used as a backbone for miniSINEUP effector-domain swapping. A conventional PCR amplification, restriction digestion, ligation, and cloning strategy was used to generate miniSINEUP-GFP mutants shown in **Fig. 1e**. The SINEB2 region of RIKEN mouse cDNA clones AK029359 (from 774–960 bp for antisense *Uxt-b* SINEB2), AK032194 (from 769–982 bp for antisense *Nars2* SINEB2), AK041742 (from 2334–2533 bp for antisense *Abhd11* SINEB2), and AK048309 (from 1203–1416 bp for antisense Epb4.9 SINEB2) was amplified by PCR (all primer sequences are shown in **Supplementary Table 5b**). To prepare the effector domain for antisense *Uchl1* SINEB2 + antisense *Uxt-b* SINEB2 (**Fig. 1e**), the SINEB2 regions of RIKEN cDNA clones AK078321 (from 521–690 bp for antisense *Uchl1* SINEB2) and AK029359 (from 774–960 bp for antisense *Uxt-b* SINEB2) were first amplified by PCR separately and then ligated to make one effector-domain insert (all primer sequences are shown in **Supplementary Table 5b**). Similarly, in the case of 3× antisense *Uchl1* SINEB2, PCR-amplified SINEB2 from AK078321 (from 521–690 bp) was consecutively ligated twice to generate three repeats of the same SINEB2. The rest of the SINEB2 effector domains originating from various mouse natural antisense lncRNAs (**Fig. 1e**, and **Supplementary Table 1**) and the consensus

sequence for SINEB2 subfamily B3 from the Repbase database[27,47] were commercially obtained from
Hokkaido System Science Co., Ltd. The backbone plasmid, miniSINEUP-GFP, and amplified/commercial
SINEB2 effector-domain inserts were digested by EcoRI and HindIII restriction enzymes, then ligated and
cloned.
The commercially available pEGFP-C2 (Clontech) plasmid was used as a sense EGFP construct. The
miniSINEUP-GFP FRAM plasmid design has been previously described[31].
610

Cell culture and transfection

Human embryonic kidney (HEK293T/17) cells (CRL-11268, ATCC) were grown in a 10-cm culture dish in
DMEM (1×) + GlutaMAX-1 (Thermo Fisher Scientific, 10569-044) supplemented with 10% fetal bovine
serum (Sigma, 172012) and 1% penicillin streptomycin solution (Wako, 168-23191) at 37°C and 5% CO_2
for 3–5 days. 70–90% confluent cells were treated with 0.05% w/v Trypsin-0.53 mmol/L EDTA 4Na Solution
with phenol red (Wako, 202-16931) for 5 min at 37°C to detach the cells, and 0.5×10^6 cells were seeded
in each well of 6-well plates (Falcon). After 24 h, cells were transfected with pEGFP-C2 and SINEUP-GFP
(long) or miniSINEUP-GFP swapping mutant plasmids at a molar ratio of 1:4.3 for long SINEUPs GFP [0.19
pmol pEGFP-C2 + 0.81 pmol of long SINEUP] and 1:4.6 for miniSINEUP-GFP [0.19 pmol pEGFP-C2 + 0.87
pmol of miniSINEUP-GFP] using 10 μL of Lipofectamine2000 (Thermo Fisher Scientific, 11668-019). After
24 h of transfection, cells were harvested.
622

Protein extraction and Western blot

Cells were washed with D-PBS (-) (Wako, 045-29795) and lysed using 140 μL of 10× Cell Lysis buffer (Cell
Signaling Technology, 9803S) with PMSF (Cell Signaling Technology, 8553S) per well in a 6-well plate.
Samples were rotated at low speed for 1 h at 4°C and then centrifuged (20,000 × g) for 10 min at 4°C. The
supernatant was collected, and the protein concentrations were measured by DC Protein Assay (BioRad,
500-0116). The absorbance (750 nm) was measured by Multimode Plate Reader ARVO X3 (PerkinElmer).
For Western blotting, 20 μg of extracted proteins mixed with 2× SDS sample buffer were separated on
10% SDS PAGE gel (Mini PROTEAN TGX Precast Gel, 10%, 12-well comb; BioRad, 456-1035) and transferred
to a nitrocellulose membrane (Amersham Protran Premium ECL 0.45 μm; GE Healthcare Life Sciences,
10600013) through semi-dry transfer at 25 V for 30 min using Trans-BLOT SD Semi-dry Transfer Cell
(BioRad). Tris-glycine buffer containing 20% methanol was used for transfer. The membranes were
blocked with 5% nonfat dry milk (Cell Signaling Technology, 9999S) in Tris Buffered Saline with Tween-20
(10× TBST, Cell Signaling Technology, 9997) for 30 min with shaking. Proteins were immunoblotted with
primary antibodies followed by horseradish peroxidase– (HRP) conjugated secondary antibodies. EGFP
was detected by anti-GFP rabbit serum A-6455 (Life Technologies), ACTINB was detected by monoclonal
anti-β-actin antibody (Sigma Aldrich, A5441). Protein signals were detected by ECL Western Blotting
Detection Reagent (GE Healthcare Life Sciences, RPN2109) with FUJI LAS-3000 system (Fujifilm) and
FUSION (Vilber-Lourmat). The band intensities were analyzed by Image J version 1.48 software (National
Institutes of Health).
642

RNA extraction, cDNA synthesis, and quantitative RT-PCR

HEK293T/17 cells were washed with D-PBS (-) (Wako, 045-29795) and detached using 0.05% w/v trypsin
(Wako, 202-16931). Cells were collected by centrifugation for 5 min (6000 × g) at 4°C, and total RNAs were
subsequently extracted using RNeasy Mini Kit (Qiagen, 74104). Extracted RNA samples were processed
through three rounds of DNase I digestion using a TURBO DNA-free Kit (Invitrogen, AM1907). RNA quality
was checked on Agilent 2100 Bioanalyzer (Agilent Technologies) using an Agilent RNA 6000 Nano Kit. Then,
500 ng of the total RNA was used for cDNA synthesis using the Prime Script 1st Strand cDNA Synthesis Kit
(TaKaRa, 6110A). Random 6-mer primers (0.3 μL) and 1.7 μL of oligodT primers were mixed to capture
RNAs. 2× diluted cDNA (1 μL) was used as a template for the quantitative real-time PCR (qRT-PCR). For
qRT-PCR, a SYBR Premix Ex Taq (Tli RNaseH Plus) Kit (TaKaRa, RR420S) was used. Each reaction contained
1 μL of cDNA, 0.4 μL of reverse primer (10 μM), 0.4 μL of forward primer (10 μM), 10 μL of SYBR premix
ExTaq, 0.4 μL of ROX reference dye (50×), and 10 μL of nuclease-free water (reaction volume was 20 μL).
EGFP and SINEUP-GFP levels were normalized by GAPDH. qRT-PCR was performed by StepOnePlus and
7900HT Real-Time PCR Systems (Applied Biosystems) under the following conditions: hold stage (1 cycle)
for 30 s at 95°C, cycling stage (40 cycles) for 5 s at 95°C and for 30 s at 60°C, melt curve stage. Reverse
transcription samples were taken to check for any plasmid DNA contamination in cDNA. Each sample was
performed in technical triplicates (n = 3, Cт standard deviation > 0.2) and with biological replicates (n ≥ 3).
2^-ΔΔCt ±SD was analyzed by StepOne v2.3 and SDS v2.4 software (Thermo Fisher Scientific). The qRT-
PCR primer sequences were as follows:

hGapdh_Fw: TCTCTGCTCCTCCTGTTC

hGapdh_Rv: GCCCAATACGACCAAATCC

EGFP_Fw: GCCCGACAACCACTACCTGAG

EGFP_Rv: CGGCGGTCACGAACTCCAG

SINEUP-GFP_Fw: CTGGTGTGTATTATCTCTTATG

SINEUP-GFP_Rv: CTCCCGAGTCTCTGTAGC

miniSINEUP-GFP_Fw: TTAATACGACTCACTATAGGGAGAC

miniSINEUP-GFP_Rev: CCGCATCTAGTGAACCGTCA

Phylogenetic analysis and hierarchical clustering

MEGA version 5 software[71] was used for sequence-based alignment, and mouse SINEB2 sequences tested
as effector domains in long and mini SINEUPs were aligned by ClustalW. SINEB2 sequence–based
molecular phylogenetic analysis was done using the Maximum Likelihood method. The evolutionary
history was inferred by the Maximum Likelihood method based on the Tamura 3-parameter model. Initial
tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ
algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood
approach and then selecting the topology with superior log likelihood value. A discrete Gamma
distribution was used to model evolutionary rate differences among sites (5 categories (+G, parameter =
4.1774)). The tree was drawn to scale, with branch lengths measured in the number of substitutions per
site.

icSHAPE library preparation

The icSHAPE standard protocol was followed[12,49]. Briefly, HEK293T/17 cells were co-transfected with pEGFP-C2 and SINEUP-GFP or miniSINEUP-GFP swapping mutants using Lipofectamine2000 reagent in 6-well plates. Cells were harvested after 24 h of transfection, pooled together, and treated with NAI-N$_3$ reagent (modified) (NAI-N$_3$ reagent was received as a gift from Dr. Howard Chang's lab at Stanford University) for in vivo SHAPE modification or with DMSO (mock control). Two biological replicates for each of the NAI-N$_3$ and DMSO libraries were made. RNA was extracted using RNeasy Mini Kit (Qiagen, 74104) and digested with DNaseI to remove DNA contamination using a TURBO DNA-free Kit (Invitrogen, AM1907). RNA was ribo-depleted using a magnetic RiboZero kit (Illumina, MRZH11124), and RNA quality was checked by Agilent 2100 Bioanalyzer (Agilent Technologies) nano and pico chip. Next, 500 ng of ribo-depleted RNA was biotin-clicked to label-probed bases followed by RNA fragmentation for 40 s using RNA fragmentation reagents (Thermo Fisher Scientific, AM8740). Fragmented RNA was end-repaired, and the icSHAPE adapter ligation standard protocol was followed[49] (adapter sequences in **Supplementary Table 6a**). Next, size selection was performed, and RNA was reverse-transcribed using icSHAPE-specific barcoded reverse-transcription primer followed by streptavidin-based selection and size selection of truncated cDNA products. Purified cDNA was circularized using CircLigase II enzyme (Epicentre, CL9025K), then library PCR, size selection, and quantification were performed. Resultant libraries were sequenced on a HiSeq2500 platform (Illumina, 100 bases, single-end). All the oligos used in the icSHAPE library preparation are noted in **Supplementary Table 6a**.

Comparison of icSHAPE profiles, secondary structure modelling, and motif analysis

The standard icSHAPE bioinformatics pipeline was used[49]. Briefly, after removing PCR duplicates and adapter trimming, raw sequencing data were mapped to EGFP and SINEUP transcripts index using Bowtie2. Next, transcript abundance and reverse transcription (RT) stop numbers were calculated before calculating the correlation between replicates and replicate merging. The mean of the RT stops of the 90–95% most reactive bases was normalized as 1 and all other RT stops were scaled proportionally. Then enrichment reactivity scores were calculated by subtracting the background signal (DMSO data), and outliers were removed by 90% Winsorization (in which the top 5th percentile is set to 1 and the bottom 5th percentile is set to 0). Valid enrichment reactivity scores were filtered and visualized in integrated genome viewer (IGV v2.11.1). The icSHAPE enrichment scores were used as a soft constraint in RNAfold (ViennaRNA Package)[72]. Linear mapping method was used to derive pairing probabilities, and a method described by Zarringhalam *et al.* (2012)[73] was used to incorporate guiding pseudo energies into the folding algorithm. To draw the secondary structure, RNAcanvas[74] (https://rna2drawer.app/) was used.

Sequence and icSHAPE-guided 2D structure of all the SINEB2s were pairwise compared with the structure and sequence of antisense *Uchl1* SINEB2 using ExpaRNA tool[50], which computes the exact matching sequence-structure motifs common to two RNAs. Thus, derived motif files from all the pairwise comparisons were then screened to extract the positions of Uchl1 SINEB2s that were present in one or multiple motifs in all of the comparisons. Since SINEB2 sequences are highly variable, in most cases only a partial sequence match could be found. Due to their repeating sequence composition, in some instances one query sequence was matched to multiple motifs, or one motif matched to multiple positions in the

723 query. In that case, the longer matching stretch was preferred, and different motif numbers were assigned
724 based on their sequence and structure neighborhood and position on SINEB2. Additionally, SINEUP 2D
725 motifs were compared with existing 3D motifs in the RNA Characterization of Secondary Structure Motifs
726 database (CossMos)[51].

Nuclear-cytoplasmic fractionation for icSHAPE

HEK293T/17 cells were co-transfected with 0.8 pmol SINEUP-GFP or miniSINEUP-GFP FRAM (0.8 pmol)
and 0.2 pmol sense pEGFP-C2 plasmids (Clontech) in each well of a 6-well culture plate. Four 6-well plates
for each sample were prepared. At 24 h post-transfection, the medium was aspirated, and the cells were
washed and collected in 1 mL of 1× D-PBS (-) (Nacalai tesque, 14249-24) per well. Cells from one 6-well
plate (~1×10^7 cells) were pooled together in a 15-mL tube and centrifuged at 300 × g for 2 min at room
temperature. The resultant supernatant was removed, and the pelleted cells were resuspended in either
250 μL of 100 mM NAI-N3 reagent (Enamine, EN300-247313) for modification or in 5% DMSO (Sigma
Aldrich, D2650) solution in 1× PBS (Gibco) for mock treatment. The reaction was incubated on a rotating
wheel at 37°C for 5 min at room temperature and then stopped by centrifugation at 2500 × g for 1 min at
4°C and further removal of the supernatant. Next, the protocol by Pandya-Jones and Black (2009)[75] was
utilized for cellular fractionation. In brief, the NAI-N3- and DMSO-treated cell pellets were resuspended
in ice-cold NP-40 lysis buffer (10 mM Tris-HCl pH 7.5, 0.15% NP-40, 150 mM NaCl), then the lysate was
added on top of 2.5 volumes of chilled 25% sucrose cushion in lysis buffer without NP-40 followed by
centrifugation at 19,000 × g for 10 min at 4°C. The resultant supernatant containing the cytoplasmic
fraction was collected. The pellet was further gently rinsed with 200 μL chilled 1× PBS/1 mM EDTA and
centrifuged at 9,730 × g for 2 min at 4°C to remove the supernatant. Next, to extract the nuclear fraction,
the protocol by Brugiolo *et al.* (2017)[76] was followed. The pellet was resuspended in 100 μL nuclear buffer
I (50% glycerol v/v, 20 mM Tris-HCl pH 7.9, 75 mM NaCl, 0.5 mM NaCl, 0.85 mM DTT, 0.125 mM PMSF)
and then in 900 μL nuclear buffer IIA (20 mM HEPES pH 7.6, 300 mM NaCl, 0.2 mM EDTA, 1 mM DTT, 7.5
mM $MgCl_2$, 1M Urea, 1% NP-40, 400 U of RNase inhibitor). The solution was mixed thoroughly by vortexing
and then pipetting up and down, and then incubated on ice for 20 min. Finally, the nucleoplasmic fraction
was extracted by centrifuging at 14,000 × g for 5 min at 4°C and collecting the supernatant. The success
of fractionation was checked by Western blot and qRT-PCR for known nuclear and cytoplasmic marker
genes. Four nucleoplasmic or cytoplasmic fractions, each coming from four 6-well plates of the same
sample, were pooled together for further RNA extraction and icSHAPE library preparation as described
above. All the steps are the same as in the original protocol[49], except the removal of free 3' RNA linkers
after ligation, for which the completed ligation reaction was treated with 4 μL RecJ$_f$ (New England Biolabs,
M0264L) and 3 μL 5' Deadenylase (New England Biolabs, M0331S) at 37°C for 1 h and further cleaned up
by using a Zymo RNA Clean and Concentrator-5 purification kit (Zymo Research, R1016). All the oligos used
in the library preparation are noted in **Supplementary Table 6a**.

seCLIP library preparation from nuclear-cytoplasmic fractionated samples

Approximately 0.5×10^7 HEK293T/17 cells were plated on a 10-cm culture dish and after 24 h, transfected
with SINEUP-GFP and pEGFP-C2 plasmids as described above. After 24 h of transfection, 2×10^7 cells were
gently washed with 1× PBS and then crosslinked by 400 mJ/cm, 254 nm UV light irradiation on ice with

the lid off for 15 min. Next, the cells were collected by scraping in 1 mL chilled PBS followed by
centrifugation at 200 × g for 5 min at 4°C. The resultant supernatant was discarded, and the pellet was
used for nuclear-cytoplasmic fractionation following the protocol by Brugiolo *et al.* (2017)[76]. The pellet
was first resuspended in 2 mL hypotonic buffer (10 mM Tris-HCl 7.5 pH, 10 mM KCl, 1.5 mM MgCl$_2$, 0.5
mM DTT, 1:200 proteinase inhibitor just prior to use) and incubated on ice for 5 min followed by
centrifugation at 500 × g for 10 min at 4°C. The supernatant containing hypotonic fraction was discarded,
and the pellet was resuspended in 1 mL lysis buffer 0.3 (50 mM Tris-HCl 7.5 pH, 150 mM NaCl, 2 mM
MgCl$_2$, 0.3% NP-40, 1:200 proteinase inhibitor just prior to use) and incubated on ice for 10 min. The mix
was then centrifuged at 1000 × g for 5 min at 4°C, and the supernatant containing the cytoplasmic fraction
was collected in a new 1.5-mL tube. The pellet was mixed with 1 mL lysis buffer 0.5 (50 mM Tris-HCl 7.5
pH, 150 mM NaCl, 2 mM MgCl$_2$, 0.5% NP-40, 1:200 proteinase inhibitor just prior to use) and incubated
on ice for 10 min followed by centrifugation at 1000 × g for 5 min at 4°C. The supernatant was discarded,
and the pellet was first resuspended in 100 µL nuclear buffer I (as used for icSHAPE fractionation, without
addition of PMSF) and then in 900 µL nuclear buffer IIA (as used for icSHAPE fractionation) with 1:200
proteinase inhibitor just prior to use. The pellet was mixed well by vortexing and then pipetting up and
down, and then incubated on ice for 15 min. The reaction mix was then centrifuged at 15,000 × g for 5
min at 4°C, and the nucleoplasm containing supernatant was transferred to a new 1.5-mL tube. All
fractions were separately centrifuged at 20,000 × g for 5 min at 4°C and tested for the presence of markers
specific to nuclear or cytoplasmic fractions by Western blot.

Next, the seCLIP protocol described by Van Nostrand *et al*. (2017)[46] was utilized to prepare nuclear and
cytoplasmic seCLIP libraries. In brief, nucleoplasmic and cytoplasmic lysates were treated with 4 U of
DNase I and 10 µL of 1:25 diluted RNase I (100 U/µL) in a thermomixer at 37°C and 1200 rpm for 5 min.
The reaction was then cooled on ice, 7 µL Murine RNase Inhibitor (New England Biolabs, M0314) was
added, and the mix was centrifuged at 15,000 × g for 15 min at 4°C. The resultant supernatant was mixed
with 15 µg of anti-hnRNPK antibody (Abcam, ab39975) coupled to 200 µL magnetic beads (Dynabeads M-
280 Sheep Anti-mouse IgG, Invitrogen, 11201D) and incubated at 4°C overnight. Around 2% of the input
samples (containing all RNA-protein complexes) were saved before immunoprecipitation.
Immunoprecipitated samples were first end-treated with FastAP thermo-sensitive alkaline phosphatase
(Thermo Scientific, EF0651) and T4 poly nucleotide kinase (New England Biolabs, M0201L) and then
ligated with 3' RNA linker (with sample barcodes, see sequence in **Supplementary Table 6b**). Next, both
immunoprecipitated and input RBP–RNA complexes were visualized by Western blot imaging and the
region corresponding to HNRNPK and above (from 55–180 kDa) was sliced from the preparative
nitrocellulose membrane blot. The RNA was extracted by treating the membrane slices with urea and
proteinase K followed by acid phenol–chloroform extraction and purification by using a Zymo RNA Kit
(R1013). Further treatment of samples and cDNA library preparation are the same as described in the
original protocol[46]. Libraries were sequenced on the HiSeq2500 platform (150 cycles, single-end).

Analysis of nuclear-cytoplasmic fractionated seCLIP data

Fastq files were processed with two rounds of cutadapt (which trims off adapters and controls for double
ligation events) before being mapped to a custom genome (hg38 + SINEUP construct) using STAR v2.7.8a
with parameters as in the ENCODE eCLIP project pipeline (https://www.encodeproject.org/eclip/)
(namely, --outFilterMultimapNmax 1, --outFilterMultimapScoreRange 1, --outFilterScoreMin 10, --

alignEndsType EndToEnd). Reads mapped on the sense strand of the SINEUP construct were removed.
Peaks were called on mapped reads using clipper v0.2.0 (https://github.com/YeoLab/clipper) with custom
data files (gff, exons bed file) containing the SINEUP region, with default parameters. Scripts from the
merge_peaks github repository (https://github.com/YeoLab/merge_peaks) were used to normalize
seCLIP reads over input reads for each replicate (overlap_peakfi_with_bam.pl) before merging input-
normalized peaks into a final set of significant binding regions
(compress_l2foldenrpeakfi_for_replicate_overlapping_bedformat.pl).

Preparation of chemically modified SINEUP-GFP RNA transcribed in vitro

Transcribed SINEUP-GFP RNA was synthesized in vitro as described by Toki *et al.* (2020)[77] by using a
mMESSAGE mMACHINE SP6 Transcription Kit (Thermo Fisher Scientific) and was then modified by
following the protocol by Mandal and Rossi (2013)[78]. For the 20% Ψ modification, UTP was replaced with
20% pseudouridine-5'-triphosphate (TriLink, USA; final concentration, 7.5 mM) by mixing the reagents
with 40 ng/μL of linearized SINEUP RNA. To add a poly-A tail, 1–10 μg of RNA transcribed in vitro was
treated with Escherichia coli poly(A) polymerase (5000 U/mL, New England Biolabs, M0276) at 37°C for
30 min followed by clean-up using an RNeasy Mini kit (Qiagen) and the modified transcripts of SINEUP-
GFP RNA were recovered.

Extraction of modified SINEUP-GFP RNA transcribed in-cell

825   HEK293T/17 cells were plated on a 10-cm dish and SINEUP-GFP plasmids were transfected for 24 h; then
826   2×10^7 cells were harvested by Trizol/chloroform extraction, and total RNAs containing SINEUP-GFP RNAs
827   at the aqueous phase were extracted by using an RNeasy mini Kit (Qiagen). The RNA was then digested
828   with 4 U of TURBO DNase (TURBO DNA-free kit, Invitrogen, AM1907) at 37°C for 90 min followed by DNase
829   inactivation.

Pull-down of SINEUP-GFP RNAs transcribed in-cell

The total RNA volume was adjusted to 100 μL with lysis buffer (50 mM Tris-HCl pH 7.0, 10 mM EDTA, 1%
SDS, 1:100 RNase inhibitor) and mixed with two volumes of hybridization buffer (750 mM NaCl, 50 mM
Tris-HCl pH 7.0, 1 mM EDTA, 1% SDS, 15% formamide, 1:200 RNase inhibitor) with additional 6.6%
formamide. Then, 100 pmol of SINEUP-GFP-specific total probe (2 μL of 50 μM probe) was added to this
mix (probe design as described by Toki *et al.*, 2020)[38], and the reaction was incubated at 37°C overnight
with vertical rotation. Next, 100 μL of MagCapture Tamavidin 2-REV magnetic beads (WAKO) were washed
and resuspended in 100 μL lysis buffer. This bead suspension was added to the completed hybridization
reaction and incubated at 37°C for 30 min with mixing. The solution was then magnetically separated, and
the supernatant was discarded. The beads were washed four times with pre-warmed wash buffer (2×
saline sodium citrate, 0.5% SDS, 1:200 RNase inhibitor), and the supernatant was discarded. After this, the
beads were resuspended in 100 μL ProK buffer (10 mM Tris-HCl pH 7.5, 0.8% SDS, 15 mM EDTA) without
proteinase K and incubated at 65°C for 5 min with end-to-end mixing at 450 rpm. The first pull-down RNA

was then extracted by the Trizol/chloroform method and purified by miRNeasy mini column (Qiagen,
217004). This pull-down and RNA extraction was repeated once again to enrich SINEUP-GFP RNAs.

Library preparation of Oxford Nanopore direct-RNA sequencing

848 Libraries of the direct SINEUP-GFP RNA transcribed in vitro and in-cell were prepared with an SQK-RNA002
849 kit (Oxford Nanopore Technology). The original reverse-transcription adapter (RTA) was used for in vitro
850 samples and four barcoded RTA adapters described by Leger *et al.* (2021)[79] were used for in-cell samples.
851 The libraries were applied to MinION Mk1C sequencer (Oxford Nanopore Technologies) and sequenced
852 for 72 h.

Modification detection using long-read sequencing

Raw fast5 files from direct-RNA sequencing of modified SINEUP RNA transcribed in-cell and in vitro were
base-called using guppy v5.0.7 with RNA-centric options (--trim_strategy rna, --u_substitution true, --
reverse_sequence true). After preparing reference transcriptome annotation files (bed and fasta
annotation files from bedparse gtf2bed and bedtools getfasta, respectively), the base-called data were
mapped to the reference using minimap2 v2.17. To detect modified k-mers in the data we followed the
nanocompore pipeline (https://github.com/tleonardi/nanocompore/). Briefly, this involved indexing the
raw data, followed by running nanopolish eventalign to realign the raw signal-level data to the k-mers of
the reference. After this, the signal was collapsed to the k-mer level using NanopolishComp, and
nanocompore was used to compare the sample transcribed in-cell with the different transcription
conditions in vitro. Results of this comparison were filtered by both log odds ratio (absolute value > 1) and
p value (<0.01). Finally, a peak-calling algorithm
(https://gist.github.com/tleonardi/0bb31e6a380e5766f04f4e197d36b38e) was used to get rid of multiple
adjacent significant sites, because a single modified nucleotide can often cause multiple neighbouring k-
mers to be measured as significant sites, when in reality there is just one modification. The peak-calling
algorithm merges these adjacent sites into a single k-mer, which helps to highlight true-positive sites.

PARIS library preparation

HEK293T/17 cells were transfected with pEGFP-C2 and SINEUP-GFP (SINEB2) or miniSINEUP-GFP FRAM
plasmids in 6-well plates as described above. After 24 h of transfection, cells were gently washed with 1×
PBS and treated with 150 μL 1× PBS (control) or 1 mg/mL Amotosalen (MedChemExpress, HY-107004A)
crosslinking solution added to each well of the 6-well plates, followed by incubation under standard cell-
culture conditions (37°C, 5% CO_2) for 30 min. Next, cells were irradiated for 30 min by 365-nm UV light
while on ice with swirling of the plate every 10 min. Then, the crosslinking solution was removed, and the
cells were scraped off into 1 mL chilled 1× PBS in each well and transferred to a 15-mL tube. Crosslinked
cells were pelleted by centrifuging at 300 × *g* for 5 min at 4°C, and the supernatant was discarded. Next,
for the total nucleic acid (TNA) extraction and PARIS library preparation, the protocol by Zhang *et al.*
(2021)[41] was used with some modifications. Briefly, the cell pellet was lysed by 6 M GuSCN with pipetting
followed by treatment with 0.5 M EDTA and passed through a 25G needle 20 times. After this, the proteins

were digested by proteinase K (final concentration 1 mg/mL) at 37°C for 1 h on a thermomixer with
shaking, and solution was again passed through the 25G needle. After TNA was precipitated by phenol-
isopropanol precipitation as described in Zhang *et al.* (2021)[41], 100 µg of TNA sample was treated with 50
units of TURBO DNase (Invitrogen, AM2239) at 37°C for 20 min. RNA was precipitated using phenol-
isopropanol and further purified using RNA clean and concentrator kit (Zymo Research, R1015). The RNA
quality was measured with Bioanalyzer Nano Kit (Agilent). DNase-treated RNA (10 µg) was fragmented by
using 5 units of ShortCut RNase III (New England Biolabs, M0245) for 5 min at 37°C and the reaction was
stopped by adding 10 µL nuclease free water, 100 µL of RNA binding buffer (Zymo Research, R1015) and
150 µL of 100% ethanol and then purified through a column from the purification kit (Zymo Research,
R1015) followed by quality check with a Bioanalyzer small RNA kit (Agilent).

For 2D gel purification of RNA duplexes, 8% 1-mm thick denatured first-dimension gel was prepared using
the UreaGel System (National Diagnostics, EC-833). A 10-µL sample of fragmented RNA was run on the
first-dimension gel in 0.5x Novex TBE running buffer (Invitrogen, LC6675) at 30 W for 9 min. The gel was
visualized under 300 nm transillumination after staining with SYBR Gold (Invitrogen, S11494) and each
lane was sliced between 50 bp to topside from the first-dimension gel. Next, 16% 1-mm thick denatured
second-dimension gel was prepared using the UreaGel System (National Diagnostics, EC-833). The second-
dimension gel was poured to encapsulate the gel slices from the first-dimension gel, and 0.5x TBE buffer
warmed to 60°C was used to facilitate the denaturing of crosslinked RNA. The gel was run at 30 W for 50
min and then stained with SYBR Gold, and crosslinked RNA was sliced out from the upper diagonal of the
2D gel. The gel slices were cut into smaller pieces and transferred to a 0.5-mL tube with a hole drilled in
the bottom with a 20G needle. This 0.5-mL tube was placed inside a 2-mL tube and centrifuged at 16,000
x *g* for 5 min to crush the gel through the hole. Then 1.5 mL of crushing buffer (20 mM Tris-HCl pH 7.5,
0.25 M sodium acetate, 1 mM EDTA, 0.25% SDS) was added to the crushed gel and rotated at room
temperature overnight. After this, 500 µL of gel slurry was transferred to a Micro Spin column (Cytiva,
27356501) and centrifuged at 3400 x *g* for 1 min or until all gel slurry was filtered. This filtered gel was
then passed through an Amicon Ultra 10K column (Millipore, UFC501096) and further purified using the
Zymo RNA Clean and Concentrator-5 column.

Next, in order to perform proximity ligation, 10 µL of the purified double-stranded RNA was treated with
7.5 unit/µL of T4 RNA Ligase I, high concentration (New England Biolabs, M0437) in a reaction mix
containing 1× T4 RNA Ligase Buffer (New England Biolabs, M0437M), 1U SUPERase•In RNase inhibitor
(Invitrogen, AM2694), and 0.5 mM ATP at room temperature overnight. The proximity-ligated RNA was
purified by using a Zymo RNA clean and concentrator-5 column.

To reverse crosslink the RNA, the samples were mixed with 3 µL of 25 mM acridine orange on ice and
irradiated with 254-nm UV light for 30 min. The reverse crosslinked sample was transferred to a new 1.5-
mL tube and purified using a Zymo RNA Clean and Concentrator-5 column. Further, the reverse crosslinked
RNA was heated at 80°C for 90 s and then snap-cooled on ice. The adapter was ligated by adding 14 µL of
adapter ligation mix (1.5 units/µL T4 RNA ligase I high concentration, 1.5 µM adapter (5' Biotin), 10%
DMSO, 12.5% PEG 8000, 5 mM DTT, 1× T4 RNA Ligase buffer) to 6 µL of reverse crosslinked RNA and
incubating the reaction for 3 h at room temperature. The sequences of all the oligos used in the PARIS
library preparation are listed in **Supplementary Table 6c**. After adapter ligation, free adapters were
removed by adding 3 units of RecJf (New England Biolabs, M0264) and 1.7 units of 5' deadenylase (NEB,
M0331S) and incubating at 37°C for 1 h. The adapter-ligated RNA was purified using a Zymo RNA Clean
and Concentrator-5 column and eluted in 10 µL of nuclease free water. Next, 2 µL of custom reverse

transcription primer (with barcode, 1 µM, sequence in **Supplementary Table 6c**) and 1 µL of 10 mM dNTPs
were added to the purified RNA and the sample was heated at 65°C for 5 min in a thermal cycler followed
by rapid cooling on ice. To this mix, 8 µL reverse transcriptase mix was added, which contained 4 µL of 5x
SSIV buffer (Mn2+) (250 mM Tris-HCl, 375 mM CH3COOK, 7.5mM $MnCl_2$), 10 mM DTT, 1 unit/µL
SUPERaseIn RNase inhibitor (Invitrogen, AM2694), 10 units/µL SuperScript IV (Invitrogen, 18090010). The
reaction was incubated at 25°C for 15 min, 42°C for 10 h; hold at 10°C.

The reverse transcribed RNA was captured on streptavidin-coated magnetic beads. For this purpose, 10
µL of Dynabeads M-270 Streptavidin (Invitrogen, 65305) were washed in 20 µL LiCl buffer (7 M LiCl, 20
mM Tris-HCl (pH7.5), 2 mM EDTA, 0.1% Tween20) and then resuspended in 45 µL of LiCl buffer. This beads
suspension was added to the reverse transcribed RNA and incubated at 37°C for 30 min with pipette
mixing in between. The supernatant was removed by separation on a magnetic stand, and beads were
washed three times with 100 µL TE wash buffer (10 mM Tris-HCl (pH7.5), 1 mM EDTA, 0.1% Tween20) and
finally resuspended in 30 µL elution buffer (1x RNaseONE buffer, 1% Tween20, 10 µM P3-tall primer).
Next, the beads solution was heated at 95°C for 5 min then immediately cooled on ice and magnetically
separated to recover the released cDNA containing supernatant in a new 1.5-mL tube. The beads were
again resuspended in 30 µL elution buffer, separated on a magnet, and supernatant was recovered and
transferred to the released cDNA tube from the previous step. Next, the cDNA was treated with 1 µL of
RNaseONE (Promega, M4261), 1 µL of RNaseH (TaKaRa, 2150A), and 1 µL of RNase cocktail (Invitrogen,
AM2286) for 30 min at 37°C and further purified by using a Zymo DNA Clean and Concentrator column
(Zymo Research, D4033).

After this, 4 µL of circularization reaction mix containing 5 units/µL CircLigase II (Lucigen, CL9021K), 1×
CircLigase II buffer (Lucigen, CL9021K), and 2.5 mM $MnCl_2$ was added to 16 µL of cDNA and incubated at
60°C for 100 min followed by 80°C for 10 min. Circularized cDNA (1 µL) was used to determine the PCR
cycles required for the library amplification. For this, 39 µL of qPCR mix containing 1× Phusion mix (New
England Biolabs, M0531S), 0.125 µM P6 Tall v4 primer, 0.125 µM P3 Tall v4 primer, 0.15 µL ROX, 1/3800
SYBR (Invitrogen, S-7563) was added to 1 µL circularized cDNA and amplified at 95°C for 45 s; 20 cycles of
95°C for 15 s, 65°C for 30 s, 72°C for 45 s on a StepOnePlus Real-Time PCR System (Applied Biosystems,
4376598). The qPCR was stopped just before the amplification plot reached the plateau and the
corresponding cycle number was noted. For the final library amplification, 6 µL of circularized cDNA was
added to 44 µL PCR mix (1× Phusion mix, 0.15 µM P6 Tall v4 primer, 0.15 µM P3 Tall v4 primer) and
amplified at 95°C for 45 s; pre-determined cycles of 95°C for 15 s, 65°C for 30 s, 72°C for 45 s on a routine
thermocycler. The PCR product was purified using Zymo DNA clean and concentrator column (Zymo
Research, D4013) and ran on 6% native TBE gel. DNA from 175 to 300 bp was excised, crushed, and
purified. Library quality was checked by using a Bioanalyzer High-sensitivity DNA chip and concentration
was quantified by using a qPCR KAPA library quantification kit (KAPABiosystems, KK4835). Four barcoded
libraries were pooled together and sequenced on HiSeq2500 platform (150 nt, single-end, rapid mode).

PARIS data analysis

To check the SINEUP regions interacting with rRNAs, the PARIS2 analysis strategy was used as described
by Zhang *et al.* (2021)[41] (https://github.com/minjiezhang-usc/PARIS2). In short, the 3' end adapters were
removed by Trimmomatic v0.39, and the sequencing reads were split according to the sample barcodes
using splitFastq script from icSHAPE pipeline[49]. Next, PCR duplicates were removed using the readCollapse

script from the icSHAPE pipeline[49] followed by removal of the 5' header using Trimmomatic v0.39. The
reads were then mapped to a custom genome generated by adding SINEUP-GFP (SINEB2), miniSINEUP-
GFP (FRAM), and sense EGFP genes as separate chromosomes to the hg38 masked genome[41], which were
previously engineered to contain a single copy of multicopy genes such as rRNA and snRNAs. Mapping
was done by STAR v2.7.10a following the parameters described in the PARIS2 paper[41]. To extract SINEUP-
rRNA interactions, first the reads that were mapped to rRNA were extracted by samtools (as in the PARIS2
analysis strategy) and then rRNA interacting reads in which one of the arms of the RNA-rRNA pair was
mapped to SINEUPs were extracted by using the grep command. The extracted file was sorted by samtools
and visualized on the integrated genome viewer (IGV v2.11.1). VfoldMCPX[80] was used to predict the
structure of the RNA-rRNA pair. Human 18S rRNA structure (ID: URS00005A14E2) and 28S rRNA structure
(ID: URS0000C873C2) were taken from the RNAcentral database (https://rnacentral.org/).

3D structure modelling

SINE RNAs 2D structure information derived from icSHAPE was machine translated into 3D structure
models by RNAComposer[81] (https://rnacomposer.cs.put.poznan.pl/). The 3D models were visualized by
using Mol* tool[82] of RCSB PDB[83] (https://www.rcsb.org/) and analyzed by RNApdbee 2.0[84]
(http://rnapdbee.cs.put.poznan.pl/) to recognize the positions with base-stacking and non-canonical
interactions.

References
988
1.   Diederichs, S. The four dimensions of noncoding RNA conservation. *Trends Genet* **30**, 121-3
(2014).
2.   Novikova, I.V., Hennelly, S.P. & Sanbonmatsu, K.Y. Structural architecture of the human long
non-coding RNA, steroid receptor RNA activator. *Nucleic Acids Res* **40**, 5034-51 (2012).
3.   Somarowthu, S. et al. HOTAIR forms an intricate and modular secondary structure. *Mol Cell* **58**,
353-61 (2015).
4.   Xue, Z. et al. A G-Rich Motif in the lncRNA Braveheart Interacts with a Zinc-Finger Transcription
Factor to Specify the Cardiovascular Lineage. *Mol Cell* **64**, 37-50 (2016).
5.   Guttman, M. & Rinn, J.L. Modular regulatory principles of large non-coding RNAs. *Nature* **482**,
339-46 (2012).
6.   Johnson, R. & Guigo, R. The RIDL hypothesis: transposable elements as functional domains of
long noncoding RNAs. *RNA* **20**, 959-76 (2014).
7.   Wilusz, J.E. et al. A triple helix stabilizes the 3' ends of long noncoding RNAs that lack poly(A)
tails. *Genes Dev* **26**, 2392-407 (2012).
8.   Brown, J.A., Valenstein, M.L., Yario, T.A., Tycowski, K.T. & Steitz, J.A. Formation of triple-helical
structures by the 3'-end sequences of MALAT1 and MENbeta noncoding RNAs. *Proc Natl Acad
Sci U S A* **109**, 19202-7 (2012).
9.   Uroda, T. et al. Conserved Pseudoknots in lncRNA MEG3 Are Essential for Stimulation of the p53
Pathway. *Mol Cell* **75**, 982-995 e9 (2019).
10. Carrieri, C. et al. Long non-coding antisense RNA controls Uchl1 translation through an
embedded SINEB2 repeat. *Nature* **491**, 454-7 (2012).

11.   Podbevsek, P. et al. Structural determinants of the SINE B2 element embedded in the long non-
1011   coding RNA activator of translation AS Uchl1. *Sci Rep* **8**, 3189 (2018).
1012   12.   Ohyama, T. et al. An NMR-based approach reveals the core structure of the functional domain of
1013   SINEUP lncRNAs. *Nucleic Acids Res* **48**, 9346-9360 (2020).
1014   13.   Kramerov, D.A. & Vassetzky, N.S. Short retroposons in eukaryotic genomes. *Int Rev Cytol* **247**,
1015   165-221 (2005).
1016   14.   Ohshima, K. & Okada, N. SINEs and LINEs: symbionts of eukaryotic genomes with a common tail.
1017   *Cytogenet Genome Res* **110**, 475-90 (2005).
1018   15.   Su, M., Han, D., Boyd-Kirkup, J., Yu, X. & Han, J.J. Evolution of Alu elements toward enhancers.
1019   *Cell Rep* **7**, 376-385 (2014).
1020   16.   Bourque, G. et al. Evolution of the mammalian transcription factor binding repertoire via
1021   transposable elements. *Genome Res* **18**, 1752-62 (2008).
1022   17.   Sundaram, V. et al. Widespread contribution of transposable elements to the innovation of gene
1023   regulatory networks. *Genome Res* **24**, 1963-76 (2014).
1024   18.   Roy-Engel, A.M. et al. Human retroelements may introduce intragenic polyadenylation signals.
1025   *Cytogenet Genome Res* **110**, 365-71 (2005).
1026   19.   Policarpi, C. et al. Enhancer SINEs Link Pol III to Pol II Transcription in Neurons. *Cell Rep* **21**, 2879-
1027   2894 (2017).
1028   20.   Zhang, X.O., Gingeras, T.R. & Weng, Z. Genome-wide analysis of polymerase III-transcribed Alu
1029   elements suggests cell-type-specific enhancer function. *Genome Res* **29**, 1402-1414 (2019).
1030   21.   Faulkner, G.J. et al. The regulated retrotransposon transcriptome of mammalian cells. *Nat Genet*
1031   **41**, 563-71 (2009).
1032   22.   Fort, A. et al. Deep transcriptome profiling of mammalian stem cells supports a regulatory role
1033   for retrotransposons in pluripotency maintenance. *Nat Genet* **46**, 558-66 (2014).
1034   23.   Lubelsky, Y. & Ulitsky, I. Sequences enriched in Alu repeats drive nuclear localization of long
1035   RNAs in human cells. *Nature* **555**, 107-111 (2018).
1036   24.   Gong, C. & Maquat, L.E. lncRNAs transactivate STAU1-mediated mRNA decay by duplexing with
1037   3' UTRs via Alu elements. *Nature* **470**, 284-8 (2011).
1038   25.   Mouse Genome Sequencing, C. et al. Initial sequencing and comparative analysis of the mouse
1039   genome. *Nature* **420**, 520-62 (2002).
1040   26.   Jurka, J. et al. Repbase Update, a database of eukaryotic repetitive elements. *Cytogenet Genome*
1041   *Res* **110**, 462-7 (2005).
1042   27.   Bao, W., Kojima, K.K. & Kohany, O. Repbase Update, a database of repetitive elements in
1043   eukaryotic genomes. *Mob DNA* **6**, 11 (2015).
1044   28.   Karijolich, J., Zhao, Y., Alla, R. & Glaunsinger, B. Genome-wide mapping of infection-induced
1045   SINE RNAs reveals a role in selective mRNA export. *Nucleic Acids Res* **45**, 6194-6208 (2017).
1046   29.   Zucchelli, S. et al. SINEUPs are modular antisense long non-coding RNAs that increase synthesis
1047   of target proteins in cells. *Front Cell Neurosci* **9**, 174 (2015).
1048   30.   Takahashi, H. et al. Identification of functional features of synthetic SINEUPs, antisense lncRNAs
1049   that specifically enhance protein translation. *PLoS One* **13**, e0183229 (2018).
1050   31.   Schein, A., Zucchelli, S., Kauppinen, S., Gustincich, S. & Carninci, P. Identification of antisense
1051   long noncoding RNAs that function as SINEUPs in human cells. *Sci Rep* **6**, 33605 (2016).
1052   32.   Sun, L. et al. RNA structure maps across mammalian cellular compartments. *Nat Struct Mol Biol*
1053   **26**, 322-330 (2019).
1054   33.   Kierzek, E. et al. The contribution of pseudouridine to stabilities and structure of RNAs. *Nucleic*
1055   *Acids Res* **42**, 3492-501 (2014).
1056   34.   Schwartz, S. et al. Transcriptome-wide mapping reveals widespread dynamic-regulated
1057   pseudouridylation of ncRNA and mRNA. *Cell* **159**, 148-162 (2014).

35. Lin, T.Y., Mehta, R. & Glatt, S. Pseudouridines in RNAs: switching atoms means shifting
paradigms. *FEBS Lett* **595**, 2310-2322 (2021).
36. Rinn, J.L. et al. Functional demarcation of active and silent chromatin domains in human HOX
loci by noncoding RNAs. *Cell* **129**, 1311-23 (2007).
37. Tsai, M.C. et al. Long noncoding RNA as modular scaffold of histone modification complexes.
Science **329**, 689-93 (2010).
38. Toki, N. et al. SINEUP long non-coding RNA acts via PTBP1 and HNRNPK to promote translational
initiation assemblies. *Nucleic Acids Res* **48**, 11626-11644 (2020).
39. (!!! INVALID CITATION !!! 38).
40. Sharma, E., Sterne-Weiler, T., O'Hanlon, D. & Blencowe, B.J. Global Mapping of Human RNA-RNA
Interactions. *Mol Cell* **62**, 618-26 (2016).
41. Zhang, M. et al. Optimized photochemistry enables efficient analysis of dynamic RNA
structuromes and interactomes in genetic and infectious diseases. *Nat Commun* **12**, 2344
(2021).
42. Meng, Z., Jackson, N.L., Shcherbakov, O.D., Choi, H. & Blume, S.W. The human IGF1R IRES likely
operates through a Shine-Dalgarno-like interaction with the G961 loop (E-site) of the 18S rRNA
and is kinetically modulated by a naturally polymorphic polyU loop. *J Cell Biochem* **110**, 531-44
(2010).
43. Chen, C.K. et al. Structured elements drive extensive circular RNA translation. *Mol Cell* **81**, 4300-
4318 e13 (2021).
44. Reisser, S., Zucchelli, S., Gustincich, S. & Bussi, G. Conformational ensembles of an RNA hairpin
using molecular dynamics and sparse NMR data. *Nucleic Acids Res* **48**, 1164-1174 (2020).
45. Spitale, R.C. et al. Structural imprints in vivo decode RNA regulatory mechanisms. *Nature* **519**,
486-90 (2015).
46. Van Nostrand, E.L. et al. Robust, Cost-Effective Profiling of RNA Binding Protein Targets with
Single-end Enhanced Crosslinking and Immunoprecipitation (seCLIP). *Methods Mol Biol* **1648**,
177-200 (2017).
47. Kalb, V.F., Glasser, S., King, D. & Lingrel, J.B. A cluster of repetitive elements within a 700 base
pair region in the mouse genome. *Nucleic Acids Res* **11**, 2177-84 (1983).
48. Lunyak, V.V. & Atallah, M. Genomic relationship between SINE retrotransposons, Pol III-Pol II
transcription, and chromatin organization: the journey from junk to jewel. *Biochem Cell Biol* **89**,
495-504 (2011).
49. Flynn, R.A. et al. Transcriptome-wide interrogation of RNA secondary structure in living cells
with icSHAPE. *Nat Protoc* **11**, 273-90 (2016).
50. Heyne, S., Will, S., Beckstette, M. & Backofen, R. Lightweight comparison of RNAs based on
exact sequence-structure matches. *Bioinformatics* **25**, 2095-102 (2009).
51. Vanegas, P.L. et al. RNA CoSSMos: Characterization of Secondary Structure Motifs--a searchable
database of secondary structure motifs in RNA three-dimensional structures. *Nucleic Acids Res*
40, D439-44 (2012).
52. Spahn, C.M. et al. Cryo-EM visualization of a viral internal ribosome entry site bound to human
ribosomes: the IRES functions as an RNA-based translation factor. *Cell* **118**, 465-75 (2004).
53. Matsuda, D. & Mauro, V.P. Base pairing between hepatitis C virus RNA and 18S rRNA is required
for IRES-dependent translation initiation in vivo. *Proc Natl Acad Sci U S A* **111**, 15385-9 (2014).
54. Lu, Z., Gong, J. & Zhang, Q.C. PARIS: Psoralen Analysis of RNA Interactions and Structures with
High Throughput and Resolution. *Methods Mol Biol* **1649**, 59-84 (2018).
55. Lu, Z. & Chang, H.Y. The RNA Base-Pairing Problem and Base-Pairing Solutions. *Cold Spring Harb*
Perspect Biol **10**(2018).

56.   Vicens, Q. & Kieft, J.S. Thoughts on how to think (and talk) about RNA structure. *Proc Natl Acad*
1106   *Sci U S A* **119**, e2112677119 (2022).
1107   57.   Vassetzky, N.S. & Kramerov, D.A. SINEBase: a database and tool for SINE analysis. *Nucleic Acids*
1108   *Res* **41**, D83-9 (2013).
1109   58.   Indrieri, A. et al. Synthetic long non-coding RNAs [SINEUPs] rescue defective gene expression in
1110   vivo. *Sci Rep* **6**, 27315 (2016).
1111   59.   Bon, C. et al. SINEUP non-coding RNAs rescue defective frataxin expression and activity in a
1112   cellular model of Friedreich's Ataxia. *Nucleic Acids Res* **47**, 10728-10743 (2019).
1113   60.   Espinoza, S. et al. SINEUP Non-coding RNA Targeting GDNF Rescues Motor Deficits and
1114   Neurodegeneration in a Mouse Model of Parkinson's Disease. *Mol Ther* **28**, 642-652 (2020).
1115   61.   Kramerov, D.A. & Vassetzky, N.S. Origin and evolution of SINEs in eukaryotic genomes. *Heredity*
1116   *(Edinb)* **107**, 487-95 (2011).
1117   62.   Davis, D.R. Stabilization of RNA stacking by pseudouridine. *Nucleic Acids Res* **23**, 5020-6 (1995).
1118   63.   Evans, J.R. et al. Members of the poly (rC) binding protein family stimulate the activity of the c-
1119   myc internal ribosome entry segment in vitro and in vivo. *Oncogene* **22**, 8012-20 (2003).
1120   64.   Bou-Nader, C. & Zhang, J. Structural Insights into RNA Dimerization: Motifs, Interfaces and
1121   Functions. *Molecules* **25**(2020).
1122   65.   Spahn, C.M. et al. Structure of the 80S ribosome from Saccharomyces cerevisiae--tRNA-
1123   ribosome and subunit-subunit interactions. *Cell* **107**, 373-86 (2001).
1124   66.   Simonetti, A., Guca, E., Bochler, A., Kuhn, L. & Hashem, Y. Structural Insights into the
1125   Mammalian Late-Stage Initiation Complexes. *Cell Rep* **31**, 107497 (2020).
1126   67.   Unbehaun, A. et al. Position of eukaryotic initiation factor eIF5B on the 80S ribosome mapped
1127   by directed hydroxyl radical probing. *EMBO J* **26**, 3109-23 (2007).
1128   68.   Armache, J.P. et al. Cryo-EM structure and rRNA model of a translating eukaryotic 80S ribosome
1129   at 5.5-A resolution. *Proc Natl Acad Sci U S A* **107**, 19748-53 (2010).
1130   69.   Heuer, A. et al. Cryo-EM structure of a late pre-40S ribosomal subunit from Saccharomyces
1131   cerevisiae. *Elife* **6**(2017).
1132   70.   Kloth, K. et al. ANK3 related neurodevelopmental disorders: expanding the spectrum of
1133   heterozygous loss-of-function variants. *Neurogenetics* **22**, 263-269 (2021).
1134   71.   Tamura, K. et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood,
1135   evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**, 2731-9 (2011).
1136   72.   Lorenz, R. et al. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011).
1137   73.   Zarringhalam, K., Meyer, M.M., Dotu, I., Chuang, J.H. & Clote, P. Integrating chemical
1138   footprinting data into RNA secondary structure prediction. *PLoS One* **7**, e45160 (2012).
1139   74.   Johnson, P.Z., Kasprzak, W.K., Shapiro, B.A. & Simon, A.E. RNA2Drawer: geometrically strict
1140   drawing of nucleic acid structures with graphical structure editing and highlighting of
1141   complementary subsequences. *RNA Biol* **16**, 1667-1671 (2019).
1142   75.   Pandya-Jones, A. & Black, D.L. Co-transcriptional splicing of constitutive and alternative exons.
1143   *RNA* **15**, 1896-908 (2009).
1144   76.   Brugiolo, M., Botti, V., Liu, N., Muller-McNicoll, M. & Neugebauer, K.M. Fractionation iCLIP
1145   detects persistent SR protein binding to conserved, retained introns in chromatin, nucleoplasm
1146   and cytoplasm. *Nucleic Acids Res* **45**, 10452-10465 (2017).
1147   77.   Toki, N., Takahashi, H., Zucchelli, S., Gustincich, S. & Carninci, P. Synthetic in vitro transcribed
1148   lncRNAs (SINEUPs) with chemical modifications enhance target mRNA translation. *FEBS Lett* **594**,
1149   4357-4369 (2020).
1150   78.   Mandal, P.K. & Rossi, D.J. Reprogramming human fibroblasts to pluripotency using modified
1151   mRNA. *Nat Protoc* **8**, 568-82 (2013).

79. Leger, A. et al. RNA modifications detection by comparative Nanopore direct RNA sequencing.
Nat Commun **12**, 7198 (2021).
80. Zhang, S., Cheng, Y., Guo, P. & Chen, S.J. VfoldMCPX: predicting multistrand RNA complexes.
RNA **28**, 596-608 (2022).
81. Biesiada, M., Purzycka, K.J., Szachniuk, M., Blazewicz, J. & Adamiak, R.W. Automated RNA 3D
Structure Prediction with RNAComposer. *Methods Mol Biol* **1490**, 199-215 (2016).
82. Sehnal, D. et al. Mol* Viewer: modern web app for 3D visualization and analysis of large
biomolecular structures. *Nucleic Acids Res* **49**, W431-W437 (2021).
83. Berman, H.M. et al. The Protein Data Bank. *Nucleic Acids Res* **28**, 235-42 (2000).
84. Zok, T. et al. RNApdbee 2.0: multifunctional tool for RNA structure annotation. *Nucleic Acids Res*
46, W30-W35 (2018).

1163

1164

1165

1166

1167

1168

1169

1170

1171

Figure 1

1172

Figure 1. SINEB2 RNAs from different mouse antisense lncRNAs can act as SINEUP effector domain and upregulate GFP protein expression in HEK293T/17. (**a**) Schematic representation of the sense/antisense EGFP system used to validate the effect of different SINEB2 RNAs on translation. The SINEUP binding domain (BD) is shown in green. The inverted SINEB2 region is the effector domain (ED) shown in blue. Alu repeat and 3' tail of SINEUP are shown in magenta and gray respectively. (**b**) SINEUP construct design to verify independent and combined effects of two SINEB2 elements from the same antisense lncRNA. (**c**) Western blot–based EGFP band intensities normalized to ACTB expression as a measurement of SINEUP-induced protein fold change observed in the HEK293T/17 cell line at 24 h post-transfection. Empty vector, Direct AS Uchl1 SINEB2, and ΔAS Uchl1 SINEB2 (SINEB2 deletion) wereused as controls. (**d**) Western blot images with anti-GFP and anti-ACTB antibodies. ACTB was used as a loading control. Abbreviations: AS, antisense; Direct, direct orientation of SINEB2; Δ, deletion; a and b, two different SINEB2 elements from the same antisense lncRNA; ab and ba, the two SINEB2 elements in original and reversed order, respectively; AS Uchl1 SINEB2–RS, antisense *Uchl1* SINEB2 construct without any restriction enzyme sites around the SINEB2 elements. (**e**) GFP protein fold change in HEK293T cells after co-transfection with sense

EGFP and miniSINEUP (at 24 h post-transfection). Western blot images and corresponding GFP band
intensities (normalized to ACTB expression level) calculated using ImageJ software. An empty vector was
used as the negative control and antisense *Uchl1* SINEB2 containing miniSINEUP (shown as AS *Uchl1*) as
the positive control. Sample labels indicate names of antisense lncRNAs from which the SINEB2 sequences
were isolated, which represent four sub-families (based on RepeatMasker annotation): B3, B3A, Mm2,
and Mm1t. SINEB2 elements are from the B3 sub-family unless specifically stated. AS *Uchl1* + AS *Uxt*-b is
the combination of AS *Uchl1* SINEB2 and the second SINEB2 of AS *Uxt*; 3× indicates three repeats of
SINEB2; and SINEB2/B3 consensus is the B3 sub-family consensus sequence taken from the RepBase
database. N = 3 (for c) and n = 5 (for e); error bars ± standard error mean; **** $P < 0.00005$; *** $P <$
0.0005; ** $P < 0.005$; * $P < 0.05$ by two-tailed Student's *t*-test.

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

Figure 2. In-cell SINE structures and conserved motifs in functionally active SINEUPs. (a) Antisense (AS) *Uchl1* SINEB2 RNA secondary structure based on icSHAPE data (right) is compared with the one previously identified using chemical footprinting[11]. In chemical footprinting-based structure, DNA primer hybridization site is highlighted in gray and indicated by an arrow; light and dark gray circles represent DMS and CMCT reactive bases, respectively; non-reactive bases are outlined by a circle. In icSHAPE-based structure, squares highlight the common sequences and structure motifs in in-cell structures of SINE, with

their corresponding motif number. AGG (blue) and UGG (red) motifs are shown. (**b**–**d**) Location of SINEUP
motifs in icSHAPE-derived structures of functionally active SINE RNAs from (**b**) antisense *Txnip* SINEB2, (**c**)
antisense *Uxt* SINEB2-b and, (**d**) human SINE element FRAM. (**e**) icSHAPE-derived RNA secondary
structures and local SINEUP motifs (colors as in Figure 2a) of functionally dormant wild-type antisense
Gadd45α SINEB2-a and its deletion mutant antisense *Gadd45α* SINEB2-a [Δ27–36] with gained SL1. (**f**)
Effect of antisense *Gadd45α* SINEB2-a deletion mutants on GFP protein expression as measured by
Western blot. An empty vector was used as the negative control. Four mutants with random deletion, as
listed in the Figure, were compared with the full-length construct (Full). N = 3, error bars ± standard
deviation, * $P < 0.05$ by two-tailed Student's *t*-test. AS, antisense; Δ, deletion; SL1, stem-loop 1. Nucleotide
color indicates normalized icSHAPE reactivity score: red, ≥ 0.8; yellow, from above 0.5 to 0.8; white
encircled with grey, ≤ 0.5; bases not encircled, no icSHAPE data available.

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

Figure 3. SINEUP structure dynamics and interaction with HNRNPK in the nucleus and cytoplasm. (**a**) Normalized icSHAPE reactivity score profiles of mouse antisense (AS) *Uchl1* SINEB2 RNA in the nucleus and cytoplasm and (**b**) related 2D structure models created from icSHAPE data. SINEUP motifs from M1 to M7 are indicated by dashed squares. (**c**) Normalized icSHAPE reactivity score profiles of human SINE RNA FRAM in the nucleus and cytoplasm and (**d**) their respective 2D structure models. Different structural regions between the compartments are shaded in blue boxes. (**e**) seCLIP data for HNRNPK binding to SINEUP-GFP RNA (of AS *Uchl1* SINEB2) in the nuclear and cytoplasmic fractions of HEK293T cells. IP: immunoprecipitation against HNRNPK; input: not immunoprecipitated, contains all RNA-protein complexes in the given fraction. Normalized peak regions representing significant enrichment of HNRNPK binding site in IP compared to input are indicated as horizontal bars. The SINEB2 region is highlighted in blue. BD, binding domain; ED, effector domain. Identified HNRNPK binding sites are marked by a red line on 2D structure models of AS *Uchl1* SINEB2 in panel b.

a   SINEB2 interaction site on human 18S rRNA

c   rRNA binding sites on AS *Uchl1* SINEB2

b   SINEB2 interaction sites on human 28S rRNA

1264

Figure 4. Mouse SINEB2 RNA interacts with human rRNA in cell. Antisense (AS) *Uchl1* SINEB2 interacting
regions (Int) in human (**a**) 18S rRNA (Int1) and (**b**) 28S rRNA (Int2 and 3) identified by PARIS2. Different
interaction sites are color-coded, and rRNA-SINEB2 pairs are shown in the corresponding color. Possible

1268   base-pairing based on sequence complementarity of interacting regions is shown. (**c**) SINEB2 regions
1269   interacting with 18S and 28S rRNA regions marked in (a) and (b) on icSHAPE data-driven SINEB2 structure
1270   from whole-cell lysate. SINEUP motifs from M1 to M7 are indicated by dashed squares.

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

Figure 5. In-cell interaction of functional human SINE element FRAM with rRNA. (**a**) Regions of 18S rRNA
found to pair with the SINEUP FRAM region in PARIS2 data. (**b**) Corresponding binding regions to (a) are
highlighted on FRAM RNA. (**c**) Description of FRAM-18S rRNA pairs and interaction sites marked in (a) and

(b). (**d**) 28S rRNA regions identified as FRAM interaction sites by PARIS2 and (**e**) their corresponding binding regions on FRAM. (**f**) Details of FRAM-28S rRNA interactions highlighted in (d) and (e). Different interaction sites (Int) are represented by different colors. Possible base-pairing based on sequence complementarity of RNA-RNA pairs is shown in (c) and (f). In (b) and (e) the icSHAPE-guided FRAM structure from whole cell lysate is shown. Matching regions to SINEUP motifs from M1 to M7 are indicated by dashed squares.

1331

1332

1333 **Figure 6. Machine-translated 3D RNA structure models based on in vivo SINE RNA 2D structures.** 3D
1334 structure models for (**a**–**d**) mouse antisense (AS) *Uchl1* SINEB2 RNA in the nucleus (**a**,**b**) and cytoplasm
1335 (**c**,**d**) and for (**e**–**h**) FRAM RNA in the nucleus (**e**,**f**) and cytoplasm (**g**,**h**). SINEUP structure motifs M1–M7
1336 are shown. Red frames show enlarged views of the SL1 region. Panels b, d, f, and h show the molecular
1337 surface view of the 3D models.

1338

1339

1340

1341

1342

1343

1344

1345

Extended Data Figure 1. Effect of mouse natural antisense *Gadd45α* lncRNA on sense *Gadd45α* protein expression. (**a**) Molecular features of mouse sense/antisense *Gadd45α* RNAs. Antisense (AS) *Gadd45α* contains a sequence complementary to intron 2, exon 3, and intron 3 of sense Gadd45α mRNA,two inverted SINEB2s, three LINE, and one SINE-Alu elements. (**b**) Western blot analysis of sense Gadd45α protein expression in HEK293T/17 cells co-transfected with sense/antisense *Gadd45α* for 24 h. Empty vector, negative control; plasmid transfection at + 0.7 pmol; ++ 0.8 pmol; +++ 0.9 pmol; ++++ 1.0 pmol; - not transfected. ACTB is the loading control.

1353

1354

1355

1356

1357

a

b

1358

Extended Data Figure 2. qRT-PCR results for GFP mRNA (black bars) and SINEUP-GFP RNA (grey bars) expression. Expression values are normalized to human GAPDH mRNA. Data were analyzed by using the $\Delta\Delta C_T$ method. RNA was extracted from (**a**) long SINEUPs (see description of terms in Figure 1c), with ΔAS *Uchl1* SINEB2 = 1 and from (**b**) MiniSINEUP-GFP (described in Figure 2b), with AS *Uchl1* = 1. Error bars are ± standard error mean for three biological replicates. AS, antisense.

1364

1365

1366

1367

1368

Extended Data Figure 3

1369

Extended Data Figure 3. Multiple sequence alignment and phylogenetic analysis of SINEB2 RNA
sequences. (a) ClustalW sequence alignment of mouse SINEB2 RNA sequences used as effector domain
in long and mini SINEUPs. Gaps in the alignment are indicated by -. Bases are highlighted in four different

colors. Black squares denote A and B boxes. Alignment is showing the sequence variation within members
of the same family of SINE. (**b**) SINEB2 sequence alignment–based molecular phylogenetic analysis by the
Maximum Likelihood method (Tamura 3-parameter model) to infer the evolutionary history. The tree with
the highest log likelihood (−2415.7806) is shown. The tree is drawn to scale, with branch lengths measured
in the number of substitutions per site. AS, antisense; B3 SINE2 Rodentia RC, the consensus sequence for
SINEB2 subfamily B3 from the Repbase database.

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

Extended Data Figure 4. icSHAPE 2D structure models of mouse SINEB2 RNAs from natural antisense
transcripts that are tested as effector domain in SINEUPs. Regions that match the SINEUP motifs are
marked by squares with their corresponding motif number. Blue square, AGG type motif; red square, UGG
type motif. FC, GFP protein fold-change induced by the respective SINEB2. Nucleotide color indicates
normalized icSHAPE reactivity score: red, ≥ 0.8; yellow, from above 0.5 to 0.8; white encircled with grey,
≤ 0.5; bases not encircled, no icSHAPE data available. AS, antisense.

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

a

S. No.	SINEUP Motif	CoSSMos motif match	PDB ID	Positions	Molecule
1.	Motif 6	GGUGG UCCC	5A2Q	G1324-G1325-U1326-G1327-G1328 U1504-C1503-C1502-C1501	18S rRNA
2.	Motif 2	GGAGG CCCC	5JB3	G999-G1000-A1001-G1002-G1003 C1170-C1169-C1168-C1167	16S rRNA

b

1434

Extended Data Figure 5. Similarity of SINEUP motifs to rRNA 3D structure motifs. (**a**) Hits found in the Characterization of Secondary Structure Motifs database for 3D motifs that match the SINEUP structure motifs. (**b**) The region similar to SINEUP motif 6 in helix 34 of human 18S rRNA is framed in black and marked with an arrow.

1439

Extended Data Figure 6. SINEUP structure motifs in functionally inactive antisense (AS) *Gadd45α* **SINEB2-a deletion mutants.** icSHAPE derived structures of AS *Gadd45α* SINEB2-a deletion mutants [Δ40–50], [Δ66–76], and [Δ92–101]. Regions that match conserved SINEUP structure motifs are marked by different colors and numbers. Δ = deletion. Nucleotide color indicates normalized icSHAPE reactivity score: red, ≥ 0.8; yellow, from above 0.5 to 0.8; white encircled with grey, ≤ 0.5; bases not encircled, no icSHAPE data available. AS, antisense.

Extended Data Figure 7

Extended Data Figure 7. Discovery of modified bases in SINEB2 RNA. (**a**) An overview of SINEUP RNA modification analysis. SINEUP-GFP RNA (containing antisense Uchl1 SINEB2) transcribed in vitro was modified with 20% pseudouridine (Ψ), and a non-modified (0% Ψ) sample was used as control. In parallel, SINEUP-GFP and sense EGFP plasmids were co-transfected in HEK293T/17 cells, then SINEUP RNA that was transcribed in-cell was captured and enriched using specific biotinylated oligo probes and pull-down (twice) on streptavidin-coated magnetic beads. Purified samples of SINEUP-GFP RNA transcribed in-cell or transcribed and modified in vitro were sequenced by Nanopore direct RNA sequencing. Profiles of non-modified (0% Ψ) and modified SINEUP-GFP RNA transcribed in vitro were compared with that transcribed in-cell, and k-mers corresponding to the modified positions were identified in the in-cell transcripts. The red peaks in the graph mark the position of modified k-mers on the SINEUP transcript. The SINEB2 region is shaded in blue, and verified Ψ sites are marked. BD, binding domain. (**b**) Modified k-mers overlaid on 2D structure of antisense *Uchl1* SINEB2. k-mers are highlighted in yellow, Ψ sites are in red. SINEUP motifs are marked in black dashed squares with motifs 1–7 labelled as M1–M7.

1471

Extended Data Figure 8. SINEUP-rRNA duplex reads in PARIS sequencing data. (**a**) SINEUP-GFP (of
antisense *Uchl1* SINEB2) RNA duplex reads with 18S, 5.8S, and 28S rRNA. Two biological replicates of the
PARIS experiment are shown as separate tracks. SINEB2-rRNA interactions shown in Fig. 4 are indicated
by respective colored arcs. (**b**) miniSINEUP-GFP FRAM RNA-rRNA duplex reads. PARIS rep1 and rep 2 are
two biological replicates. The FRAM RNA-rRNA interactions displayed in Fig. 5 are marked by arcs in their
respective colors.

1478

1479

1480

1481

1482

1483

Extended Data Figure 9. SINE RNA predicted 3D structure models based on in-cell 2D structures of mouse SINEB2 RNAs. SINEUP structure motifs are denoted from M1 to M7. AS: antisense.

1488

Extended Data Figure 10. Comparison of icSHAPE data-driven predicted 3D structure models of wild-
type AS *Gadd45α* SINEB2-a and its deletion mutants with functional AS *Uchl1* SINEB2. (**a**) AS *Uchl1*
SINEB2 3D model predicted from whole cell icSHAPE data. Predicted 3D models with enlarged view of the
corresponding SL1 regions for AS *Gadd45α* SINEB2-a (**b**) wild-type (functionally inactive), its (c)
functionally active deletion mutant [Δ27–36], and non-functional mutants (d) [Δ40–50], (e) [Δ66–76], and
(f) [Δ92–101]. SINEUP structure motifs M1–M7 are marked. Red frame shows zoomed-in view of the
region corresponding to SL1. The molecular surface views of the 3D models are shaded in gray. Δ =
deletion.

1497

1498 **Supplementary Table 1:** SINEB2 sequences tested as effector domain (ED) in (**a**) long SINEUP and (**b**) in
1499 miniSINEUP-GFP.

1500

1501 **Supplementary Table 2:** List of base-stacking positions in SINE RNA 3D models.

1502

1503 **Supplementary Table 3:** List of non-canonical interactions identified in SINE RNA 3D models.

1504

1505 **Supplementary Table 4:** Intermolecular interactions of SINEUPs with RNAs other than rRNA in PARIS2
1506 data.

1507

1508 **Supplementary Table 5:** (**a**) Primers used for QuikChange II site-directed mutagenesis. (**b**) Primers used
1509 for PCR-cloning of miniSINEUP-GFP constructs.

1510

1511 **Supplementary Table 6:** Oligos used in (**a**) icSHAPE, (**b**) seCLIP, and (**c**) PARIS library preparation.

1512

1513

1514

1515