Toxins in plastic: Evidence for the cardiodepressive effects of di-2-ethylhexylphthalate (DEHP)

*Luther M. Swift[1,2], *Anysja Roberts[1,2], Jenna Pressman[1-3], Devon Guerrelli[1-3], Samuel Allen[1,2], Kazi T. Haq[1,2], Julie A Reisz[4], Angelo D'Alessandro[4], Nikki Gillum Posnack[1,2,5]

* Authors contributed equally

[1] Sheikh Zayed Institute for Pediatric Surgical Innovation, Children's National Health System, Washington DC 20010

[2] Children's National Heart Institute, Children's National Hospital, Washington DC 20010

[3] Department of Biomedical Engineering, School of Engineering and Applied Sciences: George Washington University, Washington DC 20037

[4] Department of Biochemistry and Molecular Genetics, University of Colorado, Anschutz Medical Campus, Aurora, CO

[5] Department of Pediatrics, Department of Pharmacology & Physiology, School of Medicine and Health Sciences: George Washington University, Washington DC 20037

Corresponding author:
Nikki Gillum Posnack, Ph.D.
Sheikh Zayed Institute, 6th floor, M7707
111 Michigan Avenue, NW
Washington, DC, USA 20010
Tel: (202) 476-2475
Email: nposnack@childrensnational.org

Conflicts of interest: The authors declare they have nothing to disclose.

ABSTRACT

Background: Di-2-ethylhexylphthalate (DEHP) is commonly used in the manufacturing of plastic materials, including intravenous bags, blood storage bags, and medical-grade tubing. Prior studies demonstrated that DEHP can leach from plastic medical products, which can result in inadvertent patient exposure. Further, in vitro studies suggest that DEHP may act as a cardiodepressive agent by slowing the beating rate of isolated cardiomyocytes.

Objective: In this study, we investigated the direct effects of acute DEHP exposure on cardiac electrophysiology.

Methods: DEHP concentrations were measured in red blood cell (RBC) units stored between 7-42 days (23-119 µg/mL). Using these concentrations as a guide, Langendorff-perfused heart preparations were exposed to DEHP (15-90 minutes) and the impact on cardiac electrophysiology metrics were quantified. Secondary studies used human induced pluripotent stem cell-derived cardiomyocytes (hiPSC-CM) to measure the effect of DEHP exposure on conduction velocity over a prolonged period of time (15-180 minutes).

Results: In intact rat heart preparations, sinus activity remained stable following acute exposure to lower doses of DEHP (25-50 µg/mL), but sinus rate declined by 43% and sinus node recovery time prolonged by 56.5% following 30-minute exposure 100 µg/ml DEHP. DEHP exposure also exerted a negative dromotropic response, as indicated by a 69.4% longer PR interval, 108.5% longer Wenckebach cycle length, and increased incidence of atrioventricular uncoupling. Pretreatment with doxycycline, a matrix metalloproteinase inhibitor, partially rescued the effects of DEHP on sinus activity, but did not ameliorate the effects on atrioventricular conduction. DEHP exposure prolonged the ventricular action potential and effective refractory period but had no measurable effect on intracellular calcium transient duration. Follow-up studies using hiPSC-CM demonstrated that DEHP slows electrical conduction in a time (15 min – 3 hours) and dose-dependent manner (10-100 µg/mL).

Discussion: DEHP exposure perturbs cardiac electrophysiology in a dose- and time-dependent manner. Future studies are warranted to investigate the impact of DEHP exposures on human health, with special consideration for clinical procedures that employ plastic materials.

INTRODUCTION

Phthalate plasticizer chemicals are widely used in the manufacturing of both consumer goods and medical products, including endotracheal tubes, intravenous bags, blood storage bags, and medical-grade tubing[1,2]. Di(2-ethylhexyl) phthalate (DEHP) is frequently used as a plasticizer additive, wherein it embeds between polyvinyl chloride polymers to soften and impart flexibility to plastic materials[3,4]. DEHP is highly susceptible to leaching because it is not covalently bound to the polyvinyl chloride matrix; phthalate leaching is increased when lipophilic solutions come into contact with plastic materials that have a large surface area, at an elevated temperature, or for a prolonged period of time[5–10]. Historically, DEHP leaching has been considered beneficial in the blood banking field, since DEHP acts as a preservative agent to reduce red blood cell (RBC) hemolysis and extend the shelf life of donated blood products[11–14]. Although, several studies have shown that DEHP can accumulate in stored blood products, which can result in inadvertent patient exposure[9,15–17]. Clinical exposure to phthalate chemicals has also been documented in patients undergoing multiple medical interventions that employ plastic products[18–21]. Accordingly, safety concerns have been raised regarding phthalate exposure and the potential negative implications on human health[22–24].

Epidemiological studies have reported associations between phthalate exposure and an increased risk of all-cause and cardiovascular mortality, which has prompted a renewed interest in understanding the direct effect of DEHP on the heart[25,26]. Identifying the consequences of clinical phthalate exposure on patient outcomes is inherently difficult, given the complexity of clinical care and the heterogenous nature of patient populations. Nevertheless, experimental studies have shown that phthalate chemicals can exert cardiotoxic effects[27–29]. As examples, DEHP exposure slows the spontaneous beating rate of isolated cardiomyocyte preparations[30,31], while MEHP exposure (primary metabolite of DEHP) delays atrioventricular conduction and precipitates cardiac arrest[29,32]. Although the underlying mechanisms are incompletely understood, studies suggest that phthalates may exert immediate effects via interaction with cholinergic receptors[33,34], while longer exposure times can disrupt intercellular coupling (likely via oxidative stress and/or increased matrix metalloproteinase activity)[28,35,36]. For the latter, comparative studies conducted in liver cells estimated that DEHP reduces gap junction intercellular coupling by approximately 20% within 1 hour and >80% within 16 hours of exposure[37].

In the current experimental study, we aimed to further understand the potential risks of clinical phthalate exposure on cardiac physiology. Accordingly, we first measured DEHP concentrations in stored RBC units and then used those concentrations as a guide for cardiotoxicity assessment. Specifically, we quantified the acute effects of DEHP exposure on cardiac electrophysiology using intact rodent heart

preparations (15-90 minutes, 25-100 µg/mL) and human induced pluripotent-stem cell derived

cardiomyocytes (15-180 minutes, 10-100 µg/mL). Further, we investigated whether pretreatment with a

cholinergic receptor antagonist (atropine) or matrix metalloproteinase inhibitor (doxycycline) could

abrogate the cardiotoxic effects of DEHP.

METHODS

<u>Animal model</u>

Animal protocols were approved by the Institutional Animal Care and Use Committee within the

Children's National Research Institute and followed the Nationals Institutes of Health's *Guide for the*

Care and Use of Laboratory Animals. Experiments were performed using 2–4-month-old, mixed sex

Sprague Dawley rats (strain NTac: SD, from NIH Genetic Resource stock, Taconic Biosciences:

Germantown New York). Animals were housed in conventional acrylic cages in the Research Animal

Facility, under standard environmental conditions (12:12 hours light:dark cycle, 18-25º C, 30-70%

humidity).

<u>Isolated heart preparation and Langendorff perfusion</u>

Animals were anesthetized with isoflurane (2-4%) for approximately 10 minutes until a surgical depth of

anesthesia was reached, as determined by a tail pinch test. Animals were then euthanized by

exsanguination following heart excision. Briefly, the heart was excised and placed in ice-cold

cardioplegia; thereafter, the aorta was cannulated and the intact heart was transferred to temperature

controlled (37°C), constant pressure (70 mm/Hg) Langendorff-perfusion system. The heart was

retrograde perfused with a modified Krebs-Henseleit buffer (mM: 118.0 NaCl, 3.3 KCl, 1.2 $MgSO_4$, 1.2

KH_2PO_4, 24.0 $NaHCO_3$, 10.0 glucose, 2.0 sodium pyruvate, 10.0 HEPES buffer, 2.0 $CaCl_2$) bubbled

with carbogen (95% O_2, 5% CO_2). Intact heart preparations were allowed to equilibrate for 10-15

minutes on the perfusion system; once the myocardium was stabilized, blebbistatin was added to the

perfusate to reduce metabolic demand and suppress contractile motion[38,39]. A low dose of blebbistatin

(1.25 µg/mL) was used for electrophysiology studies, while a slightly higher dose of blebbistatin (3.75

µg/ml) was used for imaging studies[40]. To take into account inter-animal variability, each animal served

as its own control with baseline electrophysiology recordings collected before treatment. Time control

experiments were also performed to account for any slight deviation in myocardial physiology that might

occur during prolonged Langendorff-perfusion. DEHP experiments were conducted by diluting the neat

stock liquid into 0.1% DMSO (to aid in solubility), which was diluted in Krebs-Henseleit buffer to reach a

final circulating concentration of 25-100 µg/mL. This concentration was selected to mimic a clinical

exposure to DEHP following contact with plastic medical products, similar to exposure via blood

transfusion[41,42]. In a subset of studies, intact heart preparations were pretreated with either 1 mM

atropine (muscarinic antagonist) or 20 mM doxycycline (matrix metalloproteinase inhibitor).

Electrophysiology measurements

Ex vivo electrocardiograms (ECG) were recorded continuously from excised, intact heart preparations

by placing needle electrodes within the perfusate bath. ECGs were recorded using a PowerLab

acquisition system, in conjunction with a differential amplifier (Warner Instruments, Holliston MA) and a

60 Hz noise eliminator (Hum Bug: Digitimer, Fort Lauderdale FL). From each ECG recording, the heart

rate, P-wave duration, and PR interval were measured during sinus rhythm. Coronary flow rate was

monitored continuously using an inline flow sensor (Transonic, Ithaca NY) positioned above the aortic

cannula. In a subset of studies, microelectrode arrays (MEA, MappingLabs Ltd., United Kingdom) were

positioned in two configurations to measure activation time: one MEA on the right atrium and the

second MEA positioned on either the left atrium (to measure atrial conduction) or the left ventricle (to

measure atrioventricular conduction). Electrical signals were acquired with a multi-channel amplifier

using eMapRecord software and analyzed using EMapScope (MappingLabs Ltd.).

Programmed electrical stimulation was implemented by placing two small stimulation electrodes

(Harvard Biosciences, Holliston MA) on the right atrium and left ventricle. An electrophysiology

stimulator (Bloom: Fisher Medical, Wheat Ridge CO) was set to 1 ms pulse width at 1.5x the minimum

threshold current (~0.7-1.2 mA)[43]. To assess sinus node recovery time (SNRT), spontaneous sinus

activity was suppressed via fast atrial pacing (20 beats, S1-S1) and SNRT was measured as the time

delay between the last paced beat and the first spontaneous P-wave. Corrected SNRT (cSNRT) was

measured as the difference between SNRT and the RR-interval during sinus rhythm. To identify the

Wenckebach cycle length (WBCL), atrial pacing was initiated (S1-S1) and the pacing cycle length

(PCL) was decremented by 10 ms and then 1 ms increments until atrioventricular (AV) conduction

failed. WBCL was defined as the shortest S1-S1 interval that resulted in 1:1 AV conduction, and 2:1

capture was defined as the longest S1-S1 interval that resulted in 2:1 capture of AV conduction. The

AV node effective refractory period (AVNERP) was determined by introducing an extrastimulus (atria

pacing) and pinpointing the shortest S1-S2 interval that resulted in 1:1 AV conduction. The ventricular

effective refractory period (VERP) was determined by decrementing the S1-S2 interval (ventricular

pacing) to identify the shortest S1-S2 interval that resulted in ventricular depolarization.

Fluorescence imaging and analysis

Isolated hearts were loaded with a calcium indicator dye (100 µg Rhod-2, AM; AAT Bioquest,

Sunnyvale CA) and/or a potentiometric dye (62 µg RH237; AAT Bioquest). Fluorescence signals were

acquired by illuminating the epicardial surface with an LED light with excitation filter (535 ± 25 nm;

ThorLabs, Sterling VA); emission filters were used to collect the calcium signal (585 ± 20 nm) or voltage

signal (>710 nm)[44]. Images were collected using high-speed cameras at 400-600 fps (Dimax CS4:

PCO-Tech, Kelheim Germany; Optical Mapping System from MappingLabs Ltd equipped with Prime

BSI cameras, Teledyne Photometrix). Optical action potentials and calcium transients were measured

at the apex of the left ventricle; fluorescence signals were analyzed using custom software (*CaMat*[45],

Kairosight[46]).

<u>Human induced pluripotent stem cell derived-cardiomyocyte (hiPSC-CM) recordings</u>

Cryopreserved hiPSC-CMs (iCell cardiomyocytes[2], female donor #01434, Fujifilm) were thawed and

plated onto fibronectin-coated microelectrode arrays (MEA) at a density of 40-65,000 cells/well (Axion

BioSystems, Atlanta GA). Cells were defrosted in iCell cardiomyocyte plating media in a cell culture

incubator (37ºC, 5% CO_2) for 2 hours, thereafter cells were cultured in iCell maintenance media

(Fujifilm). hiPSC-CM formed a confluent monolayer within three days after plating and were used

thereafter for conduction velocity measurements. Briefly, hiPSC-CM were equilibrated in a temperature

(37°C) and gas controlled (5% O_2) live cell monitoring system (Maestro Edge, Axion) for 20-minutes.

Electrical stimulation was applied (1.5 Hz frequency, 800 mV, 400 µs duration) and conduction velocity

was measured across a 16-electrode array under baseline conditions and again after chemical

exposure (0.1% DMSO, 10-100 µg/mL DEHP). In a subset of studies, hiPSC-CM were pretreated with

either 1 µM atropine (antimuscarinic) or 20 µM doxycycline (matrix metalloproteinase inhibitor) before

DEHP exposure.

<u>Phthalate Chemical Extraction and Quantitation</u>

Phthalate chemical concentrations were measured in supernatant samples collected from refrigerated

RBC units, stored for 7-42 days. Briefly, samples were shipped on dry ice to the University of Colorado

School of Medicine Metabolomics Core. Samples were thawed on ice and phthalates were extracted

from a 10 µL aliquot via vigorous vortexing for 30 min at 4°C in the presence of 5:3:2

methanol:acetonitrile:water (240 µL) containing an isotope labeled standard (1 µM DEHP ring-1,2-$^{13}C_2$,

Cambridge Isotope Laboratories). Extraction supernatants were clarified via centrifugation at 18,000

rpm for 10 min at 4°C and analyzed immediately by ultra high-pressure liquid chromatography coupled

to mass spectrometry (UHPLC-MS). Sample injection volume was 20 µL. Samples were analyzed on a

Thermo Vanquish UHPLC coupled to a Thermo Q Exactive high resolution mass spectrometer. The LC

was equipped with a Phenomenex Kinetex C18 column (2.1 x 150 mm, 1.7 µm) held at 45 °C (gradient

information below). The mass spectrometer scanned in MS1 mode in the range of 65 to 975 m/z at a

resolution of 70,000. Instrument-generated .raw files were converted to .mzXML format via

RawConverter and peaks for phthalates and accompanying standards were extracted and integrated

using Maven (Princeton University). Absolute concentrations were obtained using the following

equation where dilution factor (DF) is 25: [Phthalate] = (Peak Area Phthalate) / (Peak Area Standard) *

[Standard] * DF

DEHP was quantified using a 5 min C18 gradient at a flow rate of 450 µL per min as previously

described[47] with a single change – eluant was introduced to the MS using atmospheric pressure

chemical ionization (APCI) in positive mode. APCI was performed using 10 sheath gas, 20 aux gas

(both N_2), and 350°C vaporizer temperature. The calculations of DEHP concentrations utilized the

average peak area of the stable isotope labeled DEHP standard across all samples within the biological

group.

Statistical analysis

Results are reported as the mean ± standard deviation. Normality was assessed for each data set

(Shapiro-Wilk test). Two group comparisons (e.g., baseline vs 60-minute control) were analyzed using

a two-tailed T-test. Three or more group comparisons (e.g., dose and time-course studies) were

analyzed using a one-way ANOVA test with Dunnett's (parametric data sets) or Dunn's (nonparametric)

multiple comparisons test. Comparisons with two independent factors (e.g., APD measurements across

multiple pacing frequencies) were analyzed using two-way ANOVA with Holm-Sidak test for multiple

comparisons. Significance is denoted in each figure with an asterisk (p-value <0.05).

220
221

RESULTS

DEHP exerts a negative chronotropic effect in a dose-dependent manner

Phthalate plasticizers have been shown to leach from medical plastic devices, resulting in unintentional

patient exposure. Prior studies demonstrated that DEHP can accumulate in stored blood or migrate

from polyvinyl chloride tubing, reaching concentrations as high as 400 µg/mL[7,8,21,41]. In agreement, we

found that DEHP concentrations increased in RBC units during refrigerated storage, beginning at 23-34

µg/mL in fresh units (7 days storage) and increasing to 42-119 µg/mL in units near expiry (35-42 days;

p<0.05; **Figure 1**). Using these concentrations as a guide, we next performed dose-response studies to

measure the direct effect of DEHP exposure on cardiac function using Langendorff-perfused heart

preparations (**Figure 2**). For these initial studies, we focused on cardiac automaticity as a primary
endpoint, since prior work has shown that DEHP slows the beating rate of isolated cardiomyocytes[30,31].
Sinus rate remained stable under control conditions, declining by only 3.5% during prolonged time
course studies (baseline: 322±43, time control: 311±38 BPM, p=0.6, **Figure 2A**). Sinus rate also
remained stable following an acute 30-minute exposure to lower doses of DEHP (baseline: 292±26; 25
µg/mL DEHP: 299±41; 50 mg/mL DEHP: 272±52 BPM; **Figure 2A**). However, a negative chronotropic
effect was observed soon after increasing the circulating concentration to 100 µg/mL DEHP, which
decreased the sinus rate by 43% within 30-minutes (baseline: 292±26, 100 µg/mL DEHP: 167±66 BPM,
p<0.001). Only minor variations in sinus rhythm were observed under control conditions (baseline
RMSSD: 1.13±0.75, time control: 1.86±1.31 ms, p=0.1) and following acute exposure to a low
concentration of DEHP (baseline RMSSD: 1.12±0.64; 25 µg/mL DEHP: 2.91±3.0 ms; **Figure 2A**). While
more extreme fluctuations and abnormal sinus rhythms were observed after increasing the circulating
concentration to 50 µg/mL (RMSSD: 8.3±9.7 ms, p<0.01) or 100 µg/mL DEHP (RMSSD: 22.6±22.5 ms,
p<0.001).

We also quantified sinus node recovery time (SNRT), a measurement of automaticity that is evaluated
using overdrive pacing. SNRT remained stable under control conditions (baseline: 229±28; time control:
235±28 ms) and following an acute 30-minute exposure to lower doses of DEHP (baseline: 244±21; 25
µg/mL DEHP: 242±33; 50 µg/mL DEHP: 280±66 ms; **Figure 2B**). While acute exposure to 100 µg/mL
DEHP significantly impaired sinus function, as demonstrated by a longer SNRT (baseline: 244±21; 100
µg/mL DEHP: 382±117 ms, p<0.05). Similar results were obtained when SNRT measurements were
corrected for heart rate, wherein the baseline measurement (cSNRT: 39.6±8.2) was significantly shorter
than DEHP-exposed hearts (50 µg/mL DEHP: 53.1±16.2; 100 µg/mL DEHP: 60.0±9.4 ms, p<0.05;
Figure 2B). Further, in one-quarter of the high dose studies (2 of 8 experiments), SNRT could not be
determined as atrial pacing either failed to elicit a ventricular response or sinus activity was completely
absent as discerned by the lack of a p-wave.

<u>DEHP exerts a negative chronotropic effect in a time-dependent manner</u>
Prolonged exposure to phthalates can occur in the clinical setting over multiple hours, as is the case in
patients receiving a blood transfusion, exchange transfusion, circulatory support (cardiopulmonary
bypass, extracorporeal membrane oxygenation), or those receiving intensive care[2]. Accordingly, we
investigated the time-dependent effects of 100 µg/mL DEHP exposure on cardiac electrophysiology. In
this subset of studies, we observed a progressive decline in sinus rate by 27.3% at 60 minutes
(baseline: 326.3±40.8, DEHP 60 min: 237.1±47.7 BPM, p<0.0001) and 46.7% at 90 minutes (DEHP 90

min: 173.6±99.3 BPM, p<0.0001; **Figure 3A,B**). Atrioventricular uncoupling was more commonly

observed with a longer exposure time, which resulted in divergent beating rates between the atria and

ventricles. As such, the ventricular rate declined, on average, by 55.6% at 90 minutes exposure

(baseline: 326.3±40.8, DEHP 90 min: 144.8±89.5 BPM, p<0.0001; **Figure 3B**). Prolonged exposure

time also impacted the ventricular rhythm, as denoted by a marked increase in heart rate variability

(baseline RMSSD: 0.86±0.2, DEHP 60 min: 17.7±14.8, DEHP 90 min: 83.6±71.3 ms, p<0.0001; **Figure**

3C). SNRT was also delayed with a longer exposure time, increasing by 52% at 60 minutes and 111%

at 90 minutes (baseline: 233.1±28.9, DEHP 60 min: 353.6±147.1, DEHP 90 min: 492.6±329.6 ms,

p<0.001; **Figure 3D**). A similar delay was observed when SNRT was corrected for sinus rate (cSNRT),

with recovery time prolonged by 103% at 60 minutes and 387% at 90 minutes (baseline: 36.6±9.6,

DEHP 60 min: 75.8±59.6, DEHP 90 min: 178.2±228.4 ms, p<0.01). SNRT and cSNRT became more

difficult to measure with prolonged exposure, as atrial pacing either failed to elicit a ventricular response

or sinus activity did not recover after pacing. Finally, it was observed that coronary flow dropped

precipitously during the course of each DEHP exposure study, decreasing by 56% at 60 minutes and

66% at 90 minutes (baseline: 5.7±1.3, DEHP 60 min: 2.5±1.7, DEHP 90 min: 1.9±1.7 ml/min, p<0.0001;

Figure 3E).

<u>DEHP exposure delays inter-atrial and atrioventricular (AV) conduction</u>

We previously reported that DEHP exposure slows conduction velocity across cardiomyocyte

monolayers[28]. Herein, we investigated the impact of DEHP on electrical conduction in intact heart

preparations, a more complex model that preserves the cardiac conduction system and three-

dimensional cell-cell interactions. Using MEAs positioned on the surface of the heart, we measured a

three-fold increase in inter-atrial activation time (baseline: 7.2±3.7, 100 μg/mL DEHP 60 min: 21.3±4.6

ms, p<0.05), and a two-fold increase in AV activation time (baseline: 37.6±9.6, 100 μg/mL DEHP 60

min: 85.2±35.7 ms; p<0.005; **Figure 4B-C**). Delayed AV conduction was readily observed using optical

mapping techniques (**Figure 4A**); in the presented example, atrial activation precedes ventricular

activation by ~40 ms at baseline and ~80 ms after DEHP exposure (100 μg/mL, 60 minutes).

AV conduction slowing was also observed in ECG recordings during sinus rhythm **(Figure 5A)**. The PR

interval lengthened by 23.7% after 15-minutes exposure to DEHP (baseline: 39.3±7.5, 100 μg/mL

DEHP: 48.6±15.9 ms, p<0.05) and 69.4% after 60-minutes DEHP exposure (100 μg/mL DEHP:

66.6±19.3 ms, p<0.0001). With prolonged DEHP exposure, PR interval measurements became

inconsistent, as AV conduction delays progressed to uncoupling and/or heart block in 64% of heart

preparations at 60-minutes **(Figure 5A).** Accordingly, PR interval measurements were only reported for
coupled beats.

Programmed electrical stimulation was used to further interrogate the effects of DEHP on AV nodal
function **(Figure 5B-D)**. Wenckebach cycle length (WBCL) was identified by dynamically pacing the
right atrium at an increasing rate, until ventricular conduction failed. Following DEHP exposure, AV
conduction failure occurred more readily at slower pacing frequencies (longer pacing cycle lengths), as
indicated by a longer WBCL. For example, WBCL increased by 44.8% after 30-minutes and 108.5%
after 60-minutes exposure to 100 µg/mL DEHP (baseline: 92.2±8.2, 30-min: 133.5±21.2, 60-min:
192.2±71.5 ms cycle length, $p<0.001$; **Figure 5B**). In a similar manner, dynamic pacing was applied to
the right atrium at an increasing rate until 2:1 AV conduction block occurred. Following DEHP exposure,
2:1 AV block occurred at slower pacing frequencies within ~45 minutes (baseline: 62.6±3.1, 45-min:
114.4±41.3 ms cycle length, $p<0.05$; **Figure 5C**). To determine the AV node effective refractory period
(AVNERP), an extrastimulus pacing protocol was applied to the right atrium, wherein the cycle length of
the last stimulus (S2) was progressively shortened until ventricular conduction failed. At baseline,
AVNERP was reached at an S1-S2 interval of 150 ms (S1) and 73.8±9.5 ms cycle length (S2), which
increased to 117.5±25.4 ms cycle length (S2) after 30-minutes exposure to 100 µg/mL DEHP. With
longer exposure times, external pacing failed to stimulate the ventricles until exceedingly slow intervals
were used (90-min: 170.3±60.8 ms cycle length (S2), $p<0.0001$; **Figure 5D**).

<u>DEHP exposure slows epicardial conduction velocity and prolongs action potential duration</u>
Our lab, and others, have previously reported that DEHP disrupts gap junction intercellular
communication[28,37,48]. Reduced gap junction intercellular communication can slow both atrioventricular
conduction velocity (described earlier) and epicardial conduction velocity. Using a voltage-sensitive dye
and high-speed optical mapping, we observed that apicobasal conduction velocity remained stable in
heart preparations perfused with control media, between baseline (64.8±10.3 cm/sec) and 60-minutes
perfusion (64.1±14.9 cm/sec; **Figure 6A**). While apicobasal conduction velocity slowed by 15.6%
between baseline (58.9±5.8 cm/sec) and 60-minutes perfusion with 100 µg/ml DEHP (49.7±5.5
cm/sec). Optical mapping of transmembrane voltage signals was also performed to evaluate ventricular
action potential duration (APD), using a dynamic pacing protocol (S1-S1). Relative to time-matched
controls (60-min perfusion), 100 µg/mL DEHP exposure prolonged the ventricular APD at 30% and
80% repolarization at both slower and faster pacing frequencies (**Figure 6B, D, E**). As an example, at
200 PCL DEHP exposure increased APD_{30} by 29.9% (control: 18.4±2.3, 100 µg/mL DEHP: 23.9±3.5
ms) and APD_{80} by 29.4% (control: 54.1±5.8, 100 µg/mL DEHP: 70.0±5.6 ms). Using an extrastimulus

pacing protocol (S1-S2), we also observed that 60-minutes perfusion with 100 µg/mL DEHP lengthened

the ventricular effective refractory period (55.9±11.1 ms) as compared to time-matched controls (40.5±7

ms; **Figure 6C**). Finally, we monitored the effects of DEHP exposure on intracellular calcium handling,

but did not detect any measurable difference in the ventricular calcium transient duration time at 30% or

80%. At 200 PCL, CaD_{30} (control: 51.4±7 ms, 100 µg/mL DEHP: 53.1±5.5 ms) and CaD_{80} (control:

92.5±7.1 ms, 100 µg/mL DEHP: 100.0±8.9 ms) were comparable after 60-minutes perfusion (**Figure**

6F, G).

<u>Effects of DEHP on cardiac electrophysiology are not abolished by muscarinic antagonism</u>

Previous studies implicated cardiac muscarinic receptors as a potential target for phthalate plasticizers,

since atropine-treatment blunted the cardiodepressive effects of phthalates in both goldfish and human

atrial trabecular preparations[33,34]. We tested this possible mechanism by pretreating Langendorff-

perfused heart preparations for 10-15 minutes with 1 µM atropine, a concentration that has been shown

to augment the effects of acetylcholine[49], followed by exposure to 100 µg/mL DEHP. Across all the

tested variables, the effects of DEHP on sinus activity and atrioventricular conduction persisted in the

presence of atropine **(Figure 7A-F)**. In this set of studies, ventricular rate slowed by 51.5% and SNRT

increased by 123.5% in hearts with 100 µg/mL DEHP plus atropine, versus atropine alone.

<u>Effects of DEHP are partially relieved by inhibition of matrix metalloproteinases</u>

Previous work indicates that DEHP can disrupt intercellular coupling[28,50], and this effect may be

mechanistically linked to an upregulation of matrix metalloproteinase activity[36,51] that modulates gap

junction proteins and influences the electrical properties of the heart[52,53]. We tested this possible

mechanism by pretreating heart preparations for 10-15 minutes with 20 µM doxycycline, a synthetic

matrix metalloproteinase inhibitor[54]. Doxycycline reduced the depressive effects of DEHP on sinus

activity, as demonstrated by SNRT (Doxy alone: 218.5±43.3, DEHP+doxy: 253.3±10.1 ms, p=0.4),

cSNRT (doxy alone: 41±12.1 ms, DEHP+doxy: 46±18.9 ms, p=0.7), and beating rate measurements

(Doxy alone: 353.8±56.1, DEHP+doxy: 291.2±37.5 BPM, p=0.051; **Figure 7A-F)**. However, doxycycline

pretreatment did not ameliorate the effects of DEHP on atrioventricular conduction, as the PR interval

lengthened by 79.8%, WBCL increased by 87.2%, and AVNERP increased by 121.5% in hearts with

100 µg/mL DEHP plus doxycycline, versus doxycycline alone.

<u>DEHP slows cardiac conduction velocity in hiPSC-CM monolayers</u>

DEHP has been shown to reduce gap junction intercellular communication, but in select tissues (e.g.,

liver) this effect may be species-specific[55,56]. To test the applicability of our findings on human

cardiomyocytes, we measured conduction velocity using hiPSC-CM monolayers plated atop a
microelectrode array (paced at 1.5 Hz). DEHP-treatment slowed conduction velocity in a dose- and
time-dependent manner. Within the first 15-minutes, conduction velocity was unchanged in cell
preparations treated with 10 µg/mL DEHP, but declined by 14.6-18.3% in cells treated with higher
concentrations (50 µg/mL DEHP: 0.215±0.06; 100 µg/mL DEHP: 0.225±0.04 mm/ms) relative to control
cells (0.263±0.02 mm/ms; **Figure 8A,C).** With a longer 3 hour exposure, conduction velocity slowed by
12.9% in cells treated with 10 µg/mL DEHP (0.210±0.02 mm/ms) and 35.3-37% in cells treated with
higher concentrations (50 DEHP: 0.156±0.05; 100 DEHP: 0.152±0.04 mm/ms) relative to control cells
(0.241±0.02 mm/ms). Although conduction velocity measurements were comparable between 50 and
100 µg/mL DEHP, it became noticeably harder to pace hiPSC-CM at the higher DEHP concentration
(**Figure 8B**). As such, the subsequent inhibitor studies were performed using only the intermediate
dose of 50 µg/mL DEHP. Pretreatment with either atropine or doxycycline appeared to abrogate the
DEHP-induced conduction velocity slowing, up to 1 hour of DEHP exposure **(Figure 8D).** However, with
a longer 3-hour DEHP exposure time, hiPSC-CM conduction velocity slowed by 22.7% despite
pretreatment with atropine (atropine: 0.229±0.02; 50 DEHP+atropine: 0.177±0.03 mm/ms) and 21.8%
despite pretreatment with doxycycline (doxy: 0.243±0.02, 50 DEHP+doxy: 0.19±0.03 mm/ms).

DISCUSSION
Herein, we demonstrate that DEHP is a cardiotoxic chemical that slows sinus rate, delays sinus node
recovery time, and prolongs AV conduction in a dose- and time-dependent manner. Our findings build
upon previous studies by others, which report that low dose DEHP exposure (4 µg/ml) halts the
spontaneous beating activity of chick cardiomyocytes within 30-minutes and elicits cell death within 24-
hour exposure[30]. DEHP exposure has also been shown to precipitate bradycardia in the rat heart after
15-60 minutes[57] – while *in vivo* MEHP exposure (primary metabolite of DEHP) results in cardiac arrest
within minutes[32]. In the current study, we also report that DEHP slows atrioventricular conduction, as
measured by optical mapping, microelectrode array recordings, and electrocardiograms (PR interval) –
wherein the later indicates AV conduction delay that progresses to heart block with longer exposure
time (60-90 minutes). Previous work by Pfuderer et al. and Barry et al. suggested that phthalates exert
cardiodepressive effects likely through interaction with cholinergic receptors, which can be rescued with
atropine[33,34]. Although in our current study, pretreatment with 1 µM atropine did not prevent the effects
of DEHP on either sinus rate or AV conduction.

Phthalate chemicals have also been shown to inhibit gap junction intercellular communication (GJIC)
across multiple cell types[28,35,58–60], wherein the degree of inhibition is influenced by the length of the

carbon chains[37]. As an example, a comparative study conducted in liver cells reported that 80 μM

DEHP reduced GJIC by ~20% within 1 hour and ~80% within 16 hours of exposure[37]. Although a few

reports have suggested that phthalate-disruption of intercellular coupling is species-dependent, at least

in liver cells, wherein GJIC is impaired in mouse and rat hepatocytes, but not hamster, primate, or

human hepatocytes[55,56]. Notably, in heart tissue, intercellular gap junction coupling is a key determinant

of electrical conduction speed. In the current study, we report that 60-minute DEHP exposure slows

epicardial conduction velocity by ~15% in intact rat heart preparations. Further, we observed a similar

effect in a human cardiomyocyte model (hiPSC-CM monolayer), wherein conduction velocity slowing

was exaggerated with prolonged DEHP exposure (~37% at 3 hours). To date, the mechanisms by

which phthalate chemicals impair GJIC appear to be multifactorial. First, the lipophilic nature of DEHP

(and other phthalates) facilitate its physical incorporation into cell membranes – as demonstrated by

radiolabeling studies whereby DEHP localizes to the membrane fraction of RBCs[11]. This physical

disruption of cell membrane fluidity and organization has been confirmed by molecular dynamic

simulation studies, which report that DEHP incorporates and aggregates in lipid bilayers[61]. In this way,

DEHP and other phthalates may act in a manner similar to long-chain alcohols or fatty acids to disrupt

GJIC[62,63]. Second, DEHP and other phthalates can disrupt the trafficking and/or stabilization of gap

junction proteins at the cell membrane, by way of increased oxidative stress and MMP activity[35,36].

Intercellular coupling is impaired under conditions of oxidative stress, due to the perturbation of forward

trafficking and delivery of connexons to the cell membrane [64]. Intercellular coupling is also impaired with

increased MMP activity, as these enzymes can modulate electrical activity through the cleavage of

connexin proteins[52,53]. We found that pretreatment with an MMP inhibitor (doxycycline) blocked the

effects of DEHP on heart rate and sinus node recovery time in intact heart preparations, but AV

conduction slowing persisted. Further, the use of an MMP inhibitor reduced the effect of DEHP on

conduction velocity in hiPSC-CM exposed for 15-60 minutes, but conduction velocity slowing persisted

with longer exposure time (2-3 hours DEHP). Additional work is warranted to fully elucidate the

mechanisms by which DEHP perturbs cardiac electrophysiology, and to clarify the degree to which its

effects vary between different species and cell types.

DEHP has been widely used as a plasticizer additive in PVC-based medical products since the

1950s[65,66]; nevertheless, several studies have documented DEHP leaching from plastic tubing circuits

and intravenous or blood storage bags [7–9,15,41,67,68]. As such, patients can incur phthalate exposure when

undergoing multiple medical interventions that employ plastic products, including blood transfusions,

cardiopulmonary bypass, and extracorporeal membrane oxygenation[16–21,69]. Given the safety concerns

associated with phthalate exposure, there is renewed interest in reducing the use of DEHP-containing

plastics during blood donation, storage, and transfusion procedures[70,71]. Although, to date, the adoption

of DEHP-free plastics has been limited due to availability, cost, and concern that alternative materials

are inferior for blood storage applications[72,73]. Our study highlights the need for future work to

investigate the impact of phthalate chemical exposure on vulnerable patient groups. Notably, the

negative effects of DEHP on cardiac electrophysiology occurred within 15-90 minutes, which is well

within the timeframe of clinical exposure. Further, we quantified DEHP concentrations in stored RBC

units (23-119 µg/mL) and used these levels as a guide for our *in vitro* experiments (10-100 µg/mL).

Accordingly, our study is both clinically relevant and timely, as the results of this work may incentivize

the development and adoption of DEHP-free materials for cardiac surgery, transfusion, and intensive

care.

LIMITATIONS

Since DEHP is a non-persistent chemical, the scope of our study was limited to acute DEHP exposure

(15 min – 3 hours). An isolated, intact heart preparation was perfused with crystalloid buffer to optically

measure cardiac action potentials and intracellular calcium transients – as blood perfusion interferes

with fluorescent dye measurements. As such, the results of these *ex vivo* experiments may differ from

the *in vivo* environment. To address human applicability, we also employed an hiPSC-CM model for

conduction velocity measurements; however, this simplified two-dimensional model lacks the

complexity of the internal cardiac conduction system and prevents any electrical measurements

between chambers of the heart (e.g., interatrial or atrioventricular conduction). hiPSC-CM are

frequently used for preclinical studies of drug efficacy and safety, as these cells express the same ion

channels that make up the human cardiac action potential. However, it is important to note that hiPSC-

CM have an immature phenotype that more closely resembles a fetal-infant cardiomyocyte versus an

adult cardiomyocyte[74,75]. Finally, the reported results are limited to the effect of DEHP, and therefore,

do not take into account additional effects caused by DEHP metabolites.

ACKNOWLEDGEMENTS

The authors acknowledge Blake Cooper for assistance with hiPSC-CM experiments, and Francesca

Cendali and Daniel Stephenson PhD for assistance with phthalate chemical quantitation. We

additionally acknowledge the Pathology & Laboratory Medicine Research Core at Children's National

Hospital for assistance with RBC unit storage and sample collection, including Meghan Delaney MD,

Willian Suslovic, Casey Yarbrough, Aszia Burrell, Eric Freeman, Abigail Williams.

FUNDING SUPPORT

NGP was supported by the National Heart, Lung, and Blood Institute (R01HL139472), Children's
Research Institute, and Children's National Heart Institute. DG was supported by the American Heart
Association (23PRE1021149). AD was supported by the National Institute of General and Medical
Sciences (RM1GM131968), and from the National Heart, Lung, and Blood Institute (R01HL146442,
R01HL149714, R01HL148151, R01HL161004).

References

1.   Shelby MD. NTP-CERHR monograph on the potential human reproductive and developmental
effects of di (2-ethylhexyl) phthalate (DEHP). *NTP CERHR MON.* 2006;(18):v, vii-7, II-iii-xiii
passim. Accessed May 31, 2020. http://www.ncbi.nlm.nih.gov/pubmed/19407857

2.   Ramadan M, Cooper B, Posnack NG. Bisphenols and phthalates: Plastic chemical exposures
can contribute to adverse cardiovascular health outcomes. *Birth Defects Res.* Published online
2020. doi:10.1002/bdr2.1752

3.   DiGangi J. *Phthalates in PVC Medical Products from 12 Countries.*; 1999.
https://wayback.archive-it.org/9650/20200516064437/http://p3-
raw.greenpeace.org/international/Global/international/planet-2/report/1999/9/phthalates-in-pvc-
medical-prod.pdf

4.   Jaeger RJ, Rubin RJ. Extraction, localization, and metabolism of di-2-ethylhexyl phthalate from
PVC plastic medical devices. *Environ Health Perspect.* 1973;3:95-102. doi:10.1289/ehp.730395

5.   Rose RJ, Priston MJ, Rigby-Jones AE, Sneyd JR. The effect of temperature on di(2-ethylhexyl)
phthalate leaching from PVC infusion sets exposed to lipid emulsions. *Anaesthesia.*
2012;67(5):514-520. doi:10.1111/J.1365-2044.2011.07006.X

6.   Bagel-Boithias S, Sautou-Miranda V, Bourdeaux D, Tramier V, Boyer A, Chopineau J. Leaching
of diethylhexyl phthalate from multilayer tubing into etoposide infusion solutions. *Am J Health*
Syst Pharm. 2005;62(2):182-188. doi:10.1093/AJHP/62.2.182

7.   Karle VA, Short BL, Martin GR, et al. Extracorporeal membrane oxygenation exposes infants to
the plasticizer, di(2-ethylhexyl)phthalate. *Crit Care Med.* 1997;25(4):696-703. Accessed October
20, 2016. http://www.ncbi.nlm.nih.gov/pubmed/9142038

8.   Loff S, Kabs F, Subotic U, Schaible T, Reinecke F, Langbein M. Kinetics of diethylhexyl-
phthalate extraction From polyvinylchloride-infusion lines. *JPEN J Parenter Enteral Nutr.*
2002;26(5):305-309. doi:10.1177/0148607102026005305

9.   D'alessandro A, Nemkov T, Hansen KC. Rapid detection of DEHP in packed red blood cells
stored under European and US standard conditions. *Blood Transfus.* 2016;14(2):140-144.
doi:10.2450/2015.0210-15

10. Rock G, Labow RS, Tocchi M. Distribution of di(2-ethylhexyl) phthalate and products in blood
and blood components. *Environ Health Perspect.* 1986;65:309. doi:10.1289/EHP.8665309

11. Rock G, Tocchi M, Ganz PR, Tackaberry ES. Incorporation of plasticizer into red cells during
storage. *Transfusion (Paris).* 1984;24(6):493-498. doi:10.1046/j.1537-2995.1984.24685066808.x

12. Labow RS, Card RT, Rock G. The effect of the plasticizer di(2-ethylhexyl)phthalate on red cell
deformability. *Blood.* 1987;70(1):319-323. Accessed January 23, 2017.
http://www.ncbi.nlm.nih.gov/pubmed/3593970

13. AuBuchon JP, Estep TN, Davey RJ. The effect of the plasticizer di-2-ethylhexyl phthalate on the
survival of stored RBCs. *Blood.* 1988;71(2):448-452. Accessed November 17, 2016.
http://www.ncbi.nlm.nih.gov/pubmed/3337906

14.   Horowitz B, Stryker MH, Waldman AA, Woods KR, Gass JD, Drago J. Stabilization of red blood
cells by the plasticizer, diethylhexylphthalate. *Vox Sang.* 1985;48(3):150-155. Accessed January
23, 2017. http://www.ncbi.nlm.nih.gov/pubmed/3984301

15.   Inoue K, Kawaguchi M, Yamanaka R, et al. Evaluation and analysis of exposure levels of di(2-
ethylhexyl) phthalate from blood bags. *Clin Chim Acta.* 2005;358(1-2):159-166.
doi:10.1016/J.CCCN.2005.02.019

16.   Kaestner F, Seiler F, Rapp D, et al. Exposure of patients to di(2-ethylhexy)phthalate (DEHP) and
its metabolite MEHP during extracorporeal membrane oxygenation (ECMO) therapy. Meliker J,
ed. 2020;15(1):e0224931. doi:10.1371/journal.pone.0224931

17.   Huygh J, Clotman K, Malarvannan G, et al. Considerable exposure to the endocrine disrupting
chemicals phthalates and bisphenol-A in intensive care unit (ICU) patients. *Environ Int.*
2015;81:64-72. doi:10.1016/j.envint.2015.04.008

18.   Green R, Hauser R, Calafat AM, et al. Use of di(2-ethylhexyl) phthalate-containing medical
products and urinary levels of mono(2-ethylhexyl) phthalate in neonatal intensive care unit
infants. *Environ Health Perspect.* 2005;113(9):1222-1225. doi:10.1289/ehp.7932

19.   Weuve J, Sánchez BN, Calafat AM, et al. Exposure to phthalates in neonatal intensive care unit
infants: urinary concentrations of monoesters and oxidative metabolites. *Environ Health
Perspect.* 2006;114(9):1424-1431.

20.   Calafat AM, Needham LL, Silva MJ, Lambert G. Exposure to di-(2-ethylhexyl) phthalate among
premature neonates in a neonatal intensive care unit. *Pediatrics.* 2004;113(5):e429-34.
doi:10.1126/science.170.3956.460

21.   Mallow EB, Fox MA. Phthalates and critically ill neonates: device-related exposures and non-
endocrine toxic risks. *Journal of Perinatology.* 2014;34(12):892-897. doi:10.1038/jp.2014.157

22.   Braun JM, Sathyanarayana S, Hauser R. *Phthalate Exposure and Children's Health.* Vol 25. Curr
Opin Pediatr; 2013:247-254.

23.   Posnack NG. Plastics and cardiovascular disease. *Nat Rev Cardiol.* 2021;18(2):69-70.
doi:10.1038/s41569-020-00474-4

24.   Tereshchenko LG, Posnack NG. Does plastic chemical exposure contribute to sudden death of
patients on dialysis? *Heart Rhythm.* 2019;16(2):312-317. doi:10.1016/j.hrthm.2018.08.020

25.   Trasande L, Liu B, Bao W. Phthalates and attributable mortality: A population-based longitudinal
cohort study and cost analysis. *Environ Pollut.* 2022;292(Pt A):118021.
doi:10.1016/J.ENVPOL.2021.118021

26.   Zeng G, Zhang Q, Wang X, Wu KH. Low-level plasticizer exposure and all-cause and
cardiovascular disease mortality in the general population. *Environ Health.* 2022;21(1):1-11.
doi:10.1186/S12940-022-00841-3/FIGURES/2

27.   Posnack NG. The Adverse Cardiac Effects of Di(2-ethylhexyl)phthalate and Bisphenol A.
Cardiovasc Toxicol. 2014;14(4):339-357. doi:10.1007/s12012-014-9258-y

28. Gillum N, Karabekian Z, Swift LM, Brown RP, Kay MW, Sarvazyan N. Clinically relevant
concentrations of di (2-ethylhexyl) phthalate (DEHP) uncouple cardiac syncytium. *Toxicol Appl*
Pharmacol. 2009;236(1):25-38. doi:10.1016/j.taap.2008.12.027

29. Jaimes R, McCullough D, Siegel B, et al. Plasticizer Interaction with the Heart: Chemicals Used
in Plastic Medical Devices Can Interfere with Cardiac Electrophysiology. *Circ Arrhythm*
Electrophysiol. 2019;12(7). doi:10.1161/CIRCEP.119.007294

30. Rubin RJ, Jaeger RJ. Some pharmacologic and toxicologic effects of di-2-ethylhexyl phthalate
(DEHP) and other plasticizers. *Environ Health Perspect*. 1973;3:53-59. doi:10.1289/ehp.730353

31. Posnack NG, Idrees R, Ding H, et al. Exposure to phthalates affects calcium handling and
intercellular connectivity of human stem cell-derived cardiomyocytes. *PLoS One*.
2015;10(3):e0121927-e0121927. doi:10.1371/journal.pone.0121927

32. Rock G, Labow RS, Franklin C, Burnett R, Tocchi M. Hypotension and cardiac arrest in rats after
infusion of mono(2-ethylhexyl) phthalate (MEHP), a contaminant of stored blood. *N Engl J Med*.
1987;316(19):1218-1219. doi:10.1056/NEJM198705073161915

33. Barry YA, Labow RS, Keon WJ, Tocchi M. Atropine inhibition of the cardiodepressive effect of
mono(2-ethylhexyl)phthalate on human myocardium. *Toxicol Appl Pharmacol*. 1990;106(1):48-
52. doi:10.1016/0041-008X(90)90104-3

34. Pfuderer P, Francis AA. Phthalate esters: Heartrate depressors in the goldfish. *Bull Environ*
Contam Toxicol. 1975;13(3):275-279. doi:10.1007/BF01685335

35. Sobarzo CM, Rosana N de M, Livia L, Berta D, Schteingart HF. Mono-(2-ethylhexyl) phthalate
(MEHP) affects intercellular junctions of Sertoli cell: A potential role of oxidative stress. *Reprod*
Toxicol. 2015;58:203-212. doi:10.1016/J.REPROTOX.2015.10.010

36. Yao PL, Lin YC, Richburg JH. Mono-(2-Ethylhexyl) Phthalate-Induced Disruption of Junctional
Complexes in the Seminiferous Epithelium of the Rodent Testis Is Mediated by MMP21. *Biol*
Reprod. 2010;82(3):516-527. doi:10.1095/biolreprod.109.080374

37. Čtveráčková L, Jančula D, Raška J, Babica P, Sovadinová I. Structure-Dependent Effects of
Phthalates on Intercellular and Intracellular Communication in Liver Oval Cells. *Int J Mol Sci*.
2020;21(17):1-21. doi:10.3390/IJMS21176069

38. Fedorov V V., Lozinsky IT, Sosunov EA, et al. Application of blebbistatin as an excitation-
contraction uncoupler for electrophysiologic study of rat and rabbit hearts. *Heart rhythm : the*
official journal of the Heart Rhythm Society. 2007;4(5):619-626. doi:10.1016/j.hrthm.2006.12.047

39. Swift LM, Kay MW, Ripplinger CM, Posnack NG. Stop the beat to see the rhythm: excitation-
contraction uncoupling in cardiac research. *Am J Physiol Heart Circ Physiol*. 2021;321(6):H1005-
H1013. doi:10.1152/AJPHEART.00477.2021

40. Swift LM, Asfour H, Posnack NG, Arutunyan A, Kay MW, Sarvazyan N. Properties of blebbistatin
for cardiac optical mapping and other imaging applications. *Pflugers Arch Eur J Physiol*.
2012;464(5):503-512. doi:10.1007/s00424-012-1147-2

41. Rael LT, Bar-Or R, Ambruso DR, et al. Phthalate esters used as plasticizers in packed red blood
cell storage bags may lead to progressive toxin exposure and the release of pro-inflammatory

cytokines. *Oxid Med Cell Longev*. 2009;2(3):166-171. Accessed January 1, 2019.
http://www.ncbi.nlm.nih.gov/pubmed/20592772

42.   Sjoberg PO, Bondesson UG, Sedin EG, Gustafsson JP. Exposure of newborn infants to
plasticizers. Plasma levels of di-(2-ethylhexyl) phthalate and mono-(2-ethylhexyl) phthalate
during exchange transfusion. *Transfusion (Paris)*. 1985;25(5):424-428. doi:10.1046/j.1537-
2995.1985.25586020115.x

43.   Swift LM, Jaimes R, McCullough D, et al. *Optocardiography and Electrophysiology Studies of Ex
Vivo Langendorff-Perfused Hearts*. Vol 153.; 2019. Accessed September 21, 2019.
https://www.jove.com/video/60472/

44.   Jaimes R, McCullough D, Siegel B, et al. Lights, camera, path splitter: a new approach for truly
simultaneous dual optical mapping of the heart with a single camera. *BMC Biomed Eng*.
2019;1(25):25. doi:10.1186/s42490-019-0024-x

45.   Jaimes R, Walton RD, Pasdois PLCP, Bernus O, Efimov IR, Kay MW. A technical review of
optical mapping of intracellular calcium within myocardial tissue. *American Journal of Physiology-
Heart and Circulatory Physiology*. 2016;310(11):H1388-401. doi:10.1152/ajpheart.00665.2015

46.   Cooper BL, Gloschat C, Swift LM, et al. KairoSight: Open-Source Software for the Analysis of
Cardiac Optical Data Collected From Multiple Species. *Front Physiol*. 2021;0:1820.
doi:10.3389/FPHYS.2021.752940

47.   Nemkov T, Reisz JA, Gehrke S, Hansen KC, D'Alessandro A. High-throughput metabolomics:
Isocratic and gradient mass spectrometry-based methods. In: *Methods in Molecular Biology*. Vol
1978. Humana Press Inc.; 2019:13-26. doi:10.1007/978-1-4939-9236-2_2

48.   Di Lorenzo M, Winge SB, Svingen T, De Falco M, Boberg J. Intrauterine exposure to diethylhexyl
phthalate disrupts gap junctions in the fetal rat testis. *Curr Res Toxicol*. 2020;1:5-11.
doi:10.1016/J.CRTOX.2020.02.002

49.   Hoover DB, Neely DA. Differentiation of Muscarinic Receptors Mediating Negative Chronotropic
and Vasoconstrictor Responses to Acetylcholine in Isolated Rat Hearts. *Journal of
Pharmacology and Experimental Therapeutics*. 1997;282(3).

50.   Zhao YX, Tang YX, Sun XH, et al. Gap Junction Protein Connexin 43 as a Target Is Internalized
in Astrocyte Neurotoxicity Caused by Di-(2-ethylhexyl) Phthalate. *J Agric Food Chem*.
2022;70(19):5921-5931.
doi:10.1021/ACS.JAFC.2C01635/SUPPL_FILE/JF2C01635_SI_001.PDF

51.   Posnack NG, Lee NH, Brown R, Sarvazyan N. Gene expression profiling of DEHP-treated
cardiomyocytes reveals potential causes of phthalate arrhythmogenicity. *Toxicology*.
2011;279(1-3):54. doi:10.1016/J.TOX.2010.09.007

52.   Lindsey ML, Escobar GP, Mukherjee R, et al. Matrix metalloproteinase-7 affects connexin-43
levels, electrical conduction, and survival after myocardial infarction. *Circulation*.
2006;113(25):2919-2928. doi:10.1161/CIRCULATIONAHA.106.612960

53.   Wang J, Li JS, Liu HZ, et al. Dynamic alterations of connexin43, matrix metalloproteinase-2 and
tissue inhibitor of matrix metalloproteinase-2 during ventricular fibrillation in canine. *Mol Cell
Biochem*. 2014;391(1-2):259-266. doi:10.1007/S11010-014-2012-8/FIGURES/5

54.   Scannevin RH, Alexander R, Haarlander TM, et al. Discovery of a highly selective chemical
634   inhibitor of matrix metalloproteinase-9 (MMP-9) that allosterically inhibits zymogen activation.
635   *Journal of Biological Chemistry*. 2017;292(43):17963-17974. doi:10.1074/jbc.M117.806075

636   55.   Isenberg JS, Kamendulis LM, Smith JH, et al. Effects of Di-2-Ethylhexyl Phthalate (DEHP) on
637   Gap-Junctional Intercellular Communication (GJIC), DNA Synthesis, and Peroxisomal Beta
638   Oxidation (PBOX) in Rat, Mouse, and Hamster Liver. *Toxicological Sciences*. 2000;56(1):73-85.
639   doi:10.1093/toxsci/56.1.73

640   56.   Kamendulis LM, Isenberg JS, Smith JH, et al. Comparative effects of phthalate monoesters on
641   gap junctional intercellular communication and peroxisome proliferation in rodent and primate
642   hepatocytes. *J Toxicol Environ Health A*. 2002;65(8):569-588.
643   doi:10.1080/152873902317349736

644   57.   Aronson CE, Serlick ER, Preti G. Effects of di-2-ethylhexyl phthalate on the isolated perfused rat
645   heart. *Toxicol Appl Pharmacol*. 1978;44(1):155-169. doi:10.1016/0041-008X(78)90295-8

646   58.   Sobarzo CM, Lustig L, Ponzio R, Suescun MO, Denduchis B. Effects of di(2-ethylhexyl)
647   phthalate on gap and tight junction protein expression in the testis of prepubertal rats. *Microsc
648   Res Tech*. 2009;72(11):868-877. doi:10.1002/jemt.20741

649   59.   Cruciani V, Mikalsen S, Vasseur P, Sanner T. Effects of Peroxisome Proliferators and 12-O-
650   tetradecanoyl phorbol-13-acetate on Intercellular Communication and connexin43 in Two
651   Hamster Fibroblast Systems. *Int J Cancer*. 1997;73(2). doi:10.1002/(SICI)1097-
652   0215(19971009)73:2<240::AID-IJC14>3.0.CO;2-J

653   60.   Isenberg JS, Kamendulis LM, Ackley DC, et al. Reversibility and persistence of di-2-ethylhexyl
654   phthalate (DEHP)- and phenobarbital-induced hepatocellular changes in rodents. *Toxicol Sci*.
655   2001;64(2):192-199. doi:10.1093/TOXSCI/64.2.192

656   61.   Bider RC, Lluka T, Himbert S, et al. Stabilization of Lipid Membranes through Partitioning of the
657   Blood Bag Plasticizer Di-2-ethylhexyl phthalate (DEHP). Published online 2020.
658   doi:10.1021/acs.langmuir.0c01964

659   62.   Burt JM, Massey KD, Minnich BN. Uncoupling of cardiac cells by fatty acids: structure-activity
660   relationships. *https://doi.org/101152/ajpcell19912603C439*. 1991;260(3 29-3).
661   doi:10.1152/AJPCELL.1991.260.3.C439

662   63.   Bastiaanse EML, Jongsma HJ, van der Laarse A, Takens-Kwak BR. Heptanol-induced decrease
663   in cardiac gap junctional conductance is mediated by a decrease in the fluidity of membranous
664   cholesterol-rich domains. *J Membr Biol*. 1993;136(2):135-145.
665   doi:10.1007/BF02505758/METRICS

666   64.   Smyth JW, Hong TT, Gao D, et al. Limited forward trafficking of connexin 43 reduces cell-cell
667   coupling in stressed human and mouse myocardium. 2010;120(1):266-279.
668   doi:10.1172/JCI39740

669   65.   Walter CW. Invention and development of the blood bag. *Vox Sang*. 1984;47(4):318-324.
670   doi:10.1111/J.1423-0410.1984.TB01603.X

671   66.   Walter CW, MurphyY WP. A closed gravity technique for the preservation of whole blood in ACD
672   solution utilizing plastic equipment. *Surg Gynecol Obstet.* 1952;94(6):687-692. Accessed April 9,
673   2023. https://pubmed.ncbi.nlm.nih.gov/14931176/

674   67.   Loff S, Subotic U, Reinicke F, Wischmann H, Brade J. Extraction of di-ethylhexyl-phthalate from
675   perfusion lines of various material, length and brand by lipid emulsions. *J Pediatr Gastroenterol
676   Nutr.* 2004;39(4):341-345. doi:10.1097/00005176-200410000-00008

677   68.   Loff S, Kabs F, Witt K, et al. Polyvinylchloride infusion lines expose infants to large amounts of
678   toxic plasticizers. *J Pediatr Surg.* 2000;35(12):1775-1781. doi:10.1053/jpsu.2000.19249

679   69.   Eckert E, Müller J, Höllerer C, et al. Plasticizer exposure of infants during cardiac surgery.
680   *Toxicol Lett.* 2020;330:7-13. doi:10.1016/J.TOXLET.2020.04.004

681   70.   van der Meer PF, Reesink HW, Panzer S, et al. Should DEHP be eliminated in blood bags? *Vox
682   Sang.* 2014;106(2):176-195. doi:10.1111/vox.12099

683   71.   Prowse C V., de Korte D, Hess JR, van der Meer PF. Commercially available blood storage
684   containers. *Vox Sang.* 2014;106(1):1-13. doi:10.1111/vox.12084

685   72.   van der Meer PF, Devine D V., Irving DO, et al. Alternatives in blood operations when choosing
686   non-DEHP bags. *Vox Sang.* 2017;112(2):183-184. doi:10.1111/VOX.12479

687   73.   Razatos A, Acker JP, de Korte D, et al. Survey of blood centre readiness regarding removal of
688   DEHP from blood bag sets: The BEST Collaborative Study. *Vox Sang.* 2022;117(6):796-802.
689   doi:10.1111/VOX.13258

690   74.   Lundy SD, Zhu WZ, Regnier M, Laflamme MA. Structural and functional maturation of
691   cardiomyocytes derived from human pluripotent stem cells. *Stem Cells Dev.* 2013;22(14):1991-
692   2002. doi:10.1089/SCD.2012.0490

693   75.   Karbassi E, Fenix A, Marchiano S, et al. Cardiomyocyte maturation: advances in knowledge and
694   implications for regenerative medicine. *Nat Rev Cardiol.* Published online February 3, 2020.
695   doi:10.1038/s41569-019-0331-x

696
697

698
699

700
701 **Figure 1. DEHP concentration increases in refrigerated RBC units with longer storage time.** RBC
702 unit supernatant samples were collected from units stored 7-42 days; DEHP levels were quantitated
703 using ultra high-pressure liquid chromatography coupled to mass spectrometry. *One-way ANOVA with*
704 *Dunnett's multiple comparisons test. Significance compared to day 7 RBC unit, denoted by *p<0.05*
705

Figure 2. DEHP slows sinus rate and recovery time in a dose-dependent manner. A) Right: Pseudo-ECG recordings from Langendorff-perfused hearts at baseline and after exposure to 100 µg/mL DEHP. Middle: Sinus rate slows and sinus rhythm becomes more erratic following acute exposure to higher concentrations of DEHP. Left: Sinus rate and rhythm remain stable in time-controlled experiments with control media perfusion. **B)** Right: Measurements of sinus note recovery time (SNRT) at baseline and after exposure to 100 µg/mL DEHP. Middle: DEHP exposure slows sinus node recovery. Left: Sinus node recovery remains unchanged in time-controlled experiments with control media perfusion. *Cardiac preparations were omitted from analysis if external pacing with 1:1 capture could not be achieved. Comparisons were analyzed by two-tailed T-test (baseline vs control) or one-way ANOVA (dose response) with multiple comparisons test. Significance compared to baseline denoted by *p<0.05.*

Figure 3. DEHP slows sinus rate and recovery time in a time-dependent manner. A) Pseudo-ECG recordings from Langendorff-perfused hearts at baseline and after 90-minute exposure to 100 µg/mL DEHP (note atrioventricular uncoupling). Prolonged exposure to 100 µg/mL DEHP resulted in **B)** slower spontaneous sinus rate and ventricular rate, **C)** greater heart rate variability (RMSSD), **D)** longer sinus node recovery time (SNRT), **E)** a longer rate-corrected SNRT (cSNRT), and **F)** reduced coronary flow. *Cardiac preparations were omitted from analysis if external pacing with 1:1 capture could not be achieved. Comparisons were analyzed by one-way ANOVA with multiple comparisons test. Significance compared to baseline denoted by *p<0.05.*

727
Figure 4. DEHP exposure delays inter-atrial and atrioventricular conduction. A) Time series
images show the activation time (red) progressing from the atria to the ventricles, during sinus rhythm,
in control and DEHP-treated heart preparations (60 minutes). **B)** Inter-atrial activation time measured
using two microelectrode arrays positioned on the right atria (RA) and left atria (LA). **C)** Atrioventricular
activation time measured using two microelectrode arrays positioned on the right atria (RA) and left
ventricle (LV). Note the delay in activation time after DEHP exposure. AU: arbitrary units. *Comparisons
were analyzed by two-tailed T-test (baseline vs DEHP); significance denoted by *p<0.05.*
735

Figure 5. DEHP exposure increases atrioventricular conduction time and refractory period. A)
DEHP exposure slows atrioventricular conduction in a time-dependent manner, first observed as
lengthening of the PR interval, then occasional dropped beats that fail to propagate to the ventricles,
and then complete atrioventricular block with dissociation between atrial and ventricular signals. **B)**
Programmed electrical stimulation at the atria shows a progressive increase in WBCL, **C)** 2:1
(atria:ventricular) capture at slower pacing frequencies, and **D)** increased AV node refractoriness at
slower pacing frequencies. *Cardiac preparations were omitted from analysis if external pacing could not
be achieved. Comparisons were analyzed by one-way ANOVA with multiple comparisons test.
Significance compared to baseline denoted by *p<0.05.*

Figure 6. DEHP exposure slows epicardial conduction velocity and prolongs action potential
duration. A) DEHP exposure slows apico-basal conduction velocity, compared to time-matched
controls. **B-E)** 60-minutes DEHP exposure lengthens the ventricular action potential duration at 30%
(APD_{30}) and 80% repolarization (APD_{80}) and ventricular effective refractory period (VERP). **F)** DEHP
exposure had no measurable effect on the calcium transient duration time at 30% (CaD_{30}) or 80%
(CaD_{80}). *Cardiac preparations were omitted from analysis if external pacing could not be achieved, or*
poor signal:noise prevented accurate measurements. VERP comparisons were analyzed by two-tailed
T-test. Action potential and calcium transient comparisons were analyzed by two-way ANOVA with
Holm-Sidak test for multiple comparisons. Significance between time-matched controls and DEHP (60-
minutes) denoted by *$p < 0.05$.

758
759
Figure 7. Effects of DEHP are partially relieved by inhibition of matrix metalloproteinases but not
muscarinic antagonism. Langendorff-perfused hearts were exposed to 100 µg/mL DEHP for 60-
minutes, without or with pretreatment using atropine (muscarinic antagonist) or doxycycline (matrix
metalloprotinease inhibitor). Cardiac parameters were assessed relative to the vehicle (control,
atropine, or doxycycline alone), including: **A)** ventricular beating rate, **B)** sinus node recovery time
(SNRT), **C)** heart rate corrected SNRT, **D)** PR interval, **E)** Wenckebach cycle length (WBCL), **F)**
atrioventricular nodal effective refractory period (AVNERP). *Cardiac preparations were omitted from*
analysis if external pacing could not be achieved. Comparisons to respective vehicle were analyzed by
*two-tailed T-test. Significance denoted by *p<0.05.*
769

Figure 8. DEHP slows conduction velocity in hiPSC-CM monolayers. Conduction velocity was
measured across hiPSC-CM monolayers plated atop microelectrode arrays, in response to external
pacing (1.5 Hz). **A)** Activation maps show electrical signal propagating from lower right corner (pacing
electrode) across the hiPSC-CM monolayer after 1-3 hours treatment with control media or media
supplemented with 50-100 µg/mL DEHP. **B)** hiPSC-CM treated with higher concentrations of DEHP
were less likely to pace at the desired frequency and were omitted from analysis. **C)** Conduction
velocity slowing in the presence of DEHP is time- and dose-dependent. **D)** After 15-60 minutes DEHP
treatment, the effect on conduction velocity is reduced in the presence of either atropine (muscarinic
antagonist) or doxycycline (matrix metalloproteinase). DEHP-induced conduction velocity slowing
persists after prolonged 2-3 hour treatment. *Comparisons were analyzed by two-way ANOVA with*
Holm-Sidak test for multiple comparisons. Significance between control and DEHP denoted by *p<0.05.