**Dnmt3a1 regulates hippocampus-dependent memory via the downstream target Nrp1**

Janina Kupke[1], Julien Klimmt[1,2,#], Franziska Mudlaff[1,#], Maximilian Schwab[1] Carsten Sticht[3]
and Ana M.M. Oliveira[1,4,*]

[1]Department of Neurobiology, Interdisciplinary Center for Neurosciences (IZN), Heidelberg
University, 69120 Heidelberg, Germany

[2]Present address: Institute for Stroke and Dementia Research, University Hospital, Ludwig-
Maximilians-University Munich, 81377 Munich, Germany.

[3]Next Generation Sequencing Core Facility, Medical Faculty Mannheim, Heidelberg
University, 68167, Mannheim, Germany.

[4]Department of Molecular and Cellular Cognition Research, Central Institute of Mental Health,
Medical Faculty Mannheim, Heidelberg University, 68159 Mannheim, Germany

[#]equal contribution

*Ana M.M. Oliveira. Department of Molecular and Cellular Cognition Research, Central
Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, 68159
Mannheim, Germany. Phone: +49(0)62215416510. Email: oliveira@nbio.uni-heidelberg.de,
ana.oliveira@zi-manheim.de

**Author Contributions:** JKupke, AMMO designed research; JKupke, JKlimmt, FM, AMMO
performed experiments; JKupke, JKlimmt, FM, CS, AMMO analyzed data; JKupke, AMMO
wrote the paper. All authors edited the manuscript.

**Abstract**

Epigenetic factors are well established players in memory formation. Specifically, DNA methylation is necessary for the formation of long-term memory in multiple brain regions including the hippocampus. Despite the demonstrated role for DNA methyltransferases (Dnmts) in memory formation, it is unclear whether individual Dnmts have unique or redundant functions in long-term memory formation. Furthermore, the downstream processes controlled by Dnmts during memory consolidation have not been investigated. In this study, we examined the requirement of the predominant Dnmt in the adult brain, Dnmt3a1, for hippocampus-dependent long-term memory formation. Using RNA-sequencing, we identified an activity-regulated Dnmt3a1-dependent genomic program in which several genes were functionally associated with functional and structural plasticity. Our data showed that Dnmt3a1, similarly to its isoform Dnmt3a2, plays a critical role in memory formation and identified Neuropilin 1 (Nrp1) as a downstream target of Dnmt3a1 in this process. Intriguingly, we found that Nrp1 expression is selectively regulated by and a specific downstream effector of Dnmt3a1, but not Dnmt3a2. Taken together, our study uncovered a Dnmt3a-isoform-specific mechanism in memory formation, identified a novel regulator of memory, and further highlighted the complex and highly regulated functions of distinct epigenetic regulators in brain function.

## Introduction

Precise control of gene transcription is required for the functional and structural neuronal remodelling that underlies memory formation and maintenance[1–3]. Gene transcription is orchestrated by the activity of transcription factors[4,5], epigenetic factors, and 3D-genomic factors such as chromatin structure or nuclear organization[6–9]. Indeed, epigenetic mechanisms such as DNA methylation have emerged as important regulators of cognitive processes[8].

DNA methylation has traditionally been viewed as a static mechanism with important roles in transcription repression, cell fate determination, imprinting and silencing of transposons during early development [10]. However, it is now well accepted that DNA methylation plays a critical role in the regulation of genomic responses in mature neurons. Several lines of evidence demonstrated dynamic and quick changes in DNA methylation in response to neuronal activity[11,12] and pharmacological and genetic studies established a causal link between DNA methyltransferase (Dnmt) activity and memory formation. Specifically, the *de novo* DNA methyltransferase, Dnmt3a, plays a documented function in cognitive abilities[13–17]. Furthermore, it has recently been shown that *DNMT3A* variants are the underlying cause of the neurodevelopmental disorder Tatton-Brown-Rahman syndrome, which is characterized by intellectual disability[18,19]. Despite this evidence, the downstream processes regulated by Dnmt3a that contribute to memory formation are unknown. Moreover, the genomic program regulated by Dnmt3a and required for the formation of long-term memory remains uncovered.

The Dnmt3a genomic locus contains two genes, Dnmt3a1 and Dnmt3a2[20], that exhibit different genomic binding[21] and expression regulation in response to neuronal activity. Dnmt3a2 is transcribed from an intronic promoter and behaves like an immediate early gene, whereas the expression of Dnmt3a1 is not regulated by neuronal activity[22–24]. The requirement for Dnmt3a2 in memory formation has been demonstrated[23–25]; however, the contribution of Dnmt3a1, despite its high abundance in the adult brain[26–28] has not been investigated. The differential regulation of expression of the two isoforms by neuronal activity and the studies that showed that normal Dnmt3a1 expression levels do not compensate the memory impairments found in Dnmt3a2 knockdown mice [23,24,29,30], suggest nonredundant functions by the two isoforms. However, whether the two enzymes are differentially required for and regulate distinct downstream processes during memory formation remains to be investigated.

In this study, we demonstrated that Dnmt3a1 is crucial for memory formation and for controlling an activity-regulated genomic program. We found that acute hippocampal reduction of Dnmt3a1 led to memory impairments in adult mice. Moreover, using RNA Sequencing analysis, we identified activity-regulated genes dependent on Dnmt3a1 levels. Intriguingly, we found that some of the identified genes are selectively dependent on Dnmt3a1, but not

Dnmt3a2. Among these genes was Neuropilin-1 (Nrp1), a member of the semaphorin complex
with an established function in synaptic plasticity[31,32]. In a next step, we demonstrated the
involvement of Nrp1 in hippocampus-dependent memory formation. Furthermore, we showed
that Dnmt3a1 regulates hippocampus-dependent memory via Nrp1. In contrast, Nrp1
overexpression did not rescue memory impairments triggered by reduced Dnmt3a2 levels,
suggesting an isoform-specific mechanism for memory processes. Taken together these
findings uncovered a Dnmt3a1-dependent genomic program and identified a downstream
effector during memory formation.

**Materials and Methods**

Animals.

In this study, 3-months-old C57BL/6N male mice (Charles River, Sulzfeld, Germany) were

used. Mice were housed in groups of 2-3 per cage (except if severe fighting occurred), under

a 12h light/dark cycle with *ad libitum* access to water and food, 22 ± 1°C and 55 ± 10% relative

humidity. The behavioural experiments were conducted during the light phase. Mice that were

sick or injured were excluded. Animals were randomly assigned to experimental groups and

analysis was conducted blindly. The procedures complied with German animal care guidelines

and European Community Council Directive 86/609/EEC.

Recombinant Adeno Associated Virus (rAAV) production.

Viral particles were produced and purified as described previously[33] and with minor

modifications. Briefly, rAAVs were generated by co-transfecting AAV-293 cells (Stratagene

#240073, California, USA) with target AAV plasmid and helper plasmids (pFΔ6, pRV1, and

pH21) using calcium phosphate precipitation. 60 hours after transfection, the HEK 293 cells

were collected and lysed. Remaining viral particles in media were concentrated using 40%

Polyethylene glycol (PEG) solution (pH 7,4). For this, the culture medium was added to 40%

PEG and stirred for 3h at 4°C. After centrifugation, the viral pellet was added to lysed HEK

cells and this was purified through heparin affinity columns (HiTrap Heparin HP, GE

Healthcare, Uppsala, Sweden) and concentrated using Amicon Ultra-4 centrifugal filters

(Millipore, Bedford, MA). For expression of shRNAs, we used a vector containing the U6

promoter upstream of the shRNA sequence and a chicken β-actin promoter to drive GFP

expression. The sequences are as following: Dnmt3a1-shRNA1:

GCAGACCAACATCGAATCCAT; Dnmt3a1-shRNA2:

GGGAGGATGATCGAAAGGAAGGAGA; Dnmt3a2-shRNA:

ACGGGCAGCTATTTACAGAGC; Nrp1-shRNA1: GGAAACCAAGAAGAAATATTA; Nrp1-

shRNA2: GGGAGAGGAAATCGGAGCTAA; Control-shRNA: ACTACCGTTGTTATAGGTG.

For temporally controlled knockdown of Dnmt3a1, we used a dual-component TetON-based

system. In detail, this system consisted of a driver plasmid that expresses the transactivator

(rtTA), the tetracycline repressor (TetR) and the fluorescent protein Kusabira Orange (KO)

that acts as an infection marker under the control of a neuron-specific promoter (hSynapsin).

The second construct contains GFP and a miR30-based shRNA targeting Dnmt3a1 under the

control of the tetracycline-responsive promoter (TRE). For the generation of this vector, the

plasmids pPRIME-CMV-GFP-FF3[34] (a gift from Stephen Elledge (Addgene plasmid # 11663;

http://n2t.net/addgene:11663;   RRID:Addgene_11663))   and   pAAV-PTRE-tight-hM3Dg-mCherry[35]   (a   gift   from   William   Wisden   [Addgene   plasmid   #   66795; http://n2t.net/addgene:66795; RRID:Addgene_66795]) were used. Specifically, the Dnmt3a1-shRNA1 sequence was inserted into the pPRIME vector and subsequently, the GFP-miR30-Dnmt3a1-shRNA expression cassette was subcloned into the vector containing the Tet-inducible promoter by replacement of the hM3Dq-mCherry insert. For the overexpression of Nrp1-HA or a control gene, we used a vector containing the neuron-specific promoter, hSynapsin, to constitutively express HA-tagged Nrp1 or LacZ, respectively. The infection rate, toxicity, viral titer, and knockdown efficiency for each batch of generated viruses were evaluated. The titer was matched to a final working concentration of around 1-2 $\times 10^{12}$ viral particles/ml.

Fluorescent labeling.

Deeply sedated mice were perfused intracardially with ice-cold PBS followed by 4% paraformaldehyde (Sigma-Aldrich, Munich, Germany). A post-fixation of the brains with 4% paraformaldehyde solution (Sigma-Aldrich, Munich, Germany) was done overnight at 4 °C. The following day, brains were put in a PBS solution with 30% sucrose and left for three days. Brain slices were cut at a thickness of 30 µm using a cryotome (CM1950, Leica). Slices were incubated in Hoechst 33258 (2 µg/ml, Serva, Heidelberg, Germany) for 5 min, mounted on glass slides and imaged using a 20x oil objective on a TCS SP8 confocal microscope (Leica Microsystems, Oberkochen, Germany).

Stereotaxic surgery.

rAAVs were injected into the dorsal hippocampus (dHPC) at the following coordinates relative to the bregma: -2 mm anteroposterior, ± 1.5 mm mediolateral, -1.7, -1.9 and -2.1 mm dorsoventral from bregma. Each spot received 500 nl of viral solution. Injections proceeded at a speed of 200 nl min$^{-1}$ through a 33G needle. Following injections at each individual site, the needle was left in place for 60 s. At the time of behavioural experiments, the experimenter was blind to the identity of the virus injected into each mouse. Behavioural experiments started 3 weeks after rAAV delivery.

Behavioural testing.

Before behavioural testing started, mice were habituated to the experimenter and behavioural room by handling for 3 consecutive days, 1 min/mouse. Different mouse cohorts were used to

test short-term (1h) or long-term memory (24h). The spatial object recognition test and
contextual fear conditioning were performed as previously described[3]. Briefly, in the object
recognition test, mice were habituated to an open arena (50 cm × 50 cm × 50 cm) with a visual
cue placed on the arena wall for 6 min, followed by a training session in which mice were
exposed to two objects for 6 min over a course of three trials with a 3 min intertrial interval.
During the testing session, which occurred 1h or 24h later, one object was moved to a new
location and exploration of objects was scored for 6 min. If in one experimental batch during
the spatial object recognition test control animals did not show a preference, due to day
effects, the whole set of animals was excluded from this analysis but was still included in
contextual fear conditioning data. Further, if an animal did not explore the objects during the
training for at least 2 seconds, that animal was excluded. One week later, contextual fear
conditioning (CFC) was performed. In CFC, mice were allowed to explore the conditioning
chamber (23 × 23 × 35 cm, TSE, Bad Homburg, Germany) for 148 s followed by a 2 s foot
shock (0.5 mA). They remained for additional 30 s in the conditioning chamber before returning
to their home cage (HC). The testing session consisted of exposing the animals to the
conditioning chamber for 5 min. For the temporally controlled knockdown of Dnmt3a1, mice
received intraperitoneal doxycycline hyclate (2.5 mg in 500 µL saline solution, 100mg/kg,
Sigma-Aldrich, Munich, Germany) 3 days prior to CFC.

Primary hippocampal cultures and pharmacological treatments.

Hippocampal cultures from newborn C57Bl/6N mice (Charles River, Sulzfeld, Germany) were
prepared and maintained as previously described[3]. In brief, mice hippocampi were dissociated
at P0 by papain digestion and plated onto tissue culture dishes coated with poly-D-lysine and
laminin (Sigma-Aldrich, Munich, Germany). The primary cultures were maintained for 8 days
in Neurobasal-A medium (Gibco™) supplemented with 1% rat serum (Biowest), 0.5 mM L-
glutamine (Sigma-Aldrich, Munich, Germany) and B27 (Gibco™), followed by incubation in
transfection media: salt-glucose-glycine solution (10 mM HEPES, pH 7.4, 114 mM NaCl,
26.1 mM NaHCO3, 5.3 mM KCl, 1 mM MgCl2, 2 mM CaCl2, 30 mM glucose, 1 mM glycine,
0.5 mM C3H3NaO3, and 0.001% phenol red) and phosphate-free Eagle's minimum essential
medium (9:1 v/v), supplemented with insulin (7.5 µg/ml), transferrin (7.5 µg/ml), and sodium
selenite (7.5 ng/ml) (ITS Liquid Media Supplement, Sigma-Aldrich, Munich, Germany) and
penicillin-streptomycin. rAAV infection occurred on day in vitro (DIV) 4. Experiments were
performed on DIV 9-11. To induce action potential bursting, cultures were treated with 50 µM
bicuculline (Alexis Biochemicals, Farmingdale, NY, USA). DNA transfection was performed
on DIV 8 using Invitrogen Lipofectamine 2000 (Invitrogen, CA, USA). The DNA (µg):
Lipofectamine (µl) ratio was 1.6:5 in 1 ml of medium. The Lipofectamine/DNA mixture was left

on the cells for 3 h before it was replaced with a transfection medium. Doxycycline hyclate (25 µM, Sigma-Aldrich, Munich, Germany) was introduced in the medium at DIV 8. To monitor doxycycline dependency, images of infected primary neurons with the TetOn-based miR30 system were acquired (Axio Vert.A1, Zeiss, Oberkochen, Germany).

qRT-PCR primer design.

The qRT-PCR primers were designed with Primer3 (https://primer3.ut.ee/) using either the RefSeq curated annotation or the GENCODE VM23 comprehensive transcript annotation, along with the GRCm38/mm10 mouse genome assembly. The specificity and amplicon product size of the primers were verified by BLAST search and *in silico* PCR (UCSC Genome Browser, mm10). Primer pair efficiencies and product melting curves were validated by qRT-PCR on serially diluted complementary DNA (cDNA) from primary mouse hippocampal cultures (refer to the "qRT-PCR" section). The list of all primers used in the study is provided in Sup Table 1.

qRT-PCR.

RNA from hippocampal cultures or mouse hippocampal tissue was extracted using the RNeasy Plus Mini Kit (Qiagen, Hilden, Germany) with extra DNase I digestion on the column, following the manufacturer's instructions. For isolation from mouse tissue, the brain was quickly isolated and sliced and the infected region (identified by GFP expression) isolated for RNA extraction. RNA was then transcribed into cDNA using the High-Capacity cDNA reverse-transcription kit (Applied Biosystems, Foster City, CA, USA). Quantitative reverse-transcription PCR (qRT-PCR) was performed on Step One Plus Real-Time PCR System (Applied Biosystems, Foster City, CA, USA) using TaqMan gene expression assays (Applied Biosystems, Foster City, CA, USA) or the Power SYBR Green PCR Master Mix (Applied Biosystems) assays. TaqMan gene expression assays were used for the following target genes: Dnmt1 (Mm00599763_m1), Dnmt3a1 (Mm00432870_m1), Dnmt3a2 (Mm00463987_m1), Dnmt3b (Mm00599800_m1) and the housekeeping gene Gusb (Mm00446953_m1). Power SYBR Green PCR Master mix was used for the remaining target genes. PCR reactions were run as technical triplicates in 10 µL reactions (96-well format) using 0.5 µM of each primer. 2 µL of diluted cDNA (about 1.25 ng) was added to each reaction. The thermal cycling was conducted with the following settings: a 10-minute incubation at 95 °C, 40 cycles of 10 seconds each at 95 °C, 60 °C, and 72 °C, followed by a 15-second incubation at 95 °C. Melt curves were generated by heating from 60 °C to 90 °C at a ramp

rate of 0.6 °C/min. Relative expression levels of each target transcript were determined by the

ΔΔCt method using beta-Actin mRNA levels as a reference[36].

Western Blotting.

Primary hippocampal neurons infected on day *in vitro* (DIV) 4 were lysed on DIV 10 in boiling

SDS sample buffer (160 mM Tris-HCl (pH 6.8), 4% SDS, 30% glycerol, 10 mM dithiothreitol,

and 0.02% bromophenol blue). In the case of western blotting of tissue samples, the dorsal

hippocampus was quickly dissected in ice-cold PBS and homogenized in RIPA buffer (150

mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris, pH 8.0)

supplemented with 1% protease inhibitor cocktail (Sigma-Aldrich, Munich, Germany). Protein

concentration was measured by Bradford assay and 20 μg of denatured protein (in Laemmli

buffer at 95°C for 5 min) was loaded in a 7.5% polyacrylamide gel and blotted onto a

nitrocellulose membrane (GE Healthcare, Buckinghamshire, UK). The membranes were

blocked in 5% milk in phosphate-buffered saline (PBS) with 0.01% Tween (PBST) and probed

with the following antibodies (diluted in 5% milk in PBST) overnight at 4 °C: α-Tubulin

(1:400000, Sigma t9026), α-Dnmt3a (1:5000, H-295, Santa Cruz, SC-20703), α-HA-tag

(1:7500, Covance, MMS101R). The next day, membranes were incubated with horseradish

peroxidase-conjugated secondary antibodies (1:5000 diluted in 5% milk in PBST) for 1h at

room temperature. Finally, they were analyzed using ChemiDoc™ Imaging System (Bio-Rad,

California, USA).

Immunocytochemistry.

Primary hippocampal neurons plated on coverslips were rinsed with PBS and fixed with a

prewarmed solution of 4% paraformaldehyde and 4% sucrose for 15 min at room temperature.

After permeabilizing the cells in ice-cold methanol for 6 min, blocking was performed with 10%

normal goat serum in PBS for 1 h at room temperature. Next, cells were incubated with primary

antibody α-Dnmt3a (1:500, H-295, Santa Cruz, SC-20703diluted in PBS containing 2% BSA,

0,1% Triton X-100) overnight at 4 °C, which continued with secondary antibody incubation

(1:500 goat anti-mouse Alexa488 [Life Technologies, Eugene, OR, USA]) diluted in PBS

containing 2% BSA, 0,1% Triton X-100) for 1 h at room temperature. Finally, coverslips were

treated with Hoechst 33258 (2 μg/ml, Serva, Heidelberg, Germany) for 5 min and mounted on

glass slides. Images were acquired with a 40x oil objective on a fluorescence microscope

(Leica Microsystems, Oberkochen, Germany).

Bulk RNA Sequencing.

Total RNA from mouse hippocampal cultures was isolated as above described. 500ng of total

RNA from each sample was used for bulk RNA-sequencing analysis. Data processing was

performed with R (version 3.6.3) and bioconductor (version 3.9) in Rstudio (version 1.1.463).

Quality control of clean sequencing reads was performed using FastQC (Babraham

Bioinformatics). Low-quality reads were removed using trim_galore (version 0.6.4). The

resulting reads were aligned to the mouse genome version GRCm38.p6 and counted using

kallisto version 0.46.1 [37]. The count data were transformed to log2-counts per million (logCPM)

using the voom-function from the limma package[38]. Differential expression analysis was

performed using the limma package in R. A false positive rate of $\alpha= 0.05$ with FDR correction

was taken as the level of significance. Volcano plots and heatmaps were created using ggplot2

package (version 2.2.1) and the complexHeatmap package (version 2.0.0)[39]. For enrichment

analysis, we used the fgsea[40], the enrichmentbrowser[41], and the Enrichr packages[42].

Statistical analysis.

Each data set was subjected to a normality test prior to further comparisons (Shapiro-Wilk

normality test; alpha = 0.05). For normally distributed data sets, a two-tailed unpaired

Student's t-test was performed to compare two groups. If more than two groups were analyzed

simultaneously, a one-way ANOVA was used followed by appropriate multiple comparison

*post hoc* tests to control for multiple comparisons as specified. In case of a non-Gaussian

distribution, two-tailed Mann-Whitney tests were used to compare two distinct groups, or a

Kruskal-Wallis test followed by Dunn´s *post hoc* test to compare more than two groups. The

sample size was determined based on similar experiments carried out in the past and the

literature. All plotted data represent mean ± SEM. Statistical analysis was performed using

GraphPad Prism 7 (GraphPad Software). All behavioural sessions were video recorded and

manually scored to determine the exploration of objects during training and testing phases or

freezing behaviour by an experimenter blind to the group identity. The TSE Systems Fear

Conditioning program was used to score the mean velocity during the training for CFC.

**Results**

**Dnmt3a1 is required for long-term memory formation**

In this study, we sought to investigate the role of DNA methyltransferase 3a1 (Dnmt3a1) in neuronal function and memory formation. We employed stereotaxic surgery and recombinant adeno-associated viruses (rAVV) to deliver a control shRNA sequence (Control-shRNA)[3] or two independent shRNAs to knockdown Dnmt3a1 into the adult (8 weeks) dorsal hippocampus (dHPC) of mice (Figure 1A, B). We used the rAAV1/2 serotype given its well established neuronal tropism[43]. To monitor viral expression upon infection, the viral constructs contained a GFP cassette under the control of the chicken-beta-actin promoter (Figure 1A, B). We confirmed knockdown efficiency of our shRNAs by transfection of primary hippocampal neurons *in vitro* and immunostaining against endogenous Dnmt3a1 (Sup Figure 1A). To further validate the knockdown efficiency *in vivo*, we performed qRT-PCR and western blot analysis of dHPC of mice infected with the viral constructs (Sup Figure 1B-G). We confirmed that both sequences selectively knockdown Dnmt3a1 without affecting other *Dnmts*. shRNA1 and shRNA2 sequences effectively reduced Dnmt3a1 expression at both the mRNA (Sup Figure 1B, C) and protein (Sup Figure 1D-G) levels. To evaluate a requirement for Dnmt3a1 in memory performance, we conducted hippocampus-dependent memory tests on mice three weeks after stereotaxic surgery (Figure 1C). We found that, when tested 24 hours after training in the spatial object recognition test, mice injected with either Dnmt3a1-shRNA1 or -shRNA2 exhibited no preference for the displaced object (Figure 1D). However, when evaluated 1 hour after training, mice in shRNA groups explored the displaced object for a length of time comparable to the control group (Figure 1F). Importantly, Dnmt3a1 knockdown did not affect the overall amount of time that mice spent exploring the objects during training (Sup Figure 1H, I). To assess further the role of hippocampal Dnmt3a1 in memory formation, we trained mice in contextual fear conditioning. Knockdown of Dnmt3a1 with either shRNA1 or shRNA2 impaired contextual fear memory response 24 hours after training (Figure 1E) without decreasing freezing levels compared to the Control-shRNA when tested 1 hour after training (Figure 1G). This decrease in freezing was not due to altered responsiveness to the shock administration during the training session (Sup Figure 1 J, K). These results show that Dnmt3a1 knockdown impairs long-term memory formation without affecting short-term memory.

The behavioural testing in this series of experiments was carried out three weeks following stereotaxic delivery of rAAVs to achieve high viral expression and, as a result, high knockdown efficiency. However, the persistent reduction of Dnmt3a1 levels may lead to network-wide changes that could impact learning and memory. Consequently, in order to more directly

assign a role for Dnmt3a1 in mature neurons during the process of memory formation, we
adopted a temporally controlled method that limited Dnmt3a1 knockdown to the time of
behavioural testing. We used a TetON-based system that consisted of two components: for
tight control of Tet-dependent transgene expression in neurons, one viral construct contained
the human synapsin (hSyn) promoter controlling the expression of the reverse tetracycline-
controlled transactivator (rtTA) together with the Tet repressor (TetR) and the fluorescent
protein Kusabira Orange (KO) that serves as an infection marker. In the second construct, the
Dnmt3a1-specific shRNA sequence 1 or the Control-shRNA was embedded in a microRNA-
30-based expression cassette[34], which is under the control of the tetracycline responsive
promoter (TRE) (Figure 2A-B). In the absence of doxycycline, TetR is bound to TRE inhibiting
its activity. Upon doxycycline administration, TetR releases from the TRE promoter allowing
the binding of the activator rtTA. Administration of doxycycline to hippocampal primary cultures
verified tight control of transgene expression via the Tet-On system (Sup Figure 2 A-D). The
expression of GFP after doxycycline administration increased over a time course of 72 hours
(Sup Figure 2B). Western blot analysis of Dnmt3a1 at the same time points revealed a
significant knockdown of Dnmt3a1 protein levels at 72 hours (Sup Figure 2C, D). Thus,
doxycycline was injected intraperitoneal 72 hours before behavioural testing (Figure 2C). We
found that knocking down hippocampal Dnmt3a1 around the time of fear conditioning reduced
freezing levels when mice were tested 24 hours after training (Figure 2D), however, mice froze
to a similar extent than the control mice when testing occurred 1 hour after training (Figure
2E). The responsiveness to the shock administration during the training session did not cause
this decrease in freezing (Sup Figure 2E). Taken together, these findings further support a
function of Dnmt3a1 during long-term memory formation and suggest that Dnmt3a1
participates in the regulation of learning-dependent gene expression required for long-term
memory.

**Dnmt3a1 regulates the expression of activity-dependent genes involved in synaptic**
**plasticity**

Next, to elucidate the molecular mechanism by which Dnmt3a1 regulates memory
consolidation we sought to identify activity-dependent genes regulated by Dnmt3a1. To this
end, we performed RNA sequencing of hippocampal neurons infected with Dnmt3a1-shRNA2
or a Control-shRNA after inducing neuronal activity by bicuculline treatment (Figure 3A, B,
Sup Figure 3A, B). In control conditions, we found that neuronal activity leads to the
downregulation of 1981 genes and upregulation of 1935 genes including established activity-
regulated genes, e.g., Arc, Fos, Npas4 (Figure 3C, Sup File 1).

To identify activity-regulated genes dependent on Dnmt3a1, we focused on genes that are

differentially expressed in Dnmt3a1 knockdown versus control conditions upon bicuculline

treatment. These included 1241 genes of which 436 were upregulated and 805 downregulated

(Figure 3D, Sup File 2). Notably, the expression of classical immediate early genes, such as

Arc, Fos, Npas4 was not altered in Dnmt3a1-shRNA conditions (Sup File 2), suggesting that

the identified changes are not a result of an overall disruption of neuronal responses. We

hypothesised that Dnmt3a1-regulated genes that are involved in memory formation should be

both a) activity-regulated and b) differentially expressed in response to Dnmt3a1 reduction.

Thus, we overlapped the activity-regulated (Figure 3C, Sup File 1) and the Dnmt3a1-regulated

gene sets (Figure 3D, Sup File 2) in order to find potential activity-regulated Dnmt3a1 target

genes. This intersection identified 327 candidates (Figure 3E, Sup Figure 3B, Sup File 3).

Gene ontology (GO) analysis of the 327 genes yielded a strong enrichment for terms related

to structural and functional plasticity, namely *"Regulation of neurotransmitter receptor activity"*,

*"Regulation of signaling receptor activity"* and *"Regulation of cell-substrate adhesion"* (Figure

3F). This suggests that genes downstream of Dnmt3a1 may play an important role in synaptic

plasticity and learning and memory. We confirmed some of the identified Dnmt3a1 target

genes in independent biological samples using qRT-PCR (Figure 3G, H). To further rule out

the possibility of off-target effects, this validation was performed using the two independent

shRNA sequences. We found that the expression of *Alkal2*, *Cacng5*, and *Nrp1* is regulated by

neuronal activity and reduced upon Dnmt3a1 depletion achieved by shRNA1 or shRNA2

(Figure 3G), whereas *Crhbp, Npy and Trpc6* failed to show a significant effect with one of the

shRNA sequences (Figure 3H). *Dnmt3a* encodes for two distinct isoforms Dnmt3a1 and

Dnmt3a2, which are differentially regulated by neuronal activity[22,23]. We further investigated

whether the two isoforms regulate distinct targets. To test this, we infected primary

hippocampal cultures with a previously validated shRNA to reduce Dnmt3a2 levels[23,24] (Sup

Figure 3C) and investigated the impact of this manipulation on the expression of the identified

Dnmt3a1-regulated genes. Reduction of Dnmt3a2 did not change the expression of *Alkal2*,

*Cacng5* or *Nrp1* neither in basal conditions nor upon neuronal activity (Sup Figure 3D). Taken

together these findings demonstrate that Dnmt3a1 is required for the regulation of a synaptic

plasticity-related transcriptional program and further suggest that the Dnmt3a2 isoform may

regulate a distinct gene pool.

**Neuropilin-1 is a Dnmt3a1 downstream target required for memory formation**

Next, we aimed at investigating whether Dnmt3a1 role in memory consolidation requires the

activity of identified target genes. Among the genes whose expression is controlled by

Dnmt3a1, we focused on neuronal genes which have been identified to be regulated by novel

environment exposure in the mouse hippocampus[44]. These included, among others, the

previously validated gene Neuropilin-1 (*Nrp1)* (Figure 3G, H). Nrp1 is known to build a complex

with Semaphorin 3a (Sema3A) and Plexin A4 and is implicated in signalling pathways

regulating neuronal morphology[31,45]. Although it has been demonstrated that this complex is

important for AMPA receptor trafficking[46], whether Nrp1 plays a role in hippocampus-

dependent memory formation is still unknown. We performed a shRNA-mediated loss-of-

function approach to assess the role of Nrp1 in memory formation. We generated two

independent shRNA constructs to reduce the expression of *Nrp1* (Figure 4A) and validated

their knockdown efficiency *in vivo* after stereotaxic delivery of the viral constructs or a Control-

shRNA into the dHPC. We confirmed that Nrp1-shRNA1 and -shRNA2 sequences led to a

robust reduction of Nrp1 mRNA levels (Sup Figure 4A, B). Mice were subjected to a

hippocampus-dependent behavioural memory task three weeks after viral delivery (Figure

4B). Knockdown of Nrp1 with either shRNA resulted in exploration of the displaced object at

chance levels (Figure 4C), as well as in decreased freezing responses in the contextual fear

conditioning test compared to mice injected with the Control-shRNA (Figure 4D). Notably,

Nrp1 knockdown did not affect the overall object exploration time (Sup Figure 4C) or the

responsiveness to the shock administration (Sup Figure 4D). In conclusion, this set of

experiments uncovered a novel function for Nrp1 in long-term memory formation.

**Dnmt3a1, but not Dnmt3a2, regulates long-term memory formation via its downstream**

**target Nrp1**

To evaluate a function of Nrp1 as a downstream effector of Dnmt3a1 during memory

formation, we performed a rescue experiment. To this end, we overexpressed HA-tagged

Nrp1 or a control protein (LacZ) under the regulation of hSyn promoter together with either

Dnmt3a1-shRNA2 or Control-shRNA sequences in the mouse hippocampus (Figure 5A, B).

We first confirmed successful overexpression of Nrp1 in the dHPC by western blot analysis

(Sup Figure 5A, B). After stereotaxic delivery of the constructs, mice underwent the spatial

object recognition task (Figure 5B). The different groups did not exhibit differences in the time

spent exploring the objects during the training session (Sup Figure 5C). As shown above, the

reduction of Dnmt3a1 levels caused memory impairments which are manifested by the

decreased preference for the displaced object (Figure 1D). Strikingly, this impairment was

rescued upon Nrp1 overexpression (Figure 5C). Mice that overexpressed Nrp1 and expressed

a control shRNA sequence showed memory performance similar to the control group (Figure

5C). Taken together, these results demonstrate that Nrp1 acts downstream of Dnmt3a1, is

required for memory formation and can per se rescue Dnmt3a1-dependent memory

impairments. Thus, indicating that Dnmt3a1 regulates memory formation, at least in part, via
its downstream target Nrp1.

Our gene expression analysis suggested that the isoforms Dnmt3a1 and Dnmt3a2 regulate
distinct genes. Specifically, we showed that Nrp1 expression is regulated by Dnmt3a1, but it
is not altered upon knockdown of Dnmt3a2 (Sup Figure 3D). Next, to investigate a specific
functional link between Dnmt3a1, Nrp1 and memory formation, we tested if Nrp1
overexpression rescues Dnmt3a2-knockdown-dependent memory impairments. We
overexpressed HA-tagged Nrp1 with a Dnmt3a2-specific shRNA sequence in the mouse
dorsal hippocampus and performed the spatial object recognition task in mice (Figure 5D, E).
Importantly, none of the conditions showed differences in overall time exploring the objects
during training (Sup Figure 5D). As previously described[23], knockdown of Dnmt3a2 impaired
spatial object recognition memory and remarkably, this impairment was not rescued upon
Nrp1 overexpression (Figure 5F). This finding shows that the rescue obtained in response to
Nrp1 overexpression is not a result of a generalized memory enhancement effect and further
supports that Nrp1 acts downstream of Dnmt3a1 but not Dnmt3a2. Thus, this data uncovers
a *Dnmt3a* isoform specific mechanism in memory formation.

**Discussion**

In this study we demonstrated a critical role for Dnmt3a1 in long-term memory formation by showing that acute knockdown of Dnmt3a1 in the adult hippocampus selectively impairs long-term memory formation and further identified activity-dependent genes regulated by Dnmt3a1 levels. Furthermore, our data suggests that despite a requirement for both Dnmt3a1 and Dnmt3a2 for memory formation, the two isoforms regulate this process via distinct mechanisms; Nrp1 overexpression rescued Dnmt3a1, but not Dnmt3a2, knockdown-driven impairments in memory formation. Taken together these findings have advanced our understanding of the requirement for distinct Dnmts in mnemonic processes as well as of the identity of downstream effector molecules.

Most studies that aimed at investigating the role of Dnmt3a in adult brain cognition have utilized heterozygous constitutive[47] or conditional knock-out mice[15,16,48]. These models delete *Dnmt3a* during critical prenatal and postnatal neurodevelopmental phases, which may confound the interpretation of the function of *Dnmt3a*-coded proteins in memory formation in the adult. Mitchnick and colleagues (2015) used small interfering RNAs to acutely disrupt *Dnmt3a* expression in the adult hippocampus and provided compelling evidence for a role of *Dnmt3a*-coding genes in long-term hippocampal memory formation[14]. In addition, our present and previous studies[23–25] have corroborated these findings and demonstrated that both Dnmt3a isoforms, Dnmt3a1 and Dnmt3a2, are required for memory formation in adult mature neurons. This study has further identified for the first time an activity-regulated genomic program regulated by Dnmt3a1 levels in hippocampal neurons. We found that the knockdown of Dnmt3a1 altered the expression of several genes. Even if counterintuitive at first, the reduction of Dnmt expression resulted not only in de-repression of gene expression, but also downregulation. Although these gene expression changes are likely a combination of direct effects on the altered genes and indirect mechanisms via the disruption of expression of other transcriptional regulatory processes, these findings corroborate previous studies in which a link between transcription activation and DNA methylation has been shown[10,28,49]. In our transcriptomic analysis we found that Dnmt3a1 regulates the expression of several genes with key roles in synaptic plasticity and memory. Namely, we found changes in the expression of *Npy*, *Cort*, *Trpc6*, which have been implicated in memory consolidation[50–52]. Upon the confirmation with two independent shRNA sequences, we focused on Nrp1 which has been shown to be regulated by enriched environment exposure[53] and within dentate gyrus memory engrams[44]. Besides the established function for the semaphorins and their receptors (Nrp and plexin family members) in neurodevelopment[32,54], only a few lines of evidence demonstrated their involvement in homeostatic and Hebbian forms of plasticity in the adult

hippocampus[31,46,55,56]. Notably, Nrp1 is localised at synapses of the adult rat hippocampus[56]

and secreted semaphorins acting on the sema-Nrp-plexin complex modulate synaptic

connectivity in granule cells of dentate gyrus and pyramidal neurons of CA1[56] as well as AMPA

receptor trafficking in CA3-CA1 synapses[46]. The present study adds to this body of literature

by demonstrating that Nrp1 is required for the formation of hippocampal long-term memory.

Intriguingly, we showed that overexpression of hippocampal Nrp1 rescued Dnmt3a1

knockdown-dependent memory impairments. Thus, our findings suggest that the regulation of

Nrp1-dependent mechanisms during memory consolidation is a mechanism via which

Dnmt3a1 contributes to memory formation.

This study uncovered a dissociation in the downstream processes regulated by the Dnmt3a

isoforms during hippocampal long-term memory formation. Notably, we showed that Nrp1

overexpression selectively rescued memory impairments promoted by the reduction of

Dnmt3a1, but not Dnmt3a2, levels in the adult hippocampus. Our previous work had already

demonstrated differential regulation of the two enzymes upon neuronal activity and learning.

In contrast to Dnmt3a1, Dnmt3a2 mRNA levels are induced in response to action potential

bursting and learning[23], cocaine administration[30] and an inflammatory response[29] in different

regions of the central nervous system. Despite the lack of regulation at the level of

transcription, Dnmt3a1 protein levels appear to be regulated upon neuronal activity[22]. It has

also been shown that Dnmt3a isoforms target unique genomic loci[21,57]. Collectively, these

findings indicate distinct regulatory functions. Specifically, during the formation of memory,

lack of redundancy is further supported by our studies that showed that impairments in long-

term memory formation promoted by the acute, reduced hippocampal expression of Dnmt3a1

or Dnmt3a2 are not compensated by the other isoform. Thus, the emerging picture is that the

distinct Dnmts are required for cognitive abilities through the regulation of complementary but

non-overlapping processes. The recruitment of Dnmts to DNA is thought to be guided by

chromatin marks, namely histone modifications[58] and non-coding RNA species that direct

gene-specific methylation patterns[59–63]. Notably, differential DNA recruitment between

Dnmt3a1 and Dnmt3a2 may be at least in part driven by distinct recognition of histone tail

marks[21,28,64,65]. Moreover, the N-terminus unique to Dnmt3a1 has been shown to confer

specific chromatin recognition properties to this isoform[28,65.] Future studies should elucidate

the mechanisms that dictate the regulation of distinct downstream targets for Dnmt3a1 and

Dnmt3a2 in response to neuronal activity.

Our findings highlight the complex and highly regulated role of distinct epigenetic regulators

in brain function. The relevance of investigating DNA methylation processes in the nervous

system is further underscored by the multiple lines of evidence that demonstrated that DNA

methylation dysregulation underlies several pathological conditions, such as

neurodevelopmental[66] and neurodegenerative[67] diseases as well as psychiatric conditions[68,69]
and chronic pain[70]. Therefore, the elucidation of the regulatory mechanisms controlled by the
various epigenetic factors is required to gain further understanding of the pathophysiological
mechanisms underlying these conditions.

**Acknowledgements**

We thank David V.C. Brito, Kübra Gülmez Karaca and Stefanos Loizou for their critical comments on the manuscript. Celia García-Vilela, Carmen Leibold, Stefanos Loizou, and Benjamin Zeuch for technical support. hSyn-LacZ construct was kindly gifted by Prof. Dr. Daniela Mauceri. This work was supported by the Deutsche Forschungsgemeinschaft (DFG) [grant numbers OL 437/1, OL 437/2, OL 437/3 and OL 437/4 to A.M.M.O.], the Chica and Heinz Schaller Foundation [fellowship and research award to A.M.M.O.], and the Joachim Herz Stiftung [Add-on Fellowships for Interdisciplinary Life Science to J.K.].

**Conflict of Interest**

The authors declare no conflicts of interest.

**Data availability**

The RNA-seq data analyzed in this study is publicly available at the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) with the accession number GSE232630.

**References**

1. Benito, E. & Barco, A. The Neuronal Activity-Driven Transcriptome. *Mol Neurobiol* **51**, 1071–1088 (2015).
2. Yap, E. L. & Greenberg, M. E. Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. *Neuron* vol. 100 Preprint at https://doi.org/10.1016/j.neuron.2018.10.013 (2018).
3. Brito, D. V. C., Kupke, J., Karaca, K. G., Zeuch, B. & Oliveira, A. M. M. Mimicking age-associated Gadd45γ dysregulation results in memory impairments in young adult mice. *Journal of Neuroscience* **40**, (2020).
4. Alberini, C. M. Transcription Factors in Long-Term Memory and Synaptic Plasticity. *Physiol Rev* **89**, 121–145 (2009).
5. Alberini, C. M. & Kandel, E. R. The Regulation of Transcription in Memory Consolidation. *Cold Spring Harb Perspect Biol* **7**, a021741 (2015).
6. Cholewa-Waclaw, J. *et al.* The Role of Epigenetic Mechanisms in the Regulation of Gene Expression in the Nervous System. *The Journal of Neuroscience* **36**, 11427–11434 (2016).
7. Day, J. J. & Sweatt, J. D. Epigenetic Mechanisms in Cognition. *Neuron* **70**, 813–829 (2011).
8. Oliveira, A. M. M. DNA methylation: a permissive mark in memory formation and maintenance. *Learning & Memory* **23**, 587–593 (2016).
9. Medrano-Fernández, A. & Barco, A. Nuclear organization and 3D chromatin architecture in cognition and neuropsychiatric disorders. *Mol Brain* **9**, 83 (2016).
10. Suzuki, M. M. & Bird, A. DNA methylation landscapes: provocative insights from epigenomics. *Nat Rev Genet* **9**, 465–476 (2008).
11. Halder, R. *et al.* DNA methylation changes in plasticity genes accompany the formation and maintenance of memory. *Nat Neurosci* **19**, (2015).
12. Guo, J. U. *et al.* Neuronal activity modifies the DNA methylation landscape in the adult brain. *Nat Neurosci* **14**, 1345–1351 (2011).
13. Miller, C. A. & Sweatt, J. D. Covalent Modification of DNA Regulates Memory Formation. *Neuron* **53**, 857–869 (2007).
14. Mitchnick, K. A., Creighton, S., O'Hara, M., Kalisch, B. E. & Winters, B. D. Differential contributions of *de novo* and maintenance DNA methyltransferases to object memory processing in the rat hippocampus and perirhinal cortex - a double dissociation. *European Journal of Neuroscience* **41**, 773–786 (2015).
15. Feng, J. *et al.* Dnmt1 and Dnmt3a maintain DNA methylation and regulate synaptic function in adult forebrain neurons. *Nat Neurosci* **13**, 423–430 (2010).
16. Morris, M. J., Adachi, M., Na, E. S. & Monteggia, L. M. Selective role for DNMT3a in learning and memory. *Neurobiol Learn Mem* **115**, 30–37 (2014).
17. Zocher, S. *et al. De novo* DNA methylation controls neuronal maturation during adult hippocampal neurogenesis. *EMBO J* **40**, (2021).
18. Tatton-Brown, K. *et al.* The Tatton-Brown-Rahman Syndrome: A clinical study of 55 individuals with de novo constitutive DNMT3A variants. *Wellcome Open Res* **3**, 46 (2018).
19. Tatton-Brown, K. *et al.* Mutations in the DNA methyltransferase gene DNMT3A cause an overgrowth syndrome with intellectual disability. *Nat Genet* **46**, 385–388 (2014).
20. Chen, T., Ueda, Y., Xie, S. & Li, E. A Novel Dnmt3a Isoform Produced from an Alternative Promoter Localizes to Euchromatin and Its Expression Correlates with

Activede Novo Methylation. *Journal of Biological Chemistry* **277**, 38746–38754 (2002).

21. Manzo, M. *et al.* Isoform-specific localization of DNMT3A regulates DNA methylation fidelity at bivalent CpG islands. *EMBO J* **36**, 3421–3434 (2017).

22. Bayraktar, G. *et al.* Synaptic control of DNA methylation involves activity-dependent degradation of DNMT3A1 in the nucleus. *Neuropsychopharmacology* **45**, (2020).

23. Oliveira, A. M. M., Hemstedt, T. J. & Bading, H. Rescue of aging-associated decline in Dnmt3a2 expression restores cognitive abilities. *Nat Neurosci* **15**, (2012).

24. Oliveira, A. M. M., Hemstedt, T. J., Freitag, H. E. & Bading, H. Dnmt3a2: A hub for enhancing cognitive functions. *Mol Psychiatry* **21**, (2016).

25. Gulmez Karaca, K. *et al.* Neuronal ensemble-specific DNA methylation strengthens engram stability. *Nat Commun* **11**, (2020).

26. Feng, J., Chang, H., Li, E. & Fan, G. Dynamic expression of de novo DNA methyltransferases Dnmt3a and Dnmt3b in the central nervous system. *J Neurosci Res* **79**, 734–746 (2005).

27. Guo, J. U. *et al.* Genome-wide antagonism between 5-hydroxymethylcytosine and DNA methylation in the adult mouse brain. *Front Biol (Beijing)* **9**, 66–74 (2014).

28. Gu, T. *et al.* The disordered N-terminal domain of DNMT3A recognizes H2AK119ub and is required for postnatal development. *Nat Genet* **54**, 625–636 (2022).

29. Oliveira, A. M. *et al.* Epigenetic control of hypersensitivity in chronic inflammatory pain by the de novo DNA methyltransferase Dnmt3a2. *Mol Pain* **15**, 174480691982746 (2019).

30. Cannella, N. *et al.* Dnmt3a2 in the Nucleus Accumbens Shell Is Required for Reinstatement of Cocaine Seeking. *The Journal of Neuroscience* **38**, 7516–7528 (2018).

31. Carulli, D., de Winter, F. & Verhaagen, J. Semaphorins in Adult Nervous System Plasticity and Disease. *Frontiers in Synaptic Neuroscience* vol. 13 Preprint at https://doi.org/10.3389/fnsyn.2021.672891 (2021).

32. Pasterkamp, R. J. Getting neural circuits into shape with semaphorins. *Nat Rev Neurosci* **13**, 605–618 (2012).

33. Gulmez Karaca, K., Brito, D. V. C., Kupke, J., Zeuch, B. & Oliveira, A. M. M. Engram reactivation during memory retrieval predicts long-term memory performance in aged mice. *Neurobiol Aging* **101**, 256–261 (2021).

34. Stegmeier, F., Hu, G., Rickles, R. J., Hannon, G. J. & Elledge, S. J. A lentiviral microRNA-based system for single-copy polymerase II-regulated RNA interference in mammalian cells. *Proc Natl Acad Sci U S A* **102**, (2005).

35. Zhang, Z. *et al.* Neuronal ensembles sufficient for recovery sleep and the sedative actions of α 2 adrenergic agonists. *Nat Neurosci* **18**, (2015).

36. Brito, D. V. C., Gulmez Karaca, K., Kupke, J., Frank, L. & Oliveira, A. M. M. MeCP2 gates spatial learning-induced alternative splicing events in the mouse hippocampus. *Mol Brain* **13**, 156 (2020).

37. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, (2016).

38. Ritchie, M. E. *et al.* Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, (2015).

39. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, (2016).

40. Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* 060012 (2021) doi:10.1101/060012.

757    41.    Geistlinger, L., Csaba, G. & Zimmer, R. Bioconductor's EnrichmentBrowser:
758           seamless navigation through combined results of set- & network-based
759           enrichment analysis. *BMC Bioinformatics* **17**, 45 (2016).

760    42.    Chen, E. Y. *et al.* Enrichr: Interactive and collaborative HTML5 gene list enrichment
761           analysis tool. *BMC Bioinformatics* **14**, (2013).

762    43.    Xu, R. *et al.* Quantitative comparison of expression with adeno-associated virus
763           (AAV-2) brain-specific gene cassettes. *Gene Ther* **8**, (2001).

764    44.    Jaeger, B. N. *et al.* A novel environment-evoked transcriptional signature predicts
765           reactivity in single dentate granule neurons. *Nat Commun* **9**, (2018).

766    45.    Simonetti, M. *et al.* The impact of Semaphorin 4C/Plexin-B2 signaling on fear
767           memory via remodeling of neuronal and synaptic morphology. *Mol Psychiatry* **26**,
768           (2021).

769    46.    Jitsuki-Takahashi, A. *et al.* Activity-induced secretion of semaphorin 3A mediates
770           learning. *European Journal of Neuroscience* **53**, (2021).

771    47.    Christian, D. L. *et al.* DNMT3A Haploinsufficiency Results in Behavioral Deficits and
772           Global Epigenomic Dysregulation Shared across Neurodevelopmental Disorders. *Cell
773           Rep* **33**, 108416 (2020).

774    48.    Li, J. *et al.* Dnmt3a knockout in excitatory neurons impairs postnatal synapse
775           maturation and increases the repressive histone modification H3K27me3. *Elife* **11**,
776           (2022).

777    49.    Wu, H. *et al.* Dnmt3a-Dependent Nonpromoter DNA Methylation Facilitates
778           Transcription of Neurogenic Genes. *Science (1979)* **329**, 444–448 (2010).

779    50.    Jiang, J. *et al.* Intrahippocampal injection of Cortistatin-14 impairs recognition
780           memory consolidation in mice through activation of sst 2 , ghrelin and GABA A/B
781           receptors. *Brain Res* **1666**, 38–47 (2017).

782    51.    Xie, R. *et al.* AAV Delivery of shRNA Against TRPC6 in Mouse Hippocampus
783           Impairs Cognitive Function. *Front Cell Dev Biol* **9**, (2021).

784    52.    Gøtzsche, C. R. & Woldbye, D. P. D. The role of NPY in learning and memory.
785           *Neuropeptides* **55**, 79–89 (2016).

786    53.    Cao, L. *et al.* VEGF links hippocampal activity with neurogenesis, learning and
787           memory. *Nat Genet* **36**, 827–835 (2004).

788    54.    Kolodkin, A. L. & Tessier-Lavigne, M. Mechanisms and Molecules of Neuronal
789           Wiring: A Primer. *Cold Spring Harb Perspect Biol* **3**, a001727–a001727 (2011).

790    55.    Wang, Q. *et al.* Neuropilin-2/PlexinA3 Receptors Associate with GluA1 and Mediate
791           Sema3F-Dependent Homeostatic Scaling in Cortical Neurons. *Neuron* **96**, 1084-
792           1098.e7 (2017).

793    56.    Sahay, A. *et al.* Secreted Semaphorins Modulate Synaptic Transmission in the Adult
794           Hippocampus. *The Journal of Neuroscience* **25**, 3613–3620 (2005).

795    57.    Sae-Lee, C. *et al.* Genomic targets and selective inhibition of DNA methyltransferase
796           isoforms. *Clin Epigenetics* **14**, 103 (2022).

797    58.    Jurkowska, R. Z. & Jeltsch, A. Enzymology of Mammalian DNA Methyltransferases.
798           in 87–122 (2016). doi:10.1007/978-3-319-43624-1_5.

799    59.    Holz-Schietinger, C. & Reich, N. O. RNA modulation of the human DNA
800           methyltransferase 3A. *Nucleic Acids Res* **40**, 8550–8557 (2012).

801    60.    Mohammad, F. *et al.* Long noncoding RNA-mediated maintenance of DNA
802           methylation and transcriptional gene silencing. *Development* **139**, 2792–2803 (2012).

803    61.    Watanabe, T. *et al.* Role for piRNAs and Noncoding RNA in de Novo DNA
804           Methylation of the Imprinted Mouse *Rasgrf1* Locus. *Science (1979)* **332**, 848–852
805           (2011).

806    62.    Di Ruscio, A. *et al.* DNMT1-interacting RNAs block gene-specific DNA methylation.
807           *Nature* **503**, 371–376 (2013).
808    63.    Savell, K. E. *et al.* Extra-coding RNAs regulate neuronal DNA methylation dynamics.
809           *Nat Commun* **7**, 12091 (2016).
810    64.    Ooi, S. K. T. *et al.* DNMT3L connects unmethylated lysine 4 of histone H3 to de novo
811           methylation of DNA. *Nature* **448**, 714–717 (2007).
812    65.    Suetake, I. *et al.* Characterization of DNA-binding activity in the N-terminal domain
813           of the DNA methyltransferase Dnmt3a. *Biochemical Journal* **437**, 141–148 (2011).
814    66.    Pohodich, A. E. & Zoghbi, H. Y. Rett syndrome: disruption of epigenetic control of
815           postnatal neurological functions. *Hum Mol Genet* **24**, R10–R16 (2015).
816    67.    De Jager, P. L. *et al.* Alzheimer's disease: early alterations in brain DNA methylation
817           at ANK1, BIN1, RHBDF2 and other loci. *Nat Neurosci* **17**, 1156–1163 (2014).
818    68.    Feng, J. & Nestler, E. J. Epigenetic mechanisms of drug addiction. *Curr Opin*
819           *Neurobiol* **23**, 521–528 (2013).
820    69.    Nestler, E. J., Peña, C. J., Kundakovic, M., Mitchell, A. & Akbarian, S. Epigenetic
821           Basis of Mental Illness. *The Neuroscientist* **22**, 447–463 (2016).
822    70.    Denk, F. & McMahon, S. B. Chronic Pain: Emerging Evidence for the Involvement of
823           Epigenetics. *Neuron* **73**, 435–444 (2012).

**Figure 1: Reduced hippocampal Dnmt3a1 levels impair memory formation.** (**A**) Schematic representation of viral constructs. (**B**) Representative images of the dorsal hippocampus of mice infected with Control-shRNA, Dnmt3a1-shRNA1 or Dnmt3a1-shRNA2. Scale bar: 100 μm. (**C**) Schematic representation of experimental design. (**D, E**) Long-term (D) or short-term (E) spatial object recognition memory of mice expressing Control-shRNA (n=7-12), Dnmt3a1-shRNA1 (n=5-10) or Dnmt3a1-shRNA2 (n=9-12). Dashed lines represent equal preference for either object (chance exploration). *p<0.05, **p<0.01 by two-tailed, unpaired t-test. (**F, G**) Long-term (F) or short-term (G) contextual fear memory of mice expressing Control-shRNA (n=9-11), Dnmt3a1-shRNA1 (n=8-11) or Dnmt3a1-shRNA2 (n=11-13). *p<0.05 by two-tailed, unpaired t-test.

**Figure 2: Temporally-restricted Dnmt3a1 knockdown impairs memory consolidation.**
(**A**) Schematic representation of the viral constructs encoding the TetON-based system. The driver construct expresses under the hSyn promoter the Tet repressor (TetR), the reverse tetracycline-controlled transactivator (rtTA) and the infection marker Kusabira Orange (mKO). The Tet response element (TRE)-dependent construct contains the TRE promoter driving the miR30-based shRNA cassettes. In the absence of doxycycline, TRE is inhibited by TetR which results in suppression of the miR30 system. In the presence of doxycycline, TetR loses its affinity thus enabling rtTA to bind to TRE and activate the expression of the miR30 system. (**B**) Representative images of the dHPC of mice infected with rAAVs expressing the miR30-Control shRNA or miR30- Dnmt3a1-shRNA2 sequence. Scale bar: 100 μm. (**C**) Schematic representation of experimental design. (**D, E**) Long-term (D) or short-term (E) contextual fear memory of mice expressing miR30-Control shRNA (n=12-17) or miR30-Dnmt3a1 shRNA (n=10-14). *p<0.05, by two-tailed, unpaired t-test.

**Figure 3: Dnmt3a1 regulates transcription of genes involved in synaptic plasticity processes.** (**A**) Schematic representation of viral constructs. (**B**) Experimental design used to identify activity-regulated genes whose expression is altered upon Dnmt3a1 reduction. DIV: Day *in-vitro*. (**C**) Volcano plot of differentially expressed genes (DEGs) in control condition (infected with rAAVs expressing Control-shRNA) in response to neuronal activity via Bicuculline treatment; n=4 independent neuronal preparation per condition. Log2 fold change cut-off: ± 0.5; adjusted p-value cut-off: 0.05. (**D**) Volcano plot of DEGs upon infection with Control-shRNA or Dnmt3a1-shRNA2 in stimulated condition (4h Bicuculline); n=4 independent neuronal preparation per condition. Log2 fold change cut-off: ± 0.5; adjusted p-value cut-off: 0.05. (**E**) Venn-Diagram of activity-regulated DEGs (C) and Dnmt3a1-regulated DEGs (D). (**F**)

GO-Term analysis of overlapping genes between activity-regulated and Dnmt3a1-regulated DEGs. Dot plot illustrates Top 10 GO term enrichment of biological processes. (**G**) qRT-PCR analysis of Alkal2, Cacng5, Nrp1 expression levels in hippocampal cultures infected with Control–shRNA or Dnmt3a1-shRNA1 or -shRNA2 and stimulated 4h with Bicuculline. Expression levels were normalised to the uninfected controls in baseline conditions (dashed line); (n=8-9 independent neuronal cultures). *p<0.05, ***p≤0.001, ****p≤0.0001 by one-way ANOVA test followed by Sidak's multiple comparisons test. (**H**) qRT-PCR analysis of Crhbp, Npy, Trpc6 expression levels in hippocampal-cultured cells infected with Control–shRNA or Dnmt3a1-shRNA1 or -shRNA2 and stimulated 4h with Bicuculline. Expression levels were normalised to the uninfected controls in baseline conditions (dashed line); (n=7-9 independent neuronal cultures). *p<0.05, **p≤0.01, ns: not significant by one-way ANOVA test followed by Sidak's multiple comparisons test.

**Figure 4: Hippocampal Neuropilin-1 is required for memory formation.** (**A**) Schematic
representation of viral constructs. (**B**) Schematic representation of experimental design. (**C**)
Long-term spatial object recognition memory of mice expressing Control-shRNA (n=12), Nrp1-
shRNA1 (n=10) or Nrp1-shRNA2 (n=10). Dashed lines represent equal preference for either
object (chance exploration). *p<0.05 by one-way ANOVA test followed by Dunnett's multiple
comparisons test. (**D**) Long-term contextual fear memory of mice infected with rAAVs
expressing Control-shRNA (n=11), Nrp1-shRNA1 (n=12) or Nrp11-shRNA2 (n=10). *p<0.05
by one-way ANOVA test followed by Dunnett's multiple comparisons test. ##p≤0.01 by Kruskal-
Wallis test followed by Dunn's multiple comparisons test.

**Figure 5: Neuropilin-1 rescues Dnmt3a1-, but not Dnmt3a2-, dependent memory impairments.** (**A**) Schematic representation of viral constructs. (**B**) Schematic representation of experimental design. (**C**) Long-term spatial object recognition memory of mice expressing Control-shRNA + LacZ (n=13), Dnmt3a1-shRNA2 + LacZ (n=13), Dnmt3a1-shRNA2 + Nrp1-HA (n=12) or Control-shRNA + Nrp1-HA (n=12). Dashed lines represent equal preference for either object (chance exploration). **p≤0.01, ***p≤0.001 by one-way ANOVA test followed by Sidak's multiple comparisons test. (**D**) Schematic representation of viral constructs. (**E**) Schematic representation of experimental design. (**F**) Long-term spatial object recognition memory mice expressing Control-shRNA + LacZ (n=11), Dnmt3a2-shRNA + LacZ (n=12) or Dnmt3a2-shRNA + Nrp1-HA (n=12). Dashed lines represent equal preference for either object

928    (chance exploration). **p≤0.01, ***p≤0.001 by one-way ANOVA test followed by Sidak's

929    multiple comparisons test.

**Sup. Table 1. List of primers used for qRT-PCR.**

| Name | | Sequence |
|---|---|---|
| **Actin** | 5' Primer | TATCCTGACCCTGAAGTACC |
| | 3' Primer | CTCGGTGAGCAGCACAGGG |
| **Alkal2** | 5' Primer | TGAAACATCTCACAGGTCCTCT |
| | 3' Primer | GTGCAGTCTCTCGTGTTGTG |
| **Cacng5** | 5' Primer | CCCTCGTCAGCCTCTTCTTC |
| | 3' Primer | ACCACAAGAGAGAGGCCAGA |
| **Crhbp** | 5' Primer | CTGAAGGTATTTGATGGTTGGA |
| | 3' Primer | TCTTCATAGTGGGCAGAGGG |
| **Nrp1** | 5' Primer | GCAAGACTCGAATCCTCCC |
| | 3' Primer | CCAATGTGAGGGCCAACTTC |
| **Npy** | 5' Primer | CGTGTGTTTGGGCATTCTGG |
| | 3' Primer | TGCCATATCTCTGTCTGGTGA |
| **Trpc6** | 5' Primer | CCAGCTTCCGGGGTAATGAA |
| | 3' Primer | ACATGTATGCTGGTCCTCGA |

**Sup. Figure 1: Validation of Dnmt3a1 shRNA-dependent knock down and behavioural control parameters.** (**A**) Representative image of primary hippocampal cultures transfected with Dnmt3a1-shRNA1 or Dnmt3a1-shRNA2 (green) and immunostained against endogenous Dnmt3a levels (red). Scale bar: 50 μm. (**B, C**) Quantitative reverse-transcription PCR (qRT-PCR) analysis of the expression of DNA methyltransferases (Dnmt) in the dHPC of mice infected with rAAVs against Control-shRNA (n=4-5) or either (B) Dnmt3a1-shRNA1 (n=4) or (C) Dnmt3a1-shRNA2 (n=4). Dashed lines represent normalised expression levels to Control-shRNA. ***$p \leq 0.001$ by two-tailed, unpaired t-test. (**D, E**) Representative immunoblot scans of the expression of Dnmt3a1 or Tubulin in the dHPC of mice infected with rAAVs against Control-shRNA or either (D) Dnmt3a1-shRNA1 or (E) Dnmt3a1-shRNA2. (**F, G**) Immunoblot quantification of Dnmt3a1 protein levels in the dHPC of mice infected with rAAVs against Control-shRNA (n= 4-5) or either (F) Dnmt3a1-shRNA1 (n=4) or (G) Dnmt3a1-shRNA2 (n=4). Dashed lines represent normalised Dnmt3a1 to Tubulin expression of control group. *$p < 0.05$, ***$p \leq 0.001$ by two-tailed, unpaired t-test. (**H, I**) Total object exploration time of mice infected with rAAVs against Control-shRNA (n=9-12) or either (H) Dnmt3a1-shRNA1 (n=10) or (I) Dnmt3a1-shRNA2 (n=12) during the three sessions of spatial object recognition training. No significant changes by two-tailed, unpaired t-test. (**J, K**) Mean velocities during CFC training pre-shock (Pre-Shock) or during shock (Shock) administration of the mice expressing rAAVs against Control-shRNA (n=11-13) or either (J) Dnmt3a1-shRNA1 (n=11) or (K) Dnmt3a1-shRNA2 (n=11). No significant changes by two-tailed, unpaired t-test.

**Sup. Figure 2: Validation of mir30-based Dnmt3a1 shRNA-dependent knock down and behavioural control parameters.** (**A**) Experimental design used to identify the time-point of knock-down using the TetON-based miR30 system. DIV: Day *in-vitro*. (**B**) Representative images of primary hippocampal cells infected with miR30-Control or miR30-Dnmt3a1-shRNA2 sequence. Cultures were treated with doxycycline for 0h, 24h, 48, or 72h. (**C**) Representative immunoblot scans of the expression of Dnmt3a1 and Tubulin of primary hippocampal cultures infected with miR30-Control or miR30- Dnmt3a1-shRNA2 sequence and treated with doxycycline for either 0h, 24h, 48h or 72h. (**D**) Immunoblot quantification of Dnmt3a1 protein levels of primary hippocampal neurons infected with miR30-Control or miR30- Dnmt3a1-shRNA2 sequence and treated with doxycycline for either 0h, 24h, 48h or 72h. Expression levels were normalised to the uninfected controls (dashed line); (n=6 independent neuronal cultures). **$p \leq 0.01$, ns: not significant by two-tailed, unpaired t-test. NS: not significant by Mann-Whitney test. (**E**) Mean velocities during CFC training pre-shock (Pre-Shock) or during shock (Shock) administration of the mice expressing miR30-Control (n=8) or miR30-Dnmt3a1 (n=8) shRNAs. No significant changes by two-tailed, unpaired t-test.

**Sup. Figure 3: PCA analysis and heat map of RNA sequencing data and effect of Dnm3a2 levels on the expression of selected genes.** (**A**) Principle-component analysis (PCA) of DEGs in primary hippocampal cultures infected with Control-shRNA or Dnmt3a1-shRNA2, that were either stimulated with Bicuculline for 4h or unstimulated; (n=4 independent neuronal culture preparations). (**B**) Heatmap of activity-regulated Dnmt3a1-dependent genes (Figure 3E) (n=4 independent neuronal cell preparations). Hippocampal cultures infected with rAAVs expressing Control-shRNA or Dnmt3a1-shRNA2, that were either stimulated with Bicuculline for 4h or unstimulated. (**C**) Schematic representation of viral constructs. (**D**) qRT-PCR analysis of Alkal2, Cacng5 and Nrp1 expression levels in hippocampal-cultured cells infected with Control–shRNA or Dnmt3a2-shRNA and stimulated 4h with Bicuculline. Expression levels were normalised to the uninfected controls in baseline conditions (dashed line); (n=6 independent neuronal cultures). No significant differences between Control-shRNA and Dnmt3a2-shRNA by two-tailed, unpaired t-test.

**Sup. Figure 4: Validation of Nrp1 shRNA-dependent knock down and behavioural control parameters. (A, B)** qRT-PCR analysis of Nrp1 expression in dHPC of mice infected with rAAVs against Control-shRNA (n=6-9), (A) Nrp1-shRNA1 (n=6) or (B) Nrp11-shRNA2 (n=9). Dashed lines represent normalised expression levels to Control-shRNA. ***p≤0.001, ***p≤0.0001, by two-tailed, unpaired t-test. (**C**) Total object exploration time of mice infected with rAAVs against either Control-shRNA (n=12) or Nrp1-shRNA1 (n=12) or Nrp1-shRNA2 (n=11) during the three sessions of spatial object recognition training. No significant changes between Control shRNA and either Nrp1-shRNA1 or Nrp1-shRNA2 by one-way ANOVA test followed by Dunnett's multiple comparisons test. (**D**) Mean velocities during CFC training pre-shock (Pre-Shock) or during shock (Shock) administration of the mice infected with either Control-shRNA (n=12) or Nrp1-shRNA1 (n=12) or Nrp1-shRNA2 (n=11). No significant changes between Control shRNA and either Nrp1-shRNA1 or Nrp1-shRNA2 by one-way ANOVA test followed by Dunnett's multiple comparisons test.

**Sup. Figure 5: Validation of Nrp1 overexpression and behavioural control parameters.** (**A**) Representative immunoblot scans of the expression of Dnmt3a1, HA-tagged Nrp1 or HA-tagged LacZ and Tubulin in the dHPC of mice infected with rAAVs as indicated. (**B**) Immunoblot quantification of Dnmt3a1 protein levels in the dHPC of mice infected with rAAVs against Control-shRNA (n= 6) or Dnmt3a1-shRNA2 (n=6). Dashed lines represent normalised Dnmt3a1 to Tubulin expression of control group. **p≤0.01 by two-tailed, unpaired t-test. (**C**) Total object exploration time of mice infected with Control-shRNA + LacZ (n=13), Dnmt3a1-shRNA2 + LacZ (n=13), Dnmt3a1-shRNA2 + Nrp1-HA (n=13) or Control-shRNA + Nrp1-HA (n=13) during the three sessions of spatial object recognition training. No significant changes by one-way ANOVA test followed by Sidak's multiple comparisons test. (**D**) Total object exploration time of mice infected with Control-shRNA + LacZ (n=11), Dnmt3a2-shRNA + LacZ (n=12), Dnmt3a2-shRNA + Nrp1-HA (n=12) during three sessions of spatial object recognition training. No significant changes by one-way ANOVA test followed by Sidak's multiple comparisons test.