Cell type-specific *cis*-regulatory divergence in gene expression and chromatin accessibility revealed by

human-chimpanzee hybrid cells

Ban Wang[1,*], Alexander L. Starr[1,*], Hunter B. Fraser[1,2]

*Indicates equal contribution

1. Department of Biology, Stanford University, Stanford, CA, USA

2. Correspondence: hbfraser@stanford.edu

Abstract:

Although gene expression divergence has long been postulated to be the primary driver of human

evolution, identifying the genes and genetic variants underlying uniquely human traits has proven to be

quite challenging. Theory suggests that cell type-specific *cis*-regulatory variants may fuel evolutionary

adaptation due to the specificity of their effects. These variants can precisely tune the expression of a

single gene in a single cell type, avoiding the potentially deleterious consequences of *trans*-acting

changes and non-cell type-specific changes that can impact many genes and cell types, respectively. It

has recently become possible to quantify human-specific *cis*-acting regulatory divergence by measuring

allele-specific expression in human-chimpanzee hybrid cells—the product of fusing induced pluripotent

stem (iPS) cells of each species *in vitro*. However, these *cis*-regulatory changes have only been explored

in a limited number of tissues and cell types. Here, we quantify human-chimpanzee *cis*-regulatory

divergence in gene expression and chromatin accessibility across six cell types, enabling the

identification of highly cell type-specific *cis*-regulatory changes. We find that cell type-specific genes and

regulatory elements evolve faster than those shared across cell types, suggesting an important role for

genes with cell type-specific expression in human evolution. Furthermore, we identify several instances

of lineage-specific natural selection that may have played key roles in specific cell types, such as

coordinated changes in the *cis*-regulation of dozens of genes involved in neuronal firing in motor

neurons. Finally, using novel metrics and a machine learning model, we identify genetic variants that

likely alter chromatin accessibility and transcription factor binding, leading to neuron-specific changes in

the expression of the neurodevelopmentally important genes *FABP7* and *GAD1*. Overall, our results

demonstrate that integrative analysis of *cis*-regulatory divergence in chromatin accessibility and gene

expression across cell types is a promising approach to identify the specific genes and genetic variants

that make us human.

Introduction:

In the past few million years, humans have evolved a multitude of unique phenotypes (Shave et al.,

2019; Vanderhaeghen & Polleux, 2023). For example, our cardiovascular system has evolved to enable

extended periods of physical exertion and the unique aspects of our nervous system enable human

language and toolmaking (Shave et al., 2019; Vanderhaeghen & Polleux, 2023). Previous research

suggests that much of human adaptation may be caused by changes in gene expression (Fraser, 2013;

King & Wilson, 1975; Reilly & Noonan, 2016; Romero et al., 2012). To catalog these changes, studies

have compared gene expression in post-mortem tissues of humans and our closest living relatives,

chimpanzees (Blake et al., 2020; Kelley & Gilad, 2020; Ma et al., 2022). Although thousands of

differentially expressed genes have been identified in post-mortem samples, it is generally not possible

to determine whether genetic differences cause a gene to be differentially expressed. In post-mortem

studies, genetic differences cannot be disentangled from sources of variation such as differences in diet,

environment, cell type abundances, age, post-mortem interval, and other confounding factors. In

addition, for many traits the relevant gene expression differences may be specific to early development,

but it is impossible to study fetal development *in vivo* in non-human great apes due to both ethical and

technical difficulties. To circumvent these issues, several groups have used great ape iPS cells to study

differences in gene expression in cell types present in early development (Benito-Kwiecinski et al., 2021;

Field et al., 2019; Kanton et al., 2019; Pavlovic et al., 2018). While the use of iPS cells addresses many of

the confounding factors present in post-mortem comparisons, they also introduce new issues such as

interspecies differences in iPS cell differentiation kinetics, efficiency, and maturation. Overall, it remains

tremendously challenging to identify human-specific changes in gene expression, which limits our ability

to link expression differences to either phenotypic differences or natural selection on the human

lineage.

One particularly powerful means of studying the evolution of gene expression is through the

measurement of allele specific expression (ASE) in hybrids between two species (Combs et al., 2018;

Fraser, 2011; Hu et al., 2022; Mack & Nachman, 2017; Wittkopp et al., 2004). This approach has the

advantage of eliminating many confounding factors inherent to interspecies comparisons, including

differences in cell type composition, environmental factors, developmental stage, and response to

differentiation protocols. Because the trans-acting environments of the two alleles in a hybrid are

identical, ASE has the additional benefit of reflecting only *cis*-regulatory changes, which are thought to

be less pleiotropic and more likely to drive evolutionary adaptation than broader *trans*-acting changes

(Agoglia et al., 2021; Gokhman et al., 2021; Prud'homme et al., 2007; Wittkopp & Kalay, 2012).

Furthermore, ASE enables the use of powerful methods that can detect lineage-specific natural selection

and, as a result, contribute to our understanding of the selective pressures that have shaped the

evolution of a wide variety of species (Fraser, 2011). Until recently, it has not been possible to

disentangle *cis*- and *trans*-acting changes fixed in the human lineage since humans cannot hybridize with

any other species. However, the development of human-chimpanzee hybrid iPS cells enables

measurement of ASE in a wide variety of tissues and developmental contexts (Agoglia et al., 2021;

Gokhman et al., 2021; Song et al., 2021). This provides an effective platform to investigate general

principles of hominid gene expression evolution, detect lineage-specific selection, and identify candidate

gene expression changes underlying human-specific traits.

While gene expression differences between humans and chimpanzees are well-studied, there has been

less focus on epigenetic changes, many of which are likely to underlie differences in gene expression

(García-Pérez et al., 2021; Kozlenkov et al., 2020; Netherlands Brain Bank et al., 2016; Trizzino et al.,

2017). Furthermore, these studies, regardless of whether they utilize post-mortem tissues or cell lines,

are subject to the same confounding factors mentioned above. Analogous to ASE, one can use the assay

for transposase accessible chromatin using sequencing (ATAC-seq) in interspecies hybrids to measure

allele-specific chromatin accessibility (ASCA) (Buenrostro et al., 2013; Corces et al., 2017; Liang et al.,

2021; S. Zhang et al., 2020). As with ASE, ASCA is unaffected by many confounders inherent to between-

species comparisons and only measures *cis*-regulatory divergence. Perhaps most importantly, ASCA can

implicate specific regulatory elements that likely underlie gene expression differences. These regulatory

elements can then be more closely studied to identify the likely causal genetic variants and the

molecular mechanisms by which those variants alter gene expression.

Here, we generated RNA-seq and ATAC-seq data from six human-chimpanzee hybrid iPS cell-derived cell

types and quantified ASE and ASCA. Using this dataset, we identified thousands of genes and *cis*-

regulatory elements showing cell type-specific ASE and ASCA. We found that cell type-specific genes and

cis-regulatory elements are more likely to have divergent expression and accessibility than their more

broadly expressed/accessible counterparts. Furthermore, we provide evidence for polygenic selection

on the expression level of genes associated with physiologically relevant gene sets including sodium

channels and syntaxin-binding proteins in motor neurons. Finally, we use newly developed metrics and

machine learning algorithms to link cell type-specific differences in chromatin accessibility and gene

expression and identify putative causal mutations underlying these differences. Using this pipeline we

identified motor neuron-specific increases in promoter chromatin accessibility and gene expression for

FABP7, which plays a key role in neurodevelopment but is not well-studied in neurons. In addition, we

focus on a human-accelerated region (HAR) near the promoter of *GAD1*. While this region is accessible

in all cell types, both the accessibility of the HAR and the expression of *GAD1* are only chimpanzee-

biased in motor neurons. Analysis of scRNA-seq from human and chimpanzee brain organoids showed

that increased expression of *GAD1* also occurs in ventral forebrain inhibitory neurons. Overall, this study

provides insight into the evolution of gene expression in great apes as well as a resource that will inform

functional dissection of human-specific molecular changes.

Results

Cis-regulatory divergence of gene expression in six cell types is largely cell type-specific or shared

across all cell types

To measure genome-wide cis-regulatory divergence in gene expression, we performed RNA-seq on six

cell types derived from human-chimpanzee hybrid iPS cells (Fig. 1a). The cell types profiled were from six

diverse developmental lineages including the motor neuron (MN), cardiomyocyte (CM), hepatocyte

progenitor (HP), pancreatic progenitor (PP), skeletal myocyte (SKM), and retinal pigment epithelium

(RPE) lineages. These represent all three germ layers and a variety of organs (Fig. 1a). It is worth noting

that these differentiations do not necessarily lead to a pure population of cells, but rather a population

of cells with different levels of maturity along a particular developmental lineage. For example, the SKM

population likely contains fully differentiated muscle fibers as well as a small population of proliferating

satellite cells; for clarity we refer to this as the SKM cell type. As these different cell types are not shared

between tissues, we use cell type-specific and tissue-specific interchangeably throughout the

manuscript.

Two independently generated hybrid lines were differentiated for each cell type and at least two

biological replicates per hybrid line per cell type were collected (see Methods). Each cell type was

sequenced to an average depth of 134 million paired-end reads (Supp Fig. 1). We used a computational

pipeline to quantify ASE adapted from the pipeline introduced by Agoglia et al (Agoglia et al., 2021).

Briefly, we computed ASE by mapping reads to both the human and chimpanzee genomes, correcting

for mapping bias, and assigning reads to the human or chimpanzee genome if a read contained one or

more human-chimpanzee single nucleotide differences (see Methods).

As expected, the samples clustered predominantly by cell type (Fig. 1b-c). Within four of the six cell type

clusters, individual samples clustered by line rather than species of origin, potentially indicating line to

line variability in differentiation (Fig. 1b). This highlights the importance of measuring ASE which, by

definition, is measured within each line and so robust to variability between lines. Indeed, when

performing PCA within cell types using allelic counts (i.e. counting reads from the human allele and

chimpanzee allele separately), human and chimpanzee species differences were clearly separated by

principal component (PC) 1 or PC2 in each cell type (Fig. 1d, Supp Fig. 2). To assess the success of our

differentiations, we examined each cell type for the expression of known marker genes in our RNA-seq

data (Fig. 1e, Supp Fig. 3). All cell types express canonical marker genes and do not express pluripotency

markers, indicating that the differentiations were successful (Fig. 1e, Supp Fig 3, Additional file 1)

(Burridge et al., 2014; Chal et al., 2016; Korytnikov & Nostro, 2016; Mallanna & Duncan, 2013; Maury et

al., 2015; Sharma et al., 2019).

Because our hybrid cells were grown concurrently with their human and chimpanzee diploid "parental"

cells, we performed an additional check for purity of the hybrid lines by quantifying genome-wide ASE.

We noticed that among our 25 RNA-seq samples, the two PP hybrid2 samples had a slight bias towards

higher expression from the chimpanzee alleles across all chromosomes. This is likely due to a small

fraction of contaminating chimpanzee cells in these samples. We corrected for this by reducing the

chimpanzee allele counts such that the number of reads assigned to the human and chimpanzee alleles

was equal. By simulating contamination of a hybrid sample with chimpanzee cells, we found that this

correction was conservative and that the log fold-change estimates were largely unaffected by

contamination after this correction (Methods, Supp Fig. 4).

We next investigated which genes were differentially expressed between the human and chimpanzee

allele in each cell type (Fig. 2a). We identified thousands of genes showing significantly biased ASE in

each cell type at a false discovery rate (FDR) cutoff of 0.05 (Fig. 2b). We detected a comparable number

of ASE genes in all cell types except SKM. As a result, we repeated all subsequent analyses both including

and excluding SKM and obtained qualitatively similar results regardless of whether SKM was included.

While a considerable number of genes had significant ASE in all cell types, many more genes only had

significant ASE in a single cell type, suggesting cell type-specific *cis*-regulatory divergence (Fig. 2b). A

notable family of developmentally important genes that exemplifies differences in ASE across cell types

is the neurotrophins and their receptors (Fig. 2c) (Caporali & Emanueli, 2009; Huang & Reichardt, 2001).

For example, *NTRK3*, which plays a key role in the development of the nervous system, is only

differentially expressed in RPE and MN but is chimpanzee-biased in RPE and human-biased in MN (Supp

Fig. 5) (Ichim et al., 2012; Naito et al., 2017). In addition, the gene coding for its primary ligand (*NTF3*) is

differentially expressed in a variety of cell types yet is human-biased in all cell types except MN in which

it is chimpanzee-biased (Supp Fig. 5). *NTRK1* differential expression is similarly tissue-specific as it is

strongly chimpanzee-biased in MN, but human-biased in CM and SKM (Supp Fig. 5). These results

indicate that the regulatory landscape of these genes has undergone many complex *cis*-regulatory

changes as the human and chimpanzee lineages have diverged.

To further investigate the relationship between tissue-specificity and ASE, we asked whether genes with

variable expression across tissues are more likely to show ASE. Using a standard definition of cell type-

specific genes—those with detectable expression in only one cell type in our study—we found that cell

type-specific genes were typically enriched for ASE in the one cell type where they are expressed (Supp

Fig. 6) (GTEx Consortium, 2017; Jain & Tuteja, 2019). However, other cell type-specific expression

patterns such as uniquely low expression in a particular cell type may also indicate an important dosage-

sensitive function in that cell type. We therefore focused on a broader definition of cell type-specificity

in which genes that are differentially expressed between one cell type and all others in our study (FDR <

0.05 for each pairwise comparison) are considered cell type-specific for that cell type. We found that

this more inclusive definition, which assigned many more genes to be cell type-specific, showed an even

more significant ASE enrichment than the narrower definition (Fig. 2d). This result is not sensitive to the

choice of FDR cutoff nor driven solely by a subgroup of highly expressed genes as it is well-preserved

across FDR cutoffs (Supp Fig. 7) and different gene expression levels (Supp Fig. 8). This trend is also

robust to separating samples into two groups and using one to define cell type-specific genes and the

other to identify differentially expressed genes. This controls for spurious relationships that can result

when the same data are used to define two different quantities which are then compared (Supp Fig. 9)

(Fraser, 2019).

This enrichment (Fig. 2d) suggests that tissue-specific genes may have less constraint and/or more

frequent positive selection on their expression. We reasoned that if the trend was solely driven by

constraint, then controlling for constraint—even if imperfectly—would be expected to reduce the

strength of the relationship. To investigate this, we binned genes by their variance in ASE across a large

cohort of human samples which we have previously shown acts as a reasonable proxy for evolutionary

constraint on gene expression (Castel et al., 2020; Starr et al., 2023). Across cell types, we generally

observe significant enrichments in each bin and little difference in enrichment between bins, suggesting

that differences in constraint on expression of cell type-specific vs. ubiquitously expressed genes are not

solely responsible for our observations (Supp Fig. 10). Furthermore, we observe even stronger

enrichments using an alternative constraint metric, the probability of haploinsufficiency score (pHI)

likely due to the larger number of genes for which pHI can be calculated (Supp Fig. 11) (Collins et al.,

2022). Overall, our analysis suggests that differences in constraint are unlikely to fully explain these

trends, suggesting a potential role for positive selection.

Lineage-specific selection has acted on tissue-specific gene expression divergence

Next, we sought to use our RNA-seq data to identify instances of lineage-specific selection. We used a

previously published method that incorporates a tissue-specific measure of constraint on gene

expression to test whether groups of genes have likely been under selection (Starr et al., 2023). Using

this strategy, we detected several signals of putative lineage-specific selection, some of which were cell

type-specific (Additional file 2). Notably, the four most significant enrichments were found in motor

neurons and cardiomyocytes and are highly relevant to those cell types (Fig. 2e; Additional file 2). In

cardiomyocytes, the top pathway was "PPAR signaling pathway" which has been shown to play an

important role in the regulation of heart morphology and lipid metabolism (Montaigne et al., 2021). For

example, *NR1H3* (also known as *LXRA*) is strongly upregulated in human cardiomyocytes as well as all

other cell types (Supp Fig. 12a). Furthermore, this upregulation appears to have occurred in the human

lineage based on data from non-hybrid cardiomyocytes as well as adult hearts (Supp Fig. 12b) (Blake et

al., 2020; Pavlovic et al., 2018). Hybrid cells are essential in determining that the human-specific

upregulation of *NR1H3* in cardiomyocytes has a strong genetic component as *NR1H3* expression is very

responsive to diet and other environmental factors (Wang & Tontonoz, 2018).

In motor neurons, multiple categories showed a strong bias toward higher chimpanzee expression

including "sodium ion transmembrane transport" and "syntaxin binding". The genes in these categories

are of fundamental importance to the function of motor neurons as sodium ion transporters control

excitability and syntaxin binding proteins play a major role in controlling the release of

neurotransmitters from synaptic vesicles (Brose et al., 2019; Meisler et al., 2021). Interestingly, several

key genes in these sets appear to have human-derived differences in expression that extend beyond

motor neurons to other neuronal types. For example, *SCN1B, SCN2B,* and *SYT2* have chimpanzee-biased

ASE in our MN data. Similarly, these genes have human-biased expression when comparing human,

chimpanzee, and rhesus macaque glutamatergic cortical neurons in a previously published dataset (Supp

Fig. 13) (Kozlenkov et al., 2020). We note that several other genes in these genes sets are not

differentially expressed between humans and chimpanzees in this cortical neuron dataset, emphasizing

the importance of studying individual neuron types (Kozlenkov et al., 2020). Overall, the strong bias in

gene expression of sodium ion transporters and syntaxin binding proteins we observe suggests lineage-

specific selection that may have altered the electrophysiological properties of human motor neurons.

Patterns of allele-specific chromatin accessibility reveal divergent *cis*-regulatory elements

While ASE provides insight into what gene expression changes might underlie phenotypic differences

between humans and chimpanzees, in the absence of additional data it is very difficult to prioritize

which specific mutations might cause expression divergence. To begin to fill this gap, we generated

ATAC-seq data from five of the six cell types (all except RPE), with each cell type sequenced to an

average depth of 184 million paired-end reads (Supp Fig. 14). ATAC-seq uses a hyperactive Tn5

transposase to cleave DNA that is not bound by nucleosomes to enrich for accessible chromatin (Fig. 3a),

a hallmark of active *cis*-regulatory elements (CREs) (Buenrostro et al., 2013; Corces et al., 2017). We

estimated ASCA for individual open chromatin peaks by mapping reads to both species' reference

genomes, correcting for mapping bias, generating a unified list of peaks across all samples, and then

counting reads supporting each allele in each peak (see Methods). Cell types clustered well using PC1

and PC2 of the ATAC-seq data, except for the HP sample which clustered closely with PP samples (Fig.

3b, Supp Fig. 15). However, performing PCA on just the HP and PP samples clearly separates the two cell

types (Supp Fig. 16). Within each cell type, species differences were clearly separated by PC1 or PC2

(Supp Fig. 17). As an example of ASCA, the accessibility of the promoter of *CTSF* was strongly

chimpanzee-biased, mirroring the chimpanzee-biased ASE for this gene (Fig. 3c).

As a first step in analyzing the ATAC-seq data, we intersected the peaks we identified with the genomic

annotations of chromatin states. These fifteen categories, predicted across many tissues and cell types

by the chromHMM model (Vu & Ernst, 2022), include terms such as "TSS" (transcription start site) and

"enhancer". We then plotted the median of the absolute human-chimpanzee ASCA log fold-changes for

each chromatin state and cell type (Fig 3d). The TSS and promoter annotations were the least divergent

in their accessibility, whereas regions of heterochromatin were the most divergent (Fig. 3d). To explore

the relationship between interspecies differences in ASCA and ASE, we assigned peaks to the nearest

TSS and computed the Pearson correlation between the allelic log fold-change of chromatin accessibility

and expression within each cell type and chromatin state. As expected, TSS and promoter annotations

showed the strongest correlation between ASCA and ASE, and correlations were stronger when

including only differentially accessible peaks (Fig. 3e, Methods). Intriguingly however, ASCA of regions

annotated as heterochromatin, polycomb repressed, or quiescent were as strongly correlated with ASE

as elements identified as enhancers or DNase hypersensitivity sites (Fig. 3e). Notably this result is robust

to how peaks and chromHMM annotations were intersected (Supp Fig. 18a), as well as to removal of all

peaks even slightly overlapping TSS or promoter-related annotations (Supp Fig. 18b). Given that

heterochromatin/quiescent regions are highly divergent in ASCA (Fig. 3d), this suggests that changes in

accessibility of these regions may be particularly important in the evolution of gene expression.

Next, we investigated whether the analog of the relationship between cell type-specific gene expression

and ASE (Fig. 2d) holds for chromatin accessibility. Since the number of called peaks is largely dependent

on sequencing depth (Supp Fig. 14, 19a), we performed down-sampling to equalize power to detect

peaks across cell types (Supp Fig. 19b, Methods). We then called peaks on the down-sampled data and

identified peaks as cell type-specific if they were called as peaks in only one cell type. In agreement with

the gene expression data, we observed that cell type-specific peaks are enriched for ASCA across all cell

types and this enrichment generally holds when using varying log_2 fold-change or p-value cutoffs (Supp

Figs. 20-21). Analogous to our analysis of gene expression, we also applied a broader definition of cell

type-specificity to the ATAC data, in which a peak was considered specific to a cell type if that peak had

an absolute log_2 fold-change greater than a chosen threshold (e.g. 0.5 or 1) across all pairwise

comparisons with other cell types. We observe strong enrichment for ASCA in cell type-specific peaks

using this definition except when using the most stringent cutoffs due to the very low number of peaks

meeting these criteria (Supp Figs. 22-23, Methods). Finally, we observe the same enrichments when

controlling for a recently published metric for constraint on non-coding elements, suggesting that

differences in evolutionary constraint may not be solely responsible for the observed trends (Supp Fig.

24) (S. Chen et al., 2022).

We next explored the relationship between cell type-specific ASCA and ASE. To do this, we developed a

novel metric called differential expression enrichment (dEE) to quantify how specific the log fold-change

is to a particular cell type or tissue. Our method is based on expression enrichment (EE) (Yu et al., 2006),

a metric that measures how specific gene expression is to a certain cell type/tissue. dEE estimates how

cell type-specific ASE is for a gene (Supp Fig. 25, Methods) and, analogously, dCAE (differential

chromatin accessibility enrichment) measures how cell type-specific ASCA is for a *cis*-regulatory element

(Supp Fig. 26, Methods). dEE and dCAE are high in a cell type if there is a high absolute log fold-change in

that cell type and much lower absolute log-fold changes or log fold-changes in the opposite direction in

the other cell types. For example, dEE would be close to one for a gene in a cell type if the gene had

strongly human-biased ASE in that cell type and very weakly human-biased or chimpanzee-biased ASE in

the other cell types. On the other hand, if a gene did not have any strong allelic bias, that gene would

have dEE close to zero. dEE is conceptually related to a metric we have previously introduced, diffASE,

and generalizes diffASE to an arbitrary number of cell types and any assay that produces log fold-

changes (Combs et al., 2018; Hu et al., 2022; York et al., 2018). Using these metrics, we identified 154

instances in which a gene with cell type-specific ASE (i.e., high dEE) had a peak with cell type-specific

ASCA in the same cell type (i.e., high dCAE, see Additional file 3 for the full list). Of these, 95 showed

ASCA and ASE in the same direction which is more than expected by chance (77 expected; $p < 0.005$,

binomial test). These results suggest that tissue-specific *cis*-regulatory divergence in chromatin

accessibility may often impact tissue-specific gene expression, though this divergence is neither

necessary nor sufficient to do so.

Identifying candidate causal *cis*-regulatory variants by integrating ASE and ASCA across cell types

As high dEE and dCAE in a given cell type might be indicative of a causal link between the change in

chromatin accessibility and the change in expression, we focused on the 95 peak-gene pairs with

matching direction and used two different strategies to identify examples to investigate in detail. First,

we prioritized genes known to play important roles in development. For example, we found that the

promoter of *FABP7* has human-biased ASCA specifically in motor neurons (Fig. 4a-b) and that the *FABP7*

gene has human-biased ASE in motor neurons (Fig 4c). *FABP7* is used as a marker of glial cells and neural

progenitor cells (NPCs) and plays a key role in NPC proliferation and astrocyte function (Arai et al., 2005;

De Rosa et al., 2012; Ebrahimi et al., 2016; Watanabe et al., 2007). Using previously published single-

nucleus RNA-seq data from humans, chimpanzees, and rhesus macaques, we confirmed that *FABP7*

shows a human-derived up-regulation in several neuronal subtypes but not glial cells (Supp Fig. 27) (Ma

et al., 2022). To investigate the genetic basis of this cell type-specific divergence, we leveraged a

machine learning model, Sei (K. M. Chen et al., 2022), to nominate potentially causal variants in the

promoter of *FABP7* (see Methods). Sei is a deep neural network that takes DNA sequence as input and

predicts the probability that the sequence has a particular epigenetic state in a variety of cell types and

tissues (Fig. 4d) (K. M. Chen et al., 2022). While single-base substitutions differing between human and

chimpanzee had only minor impacts on predicted *cis*-regulatory activity, "chimpanizing" the human

sequence of the *FABP7* promoter at two small indels (by deleting one base at chr6: 122,779,291 and

inserting three bases at chr6: 122,779,115) was sufficient to make the Sei predictions for the

chimpanized human sequence closely match the predictions for the complete chimpanzee sequence

(Fig. 4e-g). Making only one of these changes had substantial but weaker effects in both cases,

suggesting that both mutations might be functionally important (Fig. 4f,g). The 1-base insertion in the

human lineage introduces a binding site for the neuronally expressed transcription factors GLIS2/3,

suggesting a potential molecular mechanism (Calderari et al., 2018; Castro-Mondragon et al., 2022; Ke

et al., 2015).

As another approach to ranking the 95 peaks, we searched for peaks containing human-chimpanzee

sequence differences in otherwise highly conserved genomic positions, since these could reflect changes

in selective pressure. Using PhyloP scores for 241 placental mammals (Sullivan et al., 2023) to assess

conservation, one of the top-ranked peaks was a putative enhancer six kilobases away from the TSS of

GAD1, which plays a key role in the synthesis of the neurotransmitter GABA (Feldblum et al., 1993).

Notably, part of this peak has been classified as a human accelerated region (HAR) (Girskis et al., 2021;

Hubisz & Pollard, 2014; Pollard et al., 2006)—a short sequence that is highly conserved in mammals yet

contains an unusual number of human-specific mutations. Both the accessibility in the peak and *GAD1*

expression are only chimpanzee-biased in motor neurons (Fig. 4a-c, Supp Fig. 28). Applying Sei to

estimate the predicted effect of every variant in this region, we found that a single-nucleotide

substitution within the HAR (chr2: 170,823,193) has by far the largest predicted cis-regulatory effect and

most closely matches the differences in Sei predictions between the full human and chimpanzee

haplotypes (Fig. 4d-g). Interestingly, this mutation is predicted to disrupt a binding site for several basic

helix-loop-helix transcription factors that play essential roles in neuronal differentiation such as Ascl1

(Supp Fig. 29) (Castro-Mondragon et al., 2022; Mizuguchi et al., 2006; Yang et al., 2017).

As *GAD1* is only highly expressed in GABAergic neurons (and was therefore lowly expressed in the cell

types studied here, Supp Fig. 30a), we investigated whether this reduced expression of human *GAD1*

also occurs in cortical organoids which contain GABAergic neurons together with other cell types in

which *GAD1* is not highly expressed. We analyzed our previously published data from human-

chimpanzee hybrid cortical organoids (Agoglia et al., 2021) and found that the expression of *GAD1* from

the chimpanzee allele spikes higher than that of the human allele around day 50 of hybrid cortical

organoid differentiation before dropping in expression over time to match the human expression level

(Supp Fig. 30b). Because ASE in the hybrid cells controls for any potential interspecies differences in

differentiation kinetics or cell type composition, this difference must be the result of *cis*-regulatory

divergence between humans and chimpanzees. This expression difference is also more pronounced in

comparisons of human and chimpanzee parental cortical organoids, with a higher absolute log fold-

change at day 50, day 100, and day 150, only returning to equal expression at day 200 (Supp Fig. 30c).

While this could be due to differences in cell type proportion between human and chimpanzee

organoids, it might also be due to a reinforcing *trans*-acting effect.

To test whether this difference in expression also occurs specifically during GABAergic neuron

differentiation we examined *GAD1* expression in single cell RNA-seq data from human and chimpanzee

cortical organoids (Kanton et al., 2019). We observed a spike in *GAD1* expression in less mature

chimpanzee GABAergic neurons that is absent in the corresponding part of the trajectory in human

neurons (Supp Fig. 31). Notably, a similar trend holds regardless of which GABAergic sub-trajectory (i.e.,

equivalent to GABAergic neurons from the caudal, lateral, or medial ganglionic eminences) is examined

suggesting this difference is not unique to a particular type of GABAergic neuron (Supp Fig. 31). Finally,

we examined the accessibility of the putative *GAD1* enhancer more closely. Consistent with a potential

role for this enhancer in the spike in *GAD1* expression during development, the accessibility of this

enhancer mirrors the expression of *GAD1* in human cortical and striatal organoids (Supp Fig. 32) with

high accessibility between day 50 and day 100 before decreasing somewhat near day 150 (Trevino et al.,

2020). Overall, our results demonstrate how the combination of RNA-seq, ATAC-seq, and machine

learning models can nominate variants that may be responsible for cell type-specific changes in gene

expression and chromatin accessibility.

Discussion

In this study, we quantified human-chimpanzee *cis*-regulatory divergence in gene expression and

chromatin accessibility in six different cell types representing diverse developmental lineages. Across the

thousands of genes with ASE, we found that most *cis*-regulatory divergence is specific to one or a few

cell types. Furthermore, we found that divergent *cis*-regulation is linked to tissue-specificity, with tissue-

specific genes being enriched for ASE and tissue-specific regulatory elements being enriched for ASCA.

As this result was largely unchanged when stratifying by evolutionary constraint, our results suggest that

changes in the expression of genes with more cell type-specific expression patterns may be less

deleterious than changes in more broadly expressed genes, supporting the idea that cell-type specific

divergence may be less pleiotropic (Wittkopp & Kalay, 2012). Overall, this suggests that broad changes

in expression in cell type-specifically expressed genes may be an important substrate for evolution.

We also identified several sets of genes evolving under lineage-specific selection that may have played a

role in establishing unique facets of human physiology and behavior. Most interestingly, we found

evidence for selection on sodium ion transporters and syntaxin binding proteins that may alter the

electrophysiological properties of motor neurons, and potentially other types of neurons as well (Brose

et al., 2019; Meisler et al., 2021). The complexity of the molecular machinery regulating neuronal

excitability and synaptic vesicle release make it difficult to say what the effects of these gene expression

changes are on the excitability of motor neurons without electrophysiology data from human and great

ape neurons coupled with perturbation of candidate genes. However, given the divergence in

locomotion and motor skills between humans and chimpanzees, it is tempting to speculate that these

changes may have had some role in the evolution of motor control and learning in humans.

In this work, we developed two metrics—dEE and dCAE—to quantify the degree of cell type-specific

differential expression and accessibility. These metrics are largely analogous to widely used metrics that

quantify tissue- or cell type-specific expression level and applicable to any comparison of log fold-

changes across conditions. They markedly improved our ability to identify matching cell type-specific

ASE and ASCA and led to the identification of 95 peak-gene pairs that had highly cell type-specific

concordant changes in accessibility and expression.

One such example is a potentially human-derived increase in *FABP7* expression in several types of

human neurons. As *FABP7* is not highly expressed in adult mouse neurons, the functional consequences

of its higher expression in human neurons are difficult to predict (Yao et al., 2021). *FABP7* plays a role in

the uptake of the fatty acid Docosahexaenoic Acid (DHA), an important component of neuronal

membranes (Akbar et al., 2005; Choi et al., 2021). DHA promotes neuronal survival through

phosphatidylserine accumulation, so it is possible that the human-specific *FABP7* expression increases

neuronal DHA uptake leading to reduced apoptosis in human neurons during development and

ultimately contributing to a larger number of neurons in humans (Akbar et al., 2005; Choi et al., 2021).

In addition, we identified a highly conserved developmentally dynamic enhancer near *GAD1* that may

have partially lost activity in the human lineage resulting in a decrease in *GAD1* expression early in

GABAergic neuron development. By integrating with the deep learning model Sei, we identified a variant

that may account for the chimpanzee-biased ASCA in this region (K. M. Chen et al., 2022). Interestingly,

the ASE of *GAD1* was coupled with a relatively small (though significant) magnitude of ASCA. This could

potentially reflect divergence in transcription factor binding that leaves a "footprint" resulting in subtle

ASCA (Vierstra et al., 2020). Overall, our data suggest that this enhancer has lost activity in the human

lineage, potentially altering the expression pattern of *GAD1* during neuronal development. *GAD1* is the

rate-limiting enzyme for GABA synthesis so GABA levels are likely responsive to changes in *GAD1*

expression (Feldblum et al., 1993). GABA release has complex context-specific effects on

neurodevelopment making it difficult to speculate as to what the phenotypic effects of reduced GABA

synthesis during human neurodevelopment might be (Ben-Ari et al., 2012). However, the high

conservation of this *cis*-regulatory element in placental mammals implies that its human-specific

disruption is likely to have important neurodevelopmental effects. Careful perturbation of this enhancer

and *GAD1* expression in mouse models will be required to explore this further.

Overall, our study provides foundational data, insight, and computational tools that will improve our

understanding of cell type-specific *cis*-regulatory evolution and the role it has played in the

establishment of human-specific traits.

Figure 1: Allele-specific expression across diverse human-chimpanzee hybrid cell types. a) Six cell types

were differentiated from human-chimpanzee hybrid induced pluripotent stem cells. These six cell types

represent diverse body systems, including motor neurons for the central nervous system, retinal

pigment epithelium for eye, skeletal myocytes for skeletal muscle, cardiomyocytes for the heart,

hepatocyte progenitors for the liver, and pancreatic progenitors for the pancreas. **b)** Heatmap showing

the result of hierarchical clustering performed on genes with highly variable normalized allele counts. **c)**

Result of running PCA on normalized allelic counts for all samples and cell types. **d)** Result of PCA

performed on normalized allele counts for each individual cell type separately. Cardiomyocytes and

motor neurons are shown here. **e)** Expression of marker genes for each cell type.

a

Human Allele   Chimp Allele

Gene A

Expression level

Gene B

Expression level

b

Number of genes with ASE shared in x cell type(s)

6 (1361)
5 (1304)
4 (1597)
3 (2208)
2 (2594)
1 (3047)

Number of ASE Genes Intersection

Number of ASE Genes

6000 4000 2000 0

SKM
MN
RPE
CM
HP
PP

c

Log2(Human/Chimp)

-2 -1 0 1 2

BDNF
NGF
NGFR
NTF3
NTF4
NTRK1
NTRK2
NTRK3

CM HP MN PP RPE SKM

d

ASE Gene Enrichments for Cell Type Specifically Expressed Genes

Cardiomyocyte   ***
Hepatocyte Progenitor   ***
Motor Neuron   ***
Pancreatic Progenitor   ***
Retinal Pigment Epithelium   ***
Skeletal Myocyte   **

0.5 1.0 1.5 2.0 2.5

Odds Ratio

e

Cardiomyocyte

PPAR signaling pathway

Oxidoreductase activity, acting on the CH–CH group of donors

Antiporter activity

Antigen processing and presentation

20 10 0 10 20

Number of Genes Biased

Motor neuron

Sodium ion transmembrane transport

Syntaxin binding

Hyperactive deep tendon reflexes

NAD+ ADP–ribosyltransferase activity

20 10 0 10 20

Number of Genes Biased

Chimp ← → Human

Figure 2: Human-chimpanzee ASE is largely cell type-specific. a) Outline of measurement of allele-

specific expression. Reads from the human and chimpanzee alleles are counted and differences in read

counts identified. **b)** Thousands of genes with ASE were identified for each cell type, with many genes

only showing ASE in a single cell type (Conway et al., 2017). **c)** The neurotrophins and their receptors as

examples of genes showing cell type-specific ASE patterns. DESeq2 estimate of $\log_2$ fold-change

(human/chimpanzee) are shown in the heatmap and significance is indicated by asterisks where ***

indicates FDR < 0.005, ** indicates FDR < 0.01, and * indicates FDR < 0.05. Zero asterisks (i.e. a blank

box) indicates FDR > 0.05. **d)** Plot showing that genes with ASE are enriched for genes showing cell type-

specific expression patterns across all cell types. Asterisks indicate p-values rather than FDR using the

same system as in 2c. **e)** Top gene sets with evidence for lineage-specific selection in cardiomyocytes

and motor neurons are shown. The length of the bars indicates the number of genes in a category with

biased expression in each cell type.

Figure 3: Allele-specific chromatin accessibility across diverse human-chimpanzee hybrid cell types. a)

Schematic outlining the ATAC-seq protocol. A hyperactive transposase cleaves accessible DNA and adds

adapters enabling measurement of chromatin accessibility. **b)** PCA on normalized allelic counts from

ATAC-seq. **c)** ASCA in the promoter of *CTSF,* and ASE for the *CTSF* gene. **d)** Differences in ASCA were

quantified and plotted separately based on chromHMM annotation. The order is based on the median

of z-score transformed absolute log fold-change between human and chimpanzee across all cell types,

with higher z-scores indicating greater divergence in accessibility. **e)** Pearson correlation between ASE

and ASCA for all cell types with all peaks (left) or only differentially accessible peaks (right, defined as

peaks with nominal binomial p-value less than 0.05). Pearson's r values are shown in the heatmap and

significance is indicated by asterisks where *** indicates $p < 0.005$, ** indicates $p < 0.01$, and * indicates

$p < 0.05$. Zero asterisks (i.e. a blank box) indicates $p > 0.05$.

Figure 4: Motor neuron-specific human-biased ASE and ASCA for *FABP7* and the promoter of *FABP7*. a)

Allelic ATAC-seq tracks are shown in the peak containing the annotated *FABP7* promoter (highlighted in

yellow). **b)** The top panel shows allelic CPM of the *FABP7* promoter across cell types and the bottom

panel shows the log fold-change across cell types. **c)** The top panel shows allelic CPM of the *FABP7* gene

across cell types and the bottom panel shows the log fold-change across cell types. **d)** Outline of the

process for variant effect prediction with Sei for *FABP7*. All sequences input to Sei were centered at the

FABP7 promoter. The human sequence, chimpanzee sequence, and partially "chimpanized" human

sequences (modified by systematically switching the human allele to the chimpanzee allele separately

for each human-chimpanzee difference) were fed into Sei to predict the effects of these variants on

chromatin state. **e)** Histogram of the probabilities of various chromatin states and transcription factor

binding predicted by Sei was plotted for the human sequence, the chimpanzee sequence, and the

human sequence with one human-chimpanzee difference swapped to match the chimpanzee sequence.

The human sequence with SNV26 changed to the chimpanzee allele and the human sequence switched

at both indels are shown as examples. A histone modification (H3K4me3) predicted in two cell types was

labeled to illustrate how the predictions depend on both input sequence and cell type. **f)** Plot of the

predicted effects of all single nucleotide differences and indels between the human and chimpanzee

genomes in the Sei input window (the *FABP7* promoter is highlighted in yellow). **g)** Scatterplots showing

the correlation of the effects of both indels (left panel), each individual indel (middle two panels), and a

representative SNV (right panel) on Sei predictions with the difference in Sei predictions between the

human and chimpanzee sequences. The root mean square error (RMSE) was computed and shown in

each figure.

Figure 5: Motor neuron-specific chimpanzee-biased ASE for *GAD1* and ASCA for a HAR near the *GAD1*

TSS. a) Allelic ATAC-seq tracks are shown for the peak near the *GAD1* TSS that contains a HAR

(highlighted in yellow). **b)** The top panel shows allelic CPM of the CRE near the *GAD1* TSS across cell

types and the bottom panel shows the log fold-change across cell types. **c)** The top panel shows allelic

CPM of the *GAD1* gene across cell types and the bottom panel shows the log fold-change across cell

types. **d)** Outline of the process for variant effect prediction with Sei for *GAD1*. All input sequences to Sei

were centered at the HAR. The human sequence, the chimpanzee sequence, and modified sequences

with the human sequence altered at each substitution to match the chimpanzee sequence were fed into

the Sei sequence model to predict the effects of these variants on the chromatin state. **e)** Histogram of

the probabilities of various chromatin states and transcription factor binding predicted by Sei was

plotted for the human sequence, the chimpanzee sequence, and two examples in which the human

sequence with only one SNV "chimpanized" (human w/SNV) was input to Sei. The histogram of the

probability of the sequence having a particular epigenomic annotation (predicted by Sei) was plotted for

human, chimpanzee, human w/SNV1 changed to match the chimpanzee sequence, and human w/SNV21

changed to match the chimpanzee sequence. Two epigenomic annotations were labeled as examples

that show the different values output by Sei with these two different sequence inputs. **f)** Plot of the

predicted effects of all single nucleotide differences between human and chimpanzee in the Sei input

window centered at the HAR (highlighted in yellow). Positions were switched to the chimpanzee allele

individually. **g)** Scatterplots showing the correlation of the effects of four SNVs on Sei predictions with

the difference in Sei predictions between the human and chimpanzee sequences. The root mean square

error (RMSE) was computed and shown in each figure.

Methods:

Generation of multiple human-chimpanzee hybrid cell types

We used two previously described human-chimpanzee hybrid iPS cell lines (hybrid1 and hybrid2,

previously denoted Hy1-25 and Hy1-30 respectively) (Agoglia et al., 2021). Before differentiation, cells

were routinely cultured on matrigel in mTeSR1 or mTeSR Plus (Stem Cell Technologies cat #85850 or cat

#100-0276). Culture and *in vitro* differentiation of iPS cells into six cell types (motor neurons (MN),

cardiomyocytes (CM), hepatocyte progenitors (HP), pancreatic progenitors (PP), skeletal myocytes

(SKM), and retinal pigment epithelium (RPE)) was carried out by the Columbia Stem Cell Core Facility

using published protocols (Burridge et al., 2014; Chal et al., 2016; Korytnikov & Nostro, 2016; Mallanna

& Duncan, 2013; Maury et al., 2015; Sharma et al., 2019).

Preparation of RNA-seq libraries

All samples were cryopreserved in liquid nitrogen before RNA extraction (Milani et al., 2016). Cells were

gently thawed and then washed with PBS and cell pellets were collected via centrifugation at 1,000 RPM

for 5 minutes. Cell pellets were loosened by flicking the tube and an appropriate volume of Buffer RLT

based on the cell count were added following the RNeasy Mini Kit (Qiagen, 74104) protocol. Total RNA

extraction and on-column DNAse digestion were performed using RNeasy Mini Kit (Qiagen, 74104) and

RNase-Free DNase Set (Qiagen, 79254). RNA quality was assessed using the Agilent Bioanalyzer RNA Pico

assay. Only samples with an RNA integrity number (RIN) greater than or equal to 7 were used to prepare

cDNA libraries. All RNA-seq libraries except 3 motor neuron libraries (2 hybrid2 and 1 hybrid1) were

prepared using the TruSeq Stranded mRNA kit (Illumina, 20020594) and the TruSeq RNA CD Index Plate

(Illumina, 20019792) from between 100 ng and 1 ug total RNA following the manufacturer's protocols.

The other three motor neuron libraries were prepared using Illumina Stranded mRNA Prep (Illumina,

20040532) and IDT for Illumina RNA UD Indexes Set A, Ligation (Illumina, 20040553) due to low yield of

total RNA. Notably, the four motor neuron libraries did not cluster by library preparation method. All

libraries were normalized, pooled at an equimolar ratio using Qubit measurements, and sequenced on

an Illumina HiSeq 4000 to generate 2x150bp paired-end reads.

Identification of confident human-chimpanzee SNVs

To identify a confident list of human-chimpanzee SNVs that could be used to quantify allele-specific

expression and chromatin accessibility, we first downloaded hg38-panTro6 MAF files from UCSC and

whole-genome sequencing data generated from the parental human and chimpanzee iPS cells (in the

form of bam files aligned to hg38 and panTro5, generously provided by the Gilad lab). We first

converted them back to fastq files and then mapped reads to panTro6 and hg38 (we mapped both

human and chimpanzee to both reference genomes) using bowtie2 with the flags —very-sensitive-local -

p 16 (Langmead & Salzberg, 2012). We then used a modified version of our previous approach to filter

out SNVs that could not be confidently identified as homozygous in the human and chimpanzee parental

lines (Agoglia et al., 2021). Briefly, we extracted SNVs and indels from both human and chimpanzee MAF

files, counted reads in the WGS data that supported the human, chimpanzee, or an alternative base at

that position, then filtered out any SNVs with < 2 reads or < 90% of reads supporting that species' base.

We then reformatted files, merged with indels for use in Hornet, and generated a modified bed file of

SNVs that includes the human and chimpanzee base at the SNV position (van de Geijn et al., 2015).

Generation of allele-specific count tables

An allele specific expression mapping and calling pipeline adapted from Agoglia et al. and our updated

high-confidence SNV list was used. The whole pipeline was carried out twice independently using hg38

or panTro6 as the reference genome. This approach was taken to eliminate genes showing strong

mapping bias, defined here as genes with an absolute difference in log fold-change between the

panTro6 referenced and hg38 referenced runs greater than one. All sequencing reads were trimmed

with SeqPrep (adapters specified by the manufacturer for the different library preparation kits) and

mapped using STAR with two passes and the following parameters: --outFilterMultimapNmax 1 (Dobin

et al., 2013; John St. John, n.d.). Uniquely aligned reads were deduplicated with Picard and Hornet (an

implementation of WASP which first removes reads overlapping indels) was used to correct for mapping

bias (Broad Institute, n.d.; van de Geijn et al., 2015). Reads were assigned to either the human allele if

they contained one or more human-chimpanzee single nucleotide differences that matched the human

sequence and zero positions that matched the chimpanzee sequence (and vice versa for assigning reads

to the chimpanzee allele) and counted per gene as previously described (Agoglia et al., 2021).

Detection of aneuploidy on chromosome 20 and slight chimpanzee parental contamination in PP

hybrid2 samples

In our quality control process, we plotted the log fold-change for each gene along every chromosome

and inspected the results. This revealed a clear bias toward the human allele on a part of chromosome

20 for hybrid2 samples, suggesting chromosome 20 aneuploidy which was also reported by Agoglia et al

(Agoglia et al., 2021). As a result, we excluded chromosome 20 from all downstream analyses. In

addition, we found that PP hybrid2 samples had a slight bias toward the chimpanzee allele across every

chromosome which was most likely due to a small fraction of contaminating chimpanzee cells in these

samples. Rather than removing these samples, we normalized the allele-specific count tables by

subtracting a small number of reads from the chimpanzee allele counts calculated based on the biased

ratio summarized from genome wide human and chimpanzee allele counts, to force a global log fold-

change (across all autosomes except chromosome 20) of zero between the human and chimpanzee

alleles. We applied this normalization to all other samples as well. To evaluate the success of this

strategy, we took hybrid and chimpanzee parental iPSC RNA-seq from Agoglia et al (Agoglia et al., 2021)

from the same iPS cell lines as used in this study, and then simulated chimpanzee parental

contamination by mixing chimpanzee parental data into hybrid 2 data to reach similar a similar degree

of chimpanzee bias level to that observed in the PP hybrid2 samples. We then identified genes showing

ASE (see "Identifying genes with ASE") using the counts from the original hybrid samples, simulated

contaminated samples, and corrected simulated contaminated samples respectively and compared the

outputs (Supp Fig. 4).

PCA and hierarchical clustering

Allelic counts were normalized by DESeq2 rlog and principal components analysis (PCA) was performed

on rlog normalized allelic counts with default centering and scaling (Love et al., 2014). The top 1,000

variable genes with the highest variance of normalized allelic counts across all cell types were used to

compute Euclidean distance matrices. The R package pheatmap was used to do hierarchical clustering

and heatmap plotting.

Identifying genes with ASE

DESeq2 was used to measure allele-specific expression (ASE) in each cell type (Love et al., 2014). All

reads from chromosome 20 were removed (as mentioned above). Two replicates per hybrid line per cell

type (plus one additional replicate for SKM hybrid2 for a total of 3 samples) were used by DEseq2 with

model ~hybLine+Species to measure differential expression level. A likelihood ratio test (test="LRT",

betaPrior=FALSE) was used to compute p-values. P-values were then false discovery rate adjusted using

an implementation of the Benjamini-Hochberg correction in the R package qvalue (Benjamini &

Hochberg, 1995; Storey Lab, n.d.). Log fold-changes were shrunk as recommended by the DESeq2

pipeline (Zhu et al., 2019). Differentially expressed genes were defined as those with FDR < 0.05 when

aligned to hg38 and panTro6 as well as an absolute difference in log fold-change <= 1 when comparing

the results from the two alignments.

Identifying cell type-specifically expressed genes

For the more traditional definition of cell type-specific genes, we required transcripts per million (TPM)

< 1 for a gene in every cell type except one. In the cell type with TPM > 1, we varied how highly

expressed the gene had to be in that cell type (again using a TPM cutoff, varying between one and five)

to consider that gene to be specific to that cell type. A similar process to the one described in

"Identifying differentially expressed genes" was used to identify cell type-specific genes based on the

broader definition described in the main text. Rather than using allelic counts, total counts for each

sample (i.e. all uniquely mapping deduplicated reads regardless of their allelic origin) were computed by

summing all allelic and non-allelic counts. These counts were inputted to DESeq2 and the expression of

each gene was compared pairwise between all cell types (Love et al., 2014). Genes were defined as cell

type-specifically expressed in a cell type only if all pairwise comparisons between that cell type and

other cell types resulted in an FDR < 0.05 using both hg38 and panTro6 aligned counts. Due to the

markedly lower number of differentially expressed genes identified in SKM, results were computed both

including and excluding SKM. An analogous procedure was used to identify more broadly defined cell

type-specific peaks in the down-sampled ATAC-seq dataset. Peaks were defined as specific to a cell type

if the absolute log fold-change was greater than 0.5 across all pairwise comparisons with the other cell

types. We also tested an absolute log fold-change threshold of 1 to ensure that our results were not

sensitive to the choice of cutoff.

Enrichment test for genes with cell type-specific expression patterns and genes showing ASE

Odds ratios were calculated using the unconditional maximum likelihood estimate implemented in the R

package epitools function oddsratio.wald (), and 95% confidence intervals and p-values were calculated

using the normal approximation. A directly analogous procedure was performed to test for enrichment

of peaks with ASCA and cell type-specific peaks.

Enrichment test stratified by expression level or evolutionary constraint

Enrichment tests were carried out as in 'Enrichment test of cell type-specific expression patterns and

genes showing ASE' except that genes were split into five equal size bins depending on which factors

were used to stratify genes, and tests were done in each bin. When stratifying by expression level, genes

were ordered in ascending order based on expression level (TPM) and then split into 5 equal size bins

where genes in the 0-20% bin are the most lowly expressed genes and genes in the 80-100% bin are the

most highly expressed genes. When stratifying by constraint metrics such as ASE variance or pHI, genes

were ordered in ascending order based on ASE variance values and then split into five equal size bins

where the 0-20% bin contains genes with the lowest ASE variance (i.e. most evolutionarily constrained)

and the 80-100% bin contains genes with highest ASE variance (i.e. least evolutionarily constrained). To

stratify the ATAC data by constraint, we used the "QCed genomic constraint by 1kb regions" computed

by the gnomAD consortium (S. Chen et al., 2022). We further removed any regions that overlapped

protein coding exons from the human gtf file using bedtools subtract (Quinlan & Hall, 2010). If a peak

overlapped two or more 1 kilobase windows, it was assigned to the window with the highest constraint,

mirroring the procedure used by the gnomAD consortium to assign peaks to ENCODE regulatory

elements (S. Chen et al., 2022). Once the peaks were ranked by this metric, a procedure identical to that

for the gene expression constraint metrics outlined above was performed.

Identification of lineage-specific selection on gene expression

We used a modified version of our previously published pipeline, which uses ASE values from many

individuals of a single species to estimate cis-regulatory constraint of each gene (Starr et al., 2023). We

restricted these ASE values to GTEx samples from the tissue (s) of origin for each cell type (with the

exceptions of RPE which was compared to all GTEx samples and MN which was compared to all brain

and peripheral nerve samples as more closely matches tissues such as eyes for RPE are not available in

GTEx). We then used the Mann-Whitney U test to compare the human population ASE distribution to

the human-chimpanzee ASE distribution as previously described (Starr et al., 2023). We then used the

previously described signed ranking by Mann-Whitney p-value that incorporates whether a gene has

human or chimpanzee-biased ASE with GSEAPY and the binomial test to identify instances of lineage-

specific selection (Starr et al., 2023). Due to the focus on tissue-specificity, we did not filter redundant

gene sets with GSEAPY FDR < 0.25 in multiple cell types (Starr et al., 2023; Subramanian et al., 2005).

Preparation of ATAC-seq libraries

We used the OmniATAC protocol with the only modification being the use of 25,000 cells instead of

50,000 since the fused iPS cells are tetraploid (Corces et al., 2017). All samples for ATAC-seq prep were

from the same vials used in RNA-seq library preparation except for the motor neuron libraries due to the

low yield of total RNA extracted from motor neurons. After library preparation and running samples on a

Bioanalyzer, we noticed a considerable number of fragments greater than 1000 bases in length. To

reduce these fragments, we size selected with Ampure beads using the protocol from the Kaestner lab

available here: https://www.med.upenn.edu/kaestnerlab/assets/user-content/documents/ATAC-

seq-Protocol- (Omni)-Kaestner-Lab.pdf.

After size selection and rerunning on the Bioanalzyer, we pooled the libraries together and sequenced

them to compute quality control metrics. We used the R package ChrAccR to compute TSS enrichment

scores (Mueller, Fabian, n.d.). We pooled all libraries with TSS enrichment score greater than 3.5. This

resulted in 2 CM libraries, 2 MN libraries, 2 PP libraries, 1 SKM library, and 1 HP library. After pooling,

libraries were sequenced on an Illumina Hiseq 4000 to produce 2x150 paired-end reads.

Mapping the ATAC-seq data

We trimmed reads using SeqPrep and then mapped them to the hg38 and panTro6 reference genomes

with bowtie2 in paired-end mode (John St. John, n.d.; Langmead & Salzberg, 2012). The following

parameters were used: -X 2000 --very-sensitive-local -p 16. After mapping, duplicates were removed via

Picard MarkDuplicates (Broad Institute, n.d.). We then removed multi-mapping reads with the command

samtools view -b -q 10 (Li et al., 2009). Due to the format of bowtie2's output, running Hornet on all

reads at once was excessively RAM intensive. Therefore, we split the bam files by chromosome and ran

Hornet on each of the chromosomes separately. We used the files of SNVs and indels generated as

described above as input to Hornet. After Hornet finished running, we used samtools merge to merge all

autosomes and sex chromosomes (we excluded the mitochondrial genome) to create a final bam file for

downstream analysis (Li et al., 2009).

Peak calling and filtering

As only one replicate was available for SKM and HP, we generated two pseudo-replicates by randomly

assigning reads to one of two files using Picard SplitSamByNumberOfReads (Broad Institute, n.d.). We

then called peaks on each file separately, as well as a merged file containing all the reads from a

particular cell type. For example, for MN, both replicates were pooled and peaks were called on that file

as well as the two replicates separately. Before peak calling, all bam files were converted to bed files.

We called peaks using MACS2 callpeak with the following arguments: -f BED -p 0.01 --nomodel --shift 75

--extsize 150 -B --SPMR --keep-dup all --call-summits (Y. Zhang et al., 2008, p. 2). We called peaks on

both the chimpanzee-referenced and human-referenced bam files. After peak calling, we sought to filter

peaks using a modified version of the ENCODE pipeline designed to eliminate peaks that lack a one-to-

one ortholog between humans and chimpanzees. The following pipeline was run on each cell type

separately. We first filtered peaks that were not called in both replicates as well as the pooled file using

code from the ENCODE pipeline based on bedtools and awk (Quinlan & Hall, 2010). We then used a

custom Python script to merge overlapping peaks and used UCSC LiftOver to lift the peaks from hg38 to

the panTro6 and back to hg38 as well as from panTro6 to hg38 (Kuhn et al., 2013). We then used

bedtools to intersect the resulting human referenced files and filtered out any peaks that did not have at

least 25% overlap with a peak in the other file (Quinlan & Hall, 2010). After filtering out peaks

overlapping ENCODE blacklisted regions and merging overlapping peaks again, we lifted the file that was

originally chimpanzee-referenced back to the chimpanzee genome (Amemiya et al., 2019). Finally, we

removed human-referenced peaks if their chimpanzee-referenced counterpart failed to LiftOver (Kuhn

et al., 2013).

Annotating the peak lists

To annotate the peaks, we used the list of TSS defined by Horlbeck et al. to annotate peaks (Horlbeck et

al., 2016). We lifted over each TSS to hg38, expanded 1000 bases on either side of the midpoint of each

TSS to generate promoters, and merged any promoters that overlapped while retaining all unique gene

names associated with the promoter (Kuhn et al., 2013). We then used reciprocal LiftOver with panTro6

to filter out non-orthologous promoters and used bedtools intersect to link peaks to promoters and

expanded the peak to include the entirety of the promoter if necessary (Quinlan & Hall, 2010). Through

this process, we also outputted a list of non-promoter CREs (sometimes labeled as enhancers as

enhancers are thought to be the most common type of CRE). We took this list and used bedtools closest

to link them to the two closest protein coding genes (Quinlan & Hall, 2010). Notably, the gene naming

conventions differ for the Horlbeck et al. TSS list and the GTF file used for RNA-seq processing. We

altered all gene names in peaks to match those found in the GTF file. In some cases, the gene no longer

existed in the updated hg38 GTF in which case the gene name was replaced with NAN.

Generating a unified peak list

We next merged our cell type-specific peak list across all five cell types to create a unified peak list. To

do this, we iteratively intersected all the peaks with bedtools and then merged any overlapping peaks

(Quinlan & Hall, 2010). Finally, we added back any peaks that did not intersect a peak found in any other

cell types. We then took the chimpanzee and human-referenced versions of these peak lists and ran

them through the LiftOver-based non-homologous peak filtering pipeline described above to generate a

final file of all identified peaks as well as which cell type (s) they were called in (Kuhn et al., 2013). Then,

we reran the annotation pipeline described in 'Annotating the peak lists' on this new set of peaks.

Counting reads in peaks and further peak filtering

First, we split the bam files into reads that we could confidently assign to the chimpanzee genome and

reads we could assign to the human genome. We used our bed file of high-confidence SNVs and

required at least one SNV matching the human genome as well as no SNVs matching the chimpanzee

genome for a read to be assigned to human (and vice versa for chimp). We then used a custom Python

script to reformat the peak list bed files as GTF files and used HTSeq to count reads in peaks using the

following parameters: -s no -m union -r pos (Anders et al., 2015). We only kept peaks if they had a mean

read count across replicates within a cell type of at least 25 from either allele. For example, if a peak has

an average of 27 reads from the human allele and an average of 10 reads from the chimpanzee allele in

MN, that peak would be kept in MN. On the other hand, if the same peak had an average of 24 reads

from the human allele and an average of 10 reads from the chimpanzee allele in CM, that peak would be

discarded for CM.

We next filtered the reads to remove peaks that might be differentially accessible but show evidence of

mapping bias or do not agree between replicates. To do this, we removed any peaks with an absolute

log fold-change greater than one in one replicate but with a fold-change of any magnitude in the

opposite direction in the other. This was not done for SKM or HP as we had only one replicate. We then

removed any peaks that had a log fold-change in opposite directions with an absolute difference greater

than 1 in at least one replicate when comparing the human-referenced and chimpanzee-referenced

counts. Finally, as described in section ''Detection of aneuploidy on chromosome 20 and slight

chimpanzee parental contamination in PP hybrid2 samples' for RNA-seq data analysis, we removed any

peaks on chr20 and took this as our final list of peaks for downstream analyses. Allelic counts were

normalized as described in the RNA-seq data analysis. We tested for allele-specific chromatin

accessibility (ASCA) using the binomial test applied to the normalized allelic counts (summed by species

within a cell type). We considered any peaks with a binomial p-value less than 0.05 to be nominally

differentially accessible.

Down-sampling to identify cell type-specific ATAC-seq peaks

As the number of peaks detected by ATAC-seq is generally a function of read depth and our read depth

varied widely across cell types, we restricted to one replicate (always hybrid1 if two replicates were

available) and down-sampled reads to match the SKM sample with lowest sequencing depth. We then

called peaks for cell types with a single ATAC replicate as described above.

Allelic chromatin accessibility tracks

Allelic bam files with reads originating from the human allele and the chimpanzee allele (respectively)

were obtained as described in 'Counting reads in peaks and further peak filtering'. Two replicates in CM,

MN, and PP were pooled by cell type. Bam files were converted into bigWig files using python package

deepTools bamCoverage with options: --binSize 1 --normalizeUsing CPM --effectiveGenomeSize

2862010578 --ignoreForNormalization chr20 --extendReads (Ramírez et al., 2014). Tracks were

visualized and plotted using the python package pyGenomeTracks (Lopez-Delisle et al., 2021). When

comparing human and chimpanzee log fold-change track differences in each cell type, deepTools

bigwigCompare was used to compare between human bigWig and chimpanzee bigWig with options: --

pseudocount 1 --skipZeroOverZero --operation log2 -bs 1 (Ramírez et al., 2014).

ChromHMM annotation and correlation with ASE

A universal chromHMM annotation was obtained for each peak based on overlap with any of the 15

categories in chromHMM (excluding the blacklist category, for which peaks had already been removed)

(Ernst & Kellis, 2017; Vu & Ernst, 2022). Divergence was measured as the z-score transformed median of

the absolute log fold-change of human and chimpanzee normalized counts in each peak. Each peak was

assigned to the closest gene and then Pearson correlation was computed between the chromatin

accessibility log fold-change and the expression log fold-change for each peak and its nearest gene.

Pearson correlation was computed only on categories including at least 15 peaks. When showing results

for differentially accessible peaks, only peaks with binomial p-values less than 0.05 were kept and used

in computing the Pearson correlation. When assigning a unique chromHMM to each peak, the

chromHMM category that covered the largest portion of each peak was used. When filtering out

promoter-related annotations, peaks covering any promoter-related chromHMM categories ("TSS",

"flanking promoter" and "bivalent promoter") were filtered out and the analysis described above was

repeated.

Computation of differential expression enrichment (dEE) and differential chromatin accessibility

enrichment (dCAE)

For each target cell type, taking CM as an example, the log fold-change for gene A was fixed as target log

fold-change, and the log fold-changes for gene A in the remaining cell types with an opposing sign

(compared to the target log fold-change) were set to zero. Then, the dEE value was calculated as the

proportion of the target log fold-change in the sum of the zeroed log fold-changes across all cell types.

For example, the dEE for gene A in CM would be abs (target LFC)/sum (abs (LFC after zeroing))). dEE

ranges from zero to one and low dEE value indicates differential expression with similar magnitude and

direction across cell types, and/or the gene does not have any strong allelic bias, whereas a high dEE

value indicates that this gene is only strongly differentially expressed (with the sign the log fold-change

has in that cell type) in a particular cell type. dCAE uses the same procedure as dEE except the table is

populated with the log fold-changes derived from chromatin accessibility measurements. dEE and dCAE

are sensitive to the inclusion or exclusion of cell types (by definition), so we excluded RPE when

integrative analysis combining results from dEE and dCAE was performed (to match the cell types for

which dCAE could be computed, Supp Fig. 25b). After restricting to genes defined as having significant

ASE or significant ASCA, we defined genes with dEE >= 0.75 in a particular cell type as showing cell type-

specific ASE and peaks with dCAE >= 0.75 in one cell type as showing cell type-specific ASCA. We used

bedtools intersect to intersect the peaks with our list of human-chimpanzee single nucleotide

differences and the 241-way placental mammal PhyloP scores (Quinlan & Hall, 2010; Sullivan et al.,

2023). We also checked whether peaks that contained human-chimpanzee differences in sites with high

PhyloP scores were in the list of HARs described in Girskis et al (Girskis et al., 2021).

Predicting regulatory activity with single-variant resolution

We used sequences in fasta format as input to the deep neural network model Sei (K. M. Chen et al.,

2022). Sei requires a 4096 base pair input sequence, so we put the center of our region of interest at the

center of the input window and expanded equally on either side to contain 4096 base pairs. The human

sequence was retrieved from hg38 and the corresponding chimpanzee sequence was retrieved from

panTro6. The effect size when comparing the probabilities of each sequence having a particular

chromatin state was computed as the log of the human sequence probability divided by the chimpanzee

sequence probability. Only annotations for which either the chimpanzee sequence or the human

sequence had a probability value greater than or equal to 0.5 were kept for downstream analysis. All

SNVs between human and chimpanzee in this input window were identified and ordered based on

coordinates. For each SNV position, the human sequence was changed to the chimpanzee allele at that

position to generate a new sequence that was input to Sei. The log fold-change for each chromatin

annotation was computed for each input sequence as described above and used as a measure of the

effect of this change on the sequence. Similarly, an indel can be introduced to modify the human

sequence and input to Sei. With indels, the center of the regions of interest (promoter or HAR) were

always at the center of the input window and the start or end of the sequence inputted to Sei could

possibly lose or gain base pairs. However, we found that for the small indels shown here this had

essentially no effect on the Sei output.

Processing of publicly available datasets

The data from Blake et al. and Pavlovic et al. were processed as previously described (Blake et al., 2020;

Pavlovic et al., 2018; Starr et al., 2023). For the Pavlovic et al. data, log fold-changes were computed in

DESeq2 with the scaled proportion of cardiomyocytes present in each sample (available in the

supplemental materials of Pavlovic et al.), sex, and whether cardiomyocytes were treated with T3 as

covariates (i.e. using the model ~sex+scaled_proportion_cardiomyocytes+T3_Treatment+species)

(Pavlovic et al., 2018). No covariates were included for Blake et al. as they had little impact on the data

(Blake et al., 2020). The log fold-changes and FDR corrected p-values were directly downloaded from the

supplemental materials of Kozlenkov et al. (Kozlenkov et al., 2020).

The processed data from Ma et al. (Ma et al., 2022) were downloaded from

http://resources.sestanlab.org/PFC/. We pseudobulked the data by cell type by summing counts within

each individual. We then separately input each pairwise comparison of two species (human to

chimpanzee or human to rhesus macaque) into DESeq2 with no covariates to test for differential

expression and compute log fold-changes.

The counts tables from Kanton et al.[12] were downloaded from

https://www.ebi.ac.uk/biostudies/arrayexpress/studies/E-MTAB-7552 and processed with SCANPY

(Kanton et al., 2019, p. 29409532; Wolf et al., 2018). The data were filtered by removing cells with

n_genes_by_counts > 2500 and >5% mitochondrial reads. We also removed cells with fewer than 200

unique genes and genes that had non-zero counts in fewer than 3 cells. After filtering, any chimpanzee

cells not falling in the category (defined by Kanton et al. (Kanton et al., 2019)) "ventral forebrain

progenitors and neurons" were eliminated and human cells not in the categories "ventral progenitors

and neurons 1", "ventral progenitors and neurons 2", or "ventral progenitors and neurons 3" were

similarly eliminated. We then merged the two counts tables, normalized/logarithmized the counts,

computed PCA, used harmony to integrate cells from different species (human and chimpanzee), and

found nearest neighbors with the harmonized principal components (Korsunsky et al., 2019). We then

ran Leiden clustering with resolution = 0.5 to identify 7 subclusters (one of which appeared to be a

technical artifact with very low counts that was removed) (Traag et al., 2019).

We identified cell types and lineages using canonical marker genes (*MKI67* and *HES5* for progenitors,

NKX2-1 and *LHX6* for the medial ganglionic eminence or MGE, *MEIS2* and *ZFHX3* for the lateral

ganglionic eminence or LGE, and *SCGN* and *NR2F1* for the caudal ganglionic eminence or CGE) (Su-Feher

et al., 2022). We then used the implementation of PAGA in SCANPY to compute pseudotime using the

first cell in the progenitor subcluster as the root (Wolf et al., 2019). We binned cells into five equal bins

along pseudotime and compared the expression of cells with non-zero counts for *GAD1* in each

pseudotime bin. Within each bin, we used a Wilcoxon test to test for higher expression of *GAD1* in

chimpanzee cells compared to human cells. We repeated the pseudotime analysis, binning, and

comparing of *GAD1* gene expression for each subtrajectory (MGE, LGE, and CGE).

Description of Additional file 1: This file contains an extended evaluation of the success of the

differentiations in generating the desired cell type and other cell types likely to be present in each

sample.

Description of Additional file 2: This file contains the results of running the test for selection described

by Starr et al on each cell type.

Description of Additional file 3: This file contains the peak-gene pairs that had concordant high dCAE

and high dEE in each cell type.

Declarations: The authors have nothing to declare.

Ethics approval and consent to participate: Not relevant to our study.

Consent for publication: Not relevant to our study.

Availability of data and materials: Raw and processed data generated by this study are publicly

available through the Gene Expression Omnibus under accession GSE232949:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE232949. The snRNAseq data from Ma et al.

are available here: http://resources.sestanlab.org/PFC/. The scRNAseq data from Kanton et al. are

available here: https://www.ebi.ac.uk/biostudies/arrayexpress/studies/E-MTAB-7552. The bulk RNA-seq

data from Blake et al. and Pavlovic et al. are available here:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE112356 and here:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE110471 respectively. The log fold-changes

and associated statistics were used directly from the supplemental materials of Kozlenkov et al. The

human-chimp pairwise alignment used to identify SNVs and indels is available here:

https://hgdownload.soe.ucsc.edu/goldenPath/hg38/vsPanTro6/. The human and chimpanzee genomes

used are available here: https://hgdownload.soe.ucsc.edu/goldenPath/hg38/bigZips/ and here:

https://hgdownload.soe.ucsc.edu/goldenPath/panTro6/bigZips/ respectively. The 241-way PhyloP

scores were downloaded from here:

https://hgdownload.soe.ucsc.edu/goldenPath/hg38/cactus241way/. The "QCed genomic constraint by

1kb regions" from the gnomAD consortium are available here:

https://gnomad.broadinstitute.org/downloads#v3. All scripts for performing analyses and making

figures in this manuscript is publicly available at https://github.com/banwang27/multi-celltypes.

Competing interests: The authors have no competing interests to declare.

Funding: Funding for this work came from NIH R01HG012285. A.L.S was supported by an NDSEG

fellowship.

Author contributions: HBF conceived of the study. ALS and BW performed all analysis, visualization,

validation, and writing of software. ALS wrote the manuscript with input from BW and HBF. BW made all

figures with input from ALS and HBF. All authors approved the publication of the manuscript.

Acknowledgements: The authors wish to acknowledge the by the Columbia Stem Cell Core for their

hard work in differentiating these cells. We also acknowledge members of the Fraser lab past and

present for helpful discussion. BioRender was used to generate Figures 1a, 2a, and 3a.

Agoglia, R. M., Sun, D., Birey, F., Yoon, S.-J., Miura, Y., Sabatini, K., Pașca, S. P., & Fraser, H.

B. (2021). Primate cell fusion disentangles gene regulatory divergence in

neurodevelopment. *Nature*, *592* (7854), 421–427. https://doi.org/10.1038/s41586-021-

03343-3

Akbar, M., Calderon, F., Wen, Z., & Kim, H.-Y. (2005). Docosahexaenoic acid: A positive

modulator of Akt signaling in neuronal survival. *Proceedings of the National Academy of*

Sciences, *102* (31), 10858–10863. https://doi.org/10.1073/pnas.0502903102

Amemiya, H. M., Kundaje, A., & Boyle, A. P. (2019). The ENCODE Blacklist: Identification of

Problematic Regions of the Genome. *Scientific Reports*, *9* (1), 9354.

https://doi.org/10.1038/s41598-019-45839-z

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq—A Python framework to work with high-

throughput sequencing data. *Bioinformatics (Oxford, England)*, *31* (2), 166–169.

https://doi.org/10.1093/bioinformatics/btu638

Arai, Y., Funatsu, N., Numayama-Tsuruta, K., Nomura, T., Nakamura, S., & Osumi, N. (2005).

Role of Fabp7, a downstream gene of Pax6, in the maintenance of neuroepithelial cells

during early embryonic development of the rat cortex. *The Journal of Neuroscience: The*

Official Journal of the Society for Neuroscience, *25* (42), 9752–9761.

https://doi.org/10.1523/JNEUROSCI.2512-05.2005

Ben-Ari, Y., Khalilov, I., Kahle, K. T., & Cherubini, E. (2012). The GABA Excitatory/Inhibitory

Shift in Brain Maturation and Neurological Disorders. *The Neuroscientist*, *18* (5), 467–

486. https://doi.org/10.1177/1073858412438697

Benito-Kwiecinski, S., Giandomenico, S. L., Sutcliffe, M., Riis, E. S., Freire-Pritchett, P., Kelava,

I., Wunderlich, S., Martin, U., Wray, G. A., McDole, K., & Lancaster, M. A. (2021). An

early cell shape transition drives evolutionary expansion of the human forebrain. *Cell*,

184 (8), 2084-2102.e19. https://doi.org/10.1016/j.cell.2021.02.050

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological), *57* (1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Blake, L. E., Roux, J., Hernando-Herraez, I., Banovich, N. E., Perez, R. G., Hsiao, C. J., Eres, I.,

Cuevas, C., Marques-Bonet, T., & Gilad, Y. (2020). A comparison of gene expression

and DNA methylation patterns across tissues and species. *Genome Research*, *30* (2),

250–262. https://doi.org/10.1101/gr.254904.119

Broad Institute. (n.d.). *Picard*.

Brose, N., Brunger, A., Cafiso, D., Chapman, E. R., Diao, J., Hughson, F. M., Jackson, M. B.,

Jahn, R., Lindau, M., Ma, C., Rizo, J., Shin, Y.-K., Söllner, T. H., Tamm, L., Yoon, T.-Y.,

& Zhang, Y. (2019). Synaptic vesicle fusion: Today and beyond. *Nature Structural &*

Molecular Biology, *26* (8), 663–668. https://doi.org/10.1038/s41594-019-0277-z

Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y., & Greenleaf, W. J. (2013).

Transposition of native chromatin for fast and sensitive epigenomic profiling of open

chromatin, DNA-binding proteins and nucleosome position. *Nature Methods*, *10* (12),

1213–1218. https://doi.org/10.1038/nmeth.2688

Burridge, P. W., Matsa, E., Shukla, P., Lin, Z. C., Churko, J. M., Ebert, A. D., Lan, F., Diecke,

S., Huber, B., Mordwinkin, N. M., Plews, J. R., Abilez, O. J., Cui, B., Gold, J. D., & Wu, J.

C. (2014). Chemically defined generation of human cardiomyocytes. *Nature Methods*, *11*

(8), 855–860. https://doi.org/10.1038/nmeth.2999

Calderari, S., Ria, M., Gérard, C., Nogueira, T. C., Villate, O., Collins, S. C., Neil, H., Gervasi,

N., Hue, C., Suarez-Zamorano, N., Prado, C., Cnop, M., Bihoreau, M.-T., Kaisaki, P. J.,

Cazier, J.-B., Julier, C., Lathrop, M., Werner, M., Eizirik, D. L., & Gauguier, D. (2018).

Molecular genetics of the transcription factor GLIS3 identifies its dual function in beta

cells and neurons. *Genomics*, *110* (2), 98–111.

https://doi.org/10.1016/j.ygeno.2017.09.001

Caporali, A., & Emanueli, C. (2009). Cardiovascular actions of neurotrophins. *Physiological*

Reviews, *89* (1), 279–308. https://doi.org/10.1152/physrev.00007.2008

Castel, S. E., Aguet, F., Mohammadi, P., GTEx Consortium, Aguet, F., Anand, S., Ardlie, K. G.,

Gabriel, S., Getz, G. A., Graubert, A., Hadley, K., Handsaker, R. E., Huang, K. H.,

Kashin, S., Li, X., MacArthur, D. G., Meier, S. R., Nedzel, J. L., Nguyen, D. T., …

Lappalainen, T. (2020). A vast resource of allelic expression data spanning human

tissues. *Genome Biology*, *21* (1), 234. https://doi.org/10.1186/s13059-020-02122-z

Castro-Mondragon, J. A., Riudavets-Puig, R., Rauluseviciute, I., Lemma, R. B., Turchi, L.,

Blanc-Mathieu, R., Lucas, J., Boddie, P., Khan, A., Manosalva Pérez, N., Fornes, O.,

Leung, T. Y., Aguirre, A., Hammal, F., Schmelter, D., Baranasic, D., Ballester, B.,

Sandelin, A., Lenhard, B., … Mathelier, A. (2022). JASPAR 2022: The 9th release of the

open-access database of transcription factor binding profiles. *Nucleic Acids Research*,

50 (D1), D165–D173. https://doi.org/10.1093/nar/gkab1113

Chal, J., Al Tanoury, Z., Hestin, M., Gobert, B., Aivio, S., Hick, A., Cherrier, T., Nesmith, A. P.,

Parker, K. K., & Pourquié, O. (2016). Generation of human muscle fibers and satellite-

like cells from human pluripotent stem cells in vitro. *Nature Protocols*, *11* (10), 1833–

1850. https://doi.org/10.1038/nprot.2016.110

Chen, K. M., Wong, A. K., Troyanskaya, O. G., & Zhou, J. (2022). A sequence-based global

map of regulatory activity for deciphering human genetics. *Nature Genetics*, *54* (7), 940–

949. https://doi.org/10.1038/s41588-022-01102-2

Chen, S., Francioli, L. C., Goodrich, J. K., Collins, R. L., Kanai, M., Wang, Q., Alföldi, J., Watts,

N. A., Vittal, C., Gauthier, L. D., Poterba, T., Wilson, M. W., Tarasova, Y., Phu, W.,

Yohannes, M. T., Koenig, Z., Farjoun, Y., Banks, E., Donnelly, S., … Karczewski, K. J.

(2022). *A genome-wide mutational constraint map quantified from variation in 76,156*

human genomes [Preprint]. Genetics. https://doi.org/10.1101/2022.03.20.485034

Choi, W.-S., Xu, X., Goruk, S., Wang, Y., Patel, S., Chow, M., Field, C. J., & Godbout, R.

(2021). FABP7 Facilitates Uptake of Docosahexaenoic Acid in Glioblastoma Neural

Stem-like Cells. *Nutrients*, *13* (8), 2664. https://doi.org/10.3390/nu13082664

Collins, R. L., Glessner, J. T., Porcu, E., Lepamets, M., Brandon, R., Lauricella, C., Han, L.,

Morley, T., Niestroj, L.-M., Ulirsch, J., Everett, S., Howrigan, D. P., Boone, P. M., Fu, J.,

Karczewski, K. J., Kellaris, G., Lowther, C., Lucente, D., Mohajeri, K., … Talkowski, M.

E. (2022). A cross-disorder dosage sensitivity map of the human genome. *Cell*, *185* (16),

3041-3055.e25. https://doi.org/10.1016/j.cell.2022.06.036

Combs, P. A., Krupp, J. J., Khosla, N. M., Bua, D., Petrov, D. A., Levine, J. D., & Fraser, H. B.

(2018). Tissue-Specific cis-Regulatory Divergence Implicates eloF in Inhibiting

Interspecies Mating in Drosophila. *Current Biology*, *28* (24), 3969-3975.e3.

https://doi.org/10.1016/j.cub.2018.10.036

Conway, J. R., Lex, A., & Gehlenborg, N. (2017). UpSetR: An R package for the visualization of

intersecting sets and their properties. *Bioinformatics*, *33* (18), 2938–2940.

https://doi.org/10.1093/bioinformatics/btx364

Corces, M. R., Trevino, A. E., Hamilton, E. G., Greenside, P. G., Sinnott-Armstrong, N. A.,

Vesuna, S., Satpathy, A. T., Rubin, A. J., Montine, K. S., Wu, B., Kathiria, A., Cho, S.

W., Mumbach, M. R., Carter, A. C., Kasowski, M., Orloff, L. A., Risca, V. I., Kundaje, A.,

Khavari, P. A., … Chang, H. Y. (2017). An improved ATAC-seq protocol reduces

background and enables interrogation of frozen tissues. *Nature Methods*, *14* (10), 959–

962. https://doi.org/10.1038/nmeth.4396

De Rosa, A., Pellegatta, S., Rossi, M., Tunici, P., Magnoni, L., Speranza, M. C., Malusa, F.,

Miragliotta, V., Mori, E., Finocchiaro, G., & Bakker, A. (2012). A radial glia gene marker,

fatty acid binding protein 7 (FABP7), is involved in proliferation and invasion of

glioblastoma cells. *PloS One*, *7* (12), e52113.

https://doi.org/10.1371/journal.pone.0052113

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*

(Oxford, England), *29* (1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Ebrahimi, M., Yamamoto, Y., Sharifi, K., Kida, H., Kagawa, Y., Yasumoto, Y., Islam, A.,

Miyazaki, H., Shimamoto, C., Maekawa, M., Mitsushima, D., Yoshikawa, T., & Owada,

Y. (2016). Astrocyte-expressed FABP7 regulates dendritic morphology and excitatory

synaptic function of cortical neurons. *Glia*, *64* (1), 48–62.

https://doi.org/10.1002/glia.22902

Ernst, J., & Kellis, M. (2017). Chromatin-state discovery and genome annotation with

ChromHMM. *Nature Protocols*, *12* (12), 2478–2492.

https://doi.org/10.1038/nprot.2017.124

Feldblum, S., Erlander, M. G., & Tobin, A. J. (1993). Different distributions of GAD65 and

GAD67 mRNAS suggest that the two glutamate decarboxylases play distinctive

functional roles. *Journal of Neuroscience Research*, *34* (6), 689–706.

https://doi.org/10.1002/jnr.490340612

Field, A. R., Jacobs, F. M. J., Fiddes, I. T., Phillips, A. P. R., Reyes-Ortiz, A. M., LaMontagne,

E., Whitehead, L., Meng, V., Rosenkrantz, J. L., Olsen, M., Hauessler, M., Katzman, S.,

Salama, S. R., & Haussler, D. (2019). Structurally Conserved Primate LncRNAs Are

Transiently Expressed during Human Cortical Differentiation and Influence Cell-Type-

Specific Genes. *Stem Cell Reports*, *12* (2), 245–257.

https://doi.org/10.1016/j.stemcr.2018.12.006

Fraser, H. B. (2011). Genome-wide approaches to the study of adaptive gene expression

evolution: Systematic studies of evolutionary adaptations involving gene expression will

allow many fundamental questions in evolutionary biology to be addressed. *BioEssays*,

33 (6), 469–477. https://doi.org/10.1002/bies.201000094

Fraser, H. B. (2013). Gene expression drives local adaptation in humans. *Genome Research*,

23 (7), 1089–1096. https://doi.org/10.1101/gr.152710.112

Fraser, H. B. (2019). Improving Estimates of Compensatory cis–trans Regulatory Divergence.

Trends in Genetics, *35* (1), 3–5. https://doi.org/10.1016/j.tig.2018.09.003

García-Pérez, R., Esteller-Cucala, P., Mas, G., Lobón, I., Di Carlo, V., Riera, M., Kuhlwilm, M.,

Navarro, A., Blancher, A., Di Croce, L., Gómez-Skarmeta, J. L., Juan, D., & Marquès-

Bonet, T. (2021). Epigenomic profiling of primate lymphoblastoid cell lines reveals the

evolutionary patterns of epigenetic activities in gene regulatory architectures. *Nature*

Communications, *12* (1), 3116. https://doi.org/10.1038/s41467-021-23397-1

Girskis, K. M., Stergachis, A. B., DeGennaro, E. M., Doan, R. N., Qian, X., Johnson, M. B.,

Wang, P. P., Sejourne, G. M., Nagy, M. A., Pollina, E. A., Sousa, A. M. M., Shin, T.,

Kenny, C. J., Scotellaro, J. L., Debo, B. M., Gonzalez, D. M., Rento, L. M., Yeh, R. C.,

Song, J. H. T., … Walsh, C. A. (2021). Rewiring of human neurodevelopmental gene

regulatory programs by human accelerated regions. *Neuron*, *109* (20), 3239-3251.e7.

https://doi.org/10.1016/j.neuron.2021.08.005

Gokhman, D., Agoglia, R. M., Kinnebrew, M., Gordon, W., Sun, D., Bajpai, V. K., Naqvi, S.,

Chen, C., Chan, A., Chen, C., Petrov, D. A., Ahituv, N., Zhang, H., Mishina, Y.,

Wysocka, J., Rohatgi, R., & Fraser, H. B. (2021). Human–chimpanzee fused cells reveal

cis-regulatory divergence underlying skeletal evolution. *Nature Genetics*, *53* (4), 467–

476. https://doi.org/10.1038/s41588-021-00804-3

GTEx Consortium. (2017). Genetic effects on gene expression across human tissues. *Nature*,

550 (7675), 204–213. https://doi.org/10.1038/nature24277

Horlbeck, M. A., Gilbert, L. A., Villalta, J. E., Adamson, B., Pak, R. A., Chen, Y., Fields, A. P.,

Park, C. Y., Corn, J. E., Kampmann, M., & Weissman, J. S. (2016). Compact and highly

active next-generation libraries for CRISPR-mediated gene repression and activation.

ELife, *5*, e19760. https://doi.org/10.7554/eLife.19760

Hu, C. K., York, R. A., Metz, H. C., Bedford, N. L., Fraser, H. B., & Hoekstra, H. E. (2022). Cis-

Regulatory changes in locomotor genes are associated with the evolution of burrowing

behavior. *Cell Reports*, *38* (7), 110360. https://doi.org/10.1016/j.celrep.2022.110360

Huang, E. J., & Reichardt, L. F. (2001). Neurotrophins: Roles in neuronal development and

function. *Annual Review of Neuroscience*, *24*, 677–736.

https://doi.org/10.1146/annurev.neuro.24.1.677

Hubisz, M. J., & Pollard, K. S. (2014). Exploring the genesis and functions of Human

Accelerated Regions sheds light on their role in human evolution. *Current Opinion in*

Genetics & Development, *29*, 15–21. https://doi.org/10.1016/j.gde.2014.07.005

Ichim, G., Tauszig-Delamasure, S., & Mehlen, P. (2012). Neurotrophins and cell death.

Experimental Cell Research, *318* (11), 1221–1228.

https://doi.org/10.1016/j.yexcr.2012.03.006

Jain, A., & Tuteja, G. (2019). TissueEnrich: Tissue-specific gene enrichment analysis.

Bioinformatics (Oxford, England), *35* (11), 1966–1967.

https://doi.org/10.1093/bioinformatics/bty890

John St. John. (n.d.). *SeqPrep*.

Kanton, S., Boyle, M. J., He, Z., Santel, M., Weigert, A., Sanchís-Calleja, F., Guijarro, P., Sidow,

L., Fleck, J. S., Han, D., Qian, Z., Heide, M., Huttner, W. B., Khaitovich, P., Pääbo, S.,

Treutlein, B., & Camp, J. G. (2019). Organoid single-cell genomic atlas uncovers human-

specific features of brain development. *Nature*, *574* (7778), 418–422.

https://doi.org/10.1038/s41586-019-1654-9

Ke, K., Song, Y., Shen, J., Niu, M., Zhang, H., Yuan, D., Ni, H., Zhang, Y., Liu, X., Dai, A., &

Cao, M. (2015). Up-Regulation of Glis2 Involves in Neuronal Apoptosis After

Intracerebral Hemorrhage in Adult Rats. *Cellular and Molecular Neurobiology*, *35* (3),

345–354. https://doi.org/10.1007/s10571-014-0130-1

Kelley, J. L., & Gilad, Y. (2020). Effective study design for comparative functional genomics.

Nature Reviews Genetics, *21* (7), 385–386. https://doi.org/10.1038/s41576-020-0242-z

King, M.-C., & Wilson, A. C. (1975). Evolution at Two Levels in Humans and Chimpanzees:

Their macromolecules are so alike that regulatory mutations may account for their

biological differences. *Science*, *188* (4184), 107–116.

https://doi.org/10.1126/science.1090005

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner,

M., Loh, P., & Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of

single-cell data with Harmony. *Nature Methods*, *16* (12), 1289–1296.

https://doi.org/10.1038/s41592-019-0619-0

Korytnikov, R., & Nostro, M. C. (2016). Generation of polyhormonal and multipotent pancreatic

progenitor lineages from human pluripotent stem cells. *Methods*, *101*, 56–64.

https://doi.org/10.1016/j.ymeth.2015.10.017

Kozlenkov, A., Vermunt, M. W., Apontes, P., Li, J., Hao, K., Sherwood, C. C., Hof, P. R., Ely, J.

J., Wegner, M., Mukamel, E. A., Creyghton, M. P., Koonin, E. V., & Dracheva, S. (2020).

Evolution of regulatory signatures in primate cortical neurons at cell-type resolution.

Proceedings of the National Academy of Sciences, *117* (45), 28422–28432.

https://doi.org/10.1073/pnas.2011884117

Kuhn, R. M., Haussler, D., & Kent, W. J. (2013). The UCSC genome browser and associated

tools. *Briefings in Bioinformatics*, *14* (2), 144–161. https://doi.org/10.1093/bib/bbs038

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, *9* (4), 357–359. https://doi.org/10.1038/nmeth.1923

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence

Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*, *25* (16), 2078–

2079. https://doi.org/10.1093/bioinformatics/btp352

Liang, D., Elwell, A. L., Aygün, N., Krupa, O., Wolter, J. M., Kyere, F. A., Lafferty, M. J., Cheek, K. E., Courtney, K. P., Yusupova, M., Garrett, M. E., Ashley-Koch, A., Crawford, G. E., Love, M. I., de la Torre-Ubieta, L., Geschwind, D. H., & Stein, J. L. (2021). Cell-type-specific effects of genetic variation on chromatin accessibility during human neuronal differentiation. *Nature Neuroscience*, *24* (7), 941–953. https://doi.org/10.1038/s41593-021-00858-w

Lopez-Delisle, L., Rabbani, L., Wolff, J., Bhardwaj, V., Backofen, R., Grüning, B., Ramírez, F., & Manke, T. (2021). pyGenomeTracks: Reproducible plots for multivariate genomic datasets. *Bioinformatics (Oxford, England)*, *37* (3), 422–423. https://doi.org/10.1093/bioinformatics/btaa692

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15* (12), 550. https://doi.org/10.1186/s13059-014-0550-8

Ma, S., Skarica, M., Li, Q., Xu, C., Risgaard, R. D., Tebbenkamp, A. T. N., Mato-Blanco, X., Kovner, R., Krsnik, Ž., de Martin, X., Luria, V., Martí-Pérez, X., Liang, D., Karger, A., Schmidt, D. K., Gomez-Sanchez, Z., Qi, C., Gobeske, K. T., Pochareddy, S., … Sestan, N. (2022). Molecular and cellular evolution of the primate dorsolateral prefrontal cortex. *Science*, eabo7257. https://doi.org/10.1126/science.abo7257

Mack, K. L., & Nachman, M. W. (2017). Gene Regulation and Speciation. *Trends in Genetics*, *33* (1), 68–80. https://doi.org/10.1016/j.tig.2016.11.003

Mallanna, S. K., & Duncan, S. A. (2013). Differentiation of hepatocytes from pluripotent stem cells. *Current Protocols in Stem Cell Biology*, *26*, 1G.4.1-1G.4.13. https://doi.org/10.1002/9780470151808.sc01g04s26

Maury, Y., Côme, J., Piskorowski, R. A., Salah-Mohellibi, N., Chevaleyre, V., Peschanski, M., Martinat, C., & Nedelec, S. (2015). Combinatorial analysis of developmental cues

efficiently converts human pluripotent stem cells into multiple neuronal subtypes. *Nature Biotechnology*, *33* (1), 89–96. https://doi.org/10.1038/nbt.3049

Meisler, M. H., Hill, S. F., & Yu, W. (2021). Sodium channelopathies in neurodevelopmental disorders. *Nature Reviews Neuroscience*, *22* (3), 152–166. https://doi.org/10.1038/s41583-020-00418-4

Milani, P., Escalante-Chong, R., Shelley, B. C., Patel-Murray, N. L., Xin, X., Adam, M., Mandefro, B., Sareen, D., Svendsen, C. N., & Fraenkel, E. (2016). Cell freezing protocol suitable for ATAC-Seq on motor neurons derived from human induced pluripotent stem cells. *Scientific Reports*, *6*, 25474. https://doi.org/10.1038/srep25474

Mizuguchi, R., Kriks, S., Cordes, R., Gossler, A., Ma, Q., & Goulding, M. (2006). Ascl1 and Gsh1/2 control inhibitory and excitatory cell fate in spinal sensory interneurons. *Nature Neuroscience*, *9* (6), 770–778. https://doi.org/10.1038/nn1706

Montaigne, D., Butruille, L., & Staels, B. (2021). PPAR control of metabolism and cardiovascular functions. *Nature Reviews Cardiology*, *18* (12), 809–823. https://doi.org/10.1038/s41569-021-00569-6

Mueller, Fabian. (n.d.). *ChrAccR*.

Naito, Y., Lee, A. K., & Takahashi, H. (2017). Emerging roles of the neurotrophin receptor TrkC in synapse organization. *Neuroscience Research*, *116*, 10–17. https://doi.org/10.1016/j.neures.2016.09.009

Netherlands Brain Bank, Vermunt, M. W., Tan, S. C., Castelijns, B., Geeven, G., Reinink, P., de Bruijn, E., Kondova, I., Persengiev, S., Bontrop, R., Cuppen, E., de Laat, W., & Creyghton, M. P. (2016). Epigenomic annotation of gene regulatory alterations during evolution of the primate brain. *Nature Neuroscience*, *19* (3), 494–503. https://doi.org/10.1038/nn.4229

Pavlovic, B. J., Blake, L. E., Roux, J., Chavarria, C., & Gilad, Y. (2018). A Comparative Assessment of Human and Chimpanzee iPSC-derived Cardiomyocytes with Primary

Heart Tissues. *Scientific Reports*, *8* (1), 15312. https://doi.org/10.1038/s41598-018-

33478-9

Pollard, K. S., Salama, S. R., Lambert, N., Lambot, M.-A., Coppens, S., Pedersen, J. S.,

Katzman, S., King, B., Onodera, C., Siepel, A., Kern, A. D., Dehay, C., Igel, H., Ares, M.,

Vanderhaeghen, P., & Haussler, D. (2006). An RNA gene expressed during cortical

development evolved rapidly in humans. *Nature*, *443* (7108), 167–172.

https://doi.org/10.1038/nature05113

Prud'homme, B., Gompel, N., & Carroll, S. B. (2007). Emerging principles of regulatory

evolution. *Proceedings of the National Academy of Sciences*, *104* (suppl_1), 8605–8612.

https://doi.org/10.1073/pnas.0700488104

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic

features. *Bioinformatics (Oxford, England)*, *26* (6), 841–842.

https://doi.org/10.1093/bioinformatics/btq033

Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A., & Manke, T. (2014). deepTools: A flexible

platform for exploring deep-sequencing data. *Nucleic Acids Research*, *42* (Web Server

issue), W187-191. https://doi.org/10.1093/nar/gku365

Reilly, S. K., & Noonan, J. P. (2016). Evolution of Gene Regulation in Humans. *Annual Review

of Genomics and Human Genetics*, *17*, 45–67. https://doi.org/10.1146/annurev-genom-

090314-045935

Romero, I. G., Ruvinsky, I., & Gilad, Y. (2012). Comparative studies of gene expression and the

evolution of gene regulation. *Nature Reviews Genetics*, *13* (7), 505–516.

https://doi.org/10.1038/nrg3229

Sharma, R., Khristov, V., Rising, A., Jha, B. S., Dejene, R., Hotaling, N., Li, Y., Stoddard, J.,

Stankewicz, C., Wan, Q., Zhang, C., Campos, M. M., Miyagishima, K. J., McGaughey,

D., Villasmil, R., Mattapallil, M., Stanzel, B., Qian, H., Wong, W., … Bharti, K. (2019).

Clinical-grade stem cell-derived retinal pigment epithelium patch rescues retinal

degeneration in rodents and pigs. *Science Translational Medicine*, *11* (475), eaat5580.

https://doi.org/10.1126/scitranslmed.aat5580

Shave, R. E., Lieberman, D. E., Drane, A. L., Brown, M. G., Batterham, A. M., Worthington, S.,

Atencia, R., Feltrer, Y., Neary, J., Weiner, R. B., Wasfy, M. M., & Baggish, A. L. (2019).

Selection of endurance capabilities and the trade-off between pressure and volume in

the evolution of the human heart. *Proceedings of the National Academy of Sciences*,

116 (40), 19905–19910. https://doi.org/10.1073/pnas.1906902116

Song, J. H. T., Grant, R. L., Behrens, V. C., Kučka, M., Roberts Kingman, G. A., Soltys, V.,

Chan, Y. F., & Kingsley, D. M. (2021). Genetic studies of human-chimpanzee divergence

using stem cell fusions. *Proceedings of the National Academy of Sciences of the United*

States of America, *118* (51), e2117557118. https://doi.org/10.1073/pnas.2117557118

Starr, A. L., Gokhman, D., & Fraser, H. B. (2023). Accounting for cis-regulatory constraint

prioritizes genes likely to affect species-specific traits. *Genome Biology*, *24* (1), 11.

https://doi.org/10.1186/s13059-023-02846-8

Storey Lab. (n.d.). *Qvalue*.

Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A.,

Paulovich, A., Pomeroy, S. L., Golub, T. R., Lander, E. S., & Mesirov, J. P. (2005). Gene

set enrichment analysis: A knowledge-based approach for interpreting genome-wide

expression profiles. *Proceedings of the National Academy of Sciences*, *102* (43), 15545–

15550. https://doi.org/10.1073/pnas.0506580102

Su-Feher, L., Rubin, A. N., Silberberg, S. N., Catta-Preta, R., Lim, K. J., Ypsilanti, A. R., Zdilar,

I., McGinnis, C. S., McKinsey, G. L., Rubino, T. E., Hawrylycz, M. J., Thompson, C.,

Gartner, Z. J., Puelles, L., Zeng, H., Rubenstein, J. L. R., & Nord, A. S. (2022). Single

cell enhancer activity distinguishes GABAergic and cholinergic lineages in embryonic

mouse basal ganglia. *Proceedings of the National Academy of Sciences*, *119* (15),

e2108760119. https://doi.org/10.1073/pnas.2108760119

Sullivan, P. F., Meadows, J. R. S., Gazal, S., Phan, B. N., Li, X., Genereux, D. P., Dong, M. X.,

Bianchi, M., Andrews, G., Sakthikumar, S., Nordin, J., Roy, A., Christmas, M. J.,

Marinescu, V. D., Wallerman, O., Xue, J. R., Li, Y., Yao, S., Sun, Q., … Lindblad-Toh, K.

(2023). *Leveraging Base Pair Mammalian Constraint to Understand Genetic Variation*

and Human Disease [Preprint]. Genomics. https://doi.org/10.1101/2023.03.10.531987

Traag, V. A., Waltman, L., & van Eck, N. J. (2019). From Louvain to Leiden: Guaranteeing well-

connected communities. *Scientific Reports*, *9* (1), 5233. https://doi.org/10.1038/s41598-

019-41695-z

Trevino, A. E., Sinnott-Armstrong, N., Andersen, J., Yoon, S.-J., Huber, N., Pritchard, J. K.,

Chang, H. Y., Greenleaf, W. J., & Paşca, S. P. (2020). Chromatin accessibility dynamics

in a model of human forebrain development. *Science (New York, N.Y.)*, *367* (6476),

eaay1645. https://doi.org/10.1126/science.aay1645

Trizzino, M., Park, Y., Holsbach-Beltrame, M., Aracena, K., Mika, K., Caliskan, M., Perry, G. H.,

Lynch, V. J., & Brown, C. D. (2017). Transposable elements are the primary source of

novelty in primate gene regulation. *Genome Research*, *27* (10), 1623–1633.

https://doi.org/10.1101/gr.218149.116

van de Geijn, B., McVicker, G., Gilad, Y., & Pritchard, J. K. (2015). WASP: Allele-specific

software for robust molecular quantitative trait locus discovery. *Nature Methods*, *12* (11),

1061–1063. https://doi.org/10.1038/nmeth.3582

Vanderhaeghen, P., & Polleux, F. (2023). Developmental mechanisms underlying the evolution

of human cortical circuits. *Nature Reviews Neuroscience*, *24* (4), 213–232.

https://doi.org/10.1038/s41583-023-00675-z

Vierstra, J., Lazar, J., Sandstrom, R., Halow, J., Lee, K., Bates, D., Diegel, M., Dunn, D., Neri,

F., Haugen, E., Rynes, E., Reynolds, A., Nelson, J., Johnson, A., Frerker, M., Buckley,

M., Kaul, R., Meuleman, W., & Stamatoyannopoulos, J. A. (2020). Global reference

mapping of human transcription factor footprints. *Nature*, *583* (7818), 729–736.

https://doi.org/10.1038/s41586-020-2528-x

Vu, H., & Ernst, J. (2022). Universal annotation of the human genome through integration of

over a thousand epigenomic datasets. *Genome Biology*, *23* (1), 9.

https://doi.org/10.1186/s13059-021-02572-z

Wang, B., & Tontonoz, P. (2018). Liver X receptors in lipid signalling and membrane

homeostasis. *Nature Reviews Endocrinology*, *14* (8), 452–463.

https://doi.org/10.1038/s41574-018-0037-x

Watanabe, A., Toyota, T., Owada, Y., Hayashi, T., Iwayama, Y., Matsumata, M., Ishitsuka, Y.,

Nakaya, A., Maekawa, M., Ohnishi, T., Arai, R., Sakurai, K., Yamada, K., Kondo, H.,

Hashimoto, K., Osumi, N., & Yoshikawa, T. (2007). Fabp7 Maps to a Quantitative Trait

Locus for a Schizophrenia Endophenotype. *PLoS Biology*, *5* (11), e297.

https://doi.org/10.1371/journal.pbio.0050297

Wittkopp, P. J., Haerum, B. K., & Clark, A. G. (2004). Evolutionary changes in cis and trans

gene regulation. *Nature*, *430* (6995), 85–88. https://doi.org/10.1038/nature02698

Wittkopp, P. J., & Kalay, G. (2012). Cis-regulatory elements: Molecular mechanisms and

evolutionary processes underlying divergence. *Nature Reviews Genetics*, *13* (1), 59–69.

https://doi.org/10.1038/nrg3095

Wolf, F. A., Angerer, P., & Theis, F. J. (2018). SCANPY: Large-scale single-cell gene

expression data analysis. *Genome Biology*, *19* (1), 15. https://doi.org/10.1186/s13059-

017-1382-0

Wolf, F. A., Hamey, F. K., Plass, M., Solana, J., Dahlin, J. S., Göttgens, B., Rajewsky, N.,

Simon, L., & Theis, F. J. (2019). PAGA: Graph abstraction reconciles clustering with

trajectory inference through a topology preserving map of single cells. *Genome Biology*,

20 (1), 59. https://doi.org/10.1186/s13059-019-1663-x

Yang, N., Chanda, S., Marro, S., Ng, Y.-H., Janas, J. A., Haag, D., Ang, C. E., Tang, Y., Flores,

Q., Mall, M., Wapinski, O., Li, M., Ahlenius, H., Rubenstein, J. L., Chang, H. Y., Buylla,

A. A., Südhof, T. C., & Wernig, M. (2017). Generation of pure GABAergic neurons by

transcription factor programming. *Nature Methods*, *14* (6), 621–628.

https://doi.org/10.1038/nmeth.4291

Yao, Z., van Velthoven, C. T. J., Nguyen, T. N., Goldy, J., Sedeno-Cortes, A. E., Baftizadeh, F.,

Bertagnolli, D., Casper, T., Chiang, M., Crichton, K., Ding, S.-L., Fong, O., Garren, E.,

Glandon, A., Gouwens, N. W., Gray, J., Graybuck, L. T., Hawrylycz, M. J., Hirschstein,

D., … Zeng, H. (2021). A taxonomy of transcriptomic cell types across the isocortex and

hippocampal formation. *Cell*, *184* (12), 3222-3241.e26.

https://doi.org/10.1016/j.cell.2021.04.021

York, R. A., Patil, C., Abdilleh, K., Johnson, Z. V., Conte, M. A., Genner, M. J., McGrath, P. T.,

Fraser, H. B., Fernald, R. D., & Streelman, J. T. (2018). Behavior-dependent *cis*

regulation reveals genes and pathways associated with bower building in cichlid fishes.

Proceedings of the National Academy of Sciences, *115* (47).

https://doi.org/10.1073/pnas.1810140115

Yu, X., Lin, J., Zack, D. J., & Qian, J. (2006). Computational analysis of tissue-specific

combinatorial gene regulation: Predicting interaction between transcription factors in

human tissues. *Nucleic Acids Research*, *34* (17), 4925–4936.

https://doi.org/10.1093/nar/gkl595

Zhang, S., Zhang, H., Zhou, Y., Qiao, M., Zhao, S., Kozlova, A., Shi, J., Sanders, A. R., Wang,

G., Luo, K., Sengupta, S., West, S., Qian, S., Streit, M., Avramopoulos, D., Cowan, C.

A., Chen, M., Pang, Z. P., Gejman, P. V., … Duan, J. (2020). Allele-specific open

chromatin in human iPSC neurons elucidates functional disease variants. *Science*, *369*

(6503), 561–565. https://doi.org/10.1126/science.aay3983

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nusbaum, C.,

Myers, R. M., Brown, M., Li, W., & Liu, X. S. (2008). Model-based analysis of ChIP-Seq

(MACS). *Genome Biology*, *9* (9), R137. https://doi.org/10.1186/gb-2008-9-9-r137

Zhu, A., Ibrahim, J. G., & Love, M. I. (2019). Heavy-tailed prior distributions for sequence count

data: Removing the noise and preserving large differences. *Bioinformatics*, *35* (12),

2084–2092. https://doi.org/10.1093/bioinformatics/bty895

1326