The carbapenem inoculum effect provides insight into the molecular mechanisms underlying carbapenem resistance in *Enterobacterales*

Alexis Jaramillo Cartagena[1†], Kyra L. Taylor[1†], Joshua T. Smith[1], Abigail L. Manson[1], Virginia M. Pierce[2,3], Ashlee M. Earl[1], Roby P. Bhattacharyya[1,4,*]

[1]Infectious Disease and Microbiome Program, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
[2]Microbiology Laboratory, Department of Pathology, Massachusetts General Hospital, Boston, MA 02114, USA
[3]Department of Pathology and Clinical Laboratories, University of Michigan, Ann Arbor, MI 48109, USA
[4]Infectious Diseases Division, Department of Medicine, Massachusetts General Hospital, Boston, MA 02114, USA
[†]These authors contributed equally to this work.

*Corresponding Author

Email: rbhatt@broadinstitute.org

Author Contributions:
Designed research: AJC, KLT, RPB
Performed research: AJC, KLT, RPB
Contributed new reagents/analytic tools: AJC, KLT, RPB, AME
Analyzed data: AJC, KLT, RPB, JTS, ALM, VMP, AME
Wrote the paper: AJC, KLT, RPB, JTS, ALM, VMP, AME

Classification: Biological Sciences / Microbiology

Keywords: Antibiotic resistance, carbapenemases, inoculum effect, bacterial genomics, clinical microbiology

Abstract

Carbapenem-resistant *Enterobacterales* (CRE) are important pathogens that can develop

resistance via multiple molecular mechanisms, including hydrolysis or reduced antibiotic influx.

Identifying these mechanisms can improve pathogen surveillance, infection control, and patient

care. We investigated how resistance mechanisms influence the carbapenem inoculum effect

(IE), a phenomenon where inoculum size affects antimicrobial susceptibility testing (AST). We

demonstrated that seven different carbapenemases impart a meropenem IE in *Escherichia coli*.

Across 110 clinical CRE isolates, the carbapenem IE strictly depended on resistance

mechanism: all carbapenemase-producing CRE (CP-CRE) exhibited a strong IE, whereas

porin-deficient CRE displayed none. Concerningly, 50% and 24% of CP-CRE isolates changed

susceptibility classification to meropenem and ertapenem, respectively, across the allowable

inoculum range in clinical guidelines. The meropenem IE, and the ratio of ertapenem to

meropenem minimal inhibitory concentration (MIC) at standard inoculum, reliably identified CP-

CRE. Understanding how resistance mechanisms affect AST could improve diagnosis and

guide therapies for CRE infections.

Main Text

Introduction

Carbapenems are a crucial class of β-lactam antibiotics with a broad spectrum of activity (1) due

to their ability to withstand hydrolysis by many β-lactamases and their inhibition of cell wall

synthesis across many bacterial species (2). The emergence of carbapenem-resistant

Enterobacterales (CRE) poses a significant public health risk, with the US Centers for Disease

Control and Prevention (CDC) designating them as an "urgent threat" due to their extensive

drug resistance, high mortality rates, and treatment challenges (3, 4). The rapid global

emergence of CRE has created an urgent clinical and epidemiological need to better

understand, detect, and limit the spread of these pathogens (5).

CRE employ two major molecular mechanisms of carbapenem resistance: 1) expression

of carbapenemases, which efficiently hydrolyze carbapenems, or 2) disruption of porins, which

reduces influx of carbapenems into the periplasm where they act (6). This reduced periplasmic

influx can potentiate β-lactamases with weak carbapenemase activity such as AmpC, CTX-M,

SHV, TEM, or OXA-10, and together cause resistance (1, 7). Efflux pumps can also contribute

to carbapenem resistance, though their role has been better characterized in non-

Enterobacterales species like *Acinetobacter baumannii* and *Pseudomonas aeruginosa* (8, 9).

Carbapenemase-producing CRE (CP-CRE) are of particular epidemiological concern due to

their propensity to cause outbreaks (4, 10) and the risk of horizontal transfer of carbapenemase

genes (11–13). Porin deficient CRE (PD-CRE) often exhibit impaired fitness (14) compared to

CP-CRE because porins play important roles in nutrient import (15), osmotic homeostasis (14,

16), and membrane lipid asymmetry (17). Determining resistance mechanisms could lead to

better infection control measures, such as patient isolation to prevent CP-CRE spread in

hospitals (4, 13, 18). Some laboratories test for resistance mechanisms in CRE (19–21), but this

requires additional time, equipment, expertise, and expenses. Capitalizing on existing

antimicrobial susceptibility testing (AST) workflows to predict CRE resistance mechanisms

would represent a clinical advance.

Standardized AST methods and interpretive criteria are crucial to determine

antimicrobial susceptibility of pathogens. Organizations like the Clinical and Laboratory

Standards Institute (CLSI) classify pathogen susceptibility based on minimum inhibitory

concentration (MIC) breakpoints (22–24) that are established and periodically revised (25)

based on microbiological, pharmacokinetic-pharmacodynamic, and clinical data. While these

breakpoints are intended to predict clinical response to therapy, not resistance mechanism,

carbapenem MICs have been repurposed to prioritize isolates for molecular carbapenemase

testing (22, 26). In the United States, only isolates that test phenotypically carbapenem resistant

are typically tested for carbapenemases (22). Yet despite recently updated breakpoints (27),

some carbapenemase producing isolates can still be misclassified as susceptible to

carbapenems (28).

The inoculum effect (IE) is a phenomenon whereby the measured MIC varies depending

on the precise bacterial inoculum used in AST assays, which can lead to susceptibility

misclassification and treatment failures (29, 30). Multiple factors may contribute to the IE,

including metabolic state of the bacteria (31), growth productivity (32), antibiotic-target ratio (33),

heteroresistance frequency (34), target proteolysis (35), and production of enzymes that

inactivate antibiotics (36–40). The IE is frequently observed for some antibiotic classes including

β-lactams (41) but not commonly for others such as aminoglycosides. While importance of the

IE in clinical infections remains uncertain (42, 43), it has important implications for diagnostic

accuracy. To minimize the impact of the IE, CLSI recommends a standard inoculum of 5×10^5

colony forming units (CFU)/mL, with an acceptable range of 2×10^5 to 8×10^5 CFU/mL, for

broth microdilution, the gold standard AST method (44). Even within this range, however,

carbapenem susceptibility classification of *Enterobacterales* isolates may change due to the IE,

underscoring the need for continued efforts to optimize AST methods (45).

Here, we sought to systematically investigate how different genotypes conferring

carbapenem resistance influence the IE and how understanding these relationships might be

leveraged to improve clinical AST. We found that expression of distinct carbapenemases was

sufficient to cause a marked meropenem IE in a laboratory strain of *Escherichia coli*. Further, in

a collection of 110 clinical CRE isolates across 12 species of *Enterobacterales* (11, 46), we

observed that the carbapenem IE was a function of resistance mechanism. Specifically, all CP-

CRE exhibited a strong carbapenem IE, whereas all PD-CRE exhibited constant carbapenem

MICs across a wide range of inocula. Isolates with both a carbapenemase and porin deficiency

generally exhibited an IE and higher MICs than strains with either mechanism alone. The

carbapenem MICs of CP-CRE varied considerably, with a large proportion changing AST

classification across the CLSI-acceptable inoculum range. Collectively, our results

demonstrated that the IE distinguished CRE by resistance mechanism, better than either

meropenem or ertapenem MIC alone. These findings suggested two phenotypic measures that

predict carbapenem resistance mechanisms better than the MIC cutoffs currently used to

prioritize molecular carbapenemase testing.

Results

Carbapenemases belonging to all four Ambler classes impart an inoculum effect.

Carbapenemases are commonly divided into four classes based on protein sequence and

enzymatic features (47, 48). To investigate if each carbapenemase class is sufficient to confer

an IE, we systematically cloned seven different carbapenemase genes, each with its native

promoter, into a low-copy pBAD33 plasmid (49): bla_{KPC-3} and bla_{SME-2} from Ambler class A;

bla_{IMP-4}, bla_{NDM-1}, and bla_{VIM-27} from class B; bla_{CMY-10} (50) from class C; and bla_{OXA-48} from class D

(Supplementary Table S1). We transformed each carbapenemase-encoding plasmid into *E. coli*

K-12 MG1655, a carbapenem-susceptible laboratory strain, then conducted meropenem broth

microdilution MIC assays across a broad range of inocula, comprising two-fold dilutions from 1.3

$\times 10^7$ to 1.6×10^3 CFU/mL. As inoculum increased, all transformants showed a significant

increase in MIC, while the MIC of the empty vector control did not (Fig. 1*A*). Most transformants

had stable meropenem MICs from $\sim 10^3$ to $\sim 10^4$ CFU/mL, and all exhibited a strong linear

increase in MIC across the $\sim 10^5$ to $\sim 10^7$ CFU/mL inoculum range, which includes the

recommended inoculum range for clinical testing (Fig. 1*B*). In this strain background, NDM-1

(orange), SME-2 (pink), and IMP-4 (brown) imparted the highest levels of resistance, followed

by VIM-27 (yellow). KPC-3 (teal) conferred high-level resistance at high inocula, but within the

CLSI inoculum range (dashed lines in Fig. 1*B*) MICs remained within the intermediate

classification (>1 µg/mL and <4 µg/mL). OXA-48 (violet) and CMY-10 (green) conferred the

lowest meropenem MICs, which remained below 1 µg/mL until the highest inoculum tested and

well within the susceptible classification in the CLSI range.

Inoculum effect of clinical CRE isolates depends on resistance mechanism. After

establishing that all carbapenemases imparted an IE in *E. coli* K-12, we investigated the IE in

clinical isolates harboring carbapenemases, porin disruptions, or both. We analyzed 110 clinical

isolates comprising six genera and 12 species (Table S2), which we assigned to three groups

based on their genotypes: CP-CRE, PD-CRE, and CP-CRE with porin deficiency (Fig. 2 and

Supplementary Fig. S1, Table S2 and Table S3). We observed that all 36 CP-CRE harboring

carbapenemases from one of six major families (KPC, NDM, IMP, VIM, OXA-48, and SME)

displayed an IE similar to the *E. coli* K-12 carbapenemase transformants (Fig. 2*A* and

Supplementary Fig. S1*A*). In contrast, 47 PD-CRE exhibited a strikingly different pattern of

resistance, with nearly constant MICs across the tested inoculum range (Fig. 2*B* and

Supplementary Fig. S1*B*). Finally, the 27 isolates with both carbapenemase production and

porin deficiencies, which we termed "hyper-CRE", exhibited an IE and higher-level resistance

than isolates containing only one mechanism of resistance (Fig. 2*C* and Supplementary Fig.

S1*C*).

Carbapenemase production and porin deficiency together cause a hyper-CRE phenotype.

To directly test the hypothesis that the combination of carbapenemase production and porin

deficiency confers a hyper-CRE phenotype, we selected two clinical *Klebsiella pneumoniae*

isolates (BIDMC46a and BIDMC91) with disruptions in both major porins OmpK36 and

OmpK35, and transformed them with a pBAD33 plasmid encoding a KPC-3 (Fig. 3*A*). Both

transformants exhibited dramatic increases in MIC at standard inoculum (256-fold and 128-fold,

respectively) compared to empty vector controls and also developed an IE (6-fold from standard

to high inoculum). In a second experiment, we disrupted the major OmpK36 porin in a *K.*

pneumoniae isolate expressing a KPC-2 (RB582) using a CRISPR-Cas9 base editor (51).

Meropenem MICs at standard inoculum increased by 32-fold in the edited strain, with a small 2-

fold IE (Fig. 3*B*). These experiments provide direct evidence that the combination of

carbapenemase production and porin deficiency leads to a hyper-CRE phenotype.

A significant proportion of carbapenemase-producing isolates change classification

within the CLSI inoculum range. Although clinical microbiological laboratories initially

identified the isolates we studied as carbapenem-resistant, the IE caused the MIC of several

isolates to vary greatly within the CLSI inoculum range (Fig. 2*A*, and Supplementary Fig. S1*A*).

This variability prompted us to quantify which carbapenemase-producing isolates changed

susceptibility classification within the acceptable CLSI inoculum range of 2×10^5 CFU/mL to $8 \times$

10^5 CFU/mL. We tested both meropenem and ertapenem, two carbapenems in common clinical

use with distinct physicochemical and pharmacokinetic properties. Notably, ertapenem has a

lower barrier to resistance (52) since it is more affected by reduced outer membrane

permeability (53), and has lower clinical susceptibility breakpoints (22). 22 of 63 (35%)

carbapenemase-producing isolates tested with meropenem and 7 of 41 (17%) tested with

ertapenem changed AST classification (Fig. 4). When excluding hyper-CRE and considering

only isolates with intact porins, 18 of 36 (50%) CP-CRE isolates changed AST classification for

meropenem and 7 of 29 (24%) for ertapenem within the CLSI inoculum range. Concerningly,

three isolates changed from meropenem susceptible to resistant within the CLSI range (Fig. 4,

dotted lines), including one *E. coli* isolate (BIDMC43b) and two *Serratia marcescens* isolates

(BWH56 and BWH57), each of which encodes a KPC-3*.* Four isolates tested fully susceptible to

meropenem throughout the CLSI inoculum range (red circle), including two *Citrobacter freundii*

harboring KPC-3 (MGH281 and MGH283), and one *E. coli* (BAA2523) and one *K. pneumoniae*

(BAA2524) each harboring an OXA-48. In all, 15 of 63 (24%) carbapenemase producers (CP-

CRE or hyper-CRE) tested meropenem susceptible, and 2 of 41 (4.9%) tested ertapenem

susceptible, at some point in the CLSI inoculum range; this represented 15 of 36 (42%) and 2 of

29 (6.9%) CP-CRE, respectively.

The inoculum effect accurately identifies carbapenemase production among CRE. The

meropenem MIC at standard inoculum poorly distinguished carbapenemase-producing isolates

(CP-CRE or hyper-CRE) from PD-CRE (Fig. 5*A*), making it an inefficient metric to trigger

carbapenemase testing. By contrast, the clear phenotypic differences across inocula between

these groups led us to investigate a diagnostic approach based on the IE. We quantified the IE

by dividing the MIC at high inoculum (1.3×10^7 CFU/mL) by the MIC at standard inoculum ($4 \times$

10^5 CFU/mL) for each isolate (Fig. 5*B*). CRE exhibited distinct meropenem IEs depending on

their mechanism of resistance, with carbapenemase-producing isolates (CP-CRE and hyper-

CRE) exhibiting a much more prominent IE, with a median increase in MIC of 10-fold, compared

to a 1.2-fold increase for PD-CRE (Fig. 5*B*). To quantify binary classification distinguishing

carbapenemase-producing isolates from porin-deficient ones, we generated receiver operating

characteristic (ROC) curves using meropenem MIC (Fig. 5C) or IE (Fig. 5D) threshold values.

The area under each ROC curve (AUC) represents an aggregate measure of classification

accuracy in distinguishing isolates by resistance mechanism across all possible binary

thresholds of the metric being plotted. The meropenem IE distinguished carbapenemase-

producing CRE from PD-CRE better than meropenem MIC at standard inoculum, illustrated by a

ROC AUC of 0.99 for the meropenem IE versus 0.69 for meropenem MIC. To efficiently identify

CRE that might carry carbapenemases, precise IE threshold values can be chosen to optimize

sensitivity (detecting carbapenemases) while maintaining reasonable specificity. At one specific

IE threshold of 2.53-fold, the sensitivity and specificity of distinguishing carbapenemase-

encoding isolates from PD-CRE was 95.2% and 95.7%, respectively. The meropenem IE was

greater in CP-CRE (median 16-fold MIC increase) than hyper-CRE (median 5.3-fold), and it also

distinguished these two groups (Fig. 5 *B* and *D*; ROC AUC 0.89).

Comparing ertapenem and meropenem MICs gives insights into resistance mechanisms.

The meropenem IE effectively distinguished carbapenemase-producing isolates from PD-CRE,

but it is not a standard workflow in clinical laboratories, which typically only perform AST at

standard inocula. To overcome this limitation, since the periplasmic permeability of ertapenem is

typically more impacted by porin mutations than that of meropenem (6, 7, 44, 54), we

hypothesized that comparing ertapenem and meropenem MICs, even at a single inoculum,

might also distinguish CRE by mechanism of resistance. We thus measured ertapenem MICs

across a broad inoculum range for 85 of the 110 CRE isolates (Fig. 6). The ratio of ertapenem

MIC to meropenem MIC better separated carbapenemase producing isolates (CP-CRE or

hyper-CRE) from PD-CRE (Fig. 6*C*) than either the ertapenem or meropenem MIC alone (Fig. 6

A and *B*). At standard inoculum, meropenem MIC performed poorly at distinguishing

carbapenemase producers (CP-CRE or hyper-CRE) from PD-CRE (ROC AUC 0.559), whereas

ertapenem MIC performed slightly better (ROC AUC 0.696; Fig. 6*D*). By contrast, the

ertapenem-to-meropenem MIC ratio yielded a ROC AUC of 0.941, with a 90% sensitivity and

84% specificity at a ratio cutoff of 7.63 for detecting carbapenemase content. At low inoculum,

ertapenem MIC performed comparably to this ratio at identifying carbapenemase producing

isolates, while at high inoculum, meropenem MIC performed comparably. However, at the

standard CLSI inoculum, neither performed as well as the MIC ratio.

Discussion

Our systematic investigations revealed that the *in vitro* carbapenem inoculum effect (IE) is not

determined by the mechanism of action of the antibiotic, but rather by the mechanism of

resistance of the isolate. In *E. coli* K-12, we showed that expression of any carbapenemase is

sufficient to produce a carbapenem IE (Fig. 1). We confirmed this result in a collection of 110

clinical CRE isolates, where all CP-CRE exhibited a strong IE, whereas PD-CRE did not (Fig. 2,

Supplementary Fig. S1, and Table S3). Isolates with both mechanisms of resistance (hyper-

CRE) tended to have higher MICs and also exhibited an IE, although smaller in magnitude than

CP-CRE (Fig. 2*C* and Fig. 5*B*).

Our findings have important mechanistic, clinical, and diagnostic implications.

Mechanistically, all carbapenemases imparted a strong inoculum effect despite minimal

sequence or structural similarity across the four Ambler classes. Furthermore, we observed a

consistently biphasic dependence of CP-CRE MIC on inoculum (Fig. 1, Fig. 2 and

Supplementary Fig. S1): at the lowest inocula tested, carbapenemases conferred a relatively

constant degree of resistance, whereas above a certain threshold, resistance of the population

increased with cell density, by up to 100-fold at the highest cell densities compared with the

standard inoculum (Fig. 1B, Fig. 2A). This is consistent with a model where carbapenemases

provide cell-autonomous resistance at low density but increasing population-level resistance as

cell density increases, such that the collective production of carbapenemases may become the

dominant determinant of resistance at high enough cell density. Since the IE is only observed

with carbapenemase producers and not PD-CRE, one intriguing possibility is that

carbapenemases serve as a "common good" in bacterial populations above a certain density

threshold, as has been proposed for other secreted enzymes (36–38, 55). One recent study of

bacterial communities in which some members produced OXA-48 supports this model (40), but

further studies with mixed populations of carbapenemase producing and non-producing isolates

are needed to rigorously test this hypothesis. Notably, PD-CRE isolates expressing extended

spectrum β-lactamases (ESBLs), which often contribute to carbapenem resistance, did not

exhibit an appreciable carbapenem IE (Fig. 2*B* and Supplementary Fig. S1*B*), indicating that the

slow rates of carbapenem hydrolysis by ESBL enzymes were insufficient to benefit neighboring

bacteria even at high culture density. The lower prevalence of carbapenemases compared to

other β-lactamases may explain why the IE is reported less frequently for carbapenems than for

other β-lactams (56), but this might change as carbapenemases continue to spread globally.

The extent of and mechanistic basis for carbapenemase release into the extracellular space

where they may benefit neighboring cells, whether via secretion systems, outer membrane

vesicles, cell lysis, or other means, warrants more investigation.

The clinical implications of the inoculum effect have been uncertain because the relevant

bacterial cell densities during most infections *in vivo* are generally below those used for AST

(45, 55). For instance, in bacteremia, the infectious burden of ≤10 CFU/mL is several orders of

magnitude below that at which the inoculum begins to affect MIC in our assays (57, 58).

Nevertheless, in *Staphylococcus aureus* bacteremia, a cefazolin IE has been associated with

treatment failure and higher mortality (59, 60). Our findings suggest that at the low inoculum

present in bacteremia, PD-CRE (Fig. 2*B* and Supplementary Fig. S1*B*) exhibit higher

carbapenem MICs than CP-CRE (Fig. 2*A*, Fig. S1*A*), yet clinical outcomes are similar or worse

for CP-CRE bacteremia (61, 62). It is possible that the increased resistance of PD-CRE at

physiologically relevant bacterial densities may be offset by fitness costs associated with

reduced permeability. Further research is needed to fully understand the relationship between

MIC, IE, resistance mechanisms, and clinical outcomes. Intriguing recent studies found that

isolates encoding carbapenemases and displaying a large inoculum effect require higher

meropenem concentration to prevent emergence of highly resistant mutants (39, 52). Since we

observed the largest IE among CP-CRE with intact porins, perhaps these extremely resistant

escape mutants reflect *de novo* porin mutations, creating hyper-CRE.

Diagnostically, accurate recognition of carbapenem resistance is crucial for infection

control strategies and patient care. Our results confirmed prior reports that minor differences in

inoculum can lead to categorical changes in susceptibility interpretation (45), particularly

impacting the detection of CP-CRE, the isolates that carry the greatest implications for infection

control. Even though the isolates we examined were selected because they tested as CRE,

many changed classification within the CLSI-acceptable inoculum range and even appeared

susceptible (Fig. 4), suggesting that clinical algorithms may miss some isolates encoding

carbapenemases. In fact, MICs of all CP-CRE isolates varied significantly in this range. Our

observations in *E. coli* K-12 (Fig. 1) reveal that strains harboring OXA-48, CMY-10, or even KPC

might only test resistant if they possess other factors, such as reduced antibiotic permeability.

The clinical significance of isolates that express a carbapenemase but test susceptible in the

standard inoculum range is uncertain and requires further study. Nonetheless, caution is

generally advised when using carbapenems in infections with carbapenemase-producers, even

if they test susceptible, due to the lack of evidence supporting clinical outcomes in these cases

(22). Our findings in a set of isolates selected for meropenem nonsusceptibility suggest that

other carbapenemase producers may have been missed by standard diagnostic workflows,

which could contribute to poor patient outcomes and the global spread of carbapenem

resistance.

The strong correlation between the carbapenem IE and genotype (Fig. 2) allows for

improved predictions about resistance mechanisms from phenotypic testing. Although MICs

were not designed to identify resistance mechanisms, carbapenem MICs are routinely used to

prioritize isolates for molecular carbapenemase testing (63), though different MIC thresholds for

testing are recommended in the US (22) and Europe (26). Our study revealed that the

meropenem IE is a more sensitive and specific indicator of carbapenemase production in CRE

than MIC at any single inoculum (Fig. 5). Although measuring the meropenem IE would require

additional testing workflows, we also found that integrating ertapenem and meropenem MICs

from routinely generated AST data can inform the likelihood of carbapenemase production (Fig.

6). The ratio of ertapenem to meropenem MIC predicts resistance mechanism better than either

MIC alone, perhaps because the periplasmic permeability of ertapenem is more strongly

affected by porin disruption than that of meropenem (6, 44, 54), while both are hydrolyzed at

similar rates by carbapenemases (64, 65). Predicting porin function has been a major challenge

for genomic resistance prediction (66–71), as polymorphisms are frequently found but

challenging to functionally interpret. Given the differential impact of porin mutations on each

carbapenem, the ertapenem-to-meropenem MIC ratio may offer a phenotypic reflection of porin

function. Specifically, CRE with a high ratio of ertapenem to meropenem MIC (more resistant to

ertapenem) are likely to be PD-CRE, whereas those with a low MIC ratio (similar resistance to

both) are likely to be CP-CRE. As with the inoculum effect, hyper-CRE appear to fall in between,

and tend to have higher MICs at any inoculum. Since the IE impacts both ertapenem and

meropenem MICs similarly (Fig. 4), this ratio predicts resistance mechanism independent of

inoculum, an advantage over either ertapenem or meropenem MICs alone (Fig. 6).

We propose that this ertapenem to meropenem MIC ratio might therefore be better

suited than either MIC alone to inform downstream testing and prioritize isolates for molecular or

enzymatic carbapenemase detection (19, 72, 73). Recognizing that MIC thresholds for

susceptibility may miss carbapenemase-producing isolates, EUCAST recommends a 16-fold

lower MIC threshold for carbapenemase screening than their clinical susceptibility breakpoint

(26). This lower threshold will lead to more sensitive carbapenemase detection, but at lower

specificity, requiring far more testing. If validated in further studies at this lower MIC threshold,

the ertapenem to meropenem MIC ratio may enable optimization of strain selection for

molecular carbapenemase testing. By more efficiently identifying carbapenemase-producing

isolates, our work could facilitate implementation of recent advances showing that molecular

detection of carbapenemases can accelerate administration of targeted therapies, such as β-

lactamase inhibitor combinations (74), and decrease mortality (21). Moreover, this phenotypic

approach could detect strains producing rare carbapenemases, such as SME or CMY-10, that

are not routinely included in testing panels, akin to the Carba NP (22, 75) and mCIM methods

(20). This finding, in addition to its therapeutic value, could serve as a surveillance tool for the

emergence and evolution of carbapenemases (76). This work does not aim to replace well-

validated clinical carbapenemase detection methods. Instead, it illustrates how biological

phenomena can offer further insights into resistance mechanism in a clinical context.

This study has several limitations and areas requiring further study. First, the clinical

isolates analyzed were mostly chosen because they tested carbapenem resistant, limiting the

ability to determine the frequency of CP-CRE isolates that may be missed by routine testing due

to the IE. Further studies are needed to assess the performance of our proposed assays on

isolates with MICs at or below the susceptible breakpoint. Second, automated AST instruments

may not provide precise MIC measurements for ertapenem and meropenem outside a narrow

range around the susceptibility breakpoints, requiring further investigation into the utility of these

lower-resolution MIC estimates in screening for CP-CRE. Third, while we found

carbapenemases sufficient to confer a carbapenem IE, other factors such as heteroresistance,

metabolic factors, quorum sensing, and target distribution may also contribute in some isolates

or circumstances. However, carbapenemases predictably cause a robust IE without requiring

other factors. Our findings reveal a clear pattern: as the inoculum increases above a certain

threshold, we observe a steady linear rise in MIC (Fig. 2 and Supplementary Fig. S1). This

pattern is less consistent with heteroresistance, an alternative explanation for the IE whereby a

rare resistant subpopulation emerges from larger inocula. With heteroresistance, one would

expect a sudden step-like increase in MIC as inoculum rises, as opposed to the gradual MIC

increase that we consistently observed. Lastly, this study did not explore the basis of the IE for

other antibiotics, nor how other processes like antibiotic efflux may contribute to carbapenem

resistance. However, the consistent relationship between resistance mechanism and the IE for

carbapenems, where hydrolytic enzymes and other resistance mechanisms are relatively few

and well characterized, may offer a useful general model for one major mechanism underlying

the IE. Future studies could involve pairing these detailed phenotypic data and genotypic

information (*SI Appendix*, Table S3) with precise enzymological parameters and direct

quantification of enzymes and metabolites in different compartments in order to enhance and

experimentally test existing quantitative models (37, 55). This would enable a more

comprehensive description of how different mechanisms of antibiotic resistance impact the

growth of bacterial populations when exposed to antibiotic selection.

Methods:

Bacterial isolates. The use and collection of bacterial strains from three major Boston-area

hospitals (Beth Israel Deaconess Medical Center, BIDMC; Brigham and Women's Hospital,

BWH; Massachusetts General Hospital, MGH) was approved by the Partners Health Care

Institutional Review Board (IRB under protocol 2015P002215) and additional IRB approval was

granted by the Massachusetts Institute of Technology Committee on the Use of Humans as

Experimental Subjects. Additional isolates were acquired through the University of California

Irvine School of Medicine (UCI), the Wadsworth Center of the New York State Department of

Health (NYSDOH), the FDA-CDC Antimicrobial Resistance Isolate Bank (CDC, Atlanta, GA),

and the American Type Culture Collection (ATCC) (Manassas, VA).

Cloning of carbapenemases. bla_{KPC-3}, bla_{SME-2}, bla_{IMP-4}, bla_{NDM-1}, bla_{VIM-27}, and bla_{OXA-48} along

with their native promoters were cloned by Gibson assembly from isolates in our existing CRE

collection (11, 12) encoding these carbapenemases into a pBAD33 plasmid which encodes

chloramphenicol resistance (*SI Appendix*, Table S1). bla_{CMY-10} with its native promoter was

synthesized in the same pBAD33 vector backbone by GeneWiz (Suzhou, China), since we did

not have a clinical isolate expressing this enzyme.

Broad-range broth microdilution. Our MIC assays were adapted from previously published

broth microdilution methods(24, 77). See Supplemental Methods for details on setup of the

broad inoculum range in this assay format.

Data analysis. MIC replicates were imported to GraphPad Prism v 9.5.1 (GraphPad Software

Inc., San Diego, CA); MIC vs inoculum plots (Figs. 1, 2, and 3, and Supplementary Fig. S1)

depict mean and standard errors of the mean. To analyze the meropenem IE, the ratio of mean

MIC at 1.3×10^7 CFU/mL to 4×10^5 CFU/mL was calculated. The IE of the 47 porin deficient

CRE strains were compared to the 63 carbapenemase-encoding CRE strains using a non-

parametric test (Mann-Whitney for comparisons between two groups or Kruskal-Wallis for

analysis of more than two groups) in GraphPad Prism. Statistical analysis comparing PD-CRE

isolates versus CP-CRE and hyper-CRE isolate at low, standard, and high inocula for

ertapenem, meropenem, and ertapenem-to-meropenem MIC ratio was conducted using Mann-

Whitney tests. All receiver operating characteristic (ROC) curves and analysis were constructed

using GraphPad Prism.

Gene editing to disrupt OmpK36 in *Klebsiella pneumoniae* isolate RB582. A CRISPR-Cas9

cytidine base-editing system(51) was used to introduce an early termination codon in the intact

ompK36 of isolate RB582. See Supplemental Methods for further details.

Acknowledgements

We thank Pierre O. Ankomah, Jonathan Chen, Alyssa DuBois, Alasdair Fletcher, James E.
Gomez, Ria Kolli, Jonathan Livny, Peijun Ma, Melanie A. Martinsen, Michelle E. Matzko, David
J. Roach, and Eleanor L. Young for helpful advice.

We thank Kimberly Musser, the Wadsworth Center, and the CDC ARBank for sharing isolates.

This work was supported in part by the Broad Institute Next Generation Fund (RPB) and federal
funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health,
Department of Health and Human Services, under Grant Number U19AI110818 to the Broad
Institute.

References:

1. M. I. El-Gamal, *et al.*, Recent updates of carbapenem antibiotics. *Eur. J. Med. Chem.* **131**, 185–195 (2017).
2. F. S. Codjoe, E. S. Donkor, Carbapenem Resistance: A Review. *Med. Sci.* **6** (2018).
3. CDC, Antibiotic resistance threats in the United States. *Centers Dis. Control Prev.*, 1–113 (2019).
4. E. S. Snitkin, *et al.*, Tracking a hospital outbreak of carbapenem-resistant Klebsiella pneumoniae with whole-genome sequencing. *Sci. Transl. Med.* **4**, 148ra116 (2012).
5. K. Bush, P. A. Bradford, Epidemiology of β-Lactamase-Producing Pathogens. *Clin. Microbiol. Rev.* **33** (2020).
6. A. Rocker, *et al.*, Global Trends in Proteome Remodeling of the Outer Membrane Modulate Antimicrobial Permeability in Klebsiella pneumoniae. *MBio* **11** (2020).
7. S. Acosta-Gutiérrez, *et al.*, Getting Drugs into Gram-Negative Bacteria: Rational Rules for Permeation through General Porins. *ACS Infect. Dis.* **4**, 1487–1498 (2018).
8. Y. Zhang, *et al.*, Overproduction of efflux pumps caused reduced susceptibility to carbapenem under consecutive imipenem-selected stress in Acinetobacter baumannii. *Infect. Drug Resist.* **11**, 457–467 (2018).
9. H. Pai, *et al.*, Carbapenem Resistance Mechanisms in Pseudomonas aeruginosa Clinical Isolates. *Antimicrob. Agents Chemother.* **45**, 480–484 (2001).
10. M. J. Schwaber, *et al.*, Containment of a Country-wide Outbreak of Carbapenem-Resistant Klebsiella pneumoniae in Israeli Hospitals via a Nationally Implemented Intervention. *Clin. Infect. Dis.* **52**, 848–855 (2011).
11. R. Salamzade, *et al.*, Inter-species geographic signatures for tracing horizontal gene transfer and long-term persistence of carbapenem resistance. *Genome Med.* **14** (2022).
12. G. C. Cerqueira, *et al.*, Multi-institute analysis of carbapenem resistance reveals remarkable diversity, unexplained mechanisms, and limited clonal outbreaks. *Proc. Natl. Acad. Sci. United States gof Am.* **114**, 1135–1140 (2017).
13. A. J. Mathers, *et al.*, Molecular dissection of an outbreak of carbapenem-resistant Enterobacteriaceae reveals intergenus KPC carbapenemase transmission through a promiscuous plasmid. *MBio* **2** (2011).
14. U. Choi, C.-R. R. Lee, Distinct Roles of Outer Membrane Porins in Antibiotic Resistance and Membrane Integrity in Escherichia coli. *Front. Microbiol.* **0**, 953 (2019).
15. A. Alva, A. Sabido-Ramos, A. Escalante, F. Bolívar, New insights into transport capability of sugars and its impact on growth from novel mutants of Escherichia coli. *Appl. Microbiol. Biotechnol.* **104**, 1463–1479 (2020).
16. J. Andersen, S. A. Forst, K. Zhao, M. Inouye, N. Delihas, The function of micF RNA: micF RNA is a major factor in the thermal regulation of OmpF protein in Escherichia coli. *J. Biol. Chem.* **264**, 17961–17970 (1989).
17. J. Abellón-Ruiz, *et al.*, Structural basis for maintenance of bacterial outer membrane lipid asymmetry. *Nat. Microbiol.* **2**, 1616–1623 (2017).
18. K. E. Goodman, P. J. Simner, P. D. Tamma, A. M. Milstone, Infection control implications of heterogeneous resistance mechanisms in carbapenem-resistant Enterobacteriaceae (CRE). *Expert Rev. Anti. Infect. Ther.* **14**, 95–108 (2016).
19. D. Gu, *et al.*, Comparison of the NG-Test Carba 5, Colloidal Gold Immunoassay (CGI) Test, and Xpert Carba-R for the Rapid Detection of Carbapenemases in Carbapenemase-Producing Organisms. *Antibiotics* **12**, 300 (2023).
20. V. M. Pierce, *et al.*, Modified carbapenem inactivation method for phenotypic detection of carbapenemase production among enterobacteriaceae. *J. Clin. Microbiol.* **55**, 2321–2333 (2017).

21. M. J. Satlin, *et al.*, Impact of a Rapid Molecular Test for Klebsiella pneumoniae
Carbapenemase and Ceftazidime-Avibactam Use on Outcomes After Bacteremia Caused
by Carbapenem-Resistant Enterobacterales. *Clin. Infect. Dis.* **75**, 2066–2075 (2022).
22. Clinical and Laboratory Standards Institute, *M100 Performance Standards for*
Antimicrobial Susceptibility Testing (2023) (May 2, 2023).
23. M. P. Weinstein, J. S. Lewis, The clinical and laboratory standards institute subcommittee
on Antimicrobial susceptibility testing: Background, organization, functions, and
processes. *J. Clin. Microbiol.* **58** (2020).
24. Clinical and Laboratory Standards Institute, Methods for Dilution Antimicrobial
Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard—Tenth
Edition. CLSI document M07-A10. Wayne, PA: Clinical and Laboratory Standards
Institute. *Clin. Lab. Stand. Inst.* (2015).
25. R. M. Humphries, *et al.*, Carbapenem-Resistant Enterobacteriaceae Detection Practices
in California: What Are We Missing? *Clin. Infect. Dis.* **66**, 1061–1067 (2018).
26. EUCAST, The European Committee on Antimicrobial Susceptibility Testing. Breakpoint
tables for interpretation of MICs and zone diameters. Version 13.0, 2023.
http://www.eucast.org.".
*http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_5.0_*
Breakpoint_Table_01.pdf (2023).
27. R. M. Humphries, A. N. Abbott, J. A. Hindler, Understanding and Addressing CLSI
Breakpoint Revisions: a Primer for Clinical Laboratories. *J. Clin. Microbiol.* **57** (2019).
28. P. Nordmann, *et al.*, Identification and screening of carbapenemase-producing
Enterobacteriaceae. *Clin. Microbiol. Infect.* **18**, 432–438 (2012).
29. A. Adler, M. Ben-Dalak, I. Chmelnitsky, Y. Carmeli, Effect of Resistance Mechanisms on
the Inoculum Effect of Carbapenem in Klebsiella pneumoniae Isolates with Borderline
Carbapenem Resistance. **59** (2015).
30. I. Brook, Inoculum Effect. *Rev. Infect. Dis.* **11**, 361–368 (1989).
31. A. Gutierrez, *et al.*, Understanding and Sensitizing Density-Dependent Persistence to
Quinolone Antibiotics. *Mol. Cell* **68**, 1147-1154.e3 (2017).
32. G. Diaz-Tang, *et al.*, Growth productivity as a determinant of the inoculum effect for
bactericidal antibiotics. *Sci. Adv.* **8** (2022).
33. K. I. Udekwu, N. Parrish, P. Ankomah, F. Baquero, B. R. Levin, Functional relationship
between bacterial cell density and the efficacy of antibiotics. *J. Antimicrob. Chemother.*
63, 745–757 (2009).
34. O. Scheler, *et al.*, Droplet-based digital antibiotic susceptibility screen reveals single-cell
clonal heteroresistance in an isogenic bacterial population. *Sci. Rep.* **10**, 1–8 (2020).
35. C. Tan, *et al.*, The inoculum effect and band-pass bacterial response to periodic antibiotic
treatment. *Mol. Syst. Biol.* **8** (2012).
36. L. A. Dugatkin, M. Perlin, J. S. Lucas, R. Atlas, Group-beneficial traits, frequency-
dependent selection and genotypic diversity: an antibiotic resistance paradigm. *Proc. R.*
Soc. B Biol. Sci. **272**, 79–83 (2005).
37. N. M. Vega, J. Gore, Collective antibiotic resistance: mechanisms and implications. *Curr.*
Opin. Microbiol. **21**, 28–34 (2014).
38. L. Geyrhofer, *et al.*, Minimal Surviving Inoculum in Collective Antibiotic Resistance. *MBio*
(2023) https://doi.org/10.1128/MBIO.02456-22 (April 24, 2023).
39. M. V. Golikova, *et al.*, Meropenem MICs at Standard and High Inocula and Mutant
Prevention Concentration Inter-Relations: Comparative Study with Non-Carbapenemase-
Producing and OXA-48-, KPC- and NDM-Producing Klebsiella pneumoniae. *Antibiot.*
(Basel, Switzerland) **12** (2023).
40. A. Pathak, D. C. Angst, R. León-Sampedro, A. R. Hall, Antibiotic-degrading resistance
changes bacterial community structure via species-specific responses. *ISME J.* **17**

523    (2023).

524 41. R. H. K. Eng, C. Cherubin, S. M. Smith, F. Buccini, Inoculum effect of β-lactam antibiotics
525    on Enterobacteriaceae. *Antimicrob. Agents Chemother.* **28**, 601–606 (1985).

526 42. Y. Harada, *et al.*, In vitro and in vivo activities of piperacillin-tazobactam and meropenem
527    at different inoculum sizes of ESBL-producing Klebsiella pneumoniae. *Clin. Microbiol.*
528    *Infect.* **20**, O831–O839 (2014).

529 43. D. G. Lee, Y. Murakami, D. R. Andes, W. A. Craig, Inoculum Effects of Ceftobiprole,
530    Daptomycin, Linezolid, and Vancomycin with Staphylococcus aureus and Streptococcus
531    pneumoniae at Inocula of 105 and 107 CFU Injected into Opposite Thighs of Neutropenic
532    Mice. *Antimicrob. Agents Chemother.* **57**, 1434 (2013).

533 44. M. Doumith, M. J. Ellington, D. M. Livermore, N. Woodford, Molecular mechanisms
534    disrupting porin expression in ertapenem-resistant Klebsiella and Enterobacter spp.
535    clinical isolates from the UK. *J. Antimicrob. Chemother.* **63**, 659–667 (2009).

536 45. K. P. Smith, J. E. Kirby, The inoculum effect in the era of multidrug resistance: Minor
537    differences in inoculum have dramatic effect on MIC Determination. *Antimicrob. Agents*
538    *Chemother.* **62**, 1–9 (2018).

539 46. J. D. Lutgring, *et al.*, FDA-CDC Antimicrobial Resistance Isolate Bank: a Publicly
540    Available Resource To Support Research, Development, and Regulatory Requirements.
541    *J. Clin. Microbiol.* **56** (2018).

542 47. R. P. Ambler, "The structure of β-lactamases" (1980).

543 48. T. Sawa, K. Kooguchi, K. Moriyama, Molecular diversity of extended-spectrum β-
544    lactamases and carbapenemases, and antimicrobial resistance (J Intensive Care, 2020).

545 49. L. M. Guzman, D. Belin, M. J. Carson, J. Beckwith, Tight regulation, modulation, and
546    high-level expression by vectors containing the arabinose P(BAD) promoter. *J. Bacteriol.*
547    **177**, 4121–4130 (1995).

548 50. J. Y. Kim, *et al.*, Structural basis for the extended substrate spectrum of CMY-10, a
549    plasmid-encoded class C β-lactamase. *Mol. Microbiol.* **60**, 907–916 (2006).

550 51. Y. Wang, *et al.*, CRISPRCas9 and CRISPR-assisted cytidine deaminase enable precise
551    and efficient genome editing in Klebsiella pneumoniae. *Appl. Environ. Microbiol.* **84**
552    (2018).

553 52. P. Ma, *et al.*, Genetic determinants facilitating the evolution of resistance to carbapenem
554    antibiotics. *Elife* **10** (2021).

555 53. M. W. Adelman, *et al.*, Distinctive Features of Ertapenem-Mono-Resistant Carbapenem-
556    Resistant Enterobacterales in the United States: A Cohort Study. *Open Forum Infect. Dis.*
557    **9** (2022).

558 54. Y. K. Tsai, C. H. Liou, C. P. Fung, J. C. Lin, L. K. Siu, Single or in Combination
559    Antimicrobial Resistance Mechanisms of Klebsiella pneumoniae Contribute to Varied
560    Susceptibility to Different Carbapenems. *PLoS One* **8**, e79640 (2013).

561 55. T. Artemova, Y. Gerardin, C. Dudley, N. M. Vega, J. Gore, Isolated cell behavior drives
562    the evolution of antibiotic resistance. *Mol. Syst. Biol.* **11**, 822 (2015).

563 56. J. R. Lenhard, Z. P. Bulman, Inoculum effect of β-lactam antibiotics. *J. Antimicrob.*
564    *Chemother.* **74**, 2825–2843 (2019).

565 57. O. Opota, A. Croxatto, G. Prod'hom, G. Greub, Blood culture-based diagnosis of
566    bacteraemia: state of the art. *Clin. Microbiol. Infect.* **21**, 313–322 (2015).

567 58. C. Zhang, *et al.*, Identification of low abundance microbiome in clinical samples using
568    whole genome sequencing. *Genome Biol.* **16** (2015).

569 59. W. R. Miller, *et al.*, The Cefazolin Inoculum Effect Is Associated With Increased Mortality
570    in Methicillin-Susceptible Staphylococcus aureus Bacteremia. *Open forum Infect. Dis.* **5**
571    (2018).

572 60. E. C. Nannini, *et al.*, Inoculum effect with cefazolin among clinical isolates of methicillin-
573    susceptible Staphylococcus aureus: frequency and possible cause of cefazolin treatment

failure. *Antimicrob. Agents Chemother.* **53**, 3437–3441 (2009).

61.   P. D. Tamma, *et al.*, Comparing the outcomes of patients with carbapenemase-producing
and non-carbapenemase- producing carbapenem-resistant enterobacteriaceae
bacteremia. *Clin. Infect. Dis.* **64**, 257–264 (2017).

62.   M. R. Hovan, N. Narayanan, V. Cedarbaum, T. Bhowmick, T. J. Kirn, Comparing mortality
in patients with carbapenemase-producing carbapenem resistant Enterobacterales and
non-carbapenemase-producing carbapenem resistant Enterobacterales bacteremia.
Diagn. Microbiol. Infect. Dis. **101** (2021).

63.   S. S. Jean, W. Sen Lee, P. R. Hsueh, Ertapenem non-susceptibility and independent
predictors of the carbapenemase production among the Enterobacteriaceae isolates
causing intra-abdominal infections in the Asia-Pacific region: results from the Study for
Monitoring Antimicrobial Resistance Trends (SMART). *Infect. Drug Resist.* **11**, 1881
(2018).

64.   N. K. Stewart, C. A. Smith, H. Frase, D. J. Black, S. B. Vakulenko, Kinetic and Structural
Requirements for CarbapenemaseActivity in GES-Type β-Lactamases. *Biochemistry* **54**,
588 (2015).

65.   T. L. Lamoureaux, H. Frase, N. T. Antunes, S. B. Vakulenko, Antibiotic Resistance and
Substrate Profiles of the Class A Carbapenemase KPC-6. *Antimicrob. Agents*
Chemother. **56**, 6006–6008 (2012).

66.   M. W. Pesesky, *et al.*, Evaluation of Machine Learning and Rules-Based Approaches for
Predicting Antimicrobial Resistance Profiles in Gram-negative Bacilli from Whole Genome
Sequence Data. *Front. Microbiol.* **7** (2016).

67.   B. B. Flury, *et al.*, Association of Novel Nonsynonymous Single Nucleotide
Polymorphisms in ampD with Cephalosporin Resistance and Phylogenetic Variations in
ampC, ampR, ompF, and ompC in Enterobacter cloacae Isolates That Are Highly
Resistant to Carbapenems. *Antimicrob. Agents Chemother.* **60**, 2383–2390 (2016).

68.   S. A. Shelburne, *et al.*, Whole-Genome Sequencing Accurately Identifies Resistance to
Extended-Spectrum β-Lactams for Major Gram-Negative Bacterial Pathogens. *Clin.*
Infect. Dis. **65**, 738–745 (2017).

69.   D. E. Foudraine, *et al.*, Exploring antimicrobial resistance to beta-lactams,
aminoglycosides and fluoroquinolones in E. coli and K. pneumoniae using
proteogenomics. *Sci. Rep.* **11** (2021).

70.   C. S. Suelter, N. D. Hanson, OmpC regulation differs between ST131 and non-ST131
Escherichia coli clinical isolates and involves differential expression of the small RNA
MicC. *J. Antimicrob. Chemother.* **75**, 1151–1158 (2020).

71.   J. L. C. Wong, *et al.*, Recurrent emergence of Klebsiella pneumoniae carbapenem
resistance mediated by an inhibitory ompK36 mRNA secondary structure. *Proc. Natl.*
Acad. Sci. U. S. A. **119**, e2203593119 (2022).

72.   J. Monteiro, R. H. Widen, A. C. C. Pignatari, C. Kubasek, S. Silbert, Rapid detection of
carbapenemase genes by multiplex real-time PCR. *J. Antimicrob. Chemother.* **67**, 906–
909 (2012).

73.   P. D. Tamma, P. J. Simner, Phenotypic detection of carbapenemase-producing
organisms from clinical isolates. *J. Clin. Microbiol.* **56** (2018).

74.   J. C. Vázquez-Ucha, J. Arca-Suárez, G. Bou, A. Beceiro, New Carbapenemase
Inhibitors: Clearing the Way for the β-Lactams. *Int. J. Mol. Sci.* **21**, 9308 (2020).

75.   B. Elawady, M. Ghobashy, A. Balbaa, Rapidec Carba NP for Detection of
Carbapenemase-Producing Enterobacteriaceae in Clinical Isolates: A Cross-Sectional
Study. *Surg. Infect. (Larchmt).* **20**, 672–676 (2019).

76.   R. C. MacLean, A. S. Millan, The evolution of antibiotic resistance. *Science (80-.).* **365**,
1082–1083 (2019).

77.   I. Wiegand, K. Hilpert, R. E. W. Hancock, Agar and broth dilution methods to determine

the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat. Protoc.* **3**,
163–175 (2008).
627

Fig. 1. All carbapenemases impart a meropenem inoculum effect when expressed in *E. coli K12*. (A) Plots of optical density (OD) at 600 nm (y-axis) after overnight incubation with varying meropenem concentrations (x-axis) across a broad range of starting inocula (grayscale per top right panel) for *E. coli* K-12 transformed with each indicated carbapenemase. **(B)** Meropenem minimum inhibitory concentration (MIC) as a function of bacterial inoculum, colored by transformed carbapenemase. Each MIC point is the mean of three replicates (error bars = standard error of the mean). Vertical dotted lines represent the CLSI-acceptable inoculum range (2 to 8 x 10^5 CFU/mL). The background is shaded by CLSI meropenem susceptibility breakpoints (S = susceptible, white; I = intermediate, light gray; R = resistant, gray).

Fig. 2. Select CRE exhibit different patterns of carbapenem inoculum effect depending on resistance mechanism. Meropenem MICs from broth microdilution assays (y-axis) across a broad range of starting inocula (x-axis) of select clinical CRE isolates encoding **(A)** a carbapenemase of each major type, **(B)** porin deficiency, or **(C)** both, with species and relevant genotype as indicated. One susceptible isolate (MGH19, gray) is also shown in panel **(A)**. Each MIC point is the mean of three replicates (error bars = standard error of the mean). *C spp = Citrobacter* spp, *Ecl = Enterobacter cloacae*, *Eco = Escherichia coli*, *Kae = Klebsiella aerogenes*, *Kpn = Klebsiella pneumoniae*, *Kqu = Klebsiella quaispneumoniae*, *Sma = Serratia marcescens*. CPase = carbapenemase. Vertical dotted lines reflect the CLSI-recommended inoculum range, background shading reflects CLSI breakpoints, and line color reflects carbapenemase content, all as in Fig. 1; dashed lines indicate porin deficiency.

Fig. 3. Carbapenemase production and porin deficiency together create a hyper-CRE phenotype. Broad-range meropenem broth microdilution assays for **(A)** two clinical PD-CRE isolates before and after transformation with pBAD33_KPC-3 and **(B)** a clinical CP-CRE isolate and three derivative colonies in which OmpK36 was disrupted by CRISPR-Cas9 based gene editing to create a premature stop at one of two glutamines as indicated. The parental KPC-expressing isolate (RB582) is shown as a solid line with green markers. Each MIC point is the mean of three replicates (error bars = standard error of the mean). Vertical dotted lines reflect the CLSI-recommended inoculum range, background shading reflects CLSI breakpoints, and line color reflects carbapenemase content as in Fig. 1; dashed lines indicate porin deficiency as in Fig. 2. *Kpn = Klebsiella pneumoniae.*

Fig 4. Many carbapenemase-producing isolates change AST classification in the CLSI range. Broad-range broth microdilution assays for **(A)** 63 carbapenemase-producing clinical CRE isolates (36 CP-CRE and 27 hyper-CRE) treated with meropenem and **(B)** 41 carbapenemase-producing clinical CRE isolates (30 CP-CRE and 11 hyper-CRE) treated with ertapenem. Each MIC measurement was the mean of three replicates. Error bars were omitted for simplicity. Isolates that tested anything other than resistant at some point in the CLSI-recommended inoculum range are colored by carbapenemase content; those that tested resistant throughout the CLSI range are in gray. Inset table shows the number and percentage of isolates by AST class across the CLSI inoculum range; red text indicates isolates that tested susceptible at some point within the CLSI range. Four CP-CRE that tested fully meropenem-susceptible throughout the CLSI range are indicated with a red circle. Isolates whose meropenem susceptibility category changed from susceptible to resistant in the CLSI range are shown as dotted lines.

Fig. 5. Meropenem inoculum effect robustly classifies CRE isolates by resistance mechanism. (A) Meropenem MICs for 110 CRE grouped by genotypic mechanism of carbapenem resistance: CP-CRE (circles), PD-CRE (triangles), or hyper-CRE (squares). Isolates are colored by carbapenemase content as in Fig. 1. Each MIC point is the mean of three replicates. **(B)** Ratio of meropenem MIC at high (1.3 x 10^7 CFU/mL) versus standard (4.0 x 10^5 CFU/mL) inoculum for 110 CRE, grouped as in **(A).** P-values were obtained by conducting statistical analyses on the three groups in **(A)** and **(B)** using a Kruskal-Wallis test. **(C)** Receiver operating characteristic (ROC) curves based on the meropenem MICs at standard inoculum from **(A)** comparing the different genotypic CRE groups. Inset tabulates each ROC area under the curve (AUC), sensitivity (Se), and specificity (Sp) at an optimal cutoff (OC) threshold, indicated by the red points on the ROC curves. **(D)** ROC curves based on the meropenem inoculum effect, displayed as in **(C)**. Any CPase = either CP-CRE or hyper-CRE.

Fig. 6. Comparing ertapenem and meropenem MICs gives insight into resistance mechanism (A-C)
MICs of ertapenem **(A)** or meropenem **(B)**, or the ratio of the ertapenem MIC to the meropenem MIC **(C),** are
shown for 85 isolates at low, standard, and high inoculum (left, middle, and right panels respectively), grouped

by resistance mechanism along the x-axis (hyper-CRE = "Hy-CRE"). Data points are colored based on carbapenemase content as in Fig. 1. Plot backgrounds are shaded by CLSI breakpoints for ertapenem **(A)** or meropenem **(B)** as in Fig. 1. P-values were obtained by conducting statistical analyses on the three groups in using a Kruskal-Wallis test. In chemical structures, the carbapenem core is red and the modifiable substituents are black. **(D)** ROC curves comparing carbapenemase producers (CP-CRE + hyper-CRE) vs PD-CRE for each metric (red = ertapenem MIC, blue = meropenem MIC, purple = MIC ratio) at low, standard, and high inoculum. Inset tabulates ROC AUC and sensitivity (Se) and specificity (Sp) at an optimal cutoff (OC, indicated by arrow on ROC curve).