Wiring between close nodes in biological networks evolves more quickly than between distant nodes

Alejandro Gil-Gomez, alejandro.gilgomez@stonybrook.edu

Joshua S. Rest*, joshua.rest@stonybrook.edu, 631-632-1916

Department of Ecology and Evolution

Stony Brook University

650 Life Sciences

Stony Brook, NY 11794-4254

*Corresponding author

Abstract

Protein-protein interaction networks and metabolic networks change across species as a result of a wide range of evolutionary processes; these processes result in the accumulation of network differences between species. The rate at which biological networks evolve is an important question in evolutionary biology. Previous empirical work has focused on interactomes from model organisms to calculate rewiring rates, but this is limited by the relatively small number of species and incomplete nature of network data across species. We present an alternative approach to study network evolution using drug-drug interactions (DDIs) as a proxy for network topology. Here, we propose the rate at which DDIs change across species as an estimate of the rate at which the underlying biological network changes as species diverge. We computed the evolutionary rates of DDIs using previously published data on a high throughput study in gram-negative bacteria. In parallel, we mapped drugs with known targets in biological networks and calculated network distance, connectivity, and adjacency between targets. Our findings indicate that the targets of synergistic DDIs are closer in biological networks than other types of DDIs and that synergistic interactions have a higher evolutionary rate, meaning that nodes that are closer in biological networks evolve at a faster rate. Future studies of network evolution may use DDI data and phylogenetic comparative methods to gain larger-scale perspectives on the details of network evolution within and between species.

Introduction

Biological networks are representations of molecular interactions in the cell. These networks are built by collecting biochemical and genetic interactions from different kinds of biological entities, an approach that has improved drastically in the last decades with the advent of modern high-

throughput methods. However, there are still many limitations to the gathering and analysis of biological networks, specifically for studying network evolution across and within species. This is because many biological networks have poor quality or are incomplete and because the number of organisms for which network data is available is still very limited (Cusick et al. 2005; Jin et al. 2013; Ghadie et al. 2018).

Biological networks evolve as nodes and edges are added or lost in the network. This can be the result of different processes affecting network content, such as gene duplication, gene loss, pseudogenization, horizontal gene transfer, or whole genome duplication (Wagner 2003; Cork and Purugganan 2004), or a result of processes affecting quantitative properties, including non-synonymous substitutions in the nodes affecting their function (Jensen 1976; Ghadie et al. 2018). These mutations may affect the binding of ligands, protein domains, or DNA motifs, which in turn may result in changes in their associated metabolic, signaling, or gene expression networks. The type of network also may affect the rewiring rates of nodes. For example, gene regulatory networks tend to rewire at faster rates than metabolic networks (Shou et al. 2011), this suggests that some network types are less constrained to network evolution as a whole, but the rate at which whole networks change and what factors determine this process remain to be studied in the future.

Moreover, evolutionary events that remove old connections and generate new ones may be random (i.e., neutral or occurring via genetic drift) or may affect fitness. We know from studying static networks that not all motifs are equally abundant in biological networks (Picard et al. 2008), suggesting the action of natural selection. Natural selection may remove deleterious connections and favor advantageous ones. Rewiring rates also vary within networks, such as between different network modules and between nodes depending on their importance for the cell's survival and reproduction, and the essentiality of a node or module may be a constraining factor for network evolution. Previous studies have also shown an inverse relationship between highly connected proteins and their rate of substitution (Fraser et al. 2002), possibly due to strong purifying selection acting on the interfacial sites of interacting proteins (Zotenko et al. 2008).

Part of the key to understanding the forces acting on networks is a characterization of evolutionary rates of network change. Despite advances in studying the connections of individual nodes and network rewiring, the rate and patterns by which quantitative differences between networks accumulate as a function of species divergence remain relatively unexplored. If non-neutral forces act, evolution in networks should be correlated with metrics of inter-node connectivity, including minimum path length (the number of steps along the shortest path between nodes), K-edge connectivity (minimum number of edges that can be removed to disconnect two nodes), and degree (number of edges connected to the node) (Jordan 1869; Erdős and Gallai 1960; Cormen 2001). However, it is unknown whether evolutionary rates of change in inter-node connectivity are associated with these inter-node network measures. This is important for understanding what aspects of network structure that we can quantitate are subject to evolutionary forces acting on variation. This allows us to ask a broader question: which aspects of network structure that we can measure are important to organisms?

In this paper, we use drug-drug interactions (DDIs) as a quantitative proxy measurement of inter-node connectivity, since it has been shown that DDIs are partially dependent on the underlying network topology between targets (Lehár et al. 2007; Yeh et al. 2009). DDIs occur when the effect of two or more drugs is significantly stronger or weaker than their expected combined effect, respectively named synergies and antagonisms (Greco et al. 1995). DDIs are used in the

development of novel pharmacological treatments with higher efficiencies at lower doses and to reduce the evolution of drug resistance (Cowen and Steinbach 2008). The reasons why a DDI occurs are varied, but it has been shown that a DDI can occur as a result of factors such as the topology of the underlying network between drug targets, or the essentiality of the metabolites blocked by the perturbation in combination with the inhibition efficiency of the drug on the drug target (Yeh et al. 2009). For example, synergies can occur if drugs act on parallel pathways, where the individual effect of each drug is small and an alternative pathway can compensate for the effect of a single drug. Alternatively, antagonistic interactions can occur mainly in two different ways: by causing a partial loss of function in two parallel pathways of an essential product, or as a result of drugs acting sequentially along the same pathway. This relation between DDIs and network topology has been further explored using experiments and metabolic flux simulations in yeast, suggesting that specific combinations of perturbations in the network result in different quantifiable interaction scores (Lehár et al. 2007). More recently, it has been shown that most synergistic interactions are the result of drugs targeting the same cellular process, while antagonistic interactions are the result of drugs targeting different processes (Brochado et al. 2018).

Only a few studies have explored interspecific variation of DDIs (Spitzer et al. 2011; Robbins et al. 2015; Brochado et al. 2018), but they have demonstrated that DDI measurement could be scalable in the number of species and strains, allowing for the evaluation of network evolution hypotheses within and between species. A high throughput study of DDIs in gram-negative bacteria has shown that synergies are more conserved across species than antagonisms and additive interactions (Brochado et al. 2018).

These previous findings suggest several important questions about the evolutionary patterns of biological networks and DDIs across species. Do different parts of biological networks, which in turn are reflected in different types of DDIs, evolve at different rates as species diverge? Do DDIs diverge consistently with neutral processes acting on the underlying network structure? Or is there heterogeneity across the network, for example with differences accumulating slowly in constrained local neighborhoods, and faster between more distant connections? Does the evolutionary rate of DDIs or biological networks depend on evolutionary forces acting on the drug targets and their connectivity?

Another way to parse this question is to divide up interactions by type. Do synergistic interactions evolve at a slower rate than other types of drug interactions, and do antagonistic interactions evolve at a faster rate? These differences in rates would be a result of synergistic interactions taking place in local neighborhoods of nodes, while antagonistic interactions act across distant network neighborhoods, which may evolve more slowly due to greater network redundancy between distant nodes. Indeed, if we were to map DDIs to protein targets (when known), are the drug targets of synergistic drug interactions closer in the network than additive and antagonistic interactions? And, more generally, are more closely connected nodes subject to higher rates of evolution than more distantly related nodes? These questions can be addressed by comparing rates of DDI evolution with measures of inter-node connectivity.

To address these questions, we used the most complete available dataset of DDIs measured across species and strains (Brochado et al. 2018). We modeled these DDIs under a phylogenetic comparative framework and applied a multivariate Brownian motion model to estimate the evolutionary rate of interaction scores for different clusters of DDIs in six strains (three species) of gram-negative bacteria. We also mapped DDIs to their putative protein targets to evaluate them in

known biological networks. We show that DDIs can be used as an effective proxy to evaluate macroevolutionary implications and the role of network evolution in interspecific variation.

Results

DDI divergence between the strains increases as a function of species divergence.

We constructed a phylogenetic tree based on highly conserved proteins (**Fig. 1A**), which highlights the close relationship between the three species (including two strains from each species) considered in this study, *E. coli*, *S. enterica*, and *P. aeruginosa*. Other species of Bacteria and Archaea were used to calibrate the nodes for phylogenetic dating and to calibrate the rate of DDI evolution. We estimated the MRCA of the three gammaproteobacteria species in our study at 1049-1128 MYA (95% HPD) and the MRCA of the two *Enterobacteriaceae* (*E. coli* and *S. enterica*) at 57-68.15 MYA (95% HPD).

We obtained DDI scores (drug-drug interaction Bliss scores) from a previous study (Brochado et al. 2018) that assessed almost 3000 combinations of 79 different compounds on six strains, two from each of the tree species. Clustering of the six strains based on their DDI scores revealed a 'chemical distance' tree consistent with the phylogeny ("Strains" in **Fig. 1B**). To test whether DDI scores ("chemical patristic distance") diverge as a function of species divergence, we used linear and logarithmic regressions, which modeled the amount of DDI variation explained by sequence divergence (Manhattan distance on the phylogeny).

We sought to estimate the evolutionary rate of DDI change. However, there is low power with only six tips to estimate rate shifts for any individual DDI. To address this limitation, we used tSNE to detect clusters of DDIs that behave similarly across species, allowing for increased power within each cluster. In order to find DDI clusters that are the most biologically relevant from among a large parameter space (after filtering solutions), we first filtered for the top five solutions with the highest degree of phylogenetic modular signal (Adams et al. 2016). From among those five, we chose the solution with the biggest differences in evolutionary rate among clusters, when fitted with a multivariate Brownian motion model. A detailed description of the filters and tests used to select the clusters is in the Methods.

The resulting solution, with a perplexity of 245 and 15 clusters, was used for the remainder of the analysis. Cluster 14 had the highest evolutionary rate measured using the Brownian motion standard deviation parameter σ^2, with a value of 0.00784 bliss score^2 per million years, while all the other clusters had rates ranging from 0.00044 to 0.00367. Cluster 15 had the second highest rate with a value of 0.00367 bliss score^2 per million years, and cluster 1 had the third highest rate with a value of 0.00354 bliss score^2 per million years.

We examined drug-drug interaction (DDI) variation among these six strains by building a phylomorphospace plot using PCA and found that *Pseudomonas* and non-*Pseudomonas* strains differentiate well along PC1, which explains 47.01% of the variance (**Fig. 1C**). PC2 distinguishes between *E. coli* and *Salmonella* strains and explains 21.54% of the variance. Strains of the same

species group closely together in the PCA space, indicating that they share similar global DDI responses. For example, the *Pseudomonas* strains appear to be closer to each other in the phylomorphospace suggesting that they have small DDI variation, alternatively, *E. coli* and *S. typhimurium* have greater intraspecific DDI variation.

We also correlated the branch length of the dendrogram obtained from clustering DDI data with the branch lengths obtained from the Bayesian phylogeny. We found thatchemical distance increases as a function of phylogenetic distance (linear regression: R^2=0.634, p-value=0.0003; **Fig. 1D**). This relationship is even stronger when phylogenetic distance is log transformed (R^2 =0.918, p-value=$2.4*10^{-8}$; **Fig. 1E**). These results suggest that DDIs, and their underlying network basis, diverge as a function of sequence divergence. DDI divergence plateaus at higher phylogenetic divergence values; this type of saturation has been previously reported in models of network evolution for rewiring rates across species (Shou et al. 2011). Our observation that DDIs diverge as a function of species divergence is consistent with neutral processes acting on the underlying network structure, although selective processes certainly also play a role, for example, clusters with high evolutionary rates may be driven by positive selection.

Synergistic interactions have the shortest average distance between targets, followed by additive and antagonistic interactions.

We observed that highly synergistic DDIs across all species tend to occur when both drugs belong to the same chemical category and target the same cellular process, as can be seen from the accumulation of green bars for category and process across cluster 14 (**Fig. 1B**; in agreement with (Brochado et al. 2018)). This motivated us to formally examine how DDIs map onto biological networks. We therefore leave discussion of DDI rates for now, to first describe our mapping of the DDI targets to known biological networks, and how connectivity on these networks relates to interaction type (antagonism, additivity, synergy); in the next section we will describe how DDI evolutionary rates map onto interaction types network connectivity. We were able to identify protein targets for 39 of the 79 drugs tested by Brochado *et al.* (2018) (see Methods). These drugs had a total of 27 target proteins as identified by their unique protein IDs in *E. coli*. The most common target category was bacterial penicillin-binding protein, a group involved in the biosynthesis of bacterial cell walls. Other target categories mapped include ribosomal RNA, DNA polymerase, topoisomerase, thymidylate synthase, and mitochondrial glycerol-3-phosphate (**Sup. Table S1**).

Figure 1: Drug-drug interaction (DDI) chemical distance scales with phylogenetic distance. **A.** Bayesian
phylogeny of an alignment of 35 highly conserved proteins in different species of bacteria and Archaea
used for phylogenetic dating. Gammaproteobacteria (red), cyanobacteria (green), bacillota (yellow), and
Archaea (purple). **B.** Heatmap of DDI scores across strains. Hierarchical clustering based on euclidean
distances across columns (strains: *E. coli* K-12 BW2952 (ebw), *Escherichia coli* O8 IAI1 (ecr), *S. enterica
subsp. enterica* serovar Typhimurium 14028S (seo), *Salmonella enterica* subsp. enterica serovar
Typhimurium LT2 (stm), *Pseudomonas aeruginosa* PAO1 (pae), *P. aeruginosa* UCBPP-PA14 (pau)), and
rows (DDIs); the latter was constrained by tSNE clusters (see text). Synergistic interactions are shown in
blue and antagonistic interactions are in red, all DDIs are measured in bliss score units. The annotations
on the right indicate whether the two drugs involved in the interaction belong to the same drug category,
target the same cellular process, or have the same use (in green) or not (black). The bottom row indicates
the evolutionary rate calculated in each cluster, measured in bliss score units^2 per million years. **C.**
Phylomorphospace plot combining a PCA of the DDI data with the phylogeny; principal components 1 and
2 are shown. **D, E.** DDI-based Euclidean distances between strains scale as a function of phylogenetic
distance. This relationship saturates at large phylogenetic distances. Intraspecific comparisons (in blue)
have the lowest amount of sequence and chemical divergence, while comparisons with *Pseudomonas* are
the most distant in terms of sequence and chemical divergence. **E.** Same comparison is shown when
phylogenetic distance is plotted on a log scale. All drug-drug interaction data is from Brochado *et al.*
(2018).

Figure 2: **A.** Graphical representations of metabolic interactions via the co-functional gene pair network of *E. coli* (EcoliNet: EcoCyc/GO-BP). This network contains 1835 nodes with an average path length of 4.8 and contains 20 proteins targeted by 36 drugs in our analysis. **B.** Graphical representations of the PPI network of *E. coli*, as determined by small and medium-scale experiments (EcoliNet: LC. Small/medium-scale PPI). This network contains 764 nodes with an average path length of 4.9, with 18 proteins targeted by 27 drugs in our analysis. In **A** and **B**, each node represents a unique protein in *E. coli* (KEGG ID); the red nodes are target proteins identified as participating in DDIs in our analysis.

We then examined the *E. coli* DDI scores as a function of the connectivity of their targets. We used a co-functional gene pair network of *E. coli* derived from EcoCyc and GO-BP data, which contains genes, proteins, pathways, and molecular interactions in *E.coli* and Gene Ontology Biological Processes, respectively (Ashburner et al. 2000; Karp et al. 2018) (**Fig. 2A**). We also used a small/medium scale protein-protein interaction (PPI) network, also from *E. coli* (Kim et al. 2015) (**Fig. 2B**). We measured the length of minimum distance paths between all DDI targets in the networks and found that synergistic interactions have the shortest average distance, followed by additive interactions, and antagonisms. Synergistic interactions have a significantly shorter average path length between protein targets than antagonisms and additive interactions in both networks (**Fig. 3A, B**). This result is consistent with our expectations, given that synergistic interactions tend to be more common between drugs that target the same cellular process and thus should be closer to each other in biological networks (Brochado et al. 2018). We next examined K-edge connectivity, where two nodes are K-edge-connected if after removing k edges or less the nodes remain connected. For the co-functional network, we found that K-edge connectivity was higher in synergies than additivities (Wilcoxon p-value=0.013), and higher in additivities than antagonisms (Wilcoxon p-value=0.045) (**Fig. 3C).** This result was not significant for the PPI network (**Fig. 3D**). We found that node degree (the number of connected edges) was significantly higher in synergies than additivities (Wilcoxon p-value=0.0025) or antagonistic interactions (Wilcoxon p-value=0.00018) in the co-functional network (**Fig. 3E**), but not in the PPI network (**Fig. 3F**). We also found that average betweenness centrality was higher for synergies than for additivities (Wilcoxon p-value=0.0015) and antagonisms (Wilcoxon p-value=0.013) in the co-functional network, but not in the PPI network.

248 Average eigenvector centrality was also higher in synergies than additivities (Wilcoxon p-
249 value=0.0002) and antagonisms (Wilcoxon p-value=0.0005) in the co-functional network, but these
250 differences were not significant in the PPI network.
251

Synergistic interactions evolve faster than additive and antagonistic interactions.

Cluster 14, which had the highest evolutionary rate (σ^2=0.00784), appears to be rich in synergies in
E. coli and *Salmonella,* while *Pseudomonas* has more additive DDIs in this cluster (**Fig. 1B**). In
contrast, cluster 15, which was the second highest rate cluster (σ^2=0.00367), appears to be rich in
synergies in *Pseudomonas* while the DDIs of *E. coli* and *Salmonella* are primarily additive. The
third cluster with the highest rate, cluster 1 (σ^2=0.00354), contains highly antagonistic DDIs in all
the species. These observations suggest that the evolutionary rate of DDIs may vary as a function of
the biological relatedness (i.e. interaction type and network connectivity) of their targets. To more
formally investigate this, we used the evolutionary rates determined for DDIs on a per-cluster basis
(via tSNE, described above) and examined how the rates vary based on the targets' connectivity (in
E. coli) and type of interaction. None of the DDIs were exclusively antagonistic across all strains and
species, although we found all other combinations (i.e., additivity, synergy, additivity↔antagonism,
additivity↔synergy, antagonism↔synergy and additivity↔antagonism↔synergy, where the arrow
indicates that some strains or species have one interaction type and other strains or species have
another interaction type).

Overall, synergistic and additive↔synergistic interactions have faster evolutionary rates than any
other interaction combination (**Fig. 4A**; all possible group comparisons were significantly different
from each other with p-values<0.001 except additivity↔antagonism↔synergy vs.
additivity↔antagonism, and excluding antagonism↔synergy, with only a single observation; **Sup.**
Table S2). We also scored each target pair's interaction type as a simple sum of its interaction types
across strains and species (+1 for synergies, -1 for antagonisms), and found again that more
synergistic interactions (sum ≤ -3) have faster evolutionary rates than additivities (3 > sum > -3) (p-
value<0.001) and antagonisms (sum ≥ 3) (p-value<0.001) (**Fig. 4B**). This result of higher rates in
synergistic interactions is consistent with the result for interaction types from a single strain, *E. coli*
K-12 EBW2952 (**Fig. 4C**; all categories are different at p<0.001, except for additivities vs
antagonisms in the PPI network**)**.

Figure 3: Differences in minimum path length, k edge connectivity, mean node degree, mean betweenness centrality, and mean eigenvector centrality between targets for different types of drug interactions in *E. coli* for co-functional and PPI networks. **A-B**. Synergistic interactions have a lower average path length than additive interactions and antagonistic interactions. **C-D.** K-edge connectivity is higher in synergies than in additive and antagonistic interactions in the co-functional network, but not in the PPI network. **E-F.** The mean node degree is higher in synergies than in additive or antagonistic interactions in the co-functional network, but not in the PPI network. **G-H**. Mean betweenness centrality is higher in synergies than in additive or antagonistic interactions in the co-functional network, but not in the PPI network. **I-J**. Mean eigenvector centrality is higher in synergies than in additive or antagonistic interactions in the co-functional network, but not in the PPI network.

In concert with this, DDIs whose targets are nearby in the network have the highest evolutionary rates (this is consistent with the fact that cluster 14 has the highest rate, and contains a majority of synergies; **Fig. 4D**). This result is significant for both the PPI network (p-value=0.008) and the co-functional (EcoCyc/GO-BP) network (p-value=0.022) (**Sup. Table S2**). We interpret this to mean that wiring between close nodes in biological networks evolves faster than between distant nodes. The relationship between evolutionary rates and other network connectivity measures such as K-edge connectivity appears to be less obvious: evolutionary rates increase sharply as a function of k edge connectivity in the co-functional (EcoCyc/GO-BP) network when connectivity is greater than 20, however, this pattern doesn't hold for the PPI network, where all the protein targets have connectivities lower than 5 (**Sup. Fig. S1K**). Evolutionary rates drop sharply and increase for node degrees greater than 20 in the co-functional (EcoCyc/GO-BP) network, but this result is not seen in the PPI network where the maximum node degree is 10 (**Sup. Fig. S1L**). Rates do not increase or decrease as a function of betweenness centrality (**Sup. Fig. S1M**), however, we see a sharp decrease in rates for interactions with higher average eigenvector centralities in the PPI network.

We also evaluated adjacency (whether two nodes are directly linked) and connectedness (whether two nodes are connected through a path). We found that most target interactions are not adjacent, however, most adjacent nodes have higher evolutionary rates than non-adjacent nodes, both in the co-functional (EcoCyc) (Wilcoxon p-value=0.00025; **Sup. Fig. S1A**) and PPI networks (Wilcoxon p-value=0.0047; **Sup. Fig. S1B**; note that this result was not significant in the EcoCyc/GO-BP network; Wilcoxon p-value=0.2). We also examined whether interaction type changes across species are constrained by different network parameters. We found that some interaction transitions, such as additive interaction to synergistic interaction, tend to have greater connectivity, degree, and betweenness centrality values (**Sup. Fig. S1DEFG**). In addition, connected targets had higher evolutionary rates than disconnected targets in the co-functional (Ecocyc) network (Wilcoxon p-value=0.00049; **Sup. Fig. S1C**), but not in the (EcoCyc/GO-BP) network (Wilcoxon p-value=0.061). At the same time, the co-functional (EcoCyc/GO-BP) network had more connected targets in antagonisms (18 connected, 4 disconnected) and additive interactions (165 connected, 14 disconnected) than disconnected targets, while all 24 synergistic interactions were connected (**Sup. Fig. S1I**). In contrast all evaluated targets were connected in the PPI network (**Sup. Fig. S1J**).

Discussion

We studied the evolution of drug-drug interactions (DDIs) as a proxy for studying the evolution of network topology. We found that the evolutionary rates of DDIs among gram-negative bacteria are correlated with phylogenetic distance, but that they plateau at the largest distances, suggestive of neutral processes. We then mapped drugs with known targets to different biological networks in Escherichia coli K-12 and found that the targets of synergistic DDIs are closer in biological networks than other types of DDIs, and that synergistic interactions have a higher evolutionary rate, meaning that nodes that are closer in biological networks evolve at a faster rate.

Figure 4: Evolutionary rates of drug-drug interactions vary by interaction type and path length. **A.** DDIs containing only synergies or synergy-antagonism combinations across all strains have faster evolutionary rates. Error bars represent the standard error of the mean. **B.** DDIs that are more synergistic across all strains have higher evolutionary rates. The score is the aggregated sum across strains, where DDIs were scored as +1 for synergies, 0 for additive, and -1 for antagonistic interactions. Error bars represent the standard error of the mean. **C.** Evolutionary rate as a function of interaction type for the co-functional (pink) and small/medium scale PPI network (blue). Synergistic interactions have higher evolutionary rates than additive and antagonistic interactions in *E. coli*. **D.** Evolutionary rate as a function of the minimum distance between targets. Wiring between close nodes in biological networks evolves more quickly than between distant nodes. See **Sup. Table S2** for statistical tests for differences among groups shown here.

Previous studies on interspecific comparisons of DDIs focused on the identification of novel
combinations of drugs that can be used to treat different bacterial (Brochado et al. 2018) and fungal
infections (Spitzer et al. 2011; Robbins et al. 2015). These pharmacological studies provide a static
view of DDIs in species of interest, not assessing the factors that cause interspecific divergence, and
how these factors are related to differences in interactions among drug targets across species (e.g.
via divergence in metabolic or signaling pathways). These studies suggested that chemogenomic
variation might rapidly accumulate; for example, in fungi very few combinations of antifungals act
across species (Brown et al. 2014). Similarly, the species of gram-negative bacteria in Brochado *et
al.* (2018) showed that 70% of DDIs detected across six strains were species-specific, and 20% were
strain specific.
349

Here we use this chemogenomic data to evaluate hypotheses of network evolution. Examining
pairwise distances of DDIs between species, we found that DDIs diverge more rapidly than protein
sequence between species (**Fig. 1D**). We observed a rapid accumulation of differences and overall
dissimilarity in DDI responses among species. Indeed the chemical distances curve appears
saturated among distantly related species; just as network rewiring rates slow down at this scale,
according to Shou *et al.* (2011).

We also clustered DDIs by their similarity across strains and species and then evaluated the rate of
evolution of these clusters under a Brownian motion model. This approach is analogous to methods
used in geometric morphometrics, in which specimen landmarks are grouped to evaluate different
features of an organ such as the human brain, and modularity tests are used to assess whether there
are differences in the evolution rate of different brain regions (Smaers and Vanier 2019). Modularity
among traits mirrors shared functional, genetic, and developmental pathways (Cheverud 1996);
observing high modularity among clusters indicates differences among groups of DDIs due to the
variance between clusters being greater than the variance within clusters. We observed that the
distribution of synergies, additivities and antagonisms among clusters partially explains different
rate estimates. Clusters with low rates are composed of mostly additive DDIs; however some of these
low-rate estimates are caused by resistance to one or both antibiotics, and a low phylogenetic signal
is expected when multiple strains are resistant (Michel et al. 2008). Other clusters with high rates
(e.g., clusters 14, 15, and 1) have some highly synergistic or antagonistic interactions in some strains
and additivities in other strains. With these rate estimates and interaction patterns, we asked
whether network-based metrics are correlated with DDI divergence, by linking a subset of individual
DDIs with their respective protein targets in *E. coli* co-functional (EcoCyc/GO-BP) and PPI
networks.

We first made several observations about the *E. coli* DDI types and network: synergies occur when
the targets are closer in the network, while additive interactions and antagonisms target on average
more distant proteins (**Fig. 3A, B**). This result is not surprising given the fact that synergies are
expected to be more prevalent in drug combinations that target the same pathway or that belong to
the same chemical class (Cowen and Steinbach 2008; Yeh et al. 2009), however, we think this is the
first time this pattern has been shown using network data. We also found that synergistic
interactions are more prevalent among targets that are part of the same cellular process or function,
while antagonisms can take place between targets that are in different parts of the cell (Yeh et al.
2006; Wang et al. 2012; Brochado et al. 2018). We also found targets involved in synergies have a
higher node degree (for metabolic networks), and K-edge connectivity than targets involved in

additive or antagonistic interactions (**Fig. 3C-F**). This result agrees with a previous study that reports on the positive contribution of node degree in synergy prediction (Liu et al. 2022). Average node betweenness and eigenvector centralities were also found to be more significant among synergistic DDIs compared to additivities and antagonisms in the metabolic network (**Fig. 3G-J**). This result suggests that targets affected by synergistic interactions tend to be "hubs" in the network (Li et al. 2011; Tilli et al. 2016), or that they are more central to different pathways than antagonistic or additive interactions.

We examined the rate of evolution of DDIs whose targets are closer in the PPI and metabolic networks and found that their rate of evolution is higher. This is correlated with our observation that synergies tend to occur closer in the PPI and metabolic networks and have a high evolutionary rate (accounting for the two DDI clusters with the highest rate). Antagonistic interactions are overrepresented in the cluster with the third highest rate of evolution. Correspondingly, as the path length between targets decreases, the evolutionary rate of the DDI increases (**Fig. 4D**) (although completely disconnected proteins do not follow this trend). This is consistent with the positive correlation observed between path length distances between protein interaction and gene interaction/congruence networks in yeast (Ye et al. 2005; VanderSluis et al. 2018), because the gene interaction and DDI networks should have a similar structure (Wang et al. 2012).

What do these observations mean about the overall pattern of network evolution? A possible result of why synergies have a higher rate of evolution is that the underlying network is also evolving at a faster rate resulting in big differences between species, meaning that a network effect can result in a higher rate of DDI evolution. This may mean that functional modules of the network, such as biological pathways for which synergistic DDI act, undergo a faster rate of rewiring than parts of the network that are more distantly connected (additive and antagonistic interactions). Different processes may be driving this pattern, for instance, positive selection and gain of function mutations may result in novel network combinations within an already existing pathway, that wouldn't be possible across proteins involved in different processes, expressed at different times or found in different cellular components. At the same time, purifying selection may constrain the rewiring of the network in co-functional modules, since connection losses would result in a deleterious effect on the metabolic pathway, this same situation may not be as deleterious if the targets are far apart in the network and they have very few interactions among each other. However, next, we highlight a different type of bias in DDI data that may contribute to our observed signal (high evolutionary rate of synergistic, well-connected targets in *E. coli*) and cause difficulty in the interpretation of DDI data.

Structural vs. network evolution: Another important explanation for our observed pattern (high evolutionary rate of synergistic, well-connected targets) lies in the fact that some strains are resistant to particular antibiotics. We suspect that some changes in DDI that we correlated with changes in network structure are actually caused by changes in resistance, rather than changes in the DDI. Changes in resistance are likely based on structural changes in target proteins, transporters, or proteins, and don't represent the types of biological network changes that we hoped to measure with the DDI. Note that Brochado *et al.* (2018) reported any strain with fitness values (e.g., ratio of drug/non-drug treated growth) of >0.7 as being resistant to that drug. However, some drugs that are known to be clinically bacteriostatic (inhibits growth) fall in this range, and strains are reported to be resistant; e.g. see the case of SMM resistance in all species and strains, below.

Indeed, Bushby (1973) reported "resistance to one of the drugs as measured by conventional tests
may not abolish synergy." To consider the effect of structural differences on our results, we,
therefore, repeated the entire analysis filtering for DDIs that are "susceptible" (fitness < 0.7) in at
least one of the species for both drugs, assuming that this decreases the proportion of true resistance
in the dataset. This set included 1127/2655 DDIs (42.2% of the total DDIs). Our results for this
smaller set are consistent with the full dataset (**Sup. Figs. S2-S4**). A minor difference in the subset
is that antagonisms have a slightly lower (instead of equivalent) rate than additivities. Although the
consistency between the full and reduced dataset is desirable, it doesn't remove the possibility that
differences in resistance among strains are a substantial part of the evolutionary signal.
438
439   **Network and Evolutionary Insights into Drug-drug interactions**
440   We used DDIs to learn about the evolution of networks, so discussion of some of the individual DDIs
441   may illustrate limitations of the method. The evolutionary and network parameters also provide
442   some insight into individual DDIs with clinical relevance.
443
A22-NOV
For example, A22 and novobiocin (A22-NOV) (**Sup. Table S3**): A22 binds the ATP-binding domain
of MreB (eco:b3251) (Bean et al. 2009), the actin homolog in prokaryotes, while novobiocin binds
gyrase subunit B (gyrB, eco:b3699) and topoisomerase IV (parC, eco:b3019), which are two type II
topoisomerases involved in DNA unwinding and DNA duplication (Gellert et al. 1976; Hardy and
Cozzarelli 2003). Both targets, MreB and the ParC subunit of topoisomerase IV are known to
physically interact (Madabhushi and Marians 2009) and are required for chromosome segregation
(Huang et al. 1998). A22 and novobiocin have previously been reported to be synergistic in *E. coli*
and *P. aeruginosa (Taylor 2011; Taylor et al. 2012)*; however, in the dataset from Brochado *et al.*
(2018), (**Sup. Table S3**), the interaction type is variable within species. Both *E. coli* strains are
susceptible to both drugs, but they have different interaction terms, one is additive, the other
synergistic. *Salmonella* also shows intraspecific differences, where one of the strains is synergistic
and the other additive, although in this case, the synergistic strain is resistant to novobiocin, with a
fitness of 0.85. Both *Pseudomonas* strains are resistant to novobiocin, with fitnesses greater than
0.86, and they have additive interactions. This high variability of interaction type within species
resulted in an evolutionary rate of 0.008 bliss^2/MYA for this DDI, within the cluster with the highest
evolutionary rate. The distance between these targets in the co-functional (EcoCyc/GO-BP) network
was 2, and in the DDI network the distance between targets was 4. So why are the rates so high
among strains, for this DDI? While not directly explanatory, strain-specific network predictions at
string-DB (not used in this study) also suggest strain-specific differences in local network content.
Of the genes all connected to each other in the local network (mreB, gyrA, parC, gyrB), parC is
missing in three strains, while a fourth strain has reduced connectivity for MreB (only connected to
gyrB). This type of gene content variation suggests the type of intraspecific variation that may lead
to high rates of network evolution.
468
CIP-MER
A DDI with targets in the same local network is between ciprofloxacin (CIP) and meropenem (MER).
CIP targets DNA gyrase subunit A (gyrA, eco:b2231), and MER targets D-alanyl-D-alanine
carboxypeptidase DacC (dacC, eco:b0839). The mechanism of action of this interaction is very
similar to A22-NOV. Both meropenem and A22 affect the bacterial cell wall, and ciprofloxacin and
novobiocin inhibit topoisomerases and gyrases, however, the interaction between meropenem and

ciprofloxacin is additive in all species studied in the dataset, and all strains were deemed
susceptible to the drug in this dataset (however, synergistic effects have been reported in
Pseudomonas (Ermertcan et al. 2001). For this interaction, we calculated an evolutionary rate of
0.002 bliss^2/MYA, part of the cluster that is in the middle of the other clusters when sorted by rate
(9/16). This rate is 4 times lower than that of SMM-TMP (below) and A22-NOV, which makes sense
given that the interaction is additive in all the species of the set, thus having a low trait value across
species. The distance between targets in the co-functional (EcoCyc/GO-BP) network was 1.5, while
in the PPI network, the distance between targets was 5.5.
483
SMM-TMP
Another illustrative example of a known DDI used in the clinic found in our data is the combination
of sulfamonomethoxine and trimethoprim (SMM-TMP) (**Sup. Table S1**) (Bushby 1973). These drugs
inhibit successive steps in the synthesis of tetrahydrofolic acid synthesis, necessary for the
biosynthesis of amino acids, purines, and thymidine. Sulfamonomethoxine competes against the
enzymatic substrate of dihydropteroate synthase (folP, eco:b3177); this blocks the production of
dihydrofolic acid, which in turn is a substrate for dihydrofolate reductase. Trimethoprim also
inhibits dihydrofolate reductase (folA, eco:b0048) by competing against the substrate for the binding
site of the enzyme. Even on its own, trimethoprim can contribute to "thymineless death" (Then and
Angehrn 1973); but while the effect of each drug on its own is bacteriostatic (prevents growth;
Ocampo et al. 2014), when used in combination, the combined effect is bactericidal (kills bacteria),
thus suggesting a strong synergistic effect between the two drugs (Bushby 1973; Ocampo et al.
2014).
497
We measured an evolutionary rate of 0.008 bliss^2/MYA for this DDI, as it belongs to the cluster with
the highest evolutionary rate. This rate can be explained by the difference in interaction type
between *Pseudomonas* and non-*Pseudomonas* strains. In the data used in our study, the interaction
was synergistic for *E. coli* and *S. typhimurium*, but the interaction was additive in *Pseudomonas*.
The DDI may be additive in *Pseudomonas* because it is resistant to both drugs due in part to both
differences in its permeability to the drugs and efflux pumps that remove the drugs (Eliopoulos and
Huovinen 2001). *E. coli* and *S. typhimurium* are only reported by Brochado *et al. (2018)* as resistant
to sulfamonomethoxine, having a fitness between 77-88% of non drug-treated.
506
Our goal was to capture the rate of evolution associated with differences in the folate biosynthesis
pathway across species. The fact that the two targets act as successive steps in all studied species
suggests several possible causes for this observed rate. There are some differences in the presence of
pathways components that are peripheral to these steps (Pribat et al. 2010), and it is possible that
such differences in the network beyond the direct connection contribute to DDI differences. Indeed,
although directly linked in the co-functional (EcoCyc/GO-BP) network, the targets are highly
connected to each other, with a K-edge connectivity of 21 (68% quantile), meaning that 21 edges
have to be removed to disconnect the two targets. It is also possible that there are differences in the
direct target enzymes and their connectivity (e.g., concentration, kmax). An alternative explanation
is resistance: resistance to sulfamonomethoxine and trimethoprim was detected soon after they were
introduced in clinical treatments, and several point mutations have been described in the target
genes across different bacteria, and the most common mechanism of resistance is due to the bacteria
having an extra copy of the target gene (Estrada et al. 2016). All of this suggests that, for SMM-
TMP, evolution of the DDI is the result of many types of biological differences, not just specific

differences in network interactions between the drug targets. This is an important limitation to the
DDI approach to studying network evolution.
523

Conclusions

Previous research on network evolution has focused primarily on the study of individual nodes,
either by comparing rewiring rates across networks or by comparing the role of sequence
conservation to the number of connections per node. Here, we introduce a novel framework to
compare evolutionary rates across nodes using network measurements such as minimum distance,
K-edge connectivity, and node degree between proteins. To do so, we examine changes in DDIs terms
among strains and species, and compare these rates of change with the interaction types and
network topology of the drug's targets (Lehár et al. 2007). This approach has some advantages over
direct network analyses, such as increasing the number of species under study, without the need of
obtaining the underlying network in each species. DDIs can be quantified using high-throughput
experiments across different species and strains (Brochado et al. 2018), and are phenotypic
quantitative traits that can be modeled in a phylogenetic comparative framework, allowing for an
independent measurement of evolutionary rates between nodes in the network. An important
limitation of our approach is that evolution of resistance to a drug may obscure underlying network-
based effects. Still, our approach suggests a general picture of network evolution, where close nodes
in the network (which tend to respond synergistically in response to targeting both at once) evolve at
faster rates than more distant nodes.

Materials and Methods

We obtained DDI scores (Bliss scores) from a previous study (Brochado et al. 2018) that assessed
almost 3000 combinations of 79 different compounds on six strains of three species of gram-negative
bacteria: *E. coli* K-12 BW2952 (ebw), *Escherichia coli* O8 IAI1 (ecr), *S. enterica subsp. enterica* serovar
Typhimurium 14028S (seo), *Salmonella enterica* subsp. enterica serovar Typhimurium LT2 (stm),
Pseudomonas aeruginosa PAO1 (pae), *P. aeruginosa* UCBPP-PA14 (pau). We used the following
datasets from Brochado (2018): interaction scores from table ED09C as input for our phylogenetic
comparative analysis, strain susceptibility to antibiotics from Sup. Table 1 (used for our **Sup. Figs.**
S2-S4), and the categories that DDI interactions belong to from Sup. Table 2. We built a maximum
likelihood phylogenetic tree of the six strains using PhySpeTree (Fang et al. 2019) to obtain
concatenated alignments of 35 highly conserved protein sequences (HCP). We use this concatenated
alignment of HCP sequences to construct a bayesian phylogenetic tree in BEAST2 (Bouckaert et al.
2014) using a birth-death tree with a strict molecular clock, and calibration nodes from the literature
to estimate divergence times between species (Battistuzzi et al. 2004).
556

Clustering DDI using tSNE.

558 To reduce data dimensionality of the trait space (i.e., DDI data), we classified DDIs into clusters
559 using tSNE in the R packages bigMap (Garriga and Bartumeus 2018) and bigmemory (Kane et al.
560 2013), with parameters: 80 threads, 80 layers, and 9 rounds. Dimensionality reduction was
561 performed to increase the robustness of rate estimators given the small number of species used.
562 DDIs were clustered by similarity across strains, treating each strain as a dimension, thus DDIs

that are more similar to each other across strains would cluster together under tSNE in the same cluster. Our tSNE analyses were performed for a range of 251 perplexity values between 5 and 2505 (**Sup. Doc. S1**). The clustering output was visualized and evaluated based on the stability and plateauing of cost and effect size curves, and the variance between threads. We obtained stable solutions between 175 and 1455 perplexity values. The pakde algorithm was applied with a perplexity of 1/3 the respective tSNE perplexity. To find which of the clustering solutions was the most modular, we tested for modularity in the data using the function phylo.modularity within the R package geomorph (Adams et al. 2016), the five most modular clustering patterns were selected due to their most negative ZCR coefficient, which indicates a higher degree of modularity. These clustering patterns had the following perplexity values: 355, 195, 255, 245, and 335; and the following number of clusters: 9, 15, 16, 10, and 14. All five clustering patterns had a strong modular signal, with multivariate effect sizes under -26.4, p-value=0.001 and covariance ratios below 0.92. Lastly, we fitted a multivariate Brownian motion model to each of these 5 clustering patterns and calculated the evolutionary rates per cluster using compare.multi.evol.rates, also using the R package geomorph (Adams et al. 2016). Out of the 5 most modular clustering models, the model with a perplexity of 245 was the one with a higher Z effect in the test, showing the biggest differences between groups. Thus, this clustering pattern was used in the following steps. In addition, the sigma rates calculated for each of the clusters were used as approximations for DDI evolutionary rates that are part of that cluster.

Identification of drug targets and their biological networks.

Each drug was identified with unique Pubchem and CHEMBL IDs using webchem (Szöcs et al. 2020), which were then used to retrieve their mode of action from IUPHAR (Armstrong et al. 2019). We also compared our targets to a previously published dataset on drugs and drug targets (Santos et al. 2017), identified unique Uniprot IDs and KO IDs for each target protein, and converted these IDs into *E. coli* Uniprot IDs using the KEGG Orthology (Kanehisa and Goto 2000). We didn't include in the analysis drugs whose mode of action was unknown, or that had non-protein molecules as targets, such as small molecules, RNA, or DNA.

Two biological networks of *E. coli* were downloaded from EcoliNet (Kim et al. 2015), small/medium-scale protein-protein interactions (LC) and the co-functional gene pair network of E. coli derived from EcoCyc and GO-BP.

Calculating inter-node network metrics.

For each of the networks, we calculated the average path length and node degree distributions. In addition, the minimum distance between each of the nodes in the network was calculated, as well as the node degree (number of connections per node), the K-edge connectivity between each pair of nodes (i.e. the minimum number of edges that can be removed to disconnect the nodes), betweenness centrality (i.e. a measure of centrality in the network based on shortest paths) and eigenvector centrality (i.e. a measure of the influence of the node in the network). We used the R package igraph (Csardi et al. 2006) to calculate these values in each one of the biological networks and for each node or pair of nodes. We also generated an adjacency matrix, which contains information on whether two nodes are connected directly by an edge or not. In addition, we used K-edge connectivity as a proxy for connectedness between proteins (i.e., a pair with K-edge

606   connectivity equal to zero is disconnected, and proteins with K-edge connectivity different than zero
607   are connected). Connectedness is used due to the limitations of adjacency in describing the overall
608   connection between nodes.

609   Data and code availability. The code used for data analysis is available from:
610   https://github.com/phylogenomic/ch3-netbio

References

Adams DC, Collyer M, Kaliontzopoulou A, Sherratt E. 2016. geomorph: Software for geometric morphometric analyses. Available from: https://rune.une.edu.au/web/handle/1959.11/21330

Armstrong JF, Faccenda E, Harding SD, Pawson AJ, Southan C, Sharman JL, Campo B, Cavanagh DR, Alexander SPH, Davenport AP, et al. 2019. The IUPHAR/BPS Guide to PHARMACOLOGY in 2020: extending immunopharmacology content and introducing the IUPHAR/MMV Guide to MALARIA PHARMACOLOGY. *Nucleic Acids Res.*:gkz951.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. 2000. Gene Ontology: tool for the unification of biology. *Nat. Genet.* 25:25–29.

Battistuzzi FU, Feijao A, Hedges SB. 2004. A genomic timescale of prokaryote evolution: insights into the origin of methanogenesis, phototrophy, and the colonization of land. *BMC Evol. Biol.* 4:44.

Bean GJ, Flickinger ST, Westler WM, McCully ME, Sept D, Weibel DB, Amann KJ. 2009. A22 Disrupts the Bacterial Actin Cytoskeleton by Directly Binding and Inducing a Low-Affinity State in MreB. *Biochemistry* 48:4852–4857.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A, Drummond AJ. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Comput. Biol.* 10:e1003537.

Brochado AR, Telzerow A, Bobonis J, Banzhaf M, Mateus A, Selkrig J, Huth E, Bassler S, Zamarreño Beas J, Zietek M, et al. 2018. Species-specific activity of antibacterial drug combinations. *Nature* 559:259–263.

Brown JCS, Nelson J, VanderSluis B, Deshpande R, Butts A, Kagan S, Polacheck I, Krysan DJ, Myers CL, Madhani HD. 2014. Unraveling the Biology of a Fungal Meningitis Pathogen Using Chemical Genetics. *Cell* 159:1168–1187.

Bushby SR. 1973. Trimethoprim-sulfamethoxazole: in vitro microbiological aspects. *J. Infect. Dis.* 128:Suppl:442–462 p.

Cheverud JM. 1996. Developmental integration and the evolution of pleiotropy. *Am. Zool.* 36:44–50.

Cork JM, Purugganan MD. 2004. The evolution of molecular genetic pathways and networks. *Bioessays* 26:479–484.

Cormen TH. 2001. Single-source shortest paths. *Introduction to Algorithms* [Internet]. Available from: https://cir.nii.ac.jp/crid/1572543025240409984

Cowen LE, Steinbach WJ. 2008. Stress, drugs, and evolution: the role of cellular signaling in fungal drug resistance. *Eukaryot. Cell* 7:747–764.

Csardi G, Nepusz T, Others. 2006. The igraph software package for complex network research. *InterJournal, complex systems* 1695:1–9.

Cusick ME, Klitgord N, Vidal M, Hill DE. 2005. Interactome: gateway into systems biology. *Hum. Mol. Genet.* 14 Spec No. 2:R171–R181.

Eliopoulos GM, Huovinen P. 2001. Resistance to Trimethoprim-Sulfamethoxazole. *Clin. Infect. Dis.* 32:1608–1614.

Erdős P, Gallai T. 1960. Gráfok előírt fokszámú pontokkal. *Mat. Lapok* .

Ermertcan S, Hoşgör M, Tünger O, Coşar G. 2001. Investigation of synergism of meropenem and ciprofloxacin against Pseudomonas aeruginosa and Acinetobacter strains isolated from intensive care unit infections. *Scand. J. Infect. Dis.* 33:818–821.

Estrada A, Wright DL, Anderson AC. 2016. Antibacterial Antifolates: From Development through Resistance to the Next Generation. *Cold Spring Harb. Perspect. Med.* [Internet] 6. Available from: http://dx.doi.org/10.1101/cshperspect.a028324

Fang Y, Liu C, Lin J, Li X, Alavian KN, Yang Y, Niu Y. 2019. PhySpeTree: an automated pipeline for reconstructing phylogenetic species trees. *BMC Evol. Biol.* 19:219.

Fraser HB, Hirsh AE, Steinmetz LM, Scharfe C, Feldman MW. 2002. Evolutionary rate in the protein interaction network. *Science* 296:750–752.

Garriga J, Bartumeus F. 2018. bigMap: Big Data Mapping with Parallelized t-SNE. *arXiv [cs.LG]* [Internet]. Available from: http://arxiv.org/abs/1812.09869

Gellert M, O'Dea MH, Itoh T, Tomizawa J. 1976. Novobiocin and coumermycin inhibit DNA supercoiling catalyzed by DNA gyrase. *Proc. Natl. Acad. Sci. U. S. A.* 73:4474–4478.

Ghadie MA, Coulombe-Huntington J, Xia Y. 2018. Interactome evolution: insights from genome-wide analyses of protein–protein interactions. *Curr. Opin. Struct. Biol.* 50:42–48.

Greco WR, Bravo G, Parsons JC. 1995. The search for synergy: a critical review from a response surface perspective. *Pharmacol. Rev.* 47:331–385.

Hardy CD, Cozzarelli NR. 2003. Alteration of Escherichia coli topoisomerase IV to novobiocin resistance. *Antimicrob. Agents Chemother.* 47:941–947.

Huang WM, Libbey JL, van der Hoeven P, Yu SX. 1998. Bipolar localization of Bacillus subtilis topoisomerase IV, an enzyme required for chromosome segregation. *Proc. Natl. Acad. Sci. U. S. A.* 95:4652–4657.

Jensen RA. 1976. Enzyme Recruitment in Evolution of New Function. *Annu. Rev. Microbiol.* 30:409–425.

Jin Y, Turaev D, Weinmaier T, Rattei T, Makse HA. 2013. The evolutionary dynamics of protein-protein interaction networks inferred from the reconstruction of ancient networks. *PLoS One* 8:e58134.

Jordan C. 1869. Sur les assemblages de lignes. 1869:185–190.

Kanehisa M, Goto S. 2000. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 28:27–30.

Kane M, Emerson JW, Weston S. 2013. Scalable Strategies for Computing with Massive Data. *J. Stat. Softw.* 55:1–19.

Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, Kothari A, Krummenacker M, Latendresse M, Midford PE, et al. 2018. The EcoCyc Database. *EcoSal Plus* [Internet] 8. Available from: http://dx.doi.org/10.1128/ecosalplus.ESP-0006-2018

Kim H, Shim JE, Shin J, Lee I. 2015. EcoliNet: a database of cofunctional gene network for Escherichia coli. *Database* [Internet] 2015. Available from:

http://dx.doi.org/10.1093/database/bav001

Lehár J, Zimmermann GR, Krueger AS, Molnar RA, Ledell JT, Heilbut AM, Short GF 3rd, Giusti LC,
Nolan GP, Magid OA, et al. 2007. Chemical combination effects predict connectivity in biological
systems. *Mol. Syst. Biol.* 3:80.

Li S, Zhang B, Zhang N. 2011. Network target for screening synergistic drug combinations with
application to traditional Chinese medicine. *BMC Syst. Biol.* 5 Suppl 1:S10.

Liu X, Song C, Liu S, Li M, Zhou X, Zhang W. 2022. Multi-way relation-enhanced hypergraph
representation learning for anti-cancer drug synergy prediction. *Bioinformatics* 38:4782–4789.

Madabhushi R, Marians KJ. 2009. Actin homolog MreB affects chromosome segregation by regulating
topoisomerase IV in Escherichia coli. *Mol. Cell* 33:171–180.

Michel J-B, Yeh PJ, Chait R, Moellering RC Jr, Kishony R. 2008. Drug interactions modulate the
potential for evolution of resistance. *Proc. Natl. Acad. Sci. U. S. A.* 105:14918–14923.

Ocampo PS, Lázár V, Papp B, Arnoldini M, Abel zur Wiesch P, Busa-Fekete R, Fekete G, Pál C,
Ackermann M, Bonhoeffer S. 2014. Antagonism between bacteriostatic and bactericidal
antibiotics is prevalent. *Antimicrob. Agents Chemother.* 58:4573–4582.

Picard F, Daudin J-J, Koskas M, Schbath S, Robin S. 2008. Assessing the exceptionality of network
motifs. *J. Comput. Biol.* 15:1–20.

Pribat A, Blaby IK, Lara-Núñez A, Gregory JF 3rd, de Crécy-Lagard V, Hanson AD. 2010. FolX and
FolM are essential for tetrahydromonapterin synthesis in Escherichia coli and Pseudomonas
aeruginosa. *J. Bacteriol.* 192:475–482.

Robbins N, Spitzer M, Yu T, Cerone RP, Averette AK, Bahn Y-S, Heitman J, Sheppard DC, Tyers M,
Wright GD. 2015. An Antifungal Combination Matrix Identifies a Rich Pool of Adjuvant
Molecules that Enhance Drug Activity against Diverse Fungal Pathogens. *Cell Rep.* 13:1481–
1492.

Shou C, Bhardwaj N, Lam HYK, Yan K-K, Kim PM, Snyder M, Gerstein MB. 2011. Measuring the
evolutionary rewiring of biological networks. *PLoS Comput. Biol.* 7:e1001050.

Smaers JB, Vanier DR. 2019. Brain size expansion in primates and humans is explained by a
selective modular expansion of the cortico-cerebellar system. *Cortex* 118:292–305.

Spitzer M, Griffiths E, Blakely KM, Wildenhain J, Ejim L, Rossi L, De Pascale G, Curak J, Brown E,
Tyers M, et al. 2011. Cross-species discovery of syncretic drug combinations that potentiate the
antifungal fluconazole. *Mol. Syst. Biol.* 7:499–499.

Szöcs E, Stirling T, Scott ER, Scharmüller A, Schäfer RB. 2020. webchem: An R Package to Retrieve
Chemical Information from the Web. *J. Stat. Softw.* 93:1–17.

Taylor P. 2011. Combating intrinsic antibiotic resistance in Gram-negative bacteria. Available from:
https://macsphere.mcmaster.ca/handle/11375/11329

Taylor PL, Rossi L, De Pascale G, Wright GD. 2012. A forward chemical screen identifies antibiotic
adjuvants in Escherichia coli. *ACS Chem. Biol.* 7:1547–1555.

Then R, Angehrn P. 1973. Nature of the Bactericidal Action of Sulfonamides and Trimethoprim,
Alone and in Combination. *J. Infect. Dis.* 128:S498–S501.

Tilli TM, Carels N, Tuszynski JA, Pasdar M. 2016. Validation of a network-based strategy for the
optimization of combinatorial target selection in breast cancer therapy: siRNA knockdown of
network targets in MDA-MB-231 cells as an in vitro model for inhibition of tumor development.
Oncotarget 7:63189–63203.

VanderSluis B, Costanzo M, Billmann M, Ward HN, Myers CL, Andrews BJ, Boone C. 2018.
Integrating genetic and protein–protein interaction networks maps a functional wiring diagram
of a cell. *Curr. Opin. Microbiol.* 45:170–179.

Wagner A. 2003. How the global structure of protein interaction networks evolves. *Proceedings of the
Royal Society of London. Series B: Biological Sciences* 270:457–466.

Wang Y-Y, Xu K-J, Song J, Zhao X-M. 2012. Exploring drug combinations in genetic interaction
network. *BMC Bioinformatics* 13 Suppl 7:S7.

Yeh PJ, Hegreness MJ, Aiden AP, Kishony R. 2009. Drug interactions and the evolution of antibiotic
resistance. *Nat. Rev. Microbiol.* 7:460–466.

Yeh P, Tschumi AI, Kishony R. 2006. Functional classification of drugs by properties of their pairwise
interactions. *Nat. Genet.* 38:489–494.

Ye P, Peyser BD, Spencer FA, Bader JS. 2005. Commensurate distances and similar motifs in genetic
congruence and protein interaction networks in yeast. *BMC Bioinformatics* 6:270.

Zotenko E, Mestre J, O'Leary DP, Przytycka TM. 2008. Why do hubs in the yeast protein interaction
network tend to be essential: reexamining the connection between the network topology and
essentiality. *PLoS Comput. Biol.* 4:e1000140.