HIV-1 Vpr-induced DNA damage activates NF-κB independent of cell cycle arrest and CRL4A[DCAF1] engagement

Carina Sandoval[1] and Oliver I. Fregoso[1,2]

[1] Molecular Biology Institute, University of California, Los Angeles, California, USA
[2] Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles, California, USA

Correspondence: Oliver I. Fregoso
Tel: 310.794.1424
Email: ofregoso@mednet.ucla.edu

ABSTRACT

Lentiviral accessory genes enhance replication through diverse mechanisms. HIV-1 accessory protein Vpr modulates the host DNA damage response (DDR) at multiple steps through the degradation of host proteins, cell cycle arrest, DNA damage, and both activation and repression of DDR signaling. Vpr also alters host and viral transcription; however, the connection between Vpr-mediated DDR modulation and transcriptional activation remains unclear. Here, we determined the cellular consequences of Vpr-induced DNA damage using Vpr mutants that allow us to separate the ability of Vpr to induce DNA damage from CRL4A^{DCAF1} complex dependent phenotypes including cell cycle arrest, host protein degradation, and repression of DDR. In both tissue-cultured U2OS cells and primary human monocyte-derived macrophages (MDMs), we found that Vpr induces DNA breaks and activates DDR signaling in the absence of cell cycle arrest and CRL4A^{DCAF1} complex engagement. Moreover, through RNA-sequencing, we found that Vpr-induced DNA damage alters cellular transcription via activation of NF-κB/RelA signaling. NF-κB/RelA transcriptional activation was dependent on ATM-NEMO, as inhibition of NEMO resulted in loss of NF-κB transcriptional upregulation by Vpr. Furthermore, HIV-1 infection of primary MDMs validated NF-κB transcriptional activation during infection. Both virion delivered and *de novo* expressed Vpr induced DNA damage and activated NF-κB transcription, suggesting that engagement of the DDR can occur during early and late stages of viral replication. Together, our data support a model where Vpr-induced DNA damage activates NF-κB through the ATM-NEMO pathway, independent of cell cycle arrest and CRL4A^{DCAF1} engagement. We propose this is essential to overcoming restrictive environments, such as macrophages, to enhance viral transcription and replication.

INTRODUCTION

The DDR is a signaling cascade activated in response to exogenous and endogenous genotoxic stress, such as DNA breaks. The DDR consists of sensor proteins that sense the damaged DNA, mediator and transducer proteins that transmit the signal of damaged DNA, and effector proteins that elicit a cellular response. Typically, the DDR is divided into three main pathways based on the mediator kinases activated: Ataxia telangiectasia and Rad3 related (ATR), ataxia telangiectasia mutated (ATM), and DNA-dependent protein kinase (DNA-PK). Single-strand DNA breaks (SSBs) are primarily sensed by RPA to activate ATR signaling, while double-strand DNA breaks (DSBs) can be sensed by the MRN (MRE11, RAD51, and NBS1) or Ku70/80 complexes to activate either ATM or DNA-PK signaling, respectively (1,2). However, a significant amount of crosstalk exists between these pathways. DDR signaling leads to various cellular responses, such as DNA repair, cell cycle arrest (3), transcriptional changes (4,5), apoptosis, or senescence (6).

Many diverse RNA and DNA viruses have evolved to modulate the DDR to enhance viral replication (7,8). Primate lentiviruses, such as HIV-1, primarily engage the DDR through the accessory protein Vpr (9,10). Vpr is evolutionarily conserved amongst extant primate lentiviruses and is required for replication *in vivo* (11,12) and in macrophage (13,14) and dendritic cells (15) *in vitro*. Vpr is largely unique among the lentiviral accessory genes in that it is delivered by the incoming virion, allowing it to act early in viral replication, and is also expressed *de novo* from the integrated provirus, allowing it to act later in viral replication (16). Although the primary conserved function of Vpr remains elusive, viruses lacking Vpr have decreased proviral transcription in monocyte derived macrophages (MDMs) and dendritic cells (15), suggesting an important role for Vpr in viral transcription.

Emerging literature suggests that Vpr engages the DDR at multiple, potentially unique, steps. ATR activation by Vpr leads to cell cycle arrest and requires recruitment of the CRL4A^{DCAF1} complex, which has a primary role in DNA repair (17). CRL4A^{DCAF1} complex recruitment by Vpr also leads to the degradation of many host proteins involved in the DDR, including CCDC137 (18), HLTF (19), UNG2 (20,21), SAMHD1 (22), Mus81/EME1 (23), EXO1 (24), TET2 (25), and MCM10 (26), as well as more global proteome remodeling (27). Moreover, CRL4A^{DCAF1} complex recruitment is required for Vpr-mediated repression of DSB repair (28). In contrast, we have previously shown that the ability of Vpr to induce DNA damage does not correlate with cell cycle

71 arrest or repression of DSB repair (28), suggesting that Vpr-induced DNA damage may have
72 unique roles in enhancing lentiviral replication.
73
Both DNA damage and HIV-1 Vpr promote transcriptional changes involving NF-κB (29,30). In
response to DSBs, ATM signaling promotes NF-κB transcriptional upregulation through the ATM-
NEMO pathway (31). ATM and NEMO interact in the nucleus before translocating to the
cytoplasm (32,33) to subsequently activate RelA nuclear translocation, promoter binding, and NF-
κB transcriptional activation. Previous literature suggests that HIV-1 Vpr modulates NF-κB
pathways through phosphorylation and ubiquitylation of TAK1 to enhance NF-κB signaling (34)
and altering the availability of the NF-κB p50-RelA heterodimer to inhibit NF-κB signaling (35).
This proposes a testable model where Vpr-induced DNA damage alters the cellular environment
to enhance viral replication by altering transcription through ATM-NEMO and NF-κB. Here, we
aimed to identify the cellular consequences of Vpr-induced DNA damage and the connection
between DNA damage and ATM activation, CRL4A^{DCAF1} engagement, and transcriptional
changes.
86
To determine the consequences of Vpr-induced DNA damage on cellular transcription, we used
Vpr mutants that allow us to uncouple DNA damage from cell cycle arrest and CRL4A^{DCAF1}
complex mediated host protein degradation. We found that Vpr does not require CRL4A^{DCAF1}
engagement to induce DNA breaks and activate DDR signaling in U2OS tissue-culture cells and
primary human MDMs. RNA-sequencing (RNA-seq) identified that wild-type HIV-1 Vpr and Vpr
mutants that do not induce cell cycle arrest or engage the CRL4A^{DCAF1} complex alter NF-κB
associated cellular transcription. In support of this, we showed that only Vpr proteins that induce
DNA damage activate RelA nuclear translocation and upregulate NF-κB target genes, such as
BIRC3 and CXCL8. We further assessed the requirement for ATM and NEMO signaling in Vpr-
mediated NF-κB activation. We found that inhibition of NEMO resulted in loss of NF-κB
transcriptional upregulation in primary MDMs. HIV-1 infection and virus-like particle (VLP) delivery
of Vpr in MDMs validated the upregulation of NF-κB target genes early during infection. Together,
our data support a model where Vpr-induced DNA damage activates NF-κB through the ATM-
NEMO pathway, independent of cell cycle arrest and CRL4A^{DCAF1} complex mediated host protein
degradation. This study further informs how lentiviral accessory proteins engage the DDR at
multiple and unique steps, which remodels host environment and immune pathways to promote
viral replication.
104

RESULTS

HIV-1 Vpr induces DNA damage and alters cellular transcription independent of CRL4A^{DCAF1} engagement and cell cycle arrest

Given the connections between the DDR and cellular transcription, we set out to understand whether the ability of Vpr to induce DNA damage gives rise to transcriptional changes, and whether these transcriptional changes are distinct from those caused by Vpr-mediated cell cycle arrest and/or degradation of host proteins. We have previously shown that a Vpr mutant that is unable to cause cell cycle arrest (HIV-1 Q23-17 S79A) causes DNA damage (28). However this mutant still engages the DCAF1 component of the CRL4A^{DCAF1} E3 ubiquitin ligase complex (36). To test the connection between transcriptional changes, cell cycle arrest, and host protein degradation, we used the HIV-1 Q23-17 Vpr mutant H71R, which does not bind the DCAF1 component of the CRL4A^{DCAF1} E3 ubiquitin ligase complex and thus loses the ability to cause cell cycle arrest and degrade host proteins (28) (Fig. S1A). U2OS cells were infected with rAAV expressing either HIV-1 Q23-17 Vpr wild type (WT), Vpr H71R, Vpr Q65R, a mutant of Vpr that is largely functionally dead (37), or etoposide (positive control). DDR activation was assessed by immunofluorescence while DNA damage was assessed by the comet assay 24 hours post infection. Consistent with our previous results, HIV-1 Vpr WT induces DNA breaks (Fig. 1A) and activates the DDR marker γH2A.x (Fig. 1B), while the nonfunctional Q65R mutant does not. In addition, we found that H71R induces DNA breaks (Fig. 1A) and activates γH2A.x (Fig. 1B) at levels similar to Vpr WT. To validate this, we knocked-down DCAF1 in U2OS cells with shRNAs and found that Vpr can still induce DNA breaks and activate DDR signaling when DCAF1 is knocked down (Fig. S1B-D). Together, our data show that Vpr does not require DCAF1 or associated cellular responses to induce DNA damage and activate the DDR, and that the H71R Vpr mutant allows us to investigate the ability of Vpr to induce DNA damage in the absence of host protein degradation and cell cycle arrest.

Next, we asked if Vpr mutants that induce DNA damage in the absence of cell cycle arrest (H71R and S79A) and CRL4A^{DCAF1} recruitment (H71R) alter cellular transcription. U2OS cells were infected with rAAV expressing HIV-1 Vpr WT, H71R, S79A, and Q65R, and RNA was collected at 12, 24, and 36 hours post infection. RNA-seq identified that Vpr WT, H71R and S79A, but not Q65R or empty vector, significantly alter cellular transcription 36 hours post infection when compared to untreated cells (Fig. 1C-E and Fig. S2A-B). Sixty-eight differentially expressed genes with pvalue <0.01 and FDR < 4.50E-05 were shared among Vpr WT, H71R, and S79A, indicative

of genes potentially altered by Vpr-induced DNA damage (Fig. 1D and 1E). Gene ontology

indicated the shared upregulated genes were enriched for positive regulation of NF-κB signaling

and NF-κB signaling pathways, and transcriptional regulatory relationships unravelled by

sentence-based text-mining (TRRUST) analysis (38) further identified RelA/NF-κB as a top

transcription factor activated by Vpr (Fig. 1F-G, Fig. S2 A-B and Supplemental Files S1 & S2).

Finally, we identified differentially expressed genes exclusively altered in Vpr WT expressing cells,

Vpr WT and S79A expressing cells, and unique to both Vpr H71R and S79A mutant expressing

cells (Supplemental Files S1 & S2). Collectively, our data show that Vpr alters cellular

transcription in the absence of cell cycle arrest and host protein degradation, and further suggest

that Vpr-induced DNA damage specifically alters RelA/NF-κB-mediated transcription.

HIV-1 Vpr-induced DNA damage activates RelA and promotes NF-κB transcription

We next directly assayed whether Vpr mutants that can only induce DNA damage activate

RelA/NF-κB. NF-κB activation was first validated by qRT-PCR for two NF-κB target genes

identified in our RNA-seq, BIRC3 and CXCL8 (Fig. 1E), which are important for innate immunity

and cell survival (39,40). U2OS cells were infected with rAAV expressing Vpr WT, H71R and

Q65R for 24 or 36 hours. Consistent with the RNA-seq, Vpr WT and H71R upregulate BIRC3 and

CXCL8 compared to untreated cells or Q65R mutant (Fig. 2A). qRT-PCR for BIRC3 and CXCL8

in DCAF1 knock-down cells further confirmed that DCAF1 recruitment is not required for HIV-1

Vpr to activate these two NF-κB target genes (Fig. S1E). To directly assess NF-κB activation, we

assayed for RelA localization by immunofluorescence. As RelA is cytoplasmic when inactive and

nuclear when active, we expect that Vpr WT and H71R will lead to nuclear translocation of RelA.

Indeed, we found that Vpr WT and H71R activate nuclear translocation of RelA similar to the

positive control, etoposide, while Vpr Q65R and empty vector do not (Fig 2B). These data

demonstrate that Vpr activates RelA and NF-κB transcription in the absence of cell cycle arrest,

DCAF1 recruitment and host protein degradation, suggesting that Vpr-induced DNA damage is

sufficient to drive NF-κB activation.

HIV-1 Vpr-induced DNA damage activates ATM-NEMO signaling

NF-κB is activated in response to many pathways and signals involved in DNA repair and innate

immunity (41). One such pathway is ATM-NEMO, where ATM stimulates NEMO to activate RelA

and thus NF-κB signaling (33). Previous work has shown that Vpr activates markers of both ATR

and ATM signaling (28,42). While ATR is required for Vpr-induced cell cycle arrest, the extent of

ATM activation and the cellular consequences of ATM activation without cell cycle arrest are

unclear. We hypothesized that Vpr-induced DNA damage activates the ATM-NEMO pathway,
which consequently promotes RelA/NF-κB transcription. To determine if Vpr-induced DNA
damage activates ATM signaling in the absence of cell cycle arrest and DCAF1 recruitment, we
measured activation of multiple DDR proteins including two DNA damage sensors, NBS1 and
MRE11, and two downstream signaling transducers, 53BP1 and γH2A.x. U2OS cells stably
expressing NBS1-GFP or 53BP1-GFP were infected with rAAV expressing Vpr WT, H71R, and
Q65R to assess DDR activation through live-cell imaging throughout infection. We found that Vpr
WT and H71R promote the formation of NBS1 and 53PB1 foci at 26, 33, and 48 hours post
infection (hpi), similar to the positive control etoposide, while the non-functional Q65R Vpr mutant
was indistinguishable from untreated control cells (Fig. 3A and Fig. S3A-C). As many viruses
activate or dysregulate DDR signaling through relocalization or sequestration of markers of the
DDR, we next assayed for colocalization of the DNA damage sensors MRE11 and NBS1, and
transducers 53BP1 and γH2A.x. Both damage sensors MRE11 and NBS1, and transducers
53PB1 and γH2A.x, colocalize in Vpr WT and H71R, similar to etoposide treated cells (Fig. 3B&C),
suggesting that Vpr-induced DNA damage activates, but does not dysregulate, classical ATM
signaling.
189
To determine if NEMO is required for Vpr to upregulate NF-κB transcription, we inhibited NEMO
using a cell-permeable NEMO binding domain (NBD) inhibitor peptide (43). U2OS cells were
pretreated with NBD peptide and infected with rAAV expressing Vpr WT and mutants. We
assessed upregulation of the NF-κB target genes BIRC3 and CXCL8 using qRT-PCR. As before,
Vpr WT and H71R upregulated both BIRC3 and CXCL8, while Q65R and empty vector did not
(Fig. 4). With the addition of the NBD, Vpr WT and H71R mutant lost the ability to upregulate
BIRC3 and CXCL8 (Fig. 4), suggesting that Vpr requires NEMO activation to upregulate
transcription of NF-κB target genes.
198
Virion delivery of Vpr, through VLPs or HIV-1 ∆ENV infection, in primary MDMs activates
ATM-NEMO signaling and promotes RelA/NF-κB transcription
Together our data proposes a model where Vpr-induced DNA damage activates ATM and NEMO
signaling resulting in RelA nuclear translocation and NF-κB transcriptional upregulation. We next
tested this model in primary human MDMs, where Vpr enhances viral transcription and replication
(14). Primary MDMs (Fig. S4A) were infected with virus-like particles (VLPs) carrying
physiological levels of Vpr WT, H71R, or Q65R (Fig. S4B), and assayed for DNA damage, NF-κB
activation, and NEMO dependence (Fig. 5A). Similar to overexpression of Vpr in U2OS tissue

culture cell lines, Vpr WT and H71R mutant induced DNA breaks and activated γH2A.x compared
to Vpr Q65R and empty VLPs (Fig. 5B-C). Consistent with our data in U2OS cells, Vpr WT and
H71R, but not Q65R or empty VLPs, upregulated BIRC3 and CXCL8 similar to the TNFα positive
control when compared to empty cells (Fig. 5D and Fig. S4C). Finally, NBD delivery prior to VLP
infection blocked Vpr WT and H71R-mediated BIRC3 and CXCL8 transcriptional activation, but
not Vpr-induced DNA damage (Fig. 5C-D and Fig. S4C). These data indicate that physiological
levels of virion-delivered Vpr induces DNA damage that activates NF-κB transcription in a NEMO
dependent manner in primary MDMs.

To determine if HIV-1 infection activates NF-κB target genes, we infected primary human MDMs
from six different donors with HIV-1 ΔEnv and assayed for transcriptional changes at early (8
hours), intermediate (16 hours), and late (24 and 48 hours) timepoints post infection. HIV-1
infection was monitored by flow cytometry for HIV-1 core proteins and qRT-PCR for unspliced
transcripts (Fig. S4D-E), while NF-κB activation was measured by qRT-PCR for CXCL8 and
BIRC3. TNFα was used as positive control (Fig. S4F). HIV-1 ΔEnv infection resulted in
upregulation of CXCL8 and BIRC3 in MDMs as early as 8hpi in all donors, despite some donor-
to-donor variability in the magnitude of upregulation (Fig. 5E and Fig. S4G-H), suggesting that
HIV-1 infection upregulates Vpr-induced NF-κB target genes early during infection, consistent
with VLP delivery of Vpr. Jointly, our data support a model where both incoming HIV-1 Vpr and
de novo produced Vpr induces DNA damage that activates ATM-NEMO signaling to upregulate
NF-κB transcription (Fig. 6).

228
DISCUSSION

In this study, we tested the hypothesis that HIV-1 Vpr alters cellular transcription through induction
of DNA damage and activation of DDR signaling. By leveraging well-characterized Vpr mutants
that separate Vpr-induced DNA damage from cell cycle arrest and CRL4A^{DCAF1} complex mediated
host protein degradation, we identified a current model where Vpr-induced DNA damage activates
ATM-NEMO signaling, which stimulates RelA and upregulates NF-κB mediated transcription.
Using U2OS and primary human MDMs, which allowed us for the first time to investigate how
HIV-1 Vpr engages the DDR in this important cell type, we showed that Vpr induces DNA breaks
and activates markers of ATM signaling independent of cell cycle arrest, CRL4A^{DCAF1} complex
recruitment, and host protein degradation. Moreover, we showed that Vpr-induced DNA damage
correlates with RelA/NF-κB activation, as assayed by bulk RNA-seq, and RelA

immunofluorescence. We validated the upregulation of two NF-κB target genes, BIRC3 and CXCL8, and showed that inhibition of NEMO ablates the ability of Vpr to upregulate NF-κB transcription. Finally, VLP delivery of Vpr and HIV-1 infection show that incoming virion-associated Vpr is sufficient to induce DNA damage and activate RelA/NF-κB transcription. In complement, rAAV expression of Vpr show that *de novo* expressed Vpr can also induce DNA damage and activate RelA/NF-κB transcription. These data suggest that Vpr-induced DNA damage can play roles in both early and late stages of viral replication. Overall, our data support a model where Vpr-induced DNA damage activates ATM-NEMO signaling and upregulates RelA/NF-κB transcription in cell lines and primary human MDMs.

Although Vpr is a multifunctional and enigmatic protein, the DNA damage response is central to many of the phenotypes associated with Vpr. Previous reports from our lab and others have worked to untangle how and why Vpr engages the DDR. A consensus is emerging that Vpr engages the DDR at multiple, potentially unique, steps. Through recruitment of the CRL4A^{DCAF1} ubiquitin ligase complex, Vpr degrades various DDR proteins, activates ATR signaling, represses double-strand DNA break repair, and causes cell cycle arrest (28,44). However, as shown here, Vpr induces DNA damage and alters cellular transcription independent of its ability to engage the CRL4A^{DCAF1} ubiquitin ligase complex and associated phenotypes. Functions independent of host protein degradation are unique among most lentiviral accessory genes, as their typical primary functions depend on subverting antiviral mechanisms through host protein degradation (45). These data further define the multifunctional nature of Vpr in enhancing lentiviral infection and support the model where Vpr engages the DDR at multiple independent steps.

Here, we have also shown that Vpr-induced DNA damage directly correlates with NF-κB transcriptional activation that is dependent on ATM signaling. We have demonstrated that Vpr activates markers of ATM signaling, such as MRE11, NBS1, γH2A.x, and 53PB1, in a manner that resembles host activation rather than dysregulation. Antagonism or dysregulation of DNA damage sensors through relocalization or sequestration is a conserved mechanism among many viruses to evade innate immune detection (46). While we have shown that Vpr-induced DNA damage activates markers of ATM signaling in a manner that seems to facilitate classical ATM activation, with the end goal of activating transcription, it remains to be seen whether Vpr-mediated activation of ATR, which correlates with CRL4A^{DCAF1} recruitment and cell cycle arrest, is classically activated or dysreglated. Moreover, how Vpr induces DNA breaks leading to both ATM and ATR activation is not understood.

Complementary studies have shown that Vpr modulates cellular (47) and viral transcription (48,49). In the context of cellular transcription, Vpr activates genes associated with innate immunity and proliferation in CD4+ T cells (50) and promotes the expression of proinflammatory cytokines in MDMs and monocyte-derived dendritic cells (MDDCs) (15,25). Together this suggests that Vpr has a vital role in transcriptional reprogramming to potentially create a proinflammatory environment that is conducive for viral replication. Consistent with these studies, we identified by bulk RNA-seq that Vpr regulates innate immune pathways that are either independent of or dependent on cell cycle arrest and CRL4A^{DCAF1} engagement and host protein degradation. Our data show that Vpr downregulates TASOR target genes (Fig. S2 D-F), which is consistent with the literature suggesting that SIV and HIV-2 Vpr counteract the human silencing hub (HUSH) repressor complex (51,52). Furthermore, we found that Vpr downregulates CXCL14, which aligns with the data reporting that human papillomaviruses downregulate CXCL14 to enhance replication (53).

Our data also indicate that Vpr activates the RelA/NF-κB immune pathway without induction of cell cycle arrest and CRL4A^{DCAF1} recruitment and host protein degradation. This is consistent with previous reports showing that Vpr activates NF-κB transcription via phosphorylation of TAK1 (34), an upstream regulator of NF-κB. We further found that Vpr-induced DNA damage activates ATM-NEMO signaling to upregulate RelA/NF-κB transcription, adding to the mechanistic understanding of NF-κB activation by Vpr. Through engagement with the CRL4A^{DCAF1} complex, Vpr has also been found to repress NF-κB activation by altering the availability of the NF-κB p50-p65 heterodimer, thus limiting proinflammatory cytokine expression (35). While it remains unclear what differentiates Vpr-mediated NF-κB activation from repression, whether Vpr engages the CRL4A^{DCAF1} complex seems to be a distinguishing factor. Furthermore, it is clear that in all cases Vpr carefully modulates NF-κB activation without globally activating interferon, which would inhibit viral replication. Thus, further studies understanding how Vpr manages to activate only aspects of NF-κB signaling (54,55) will help to define how Vpr may contribute to the ability of HIV to subvert the innate immune response.

Although various studies have shown that Vpr also alters viral transcription the mechanisms are unclear and disparate. For example, Vpr-mediated LTR activation is associated with cell cycle arrest where the LTR is most transcriptionally active (56,57), degradation of host proteins such as CCDC137 (18), and CRL4A^{DCAF1} independent mechanisms (58). However, studies have

shown that Vpr also promotes LTR transcription in noncycling cells (59,60) where induction of cell

cycle arrest is absent and the necessity for degradation of specific host proteins in LTR activation

has not been extensively examined. One benefit of Vpr activating NF-κB signaling via ATM is the

potential direct enhancement of LTR transcription, as the HIV-1 LTR contains multiple NF-κB

binding sites essential for viral gene expression. In addition to NF-κB, we further identified that

Vpr activates CEBPB and JUN transcription factors involved in LTR activation. This is similar to

Vpr-mediated de-repression of the LTR by removal of the transcriptional repressor ZBTB2

following ATR activation (61). Overall, Vpr WT and mutants that induce only DNA damage globally

alter cellular transcription, upregulate NF-κB signaling and transcription factors that promote LTR-

driven transcription, suggesting that Vpr-induced DNA damage is important for promoting LTR

transcription to enhance viral replication.

Overall, our data suggest that during HIV infection, incoming Vpr and *de novo* expressed Vpr

prime the cellular environment by activating RelA/NF-κB signaling to promote LTR transcription

and enhance viral replication in macrophages. Our data align with the growing body of literature

that supports the role of accessory proteins modulating the host environment and the DDR to

promote viral replication through mechanisms independent of host protein degradation.

	FIGURE LEGENDS

Figure 1: HIV-1 Vpr induces DNA damage and alters cellular transcription independent of cell cycle arrest and CRL4A^{DCAF1} engagement. (A) Comet assay of U2OS cells infected with rAAV expressing 3xFLAG-tagged Vpr WT, H71R, Q65R, untreated (negative control), or 50uM etoposide (positive control). Percent tail DNA was quantified at 24 hours post infection (hpi) using the OpenComet plug-in for the ImageJ software. Each circle represents one cell. N=3, one representative experiment shown. (B) Representative immunofluorescence images of U2OS cells infected under the same conditions as Fig. 1A; γH2A.x (magenta) and nuclei stained with DAPI (blue). Images were taken at 63x magnification. N=3, one representative experiment shown. EV, empty vector. (C) RNA-seq of U2OS cells at 36 hpi. Heat map displays Log$_2$ fold changes of upregulated genes in red and downregulated genes in blue. Each column represents a biological replicate, n=3 (D) Venn diagram of the top 100 upregulated and top 100 downregulated genes among Vpr WT, S79A, H71R, and Q65R. E) Dot plot of the 68 conserved differentially expressed genes among Vpr WT, S79A, and H71R. BIRC3 and CXCL8 are highlighted as the two most upregulated genes (Log$_2$ fold change) under all conditions. (F) Gene ontology and (G) TRRUST analysis of the 30 upregulated genes performed with Metascape software. Asterisk indicate statistical significance compared to untreated control, as determined by one-way ANOVA test (NS, nonsignificant; *** P< 0.0004, **** P< 0.0001). Related to Figures S1 and S2.

Figure 2: HIV-1 Vpr activates RelA and promotes NF-κB transcription independent of cell cycle arrest and CRL4A^{DCAF1} engagement. (A) qRT-PCR validation of upregulated NF-κB target genes BIRC3 and CXCL8. Normalized expression (ΔΔCt) was calculated by normalizing to GAPDH followed by calculating fold change to untreated or empty vector treated cells. Cells treated under the same conditions as Fig. 1C. n=3, one representative experiment shown. (B) Representative immunofluorescence images of U2OS cells infected under the same conditions as Fig. 1A; RelA (red) and nuclei stained with DAPI (blue). Images were taken at 63x magnification. n=3, one representative experiment shown. Asterisk indicate statistical significance compared to untreated control, as determined by one-way ANOVA test (NS, nonsignificant; *** P< 0.0003, **** P< 0.0001).

Figure 3: Vpr-induced DNA damage activates ATM signaling independent of cell cycle arrest and CRL4A^{DCAF1} engagement. (A) Live-cell imaging of U2OS cells stably expressing NBS1-GFP or 53PB1-GFP. Cells were infected under the same conditions as Fig. 1A and

representative images were taken at 33 hpi at 63x magnification. n=3, one representative

experiment shown. (B) Representative images of co-localization of DNA damage sensors MRE11

(magenta) and NBS1 (yellow) in U2OS NBS1-GFP or (C) DNA damage transducers yH2A.x

(magenta) and 53PB1 (yellow) in U2OS 53PB1-GFP cells infected under the same conditions as

Fig. 1A. Representative images taken at 24 hpi at 63x magnification. n=3, one representative

experiment shown. Related to Figure S3.

Figure 4: Vpr upregulates NF-κB transcription dependent on NEMO, yet independent of cell

cycle arrest and CRL4A^{DCAF1} engagement. qRT-PCR of BIRC3 and CXCL8 in the presence or

absence of Nemo-Binding Inhibitor peptide (NBD) at 36 hpi. U2OS cells were pretreated with

100uM of NBD for 2 hours prior to infection with rAAV expressing 3xFLAG-tagged Vpr WT, H71R,

Q65R, empty vector (negative control). Normalized expression to GAPDH. Asterisk indicate

statistical significance compared to untreated control, as determined by one-way ANOVA test

(NS, nonsignificant; **** P< 0.0001).

Figure 5: Vpr-induced DNA damage activates ATM-NEMO signaling and NF-κB

transcription in primary MDMs early during infection. (A) Experimental schematic depicting

the isolation and differentiation of primary human monocyte-derived macrophages (MDMs) from

peripheral blood mononuclear cells (PBMCs) followed by delivery of Vpr via Virus-like Particles

(VLPs) or HIV-1 infection. Infected MDMs were assayed for induction of DNA damage, activation

of DDR signaling, and the upregulation of NF-κB target genes in the context of NEMO inhibitor.

(B) Comet assay of MDMs treated with VLPs packaging 3xFLAG-tagged Vpr WT, H71R, Q65R,

empty VLPs (negative control), or 25uM Mitomycin C (positive control) for 8 hours. Comet assay

analysis was performed as in Fig. 1A. n=2, one representative experiment shown. (C)

Representative immunofluorescence images of MDMs infected with VLPs as in Fig. 5B and

treated or not with Nemo-Binding Inhibitor peptide (NBD); γH2A.x (magenta) and nuclei stained

with DAPI (blue). Images were taken at 100x magnification. n=3, one representative experiment

shown. (D) qRT-PCR for CXCL8 of MDMs treated with NBD as in Fig. 5C with TNFα as positive

control. n=3, one representative experiment shown. (E) qRT-PCR for CXCL8 of MDMs infected

with HIV-1 ΔEnv for 8, 16, 24, and 48 hpi from four separate donors. Normalized expression to

GAPDH. Asterisk indicate statistical significance compared to untreated control, as determined

by one-way ANOVA test (NS, nonsignificant; * P<0.03, ** P < 0.005, **** P< 0.0001). Related to

Figure S4.

Figure 6: Model showing Vpr-induced DNA damage activates ATM and NEMO signaling resulting in RelA nuclear translocation and NF-κB transcriptional upregulation. 1. Vpr induces double-strand DNA breaks independent of cell cycle arrest and CRL4A^{DCAF1} complex engagement. 2. Vpr-induced DNA damage activates markers of ATM signaling including MRE11, NBS1, γH2A.x, and 53PB1. 3. ATM-DDR activation results in NEMO relieving RelA inhibition from inhibitor of NF-κB (IκBα and IκBβ). 4. RelA translocates into the nucleus. 5. RelA binds to NF-κB promoters and activates transcription (txn) of NF-κB target genes such as BIRC3 and CXCL8. (Schematic made on Biorender).

SUPPLEMENTAL FIGURE LEGENDS

Figure S1: Knockdown of DCAF1 does not abrogate the ability of Vpr to induce DNA damage, activate DDR signaling, or upregulate NF-κB target genes. (A) Schematic representation of Vpr mutants used in these studies to decouple some Vpr functions. Vpr S79A and Vpr H71R do not induce cell cycle arrest, and H71R also does not bind DCAF1 of the CRL4A^{DCAF1} complex to degrade host proteins or remodel the proteome. Vpr Q65R (not shown) is a largely non-functional negative control. (B) Western blot of endogenous DCAF1 U2OS cells to test stable knockdown with shRNA. Control (ctl) shRNA and two DCAF1-specific shRNAs (shRNA1 and shRNA2) were used. Actin is used as a loading control. (C) Comet assay of DCAF1 knock-down cells infected with rAAV and analyzed as in Fig. 1A. n=1. (D) Representative immunofluorescence images of DCAF1 knock-down U2OS cells infected under the same conditions as Fig. 1A.; γH2A.x (magenta) and nuclei with DAPI (blue). Images were taken at 63x magnification. n=1, one representative image shown. (E) qRT-PCR for CXCL8 and BIRC3 in DCAF1 knock-down U2OS cells infected under the same conditions as Fig. 1A. n=1. Asterisk indicate statistical significance compared to untreated control, as determined by t-test (NS, nonsignificant; * P< 0.02, ** P<0.004, *** P<0.0003, **** P<0.0001). EV, empty vector negative control; Etop., etoposide positive control. Related to Figures 1 and 2.

Figure S2: Vpr alters cellular transcription through diverse mechanisms that are either dependent or independent of cell cycle arrest and CRL4A^{DCAF1} engagement. (A) Gene ontology using Metascape displaying TRRUST analysis for upregulated genes. (B) Principal component analysis (PCA) of the RNA-seq data showing the relatedness of the three biological replicates compared to empty vector or untreated cells. Related to Figure 1.

Figure S3: Vpr-induced DNA damage activates markers of ATM signaling. (A) Live-cell

imaging taken on the IncuCyte of U2OS NBS1-GFP or 53PB1-GFP cells infected as in Fig. 1A

for 48 hrs. n=2, one representative experiment shown. (B & C) Quantification of DNA damage

induced in the live-cell imaging experiment in Fig. 3A at 8, 26, 33, and 48hpi. Foci per cell and

foci size were quantified using ImageJ software. Images were taken at 63x magnification using

the LSM 900. n=3, one representative experiment shown. EV, empty vector; AU, arbitrary units.

Related to Figure 3.

Figure S4: Infection of primary MDMs with VLPs packaging Vpr protein or HIV-1 infection

upregulate NF-κB transcription dependent on NEMO. (A) Flow cytometry plots displaying

differentiation of PBMC-derived monocytes to macrophages. Primary MDMs are CD14+, CD45+

and CD16+. n=6, one representative experiment shown. (B) Western blot of 3X FLAG-Vpr WT

and mutants packaged in Virus-like particles (VLPs). (C) qRT-PCR for BIRC3 of MDMs treated

with NBD as in Fig. 5C with TNFα as positive control. n=3, one representative experiment shown.

(D) Histogram quantifying percent of infected MDMs with HIV-1 ΔEnv at 48hpi. n=6, one

representative experiment shown. (E) qRT-PCR for unspliced HIV-1 RNA at 8, 16, 24 and 48hpi.

n=6, one representative experiment shown. (F) qRT-PCR for TNFα of MDMs infected with HIV-1

ΔEnv at 8, 16, 24 and 48hpi. n=6, one representative experiment shown. (G & H) qRT-PCR for

BIRC3 or CXCL8 of MDMs infected with HIV-1 ΔEnv at 8, 16, 24 and 48hpi in separate donors.

n=6, (G) is all 6 donors, (H) is donors 5 and 6 in addition to those shown in Fig. 5E. Asterisk

indicate statistical significance compared to untreated control, as determined by one-way ANOVA

test (NS, nonsignificant; **** P<0.0001). Related to Figure 5.

Files S1 & S2: HIV-1 Vpr WT and mutants alter cellular transcription. Analysis of all

upregulated (File S1) and downregulated (File S2) genes in the RNA seq analysis that passes

statistical significance of Log2 (1.25), p < 0.01, and FDR < 4.50E-05. Counts per million reads

mapped (CPM) Log2 fold changes were calculated relative to untreated control. Related to Figure

1 and Figure S2.

MATERIALS AND METHODS

Plasmids

pcDNA-3xFLAG-Vpr and pscAAV-mCherry-T2A-Vpr WT and mutant plasmids were generated as

previously described (28). For rAAV production, pHelper and pAAV-2.5 capsid plasmids were

used (Addgene and (28). For VLP production, psPAX2 and pmD2.G were used (Addgene). For

HIV-1 ΔENV production, Bru-GFP ΔENV was generated as previously described (62) and

pmD2.G (Addgene). For lentivirus stable knock-down of DCAF1, pLKO.1 (Addgene),

psPAX2(Addgene) and pmD2.G (Addgene) were used.

Generation of Viruses

rAAV vectors packaging the pscAAV-mCherry-T2A-Vpr WT or mutant plasmids were generated

by transient transfection of HEK 293 cells using polyethyleneimine (PEI) as previously described

(63). Virus-like particles (VLPs) packaging Vpr WT or mutant proteins were generated by transient

transfection of HEK 293T using TransIT-LT1 (Mirus). VLPs were harvested 48hrs post

transfection, concentrated through a 25% sucrose cushion at 24,000 rpm for 3 hrs at 4°C, and

resuspended in PBS. Protein packaging was validated through western blot. HIV-1 ΔENV

pseudotyped with VSV-G were generated by transient transfection of HEK 293T using TransIT-

LT1 (Mirus). Virus was collected 48hrs post transfection. Lentivirus expressing DCAF1 shRNAs

were generated by transfection of HEK 293T using TransIT-LT1 (Mirus).

shRNA1:

AATTCAAAAAGCTGAGAATACTCTTCAAGAACTCGAGTTCTTGAAGAGTATTCTCAGC

shRNA2:

AATTCAAAAAAGCACTTCAGATTATCATCAACTCGAGTTGATGATAATCTGAAGTGCT.

Media was changed 18 hours post transfection. Lentiviruses were harvested 48 hours post media

change and filtered through a 0.45 µm PES filter.

Cell Lines and Cell Culture

Human bone osteosarcoma epithelial (U2OS), Human embryonic kidney (HEK) 293, and HEK

293T cells were cultured as adherent cells directly on tissue culture plastic (Greiner) in Dulbecco's

modified Eagle's medium (DMEM) growth medium (high glucose, L-glutamine, no sodium

pyruvate; Gibco) with 10% fetal bovine serum (FBS) (Gibco) and 1% penicillin-streptomycin

(Gibco) at 37°C and 5% CO_2. All cells were harvested using 0.05% trypsin-EDTA (Gibco). The

panel of U2OS cells stably expressing 53PB1-GFP (64) and NBS1-GFP (65) were kindly provided

by Claudia Lukas (University of Copenhagen, Denmark). Stable DCAF1 knock-down cells were

generated by delivering lentiviruses expressing DCAF1 shRNAs or empty vector to U2OS cells.

U2OS with integrated lentiviral construct were selected for with 1ug/ml of Puromycin. Knock-down

of DCAF1 was assessed through western blots.

Monocyte-Derived Macrophages (MDMs)

Human peripheral blood mononuclear cells (PBMCs) were obtained from human donors at the

UCLA/CFAR Virology Core Laboratory. Primary monocytes were isolated from PBMCs by

negative selection using the EasySep™ Human T Cell Isolation Kit (STEM CELL) and were

differentiated into monocyte-derived macrophages (MDMs) by stimulation with 20ng/mL

macrophage colony-stimulating factor (M-CSF) (R&D Systems). MDMs were cultured in Roswell

Park Memorial Institute (RPMI) 1640 growth medium (L-Glutamine) with 10% fetal bovine serum

(Gibco) at 37°C and 5% CO_2 for 7 days while replenishing media every 3 days.

Alkaline Comet Assay

The alkaline comet assay and data analysis was performed as previously described (28), with

minor changes. MDMs were infected with VLPs delivering Vpr WT and mutants at equal protein

levels or 25uM Mitomycin C (Cayman). Cells were harvested with Accutase (STEM CELL) at 10

hrs post infection and resuspended in 0.5% low-melting-point agarose at 37°C. Images were

acquired on the Zeiss Axio Imager Z1. Images were analyzed using the OpenComet plug-in for

ImageJ.

RNA-sequencing

Total RNA from U2OS cells was isolated using TRIzol® reagent (Invitrogen). RNA integrity was

assessed using the Bioanalyzer TapeStation 4200 RNA High Sensitivity (Agilent). Library

Preparation was completed using the KAPA mRNA HyperPrep Kit (Illumina) enriching for Poly(A)

RNA with magnetic oligo-dT beads and attaching unique dual indexed adapters. Quality control

of the library was done with the Bioanalyzer TapeStation 4200 D1000 High Sensitivity (Agilent).

RNA was sequenced using the Hiseq3000 1x50 at the UCLA Technology Center for Genomics &

Bioinformatics core. Reads were aligned to the Human h38 STAR genome and gene counts were

determined using edgeR. log_2 reads per kilobase of transcript per million reads mapped (rpkms)

were calculated for samples compared to untreated cells. Heat map was generated using

pheatmap hierarchical clustering on z-score log2 rpkms. Gene ontology and TRRUST analysis

were done on Metascape.

Quantitative Reverse Transcription PCR (qRT-PCR)

Total RNA was isolated using the PureLink™ RNA Mini Kit (Invitrogen). RNA was reverse transcribed using SuperScript IV First-Stand Synthesis System (Invitrogen) with Oligo(dT) primers. qRT-PCR was performed with PowerTrack SYBR Green Master Mix (Thermo Fisher Scientific) on the LightCycler 480 System (Roche) with the following primers (5' to 3') BIRC3 AAGCTACCTCTCAGCCTACTTT and CCACTGTTTTCTGTACCCGGA, CXCL8 TTTTGCCAAGGAGTGCTAAAGA and AACCCTCTGCACCCAGTTTTC, TNFα CTCTTCTGCCTGCTGCACTTTG and ATGGGCTACAGGCTTGTCACTC, HIV-1 unspliced RNA GCGACGAAGACCTCCTCAG and GAGGTGGGTTGCTTTGATAGAGA, and GAPDH CAAGATCATCAGCAATGCCT and AGGGATGATGTTCTGGAGAG. mRNA levels were quantified by calculating $\Delta\Delta$Ct. Target transcript Ct values were normalized to the Ct value of the housekeeping gene GAPDH followed by calculating fold change to untreated or empty vector treated cells.

Live-Cell Imaging

U2OS-53PB1-GFP and U2OS-NBS1-GFP cells were imaged using the IncuCyte S3 Live-Cell Analysis Instrument (Sartorius). Mean Fluorescence Intensity (MFI) was calculated using the Sartorius software. For higher resolution, U2OS-53PB1-GFP and U2OS-NBS1-GFP cells were imaged using the LSM 900. Foci per cell and foci size were analyzed using ImageJ.

Immunofluorescence

Cells were plated in 6- or 24-well tissue culture plates (Greiner) and allowed to adhere overnight, then infected with VLP (equal protein expression), rAAV-2.5 (1.4×10^8 copies/well), or etoposide (Sigma). U2OS cells were permeabilized with 0.5% Triton X-100 in PBS at 4°C for 5 min and fixed in 4% PFA for 20 min, MDM cells were permeabilized with 0.1% Saponin (Fisher) in PBS at 4°C for 15 min and fixed in 4% PFA for 15 min. Cells were washed, incubated with blocking buffer (3% BSA, 0.05% Tween 20, and 0.04 NaN_3 in PBS for U2OS cells or 3% FBS in 0.1% Saponin for MDM cells) for 30 min. Cells were probed with appropriate primary antibodies (anti-γH2A.x Ser139, anti-53BP1, or anti-RelA(p65) [Cell Signaling], anti-GFP [Takara], and anti-MRE11 [Novous]) and then washed and probed with Alexa Fluor-conjugated secondary antibodies (Life Technologies). Nuclei were stained with diamidino-2-phenylindole (DAPI; Life Technologies). Images were acquired on the LSM 980.

Western Blots

Protein was collected from cells as previously described (28). Membranes were blocked in intercept blocking buffer (Li-COR Biosciences). Immunoblotting was performed using mouse anti-FLAG M2 (Sigma-Aldrich), rabbit anti-Actin (Bethyl), or mouse anti-DCAF1 (Proteintech) for 1hr. Blots were incubated with secondary antibodies IRDye 800CW anti-Rabbit and IRDye 680RD anti-Mouse (Li-COR Biosciences) for 1 hr and then visualized using the Li-COR Odyssey M (Li-COR Biosciences).

HIV-1 ΔEnv Infection

MDMs were plated in 96-well tissue culture plates (Greiner) and allowed to adhere overnight. MDMs underwent spinfection with HIV-1 ΔEnv pseudotyped with VSV-G at 1200× g for 90 min at 37°C. Infection was assessed 48 hours after infection via qRT-PCR and Flow Cytometry.

Flow Cytometry

Isolated monocytes and infected MDMs were lifted from tissue culture plates using accutase (STEM CELL). Cells were stained for CD14-FITC, CD45-APC, or CD16-APC (STEM CELL) for 30 min at 4°C, washed with PBS, fixed in 4% PFA for 15 min, permeabilized with 0.1% Triton X-100 in PBS at 4°C for 15 min, then washed with PBS. Cells were probed for HIV-1 core antigen-FITC KC57 (Beckman Coulter) for 1hr at 4°C then washed with PBS and resuspended in FACS buffer (5% FBS in PBS). Events were assessed by flow cytometry on the AttuneNxT (Thermo Fisher Scientific). At least 10,000 events were collected per run. Data was analyzed using FlowJo software.

Statistical Analyses

All statistical analyses were performed using GraphPad Prism 9.

AUTHOR CONTRIBUTIONS

Conceptualization, C.S. and O.I.F; methodology, C.S.; writing original draft C.S and O.I.F; writing, review and editing C.S. and O.I.F; visualization C.S. and O.I.F.; supervision and funding acquisition O.I.F. All authors have read and agreed to the published version of the manuscript.

FUNDING

This work was supported by NIH NIAID grant R01AI147837 to O.I.F. C.S was supported by NIH NIAID F31 Ruth L. Kirschstein National Research Service Award Predoctoral Fellowship F31-

593   AI165286, and the NIH NIGMS Grant GM007185 T32 Ruth L. Kirschstein National Research
594   Service Award Cell and Molecular Biology Training Grant. The funders had no role in study
595   design, data collection and interpretation, or the decision to submit the work for publication.
596
ACKNOWLEDGEMENTS
We like to thank Dr. Julia Mack and Julianne Ashby for their assistance and expertise with Live-
cell imaging using the LSM900. We thank Dr. Alexander Hoffmann and Dr. Diane Lefaudeux for
her instruction on RNA-seq analysis. We thank Dr. Claudia Lukas for providing U2OS cells that
stably express 53PB1-GFP and NBS1-GFP. We thank Dr. Steven Bensinger for the LightCycler
480 System, Dr. Jesse Zamudio for the Qubit fluorometer, Dr. Peter Bradley for the Zeiss Axio
Imager Z1, Dr. Matteo Pellegrini for the TapeStation, Dr. Steve Jacobsen and Dr. Jefferey Long
for the LSM980. We thank Dr. Randilea Nichols Doyle, Karly Nisson, and Vivian Yang for
providing comments on the manuscript.
606
607   **COMPETING INTERESTS**
608   The author(s) declare(s) they have no competing interests.

REFERENCES

1. Blackford AN, Jackson SP. ATM, ATR, and DNA-PK: The Trinity at the Heart of the DNA Damage Response. Mol Cell. 2017 Jun 15;66(6):801–17.

2. Shiloh Y, Ziv Y. The ATM protein kinase: regulating the cellular response to genotoxic stress, and more. Nat Rev Mol Cell Biol. 2013 Apr;14(4):197–210.

3. Matthews HK, Bertoli C, de Bruin RAM. Cell cycle control in cancer. Nat Rev Mol Cell Biol. 2022 Jan;23(1):74–88.

4. Valerie K, Delers A, Bruck C, Thiriart C, Rosenberg H, Debouck C, et al. Activation of human immunodeficiency virus type 1 by DNA damage in human cells. Nature. 1988 May;333(6168):78–81.

5. Dunphy G, Flannery SM, Almine JF, Connolly DJ, Paulus C, Jønsson KL, et al. Non-canonical Activation of the DNA Sensing Adaptor STING by ATM and IFI16 Mediates NF-κB Signaling after Nuclear DNA Damage. Mol Cell. 2018 Sep 6;71(5):745-760.e5.

6. Gorgoulis V, Adams PD, Alimonti A, Bennett DC, Bischof O, Bishop C, et al. Cellular Senescence: Defining a Path Forward. Cell. 2019 Oct 31;179(4):813–27.

7. Ryan EL, Hollingworth R, Grand RJ. Activation of the DNA Damage Response by RNA Viruses. Biomolecules. 2016 Jan 6;6(1):2.

8. Weitzman MD, Fradet-Turcotte A. Virus DNA Replication and the Host DNA Damage Response. Annu Rev Virol. 2018 Sep 29;5(1):141–64.

9. Tachiwana H, Shimura M, Nakai-Murakami C, Tokunaga K, Takizawa Y, Sata T, et al. HIV-1 Vpr Induces DNA Double-Strand Breaks. Cancer Res. 2006 Jan 15;66(2):627–31.

10. Fregoso OI, Emerman M. Activation of the DNA Damage Response Is a Conserved Function of HIV-1 and HIV-2 Vpr That Is Independent of SLX4 Recruitment. Prasad VR, editor. mBio. 2016 Nov 2;7(5):e01433-16.

11. Lang SM, Weeger M, Stahl-Hennig C, Coulibaly C, Hunsmann G, Müller J, et al. Importance of vpr for infection of rhesus monkeys with simian immunodeficiency virus. J Virol. 1993 Feb;67(2):902–12.

12.   Beaumont T, Van Nuenen A, Broersen S, Blattner WA, Lukashov VV, Schuitemaker H.
Reversal of Human Immunodeficiency Virus Type 1 IIIB to a Neutralization-Resistant
Phenotype in an Accidentally Infected Laboratory Worker with a Progressive Clinical Course.
J Virol. 2001 Mar;75(5):2246–52.

13.   Balliet JW, Kolson DL, Eiger G, Kim FM, McGann KA, Srinivasan A, et al. Distinct
Effects in Primary Macrophages and Lymphocytes of the Human Immunodeficiency Virus
Type 1 Accessory Genes vpr, vpu, and nef: Mutational Analysis of a Primary HIV-1 Isolate.
Virology. 1994 May 1;200(2):623–31.

14.   Mashiba M, Collins DR, Terry VH, Collins KL. Vpr Overcomes Macrophage-Specific
Restriction of HIV-1 Env Expression and Virion Production. Cell Host Microbe. 2014 Dec
10;16(6):722–35.

15.   Miller CM, Akiyama H, Agosto LM, Emery A, Ettinger CR, Swanstrom RI, et al. Virion-
Associated Vpr Alleviates a Postintegration Block to HIV-1 Infection of Dendritic Cells.
Sundquist WI, editor. J Virol. 2017 Jul;91(13):e00051-17.

16.   Paxton W, Connor RI, Landau NR. Incorporation of Vpr into human immunodeficiency
virus type 1 virions: requirement for the p6 region of gag and mutational analysis. J Virol.
1993 Dec;67(12):7229–37.

17.   Roshal M, Kim B, Zhu Y, Nghiem P, Planelles V. Activation of the ATR-mediated DNA
Damage Response by the HIV-1 Viral Protein R. J Biol Chem. 2003 Jul;278(28):25879–86.

18.   Zhang F, Bieniasz PD. HIV-1 Vpr induces cell cycle arrest and enhances viral gene
expression by depleting CCDC137. eLife. 9:e55806.

19.   Lahouassa H, Blondot ML, Chauveau L, Chougui G, Morel M, Leduc M, et al. HIV-1 Vpr
degrades the HLTF DNA translocase in T cells and macrophages. Proc Natl Acad Sci. 2016
May 10;113(19):5311–6.

20.   Schröfelbauer B, Yu Q, Zeitlin SG, Landau NR. Human Immunodeficiency Virus Type 1
Vpr Induces the Degradation of the UNG and SMUG Uracil-DNA Glycosylases. J Virol. 2005
Sep;79(17):10978–87.

21.   Ahn J, Vu T, Novince Z, Guerrero-Santoro J, Rapic-Otrin V, Gronenborn AM. HIV-1 Vpr

Loads Uracil DNA Glycosylase-2 onto DCAF1, a Substrate Recognition Subunit of a Cullin

4A-RING E3 Ubiquitin Ligase for Proteasome-dependent Degradation *. J Biol Chem. 2010

Nov 26;285(48):37333–41.

22.   Lim ES, Fregoso OI, McCoy CO, Matsen FA, Malik HS, Emerman M. The ability of

primate lentiviruses to degrade the monocyte restriction factor SAMHD1 preceded the birth of

the viral accessory protein Vpx. Cell Host Microbe. 2012 Feb 16;11(2):194–204.

23.   Zhou X, DeLucia M, Ahn J. SLX4-SLX1 Protein-independent Down-regulation of

MUS81-EME1 Protein by HIV-1 Viral Protein R (Vpr) *♦. J Biol Chem. 2016 Aug

12;291(33):16936–47.

24.   Yan J, Shun MC, Hao C, Zhang Y, Qian J, Hrecka K, et al. HIV-1 Vpr Reprograms CLR4

[DCAF1] E3 Ubiquitin Ligase to Antagonize Exonuclease 1-Mediated Restriction of HIV-1

Infection. Smithgall TE, editor. mBio. 2018 Nov 7;9(5):e01732-18.

25.   Lv L, Wang Q, Xu Y, Tsao LC, Nakagawa T, Guo H, et al. Vpr Targets TET2 for

Degradation by CRL4VprBP E3 Ligase to Sustain IL-6 Expression and Enhance HIV-1

Replication. Mol Cell. 2018 Jun 7;70(5):961-970.e5.

26.   Romani B, Shaykh Baygloo N, Aghasadeghi MR, Allahbakhshi E. HIV-1 Vpr Protein

Enhances Proteasomal Degradation of MCM10 DNA Replication Factor through the Cul4-

DDB1[VprBP] E3 Ubiquitin Ligase to Induce G2/M Cell Cycle Arrest. J Biol Chem. 2015 Jul

10;290(28):17380–9.

27.   Greenwood EJD, Williamson JC, Sienkiewicz A, Naamati A, Matheson NJ, Lehner PJ.

Promiscuous Targeting of Cellular Proteins by Vpr Drives Systems-Level Proteomic

Remodeling in HIV-1 Infection. Cell Rep. 2019 Apr 30;27(5):1579-1596.e7.

28.   Li D, Lopez A, Sandoval C, Nichols Doyle R, Fregoso OI. HIV Vpr Modulates the Host

DNA Damage Response at Two Independent Steps to Damage DNA and Repress Double-

Strand DNA Break Repair. mBio. 2020 Aug 4;11(4):e00940-20.

29.   Brzostek-Racine S, Gordon C, Van Scoy S, Reich NC. The DNA Damage Response

Induces IFN. J Immunol. 2011 Nov 15;187(10):5336–45.

30. Roux P, Alfieri C, Hrimech M, Cohen EA, Tanner JE. Activation of Transcription Factors
NF-κB and NF-IL-6 by Human Immunodeficiency Virus Type 1 Protein R (Vpr) Induces
Interleukin-8 Expression. J Virol. 2000 May 15;74(10):4658–65.

31. Wu ZH, Shi Y, Tibbetts RS, Miyamoto S. Molecular Linkage Between the Kinase ATM
and NF-κB Signaling in Response to Genotoxic Stimuli. Science. 2006 Feb
24;311(5764):1141–6.

32. Huang TT, Wuerzberger-Davis SM, Wu ZH, Miyamoto S. Sequential modification of
NEMO/IKKgamma by SUMO-1 and ubiquitin mediates NF-kappaB activation by genotoxic
stress. Cell. 2003 Nov 26;115(5):565–76.

33. Miyamoto S. Nuclear initiated NF-κB signaling: NEMO and ATM take center stage. Cell
Res. 2011 Jan;21(1):116–30.

34. Liu R, Lin Y, Jia R, Geng Y, Liang C, Tan J, et al. HIV-1 Vpr stimulates NF-κB and AP-1
signaling by activating TAK1. Retrovirology. 2014 Jun 9;11(1):45.

35. Kogan M, Deshmane S, Sawaya BE, Gracely EJ, Khalili K, Rappaport J. Inhibition of
NF-κB activity by HIV-1 Vpr is dependent on Vpr binding protein. J Cell Physiol. 2013
Apr;228(4):781–90.

36. Belzile JP, Duisit G, Rougeau N, Mercier J, Finzi A, Cohen ÉA. HIV-1 Vpr-Mediated G2
Arrest Involves the DDB1-CUL4AVPRBP E3 Ubiquitin Ligase. PLOS Pathog. 2007 Jul
13;3(7):e85.

37. Le Rouzic E, Belaïdouni N, Estrabaud E, Morel M, Rain JC, Transy C, et al. HIV1 Vpr
arrests the cell cycle by recruiting DCAF1/VprBP, a receptor of the Cul4-DDB1 ubiquitin
ligase. Cell Cycle Georget Tex. 2007 Jan 15;6(2):182–8.

38. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al.
Metascape provides a biologist-oriented resource for the analysis of systems-level datasets.
Nat Commun. 2019 Apr 3;10(1):1523.

39. Russo RC, Garcia CC, Teixeira MM, Amaral FA. The CXCL8/IL-8 chemokine family and
its receptors in inflammatory diseases. Expert Rev Clin Immunol. 2014 May 1;10(5):593–619.

40.   Frazzi R. BIRC3 and BIRC5: multi-faceted inhibitors in cancer. Cell Biosci. 2021 Jan
7;11(1):8.

41.   Mitchell S, Vargas J, Hoffmann A. Signaling via the NFκB system. WIREs Syst Biol Med.
2016 May;8(3):227–41.

42.   Nakai-Murakami C, Shimura M, Kinomoto M, Takizawa Y, Tokunaga K, Taguchi T, et al.
HIV-1 Vpr induces ATM-dependent cellular signal with enhanced homologous recombination.
Oncogene. 2007 Jan;26(4):477–86.

43.   Strickland I, Ghosh S. Use of cell permeable NBD peptides for suppression of
inflammation. Ann Rheum Dis. 2006 Nov;65(Suppl 3):iii75–82.

44.   Liang Z, Liu R, Lin Y, Liang C, Tan J, Qiao W. HIV-1 Vpr protein activates the NF-κB
pathway to promote G2/M cell cycle arrest. Virol Sin. 2015 Dec 1;30(6):441–8.

45.   Strebel K. HIV Accessory Proteins versus Host Restriction Factors. Curr Opin Virol.
2013 Dec;3(6):10.1016/j.coviro.2013.08.004.

46.   Lopez A, Nichols Doyle R, Sandoval C, Nisson K, Yang V, Fregoso OI. Viral Modulation
of the DNA Damage Response and Innate Immunity: Two Sides of the Same Coin. J Mol
Biol. 2022 Mar 30;434(6):167327.

47.   Virgilio MC, Miguel Disbennett W, Chen T, Lubow J, Welch JD, Collins KL. HIV-1 Vpr
combats the PU.1-driven antiviral response in primary human macrophages. BioRxiv Prepr
Serv Biol. 2023 Mar 21;2023.03.21.533528.

48.   Joseph AM, Ladha JS, Mojamdar M, Mitra D. Human immunodeficiency virus-1 Nef
protein interacts with Tat and enhances HIV-1 gene expression. FEBS Lett. 2003 Jul
31;548(1–3):37–42.

49.   Sawaya BE, Khalili K, Gordon J, Taube R, Amini S. Cooperative Interaction between
HIV-1 Regulatory Proteins Tat and Vpr Modulates Transcription of the Viral Genome *. J Biol
Chem. 2000 Nov 10;275(45):35209–14.

50.   Bauby H, Ward CC, Hugh-White R, Swanson CM, Schulz R, Goujon C, et al. HIV-1 Vpr
Induces Widespread Transcriptomic Changes in CD4 [+] T Cells Early Postinfection. Goff SP,
editor. mBio. 2021 Jun 29;12(3):e01369-21.

51. Chougui G, Munir-Matloob S, Matkovic R, Martin MM, Morel M, Lahouassa H, et al. HIV-
2/SIV viral protein X counteracts HUSH repressor complex. Nat Microbiol. 2018
Aug;3(8):891–7.

52. Yurkovetskiy L, Guney MH, Kim K, Goh SL, McCauley S, Dauphin A, et al. Primate
immunodeficiency virus proteins Vpx and Vpr counteract transcriptional repression of
proviruses by the HUSH complex. Nat Microbiol. 2018 Dec;3(12):1354–61.

53. Cicchini L, Westrich JA, Xu T, Vermeer DW, Berger JN, Clambey ET, et al. Suppression
of Antitumor Immune Responses by Human Papillomavirus through Epigenetic
Downregulation of CXCL14. Imperiale MJ, editor. mBio. 2016 Jul 6;7(3):e00270-16.

54. Romero N, Favoreel HW. Pseudorabies virus infection triggers NF-κB activation via the
DNA damage response, but actively inhibits NFkB-dependent gene expression. J Virol. 2021
Oct 6;JVI.01666-21.

55. Harman AN, Nasr N, Feetham A, Galoyan A, Alshehri AA, Rambukwelle D, et al. HIV
Blocks Interferon Induction in Human Dendritic Cells and Macrophages by Dysregulation of
TBK1. Silvestri G, editor. J Virol. 2015 Jul;89(13):6575–84.

56. Gummuluru S, Emerman M. Cell Cycle- and Vpr-Mediated Regulation of Human
Immunodeficiency Virus Type 1 Expression in Primary and Transformed T-Cell Lines. J Virol.
1999 Jul;73(7):5422–30.

57. Goh WC, Rogel ME, Kinsey CM, Michael SF, Fultz PN, Nowak MA, et al. HIV-1 Vpr
increases viral expression by manipulation of the cell cycle: A mechanism for selection of Vpr
in vivo. Nat Med. 1998 Jan;4(1):65–71.

58. Nodder SB, Gummuluru S. Illuminating the Role of Vpr in HIV Infection of Myeloid Cells.
Front Immunol [Internet]. 2019 [cited 2023 Apr 3];10. Available from:
https://www.frontiersin.org/articles/10.3389/fimmu.2019.01606

59. Varin A, Decrion AZ, Sabbah E, Quivy V, Sire J, Van Lint C, et al. Synthetic Vpr Protein
Activates Activator Protein-1, c-Jun N-terminal Kinase, and NF-κB and Stimulates HIV-1
Transcription in Promonocytic Cells and Primary Macrophages*. J Biol Chem. 2005 Dec
30;280(52):42557–67.

60.   Fabryova H, Kao S, Sukegawa S, Miyagi E, Taylor L, Ferhadian D, et al. HIV-1 Vpr
Induces Degradation of Gelsolin, a Myeloid Cell-Specific Host Factor That Reduces Viral
Infectivity by Inhibiting the Expression and Packaging of the HIV-1 Env Glycoprotein. Goff
SP, editor. mBio. 2023 Feb 28;14(1):e02973-22.

61.   Bruce JW, Bracken M, Evans E, Sherer N, Ahlquist P. ZBTB2 represses HIV-1
transcription and is regulated by HIV-1 Vpr and cellular DNA damage responses. PLOS
Pathog. 2021 Feb 26;17(2):e1009364.

62.   Yamashita M, Emerman M. Capsid Is a Dominant Determinant of Retrovirus Infectivity in
Nondividing Cells. J Virol. 2004 Jun;78(11):5670–8.

63.   Choi VW, Asokan A, Haberman RA, Samulski RJ. Production of Recombinant Adeno-
Associated Viral Vectors for In Vitro and In Vivo Use. Curr Protoc Mol Biol [Internet]. 2007
Apr [cited 2023 May 3];78(1). Available from:
https://onlinelibrary.wiley.com/doi/10.1002/0471142727.mb1625s78

64.   Bekker-Jensen S, Lukas C, Melander F, Bartek J, Lukas J. Dynamic assembly and
sustained retention of 53BP1 at the sites of DNA damage are controlled by Mdc1/NFBD1. J
Cell Biol. 2005 Jul 18;170(2):201–11.

65.   Lukas C, Melander F, Stucki M, Falck J, Bekker-Jensen S, Goldberg M, et al. Mdc1
couples DNA double-strand break recognition by Nbs1 with its H2AX-dependent chromatin
retention. EMBO J. 2004 Jul 7;23(13):2674–83.

795
796
Figure 1: HIV-1 Vpr induces DNA damage and alters cellular transcription independent of
cell cycle arrest and CRL4A^{DCAF1} engagement. (A) Comet assay of U2OS cells infected with
rAAV expressing 3xFLAG-tagged Vpr WT, H71R, Q65R, untreated (negative control), or 50uM
etoposide (positive control). Percent tail DNA was quantified at 24 hours post infection (hpi) using
the OpenComet plug-in for the ImageJ software. Each circle represents one cell. N=3, one

representative experiment shown. (B) Representative immunofluorescence images of U2OS cells
infected under the same conditions as Fig. 1A; γH2A.x (magenta) and nuclei stained with DAPI
(blue). Images were taken at 63x magnification. N=3, one representative experiment shown. EV,
empty vector. (C) RNA-seq of U2OS cells at 36 hpi. Heat map displays Log_2 fold changes of
upregulated genes in red and downregulated genes in blue. Each column represents a biological
replicate, n=3 (D) Venn diagram of the top 100 upregulated and top 100 downregulated genes
among Vpr WT, S79A, H71R, and Q65R. E) Dot plot of the 68 conserved differentially expressed
genes among Vpr WT, S79A, and H71R. BIRC3 and CXCL8 are highlighted as the two most
upregulated genes (Log_2 fold change) under all conditions. (F) Gene ontology and (G) TRRUST
analysis of the 30 upregulated genes performed with Metascape software. Asterisk indicate
statistical significance compared to untreated control, as determined by one-way ANOVA test
(NS, nonsignificant; *** $P < 0.0004$, **** $P < 0.0001$). Related to Figures S1 and S2.

Figure 2: HIV-1 Vpr activates RelA and promotes NF-κB transcription independent of cell cycle arrest and CRL4A^{DCAF1} engagement. (A) qRT-PCR validation of upregulated NF-κB target genes BIRC3 and CXCL8. Normalized expression (ΔΔCt) was calculated by normalizing to GAPDH followed by calculating fold change to untreated or empty vector treated cells. Cells treated under the same conditions as Fig. 1C. n=3, one representative experiment shown. (B) Representative immunofluorescence images of U2OS cells infected under the same conditions as Fig. 1A; RelA (red) and nuclei stained with DAPI (blue). Images were taken at 63x magnification. n=3, one representative experiment shown. Asterisk indicate statistical significance compared to untreated control, as determined by one-way ANOVA test (NS, nonsignificant; *** P< 0.0003, **** P< 0.0001).

826
827
Figure 3: Vpr-induced DNA damage activates ATM signaling independent of cell cycle
arrest and CRL4A^DCAF1 engagement. (A) Live-cell imaging of U2OS cells stably expressing
NBS1-GFP or 53PB1-GFP. Cells were infected under the same conditions as Fig. 1A and
representative images were taken at 33 hpi at 63x magnification. n=3, one representative
experiment shown. (B) Representative images of co-localization of DNA damage sensors
MRE11 (magenta) and NBS1 (yellow) in U2OS NBS1-GFP or (C) DNA damage transducers
yH2A.x (magenta) and 53PB1 (yellow) in U2OS 53PB1-GFP cells infected under the same
conditions as Fig. 1A. Representative images taken at 24 hpi at 63x magnification. n=3, one
representative experiment shown. Related to Figure S3.

Figure 4: Vpr upregulates NF-κB transcription dependent on NEMO, yet independent of cell cycle arrest and CRL4A^{DCAF1} engagement. qRT-PCR of BIRC3 and CXCL8 in the presence or absence of Nemo-Binding Inhibitor peptide (NBD) at 36 hpi. U2OS cells were pretreated with 100uM of NBD for 2 hours prior to infection with rAAV expressing 3xFLAG-tagged Vpr WT, H71R, Q65R, empty vector (negative control). Normalized expression to GAPDH. Asterisk indicate statistical significance compared to untreated control, as determined by one-way ANOVA test (NS, nonsignificant; **** P< 0.0001).

846
847
Figure 5: Vpr-induced DNA damage activates ATM-NEMO signaling and NF-κB
transcription in primary MDMs early during infection. (A) Experimental schematic depicting
the isolation and differentiation of primary human monocyte-derived macrophages (MDMs) from
peripheral blood mononuclear cells (PBMCs) followed by delivery of Vpr via Virus-like Particles
(VLPs) or HIV-1 infection. Infected MDMs were assayed for induction of DNA damage, activation
of DDR signaling, and the upregulation of NF-κB target genes in the context of NEMO inhibitor.
(B) Comet assay of MDMs treated with VLPs packaging 3xFLAG-tagged Vpr WT, H71R, Q65R,
empty VLPs (negative control), or 25uM Mitomycin C (positive control) for 8 hours. Comet assay
analysis was performed as in Fig. 1A. n=2, one representative experiment shown. (C)
Representative immunofluorescence images of MDMs infected with VLPs as in Fig. 5B and
treated or not with Nemo-Binding Inhibitor peptide (NBD); γH2A.x (magenta) and nuclei stained
with DAPI (blue). Images were taken at 100x magnification. n=3, one representative experiment
shown. (D) qRT-PCR for CXCL8 of MDMs treated with NBD as in Fig. 5C with TNFα as positive
control. n=3, one representative experiment shown. (E) qRT-PCR for CXCL8 of MDMs infected
with HIV-1 ΔEnv for 8, 16, 24, and 48 hpi from four separate donors. Normalized expression to
GAPDH. Asterisk indicate statistical significance compared to untreated control, as determined
by one-way ANOVA test (NS, nonsignificant; * P<0.03, ** P < 0.005, **** P< 0.0001). Related to
Figure S4.

Figure 6: Model showing Vpr-induced DNA damage activates ATM and NEMO signaling resulting in RelA nuclear translocation and NF-κB transcriptional upregulation. 1. Vpr induces double-strand DNA breaks independent of cell cycle arrest and CRL4A^{DCAF1} complex engagement. 2. Vpr-induced DNA damage activates markers of ATM signaling including MRE11, NBS1, γH2A.x, and 53PB1. 3. ATM-DDR activation results in NEMO relieving RelA inhibition from inhibitor of NF-κB (IκBα and IκBβ). 4. RelA translocates into the nucleus. 5. RelA binds to NF-κB promoters and activates transcription (txn) of NF-κB target genes such as BIRC3 and CXCL8. (Schematic made on Biorender).

Figure S1: Knockdown of DCAF1 does not abrogate the ability of Vpr to induce DNA damage, activate DDR signaling, or upregulate NF-κB target genes. (A) Schematic representation of Vpr mutants used in these studies to decouple some Vpr functions. Vpr S79A and Vpr H71R do not induce cell cycle arrest, and H71R also does not bind DCAF1 of the CRL4A^{DCAF1} complex to degrade host proteins or remodel the proteome. Vpr Q65R (not shown) is a largely non-functional negative control. (B) Western blot of endogenous DCAF1 U2OS cells to test stable knockdown with shRNA. Control (ctl) shRNA and two DCAF1-specific shRNAs (shRNA1 and shRNA2) were used. Actin is used as a loading control. (C) Comet assay of DCAF1 knock-down cells infected with rAAV and analyzed as in Fig. 1A. n=1. (D) Representative immunofluorescence images of DCAF1 knock-down U2OS cells infected under the same conditions as Fig. 1A.; γH2A.x (magenta) and nuclei with DAPI (blue). Images were taken at 63x magnification. n=1, one representative image shown. (E) qRT-PCR for CXCL8 and BIRC3 in DCAF1 knock-down U2OS cells infected under the same conditions as Fig. 1A. n=1. Asterisk indicate statistical significance compared to untreated control, as determined by t-test (NS, nonsignificant; * P< 0.02, ** P<0.004, *** P<0.0003, **** P<0.0001). EV, empty vector negative control; Etop., etoposide positive control. Related to Figures 1 and 2.

892
893
894 **Figure S2: Vpr alters cellular transcription through diverse mechanisms that are either**
895 **dependent or independent of cell cycle arrest and CRL4A^DCAF1 engagement.** (A) Gene
896 ontology using Metascape displaying TRRUST analysis for upregulated genes. (B) Principal
897 component analysis (PCA) of the RNA-seq data showing the relatedness of the three biological
898 replicates compared to empty vector or untreated cells. Related to Figure 1.

899
900
Figure S3: Vpr-induced DNA damage activates markers of ATM signaling. (A) Live-cell
imaging taken on the IncuCyte of U2OS NBS1-GFP or 53PB1-GFP cells infected as in Fig. 1A
for 48 hrs. n=2, one representative experiment shown. (B & C) Quantification of DNA damage
induced in the live-cell imaging experiment in Fig. 3A at 8, 26, 33, and 48hpi. Foci per cell and
foci size were quantified using ImageJ software. Images were taken at 63x magnification using
the LSM 900. n=3, one representative experiment shown. EV, empty vector; AU, arbitrary units.
Related to Figure 3.

Figure S4: Infection of primary MDMs with VLPs packaging Vpr protein or HIV-1 infection
upregulate NF-κB transcription dependent on NEMO. (A) Flow cytometry plots displaying
differentiation of PBMC-derived monocytes to macrophages. Primary MDMs are CD14+, CD45+
and CD16+. n=6, one representative experiment shown. (B) Western blot of 3X FLAG-Vpr WT
and mutants packaged in Virus-like particles (VLPs). (C) qRT-PCR for BIRC3 of MDMs treated
with NBD as in Fig. 5C with TNFα as positive control. n=3, one representative experiment
shown. (D) Histogram quantifying percent of infected MDMs with HIV-1 ΔEnv at 48hpi. n=6, one
representative experiment shown. (E) qRT-PCR for unspliced HIV-1 RNA at 8, 16, 24 and
48hpi. n=6, one representative experiment shown. (F) qRT-PCR for TNFα of MDMs infected
with HIV-1 ΔEnv at 8, 16, 24 and 48hpi. n=6, one representative experiment shown. (G & H)
qRT-PCR for BIRC3 or CXCL8 of MDMs infected with HIV-1 ΔEnv at 8, 16, 24 and 48hpi in
separate donors. n=6, (G) is all 6 donors, (H) is donors 5 and 6 in addition to those shown in
Fig. 5E. Asterisk indicate statistical significance compared to untreated control, as determined
by one-way ANOVA test (NS, nonsignificant; **** P<0.0001). Related to Figure 5.