Aging drives cerebrovascular network remodeling and functional changes in the mouse brain

Hannah C. Bennett[1,6], Qingguang Zhang[2,6], Yuan-ting Wu[1,3], Uree Chon[1,4], Hyun-Jae Pi[1], Patrick J. Drew[2,5], Yongsoo Kim[1,2,7]

[1] Department of Neural and Behavioral Sciences, The Pennsylvania State University, Hershey, PA, 17033, USA
[2] Center for Neural Engineering, Department of Engineering Science and Mechanics, The Pennsylvania State University, University Park, PA, 16802, USA
[5] Biomedical Engineering, Biology, and Neurosurgery, The Pennsylvania State University, University Park, PA, 16802, USA
[6] Equal contribution
[7] Lead contact. Email: yuk17@psu.edu

Current address:
[3]Department of Neurosurgery, Department of Computational Biomedicine, Cedars-Sinai Medical Center, CA 90048
[4] Stanford University, Stanford, CA 94305

Abstract

Aging is the largest risk factor for neurodegenerative disorders, and commonly associated with compromised cerebrovasculature and pericytes. However, we do not know how normal aging differentially impacts the vascular structure and function in different brain areas. Here we utilize mesoscale microscopy methods (serial two-photon tomography and light sheet microscopy) and *in vivo* imaging (wide field optical spectroscopy and two-photon imaging) to determine detailed changes in aged cerebrovascular networks. Whole-brain vascular tracing showed an overall ~10% decrease in vascular length and branching density, and light sheet imaging with 3D immunolabeling revealed increased arteriole tortuosity in aged brains. Vasculature and pericyte densities showed significant reductions in the deep cortical layers, hippocampal network, and basal forebrain areas. Moreover, *in vivo* imaging in awake mice identified delays in neurovascular coupling and disrupted blood oxygenation. Collectively, we uncover regional vulnerabilities of cerebrovascular network and physiological changes that can mediate cognitive decline in normal aging.

Key Words:
Aging, pericyte, cerebrovasculature, brain, high-resolution mapping, serial two-photon tomography, light sheet fluorescence microscopy, neurovascular coupling

Highlight
- Brain-wide mapping of vasculature and pericyte changes with normal aging
- Simplified vascular network with tortuous vessels in aged brains
- Vascular rarefication in the deep cortical layers, hippocampus, and the basal forebrain
- Slowed hemodynamic response in aged animals

Introduction

Aging is the primary risk factor for the development of various neurodegenerative diseases. Notably, aging is associated with decreased cerebral blood flow and general vascular impairment [1]. A common denominator in diseases that increases the risk of dementia, such as stroke, atherosclerosis, and diabetes mellitus, is vascular perturbation and dysfunction of neurovascular coupling [2-9]. All of the disease processes mentioned above increase the risk of developing vascular dementia, which is the second leading cause of cognitive impairment in the United States. Impairment in the cerebrovascular network can have a significant impact on energy supply and metabolic waste removal processes, which can result in neuronal death linked with various clinical symptoms [10-12]. Thus, understanding the anatomical and functional changes in the brain vasculature upon normal aging is a critical first step in understanding neurodegenerative disorders.

The vessels of the cerebrovascular network are composed of endothelial cells linked by tight junctions. These blood vessels are surrounded by mural cells, such as vascular smooth muscle cells and pericytes, which wrap around vessels of the vascular tree and contribute to blood flow regulation [13]. Pericytes are essential for maintaining the blood brain barrier and play important roles in waste removal and capillary blood flow regulation [14,15]. The importance of these vascular cell types is becoming increasingly recognized in the context of brain disorders, particularly in the case of neurodegenerative diseases. Previous studies showed that aging with cognitive impairment is associated with vascular pathologies including increased arterial tortuosity, rarefaction of the vascular tree, and impairment of pericyte dynamics [3,5,14,16-18]. In addition to anatomical changes, advanced aging is associated with reduced cerebral blood flow (CBF), increased CBF pulsatility, and stiffening of the major arteries [19-21]. It is becoming increasingly recognized that disruption to the brain's vasculature may precede the neuronal damage associated with neurodegenerative disease and other types of dementia [22], implying that vascular dysfunction may play a causative role in neurodegeneration. Despite its significance, it remains unclear how the cerebrovascular network, including mural cell types, across different brain regions undergoes structural and functional changes during the aging process. Prior work has primarily focused on single brain regions without accounting for brain-wide changes in the cerebrovascular network, largely due to the complexities of visualizing and analyzing large 3D brain volumes.

Recent advances in 3D whole brain imaging methods make it possible to quantitatively examine detailed cerebrovascular networks in the entire mouse brain [23-28]. We previously showed that regional differences in pericyte density and cerebrovascular structure strongly correlate with the number of parvalbumin-expressing neuron populations in the cortex of young adult mice [28]. Here, leveraging high-resolution 3D mapping methods (light sheet and serial two-photon microscopy), we ask whether there are regional vulnerabilities within the cerebrovasculature and mural cell types upon aging. We found selective reduction of vascular length and pericyte density in deep cortical layers, as well as the basal forebrain areas where cholinergic neurons with large cell bodies reside. Advanced age causes vascular remodeling with increased arterial tortuosity in the isocortex and reduces capillary pericyte density in the entorhinal cortex. In addition to anatomical changes, *in vivo* imaging (two-photon and wide field optical intrinsic signal imaging) of the vasculature in awake aged mice indicates slowed hemodynamic responses

92   to sensory stimulation and voluntary locomotion. Collectively, our results demonstrate
93   significant cerebrovascular network changes, linked to regional vulnerabilities and reduced
94   hemodynamic responsiveness in aging.
95
96
97

Results

Early aging in the mouse brain shows overall decreased vascular length density and branching density, but increased vascular radii

To determine structural changes of the cerebrovasculature upon normal aging, we applied our cerebrovascular mapping pipelines in 18-month-old (aged) mice in order to compare 2-month-old (young adult) mice [28] (Figure 1). We labeled the brain vasculature by cardiac perfusion of fluorescein isothiocyanate (FITC)-conjugated albumin gel [26,28–30]. Then, we utilized serial two-photon tomography (STPT) imaging to image the whole mouse brain at 1x1x5 µm resolution (*x,y,z;* media-lateral, dorsal-ventral, rostral-caudal) followed by computational analysis for vasculature tracing and quantification [28,31]. All signals were registered to the Allen Common Coordinate Framework (AllenCCF) as a reference brain [32] (Figure 1).

To identify potential regional vascular vulnerabilities, we first examined the overall changes of the cerebrovasculature across the whole mouse brain, comparing 18-month-old mice to 2-month-old mice (Figure 2A). Total vessel length in most regions remained similar between 18- and 2-month-old mice (Figure 2B) but overall brain volume increased about 6% (Figure 2C). The brain volume increase was also seen with *in vivo* longitudinal MRI [33], indicating our result is not an artifact from fixation or imaging. As a result, overall vascular length density across different brain regions decreased by 5 – 10% in the aged brain (Figure 2D). In addition, we found an approximate 10-20% decrease in branching density across most brain regions (Figure 2E). In contrast, the average radius of 18-month-old mouse brain vasculature is increased by about 5 – 10% compared to 2-month-old mice, suggesting reduced basal constrictive tone (Figures 2F). Notably, we found significant changes in brain regions related to memory processing and storage (e.g., Ammon's horn; CA, lateral entorhinal cortex; ENTl, Anteromedial nucleus; AM), appetitive behavior (e.g., medial preoptic area; MPO, ventral premammillary area; PMv), body physiology and sleep (e.g., lateral preoptic area; LPO, anterior hypothalamic area; AHA), attention (e.g., substantia innominata; SI, medial septum; MS as basal forebrain areas), sensory processing and integration (e.g., zona incerta; ZI, Dorsal lateral geniculate nucleus; LGd), and executive function (e.g., medial group of the dorsal thalamus; MED) (Figure 2D-F, highlighted with magenta boxes; Supplementary Data 1).

Next, we examined isocortical areas for aging-related vascular changes. Surprisingly, isocortical areas showed no significant changes, with mostly less than 10% decrease in length and branching density, and about 5% increase in average vessel radius (Figure 2G-I). To examine vascular changes in the isocortex more intuitively, we utilized our previously developed isocortical flatmap with five distinct cortical domains marked by different colors (Figure 2J) [28]. We found a significant reduction in vessel length density only in layer 6 of aged brains compared to young brains (Figure 2K-N). Our result corroborates a previous finding showing selective vulnerability of deep cortical layers [34]. Together, these findings indicate that the vasculature of the isocortex is relatively resilient to aging, and the earliest evidence of age-related vascular degeneration occurs in layer 6.

Pericyte density in aged brains showed significant decrease in basal forebrain regions and the deep cortical layer.

Pericytes are a mural cell type that plays a key role in the regulation of the capillary network blood flow and diameter, and are known to be vulnerable in aging [15,35,36]. Our results show increased vascular radius in aged brains, which raises the possibility of dysfunction in pericytes

in the maintenance of vascular diameter. To quantitatively determine changes of pericytes, we
compared capillary pericyte densities in 2-month-old and 18-month-old PDGFRβ-Cre;Ai14 mice
[37,38], where tdTomato is expressed in pericytes and other mural cells. We used STPT imaging of
PDGFRβ-Cre;Ai14 mice with previously developed computational analyses to image, identify,
and quantify changes of capillary pericytes upon aging across the whole mouse brain [28,31] (Figure
1 bottom and Figure 3A).
Overall, pericyte density in the aged brain remained within 10% of that in young brains in
most areas, including many cortical and thalamic subregions (Figure 3B; Supplementary Data 2).
However, a significant reduction of pericyte density was found in basal forebrain areas (e.g., the
substantia innominata; SI, magnocellular nucleus; MA) and the closely related anterior amygdala
area (AAA) (Figures 3B; red boxed, C-D) [39]. Considering the basal forebrain contains cortical-
projecting cholinergic neurons, the observed significant reduction in pericyte and vascular
densities reflects the selective and early vulnerability of the basal forebrain during aging. These
results could potentially provide a link between known vascular impairment and dysfunction of
cholinergic neurons in neurodegenerative diseases such as Alzheimer's disease [40,41].
Given that we saw few vascular changes with aging in the isocortex (Figure 2), we
investigated whether this resilience extends to pericyte density. We compared 2-month- and 18-
month-old mice capillary pericyte densities by brain region using our isocortical flatmap (Figure
3E). The capillary pericyte density in aged mice overall remained similar to, or even slightly
increased as compared with young adult mice (Figure 3E-F), particularly in motor sensory
regions (white and gray arrowheads in Figure 3E). Due to reduced vessel length density, the
overall pericyte cell body coverage (capillary pericyte number per vascular length) is increased
by about 10% in sensorimotor areas in aged mice compared to young adult mice (Figure 3G). In
contrast to sensorimotor areas, relatively little or even reduced pericyte coverage was observed in
medial prefrontal areas, suggesting selective reduction of pericytes with aging in an association
area known to be related to higher cognitive functions.
We then asked whether there are selective changes across cortical layers. We noted that
the deep cortical layer (L6) showed selective reduction of pericyte density in the infralimbic
cortex, while the superficial layers (2/3 and 4) in the whisker representation of the primary
somatosensory cortex ('barrel field') showed a significant increase in the 18-month-old brain
compared to the 2-month-old brain (Figure 3H-J). When layer specific density from all
isocortical areas was combined, pericyte density was significantly reduced in deep layer 6b in the
aged brain, while layers 2/3 and 4 showed significant increases (Figures 3K). Considering layer
6b plays a role in brain state modulation [42] and the protective role of pericytes in vascular
integrity, significant reduction of the pericytes could make layer 6b and nearby white matter
tracks more vulnerable upon aging.
180
Artery specific labeling shows striking vascular remodeling in penetrating cortical
arterioles of aged brains.
Previous studies have identified age-related changes in arteries and arterioles in both rodents and
humans [43,44]. To investigate potential remodeling in main arteries and penetrating cortical
arterioles, we utilized tissue clearing, 3D immunolabeling, and high-resolution light sheet
fluorescence microscopy (LSFM) imaging (Figure 4A) (Methods for more details). We labeled
arteries with smooth muscle actin (Acta 2) and transgelin (Sm22) antibodies, pan-vasculature
with lectin, and pericytes with CD13 and PDGFRβ antibodies in the same brain. This approach

enabled us to examine different vascular compartments and mural cell types in the same intact
3D brain (Figure 4B-H).
We applied the method to 2-month-old and 24-month-old (late aging) C57BL/6 mice to
test whether late aging shows structural remodeling of different vascular compartments and
progression of capillary pericyte density reduction. We first focused our analysis on the middle
cerebral artery and anterior communicating artery branches contributing to the anterior
circulation of the circle of Willis, which is responsible for supplying the majority of cerebral
blood flow (Figure 5A). We quantified the average radius of each artery. We did not find
significant differences in young and aged mice, nor differences between sexes (Figure 5B-C),
suggesting that aging does not impact the diameters of the main feed arteries to the brain.
Next, we examined the number of cortical penetrating arterioles, which are bottlenecks in
the supply of blood to the brain [45,46]. There were no significant changes in cortical arteriole
numbers (both total arterioles and arterioles that extend into layer 6/corpus callosum) in aged
brains compared to the young adult mice (Figure 5D). However, we observed highly tortuous
(twisted) vessels across the entire cortex (Figure 5E; highlighted with red arrowheads), which is
consistent with prior observations in aged animals and humans [47,48]. Further analysis revealed
that aged animals demonstrate increased arteriole tortuosity, as measured by the arc chord ratio
(Figure 5F) (see Methods for more details). The number of branching points per arteriole
remains similar across the age group (Figure 5G). This increased tortuosity of penetrating
arterioles will result in increased blood flow resistance, leading to slowed blood flow with
decreased oxygen and nutrient delivery if there is no increase in blood pressure. This decreased
flow could make the deep cortical layers and nearby white matter tracks vulnerable during aging.
211
Advanced aging is associated with selective loss of capillary pericytes in cortical layer 6 and
the entorhinal cortex.
Different pericyte subtypes associate with different vascular branches (Figure 5H) [13]. Aging has
been shown to impair specific pericyte subtypes, such as first order (ensheathing) pericytes at the
junction between arterioles and microvessels [49]. To examine how different pericyte subtypes are
differentially impacted in advanced aging, we used a combination of artery, pan-vascular, and
mural cell immunolabeling, in order to distinguish pericyte subtypes at different vascular zones
with submicron resolution ($0.4 \times 0.4 \times 1 \ \mu m^3$) using LSFM imaging (Figure 5I). We successfully
visualized individual pericytes and their subtypes, including capillary pericytes, both mesh and
thin-strand morphologies, and ensheathing pericytes, which are located along pre-capillary
arterioles and express smooth muscle markers such as Acta2 [34] (Figure 5H-I). Examples of
different pericyte subtypes (i.e., ensheathing, mesh, and thin strand) are labeled in Figure 5I with
cyan, yellow, and purple arrows in each panel, respectively. By following individual vasculature,
each pericyte type was manually counted in a region of interest.
Consistent with our STPT data, we did not observe any significant changes in pericyte
subtype density within the primary somatosensory cortex except a significant reduction of
capillary pericyte density in layer 6 (Figure 5J-L). This result further confirmed that pericyte
density remains largely unchanged in cortical areas, including contractile ensheathing pericytes.
We further examined the entorhinal cortex, since this region is important for memory and is
known to be very sensitive to age-related diseases [4,18,35,50–52]. While this region did not show any
statistically significant decreases at 18 months of age (early aging) in STPT pericyte mapping
(Figure 3B), we found that both mesh and thin-strand pericytes, but not ensheathing pericytes,

showed significant reductions in 24-month-old (late aging) mice (Figure 5M-N). This suggests
that capillary pericytes are at higher risk of cellular density loss, particularly in advanced age.
236
In vivo imaging to examine hemodynamic changes in aged brains.

In addition to structural changes, the cerebrovasculature may undergo functional changes in
neurovascular coupling with aging. Thus, we investigated how normal aging impacts brain
hemodynamics during rest and in response to voluntary locomotion and sensory stimulation in
awake behaving mice, using wide field intrinsic optical imaging of spectroscopy (IOS) [53] and
two-photon laser scanning microscopy (2PLSM) [54–56](Figure 6 and 7). All experiments were
performed in awake mice that were head-fixed on a spherical treadmill for voluntary locomotion
[53–57]. Imaging was performed through polished and reenforced thin-skull windows (PoRTS) to
minimize the disruption of the intracranial environment [58]. We utilized two different models,
voluntary locomotion [53,55] and whisker stimulation [54,56,59], to quantify the evoked responses. We
focused our analysis on two functionally distinct cortical regions, the forelimb/hindlimb
representation of the somatosensory cortex (FL/HL) and a frontal cortical region (FC) including
the anterior lateral motor cortex (ALM). We targeted ALM because it is involved in motor
planning and performs "higher-order" cognitive functions in mice, which makes it analogous to
the human prefrontal cortex. We performed these measurements in mice of ages of 2-4 month,
18 month, and 24 month.
253
Neurovascular coupling shows slower response time in aged brains.

We first assessed the spatial extent of cortical hemodynamic responses and their relationship to
voluntary locomotion, using intrinsic optical signal imaging of spectroscopy [53]. Taking
advantage of differences in the optical absorption spectra of oxyhemoglobin (HbO) and
deoxyhemoglobin (HbR) [60,61], we collected reflectance images during rapid alternating green
(530 nm) and blue (470 nm) illumination (Figure 6A and H). When the brain is illuminated with
light of different wavelengths, increases in total hemoglobin concentration (ΔHbT) in turn report
dilations of arteries, capillaries, and veins, which correspond with increases in cerebral blood
volume (CBV). The ΔHbT observed with IOS closely tracks measurements of vessel diameter
made with two-photon microscopy [62]. The consistency of microscopic measurements of vessel
diameter, combined with its very high signal-to-noise ratio [54], and spatial resolution (less than
200 µm) [63], makes IOS suitable for detecting hemodynamic responses to locomotion. While
neurally-evoked dilations initiate in the deeper layers of the cortex, the dilations propagate up the
vascular tree to the surface arteries [64–67], where they can be easily detected with IOS.
We quantified how locomotion affected CBV in two complementary ways. We calculated
the locomotion-triggered average, generated by aligning the IOS or vessel diameter signals to the
onset or offset of locomotion using only locomotion events ≥ 5 seconds in duration (Figure 6B
and C). Using changes in ΔHbT as an indicator of CBV, we observed region-specific changes in
ΔHbT during locomotion (Figure 6B and C). In young adult mice (2-4 months old), there was a
pronounced increase in the ΔHbT (corresponding to an increase in CBV) in the
forelimb/hindlimb representation of the somatosensory cortex (FL/HL), while in the frontal
cortex (FC) there was no change, or even a slight decrease in ΔHbT (n = 7 mice) (Figure 6B and
C), consistent with previous reports [53,57,68]. This pattern was not affected by aging, as we
observed similar results in 18-month-old (n = 5 mice) and 24-month-old (n = 11 mice) (Figure
6B and C).

We also calculated the hemodynamic response function (HRF) [54,69], which is the linear kernel relating locomotion events to observed changes in CBV and vessel diameter (Figure 6D and F; see Methods), using all locomotion events. Hemodynamic response functions are used in all of fMRI analyses to extrapolate neuronal activity from a stimulus or a task from hemodynamic signals, and take into account the slower responses of the vasculature relative to neurons [70]. Using the HRFs to quantify the net CBV, we obtained the same conclusions as derived from the locomotion-triggered average, i.e., the net increase in cerebral blood volume does not change during aging (Figure 6E and G, left) in either FC or FL/HL (2-month: 0.53 ± 0.18 µM; 18-month: 0.57 ± 0.06 µM; 24-month: 0.53 ± 0.12 µM). In addition to the amplitude of the hyperemic response evoked by locomotion, HRFs also provide us information regarding the temporal dynamics of CBV responses. We found that the onset time (Figure 6E, middle, 2-month: 0.95 ± 0.15 s; 18-month: 0.95 ± 0.14 s; 24-month: 1.17 ± 0.10 s) and duration (Figure 6E, right, 2-month: 1.11 ± 0.12 s; 18-month: 1.24 ± 0.24 s; 24-month: 1.38 ± 0.24 s) of locomotion evoked hyperemic response is significantly lengthened with aged brains, especially in the late aging groups (22-26 month old) in the FL/HL (2-4 month vs 22-26 month: time to peak, unpaired t-test, $t(14) = 3.54$, $p = 0.0033$; FWHM, Wilcoxon rank sum test, $p = 0.0311$). To further validate the results from HRFs, we quantified the responses of ΔHbT in response to a brief whisker stimulation (100 ms duration) (Figure 6H). We observed that in response to contralateral whisker stimulation, the onset time and duration of ΔHbT response are significantly lengthened in the 24-month-old late aging groups in both FC and FL/HL (Figure 6I-L).

In addition to the mesoscopic level measurements using IOS, we further compared whether hemodynamics was different between age groups at individual vessel level in FL/HL, in terms of pial arterial diameter change in response to locomotion, using *in vivo* 2PLSM (Figure 6M). The locomotion-evoked arterial diameter change (Figure 6N), as well as the HRF of arterial diameter change (Figure 6O) showed a similar spatial pattern of responses as the CBV measured using IOS, i.e., a trend of delayed response during aging.

Finally, to determine whether vascular dilation capacity was intact in aged mice, we measured the mesoscopic brain hemodynamic responses using IOS (Figure 6P) and microscopic vessel diameter response to isoflurane, a potent vasodilator, using 2PLSM. In FL/HL, we observed an increase of ΔHbT (2-month: 176.8 ± 29.5 µM, 4 mice; 18-month: 146.7 ± 22.2 µM, 4 mice; 24-month: 133.8 ± 23.5 µM, 5 mice) and arteriole diameter (2-month: 65.5 ± 19.2%, 4 mice; 18-month: 51.7 ± 10.8%, 5 mice; 24-month: 54.6 ± 28.9%, 8 mice; data not shown) when animals inhale isoflurane. In FC, we observed an increase of ΔHbT (2-month: 106.6 ± 28.5 µM, 4 mice; 18-month: 85.4 ± 23.0 µM, 4 mice; 24-month: 93.6 ± 18.2 µM, 5 mice). The extent of vasodilation observed between young and old mice was not different when animals were transitioned from air to 5% isoflurane, suggesting the dilation capacity remains similar across different age groups.

Oxygenation carrying capacity is decreased in aged mice.
One of the important functions of increased blood flow/volume is to deliver oxygen to the brain. Using the cerebral oxygenation index (HbO-HbR) [53,71], a spectroscopic measurement of hemoglobin oxygenation, we saw an increase in oxygenation during locomotion and whisker stimulation in both FC and FL/HL areas in young mice. The oxygen increase in response to locomotion (Figure 7A) and whisker stimulation (Figure 7B) did not significantly differ across age groups. As vasodilation is one of the determining factors controlling brain oxygenation [53], we quantified the relationship between locomotion evoked responses of ΔHbT and ΔHbO-HbR

using linear regression. The slope and intercept of the fitting decreased with the healthy aging
process (2-month: $y = 0.8667x + 35.43$; 18-month: $y = 0.5209x + 27.37$; 24-month: $0.5885x +$
30.42; Figure 7C), which suggests that oxygen carrying capacity for the red blood cells decreases
during aging, and that the aging brain has lower baseline oxygenation, respectively.
To determine whether the oxygen exchange and oxygen delivery capacity were intact in
aged mice, we measured the brain tissue oxygenation response when mice breathed 100%
oxygen (Figure 7D-G). We observed that the oxygen delivered to the brain is significantly
smaller in the aged mouse brain, both in the FC (2-month: 43.7 ± 4.1 μM; 18-month: 32.5 ± 10.4
μM; 24-month: 27.0 ± 9.5 μM. Linear mixed effects model, $p = 0.0078$) and FL/HL (2-month:
71.0 ± 14.4 μM; 18-month: 51.0 ± 5.3 μM; 24-month: 35.5 ± 21.7 μM. Linear mixed effects
model, $p = 0.0032$).
Lastly, we quantified the functions of the brain capillary network during aging progress,
as its dynamics affect brain oxygenation responses [53,72]. We first compared whether red blood
cell (RBC) velocity differed between age groups in the capillary network. We found no
significant differences in lumen diameter between different groups (2-month: 4.7 ± 1.65 μm, 32
capillaries; 18-month: 4.9 ± 1.2 μm, 36 capillaries; 24-month: 3.6 ± 1.0 μm, 56 capillaries), a
trend toward decreased RBC velocity, but not a statistically significant difference (2-month: 0.58
± 0.33 mm/s; 18-month: 0.53 ± 0.34 mm/s; 24-month: 0.34 ± 0.25 mm/s) (Supplementary Figure
1A), no difference in hematocrit (2-month: $38.3 \pm 7.6\%$; 18-month: $33.4 \pm 9.6\%$; 24-month: 34.5
$\pm 9.7\%$; Supplementary Figure 1B). In addition to RBC flow rate and hematocrit, the "stochastic"
nature of red blood cell distribution in the capillary also affects brain oxygenation [55,72]. When we
quantified the spacing of RBC and the occurrence of "stall" events, we found no significant
difference between different aging groups (Supplementary Text 1).
Collectively, our *in vivo* recording results suggest slowed vascular response dynamics
and decreased oxygen carrying capacity in normal aging, which can create imbalances in
baseline and on demand supply of energy and oxygen in aged brains.
351
352
353
354

Discussion

Understanding structural and functional changes of the cerebrovasculature during normal aging will provide foundational information to understand altered brain energy infrastructure that can be commonly linked with many neurodegenerative disorders. Here, we provide detailed information regarding anatomical changes of the cerebrovascular network and physiological alteration of the blood flow in aged mouse brains, as summarized in Figure 8. We found overall reductions in vascular length and branching densities, along with tortuous arterioles that indicate sparser and remodeled vascular networks in aged brains. We also uncovered selective vascular and pericyte loss in cortical deep layers, basal forebrain regions, and the hippocampal network, including the entorhinal cortex, which may contribute to their regional vulnerabilities in neurodegenerative disorders [41,73]. Lastly, our *in vivo* studies showed delayed neurovascular coupling response time and inefficient oxygen delivery in aged brains. Collectively, our results advance our understanding of global changes and regional vulnerabilities associated with deteriorating vascular networks in aged brains.

Cerebrovascular structural changes with selective pericyte reduction in aged brains.
Previous studies in aged cerebral vasculature have shown stiffened arteries, microvascular rarefaction, and remodeled vascular trees in selected brain regions [16,19,74–76]. Our study showed that there is an approximate 10% decrease in overall vascular density, as well as branching density, in 18-month-old compared to 2-month-old mouse brains, suggesting a sparser vascular network to distribute the blood [33]. Moreover, aged brains showed substantially more tortuous penetrating arterioles, which impede blood flow by increasing flow resistance. This increase in resistance, unless countered by an increase in blood pressure, could result in reduced oxygen and nutrient supply, particularly in distal areas from main arteries such as the deep cortical layers and white matter tracks [19,77]. Importantly, human studies have shown similar changes with tortuous vasculature and slowed cerebral blood flow [21,75,78,79]. Such changes can lead to an increased heart rate to compensate for cerebral hypoperfusion, as frequently observed in elderly population [80]. We also found an overall increase in average vessel radii. Notably, pericytes are known to regulate the basal tone of microvessels [13,15], and a recent study showed that pericytes in superficial cortical layers have impaired recovery of cellular processes in the aged brain [34]. Therefore, while pericyte cell density does not change significantly during aging, their regulatory function may be impaired, resulting in slightly dilated cerebrovasculature.

Our data showed that the vasculature of the isocortex is more resilient to aging compared to other brain regions, as evidenced by no significant changes in both microvascular and capillary pericyte densities. However, deep cortical layers, especially layer 6b, showed reduced vessel density and pericyte density, consistent with previous studies [81]. Notably, layer 6 plays a crucially important role as the output layer to the thalamus [42,82]. Moreover, layer 6b is the only cortical layer that is responsive to sleep-wake neuropeptides such as orexin, which is produced in the lateral hypothalamus [82,83]. Considering that sleep is often dysregulated with increased age in humans [84], failing cerebrovascular network in the deep cortical layer may provide important insight to understand aging related sleep dysregulation.

Since our 3D mapping data examine vascular network changes of the whole mouse brain in an unbiased way, we identified specific brain regions with selective vulnerabilities in aged brains. For example, we found significantly reduced vascular and pericyte densities in the basal forebrain area, which contains cholinergic neurons [85]. The basal forebrain cholinergic neurons

(BFCNs) have highly extensive projections to the cortical area and have large soma size with
high energy demands [86]. Previous clinical and preclinical studies have shown that BFCNs are
highly vulnerable in Alzheimer's disease (AD) and their deterioration is linked with memory
impairment [41,87]. Impaired vascular networks with decreased pericyte density may, potentially
serve as an underlying cause of BFNC degeneration in normal aging and neurodegenerative
disorders, including AD [73,88,89]. Another notable area is the entorhinal cortex (ENT), a part of the
hippocampal network, which has been heavily implicated in AD and particular cognitive deficits
[90,91]. The lateral ENT (ENTl) showed significantly decreased vascular length, branching point,
and capillary pericyte density. The ENTl vascular density is one of lowest across the brain region
in normal adult mice [28]. With additional decreases with aging, blood supply in the ENTl is likely
to be highly limited and less able to withstand further insult, which may explain its vulnerability
to neurodegenerative disorders. Lastly, our study identified specific thalamic and hypothalamic
areas with decreases in the vascular network density, such as the medial preoptic area, which
warrant future studies for these largely understudied subcortical areas in aging research.
415
Slowed neurovascular response and decreased oxygen in aged brains.
In addition to anatomical changes, we found slowed brain hemodynamic responses during
locomotion and whisker stimulation in aged mice, while the amplitude of blood increase
remained intact. This suggests that the aged brain can still deliver enough red blood cells to the
regions with energy demands, but the timing of the delivery is perturbed. Interestingly, the
slowed hemodynamic response is also observed when noradrenergic input from the locus
coeruleus is disrupted [92], consistent with the disruption of the locus coeruleus having a role in
Alzheimer's disease and dementia [93]. Moreover, the baseline oxygenation and oxygen carrying
capacity of the red blood cells decrease with age [94,95]. Notably, respiration is an important
regulator of brain oxygenation [53], and lung function decreases during the aging process [96,97]. The
decreased ability to deliver oxygen can also be related to decreased microvessel density and its
connectivity, resulting in less effective oxygen distribution, and the shift of the oxyhemoglobin
dissociation curve with age [94]. This baseline drop in brain oxygenation will make the brain more
vulnerable to hypoxia when facing increased oxygen demand, as neurons become hyperexcitable
in aged brains [98–100]. This baseline drop in brain oxygenation, in combination with increased
blood flow resistance, due to increased vessel tortuosity and reduced vascular density, will make
brain areas in distal vascular territories, such as white matter tracks, and water shed areas
(located at the junction between main artery territories) selectively vulnerable in aged brains
[21,101]. Finally, as neurovascular coupling potentially serves many other functions besides oxygen
delivery, disruption of the normal hemodynamic response may have other adverse physiological
effects [102].
437
Limitations of the Study
In our anatomical studies, we found significant vascular loss in deep cortical areas and many
subcortical areas (e.g., basal forebrains, hypothalamus, and entorhinal cortex). However, our *in*
vivo measurement is limited to superficial cortical layers, where we did not observe dramatic
anatomical changes. Although the brain hemodynamics at the surface reflect the dynamics along
the vascular tree, future studies with emerging techniques such as functional ultrasound imaging
or three-photon microscopy imaging will help to address functional changes in these important,
yet hard-to-reach brain areas [103,104]. Moreover, our analysis mostly focuses on the arterial and

capillary compartments of the vasculature. Future studies are needed to elucidate how aging
affects the structure and function of the venous side of aged brains.
448
Summary
Taken together, our study reveals aging-related brain-wide and area-specific changes in vascular
and mural cell types. These changes can explain the vulnerability and resilience of different brain
areas in normal aging. Moreover, we identified an age-related decrease in brain oxygenation and
delayed neurovascular coupling, which can be linked with cognitive impairment in aged brains.
These aging-related changes will serve as a common factor in understanding many
neurodegenerative disorders and cognition decline in the elderly population.

Acknowledgments

We thank Josephine Liwang for constructive discussion of the manuscript, Dr. Volkhard Linder for kindly sharing PDGFRβ-Cre transgenic mice and Dr. Zhuhao Wu for sharing tissue clearing protocol. We thank Jen Minteer and Becca Betty for valuable management of our animal colony, as well as the coding support provided by Fae Kronman. We acknowledge the use of computational resources in the High Performance Computing cluster at the Penn State College of Medicine.

Funding:

National Institutes of Health grant R01NS108407 and RF1MH12460501 (YK)

National Institutes of Health grant R01NS078168 and R01NS101353 (PJD)

American Heart Association Career Development Award #935961 (QZ)

Its contents are solely the responsibility of the authors and do not necessarily represent the views of the funding agency.

Author Contributions

Conceptualization: YK, HCB

Data Collection: HCB, UC, YK

Developing Computational Analysis: YTW

Data Analysis: HCB, QZ, YTW, HP, YK

In vivo imaging: QZ, PJD

Manuscript preparation: HCB, YK, QZ with help from other authors.

Declaration of Interests

The authors declare no competing interests.

487
488 RESOURCE AVAILABILITY
489
490 **Lead contact**
491 Further information and requests for resources should be directed to and will be fulfilled by the
492 lead contact, Yongsoo Kim (yuk17@psu.edu).
493
494 **Materials availability**
495 This study did not generate new unique reagents.
496
497 **Data and code availability**
498 All dataset and codes can be used for non-profit research without any restriction. Any additional
499 information required to reanalyze the data reported in this paper is available from the lead
500 contact upon request.
501
502

503   **METHODs**
504
Animals
Animal experiments were approved by the Institutional Animal Care and Use Committee at Penn
State University. Adult male and female mice were used across all age and genotype groups in
this study. For transgenic pericyte-specific experiments, PDGFRβ-Cre mice (a kind gift from the
Volkhard Lindner Lab) [37] were crossed with female Ai14 mice which express a Cre-dependent
tdTomato fluorescent reporter (LoxP-Stop-LoxP-tdTomato). These PDGFRβ-Cre:Ai14 mice
exhibit PDGFRβ expression in two distinct mural cell types, pericytes and vascular smooth
muscle cells. Both the adult 2-month-old and 18-month-old PDGFRβ-Cre:Ai14 mice were bred
and aged in house. We used tail genomic DNA with PCR for the transgenic mouse lines
requiring genotyping. Adult 2-month-old C57BL/6J mice were bred from C57BL/6J mice
directly obtained from the Jackson Laboratory and used for vascular tracing experiments with
FITC filling (n=4) [28]. 18-month-old C57BL/6J mice utilized for FITC-fill vascular mapping
experiments were aged from a local C57BL/6 mouse colony. 24-month-old C57BL/6J mice used
for the current study were directly obtained from the National Institute of Aging at 18 months
and aged to 24 months in house. All animals were used once to generate data, and aged animals
with tumors or other appreciable abnormalities were excluded from analysis.
For the *in vivo* two-photon imaging experiments, a total of 31 C57BL/6J mice of both
sexes (2-26 months old, 18-35 g, Jackson Laboratory) were used. Recordings of cerebral blood
volume and cerebral oxygenation response to locomotion were made from 23 mice (2-4 month
old: n = 7 mice; 18 month old: n = 5 mice; 22-26 month old: n = 11 mice) using wide field
optical imaging. In a subset of the mice (2-4 month old: n = 5 mice; 22-26 month old: n = 5
mice), we also recorded cerebral blood volume and cerebral oxygenation response to whisker
stimulus using optical imaging. Recordings of stacks, capillary blood flow velocity, and
diameters of arteries and veins using two-photon laser scanning microscopy (2PLSM) were
conducted in 23 mice (15 of these 23 mice were also used for wide field optical imaging; 2-4
month old: n = 10 mice; 18 month old: n = 5 mice; 22-26 month old: n = 8 mice). Mice were
given food and water *ad libitum* and maintained on 12-hour (7:00–19:00) light/dark cycles. All
experiments were conducted during the light period of the cycle.
533
Perfusion based vascular labeling, STPT imaging, and computational analysis
Overall procedure remains similar to our previous publication [28]. The detailed procedure has
been included in a separate protocol paper [31]. Briefly, animals were deeply anesthetized with
ketamine-xylazine, and perfused with 1X PBS followed by 4% paraformaldehyde to wash out
blood and allow for tissue fixation, respectively. For vessel labeling, immediately following 4%
paraformaldehyde, 0.1% (w/v) fluorescein isothiocyanate (FITC)-conjugated albumin (Sigma-
Aldrich, cat.no.: A9771-1G) in a 2% (w/v) solution of porcine skin gelatin (Sigma-Aldrich,
cat.no: G1890-500G) was perfused to obtain vascular filling. For STPT imaging, the brain
sample was embedded in oxidized agarose and cross-linked in 0.05M sodium borohydrate at 4°C
for at least 2 days ahead of imaging. We used 910nm wavelength (UltraII, Coherent) as
excitation light for all samples. Signals in the green and red spectrum were simultaneously
collected using 560 nm dichroic mirror at x,y = 1,1 μm resolution in every 50 μm z (for pericyte
mapping) or x,y,z = 1,1,5 μm resolution (for vascular mapping).
We utilized our previously described software pipeline to perform de-aberration,
normalization, and imaging stitching steps for all STPT data collected for this study [28]. Moreover,

we used the same analytical tools to binarize the vessel signals and skeletonize for further
analysis. This pipeline also performs cleaning/reconnecting of artifacts, traces the vessel
diameter, and finally outputs the coordinates for each vessel segment and its connectivity. For
pericyte cell counting, we used previously developed Deep Learning Neural Network (DLNN)
cell counting [28]. This DLNN uses a per-cell multi-resolution-hybrid ResNet classification with
potential cell locations to reduce computational time and resources without loss of quality. While
aged mouse brains do have increased noise due to the accumulation of cellular debris, we
validated that our DLNN pipeline performed at the same level as with young adult mice and did
not incorporate cellular debris as potential cells.
558
Tissue clearing, 3D immunolabeling, and LSFM imaging
Whole brain vascular staining was performed following the iDISCO+ protocol with
modifications [24]. Brain samples were delipidated in SBiP buffer, consisting of ice-cold water,
50mM Na_2HPO_4, 4% SDS, 2-methyl-2-butanol and 2-propanol. This buffer is activated at room
temperature and is therefore made and stored at 4°C before use. Each sample was submerged in
10ml of SBiP buffer, rotated at room temperature with buffer changes at 3 hours, 6 hours and
then incubated with fresh SBiP buffer overnight. For adequate delipidation, particularly for aged
samples, each brain was then washed with SBiP for a total of 6 days, with daily buffer changes.
After delipidation, brain samples were washed with B1n buffer, which consists of 0.1% TritonX-
100, 1g of glycine, 0.01% 10N NaOH and 20% NaN3. Brain samples were washed with 10ml of
B1n buffer at room temperature for 2 days. To begin immunolabeling, brains were rinsed 3 times
for 1 hour each with PTwH buffer, consisting of 1X PBS, 0.2% Tween-20, 10mg heparin, and 2g
of NaN3. For primary antibody incubations, antibodies were diluted in antibody solution
consisting of PTwH buffer with 5% DMSO and 3% normal donkey serum. Antibodies to smooth
muscle actin (Acta2) (Rabbit anti-Acta2, Abcam, cat: ab5694, dilution 1:1000) and transgelin
(Sm22) (Rabbit anti-Sm22 Abcam, cat: ab14106, dilution 1:1500) were combined to label the
artery wall, as previously described [24]. Pan-vascular labeling was achieved through staining with
DyLight-594 labeled Lycopersicon Esculentum (Tomato) Lectin (Vector labs, cat. no.: DL-1177-
1), which was added to both primary and secondary incubations at 1:100 concentration. Pericytes
were labeled by combining PDGFRβ (Goat anti- PDGFRβ , R&D Systems, cat. no.: AF1042,
dilution: 1:100) and Mouse Aminopeptidase N/CD13 (Goat anti-CD13, R&D Systems, cat. no.:
AF2335, dilution: 1:100). Primary antibodies were incubated for 10 days at 37°C. Following
primary incubation, PTwH buffer was changed 4-5 times for each sample over the course of 24
hours. A fresh antibody solution was used to dilute all secondary antibodies to a concentration of
1:500. For secondary antibodies, Alexa Fluor® 488-AffiniPure Fab Fragment Donkey Anti-
Rabbit IgG (H+L) (Jackson ImmunoResearch laboratories, cat. no.: 711-547-003) was used to
detect artery staining and Alexa Fluor® 647-AffiniPure Fab Fragment Donkey Anti-Goat IgG
(H+L)
(Jackson ImmunoResearch laboratories, cat. no.: 705-607-003) was utilized to detect pericyte
staining. After secondary incubation for 10 days at 37°C, brains were washed 4-5 times in PTwH
buffer for 24 hours. Brain samples were then dehydrated in a series of methanol dilutions in
water (1-hour washes in 20%, 40%, 60%, 80% and 100%). An additional wash of 100%
methanol was conducted overnight to remove any remaining water. The next day, brains were
incubated in 66% dichloromethane/33% methanol for 3 hours and subsequently incubated in 100%
dichloromethane twice for at least 15 minutes each. Brains were equilibrated in dibenzyl ether
for at least two days before transitioning to ethyl cinnamate one day prior to imaging.

We used the SmartSPIM light sheet fluorescence microscope (LifeCanvas Technologies).
Brains were supported in the custom sample holder by standardized pieces of dehydrated agarose
consisting of 1% agarose in 1X TAE buffer. The sample holder arm was then submerged in ethyl
cinnamate for imaging. We used a 3.6X objective (LifeCanvas, 0.2NA, 12mm working distance,
1.8μm lateral resolution) and three lasers (488nm, 560nm, 642nm wavelengths) with a 2mm step
size. For detailed examination of pericytes, we used a 15X objective (Applied Scientific
Instrumentation, 0.4NA, 12mm working distance, 0.4mm lateral resolution) with a 1μm z step
size. Acquired data was stitched using custom Matlab codes adapted from Wobbly Stitcher [24].
603
Analysis of LSFM-based vascular and pericyte signals
For pericyte counting, prior to quantification, each stitched image stack, per signal channel, was
separately normalized, and the entire volume of each image stack was then converted to 20 μm
maximum intensity projections (MIP). Normalization of each signal channel is done by adjusting
according to the histogram-determined global mean value of the background by utilizing a 10x
downsized copy of the entire image stack. The 20μm MIP step was determined to prevent over or
under counting of cell bodies, since pericyte cell body size tended to range from 6-10μm
depending on the orientation of the cell measured within a 3D context in the original image stack.
Finally, all three image channels (artery, lectin, and mural cell labels) were merged into a
channel overlay to provide additional context, such as vascular zone information. Cells with
stereotypical pericyte morphology (i.e., ovoid shape and protruding from the vessel wall),
typically along the first through third order arteriolar branches, that also expressed smooth
muscle markers and extended processes that wrapped around the vessel were classified as
ensheathing pericytes. Capillary pericytes were classified according to cell body shape and
localized to the capillary bed without any Acta2/Transgelin expression. These cells were further
subdivided into mesh or thin strand morphologies according to their microvessel placement and
type of processes, according to the definitions of these subtypes [105]. Cell bodies along larger
veins, including the principle cortical venules were excluded from this analysis.
For arteriole analysis, 600 μm MIPs were obtained from the channel labeling of Acta2
and Transgelin (i.e., artery labeling). We cropped the supplementary somatosensory cortex from
full datasets and quantified the total number of arteries and their branches manually.
For tortuosity measurements, a centerline of the entire vessel length was first traced to
obtain the Euclidean distance (arc length) using a skeletonization tool in Clearmap 2.0 [24]. Then a
straight line connecting the start and end points of the previous length was obtained to measure
the chord length. The arc chord ratio was then determined by dividing the Euclidean distance by
the arc chord distance. We used 32 arteries from 2-month-old (n=3 animals) and 23 from 24-
month-old (n=3 animals) in the medial prefrontal, and 19 arteries from 2-month-old (n=2
animals) and 23 from 24-month-old (n=3 animals). For Circle of Willis analysis, entire brain
datasets for the artery channel were converted to 10μm isotropic. Next, a cropped volume
including the branching point of the middle cerebral artery as well as ample segments of the
anterior communicating artery and middle cerebral artery were obtained within 250x250x120 μm
(x,y,z) to fully capture the entire branch point and associated arteries in x,y,z dimensions. This
subset was re-sliced to obtain the cross-sectional area of this section of the vessel. The average
radius was obtained from the cross-sectional areas.
638
639   **Surgery, habituation, and measurement for *in vivo* recording**

Cerebral oxygenation, cerebral blood volume (CVB) and vessel diameter data were acquired from the same groups of awake, behaving mice during voluntary locomotion and whisker stimulation. All surgeries were performed under isoflurane anesthesia (in oxygen, 5% for induction and 1.5-2% for maintenance). A custom-machined titanium head bolt was attached to the skull with cyanoacrylate glue (#32002, Vibra-tite). The head bolt was positioned along the midline and just posterior to the lambda cranial suture. Two self-tapping 3/32" #000 screws (J.I. Morris) were implanted into the skull contralateral to the measurement sites over the frontal lobe and parietal lobe. For measurements using two-photon laser scanning microscopy (2PLSM), CBV measurement using intrinsic optical signal (IOS) imaging or brain oxygenation measurement using spectroscopy, a polished and reinforced thin-skull (PoRTS) window was made covering the right hemisphere or both hemispheres as described previously [53–58,62]. Following the surgery, mice were then returned to their home cage for recovery for at least one week, and then started habituation on experimental apparatus. Habituation sessions were performed 2-4 times per day over the course of one week, with the duration increasing from 5 min to 45 min.

Habituation. Animals were gradually acclimated to head-fixation on a spherical treadmill [53,55,57,106] with one degree of freedom over at least three habituation sessions. The spherical treadmill was covered with nonabrasive anti-slip tape (McMaster-Carr) and attached to an optical rotary encoder (#E7PD-720-118, US Digital) to monitor locomotion. Mice were acclimated to head-fixation for ~15 minutes during the first session and were head-fixed for longer durations (> 1 hour) in the subsequent sessions. Mice were monitored for any signs of stress during habituation. In all cases, the mice exhibited normal behaviors such as exploratory whisking and occasional grooming after being head-fixed. Heart rate fluctuations were detectable in the intrinsic optical signal [53,62] and varied between 7 and 13 Hz for all mice after habituation, which is comparable to the mean heart rate (~12 Hz) recorded telemetrically from mice in their home cage [107]. Habituation sessions were achieved 2-4 times per day over the course of one week, with the duration increasing from 5 min to 45 min. Mice that received whisker stimulation (n = 10) were acclimatized to head-fixation for 15–30 min during the first session. In subsequent sessions, they began to receive air puffs directed at the whiskers and were head-fixed for longer durations (> 60 minutes).

Physiological measurements. Data from all experiments (except two photon laser scanning microscopy) were collected using custom software written in LabVIEW (version 2014, National Instruments). *Behavioral measurement.* The treadmill movements were used to quantify the locomotion events of the mouse. The animal was also monitored using a webcam (Microsoft LifeCam Cinema®) as an additional behavioral measurement. *Vibrissa stimulation.* Animals were awake and engaged in whisking behavior during IOS data acquisition [54,56]. Brief (0.1-s duration) puffs of air were delivered to the ipsilateral and contralateral whiskers through a thin plastic tube (length 130 mm, diameter 2 mm). Air puffs were directed to the distal ends of the whiskers at an angle parallel to the face to prevent stimulation of other parts of the head or face. An additional air puffer was set up to point away from the body for use as an auditory stimulus. The puffs were delivered via solenoid actuator valves (Sizto Tech Corporation, 2V025 1/4) at constant air pressure (10 psi) maintained by an upstream regulator (Wilkerson, R03-02-000). Air puffs were separated by intervals of 30-60 s, and the order of all sensory stimulation was randomized, with a nominal ratio of three contralateral stimuli for every ipsilateral or auditory stimulation. Auditory and ipsilateral stimuli were omitted from the principal analysis because their responses were primarily related to stimulus-provoked movement. *Brain oxygen*

measurement using optical imaging. We mapped the spatiotemporal dynamics of oxyhemoglobin
and deoxyhemoglobin concentrations using their oxygen-dependent optical absorption spectra [72].
Reflectance images were collected during periods of green LED light illumination at 530 nm
(equally absorbed by oxygenated and deoxygenated hemoglobin, M530L3, Thorlabs) or blue
LED light illumination at 470 nm (absorbed more by oxygenated than deoxygenated hemoglobin,
M470L3, Thorlabs). For these experiments, a CCD camera (Dalsa 1M60) was operated at 60 Hz
with 4X4 binning (256 X 256 pixels), mounted with a VZM300i optical zoom lens (Edmund
Optics). Green and blue reflectance data were converted to changes in oxy- and
deoxyhemoglobin concentrations using the modified Beer-Lambert law with Monte Carlo-
derived wavelength-dependent path length factors [61]. We used the cerebral oxygenation index [71]
(i.e., HbO-HbR) to quantify the change in oxygenation, as calculating the percentage change
requires knowledge of the concentration of hemoglobin on a pixel-by-pixel basis, which is not
feasible given the wide heterogeneity in the density of the cortical vasculature [30]. *Measurements*
using two-photon laser scanning microscopy (2PLSM). Mice were briefly anesthetized with
isoflurane (5% in oxygen) and retro-orbitally injected with 50 µL 5% (weight/volume in saline)
fluorescein-conjugated dextran (70 kDa, Sigma-Aldrich), and then fixed on a spherical treadmill.
Imaging was done on a Sutter Movable Objective Microscope with a 20X, 1.0 NA water dipping
objective (Olympus, XLUMPlanFLN). A MaiTai HP (Spectra-Physics, Santa Clara, CA) laser
tuned to 800 nm was used for fluorophore excitation. All imaging with the water-immersion lens
was done with room temperature distilled water. All the 2PLSM measurements were started at
least 20 minutes after isoflurane exposure to avoid the disruption of physiological signals due to
anesthetics.

For navigational purposes, wide field images were collected to generate vascular maps of
brain pial vascular maps of the entire PoRTS window. We performed three different
measurements using 2PLSM. (1) To measure blood vessel diameter responses to locomotion,
individual arteries and veins were imaged at nominal frame rate of 3 Hz for 5 minutes using 10-
15 mW of power exiting the objective. Diameter of pial vessels were calculated using Radon
transform [108]. (2) To measure RBC velocity and RBC spacing, line scan images were collected
from individual capillaries (diameter range: 2-8 µm). The pixel dwell time for the line scan
segments was 1 µs and we achieved a ~1.5 kHz sampling rate. (3) To measure the vasculature
diameter under physiological conditions (i.e., awake and resting), we collected stack image every
other day for each mouse. For each mouse, we collected data from 4 different days and collected
3 different trials on each day. Shortly (within 20 minutes) after the last trial on the last day, the
mouse was perfused for future vasculature reconstruction. The resolution for each XY plane is
0.64 µm/pixel and the resolution for Z direction is 1 µm. On the Z-direction, three frames were
collected and averaged, the averaged frame was saved in the file. All the images were acquired
with increasing laser power up to 100 mW at a depth of ~200 um.

Isoflurane challenge. To compare the capability of vasodilation in both young and aged
mice, we exposed a subset of mice to short period (~ 2 minutes) of isoflurane (5% in pure
oxygen) and imaged the pial vessel (specifically, the branch of the middle cerebral artery)
diameter responses. This allowed us to assess the magnitude of diameter change of pial arteries
and veins.

Oxygen challenge experiments. In a subset of experiments, hyperoxia was induced by
substituting breathing air for 100% pure oxygen. Using optical imaging of spectroscopy, we
performed an oxygen challenge. Mice were head-fixed on a spherical treadmill, and a nose cone
was fixed ~ 1 inch in front of the nose, with care taken not to contact the whiskers. Two gases

were administered during a 5-min spectroscopy trial in the following order: 1 min breathable air
(21% oxygen), 3 min 100% oxygen, and 1 min breathable air. Mice breathed breathable air for at
least 2 min between trials, to ensure physiological parameters returned to baseline. Reflectance
images were collected during periods of green LED light illumination at 530 nm (equally
absorbed by oxygenated and deoxygenated hemoglobin, M530L3, Thorlabs) or blue LED light
illumination at 470 nm (absorbed more by oxygenated than deoxygenated hemoglobin, M470L3,
Thorlabs) or red LED light illumination at 660 nm (absorbed more by deoxygenated than
oxygenated hemoglobin, M660L2, Thorlabs).
740
Data analysis for *in vivo* recording.
All data analyses were performed in Matlab (R2019b, MathWorks) using custom code.
Locomotion event identification. Locomotion events [53,57,106] from the spherical treadmill were
identified by first applying a low-pass filter (10 Hz, 5^{th} order Butterworth) to the velocity signal
from the optical rotary encoder, and then comparing the absolute value of acceleration (first
derivative of the velocity signal) to a threshold of 3 cm/s^2. Periods of locomotion were
categorized based on the binarized detection of the treadmill acceleration:

$$\delta(t) = \theta(|a_t| - a_c) = \begin{cases} 1, & |a_t| \geq a_c \\ 0, & |a_t| < a_c \end{cases}$$

where a_t is the acceleration at time t, and a_c is the treadmill acceleration threshold.
Spontaneous activity. To characterize spontaneous (non-locomotion-evoked) activity, we defined
"resting" periods as periods started 4 seconds after the end of previous locomotion event and
lasting no less than 60 seconds.
Calculation of hemodynamic response function. We considered the neurovascular relationship to
be a linear time invariant system [69,109,110]. To provide a model-free approach to assess the
relationship between CBV or vessel diameter and neural activity, hemodynamic response
function (HRF) was calculated by deconvoluting CBV signal, oxygen signal or vessel diameter
signal to locomotion events, respectively, using the following equation:

$$H_{(k+1)\times 1} = (L^T L)^{-1} L^T V_{(m+k)\times 1}$$

757 H is the HRF, V is the tissue oxygenation signal or neural activity signal. L is a Toeplitz matrix
758 of size (m+k) x (k+1) containing binarized locomotion events (n):

$$L(\vec{n}) = \begin{pmatrix} 1 & n_1 & 0 & 0 & \cdots & 0 \\ 1 & n_2 & n_1 & 0 & \cdots & 0 \\ \vdots & \vdots & n_2 & n_1 & \cdots & \vdots \\ \vdots & n_k & \vdots & n_2 & \cdots & n_1 \\ \vdots & 0 & n_k & \vdots & \cdots & n_2 \\ \vdots & \vdots & \vdots & n_k & \ddots & \vdots \\ 1 & 0 & 0 & 0 & \cdots & n_k \end{pmatrix}$$

Comparison of HRF parameters. To quantify the temporal features of HRF, the HRF for CBV
was fitted using a gamma-variate fitting process [69,111–114] using a gamma-variate function kernel
of the following form,

$\mathrm{HRF}(t, T, W, A) = A * \left(\frac{t}{T}\right)^\alpha * e^{\left(\frac{t-T}{-\beta}\right)},$

where $\alpha = (T/W)^2 * 8.0 * \log(2.0)$, $\beta = W^2/(T * 8.0 * \log(2.0))$. For modeling HRF using a
gamma-variate function kernel, we used a downhill simplex algorithm minimizing the sum
square difference between measured and predicted hemodynamics. The goodness of fit was
quantified as $R^2 = 1 - \frac{\sum(HRF_{actual} - HRF_{model})^2}{\sum(HRF_{actual} - \overline{HRF})^2}$, where $\overline{HRF}$ is the mean value of the actual HRF.

To quantify the amplitude of each HRF, we used the value at the peak of the modeled HRF.
Time to peak (TTP) was calculated as the time at which the modeled HRF reached its maximum
amplitude. Full-width at half maximum (FWHM) was defined as the time from which the
modeled HRF rose to 50% of its peak until it fell to 50% of its peak. TTP, FWHM and HRF
amplitudes across different cortical depths were compared using a linear model to quantify
trends (robustfit, MATLAB).
2PLSM image processing. (1) To quantify blood vessel diameter responses to locomotion,
individual frames from 2PLSM imaging were aligned using a rigid registration algorithm to
remove motion artifacts in the x–y plane [59]. Visual inspection of movies indicated that there was
minimal z-axis motion. A rectangular box was manually drawn around a short segment of the
vessel and the pixel intensity was averaged along the long axis [59]. Pixel intensity was used to
calculate diameter from the full-width at half-maximum. Periods of rest were segregated using
locomotion events measured with the rotary encoder. For each 5-min trial, diameter
measurements were normalized to the average diameter during periods of rest. The diameters
were smoothed with a third-order, 15-point Savitzky–Golay filter (Matlab function: sgolayfilt).
(2) To quantify RBC velocity, blood flow velocity was calculated using Radon transform [58].
Only blood flow velocity during resting periods was reported. Capillary diameter was manually
measured using ImageJ software. To quantify RBC spacing, we utilized the method reported in
our previous study [55]. We identified RBC "stall" events as an inter-RBC spacing greater than 1
second. We only used RBCs spacing intervals during relatively long resting segments (i.e., ≥ 5
second). (3) As the perfusion procedure and brain fixation might affect the brain vasculature [115],
to compare our measurements for vessel radii in STPT and LSFM imaging datasets to vessel
parameters measured in vivo using 2PLSM, the same animals that were used for 2PLSM and
STPT imaging were reconstructed and compared, as described before [28].
791
Statistical analysis
For the STPT and LSFM dataset, we used Matlab (Mathworks) and/or Prism (Graphpad) for all
statistical analysis, including multi-region of interest (ROI) correlation analysis. We used an
averaged value of the experimented animals while treating each ROI as an individual data point.
For two group comparisons, multiple unpaired t-tests were used with multiple comparison
corrections. The p value was adjusted with the false discovery rate for multiple comparison
corrections using the Two-stage step-up method of Benjamini, Krieger and Yekutieli in
Graphpad. For multiple group comparisons, two-way ANOVA, or mixed model if including
NaN values, to generate comparison between groups using Prism (Graphpad).
For *in vivo* recording, all summary data were reported as the mean ± standard deviation (SD)
unless stated otherwise. The normality of the samples was tested before statistical testing using
the Anderson-Darling test (adtest). For comparison of multiple populations, the assumption of
equal variance for parametric statistical method was also tested (vartest2). If criteria of normality
and equal variance were not met, parametric tests (unpaired t test) were replaced with a
nonparametric method (Wilcoxon rank sum test). For comparisons of oxygen challenge and
isoflurane challenge effects on brain hemodynamics across different age groups, we used the
linear mixed effect model (MATLAB function: fitlme). Significance was accepted at $p < 0.05$.

References

1. Lu, H. *et al.* Alterations in cerebral metabolic rate and blood supply across the adult lifespan. *Cereb Cortex* **21**, 1426–1434 (2011).

2. Pfister, F. *et al.* Pericyte migration: A novel mechanism of pericyte loss in experimental diabetic retinopathy. *Diabetes* (2008) doi:10.2337/db08-0325.

3. Brown, W. R. & Thore, C. R. Review: Cerebral microvascular pathology in ageing and neurodegeneration. *Neuropathology and Applied Neurobiology* **37**, 56–74 (2011).

4. Baloyannis, S. J. & Baloyannis, I. S. The vascular factor in Alzheimer's disease: A study in Golgi technique and electron microscopy. *Journal of the Neurological Sciences* **322**, 117–121 (2012).

5. Almeida, V. M. *et al.* Pericytes Make Spinal Cord Breathless after Injury. *Neuroscientist* **24**, 440–447 (2018).

6. Hayden. Type 2 Diabetes Mellitus Increases The Risk of Late-Onset Alzheimer's Disease: Ultrastructural Remodeling of the Neurovascular Unit and Diabetic Gliopathy. *Brain Sciences* **9**, 262 (2019).

7. Abrahamson, E. E. & Ikonomovic, M. D. Brain injury-induced dysfunction of the blood brain barrier as a risk for dementia. *Experimental Neurology* **328**, 113257 (2020).

8. Uemura, M. T., Maki, T., Ihara, M., Lee, V. M. Y. & Trojanowski, J. Q. Brain Microvascular Pericytes in Vascular Cognitive Impairment and Dementia. *Front. Aging Neurosci.* **12**, 80 (2020).

9. Iadecola, C. *et al.* The Neurovasculome: Key Roles in Brain Health and Cognitive Impairment: A Scientific Statement From the American Heart Association/American Stroke Association. *Stroke* **0**, (2023).

10. Kapogiannis, D. & Mattson, M. P. Disrupted energy metabolism and neuronal circuit

dysfunction in cognitive impairment and Alzheimer's disease. *Lancet Neurol* **10**, 187–198

(2011).

11. Cunnane, S. C. *et al.* Brain energy rescue: an emerging therapeutic concept for

neurodegenerative disorders of ageing. *Nat Rev Drug Discov* **19**, 609–633 (2020).

12. Nedergaard, M. & Goldman, S. A. Glymphatic failure as a final common pathway to

dementia. *Science* **370**, 50–56 (2020).

13. Hartmann, D. A., Coelho-Santos, V. & Shih, A. Y. Pericyte Control of Blood Flow Across

Microvascular Zones in the Central Nervous System. *Annu Rev Physiol* **84**, 331–354 (2022).

14. Sweeney, M. D., Zhao, Z., Montagne, A., Nelson, A. R. & Zlokovic, B. V. Blood-Brain

Barrier: From Physiology to Disease and Back. *Physiological reviews* **99**, 21–78 (2019).

15. Hartmann, D. A. *et al.* Brain capillary pericytes exert a substantial but slow influence on

blood flow. *Nature Neuroscience* **24**, 633–645 (2021).

16. Yang, T., Sun, Y., Lu, Z., Leak, R. K. & Zhang, F. The impact of cerebrovascular aging on

vascular cognitive impairment and dementia. *Ageing Res Rev* **34**, 15–29 (2017).

17. *Pericyte Biology in Disease*. vol. 1147 (Springer International Publishing, 2019).

18. Ding, R. *et al.* Loss of capillary pericytes and the blood–brain barrier in white matter in

poststroke and vascular dementias and Alzheimer's disease. *Brain Pathol.* **30**, 1087–1101

(2020).

19. Li, Y. *et al.* Aging-associated changes in cerebral vasculature and blood flow as determined

by quantitative optical coherence tomography angiography. *Neurobiology of Aging* **70**,

148–159 (2018).

20. Tarumi, T. & Zhang, R. Cerebral Blood Flow in Normal Aging Adults: Cardiovascular

Determinants, Clinical Implications, and Aerobic Fitness. *J Neurochem* **144**, 595–608

(2018).

21. Juttukonda, M. R. *et al.* Characterizing cerebral hemodynamics across the adult lifespan with

arterial spin labeling MRI data from the Human Connectome Project-Aging. *Neuroimage*

230, 117807 (2021).

22. Sweeney, M. D., Kisler, K., Montagne, A., Toga, A. W. & Zlokovic, B. V. The role of brain

vasculature in neurodegenerative disorders. *Nature Neuroscience* **21**, 1318–1331 (2018).

23. Xiong, B. *et al.* Precise Cerebral Vascular Atlas in Stereotaxic Coordinates of Whole Mouse

Brain. *Frontiers in Neuroanatomy* **11**, 128 (2017).

24. Kirst, C. *et al.* Mapping the Fine-Scale Organization and Plasticity of the Brain Vasculature

Resource Mapping the Fine-Scale Organization and Plasticity of the Brain Vasculature.

Cell 1–16 (2020) doi:10.1016/j.cell.2020.01.028.

25. Todorov, M. I. *et al.* Machine learning analysis of whole mouse brain vasculature. *Nature*

Methods **17**, 442–449 (2020).

26. Ji, X. *et al.* Brain microvasculature has a common topology with local differences in

geometry that match metabolic load. *Neuron* **109**, 1168-1187.e13 (2021).

27. Bennett, H. C. & Kim, Y. Advances in studying whole mouse brain vasculature using high-

resolution 3D light microscopy imaging. *NPh* **9**, 021902 (2022).

28. Wu, Y. *et al.* Quantitative relationship between cerebrovascular network and neuronal cell

types in mice. *Cell Reports* **39**, (2022).

29. Blinder, P. *et al.* The cortical angiome: an interconnected vascular network with

noncolumnar patterns of blood flow. *Nat Neurosci* **16**, 889–897 (2013).

30. Tsai, P. S. *et al.* Correlations of Neuronal and Microvascular Densities in Murine Cortex

Revealed by Direct Counting and Colocalization of Nuclei and Vessels. *Journal of*

Neuroscience **29**, 14553–14570 (2009).

31. Liwang, J. K., Bennett, H. C., Pi, H.-J. & Kim, Y. Protocol for using serial two-photon

tomography to map cell types and cerebrovasculature at single-cell resolution in the whole

adult mouse brain. *STAR Protoc* **4**, 102048 (2023).

32. Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference

Atlas. *Cell* **181**, 936-953.e20 (2020).

33. Maheswaran, S. *et al.* Longitudinal regional brain volume changes quantified in normal

aging and Alzheimer's APP x PS1 mice using MRI. *Brain Res* **1270**, 19–32 (2009).

34. Berthiaume, A.-A. *et al.* Pericyte remodeling is deficient in the aged brain and contributes to

impaired capillary flow and structure. *Nat Commun* **13**, 5912 (2022).

35. Brown, L. S. *et al.* Pericytes and Neurovascular Function in the Healthy and Diseased Brain.

Front. Cell. Neurosci. **13**, 282 (2019).

36. Berthiaume, A. A. *et al.* Dynamic Remodeling of Pericytes In Vivo Maintains Capillary

Coverage in the Adult Mouse Brain. *Cell Reports* **22**, 8–16 (2018).

37. Cuttler, A. S. *et al.* Characterization of Pdgfrb-Cre transgenic mice reveals reduction of

ROSA26 reporter activity in remodeling arteries. *Genesis (New York, N.Y.: 2000)* **49**,

673–680 (2011).

38. Hartmann, D. A., Underly, R. G., Watson, A. N. & Shih, A. Y. A murine toolbox for

imaging the neurovascular unit. *Microcirculation (New York, N.Y.: 1994)* **22**, 168–182

(2015).

39. Alheid, G. F. Extended amygdala and basal forebrain. *Ann N Y Acad Sci* **985**, 185–205

(2003).

40. Sato, A., Sato, Y. & Uchida, S. Regulation of cerebral cortical blood flow by the basal

forebrain cholinergic fibers and aging. *Autonomic Neuroscience* **96**, 13–19 (2002).

41. Ballinger, E. C., Ananth, M., Talmage, D. A. & Role, L. W. Basal Forebrain Cholinergic

Circuits and Signaling in Cognition and Cognitive Decline. *Neuron* **91**, 1199–1218 (2016).

42. Zolnik, T. A. *et al.* Layer 6b Is Driven by Intracortical Long-Range Projection Neurons. *Cell*

Reports **30**, 3492-3505.e5 (2020).

43. Trache, A., Massett, M. P. & Woodman, C. R. Vascular smooth muscle stiffness and its role

in aging. in *Current Topics in Membranes* vol. 86 217–253 (Elsevier, 2020).

44. Montero, D., Pierce, G. L., Stehouwer, C. D. A., Padilla, J. & Thijssen, D. H. J. The impact

of age on vascular smooth muscle function in humans: *Journal of Hypertension* **33**, 445–

453 (2015).

45. Nishimura, N., Schaffer, C. B., Friedman, B., Lyden, P. D. & Kleinfeld, D. Penetrating

arterioles are a bottleneck in the perfusion of neocortex. *Proc Natl Acad Sci U S A* **104**,

365–370 (2007).

46. Shih, A. Y. *et al.* Robust and fragile aspects of cortical blood flow in relation to the

underlying angioarchitecture. *Microcirculation* **22**, 204–218 (2015).

47. Ciurică, S. *et al.* Arterial Tortuosity. *Hypertension* **73**, 951–960 (2019).

48. Weiss, D. *et al.* Mechanics-driven mechanobiological mechanisms of arterial tortuosity. *Sci*

Adv **6**, eabd3574 (2020).

49. Cai, C. *et al.* Impaired dynamics of precapillary sphincters and pericytes at first-order

capillaries predict reduced neurovascular function in the aging mouse brain. *Nat Aging* **3**,

173–184 (2023).

50. Khan, U. A. *et al.* Molecular drivers and cortical spread of lateral entorhinal cortex

dysfunction in preclinical Alzheimer's disease. *Nat Neurosci* **17**, 304–311 (2014).

51. Van Hoesen, G. W., Hyman, B. T. & Damasio, A. R. Entorhinal cortex pathology in

Alzheimer's disease. *Hippocampus* **1**, 1–8 (1991).

52. Rodriguez, G. A., Burns, M. P., Weeber, E. J. & Rebeck, G. W. Young APOE4 targeted

replacement mice exhibit poor spatial learning and memory, with reduced dendritic spine

density in the medial entorhinal cortex. *Learning & Memory* **20**, 256–266 (2013).

53. Zhang, Q. *et al.* Cerebral oxygenation during locomotion is modulated by respiration. *Nature*

Communications **10**, 1–15 (2019).

54. Winder, A. T., Echagarruga, C., Zhang, Q. & Drew, P. J. Weak correlations between

hemodynamic signals and ongoing neural activity during the resting state. *Nature*

Neuroscience **20**, 1761–1769 (2017).

55. Zhang, Q., Gheres, K. W. & Drew, P. J. Origins of 1/f-like tissue oxygenation fluctuations in

the murine cortex. *PLOS Biology* **19**, (2021).

56. Zhang, Q., Turner, K. L., Gheres, K. W., Hossain, M. S. & Drew, P. J. Behavioral and

physiological monitoring for awake neurovascular coupling experiments: a how-to guide.

Neurophotonics **9**, (2022).

57. Huo, B.-X., Smith, J. B. & Drew, P. J. Neurovascular coupling and decoupling in the cortex

during voluntary locomotion. *Journal of Neuroscience* **34**, 10975–10981 (2014).

58. Drew, P. J. *et al.* Chronic optical access through a polished and reinforced thinned skull. *Nat*

Methods **7**, 981–984 (2010).

59. Drew, P. J., Shih, A. Y. & Kleinfeld, D. Fluctuating and sensory-induced vasodynamics in

rodent cortex extend arteriole capacity. *Proc Natl Acad Sci U S A* **108**, 8473–8 (2011).

60. Hillman, E. M. Coupling mechanism and significance of the BOLD signal: a status report.

Annu Rev Neurosci **37**, 161–81 (2014).

61. Ma, Y. *et al.* Wide-field optical mapping of neural activity and brain haemodynamics:

considerations and novel approaches. *Philos Trans R Soc Lond B Biol Sci* **371**, (2016).

62. Huo, B. X., Gao, Y. R. & Drew, P. J. Quantitative separation of arterial and venous cerebral

blood volume increases during voluntary locomotion. *Neuroimage* **105**, 369–79 (2015).

63. Vazquez, A. L., Fukuda, M., Crowley, J. C. & Kim, S.-G. Neural and Hemodynamic

Responses Elicited by Forelimb- and Photo-stimulation in Channelrhodopsin-2 Mice:

Insights into the Hemodynamic Point Spread Function. *Cerebral cortex (New York, N.Y. :*

1991) **24**, 2908–2919 (2014).

64. Tian, P. *et al.* Cortical depth-specific microvascular dilation underlies laminar differences in

blood oxygenation level-dependent functional MRI signal. *Proc Natl Acad Sci U S A* **107**,

15246–51 (2010).

65. Longden, T. A. *et al.* Capillary K+-sensing initiates retrograde hyperpolarization to increase

local cerebral blood flow. *Nat Neurosci* **20**, 717–726 (2017).

66. Rungta, R. L., Chaigneau, E., Osmanski, B.-F. & Charpak, S. Vascular

Compartmentalization of Functional Hyperemia from the Synapse to the Pia. *Neuron* **99**,

362-375.e4 (2018).

67. Rungta, R. L. *et al.* Diversity of neurovascular coupling dynamics along vascular arbors in

layer II/III somatosensory cortex. *Commun Biol* **4**, 855 (2021).

68. Bergel, A. *et al.* Adaptive modulation of brain hemodynamics across stereotyped running

episodes. *Nat Commun* **11**, 6193 (2020).

69. Cardoso, M. M., Sirotin, Y. B., Lima, B., Glushenkova, E. & Das, A. The neuroimaging

signal is a linear sum of neurally distinct stimulus- and task-related components. *Nat*

Neurosci **15**, 1298–306 (2012).

70. Drew, P. J. Vascular and neural basis of the BOLD signal. *Curr Opin Neurobiol* **58**, 61–69

(2019).

71. Boas, D. A. & Franceschini, M. A. Haemoglobin oxygen saturation as a biomarker: the

problem and a solution. *Philos Trans A Math Phys Eng Sci* **369**, 4407–24 (2011).

72. Lyons, D. G., Parpaleix, A., Roche, M. & Charpak, S. Mapping oxygen concentration in the

awake mouse brain. *Elife* **5**, (2016).

73. Pandya, V. A. & Patani, R. Region-specific vulnerability in neurodegeneration: lessons from

normal ageing. *Ageing Res Rev* **67**, 101311 (2021).

74. Erdő, F., Denes, L. & de Lange, E. Age-associated physiological and pathological changes at

the blood-brain barrier: A review. *J Cereb Blood Flow Metab* **37**, 4–24 (2017).

75. Zimmerman, B., Rypma, B., Gratton, G. & Fabiani, M. Age-related changes in

cerebrovascular health and their effects on neural function and cognition: A comprehensive

review. *Psychophysiology* **58**, e13796 (2021).

76. Lowerison, M. R. *et al.* Aging-related cerebral microvascular changes visualized using

ultrasound localization microscopy in the living mouse. *Sci Rep* **12**, 619 (2022).

77. Gunning-Dixon, F. M., Brickman, A. M., Cheng, J. C. & Alexopoulos, G. S. Aging of

Cerebral White Matter: A Review of MRI Findings. *Int J Geriatr Psychiatry* **24**, 109–117

(2009).

78. Fulop, G. A. *et al.* Role of age-related alterations of the cerebral venous circulation in the

pathogenesis of vascular cognitive impairment. *American Journal of Physiology-Heart and*

Circulatory Physiology **316**, H1124–H1140 (2019).

79. Sun, Z. *et al.* Age-Related Tortuosity of Carotid and Vertebral Arteries: Quantitative

Evaluation With MR Angiography. *Frontiers in Neurology* **13**, (2022).

80. Oliveros, E. *et al.* Hypertension in older adults: Assessment, management, and challenges.

Clin Cardiol **43**, 99–107 (2019).

81. Schager, B. & Brown, C. E. Susceptibility to capillary plugging can predict brain region

specific vessel loss with aging. *J Cereb Blood Flow Metab* **40**, 2475–2490 (2020).

82. Hoerder-Suabedissen, A. *et al.* Subset of Cortical Layer 6b Neurons Selectively Innervates

Higher Order Thalamic Nuclei in Mice. *Cereb Cortex* **28**, 1882–1897 (2018).

83. Hoyer, D. & Jacobson, L. H. Orexin in sleep, addiction and more: Is the perfect insomnia

drug at hand? *Neuropeptides* **47**, 477–488 (2013).

84. Nixon, J. P. *et al.* Sleep disorders, obesity, and aging: the role of orexin. *Ageing Res Rev* **20**,

63–73 (2015).

85. Zaborszky, L. *et al.* Specific Basal Forebrain-Cortical Cholinergic Circuits Coordinate

Cognitive Operations. *Journal of Neuroscience* **38**, 9446–9458 (2018).

86. Li, X. *et al.* Generation of a whole-brain atlas for the cholinergic system and mesoscopic

projectome analysis of basal forebrain cholinergic neurons. *Proceedings of the National*

Academy of Sciences **115**, 415–420 (2018).

87. Grothe, M. *et al.* Reduction of basal forebrain cholinergic system parallels cognitive

impairment in patients at high risk of developing Alzheimer's disease. *Cerebral cortex*

(New York, NY⬜: 1991) **20**, 1685–1695 (2010).

88. Fischer, V. W., Siddiqi, A. & Yusufaly, Y. Altered angioarchitecture in selected areas of

brains with Alzheimer's disease. *Acta Neuropathol* **79**, 672–679 (1990).

89. Casu, M. A., Pan Wong, T., De Koninck, Y., Ribeiro-da-Silva, A. & Cuello, A. C. Aging

Causes a Preferential Loss of Cholinergic Innervation of Characterized Neocortical

Pyramidal Neurons. *Cerebral Cortex* **12**, 329–337 (2002).

90. Sagare, A. P. *et al.* Pericyte loss influences Alzheimer-like neurodegeneration in mice.

Nature Communications **4**, (2013).

91. Zhang, X. *et al.* High-resolution mapping of brain vasculature and its impairment in the

hippocampus of Alzheimer's disease mice. *National Science Review* **6**, 1223–1238 (2019).

92. Bekar, L. K., Wei, H. S. & Nedergaard, M. The locus coeruleus-norepinephrine network

optimizes coupling of cerebral blood volume with oxygen demand. *J Cereb Blood Flow*

Metab **32**, 2135–2145 (2012).

93. Jacobs, H. I. L. *et al.* In vivo and neuropathology data support locus coeruleus integrity as

indicator of Alzheimer's disease pathology and cognitive decline. *Sci Transl Med* **13**,

eabj2511 (2021).

94. Korkushko, O. V., Ivanov, L. A., Pisaruk, A. V. & Chebotarev, N. D. The respiratory

function of blood in elderly and old age and the factors that determine it. *Human*

Physiology **35**, 163–169 (2009).

95. Moeini, M. *et al.* Compromised microvascular oxygen delivery increases brain tissue

vulnerability with age. *Sci Rep-Uk* **8**, 8219 (2018).

96. Haas, C. F., Loik, P. S. & Gay, S. E. Airway clearance applications in the elderly and in

patients with neurologic or neuromuscular compromise. *Respir Care* **52**, 1362–81;

discussion 1381 (2007).

97. Schulte, H., Muhlfeld, C. & Brandenberger, C. Age-Related Structural and Functional

Changes in the Mouse Lung. *Front Physiol* **10**, 1466 (2019).

98. Haberman, R. P., Koh, M. T. & Gallagher, M. Heightened cortical excitability in aged

rodents with memory impairment. *Neurobiol Aging* **54**, 144–151 (2017).

99. Fischer, C. *et al.* Prevention of age-associated neuronal hyperexcitability with improved

learning and attention upon knockout or antagonism of LPAR2. *Cell Mol Life Sci* **78**, 1029–

1050 (2021).

100. Popescu, I. R. *et al.* Increased intrinsic excitability and decreased synaptic inhibition in

aged somatosensory cortex pyramidal neurons. *Neurobiol Aging* **98**, 88–98 (2021).

101. Kapasi, A. *et al.* Watershed Microinfarct Pathology and Cognition in Older Persons.

Neurobiol Aging **70**, 10–17 (2018).

102. Drew, P. J. Neurovascular coupling: motive unknown. *Trends Neurosci* **45**, 809–819 (2022).

103. Wang, T. *et al.* Three-photon imaging of mouse brain structure and function through the

intact skull. *Nat Methods* **15**, 789–792 (2018).

104. Brunner, C. *et al.* Whole-brain functional ultrasound imaging in awake head-fixed mice.

Nat Protoc **16**, 3547–3571 (2021).

105. Grant, R. I. *et al.* Organizational hierarchy and structural diversity of microvascular

pericytes in adult mouse cortex. *Journal of Cerebral Blood Flow and Metabolism* **39**, 411–

425 (2019).

106. Gao, Y. R. & Drew, P. J. Effects of Voluntary Locomotion and Calcitonin Gene-Related

Peptide on the Dynamics of Single Dural Vessels in Awake Mice. *J Neurosci* **36**, 2503–16

(2016).

107. Gehrmann, J. *et al.* Phenotypic screening for heart rate variability in the mouse. *Am J*

Physiol Heart Circ Physiol **279**, H733-40 (2000).

108. Drew, P. J., Blinder, P., Cauwenberghs, G., Shih, A. Y. & Kleinfeld, D. Rapid

determination of particle velocity from space-time images using the Radon transform. *J*

Comput Neurosci **29**, 5–11 (2010).

109. Boynton, G. M., Engel, S. A., Glover, G. H. & Heeger, D. J. Linear systems analysis of

functional magnetic resonance imaging in human V1. *J Neurosci* **16**, 4207–21 (1996).

110. Glover, G. H. Deconvolution of impulse response in event-related BOLD fMRI.

Neuroimage **9**, 416–29 (1999).

111. Madsen, M. T. A simplified formulation of the gamma variate function. *Phys Med Biol* **37**,

1597–1600 (1992).

112. Cohen, M. S. Parametric analysis of fMRI data using linear systems methods. *Neuroimage*

6, 93–103 (1997).

113. Sirotin, Y. B. & Das, A. Anticipatory haemodynamic signals in sensory cortex not

predicted by local neuronal activity. *Nature* **457**, 475–9 (2009).

114. Yu, X., Qian, C., Chen, D. Y., Dodd, S. J. & Koretsky, A. P. Deciphering laminar-specific

neural inputs with line-scanning fMRI. *Nat Methods* **11**, 55–8 (2014).

115. Korogod, N., Petersen, C. C. & Knott, G. W. Ultrastructural analysis of adult mouse

neocortex comparing aldehyde perfusion with cryo fixation. *Elife* **4**, (2015).

1076
1077

Figure 1

Figure 1. STPT based vessel tracing and pericyte counting pipeline
The steps of the STPT pipeline are outlined in order. **1**. Brain sample collection: vascular filling procedure via cardiac perfusion with FITC conjugated albumin gel and pericyte mapping using PDGFRβ-Cre;Ai14 reporter mice. **2**. STPT imaging: combination of 2-photon microscope 2D tile imaging with serial vibratome sectioning to obtain cellular resolution images of the whole mouse brain. These image tiles are then stitched to reconstruct tissue sections spanning the olfactory bulb to the cerebellum. **3**. Signal detection: vascular tracing with binarization of FITC filled vascular signals and skeletonization, and deep learning-based detection of capillary pericytes. **4**. Mapping signals in a reference brain: All detected signals were registered to the Allen Common Coordinate Framework (Allen CCF) and an isocortical flatmap was used to examine signals in the isocortex.

Figure 2

Figure 2. Region specific reduction of vascular length and branching density and increased vascular radii

(**A**) The averaged vasculature length density of 2-month-old (N=4) and 18-month-old (N=5) brains registered to the Allen CCF. (**B-C**) Summed vessel length (**B**) and brain volume (**C**) in 2-month-old and 18-month-old brains. (**D-F**) Scatter plots of averaged vascular length density (m/mm^3) (**D**), vascular branching density (1/mm^3) (**F**), and vascular radii (μm) between 2-month-old (x axis) and 18-month-old (y axis) brains across different brain regions. Areas with statistically significant differences were highlighted with magenta boxes. (**G-I**) Scatter plots of the isocortex data for averaged vascular length density (**G**), vascular branching density (**H**), vascular radii (**I**) between 2-month-old (x axis) and 18-month-old (y axis) brains. Isocortical areas are color coded based on grouping in **J**. The solid yellow line represents y=x and the dotted line on either side represents a 10% difference from the solid yellow line. (**J**) Isocortical flatmap with Allen CCF border lines and region-based color-coding. Y axis: Bregma anterior-posterior (A-P) coordinates, X axis: the azimuth coordinate represents the physical distance by tracing the cortical surface on the coronal cut. (**K**) Averaged vascular length density of different cortical layers in the flatmap between 2-month-old and 18-month-old brains. A white arrow highlights the significant decrease of vascular length density in the layer 6 cortical layers. (**L-M**) 250μm maximum intensity projection images of the primary somatosensory cortex (**L**) and the infralimbic cortex (**M**) with vascular tracing (green on the right side) between 2-month-old and 18-month-old brains. Note the significant reduction of vasculature in the deep layer. (**N**) Both vascular length and branching density showed significant reductions in layer 6. Brain region abbreviations can be found in Supplementary data 1 or Allen atlas at https://atlas.brain-map.org/atlas?atlas=602630314.

Figure 3

Figure 3. Aged brain showed selective reduction of capillary pericytes in the basal forebrain area and the deep cortical layer

(**A**) Averaged capillary pericyte density in 2-month-old (n=10) and 18-month-old (n=9) PDGFRβ-Cre;Ai14 mouse brains that are registered to the Allen CCF. (**B**) A scatter plot of capillary pericyte density between 2-month-old (x axis) and 18-month-old (y axis). Brain areas are color coded based on the Allen CCF ontology. Brain areas with significant changes were highlighted with magenta-colored boxes. The solid yellow line represents the value for y=x and the yellow dotted lines on either side of the solid yellow line represent a 10% difference from the

solid yellow line. (**C**) Bar graphs of capillary pericyte density in the substantia innominata and magnocellular nucleus between 2-month-old and 18-month-old brains. (**D**) Representative images of the basal forebrain (left) and higher resolution examples of the magnocellular nucleus and substantia innominata in 2-month-old and 18-month-old brains. Red dots represent detected pericyte cell bodies in each respective region. (**E**) The isocortical flatmap (left) and averaged capillary pericyte densities plotted in the flatmap from 2-month-old and 18-month-old brains. (**F-G**) Scatter plots of capillary pericyte density (**F**) and pericyte coverage (pericyte density per vascular length density; **G**) in isocortical areas. (**H**) Representative images of capillary pericyte density in the infralimbic cortex from 2-month-old and 18-month-old brains. (**I-K**) Layer specific capillary pericyte densities from the infralimbic cortex (**I**), the somatosensory cortex (**J**), and across all isocortical areas (**K**). Note the significant density reduction in the deep cortical layers. All q values obtained from multiple comparison correction by false discovery rate are reported in each graph as well as the uncorrected p-value.

Figure 4

Figure 4. Tissue clearing and 3D immunolabeling with LSFM imaging to examine different vascular compartments and mural cells in the same brain

(**A**) The steps of brain clearing, whole brain immunolabeling, and light sheet fluorescent microscopy (LSFM) pipeline are outlined in order. **1**. Brain sample collection with transcardial perfusion. **2**. Modified iDISCO protocol including delipidation, immunolabeling for arteries, whole vasculature, and pericytes, and optical clearing. **3**. LSFM imaging and data processing to visualize cleared brains at cellular resolution. **4**. Data analysis such as arteriole geometry analysis and pericyte counting. (**B**) 3D reconstruction of a brain with artery staining by LSFM imaging. (**C**) Max projection of the 500 µm thick z stack of the artery staining. (**D-G**) Zoom-in images of the red box area from (**C**). (**D**) artery staining in the green channel, (**E**) lectin based total vasculature staining in the red channel, (**F**) pericyte staining in the far-red channel, (**G**) a merged image of pseudo-colored arteries (blue), total vasculature (green), and pericyte (red). (**H**) Maximum projection of the artery channel in a brain hemisphere.

Figure 5

Figure 5. Aging induces significant arteriole remodeling and selective pericyte density reduction.

(**A**) Schema of main arteries of the circle of Willis at the ventral surface of the brain. (**B**) Artery specific labeling of the middle cerebral artery branching area (red box area in H) from 2-month-old and 24-month-old brains. (**C**) Artery radii do not show a significant difference between the two age groups. (**D**) The number of both total and deep layer 6 reaching penetrating cortical arteriole did not show a significant difference between the two age groups. (**E**) Representative 600 μm MIPs of artery labeling in the somatosensory area of a young (top) and an aged (bottom) brain. Note tortuous arterioles in the old brain (red arrowheads) compared to straight ones in the young brain (light blue arrowheads). (**F**) Old brains showed significantly tortuous arterioles in the medial prefrontal and somatosensory cortices. Data from 3 animals for both young and aged groups. (**G**) Both immediate and total arteriole branch numbers show no significant differences between the two age groups. (**H**) Different pericyte subtypes with immuno markers and their

position in the vascular order. (**I**) Submicron resolution LSFM images with artery labeling, whole vasculature labeled with lectin and mural cell labeling with PDGFRβ and CD13 antibodies. The cyan arrow for an ensheathing pericyte, the yellow arrow for a mesh capillary pericyte, and the purple arrow for a thin-strand capillary pericyte. (**J**) Manual cell counting did not show any significant difference in the somatosensory cortex between the two age groups. (**K-L**) However, layer 6 of the somatosensory cortex (**K**) showed a significant reduction in pericyte density (**L**). (**M-N**) The entorhinal cortex (**M**) showed a significant reduction of capillary pericytes (**N**). All q values obtained from multiple comparison correction by false discovery rate and uncorrected p-value are reported in each graph, except (F) with Bonferroni correction.

Figure 6

Figure 6. Delayed cortical hemodynamic responses to voluntary locomotion and whisker stimulation is delayed in normal aging.

(**A**) Left, schematic of the experimental setup for IOS imaging during voluntary locomotion. Right, an image of thin-skull window and corresponding anatomical reconstruction; scale bar = 1 mm. FC, frontal cortex; FL/HL, forelimb/hindlimb representation of the somatosensory cortex; Wh, vibrissae cortex. (**B**) Population average of locomotion onset (left) and offset (right) triggered average of ΔHbT responses in FL/HL across different age groups. (**C**) As in (**B**) but for FC. (**D**) Hemodynamic response function (HRF) of ΔHbT in the FL/HL across different age groups. (**E**) Quantification of HRF of ΔHbT in the FL/HL: amplitude (A, left), time to peak (T, middle), and full-width at half maximum (FWHM, right). (**F**) As in (**D**) but for FC. (**G**) As in (**E**) but for FC. (**H**) Schematic of the experimental setup for IOS imaging during whisker stimulation. (**I**) Average population responses of ΔHbT to contralateral whisker stimulation in the FL/HL across different age groups. (**J**) Quantification of the whisker stimulation evoked responses of ΔHbT in the FL/HL: amplitude (left), time to peak (middle), and full-width at half maximum (right). (**K**) As in (**I**) but for FC. (**L**) As in (**J**) but for FC. (**M**) Schematic of the experimental setup for 2PLSM imaging during locomotion. (**N**) Population average of locomotion onset (left) and offset (right) triggered average of arteriole diameter responses in FL/HL across different age groups. (**O**) Hemodynamic response function (HRF) of arteriole diameter changes in the FL/HL across different age groups. (**P**) Population average of ΔHbT responses to inhalation of 5% isoflurane in the FL/HL (top) and FC (bottom) across different age groups. Solid lines and shaded areas in (**B**, **C**, **D**, **F**, **I**, **K**, **N**, **O**, **P**) denote mean ± SEM, respectively. Data are shown as mean ± SD in all other graphs.

Figure 7. Oxygen carrying capacity of the blood is reduced by aging.
(**A**) Population average of locomotion onset and offset triggered average of brain oxygenation (ΔHbO-HbR) responses in FL/HL and FC across different age groups. (**B**) Average population responses of ΔHbO-HbR to contralateral whisker stimulation in the FL/HL and FC across different age groups. (**C**) Relationship between locomotion evoked change in ΔHbT and ΔHbO-HbR, 2-5 s after the onset of locomotion, across different age groups, in FL/HL. (**D**) Population average of ΔHbT (top) and ΔHbO-HbR (bottom) responses to inhalation of 100% oxygen in the FL/HL across different age groups. (**E**) As in (**E**) but for FC. (**F**) Group average of fractional changes of ΔHbT (left) and ΔHbO-HbR (right) in response to 100% oxygen in FL/HL across different age groups. (**G**) As in (**F**) but for FC. Solid lines and shaded areas in (**A**, **B**, **E**, **F**) denote mean ± SEM, respectively. Data are shown as mean ± SD in all other graphs.

Figure 8

Figure 8. Summary of changes in aged brains

Aged brains show reduced vascular length and branching density, increased radii, reduced pericyte density, slowed vascular response time, and lower oxygen carrying capacity in the blood compared to young brains.