SifiNet: A robust and accurate method to identify feature gene sets and annotate cells

Qi Gao, Zhicheng Ji, Liuyang Wang, Kouros Owzar, Qi-Jing Li,
Cliburn Chan, Jichun Xie

Abstract

SifiNet is a robust and accurate computational pipeline for identifying distinct gene sets, extracting and annotating cellular subpopulations, and elucidating intrinsic relationships among these subpopulations. Uniquely, SifiNet bypasses the cell clustering stage, commonly integrated into other cellular annotation pipelines, thereby circumventing potential inaccuracies in clustering that may compromise subsequent analyses. Consequently, SifiNet has demonstrated superior performance in multiple experimental datasets compared with other state-of-the-art methods. SifiNet can analyze both single-cell RNA and ATAC sequencing data, thereby rendering comprehensive multiomic cellular profiles. It is conveniently available as an open-source R package.

Introduction

Single-cell sequencing techniques, including single-cell RNA sequencing (scRNA-seq) and single-cell ATAC sequencing (scATAC-seq), allow researchers to quantify cellular omic phenotypes. An ideal single-cell data analysis is expected to help researchers understand cellular omic heterogeneity, extract the cell subpopulations of interest, identify feature gene sets corresponding to cell subpopulations, as well as reveal the relationship between cell subpopulations.

Among these analysis tasks, identifying feature gene sets is a crucial step. Feature gene sets are defined as gene sets that are differentially expressed between cell subpopulations. They are often used to annotate cell subpopulations and to perform gene set enrichment analysis. Existing feature gene identification methods often employ a two-step approach (called two-step methods thereafter): cells are first clustered (such as Seurat[1], simple Louvain[2], CIDR[3], and SCANPY[4]) and differentially expressed gene (DEG) analysis (such as DESeq2[5], edgeR[6], limma[7], limma-voom[8], and MAST[9]) is then performed across cell clusters to identify cluster-specific feature genes. However, this approach has questionable accuracy on data with complex or subtle heterogeneity, since an inaccurate initial clustering step could lead to subsequent erroneous feature gene identification[10]. Alternatively, some methods identify feature genes by detecting highly variable genes (HVG) based on their deviations from the population with respect to deviance of model fitting[11], dropout rates[12], and UMI count distributions[13] (called HVG methods thereafter). However, these methods

do not separate feature genes into subpopulation-specific gene sets, limiting their utility for annotating cells.

To overcome the limitations of existing methods, we present SifiNet, a unique approach for direct identification of feature gene sets that eliminates the need for prior cell clustering. Stemming from the crucial observation that genes co-differentially expressed within a cell subpopulation also exhibit co-expression patterns (Suppl. Note 1), SifiNet builds a gene co-expression network and inspects its topology to discern feature gene sets. These gene sets serve to compute cellular gene set enrichment scores and subsequently annotate cells. Additionally, networks among these gene sets imply relationships between cell subpopulations (Fig. 1). Furthermore, SifiNet can optionally integrate scATAC-seq data, as it forms a gene co-open-chromatin network and explores its topology to determine epigenomic feature gene sets. The ability of SifiNet to analyze both scRNA-seq and scATAC-seq data endows researchers with insight into cellular multiomic heterogeneity.

We demonstrate that SifiNet outperforms existing two-step methods and HVG methods in identifying feature gene sets and enhancing cell annotation accuracy. Moreover, we illustrate that SifiNet can identify complex heterogeneity among cells and reveal potential developmental lineages among cell subpopulations. SifiNet can also be scaled to analyze datasets with millions of cells. We applied SifiNet to five published experimental datasets and uncovered several potentially novel discoveries, such as potential new cell cycle markers and senescence markers, the senescent-cell enriched subpopulation, the developmental lineage of myeloid progenitor cells, and the CD8 cell subpopulations along with their possible transition paths.

Results

To benchmark SifiNet's performance, we applied SifiNet to simulated and experimental datasets, including two simulated datasets (SD1, SD2) and five experimental datasets (Monoclonal[14], IPF1[15], BoneMarrow[16], CD8[17], MassiveRNA[18]). An additional experimental dataset IPF2[15] was used for validation. Also, we compared the performance of SifiNet against 5 two-step methods, 3 HVG identification methods, and 1 trajectory analysis method. See Methods, Suppl. Table 1, and Suppl. Table 2 for details of the datasets and the competing methods.

SifiNet identifies feature gene sets with high accuracy. We applied SifiNet to SD1, Monoclonal, and IPF1, to identify the feature gene sets. SifiNet achieved a high accuracy in identifying known feature gene sets, and also identified potential new cell cycle markers and new ciliated cell senescence markers.

First, we applied SifiNet and competing methods to SD1, a dataset consisting of 100 simulated scRNA-seq data replications with rare cell type. Among all methods, SifiNet has the highest median F1 score of 0.62; which is more than twice the best score among competing methods (Fig. 2a). Among them, singleCellHaystack, limma-CIDR-TS, limma-voom-CIDR-TS, and MAST-CIDR-TS exhibit higher median sensitivity over 100 replications than SifiNet, but significantly lower median specificity. High specificity is crucial in feature gene identification studies due to the large number of candidate genes. For instance, each

Figure 1: **SifiNet pipeline.** SifiNet is a versatile pipeline with multiple functions. Taking the pre-processed feature count matrix, SifiNet can use gene co-expression network topology to identify feature gene sets (function 1). Then, based on those feature gene sets, it can extract and annotate the corresponding cell subpopulations (function 2). Last but not least, it can construct a cell subpopulation network to reveal their relationship (function 3). See Methods for details.

replication of SD1 dataset contains, on average, 83 feature genes and 3318 non-feature genes. A specificity as high as 0.9 will falsely label, on average, 332 among 3318 non-feature genes as feature genes, substantially obscuring the signal of the true feature genes. SifiNet's main advantage is its high specificity (close to 1), indicating that most identified genes are genuine feature genes.

We further investigated why the commonly used two-step methods underperform on SD1. Assuming that true cell type labels are known, we applied DEG methods to identify cell-type-specific feature genes. These methods are referred as oracle-cluster methods. All oracle-cluster methods perform well on SD1, exhibiting high sensitivity and specificity (Suppl. Fig. 1). This indicates that the key factor is the clustering accuracy. If the resulting clustering is perfect, the DEG step can effectively identify feature genes. However, in practice, exact cell types are unknown, and clustering methods can only provide a reasonable estimate. When these estimates are inaccurate, such as on SD1 (Suppl. Fig. 2), two-step methods perform poorly.

Second, we applied SifiNet to the Monoclonal dataset to identify the feature genes related to cell cycles. We derived monoclonal CT26 tumor cells from a single ancestral cell. Within this population, we reasoned that the main transcriptomic difference among individual cells should reflect their cell cycle stage. SifiNet identified two feature gene sets. Set 1 contains many G1 and S (G1/S) phase genes, such as GINS, MCM, DNA polymerase genes, and

Figure 2: **(a)-(b) Performance comparison between SifiNet and competing methods on SD1 and Monoclonal.** Sensitivity, specificity, and F1 accuracy scores of 3 types of methods: SifiNet (red), HVG methods (green), and two-step methods (blue). (a) Performance on SD1, a simulated dataset with 100 replications. (b) Performance on Monoclonal to identify cell cycle marker genes. The benchmark cell cycle marker genes were curated from existing literature (Methods). **(c) Verificiation of SifiNet-identified feature genes with independent dataset.** SifiNet identified the feature genes in IPF1. Then, DEG analysis was performed on IPF2 to show their Log2 fold changes between KRT5-/KRT17+ (senescence-enriched) epithelial cells and proliferating (non-senescent) epithelial cells in IPF2. Stars represent the significance level of the log2 fold change, *: adjusted p-value $<= 10^{-2}$, **: adjusted p-value $<= 10^{-4}$, ***: adjusted p-value $<= 10^{-6}$.

RPA complex genes. Interestingly, it also includes HAUS genes (*Haus3*, *Haus5*, *Haus6*, *Haus8*). Previously, HAUS genes were thought to be related to the centrosome cycle by maintaining the integrity of the centrosome. Although many other centrosome-associated genes are considered to be markers for G2/M phase[19], studies in Heinrichs[20] indicate that centrosome integrity might be also a checkpoint for G1 to S phase transition. Without HAUS proteins maintaining the centrosome integrity, the cells may be unable to progress into the S phase. SifiNet analysis pinpointed the differential expression of HAUS genes is strongly correlated with other G1/S phase feature genes. Set 2 features are composed with many well-known G2 and M (G2/M) phase genes, including Tubulin and SKA. The full list of identified genes is in Suppl. Table 3.

We also applied the two-step methods to Monoclonal. We compared the SifiNet-identified feature genes and the two-step-method-identified feature genes to the benchmark gene set containing 437 benchmark cell cycle marker genes collected from the GSEA MSigDB[21–23] and Uniprot[24] databases (Methods). Among them, 137 benchmark genes were identified by SifiNet. Several methods identified more genes than SifiNet; however, their specificities are much lower (Fig. 2b). For instance, MAST-CIDR-TS has the highest sensitivity, identifying all 437 benchmark genes, but it also identified 8,134 irrelevant genes, complicating the understanding of cell cycles. Overall, SifiNet has the highest F1 score among all methods.

Third, we applied SifiNet to IPF1 to identify senescence-related feature gene sets. IPF1 contains 10,375 ciliated cells from Idiopathic Pulmonary Fibrosis (IPF) human lung tissues. Prior research has demonstrated a close relationship between senescence and IPF[25,26]. Therefore, by exploring cell-type-specific feature genes, we expect to identify feature genes associated with senescence. SifiNet identified three feature gene sets (Suppl. Table 3). Set 1 includes numerous ribosomal protein genes, known to accumulate in senescent cells[27]. Set 2 contains *Jun*, *Junb*, *Fos*, and *Fosb* from the Activator Protein-1 (AP-1) transcription factor (TF) family. Other genes in Set 2, such as *Gdf15*, *Egr1*, and *Zfp36*, have been previously found to interact with the AP-1 TF family[28–30]. This family is known to be involved in proliferation, differentiation, and tumorigenesis[28], highly related to the senescence process[29]. Set 3 genes are all mitochondrial genes. Previous studies have shown that mitochondrial genes are related to the senescence-associated secretory phenotype (SASP)[31], and mitochondrial dysfunction is associated with senescence[32]. These three gene sets may relate to different aspects of senescence.

To verify the identified senescence genes, we checked their log-fold change and significance levels using an independent dataset, IPF2 (Fig. 2c). IPF2 contains two cell subpopulations: KRT5-/KRT17+ epithelial cells and proliferating epithelial cells; the former are treated as senescent cells, and the latter non-senescent. Among the 84 SifiNet identified genes, 45 are significantly differentially expressed (adjusted p-values < 0.0001, Fig. 2c) between two cell subpopulations. The analysis of independent dataset IPF2 verified the quality of the identified feature genes.

SifiNet identifies cell subpopulations with high accuracy. Here, we aim to identify cell subpopulations corresponding to different cell states, such as different cell cycles and senescence level. Cells with the same functional cell type but different states only have subtle differences; thus, existing clustering methods often fail to separate them well. However, SifiNet achieves a high accuracy in calling these cell subpopulations with subtle differences.

Figure 3: **(a)-(c) UMAPs of Monoclonal cells.** The cells were annotated by (a) SifiNet or (b) Seurat cell cycle scores, or clustered by (c) CIDR. For CIDR, the cell clusters are annotated to best map to the Seurat cell cycle score annotation. Red circles mark the major difference between CIDR annotation and the benchmark annotation. **(d)-(f) SifiNet-annotated BoneMarrow cell subpopulations.** The x- and y-axis of (d) and (e) are the first two principal components (PCs) of the GSVA scores based on the SifiNet-identified gene sets. (d) When applied to all cells, SifiNet annotated three cell subpopulations: early progenitors, megakaryocyte/erythrocyte progenitor cells, and granulocyte/macrophage progenitor cells. (e) Applied to the granulocyte/macrophage progenitor cells, SifiNet identified 4 cell subpopulations: Granulocyte progenitor cells, macrophage progenitor cells, Monocyte and dentritic progenitor cells, and T and B progenitor cells. T and B cells are not myeloid cells but all these cells share common progenitor cells, hematopoietic stem cells. These cells may get mixed up with the myeloid progenitor cells. (f) Heatmap of some marker gene expressions in the annotated cell subpopulations.

First, we applied SifiNet to Monoclonal to call cells in different cell cycles. Fig. 3a displays the Uniform Manifold Approximation and Projection (UMAP)[33] plot of the cells annotated by SifiNet. To benchmark its performance, we also annotated cells according to the Seurat cell cycle scores using known cell cycle markers[1,34] (Fig. 3b). Although SifiNet does not utilize any known marker information, the resulting cell UMAP closely resembles the benchmark UMAP. In contrast, we applied CIDR to cluster cells into three cell clusters, without using the known cell cycle markers. The clustering results are less satisfactory: notably, some G1 and G2/M phase cells are clustered into the S phase (Fig. 3c). Furthermore, CIDR, with the default resolution parameter, yielded 8 clusters for cell clustering; the results were hard to interpret. (Suppl. Fig. 3).

Second, we applied the SifiNet to label the senescent cells in IPF1 (Methods). Then, we check whether the widely known senescence marker, $Cdkn1a$[35], is differentially expressed in the SifiNet-senescent cells versus the others. It turns out that $Cdkn1a$ has log2 fold change 0.636, and adjusted p-value= 1.54×10^{-34}. As a comparison, we used SenMayo, a recently published senescence marker set[36], to label the senescence cells. We found that $Cdkn1a$ is not significantly differentially expressed in the SenMayo-senescent cells (log2 fold change 0.281, adjusted p-value= 0.169). This suggests that SifiNet-identified gene sets annotated senescent ciliated cells better than the SenMayo senescence marker set.

SifiNet revealed cellular hierarchical heterogeneity. Cellular heterogeneity is often hierarchical, especially for stem-like cells. We applied SifiNet to BoneMarrow, single cell transcriptome dataset with heterogenous cells at various development stage isolated from mouse bone marrow. And SifiNet successfully predicted the developmental lineage of the myeloid progenitor cells before they grow into the downstream cells.

Taking the gene expression data from the whole cell set, SifiNet separated them into three major populations with distinct transcriptomic features of early myeloid progenitor cells, megakaryocyte/erythrocyte progenitor cells, and granulocyte/macrophage progenitor cells (Fig. 3d). Then we applied SifiNet again to the labeled granulocyte/macrophage progenitor cells, and further divided them into 3 cell subpopulations: granulocyte progenitor cells, macrophage progenitor cells, monocyte and dendritic progenitor cells (Fig. 3e). In addition, a subset of cells do not belong to any of these 3 subpopulations. After checking their DEGs, we found that some T cell and B cell markers are over-expressed in these cells; thus, they are labeled as lymphoid progenitors. These cells may be mixed up with the myeloid progenitor cells when the original study sorted the cells[16].

SifiNet revealed accurate cell subpopulation network in SD2. We applied SifiNet to SD2, a simulated dataset with five cell subpopulations representing five lineage developmental stages (Fig. 4b). SifiNet identified five gene sets, separated the cells into five corresponding subpopulations, and constructed the network. Clearly, the cell subpopulation network built by SifiNet (Fig. 4a) is identical to the actual network (Fig. 4b). As a comparison, we applied Monocle3[33,37–41], a popular single-cell pipeline for cell lineage analysis. The resulting cell subpopulation network (Fig. 4c) is different from the actual lineage. Thus, SifiNet outperforms Monocle3 in SD2 to reveal cell developmental lineage.

Figure 4: **(a)-(c) UMAPs of SD2**, with (a) the SifiNet-built lineage, (b) the true cell subpopulation lineage, and (c) the Monocle3-built lineage. **(d)-(f) Cellular GSVA score PC plots for CD8.** The x- and y-axis are PC directions of the GSVA scores based on the SifiNet-derived gene sets using scRNA-seq and scATAC-seq data. The cells were annotated by (d) Seurat on the multiome data (e) SifiNet on the scRNA-seq data (f) SifiNet on the scATAC-seq data. Cell subpopulation networks are shown in (e) and (f).

SifiNet accurately annotated CD8 T cells and identified long-lasting T cells.
SifiNet is able to differentiate cells into subpopulations based on either single-cell transcriptomic or epigenomic data. Moreover, it can provide useful insights on the relationships between cell subpopulations.

In the CD8 scRNA-seq data, SifiNet annotated the cells with five cell subpopulations. Central memory T cells (TCM_1, TCM_2) and Naïve T cells (TNaive) were previously annotated by Seurat as Naïve cells, and the TCM cells are missing based on Seurat annotations (Fig. 4d). This suggests that SifiNet has a stronger ability in differentiating cells. Further, SifiNet identified two effector T cell (TEM_1, TEM_2) subpopulations, both exhibiting bridging and transitional characteristics between TCM and TNaive cells (Fig. 4e). These results suggest CD8 subpopulation lineage relationships.

Previous studies used gene chromatin accessibility profiles to supplement genes' expression profiles for defining T cell subpopulations.[42–44]. We adopted the same strategy. In the CD8 scATAC-seq data, SifiNet identified two cell subpopulations, showing excessive chromatin accessibility on TCM and TEM marker genes. In particular, we identified some CD8 cells with a TCM cell transcriptome profile but also excessive chromatin accessibility on TEM markers (Fig. 4e,f). They were considered as the most long-lasting T cells after yellow fever vaccination in a recent study[45]. Thus, further studying this CD8 cell subpopulation may lead to new insights of long-term immune protection.

SifiNet scales to million-cell datasets. The rapid development of single-cell technology is leading to the generation of large-scale single-cell data collection. Thus, we tested SifiNet's computation performance on a super-large-scale scRNA-seq dataset, MassiveRNA, containing 10,448 genes and 1,287,071 cells after pre-processing. SifiNet took 12.2 hours and 101 GB peak memory to complete the analysis. To learn the impact of cell numbers on time cost and peak memory usage, we down-sampled MassiveRNA dataset into smaller datasets with cell numbers ranging between 10,000 to 1,000,000. SifiNet analysis on these datasets showed that the log10 running time is roughly linear with the log10 cell numbers (Suppl. Fig. 4). When the cell number is less than 50,000, the peak memory usage was 4.1 GB, dominated by the multiple testing procedure. As the cell number increases above 50,000, the log10 peak memory usage was also roughly linear with the log10 cell number; the peak memory usage was reached in the co-expression calculation step (Suppl. Fig. 4). Overall, SifiNet is efficient with respect to computation time and memory usage. In general, the computation complexity of SifiNet is $O\left(Np^2\right)$, where N and p are the numbers of cells and genes after pre-processing.

Discussion. Cell subpopulation annotation and understanding their relationships pose challenges, particularly when cells exhibit complex and subtle heterogeneity in states such as developmental stages, cell cycles, and senescence levels. While traditional clustering methods perform well in annotating subpopulations when there are clear functional distinctions reflected in cellular transcriptome profiles, they are unreliable in the presence of nuanced heterogeneity. To address these limitations, traditional methods often rely on domain-knowledge-assisted approaches. For instance, Seurat utilizes known cell cycle markers to derive cell cycle scores. However, for less understood cell statuses like senescence, the availability of marker

genes may be limited. SifiNet demonstrates significant potential in these areas to identify new cell status markers and their associated cells.

Moving forward, there are several avenues for further methodological improvement. Firstly, SifiNet's performance is influenced by the quality of the gene co-expression network. Currently, we use quantile association networks due to their superior performance over Pearson and rank correlation networks for count data[46]. Future exploration can focus on finding better co-expression representations to optimize SifiNet's performance. Secondly, SifiNet utilizes the Louvain method with default resolution parameters to cluster feature genes, which generally works well when gene sets are well-separated. However, for gene sets with less distinct boundaries, optimizing gene set clustering becomes crucial to further enhance SifiNet's performance on more challenging datasets. Thirdly, the incorporation of domain-knowledge-assisted approaches into SifiNet is being considered. This would allow SifiNet to leverage partially known marker genes, accurately capturing more feature genes and identifying corresponding cell subpopulations.

Methods

Computation Pipeline of SifiNet

SifiNet is a versatile pipeline with multiple functions, such as identifying feature gene sets, recognizing cell subpopulations, and uncovering potential developmental lineage among cell subpopulations. Its main idea is to employ gene co-expression network topology to identify feature gene sets (Fig. 1). Its main advantage lies in its high accuracy, particularly when cells display complex heterogeneity structures. For instance, some cells may be in a transitional stage, some cell subpopulations may share feature genes, and certain cell subpopulations may exhibit hierarchical subpopulation structures. SifiNet can effectively handle these complex cases.

Background and Notations

Feature gene sets. We define feature genes as those over-expressed within a cell subpopulation. Those genes under-expressed within a cell subpopulation are treated as the feature genes of the remaining cells, due to identifiability issues.

Definition 1 (Feature gene). Gene i is called the feature gene of the cell subpopulation $\mathcal{C}$ if $P(Y_{ic} \leq y \mid c \in \mathcal{C}) \leq P(Y_{ic} \leq y \mid c \neq \mathcal{C})$ for any y, and $P(Y_{ic} \leq y_0 \mid c \in \mathcal{C}) < P(Y_{ic} \leq y_0 \mid c \neq \mathcal{C})$ for some y_0.

A set of these genes, exhibiting co-differential expression within the same cell subpopulation, constitutes a feature gene set.

Gene co-expression network. Gene co-expression network is a graph representation of gene co-expression patterns. Each node represents a gene, and each edge represents the co-expression relationship between two genes. Let Y_{ic} be feature values of gene i in cell c. For scRNA-seq data, the gene feature value could be the gene UMI count; for scATAC-seq data,

it could be the aggregated gene-level cut number. Let s_c be cell c's library size (such as total read counts). For any two genes, if $Y_{ic} \not\perp\!\!\!\perp Y_{jc} \mid s_c$, then genes i and j co-express; thus, there will be an edge between nodes i and j. s_c can be further extended to include other cell-specific covariates.

Measuring and estimating gene co-expression. We use quantile association to quantify gene co-expression: $D_{ij} = \frac{1}{N} \sum_c P(Y_{ic} \leq q_{ic}, Y_{jc} \leq q_{jc} \mid s_c) - \tau_i \tau_j$, where q_{ic} and q_{jc} are the τ_i-th and τ_j-th conditional quantiles of gene i and gene j's expressions. Our previous study shows that quantile associations can effectively measure general dependence, especially for count data such as RNA-seq data[46]. Here, we establish a standardized estimator S_{ij} for quantile associations:

$$S_{ij} = \frac{\sum_{c=1}^{N}[I(Y_{ic} \leq \hat{q}_{ic}, Y_{jc} \leq \hat{q}_{jc}) - \hat{\tau}_i \hat{\tau}_j]}{\sqrt{N \hat{\tau}_i (1 - \hat{\tau}_i) \hat{\tau}_j (1 - \hat{\tau}_j)}}, \tag{1}$$

where $\hat{q}_{ic}$ is the estimated cell specific quantile of gene i, and $\hat{\tau}_i = \sum_{c=1}^{N} I(Y_{ic} \leq \hat{q}_{ic})/N$. By default, we use the 50%-quantile among the non-zero observations $q_{ic} = \alpha_{i0} + \alpha_{i1} s_c$ to estimate $\hat{q}_{ic}$. However, when gene i has very low expression (when the median of Y_{ic} is equal to 0 or Y_{ic} are either 0 or 1), we set $\hat{q}_{ic} = 0$ for stability and robustness consideration.

Feature genes and co-expression patterns. Assume $Y_{ic} \perp\!\!\!\perp Y_{jc} \mid c$. In Suppl. Note 1, we list the mathematical proof for a few cases.

- *Case 1: two genes over-expressing in the same cell subpopulation.* If two genes i and j over-expressed in a subpopulation $\tilde{\mathcal{C}}$, then $D_{ij} > 0$, *i.e.*, they exhibit positive co-expression.

- *Case 2: two genes over-expressing in two non-overlapping cell subpopulations.* Suppose gene i over-express in $\tilde{\mathcal{C}}_1$ and gene j over-express in $\tilde{\mathcal{C}}_2$, $\tilde{\mathcal{C}}_1 \cap \tilde{\mathcal{C}}_2 = \emptyset$. When cells have similar library sizes, $D_{ij} < 0$, *i.e.*, these two genes exhibit negative co-expression.

- *Case 3: at least one of the genes is not a feature gene.* If Gene i or gene i is not a feature gene, then $D_{ij} = 0$, *i.e.*, these two genes are independent.

To sum up, two feature genes exhibit positive or negative co-expression depending on the cell subpopulations these two genes over-express. We verified those mathematical derivations in an experimental dataset containing T and B cells (Suppl. Note 1). Moreover, under Case 3, we derive the asymptotic distribution of S_{ij}, a standardized estimator of D_{ij} (Suppl. Note 2).

Co-expression network topology. The mathematical derivations assume all the genes are independent given a cell. In reality, this assumption is not true; thus, not all the co-expressed genes are feature genes. For examples, if one gene regulates another, they will also co-express. Besides, hub genes usually co-express with many other genes. However, we found that feature genes, pathway genes, and hub genes exhibit different local topologies in the co-expression network. See Suppl. Fig. 5 for an illustration. Specifically, feature genes tend to form close community, while pathway genes and hub genes do not. To capture

the community patterns of feature genes, we introduce three node-specific topology measures: first-order, second-order, and third-order connectives. Denote the gene co-expression network $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$ with node set $\mathcal{V}$ and edge set $\mathcal{E}$. Here, $\mathcal{E} = \cup \mathcal{E}_i$, where $\mathcal{E}_i$ is gene i's edge set. For any set $\mathcal{S}$, $|\mathcal{S}|$ is its cardinality.

- First-order connectivity: measuring gene i's connectivity with other genes, defined as $C_{1i} = |\mathcal{E}_i|/(p-1)$.

- Second-order connectivity: measuring connectivity among gene i's neighbors, defined as $C_{2i} = |\mathcal{E} \cap \{(j,k) : j,k \in \mathcal{N}_i\}|/|\mathcal{N}_i|(|\mathcal{N}_i| - 1)$, where $\mathcal{N}_i = \{j : (i,j) \in \mathcal{E}_i\}$ is gene i's neighbor set.

- Third-order connectivity: measuring connectivity among gene i's 2^{nd}-order neighbors (*i.e.*, neighbors' neighbors). Some genes may have too many 2^{nd}-order neighbors. To improve computation efficiency, we employ a sampling strategy: randomly sample m genes from gene i's 2^{nd}-order neighbors, with the sampling probabilities proportional to the number of the gene's connecting edges with the genes in $\mathcal{N}_i$. In other word, the weight of gene k for calculating gene i's 3^{rd}-order connectivity $w_{ik} \propto |\{j : (k,j) \in \mathcal{E}, j \in \mathcal{N}_i\}|$. The resulting gene set at repetition ℓ is denoted by $\mathcal{A}_i^{(\ell)}$. Then, C_{3i} computes the connectivity levels between the genes in $\mathcal{A}_i^{(\ell)}$ as $C_{3i} = \sum_{\ell=1}^{L} |\mathcal{E} \cap \{(j,k) : j,k \in \mathcal{A}_i^{(\ell)}\}|/Lm(m-1)$.

As we anticipate that feature genes will form communities with other feature genes, we define a gene as a feature gene if it has high C_{2i} and C_{3i} and moderate or high C_{1i}. Genes with low C_{1i}, C_{2i} and C_{3i} could be genes on the pathways or hub genes. Some feature genes may serve multi-roles as pathway genes and/or hub genes. These genes will have different topology structures from other feature genes. We will identify these multi-role feature genes in the end, after we identify other feature genes.

SifiNet function 1: Identifying feature gene sets

SifiNet rests upon a key observation: co-differentially-expressed genes show co-expression patterns, even if the expressions of those genes are conditionally independent given the cell types. Thus, feature genes, which co-differentially-express in the same cell subpopulations, tend to link with each other in the co-expression network, forming a close community.

Based on this observation, SifiNet first constructs a co-expression network based on quantile association measures. Then, based on topology measures, we choose those genes that form a close community. Next, we focus on positive co-expression network of those genes and their community patterns to cluster genes into multiple core feature gene sets. These core feature gene sets will be further enriched to add additional genes, such as shared feature genes and genes performing multi-role functions.

Suppose after pre-processing, the single-cell dataset contains p genes and N cells. Here is a short description on the default steps for identifying feature gene sets. The flowchart illustrating these steps is in Suppl. Fig. 6.

Step 1: Initiating a gene co-expression graph. Initiate gene co-expression graph by $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$ with $\mathcal{V} = [p]$. To obtain $\mathcal{E}$, we set up $p(p-1)/2$ hypotheses: $\text{H}_{\text{null},ij} : Y_{ic} \perp\!\!\!\perp Y_{jc} \mid s_c$, for all $i,j \in \mathcal{V}$ and $i < j$. Then we perform multiple testing on $\text{H}_{\text{null},ij}$ to determine the edges between genes. The testing procedure asymptotically controls the

false discovery rate (FDR) under a pre-specified level, say 5%. See Suppl. Note 3 for details.

Step 2: Eliminating problematic low-count genes and corresponding edges. Some low-count genes exhibit spurious co-expressions with many other low-count genes, and these low-count genes form a spurious community. Thus, we remove gene i from $\mathcal{V}$ if it satisfies the following three criteria.

- Gene i has a median read count ≤ 2.
- Among all S_{ij} passed the multiple testing threshold (Suppl. Note 3), the positive ones take up more than 90%.
- Among all the genes in $\widetilde{\mathcal{N}}_i^+ = \{j : S_{ij} > 0, (i,j) \in \mathcal{E}\}$, those with the same median as gene i take up more than 90%.

Step 3. Identifying feature genes. We require feature genes to have $C_{1i} \geq 5/(p-1)$, $C_{2i} \geq 0.4$ and $C_{3i} \geq 0.3$. See Suppl. Note 4 for details about how we selected these default cutoffs.

Step 4. Identifying core feature genes. First, we define positive co-expression network $\mathcal{G}_+ = \{\mathcal{V}_+, \mathcal{E}_+\}$, where $\mathcal{V}_+$ contains the feature genes identified in Step 3, and $\mathcal{E}_+ = \mathcal{E} \cap \{(i,j) : i, j \in \mathcal{V}_+, S_{ij} > 0\}$. The 1st-, 2nd-, and 3rd-order connectivities for $\mathcal{G}_+$, denoted by $C_{+,1i}$, $C_{+,2i}$, and $C_{+,3i}$, are calculated correspondingly. The core feature genes are those with $C_{+,1i} \geq 5/(|\mathcal{V}_+|-1)$, $C_{+,2i} \geq 0.7$, and $C_{+,3i} \geq 0.5$. See Suppl. Fig. 7 for an illustration.

Step 5. Clustering core feature genes into gene sets. Taking the positive co-expression network restricted to the core feature genes $\mathcal{G}_{+,\text{core}}$, we apply the Louvain algorithm to cluster these genes into K gene sets, denoted by $\mathcal{V}_k$, where $k \in [K]$.

Step 6. Assigning non-core feature genes. Feature genes identified in Step 3 but not in Step 4 are referred to as non-core feature genes. These genes may be shared feature genes or genes with weak diffential-expression signals. Thus, we check their 1st-order connectivity to the core feature genes to decide their assignments. For any gene $i \in \mathcal{V}_+ \setminus (\cup_k \mathcal{V}_k)$, it is added to $\mathcal{V}_k$ if $|\mathcal{N}_i \cap \mathcal{V}_k|/|\mathcal{V}_k| \geq 40\%$. After this process, different feature gene sets may have shared feature genes.

Step 7. Augmenting feature gene sets with additional genes. Multi-role genes will not have the same connectivities as sole feature genes. Thus, they might not be identified as feature genes in Step 3. We add them to the feature gene sets in this step. For any $i \in \mathcal{V} \setminus \mathcal{V}_+$, it is added to $\mathcal{V}_k$ if $|\mathcal{N}_i \cap \mathcal{V}_k|/|\mathcal{V}_k| \geq 90\%$.

After these seven steps, the resulting gene sets $\mathcal{V}_k$, $k \in [K]$ is the K feature gene sets identified by SifiNet. Some steps involve setting tuning parameters, such as the gene set number K, the 1st-, 2nd-, and 3nd-order connectivity thresholds, and the assignment criteria for non-core feature genes, etc. The default values of these parameters are set based on our experiences on analyzing multiple experimental datasets. In Step 5, the Louvain algorithm is applied with the default resolution parameter. Because the core feature genes usually exhibit a clear community structure, different resolution parameters usually lead to the same cluster results.

SifiNet function 2: Annotating cells with feature gene sets

Based on $\mathcal{V}_k$, $k \in [K]$, we calculate the cellular gene set enrichment scores $\alpha_{c,1}, \ldots, \alpha_{c,K}$ for any cell c. For all the data analyses in this paper, we used the gene set variation analysis (GSVA)[47] scores. We define the cell subpopulation corresponding to $\mathcal{V}_k$ by

$$\mathcal{C}_k = \{c : \alpha_{c,k} > 0, \text{ and } \alpha_{c,k} \text{ is the largest among all } \alpha_{c,\ell}, \ \ell \in [K]\}.$$

Those cells with all negative $\alpha_{c,k}$ will be assigned to $\mathcal{C}_{K+1}$, called the unassigned group. These cells could be in the transitional stage or simply outliers. To annotate these unassigned cells, we can further apply DEG analysis to find their corresponding feature gene set.

To visualize the cell subpopulations, we derive the GSVA score matrix for all the cells and all the gene sets. Then we calculate the first two principal components (PCs) of the GSVA score matrix. The first two PCs are used to plot the cell subpopulations in a 2D space. See Figs. 3d, 3e, 4e, 4f for examples.

For data with complex heterogeneity structure, after we cluster cells based on the gene set enrichment scores, SifiNet can be applied again to a subset of cells to explore their local heterogeneity structure. By sequentially applying SifiNet, we can identify the hierarchical structure of cell subpopulations. See the BoneMarrow dataset analysis for an example.

SifiNet function 3: Constructing cell subpopulation network

By using the K feature gene sets as nodes, we can construct a gene-set-level graph representing the cell subpopulation lineage. There are a total of $K(K-1)/2$ possible edges. For edge (k, ℓ), its width is defined as

$$W_{k,\ell} = |\mathcal{E} \cap (i,j) : i \in \mathcal{V}_k, j \in \mathcal{V}_\ell| / |\mathcal{V}_k| \cdot |\mathcal{V}_\ell|.$$

The width indicates the relationship between the two corresponding gene sets. Finally, the gene set network can be transferred to cell subpopulation network after identifying the cell subpopulation corresponding to each feature gene set. Then, a wide edge between two cell subpopulations suggests possible transitional or developmental relationships between them. By default, edges with width less than 0.2 will be removed from the visualization figure.

Competing methods

We compared SifiNet with the two-step methods and HVG methods. The oracle-cluster methods were also introduced to explain why SifiNet work better than the two-step methods.

The two-step methods first use clustering methods to identify cell subpopulations, and then apply DEG analysis to identify cluster-specific DEGs. They are named after the DEG methods (such as DESeq2[5], edgeR[6,48,49], limma[7,50,51], limma-voom[8], and MAST[9,52]) and the clustering methods (such as Seurat[1,53–55], simple Louvain[2], and CIDR[3]), followed by TS. For example, DESeq2-Seurat-TS means that we first apply Seurat to cluster cells, and then use DESeq2 to identify cluster-specific DEGs. The two-step methods include DESeq2-Seurat-TS, edgeR-Seurat-TS, limma-Seurat-TS, limma-voom-Seurat-TS, MAST-Seurat-TS, DESeq2-Louvain-TS, edgeR-Louvain-TS, limma-Louvain-TS, limma-voom-Louvain-TS, MAST-Louvain-TS, DESeq2-CIDR-TS, edgeR-CIDR-TS, limma-CIDR-TS, limma-voom-CIDR-TS, MAST-CIDR-TS. The clustering methods are applied at default

resolution. The DEG methods are applied in similar ways as described in[56]. False discovery rate (FDR) is controlled at 0.05 with Benjamini-Hochberg (BH) procedure if there are p-values available in the method.

The oracle-cluster methods assume the ground truth of cell clusters, and they apply DEG analysis to identify cell-type-specific feature genes. The oracle-cluster methods include DESeq2-OC, edgeR-OC, limma-OC, limma-voom-OC, and MAST-OC.

The HVG identification methods identify feature genes through their deviation from the average gene properties. The HVG identification methods do not rely on knowing cell types or clustering. The methods we include in the comparisons are the following.

scry[57]**:** scry ranks genes by binomial deviance. Scry does not provide gene p-values; it ranks the genes based on the deviance scores. When we apply scry to the datasets in this paper, we selected the top p_1 genes where p_1 is the number of the actual feature genes.

M3Drop[58]**:** M3Drop finds feature genes by testing the relationship between mean expression level and dropout rate (proportion of zero observation) with Michaelis-Menten function.

singleCellHaystack[13]**:** singleCellHaystack detects feature genes using Kullback-Leibler divergence between observed distribution and reference distribution of each gene.

Moreover, we also compared SifiNet with Monocle3[33,37–41] in their performance to construct feature gene set networks. Monocle3 is comprehensive pipeline with a function to construct graph-based single-cell trajectories. With the input of the cell clusters, it can also construct feature gene set networks.

Simulated Data

We used the R package `GeneScape`[59] to generate two numerical dataset, SD1 and SD2. `GeneScape` can simulate scRNA-seq data with customized differential expression structures, such as core and shared feature genes, sub-cell types, pathways and hub genes.

SD1. SD1 contains 100 repetitions of simulated scRNA-seq data. Each data repetition includes 3 cell types with a total of 6000 cells. One of the cell types is rare (with 100 cells), sharing 60% (30 out of 50) of its feature genes with another non-rare cell type (with 4900 cells): these rare cells mimic those cells recently developed from the non-rare cell type. For preprocessing, we removed the cells with less than 10% genes expressed and genes expressed in less than 10% cells.

Complete analysis results of SD1 are shown in Suppl. Fig. 1. Because the two-step methods with CIDR have slightly better performance than those with Seurat (Suppl. Fig. 8), for all other analyses, we use the two-step methods with CIDR.

SD2. SD2 consists of 5 equal-sized main cell subpopulations, each containing 1200 cells. Among them, two are transitional, and the other three are ending. Transitional cell types shared feature genes with their neighboring ending cell types. The cell developmental lineage has the structure: ending-transition-ending-transition-ending. Each main transitional cell subpopulation further contains 3 subpopulations with the same feature genes but varying log

fold changes. For preprocessing, we removed the cells with less than 10% genes expressed and genes expressed in less than 10% cells.

Experimental Data

Monoclonal. The dataset was collected from a monoclonal murine colorectal carcinoma cell line CT26WT. The dataset details are given in Fang et al.[14]. We preprocessed the data in the following way. We removed the cells whose mitochondrial genes expression takes more than 5% of the total gene expression, the cells whose total number of read count is greater than 1.8 times of the mean expression levels, the cells with less than 10% of the genes expressed, and the genes expressed in less than 10% of the cells.

To curate a benchmark gene set to verify the identified genes, we did the followign steps. First, we collected human cell cycle related genes using 3 libraries, Reactome[60], Biocarta[61], KEGG[62,63], and 32 datasets from single studies or pathway analyses (Suppl. Table 4) with "CELLCYCLE" or "CELL_CYCLE" in their name from GSEA MSigDB C2 curated human gene sets v2023.1. Human genes that were included in the related pathway in at least 1 library or at least 3 studies were selected and converted to their mouse equivalents using `biomaRt` R package[64,65]. Second, for the Uniprot database, we downloaded the Uniprot gene annotations on May 1, 2023, and selected those genes whose "Biological process" annotations include at least 1 of the following keywords: mitotic, centriole, centrosome, centromere, spindle, chromosome separation, cohesin, DNA replication, DNA repair, microtubule, nuclear chromosome segregation, cell cycle, cell division, cell growth. Finally, the intersection of selected genes in these two databases were chosen as the benchmark cell cycle set (475 genes). Among them, 437 genes remained in the Monoclonal data after preprocessing.

IPF1 and IPF2. These two datasets were collected from the existing study on idiopathic pulmonary fibrosis (IPF)[15]. IPF1 contains the pre-annotated "Ciliate" cells in IPF patients. IPF2 contains the pre-annotated "KRT5-/KRT17+" epithelial cells and the "Proliferating Epithelial Cells". For pre-processing, we removed the cells with less than 5% genes expressed, and the genes expressed in less than 5% cells. Then, cell counts were normalized by `SCTransform` in Seurat.

To call the SifiNet-senescent cells in IPF1, we calculated the GSVA scores of the three gene sets. Those cells with top/bottom 40% GSVA scores of all three gene sets SifiNet-senescent cells: top scores were used for up-regulated gene set (Gene Set 2), bottom scores were used for down-regulated gene sets (Gene sets 1 and 3). A total of 594 cells are annotated as SifiNet-senescent cells.

To call the SenMayo-senescent cells in IPF1, we chose 594 cells (same number of cells as SifiNet-senescent cells) with the top SenMayo-GSVA scores.

BoneMarrow. BoneMarrow was collected from existing study on myeloid differentiation[16]. We focus on the pre-annotated "Unsorted myeloid" cells. We used `SCTransform` to normalize the cell UMI counts and remove the batch effect. Then, we removed the cells with fewer than 500 genes expressed, and the genes expressed in fewer than 2% of the cells.

SifiNet identified two feature gene sets in BoneMarrow, S1 and S2 (Suppl. Table 3). S1 contains markers for lineages in the granulocyte/macrophage progenitors[66,67], such as

Cebpa. S2 contains markers for megakaryocyte/erythrocyte progenitors[68–70], such as *Gata1*, *Gfi1b*, and *Klf1* (Fig. 3f). The cells with positive S1 (S2) GSVA scores and negative S2 (S1) GSVA scores were clustered into granulocyte/macrophage progenitor cells (megakaryocyte/erythrocyte progenitor cells); those cells with both negative S1 and S2 GSVA scores were clustered into early progenitor cells. Early progenitor cell markers such as *Apoe* and *Gata2*[16] are significantly highly expressed in the early progenitor cells (*Apoe*: log2 fold change 1.75, adjusted p-value= 1.19×10^{-169}; *Gata2*: log2 fold change 0.887, adjusted p-value= 2.77×10^{-59}).

When applied to the annotated granulocyte/macrophage progenitor cells, SifiNet identified 3 feature gene sets S1a-S1c (Suppl. Table 3), dividing the cells into 4 subpopulations: granulocyte progenitor cells (corresponding to S1a), macrophage progenitor cells (corresponding to S1b), monocyte and dendritic progenitor cells (corresponding to S1c), and unassigned. DEG analysis shows that hematopoietic stem cell marker gene, *CD34*, and memory B cell and T cell marker gene, *CD27*[71–74], are significantly over-expressed in the unassigned subpopulation. Thus, we label this subpopulation as lymphoid progenitor cells (Fig. 3e).

CD8. CD8 is a subset of a 10x scRNA-seq and scATAC-seq multiome datasets[17]. An annotation of cell types is provided by Seurat. CD8 dataset contains cells with "CD8 Naïve", "CD8 TEM_1" and "CD8 TEM_2". In scATAC-seq data, peaks within the 5000-base-pair upstream of the transcription starting site (TSS) of genes were aggregated to represent the epigenomic features of the corresponding genes. Only genes available in both scRNA-seq data and scATAC-seq data were kept. In scRNA-seq data, cells with total read count greater than 1.8 times of the mean total read count were removed. In scATAC-seq data, cells with total epigenomic feature count greater than 1.8 times of the mean total epigenomic feature count were removed. We further removed cells with fewer than 2% of the genes detected in either scRNA-seq or scATAC-seq, and then the genes expressed in fewer than 2% of the cells in either scRNA-seq or scATAC-seq.

When applied to the CD8 scRNA-seq data, SifiNet identified four gene sets. Set 1 and Set 3 share *Gzma*, *Klrd1*, and *Klrg1*, markers for effector T cells[75,76]. Set 2 contains many central memory T cell (TCM) markers, including *Foxo1*, *Ccr7*, *Foxp1*, and *Tcf7*[77,78]. Set 4 contains 19 Rpl and Rps genes, which are highly expressed in Naïve T cells[79]. Then, SifiNet clustered the cells into five subpopulations, TEM_1, TCM_1, TEM_2, TNaïve, and unassigned (the remaining cells), which are between the TCM_1 and TEM_2 cells (Fig. 4e). DEG analysis shows that the expression levels of TEM and TCM markers, *Gzma*, *Klrd1*, *Klrg1*, *Foxo1*, *Ccr7*, *Foxp1*, and *Tcf7*, in the unassigned subpopulation exhibit greater similarity to the TCM cells (Suppl. Fig. 9). Thus, we annotate the unassigned subpopulation as "TCM_2".

When applied to the scATAC-seq data, SifiNet identified two feature gene sets. Set 1 includes TCM markers, such as *Klf7* and *Foxp1*[78]. Set 2 includes TEM markers, such as *Klrg1*[75]. These gene sets separated the cells into TCM and TEM cells.

MassiveRNA. MassiveRNA is a large dataset with more than 1 million cells[18]. We removed those cells with fewer than 2% of the genes expressed, and then those genes expressed in fewer than 2% of the cells. After that, 1287071 cells and 10448 genes remain in the dataset. SifiNet was applied to the whole dataset and downsampled datasets with number of

cells 10000, 20000, 50000, 100000, 200000, 400000, 600000, 800000, and 1000000. The time consumption was evaluated using the total elapsed time when running SifiNet on the dataset, and the peak memory usage was evaluated using the `gc` function in R.

Data availability

Monoclonal dataset is available at the Genome Sequencing Archive (GSA) with accession CRA008966. IPF1 and IPF2 datasets are collected from study GSE135893. Bone-Marrow dataset is collected from study GSE72857. CD8 dataset is available at 10x genomics website `https://support.10xgenomics.com/single-cell-multiome-atac-gex/datasets/1.0.0/pbmc_granulocyte_sorted_10k`, and is also included in SeuratData R package as pbmc.rna and pbmc.atac. Seurat annotations for CD8 are available at SeuratData R package v0.2.2. MassiveRNA is downloaded from 10x genomics website `https://support.10xgenomics.com/single-cell-gene-expression/datasets/1.3.0/1M_neurons`.

Code availability

SifiNet v1.9 is used for the analysis in this study. SifiNet package is available at `https://github.com/jichunxie/SifiNet`. The codes used for generating the results shown in this paper are available at `https://github.com/jichunxie/SifiNet_manu_support`. GeneScape v1.0 used for generating numerical datasets is available on R CRAN.

Acknowledgement

The authors thank the Duke CHSI for providing the computation resources.

Funding

QG, ZJ, CC, and JX's research was partially supported by NIH Common Fund, through the Office of Strategic Coordination/Office of the NIH Director under awards, 1U54AG075936-01 from the National Institute of Aging. LW's research was supported by Duke University. QJL's research was supported by NIH Award R33 CA225328 (early stage of this project) and core funds from IMCB and SIgN, A*STAR. In addition, QG, KO and JX's research was partially supported by the NIH Award 1R01HG012555-01.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

[1] Rahul Satija, Jeffrey A Farrell, David Gennert, Alexander F Schier, and Aviv Regev. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol*, 33(5):495–502, May 2015. doi: 10.1038/nbt.3192. PMCID: PMC4430369.

[2] Vincent D Blondel, Jean-Loup Guillaume, Renaud Lambiotte, and Etienne Lefebvre. Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment*, 2008(10):10008, October 2008. doi: 10.1088/1742-5468/2008/10/P10008. URL `https://dx.doi.org/10.1088/1742-5468/2008/10/P10008`.

[3] Peijie Lin, Michael Troup, and Joshua W K Ho. Cidr: Ultrafast and accurate clustering through imputation for single-cell rna-seq data. *Genome Biol*, 18(1):59, Mar 2017. doi: 10.1186/s13059-017-1188-0. PMCID: PMC5371246.

[4] F Alexander Wolf, Philipp Angerer, and Fabian J Theis. Scanpy: large-scale single-cell gene expression data analysis. *Genome Biol*, 19(1):15, Feb 2018. doi: 10.1186/s13059-017-1382-0. PMCID: PMC5802054.

[5] Michael I Love, Wolfgang Huber, and Simon Anders. Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome Biol*, 15(12):550, 2014. doi: 10.1186/s13059-014-0550-8. PMCID: PMC4302049.

[6] Mark D Robinson, Davis J McCarthy, and Gordon K Smyth. edger: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1):139–40, Jan 2010. doi: 10.1093/bioinformatics/btp616. PMCID: PMC2796818.

[7] Matthew E Ritchie, Belinda Phipson, Di Wu, Yifang Hu, Charity W Law, Wei Shi, and Gordon K Smyth. limma powers differential expression analyses for rna-sequencing and microarray studies. *Nucleic Acids Res*, 43(7):e47, Apr 2015. doi: 10.1093/nar/gkv007. PMCID: PMC4402510.

[8] Charity W Law, Yunshun Chen, Wei Shi, and Gordon K Smyth. voom: Precision weights unlock linear model analysis tools for rna-seq read counts. *Genome Biol*, 15(2):R29, Feb 2014. doi: 10.1186/gb-2014-15-2-r29. PMCID: PMC4053721.

[9] Andrew McDavid, Greg Finak, and Masanao Yajima. *MAST: Model-based Analysis of Single Cell Transcriptomics*, 2021. URL https://github.com/RGLab/MAST/. R package version 1.20.0.

[10] Lucy L. Gao, Jacob Bien, and Daniela Witten. Selective inference for hierarchical clustering. *Journal of the American Statistical Association*, pages 1–11, Aug 2022. doi: 10.1080/01621459.2022.2116331. URL https://doi.org/10.1080/01621459.2022.2116331.

[11] F William Townes, Stephanie C Hicks, Martin J Aryee, and Rafael A Irizarry. Feature selection and dimension reduction for single-cell rna-seq based on a multinomial model. *Genome Biol*, 20(1):295, Dec 2019. doi: 10.1186/s13059-019-1861-6. PMCID: PMC6927135.

[12] Tallulah S Andrews and Martin Hemberg. M3drop: dropout-based feature selection for scrnaseq. *Bioinformatics*, 35(16):2865–2867, Aug 2019. doi: 10.1093/bioinformatics/bty1044. PMCID: PMC6691329.

[13] Alexis Vandenbon and Diego Diez. A clustering-independent method for finding differentially expressed genes in single-cell transcriptome data. *Nat Commun*, 11(1):4318, Aug 2020. doi: 10.1038/s41467-020-17900-3. PMCID: PMC7455704.

[14] Jiyuan Fang, Cliburn Chan, Kouros Owzar, Liuyang Wang, Diyuan Qin, Qi-Jing Li, and Jichun Xie. Clustering deviation index (cdi): a robust and accurate internal measure for

evaluating scrna-seq data clustering. *Genome Biol*, 23(1):269, Dec 2022. doi: 10.1186/ s13059-022-02825-5. PMCID: PMC9793368.

[15] Arun C Habermann, Austin J Gutierrez, Linh T Bui, Stephanie L Yahn, Nichelle I Winters, Carla L Calvi, Lance Peter, Mei-I Chung, Chase J Taylor, Christopher Jetter, Latha Raju, Jamie Roberson, Guixiao Ding, Lori Wood, Jennifer M S Sucre, Bradley W Richmond, Ana P Serezani, Wyatt J McDonnell, Simon B Mallal, Matthew J Bacchetta, James E Loyd, Ciara M Shaver, Lorraine B Ware, Ross Bremner, Rajat Walia, Timothy S Blackwell, Nicholas E Banovich, and Jonathan A Kropski. Single-cell rna sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in pulmonary fibrosis. *Sci Adv*, 6(28):eaba1972, Jul 2020. doi: 10.1126/sciadv.aba1972. PMCID: PMC7439444.

[16] Franziska Paul, Ya'ara Arkin, Amir Giladi, Diego Adhemar Jaitin, Ephraim Kenigsberg, Hadas Keren-Shaul, Deborah Winter, David Lara-Astiaso, Meital Gury, Assaf Weiner, Eyal David, Nadav Cohen, Felicia Kathrine Bratt Lauridsen, Simon Haas, Andreas Schlitzer, Alexander Mildner, Florent Ginhoux, Steffen Jung, Andreas Trumpp, Bo Torben Porse, Amos Tanay, and Ido Amit. Transcriptional heterogeneity and lineage commitment in myeloid progenitors. *Cell*, 163(7):1663–77, Dec 2015. doi: 10.1016/j.cell.2015.11.013. PMID: 26627738.

[17] 10x Genomics. Pbmc from a healthy donor - granulocytes removed through cell sorting (10k), 2020. Single Cell Multiome ATAC + Gene Exp. Dataset by Cell Ranger ARC 1.0.0.

[18] 10x Genomics. 1.3 million brain cells from e18 mice, 2017. URL `https://support. 10xgenomics.com/single-cell-gene-expression/datasets/1.3.0/1M_neurons`. Single Cell Gene Expression Dataset by Cell Ranger 1.3.0.

[19] Yingmei Wang, Ping Ji, Jinsong Liu, Russell R Broaddus, Fengxia Xue, and Wei Zhang. Centrosome-associated regulators of the g(2)/m checkpoint as targets for cancer therapy. *Mol Cancer*, 8:8, Feb 2009. doi: 10.1186/1476-4598-8-8. PMCID: PMC2657106.

[20] Arianne Heinrichs. A centrosome-integrity checkpoint. *Nature Reviews Molecular Cell Biology*, 8(2):98–98, February 2007. doi: 10.1038/nrm2107.

[21] Aravind Subramanian, Pablo Tamayo, Vamsi K Mootha, Sayan Mukherjee, Benjamin L Ebert, Michael A Gillette, Amanda Paulovich, Scott L Pomeroy, Todd R Golub, Eric S Lander, and Jill P Mesirov. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*, 102(43): 15545–50, Oct 2005. doi: 10.1073/pnas.0506580102. PMCID: PMC1239896.

[22] Arthur Liberzon, Aravind Subramanian, Reid Pinchback, Helga Thorvaldsdóttir, Pablo Tamayo, and Jill P Mesirov. Molecular signatures database (msigdb) 3.0. *Bioinformatics*, 27(12):1739–40, Jun 2011. doi: 10.1093/bioinformatics/btr260. PMCID: PMC3106198.

[23] Arthur Liberzon, Chet Birger, Helga Thorvaldsdóttir, Mahmoud Ghandi, Jill P Mesirov, and Pablo Tamayo. The molecular signatures database (msigdb) hallmark gene set

collection. *Cell Syst*, 1(6):417–425, Dec 2015. doi: 10.1016/j.cels.2015.12.004. PMCID: PMC4707969.

[24] UniProt Consortium. Uniprot: the universal protein knowledgebase in 2023. *Nucleic Acids Res*, 51(D1):D523–D531, Jan 2023. doi: 10.1093/nar/gkac1052. PMCID: PMC9825514.

[25] D L Kellogg, D L Kellogg, Jr, N Musi, and A M Nambiar. Cellular senescence in idiopathic pulmonary fibrosis. *Curr Mol Biol Rep*, 7(3):31–40, August 2021. doi: 10.1007/s40610-021-00145-4. PMCID: PMC8358258.

[26] Kaj E C Blokland, David W Waters, Michael Schuliga, Jane Read, Simon D Pouwels, Christopher L Grainge, Jade Jaffar, Glen Westall, Steven E Mutsaers, Cecilia M Prêle, Janette K Burgess, and Darryl A Knight. Senescence of ipf lung fibroblasts disrupt alveolar epithelial cell proliferation and promote migration in wound healing. *Pharmaceutics*, 12(4), Apr 2020. doi: 10.3390/pharmaceutics12040389. PMCID: PMC7238173.

[27] Frédéric Lessard, Sebastian Igelmann, Christian Trahan, Geneviève Huot, Emmanuelle Saint-Germain, Lian Mignacca, Neylen Del Toro, Stéphane Lopes-Paciencia, Benjamin Le Calvé, Marinieve Montero, Xavier Deschênes-Simard, Marina Bury, Olga Moiseeva, Marie-Camille Rowell, Cornelia E Zorca, Daniel Zenklusen, Léa Brakier-Gingras, Véronique Bourdeau, Marlene Oeffinger, and Gerardo Ferbeyre. Senescence-associated ribosome biogenesis defects contributes to cell cycle arrest through the rb pathway. *Nat Cell Biol*, 20(7):789–799, Jul 2018. doi: 10.1038/s41556-018-0127-y. PMID: 29941930.

[28] Ines Garces de Los Fayos Alonso, Huan-Chang Liang, Suzanne D Turner, Sabine Lagger, Olaf Merkel, and Lukas Kenner. The role of activator protein-1 (ap-1) family members in cd30-positive lymphomas. *Cancers (Basel)*, 10(4), Mar 2018. doi: 10.3390/cancers10040093. PMCID: PMC5923348.

[29] Yuting Wang, Liping Liu, Yifan Song, Xiaojie Yu, and Hongkui Deng. Unveiling e2f4, tead1 and ap-1 as regulatory transcription factors of the replicative senescence program by multi-omics analysis. *Protein Cell*, 13(10):742–759, Oct 2022. doi: 10.1007/s13238-021-00894-z. PMCID: PMC9233726.

[30] Luc Rochette, Marianne Zeller, Yves Cottin, and Catherine Vergely. Insights into mechanisms of gdf15 and receptor gfral: Therapeutic targets. *Trends Endocrinol Metab*, 31 (12):939–951, Dec 2020. doi: 10.1016/j.tem.2020.10.004. PMID: 33172749.

[31] Hélène Martini and João F Passos. Cellular senescence: all roads lead to mitochondria. *FEBS J*, 290(5):1186–1202, Mar 2023. doi: 10.1111/febs.16361. PMCID: PMC9296701.

[32] Ruchi Kumari and Parmjit Jat. Mechanisms of cellular senescence: Cell cycle arrest and senescence associated secretory phenotype. *Front Cell Dev Biol*, 9:645593, March 2021. doi: 10.3389/fcell.2021.645593. PMCID: PMC8039141.

[33] Leland McInnes, John Healy, and James Melville. Umap: Uniform manifold approximation and projection for dimension reduction, September 2020.

[34] Monika S Kowalczyk, Itay Tirosh, Dirk Heckl, Tata Nageswara Rao, Atray Dixit, Brian J Haas, Rebekka K Schneider, Amy J Wagers, Benjamin L Ebert, and Aviv Regev. Single-cell rna-seq reveals changes in cell cycle and differentiation programs upon aging of hematopoietic stem cells. *Genome Res*, 25(12):1860–72, Dec 2015. doi: 10.1101/gr. 192237.115. PMCID: PMC4665007.

[35] José Alberto López-Domínguez, Sandra Rodríguez-López, Ulises Ahumada-Castro, Pierre-Yves Desprez, Maria Konovalenko, Remi-Martin Laberge, César Cárdenas, José Manuel Villalba, and Judith Campisi. Cdkn1a transcript variant 2 is a marker of aging and cellular senescence. *Aging (Albany NY)*, 13(10):13380–13392, May 2021. doi: 10.18632/aging.203110. PMCID: PMC8202863.

[36] Dominik Saul, Robyn Laura Kosinsky, Elizabeth J Atkinson, Madison L Doolittle, Xu Zhang, Nathan K LeBrasseur, Robert J Pignolo, Paul D Robbins, Laura J Niedernhofer, Yuji Ikeno, Diana Jurk, João F Passos, LaTonya J Hickson, Ailing Xue, David G Monroe, Tamara Tchkonia, James L Kirkland, Joshua N Farr, and Sundeep Khosla. A new gene set identifies senescent cells and predicts senescence-associated pathways across tissues. *Nat Commun*, 13(1):4827, Aug 2022. doi: 10.1038/s41467-022-32552-1. PMCID: PMC9381717.

[37] Cole Trapnell, Davide Cacchiarelli, Jonna Grimsby, Prapti Pokharel, Shuqiang Li, Michael Morse, Niall J Lennon, Kenneth J Livak, Tarjei S Mikkelsen, and John L Rinn. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol*, 32(4):381–386, Apr 2014. doi: 10.1038/nbt.2859. PMCID: PMC4122333.

[38] Xiaojie Qiu, Qi Mao, Ying Tang, Li Wang, Raghav Chawla, Hannah A Pliner, and Cole Trapnell. Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods*, 14(10):979–982, Oct 2017. doi: 10.1038/nmeth.4402. PMCID: PMC5764547.

[39] Junyue Cao, Malte Spielmann, Xiaojie Qiu, Xingfan Huang, Daniel M Ibrahim, Andrew J Hill, Fan Zhang, Stefan Mundlos, Lena Christiansen, Frank J Steemers, Cole Trapnell, and Jay Shendure. The single-cell transcriptional landscape of mammalian organogenesis. *Nature*, 566(7745):496–502, Feb 2019. doi: 10.1038/s41586-019-0969-x. PMCID: PMC6434952.

[40] V A Traag, L Waltman, and N J van Eck. From louvain to leiden: guaranteeing well-connected communities. *Sci Rep*, 9(1):5233, Mar 2019. doi: 10.1038/s41598-019-41695-z. PMCID: PMC6435756.

[41] Jacob H Levine, Erin F Simonds, Sean C Bendall, Kara L Davis, El-ad D Amir, Michelle D Tadmor, Oren Litvin, Harris G Fienberg, Astraea Jager, Eli R Zunder, Rachel Finck, Amanda L Gedman, Ina Radtke, James R Downing, Dana Pe'er, and Garry P Nolan. Data-driven phenotypic dissection of aml reveals progenitor-like cells that correlate with prognosis. *Cell*, 162(1):184–97, Jul 2015. doi: 10.1016/j.cell.2015.05.047. PMCID: PMC4508757.

[42] Christopher D Scharer, Alexander P R Bally, Bhanu Gandham, and Jeremy M Boss. Cutting edge: Chromatin accessibility programs cd8 t cell memory. *J Immunol*, 198(6): 2238–2243, Mar 2017. doi: 10.4049/jimmunol.1602086. PMCID: PMC5341694.

[43] Zhilin Long, Chengfang Sun, Min Tang, Yin Wang, Jiayan Ma, Jichuan Yu, Jingchao Wei, Jianzhu Ma, Bohan Wang, Qi Xie, and Jiaming Wen. Single-cell multiomics analysis reveals regulatory programs in clear cell renal cell carcinoma. *Cell Discov*, 8(1):68, Jul 2022. doi: 10.1038/s41421-022-00415-0. PMCID: PMC9296597.

[44] James R Rose, Bagdeser Akdogan-Ozdilek, Andrew R Rahmberg, Michael D Powell, Sakeenah L Hicks, Christopher D Scharer, and Jeremy M Boss. Distinct transcriptomic and epigenomic modalities underpin human memory t cell subsets and their activation potential. *Commun Biol*, 6(1):363, Apr 2023. doi: 10.1038/s42003-023-04747-9. PMCID: PMC10070634.

[45] Rama S Akondy, Mark Fitch, Srilatha Edupuganti, Shu Yang, Haydn T Kissick, Kelvin W Li, Ben A Youngblood, Hossam A Abdelsamed, Donald J McGuire, Kristen W Cohen, Gabriela Alexe, Shashi Nagar, Megan M McCausland, Satish Gupta, Pramila Tata, W Nicholas Haining, M Juliana McElrath, David Zhang, Bin Hu, William J Greenleaf, Jorg J Goronzy, Mark J Mulligan, Marc Hellerstein, and Rafi Ahmed. Origin and differentiation of human memory cd8 t cells after vaccination. *Nature*, 552(7685): 362–367, Dec 2017. doi: 10.1038/nature24633. PMCID: PMC6037316.

[46] Jichun Xie and Ruosha Li. False discovery rate control for high dimensional networks of quantile associations conditioning on covariates. *J R Stat Soc Series B Stat Methodol*, 80(5):1015–1034, Nov 2018. doi: 10.1111/rssb.12288. PMCID: PMC6497089.

[47] Sonja Hänzelmann, Robert Castelo, and Justin Guinney. Gsva: gene set variation analysis for microarray and rna-seq data. *BMC Bioinformatics*, 14:7, Jan 2013. doi: 10.1186/1471-2105-14-7. PMCID: PMC3618321.

[48] Davis J McCarthy, Yunshun Chen, and Gordon K Smyth. Differential expression analysis of multifactor rna-seq experiments with respect to biological variation. *Nucleic Acids Res*, 40(10):4288–97, May 2012. doi: 10.1093/nar/gks042. PMCID: PMC3378882.

[49] Yunshun Chen, Aaron T L Lun, and Gordon K Smyth. From reads to genes to pathways: differential expression analysis of rna-seq experiments using rsubread and the edger quasi-likelihood pipeline. *F1000Res*, 5:1438, June 2016. doi: 10.12688/f1000research. 8987.2. PMCID: PMC4934518.

[50] Belinda Phipson, Stanley Lee, Ian J Majewski, Warren S Alexander, and Gordon K Smyth. Robust hyperparameter estimation protects against hypervariable genes and improves power to detect differential expression. *Ann Appl Stat*, 10(2):946–963, Jun 2016. doi: 10.1214/16-AOAS920. PMCID: PMC5373812.

[51] Gordon K Smyth. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol*, 3:Article3, February 2004. doi: 10.2202/1544-6115.1027. PMID: 16646809.

[52] Greg Finak, Andrew McDavid, Masanao Yajima, Jingyuan Deng, Vivian Gersuk, Alex K Shalek, Chloe K Slichter, Hannah W Miller, M Juliana McElrath, Martin Prlic, Peter S Linsley, and Raphael Gottardo. Mast: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell rna sequencing data. *Genome Biol*, 16:278, Dec 2015. doi: 10.1186/s13059-015-0844-5. PMCID: PMC4676162.

[53] Yuhan Hao, Stephanie Hao, Erica Andersen-Nissen, William M Mauck, 3rd, Shiwei Zheng, Andrew Butler, Maddie J Lee, Aaron J Wilk, Charlotte Darby, Michael Zager, Paul Hoffman, Marlon Stoeckius, Efthymia Papalexi, Eleni P Mimitou, Jaison Jain, Avi Srivastava, Tim Stuart, Lamar M Fleming, Bertrand Yeung, Angela J Rogers, Juliana M McElrath, Catherine A Blish, Raphael Gottardo, Peter Smibert, and Rahul Satija. Integrated analysis of multimodal single-cell data. *Cell*, 184(13):3573–3587.e29, Jun 2021. doi: 10.1016/j.cell.2021.04.048. PMCID: PMC8238499.

[54] Tim Stuart, Andrew Butler, Paul Hoffman, Christoph Hafemeister, Efthymia Papalexi, William M Mauck, 3rd, Yuhan Hao, Marlon Stoeckius, Peter Smibert, and Rahul Satija. Comprehensive integration of single-cell data. *Cell*, 177(7):1888–1902.e21, Jun 2019. doi: 10.1016/j.cell.2019.05.031. PMCID: PMC6687398.

[55] Andrew Butler, Paul Hoffman, Peter Smibert, Efthymia Papalexi, and Rahul Satija. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*, 36(5):411–420, Jun 2018. doi: 10.1038/nbt.4096. PMCID: PMC6700744.

[56] Charlotte Soneson and Mark D Robinson. Bias, robustness and scalability in single-cell differential expression analysis. *Nat Methods*, 15(4):255–261, Apr 2018. doi: 10.1038/nmeth.4612. PMID: 29481549.

[57] Kelly Street, F. William Townes, Davide Risso, and Stephanie Hicks. *scry: Small-Count Analysis Methods for High-Dimensional Data*, 2021. URL https://bioconductor.org/packages/scry.html. R package version 1.6.0.

[58] Tallulah Andrews. *M3Drop: Michaelis-Menten Modelling of Dropouts in single-cell RNASeq*, 2021. URL https://github.com/tallulandrews/M3Drop. R package version 1.20.0.

[59] Qi Gao. Genescape: Simulation of single cell rna-seq data with complex structure, 2023. URL https://CRAN.R-project.org/package=GeneScape. R package version 1.0.

[60] Marc Gillespie, Bijay Jassal, Ralf Stephan, Marija Milacic, Karen Rothfels, Andrea Senff-Ribeiro, Johannes Griss, Cristoffer Sevilla, Lisa Matthews, Chuqiao Gong, Chuan Deng, Thawfeek Varusai, Eliot Ragueneau, Yusra Haider, Bruce May, Veronica Shamovsky, Joel Weiser, Timothy Brunson, Nasim Sanati, Liam Beckman, Xiang Shao, Antonio Fabregat, Konstantinos Sidiropoulos, Julieth Murillo, Guilherme Viteri, Justin Cook, Solomon Shorser, Gary Bader, Emek Demir, Chris Sander, Robin Haw, Guanming Wu, Lincoln Stein, Henning Hermjakob, and Peter D'Eustachio. The reactome

pathway knowledgebase 2022. *Nucleic Acids Res*, 50(D1):D687–D692, Jan 2022. doi: 10.1093/nar/gkab1028. PMCID: PMC8689983.

[61] Darryl Nishimura. Biocarta. *Biotech Software & Internet Report*, 2(3):117–120, 2023/05/17 2001. doi: 10.1089/152791601750294344. URL https://doi.org/10.1089/152791601750294344.

[62] M Kanehisa and S Goto. Kegg: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res*, 28(1):27–30, Jan 2000. doi: 10.1093/nar/28.1.27. PMCID: PMC102409.

[63] Minoru Kanehisa. Toward understanding the origin and evolution of cellular organisms. *Protein Sci*, 28(11):1947–1951, Nov 2019. doi: 10.1002/pro.3715. PMCID: PMC6798127.

[64] Steffen Durinck, Paul T Spellman, Ewan Birney, and Wolfgang Huber. Mapping identifiers for the integration of genomic datasets with the r/bioconductor package biomart. *Nat Protoc*, 4(8):1184–91, 2009. doi: 10.1038/nprot.2009.97. PMCID: PMC3159387.

[65] Steffen Durinck, Yves Moreau, Arek Kasprzyk, Sean Davis, Bart De Moor, Alvis Brazma, and Wolfgang Huber. Biomart and bioconductor: a powerful link between biological databases and microarray data analysis. *Bioinformatics*, 21(16):3439–40, Aug 2005. doi: 10.1093/bioinformatics/bti525. PMID: 16082012.

[66] Richard Dahl, Jonathan C Walsh, David Lancki, Peter Laslo, Sangeeta R Iyer, Harinder Singh, and M Celeste Simon. Regulation of macrophage and neutrophil cell fates by the pu.1:c/ebpalpha ratio and granulocyte colony-stimulating factor. *Nat Immunol*, 4(10): 1029–36, Oct 2003. doi: 10.1038/ni973. PMID: 12958595.

[67] Catherine V Laiosa, Matthias Stadtfeld, Huafeng Xie, Luisa de Andres-Aguayo, and Thomas Graf. Reprogramming of committed t cell progenitors to macrophages and dendritic cells by c/ebp alpha and pu.1 transcription factors. *Immunity*, 25(5):731–44, Nov 2006. doi: 10.1016/j.immuni.2006.09.011. PMID: 17088084.

[68] Hiromi Iwasaki, Shin-ichi Mizuno, Richard A Wells, Alan B Cantor, Sumiko Watanabe, and Koichi Akashi. Gata-1 converts lymphoid and myelomonocytic progenitors into the megakaryocyte/erythrocyte lineages. *Immunity*, 19(3):451–62, Sep 2003. doi: 10.1016/s1074-7613(03)00242-5. PMID: 14499119.

[69] Lothar Vassen, Hugues Beauchemin, Wafaa Lemsaddek, Joseph Krongold, Marie Trudel, and Tarik Möröy. Growth factor independence 1b (gfi1b) is important for the maturation of erythroid cells and the regulation of embryonic globin expression. *PLoS One*, 9(5): e96636, May 2014. doi: 10.1371/journal.pone.0096636. PMCID: PMC4011847.

[70] Cheng-Tao Yang, Rui Ma, Richard A Axton, Melany Jackson, A Helen Taylor, Antonella Fidanza, Lamin Marenah, Jan Frayne, Joanne C Mountford, and Lesley M Forrester. Activation of klf1 enhances the differentiation and maturation of red blood cells from human pluripotent stem cells. *Stem Cells*, 35(4):886–897, Apr 2017. doi: 10.1002/stem.2562. PMCID: PMC5396323.

[71] Sara E Vazquez, Matthew A Inlay, and Thomas Serwold. Cd201 and cd27 identify hematopoietic stem and progenitor cells across multiple murine strains independently of kit and sca-1. *Exp Hematol*, 43(7):578–85, Jul 2015. doi: 10.1016/j.exphem.2015.04.001. PMCID: PMC4480781.

[72] J Hendriks, L A Gravestein, K Tesselaar, R A van Lier, T N Schumacher, and J Borst. Cd27 is required for generation and long-term maintenance of t cell immunity. *Nat Immunol*, 1(5):433–40, Nov 2000. doi: 10.1038/80877. PMID: 11062504.

[73] K Agematsu, S Hokibara, H Nagumo, and A Komiyama. Cd27: a memory b-cell marker. *Immunol Today*, 21(5):204–6, May 2000. doi: 10.1016/s0167-5699(00)01605-4. PMID: 10782048.

[74] C Schmitt, S Ktorza, S Sarun, M P Verpilleux, C Blanc, M A Deugnier, A Dalloul, and P Debré. Cd34-positive early stages of human t-cell differentiation. *Leuk Lymphoma*, 17(1-2):43–50, Mar 1995. doi: 10.3109/10428199509051702. PMID: 7539659.

[75] Dietmar Herndler-Brandstetter, Harumichi Ishigame, Ryo Shinnakasu, Valerie Plajer, Carmen Stecher, Jun Zhao, Melanie Lietzenmayer, Lina Kroehling, Akiko Takumi, Kohei Kometani, Takeshi Inoue, Yuval Kluger, Susan M Kaech, Tomohiro Kurosaki, Takaharu Okada, and Richard A Flavell. Klrg1+ effector cd8+ t cells lose klrg1, differentiate into all memory t cell lineages, and convey enhanced protective immunity. *Immunity*, 48(4): 716–729.e8, Apr 2018. doi: 10.1016/j.immuni.2018.03.015. PMCID: PMC6465538.

[76] Trine Sundebo Meldgaard, Fabiola Blengio, Denise Maffione, Chiara Sammicheli, Simona Tavarini, Sandra Nuti, Roland Kratzer, Duccio Medini, Emilio Siena, and Sylvie Bertholet. Single-cell analysis of antigen-specific cd8+ t-cell transcripts reveals profiles specific to mrna or adjuvanted protein vaccines. *Front Immunol*, 12:757151, 2021. doi: 10.3389/fimmu.2021.757151. PMCID: PMC8586650.

[77] Myoungjoo V Kim, Weiming Ouyang, Will Liao, Michael Q Zhang, and Ming O Li. The transcription factor foxo1 controls central-memory cd8+ t cell responses to infection. *Immunity*, 39(2):286–97, Aug 2013. doi: 10.1016/j.immuni.2013.07.013. PMCID: PMC3809840.

[78] Hua Lu, Hui Wang, Lijun Yan, Hongwei Shao, Wenfeng Zhang, Han Shen, Huaben Bo, Changli Tao, Shengfang Xia, and Fenglin Wu. Overexpression of early t cell differentiation-specific transcription factors transforms the terminally differentiated effector t cells into less differentiated state. *Cell Immunol*, 353:104118, Jul 2020. doi: 10.1016/j.cellimm.2020.104118. PMID: 32413598.

[79] Laura S Christian, Liuyang Wang, Bryan Lim, Dachuan Deng, Haiyang Wu, Xiao-Fan Wang, and Qi-Jing Li. Resident memory t cells in tumor-distant tissues fortify against metastasis formation. *Cell Rep*, 35(6):109118, May 2021. doi: 10.1016/j.celrep.2021.109118. PMCID: PMC8204287.