Title:

Pupil size predicts the onset of exploration and changes in prefrontal dynamics

Authors:

Akram Shourkeshti[1†], Mojtaba Abbaszadeh[1†], Gabriel Marrocco[1], Katarzyna Jurewicz[1,2], Tirin
Moore[3,4], R. Becket Ebitz*[1]

Affiliations:

[1]Département de Neurosciences, Université de Montréal, Montréal, QC, Canada

[2]Department of Physiology, McGill University, Montréal, QC, Canada

[3]Department of Neurobiology, Stanford University School of Medicine, Stanford, CA, USA

[4]Howard Hughes Medical Institute, Chevy Chase, MD, USA

[†] These authors contributed equally to this work.

*Corresponding author and lead contact:

R. Becket Ebitz

Department of Neurosciences

Université de Montréal

Montréal, QC CANADA H3T 1J4

Email: becket@ebitzlab.com

Abstract

In uncertain environments, intelligent decision-makers exploit actions that have been rewarding in the past, but also explore actions that could be even better. Several neuromodulatory systems are implicated in exploration, based, in part, on work linking exploration to pupil size–a peripheral correlate of neuromodulatory tone and index of arousal. However, pupil size could instead track variables that make exploration more likely, like volatility or reward, without directly predicting either exploration or its neural bases. Here, we simultaneously measured pupil size, exploration, and neural population activity in the prefrontal cortex while two rhesus macaques explored and exploited in a dynamic environment. We found that pupil size under constant luminance specifically predicted the onset of exploration, the first exploratory trial in a sequence, beyond what could be explained by reward history. Pupil size also predicted disorganized patterns of prefrontal neural activity at both the single neuron and population levels, even within periods of exploitation. Ultimately, our results support a model in which pupil-linked mechanisms promote the onset of exploration via driving the prefrontal cortex through a critical tipping point where prefrontal control dynamics become disorganized and exploratory decisions are possible.

Significance Statement

Humans and other animals learn about the world through exploration: through making decisions that offer the opportunity to learn and discover, even when these decisions are not the best option in the moment. Neuroscience research has historically focused on understanding good choices, delivering many key insights into the neural mechanisms involved in these calculations. However, much less is known about how the brain generates exploratory decisions. This study identifies certain "early warning signs" of exploratory decisions in the brain and body, including certain signals in size of the pupil and the speed of neural activity in the prefrontal cortex. These early warning signs suggest that exploration may be the result of a critical tipping point in prefrontal brain states.

Introduction

Many decisions maximize immediate rewards. However, in uncertain or changing environments, it is important to sacrifice some immediate rewards in order to learn about the value of alternative options and discover new, more valuable strategies for interacting with the world. In

short, in complex environments, intelligent decision-makers exploit rewarding strategies, but
also explore alternative strategies that could be even better.

Because exploitation maximizes immediate reward, it can rely on the same value-based
decision-making processes that have been the subject of neurobiological studies for decades
(Ding and Hikosaka, 2006; Jurewicz et al., 2022; Platt and Glimcher, 1999; Roesch and Olson,
2007; Schultz et al., 2008). However, we are only just beginning to understand the neural bases
of exploration (Costa and Averbeck, 2020; Daw et al., 2006; Pearson et al., 2009; Wilson et al.,
2021, 2014). One clue is that many organisms seem to explore via random sampling
(Gershman, 2019; Wilson et al., 2021, 2014). Randomness is a critical component of
exploratory discovery in bird song and motor learning (Fiete et al., 2007; Wu et al., 2014), it can
perform about as well as more sophisticated strategies in many environments (Dayan and Daw,
2008), and humans and other primates tend to explore randomly even when more sophisticated
strategies are available (Ebitz et al., 2018; Wilson et al., 2014). There is some neurobiological
evidence linking random exploration to disorganized activity patterns in the prefrontal cortex
(Ebitz et al., 2019, 2018; Muller et al., 2019; Wilson et al., 2021), but we still do not understand
the proximate causes of random exploration and its neural correlates in the prefrontal cortex.

One promising hypothesis is that exploration could be under the control of some process(es)
linked to pupil size. Pupil size under constant luminance is a peripheral index of autonomic
arousal (Bradley et al., 2008; Ebitz and Moore, 2019; Loewenfeld, 1999) that also predicts
widespread changes in neural population activity (McGinley et al., 2015; Reimer et al., 2014)–
including in regions implicated in decision-making noise (Ebitz and Platt, 2015; Tervo et al.,
2014). Among other neuromodulators (Gilzenrat et al., 2010; Koss, 1986; Reimer et al., 2016),
pupil size is correlated with central norepinephrine (Costa and Rudebeck, 2016; Joshi et al.,
2016): a catecholamine that flattens neuronal tuning functions (Martins and Froemke, 2015) and
predicts "resets" in cortical networks (Aston-Jones and Cohen, 2005; Bouret and Sara, 2005).
Behaviorally, pupil size predicts decision-making noise (Aston-Jones and Cohen, 2005; Ebitz et
al., 2014; Eldar et al., 2013; Gilzenrat et al., 2010; O'Reilly et al., 2013; Wilson et al., 2021),
especially errors of reward-maximization (Jepma and Nieuwenhuis, 2011a) and task
performance (Ebitz et al., 2014; Ebitz and Platt, 2015). Some of these "errors" may be caused
by exploratory processes (Ebitz et al., 2019; Jepma and Nieuwenhuis, 2011a; Pisupati et al.,
2021).

There is a plausible alternative interpretation of this data: perhaps pupil size only tracks the variables that make exploration more likely. Pupil size under constant luminance increases with the volatility of reward environments, the surprise of reward outcomes, novelty, uncertainty, and context changes (Clewett et al., 2020; Filipowicz et al., 2020; Graves et al., 2021; Preuschoff et al., 2011; Slooten et al., 2018; Yokoi and Weiler, 2022): all variables that make exploration more likely. However, it is often unclear whether the pupil is tracking these variables or instead directly predicting behavioral changes like increased learning, decision-noise or exploration (Nassar et al., 2012; O'Reilly et al., 2013; Urai et al., 2017). Fortunately, recent results suggest that at least some exploration appears to occur tonically, regardless of these variables (Ebitz et al., 2019; Pisupati et al., 2021; Wilson et al., 2021). Further, in parallel, new computational approaches allows us to determine when exploration is occurring independently of the reward-based computations thought to drive it (Chen et al., 2021; Ebitz et al., 2020, 2019, 2018). This means that it is now possible to determine whether pupil size predicts exploration itself or instead simply tracks the variables that make exploration more likely.

Here, we measured pupil size and recorded from populations of prefrontal neurons while two rhesus macaques performed a task that encouraged exploration and exploitation. We found that pupil size under constant luminance was larger during explore choices than exploit choices. However, the temporal relationship between pupil size and exploration was both precise and complex: spontaneous oscillations in pupil size entrained the onset of exploration. Together, these results support the hypothesis that pupil-linked processes drive the prefrontal cortex through a critical tipping point that permits exploratory decisions.

Results

Two male rhesus macaques performed a total of 28 sessions of a classic explore/exploit task: a restless three-armed bandit (subject B: 10 sessions, subject O: 18 sessions; a total of 21,793 trials). Some analyses of this dataset have been reported previously (Ebitz et al., 2018), but the pupil data has not been analyzed previously and all analyses presented here are new. In this task, the reward probability (value) of three targets walks randomly and independently over time (**Figure 1A**). This means that the subjects have to take advantage of valuable options when they are available (exploit), but also occasionally sample alternative options to determine if they have become more valuable (explore).

Rather than instructing subjects to explore and exploit, this task takes advantage of the subjects' natural tendency to alternate between exploration and exploitation in a changing environment. We have previously shown that both monkeys and mice exhibit 2 behavioral modes in this task: one exploitative mode in which they repeatedly choose the same option—learning little but maximizing reward—and one exploratory mode in which they alternate rapidly between the options—choosing randomly with respect to rewards and learning rapidly (Chen et al., 2021; Ebitz et al., 2018). We infer which of these modes is driving behavior with a hidden Markov model (HMM; **Figure 1B**; see **Methods**). This approach models the exploratory and exploitative modes as latent goal states and the maximum *a posteriori* goal is taken as the state label for each choice. We have previously shown that this method identifies explore/exploit state labels that match normative definitions (Chen et al., 2021; Ebitz et al., 2018) and explain variance in prefrontal neural activity that cannot be explained by reward value, reward history, and switch/stay decisions (Ebitz et al., 2018). This task design naturally elicits exploration and exploitation, allowing us to investigate variability in pupil size and neural activity under both conditions.

Some previous studies used a different method to identify exploratory choices (Daw et al., 2006; Jepma and Nieuwenhuis, 2011a; Pearson et al., 2009). These studies fit a reinforcement learning (RL) model to the behavior and identified the choices that are not consistent with the model's subjective values as exploratory. However, this previous RL-based approach (1) equates exploration with errors of reward maximization, not a goal that is orthogonal to reward maximization, and (2) its accuracy depends on precise knowledge of the computations involved in the choice, which are highly variable, both across individuals and over time (Chen et al., 2021, 2021; Kaske et al., 2022). The HMM approach, conversely, makes no assumptions about the computations involved in the choice and identifies choices that are orthogonal to reward value, not anti-correlated with it (Chen et al., 2021; Ebitz et al., 2018). Here, we found that state labels from the HMM method explained more variance in behavior and neural activity than choice labels from the previous, RL method (**Figure 1C;** response time: both subjects, paired t-test: $p < 0.005$, $t(27) = 3.41$, the mean difference of beta weights = 0.004, 95% CI = 0.002 to 0.007; scatter index [(Ebitz et al., 2018)]: both subjects, paired t-test: $p < 0.001$, $t(27) = 3.84$, the mean difference of beta weights = 0.15, 95% CI = 0.07 to 0.24: see **Methods**). In short, we find that the HMM approach is a more robust and accurate method, with better face validity, than the RL-based method for identifying explore choices. Therefore, here, we used this more precise

approach to determine whether physiological signals, like pupil size, reliably track exploratory
behavior.

Figure 1. Task design and pupil. *A) Top: Subjects made saccadic choices between three*
identical options (T1, T2, and T3). One of the options (e.g., T1 in this example trial) was located
in the receptive field of a neuron in the frontal eye field (FEF; dotted circle). Bottom: Reward
probabilities for the 3 options (lines), with choices overlaid (dots) for 200 example trials. Gray
bars = explore-labels. B) The HMM models exploration and exploitation as latent goal states
underlying choice sequences. C) Comparison of regression coefficients for HMM-inferred and
RL-inferred explore choices, predicting either the disorganization of neural population responses
("scatter index"; see Methods; Ebitz et al., 2018) or response time. Beta weights were obtained
from session-level generalized linear models (GLMs) with explore state as the predictor (0 =

exploit, 1 = explore). Separate GLMs were fitted using explore labels from the HMM and from
an RL-based model. D) Average pupil size on explore and exploit choices. Right: Same for
individual subjects. E) The probability of explore choices as a function of pupil size quantile.
Dotted line: linear GLM fit. Solid line: quadratic fit. Right: Same for individual subjects. F)
Several behavior measures compared across median-split large- and small-pupil-size explore
choices. Left to right: reward probability, a one-trial-back learning index (see Methods), saccadic
peak velocity of saccades, the scatter index, and reaction time. No significant differences
between pupil bins. The grey line is the mean ± SEM for exploit choices. Error bars depict ±
SEM throughout.

Pupil size is larger during exploratory states

Having established that the HMM reliably distinguishes explore and exploit states, we next
asked whether pupil size changes across these behavioral states. Previous work using RL-
based labels reported that pupil size under constant luminance is larger during exploration than
exploitation (Jepma and Nieuwenhuis, 2011b). We therefore tested whether this pattern holds
using HMM-based labels in our dataset. Indeed, we found that pupil size at fixation (see
Methods) was larger on explore-labeled trials than exploit-labeled trials in both subjects (**Figure**
1D; both subjects, paired t-test: $p < 0.0001$, $t(27) = 4.95$, mean offset = 0.23, 95% CI = 0.13 to
0.32; subject B: $p < 0.001$, $t(9) = 5.50$, mean offset = 0.4, 95% CI = 0.24 to 0.57; subject O: $p <$
0.02, $t(17) = 2.85$, mean offset = 0.13, 95% CI = 0.03 to 0.23). Thus, pupil size was larger
during exploratory choices identified with the HMM method.

However, the probability of exploration did not increase linearly as a function of pupil size
(**Figure 1E**). A linear, first-order GLM confirmed that larger pupil size generally predicted more
explore choices (both subjects: $\beta = 0.084$, $p < 0.0001$, AIC = −1053.70, n = 28 sessions). This
relationship held when analyzed separately in each subject (subject B: $\beta = 0.063$, $p = 0.002$, AIC
= −368.01, n = 10 sessions; subject O: $\beta = 0.110$, $p < 0.0001$, AIC = −249.46, n = 18 sessions).
Yet the relationship was clearly nonlinear. A quadratic model provided a significantly better fit
than the linear model for the combined dataset (2nd order GLM: $\beta_1 = -0.081$, $p = 0.101$; $\beta_2 =$
0.166, $p = 0.0006$; AIC = −1063.62, AIC weight for the quadratic model = 0.993), consistent with
a U-shaped relationship. This U-shape was especially prominent in subject O ($\beta_1 = -0.13$, $p =$
0.090; $\beta_2 = 0.240$, $p = 0.001$; AIC = −258.01), although the quadratic model was not an
improvement over the linear model in subject B ($\beta_1 = -0.027$, $p = 0.709$; $\beta_2 = -0.091$, $p = 0.205$;
AIC = −367.63).

In order to determine whether this pattern was also apparent in the raw data (i.e. not the HMM-
model labels), we next examined how pupil size predicted switching behavior—i.e., choosing a

different option than in the previous trial. We again observed a U-shaped relationship between pupil size and the probability of making a switch choice in both subjects (1st-order GLM: β = 0.084, p < 0.0001, n = 28 sessions; subject B: β = 0.0800, p < 0.0001; subject O: β = 0.0807, p < 0.0001). A 2nd-order quadratic model provided a superior fit in both animals (both subjects: β_1 = −0.099, p = 0.023; β_2 = 0.184, p < 0.0001; subject B: β_1 = −0.069, p = 0.284, β_2 = 0.149, p = 0.010; subject O: β_1 = −0.093, p = 0.103, β_2 = 0.174, p = 0.001). Model comparison strongly favored the quadratic model (linear AIC = −1195.05; quadratic AIC = −1211.95; AIC weight for quadratic model = 0.9998). Thus, although pupil size tended to be larger during exploration than exploitation, its relationship with both exploration and switching was clearly U-shaped.

One possible explanation for the U-shaped pattern is that some "explore" choices—particularly those with small pupil size—reflect disengagement at low levels of arousal, rather than true exploration. However, if this were the case, then the valid, large-pupil explore choices would systematically differ from the false, small-pupil "explore" choices. They did not. To evaluate this possibility, we compared small- and large-pupil explore trials across several behavioral dimensions known to be sensitive to lapses in task engagement . These included reward rate, saccade velocity, the neural scatter index, a trial-wise learning index, reaction time—each previously associated with arousal, motivation, or task-related updating (Chen et al., 2021; Ebitz et al., 2018; Laurie et al., 2025). Small- and large-pupil explore choices (median split) were indistinguishable along several of the key dimensions that differentiate explore choices from exploit choices (**Figure 1F**). For example, both were equally likely to be rewarded (mean difference = 0.03 ± 0.24 STD) between large- and small pupil-explore choices (p > 0.4, t(1,27) = 0.75, paired t-test; AUC for discriminating explore and exploit = 0.65 ± 0.05 STD across sessions). Both had similar peak saccadic velocities (mean difference = -0.05 ± 0.23 STD, p > 0.2, t(27) = -1.08; explore/exploit AUC = 0.61 ± 0.10 STD) and both had more variability in neural population choice information ("scatter index", mean difference = 0.03 ± 0.33 STD, p > 0.6, t(27) = 0.45; explore/exploit AUC = 0.60 ± 0.07 STD). Both had similar levels of reward learning (see **Methods**; the mean difference = -0.03 ± 0.57 STD, p > 0.7, t(27) = 0.27): in both cases, learning was substantially enhanced relative to the exploit choices (small-pupil, the mean difference from exploit = 0.24 ± 0.48 STD, p < 0.02, t(27) = 2.69; large-pupil, the mean difference from exploit = 0.21 ± 0.39 STD, p < 0.01, t(27) = 2.91). Reaction times were also similar across small- and large-pupil explore choices (mean difference = 0.01 ± 0.02 STD, *p* > 0.6, *t*(27) = 1.84; explore/exploit AUC = 0.58 ± 0.06 STD). These results are incompatible with the idea that either type of explore choice reflects disengagement in the task or that small- and

225   large-pupil explore choices have different causes. Instead, we will see that the U-shape was
226   due to the complex temporal relationship between pupil size and exploration.

Pupil size, but not other measures ramp up before exploration

Pupil size ramped up across trials before exploration began in both subjects. After exploration, it
shrank to below-baseline levels when exploitation resumed (**Figure 2A**). Here, "baseline" refers
to a z-scored value of 0, computed by subtracting the session mean and dividing by the session
standard deviation of pupil size (see **Methods**).This ramping meant that pupil size was larger
not just during exploration, but also during the exploit choices immediately before exploration
(both subjects, GLM slope = 0.01, $p < 0.005$, n = 28; subject B: beta = 0.02, $p < 0.02$, n = 10;
subject O: beta = 0.01, $p < 0.05$, n = 18; average pupil size compared to the exploit choices,
post-hoc paired t-tests, 1 trial before exploration mean = 0.12, $p < 0.005$, $t(27) = 3.42$; 2 trials
mean = 0.09, $p < 0.03$, $t(27) = 2.41$; 3 trials mean = 0.03, $p > 0.1$, $t(27) = 1.42$; 4 trials mean =
0.05, $p < 0.05$, $t(27) = 2.07$). By the first exploit choice after exploration, pupil size had already
begun shrinking to below-baseline levels (post-hoc paired t-tests, 1 trial after exploration mean
= 0.03, $p = 0.09$, $t(27) = 1.73$; 2 trials after mean = -0.11, $p < 0.02$, $t(27) = -2.67$; 3 trials after
mean = -0.16, $p < 0.02$, $t(27) = -2.72$; 4 trials after mean = -0.08, $p > 0.2$, $t(27) = -1.27$; 5 trials
after mean = -0.16, $p < 0.001$, $t(27) = -3.94$; p-values are significant with a Holm-Bonferroni
correction). The shrinking to below-baseline levels could suggest a refractory mechanism that
would prevent exploration from re-occurring immediately after it happened.

To rule out potential confounds, we tested whether the pupil ramping and shrinking effects could
be explained by misaligned labels or unrelated behavioral dynamics. We saw no evidence of
ramping in peak saccadic velocity, another behavioral measure that differentiated explore trials
and exploit trials (**Figure 2B**; no significant decrease from baseline 1 trial before, paired t-test: p
> 0.7, $t(27) = 0.42$; a GLM was nonsignificant with the trend pointing in the opposite direction: 10
trials preceding exploration, beta = 0.008, $p > 0.1$) and no significant change from baseline
afterward (not greater than the baseline during the 5 trials after exploration, when pupil shrinking
was maximal, mean = -0.47 ± 0.15 STD, $p > 0.9$, one-sided $t(27) = -1.67$). We previously
reported similar results for decoded choice probability and the scatter index (see **Methods**) in
neurons of the frontal eye field (FEF), a prefrontal cortex region involved in directing gaze and
attention (Ebitz et al., 2018). Thus, while pupil size ramped before exploration began and shrank
afterward, the same was not true of other behavioral and neural variables, suggesting that these
dynamics were not some artifact of misalignment.

Pupil size generally ramps across trials, but resets with exploration

To better understand whether pupil ramping was a general feature of arousal dynamics or specific to exploration, we examined how pupil size evolved across trials with and without exploratory transitions (see **Methods**). When the subjects did not explore the pupil size increased steadily across trials (**Figure 2C**; both subjects, GLM: beta = 0.004, p < 0.0001; subject B: beta = 0.003, p < 0.0001; subject O: beta = 0.005, p < 0.0001, n = 25 lags over 28 sessions). This implies that the ramping in pupil size before explore choices may be a general dynamic of how pupil size evolves in the absence of exploration. However, a different pattern emerged when we looked at how the pupil changed between exploit trials that were separated by exploration. When two exploit trials were separated by at least one explore choice, pupil size was smaller on the second exploit trial (both subjects, GLM: beta = -0.09, p < 0.0001; subject B: beta = -0.14, p < 0.0001; subject O: beta = -0.07, p < 0.0001). Critically, passing through exploration only produced a baseline decrease in pupil size but did not alter the rate at which pupil size grew over trials (no significant interaction between slope and condition in both subjects, GLM: beta < -0.0001, p > 0.9; subject B: beta = 0.003, p < 0.05; subject O: beta = -0.002, p > 0.1; also nonsignificant on trials 5-25: both subjects: beta < 0.0005, p > 0.5). Therefore, pupil size tended to ramp across trials but exploratory choices

temporarily decreased pupil size without disrupting this ramping in the long term.

Figure 2. Pupil size ramps up before exploration and shrinks down after. *A) Average pupil*
size for 10 trials before and 10 trials after explore choices. Purple line: GLM fit. Right: Same for
each subject separately. B) Same as ***Figures 1D*** *and 2A but for peak velocity rather than pupil*
size. C) Change in pupil size between exploit trials that are either in a single bout of exploitation
(gray) or separated by explore trials (purple). Right: Same for each subject separately. D)
Change in pupil size over certain pairs of trials: starting (exploit to explore), during (explore to
*explore), and leaving (explore to exploit) exploration. *p < 0.001 E) The probability of starting to*
explore as a function of pupil size quantile. Solid line: Linear GLM fit. Error bars and shaded
regions depict mean ± SEM. Insets: Same analysis shown separately for each monkey.

Pupil size specifically predicts the onset—not the maintenance—of exploration

Pupil size tends to be smaller after exploration, but this shrinkage could either be driven by the
end of exploration (i.e. the start of exploitation) or it could begin shortly after the beginning of
exploration itself. If the pupil starts to shrink only after exploration ends, it would support models
that suggest that pupil size decreases with commitment to a new option or belief state (O'Reilly
et al., 2013). Conversely, if the pupil shrinks immediately after exploration begins, it might
suggest that pupil-linked mechanisms are important for initiating exploration, but not sustaining
it. Our results were consistent with the latter hypothesis: the pupil immediately began shrinking
as soon as exploration began, not when it ended (**Figure 2D**; mean change in pupil size
between neighboring explore choices = -0.17, t-test, $t(27)$ = -2.69, $p < 0.02$; 95% CI = -0.30 to -
0.04). This was essentially identical to the magnitude with which the pupil shrank on exploit
trials that followed explore trials (mean change = -0.17, t-test, $t(27)$ = -2.56, $p < 0.02$; 95% CI = -
0.31 to -0.03). Validating the ramping we observed with other methods, we also found that pupil
size tended to grow on explore trials that followed exploit trials here (mean change = 0.14, t-
test, $t(27)$ = 2.96, $p < 0.01$; 95% CI = 0.04 to 0.24). Together, these results suggest that pupil
size and pupil-linked mechanisms specifically predict the "onset" of exploration—the first
exploratory trial in a sequence—and may not be important for sustaining exploration after the
first explore choice.

317
The tendency of the pupil to shrink after the onset of exploration could explain the previously
noted U-shaped relationship between pupil size and exploration. In this view, the small-pupil-
size explore choices would be the later explore choices in a sequence and the larger pupil size
explore choices would tend to be the first explore choice(s). Indeed, pupil size had a primarily
linear relationship with the onset of exploration in both subjects (**Figure 2E**; 1st order GLM: β =
0.042, $p < 0.0001$, AIC = −1973.57). Adding a quadratic term did not substantially improve the

model fit (β_2 = 0.038, p = 0.073; quadratic model AIC = −1974.81; ΔAIC = −1.24; AIC weight of

quadratic model = 0.65; see **Methods**). This linear relationship was also observed in both

monkeys individually (see **Figure 2E**, right panels). For subject B, a first order GLM confirmed a

significant positive association (β = 0.044, *p* < 0.0001, AIC = −699.13), and adding a quadratic

term did not improve the model fit (β_2 = 0.049, *p* = 0.0878; ΔAIC = −0.96). Similarly, for subject

O, pupil size showed a significant linear relationship with exploration onset (β = 0.034, *p* <

0.0001, AIC = −460.29), and the quadratic model again provided no additional explanatory

power (β_2 = 0.022, *p* = 0.402; ΔAIC = +1.30). These results confirm that the linear relationship

between pupil size and the onset of exploration was robust across both subjects and not driven

by outliers or subject-specific variability. Conversely, there was no special relationship between

pupil size and probability of starting to exploit (1st order GLM: beta = 0.05, p > 0.05). Thus, pupil

size specifically predicted the onset of exploration, rather than explore choices or state switches

more generally.

If the U-shaped relationship between pupil size and exploration (**Figure 1E**) were driven

primarily by later explore trials, it should remain evident after excluding onset trials from the

analysis. Moreover, removing onsets should substantially reduce the slope of the linear effect.

To test this, we repeated the analysis using only later explore trials. As expected, the linear

slope decreased in both subjects (**Figure S1**): for subject B, β_1 dropped from 0.063 (all explore

trials) to 0.018 (excluding onsets), and for subject O, from 0.110 to 0.075. In contrast, the

quadratic terms remained relatively stable: for subject B, β_2 was 0.091 for all explore trials and

0.041 with onsets excluded; for subject O, β_2 was 0.240 and 0.215, respectively. Importantly,

the U-shaped relationship persisted when data were combined across both subjects (**Figure

S1**), with the quadratic model significantly outperforming the linear model

($\beta_2 = 0.121, p = 0.012$; AIC = −878.78 vs. −874.55;

AIC weight for the quadratic model = 0.893). These findings confirm that the nonlinearity

observed in the original analysis (**Figure 1E**) was driven by the decrease in pupil size in later

exploratory trials, whereas the onset of exploration had a largely linear relationship with pupil

size.

Exploration is gated by pupil-linked arousal, not just reward history

Although pupil size predicted the onset of exploration, it remained possible that this relationship

was driven by a shared sensitivity to recent reward outcomes, since both exploration (Daw et

al., 2006; Ebitz et al., 2018; Wilson et al., 2014) and pupil dilation (Bijleveld et al., 2009; Jepma

and Nieuwenhuis, 2011b) tend to increase following reward omission. To determine if there was
a direct effect of pupil-related processes on exploration, we compared pupil size across exploit
trials before exploration with pupil size from matched trial sequences where exploration did not
happen (see **Methods**). There was a significant increase in pupil size during the trials before
exploration compared to "matched rewards" control trials (**Figure 3B**; GLM, beta = 0.025, p <
0.01, n = 28), suggesting that pupil size predicted the onset of exploration beyond what could be
explained by reward. Again, pupil size ramped up over time (GLM, beta = 0.119, p < 0.02, n =
28), but this ramping did not differ between the traces (GLM, beta = 0.007, p > 0.5, n = 28). This
implies that either reward history or time (i.e., the number of trials) may explain the pupil
ramping before exploration, although there is still an offset in pupil size that predicts the onset of
exploration above and beyond the effect of reward history.

Visual inspection of **Figure 3A** suggested that there may be a phase difference in pupil size
between trials where exploration began and matched reward trials where exploration did not
begin. This led us to develop a novel hypothesis (**Figure 3B**): that rewards may interact with
ongoing oscillations in pupil size. Due to delays in communication between the baroreceptor
reflect and changes in heart rate, the sympathetic nervous system (Borjon et al., 2016;
Japundzic et al., 1990; Julien, 2020, 2006; Kamiya et al., 2005; Liao et al., 2018) has a natural
oscillation known as the Mayer wave, with a period of approximately 0.05–0.1 Hz (Borjon et al.,
2016b; Julien, 2006). Critically, transitions in other behavioral states can be entrained by this
oscillation, with eliciting stimuli causing transitions only at certain phases of arousal. This view
predicts that the previous trials most predictive of the onset of exploration may actually be
several trials prior to onset itself—during the periods in which the signals are most out of phase.
Indeed, the trials in which pupil size best predicted the onset of exploration were not those
immediately preceding it (e.g., trial t–1 or t–2), but rather trials t–4 and t–5 (**Figure 3C**; trial t–4,
mean difference = 0.117, p < 0.05, t(27) = 2.09; trial t–5 = 0.170, p < 0.01, t(27) = 2.84).

The view that omitted rewards only evoke exploration when they coincide with particular phases
of sympathetic arousal also implies that that the onset of exploration should be phase-locked to
the Mayer wave frequency (see **Methods**). The median trial duration was ~3 seconds (range =
[2.2, 3.2]), so a ~5-trial cycle would correspond to a 0.06–0.09 Hz oscillation, which aligns with
the frequency range of the Mayer wave. We found that pupil phase at the onset of exploration
was concentrated at the rising phase (**Figure 3D**; mean phase = 47.18°, Hodges-Ajne test, p <
0.01; vector length = 0.075; null = 0.026, 95% CI = 0.004 to 0.057, p < 0.0001). In contrast, pupil
phases during reward-matched trials pointed in the opposite direction (mean phase = 207.58°;

significantly different from onsets, $p < 0.02$, Watson's $U^2 = 0.25$, n = 2170 phases including
1135 onsets). Together, these results support the hypothesis (**Figure 3B**) that slow, rhythmic
fluctuations in arousal interact with reward history to determine the timing of exploration onset.

392
393
Figure 3. The onset of exploration is phase-locked to pupil size. *A) Average pupil size over*
sequences of exploit trials before the onset of exploration (black line) and sequences with
matched rewards, but no exploration at the end (gray line). Lines: GLM fit. B) Cartoon illustrating
how oscillations in arousal (top) could interact with reward history (middle), to regulate
exploration. The bottom panel illustrates a hypothetical pupil trace that has an additive effect of
reward omissions and by oscillating arousal. Exploration (diamond shapes) begins when pupil
size reaches a threshold (dotted line). Note that identical patterns of reward delivery and
omission have different outcomes, depending on how they align with the phase of arousal (gray
= no exploration, blue = exploration). C) Difference in pupil size between the traces in A. D)
Phase distribution of pupil size at the onset of exploration (blue) and bootstrapped null
distribution (black). The vectors at the center indicate the mean vector direction and length for
the trials before exploration (blue) and the matched reward trials (gray). Shaded areas ± SEM
throughout.
407

Pupil size predicts flattened neural tuning in prefrontal cortex during exploration

To probe the neural mechanisms linking pupil size to exploration, we examined how pupil size
predicts neural activity in the FEF (Bruce and Goldberg, 1985; Moore and Armstrong, 2003;
Moore and Fallah, 2001; Schall and Hanes, 1993) (**Figure 4A**). We previously reported that
exploration is associated with flattened tuning for choice in FEF neurons (Ebitz et al., 2018).
While FEF neurons often predict upcoming choices during exploitation, many show reduced
choice selectivity during exploration. Pupil size predicted similar changes in in FEF neurons and

did so beyond what could be explained by exploratory states themselves. Out of 155 recorded
single neurons, 88 (57%) were tuned for choice (**Figure 4B**; 57%, one sample proportion test: p
< 0.001). These are referred to as "tuned neurons," regardless of whether they were modulated
by pupil size. Among tuned neurons, 21 (24%) were also modulated by pupil size, and 16 (18%)
showed a significant interaction between choice and pupil size. On average, tuning curves
flattened as pupil size increased in both tuned and untuned neurons (**Figure 4C-D**). Among
untuned neurons, an additional 22% (15/67) were significantly modulated by pupil size (*p* <
0.05), with a median regression coefficient (β) of −0.0011 ± 0.063. This suggests that pupil-
linked mechanisms affect FEF activity even in neurons that are not directly involved in encoding
choice. This may suggest a more domain-general role for arousal in modulating prefrontal
network dynamics.
426

Figure 4. Pupil size predicts choice tuning curves and population disorganization. A) Recordings were made in the FEF. Right: The cartoon illustrates the relative positions of the receptive field target (T_{in}, red) and the ipsilateral and contralateral targets (T_{out}, blue and green). B) Percent of neurons with significant tuning for choice target, pupil size, and the interaction. C) Tuning curve for an example neuron across target locations, separated by pupil size. Lighter = larger pupil. The dashed line indicates the number of neurons expected to be significant by chance at p < 0.05. D) Same for all tuned neurons, which refers to the 88 out of 155 FEF neurons that were significantly tuned for choice, regardless of their modulation by pupil size. E) Cartoon illustrating how neural population measures consider patterns of firing rates across neurons as vectors in neural state space. Targeted dimensionality reduction is used to find the hyperplane where the distribution of neural activity across trials best predicts choice. Vectors here are the coding dimensions that separate the choices. F) The decoded choice probability (projection onto the correct coding dimension) plotted as a function of pupil size quantile. Inset: Same for exploit trials alone. G) The scatter index, a measure of the variance in choice-predictive population activity, plotted as a function of pupil size quantile. Inset: Same for exploit trials. H) Decoded choice probability for trials before the onset of exploration (in blue) and trials with matched rewards (in gray).I) Scatter index for trials before the onset of exploration (in blue) and trials with matched rewards (in gray). Error bars and shaded regions ± SEM. J) Mediation analysis between pupil size, scatter index, and the onset of exploration. Top: Direct model. Bottom: Indirect, mediated model. Asterisks marked significant paths (*p < 0.01 **p < 0.001).

Because single neurons are noisy, it is difficult to dissociate the effects of pupil size and exploration at the level of individual cells. However, changes in tuning at the single-neuron level also imply shifts in the organization of the neural population and looking at the population level can allow us to estimate these effects within smaller subsets of the data (Ebitz and Hayden, 2021); **Figure 4E**). Indeed, we found that pupil size also predicted changes in how accurately choice information could be decoded from simultaneously recorded populations of FEF neurons. Consistent with our prior results (Ebitz et al., 2018), decoded choice probability was significantly lower during exploration compared to exploitation (paired t-test: both subjects, p < 0.0001, **Figure S2A**). Critically, larger pupil size predicted weaker choice encoding both across all trials (**Figure 4F**; GLM: beta = -0.032, p < 0.0001). This was not driven by differences between the states because pupil size also predicted choice decoding accuracy within exploit trials alone (GLM: beta = -0.037, p < 0.005). There was no significant effect of pupil size on choice decoding within explore trials (GLM: β = –0.003, p = 0.77, **Figure S2B**), which could have been due to differences in trial counts (i.e., explore trials made up only ~15% of total trials) or to floor effects (decoding accuracy was already close to chance in these trials). These findings show that pupil-linked arousal predicts the strength of choice-predictive neural signals in FEF above and beyond what can be explained by differences between the states.

In our previous work, we found that decreases in choice-predictive activity were accompanied by increases in variability in FEF population responses to the same choice (Ebitz et al., 2018).

We quantified this with the "scatter index": a measure of the spread within clusters of same-
choice population activity (see **Methods**). A high scatter index indicates that neural activity on a
given trial was dissimilar to other trials where the same choice was made, whereas a low scatter
index indicates that neural activity was tightly clustered. We observed a higher scatter index
during exploration compared to exploitation (paired t-test: both subjects, $p < 0.0001$, **Figure**
S2C). Here, we also found that increasing pupil size predicted an increase in the scatter index in
both subject B and subject O (**Figure 4G**; GLM: beta = 0.04, $p < 0.0001$). This effect remained
significant and of similar magnitude within exploit trials alone (GLM: $\beta = 0.03$, $p < 0.0005$), again
suggesting that the relationship between pupil size and scatter was not an artifact of state
differences with pupil size. Pupil size again did not significantly predict the scatter index during
explore trials (GLM: $\beta = 0.0006$, $p = 0.94$, **Figure S2D**). Thus, pupil size predicted
disorganization of choice-predictive signals in the FEF, at both the level of single neurons and in
the population.

Neural disorganization mediates the relationship between pupil size and exploration

To test whether neural population activity, like pupil size, also specifically predicted the onset of
exploration, we compared its dynamics in the trials preceding exploration to those in matched-
reward control trials. While sudden changes in the decoded choice probability and scatter index
were largely aligned with the onset of exploration (as reported previously), these neural
measures were at a different average level on the trials preceding exploration, compared to
reward-matched controls (**Figure 4H-I**; choice probability, offset = -0.611, $p < 0.001$, n = 28;
scatter index = 0.258, $p < 0.001$). Reward information did not cause a change in either variable
(choice probability, slope = -0.004, $p > 0.5$; scatter index = 0.002, $p > 0.5$), while small, but
significant interaction terms suggested that both variables anticipated the onset of exploration
(choice probability interaction = -0.058, $p < 0.01$; scatter index interaction: beta = 0.040, $p <$
0.001). To determine if these neural measures might explain or mediate some of the
relationship between pupil size and exploration, we turned to structural equation modeling
(Preacher et al., 2007; Sobel, 1986). We found that the scatter index was a significant mediator
of the relationship between pupil size and the onset of exploration (**Figure 4J**; effect of
mediation, ab = 0.003, $p < 0.005$; full report in Table S1). Together, these results suggest that
pupil size predicts disruptions in the organization of prefrontal neural activity that then mediate
its relationship with the onset of exploration.

Figure 5. Pupil size predicts behavioral and neural slowing. *A) Response time on exploit trials before the onset of exploration (blue) and trials with matched rewards but no exploration (gray). B) Response time plotted as a function of pupil size quantile. Inset: Same for exploit trials alone. C) Mediation analysis between pupil size, response time, and the onset of exploration. Top: Direct model. Asterisks marked significant paths (*p < 0.01). Bottom: Indirect, mediated model. D) Neural speed on exploit trials before the onset of exploration (in blue) and trials with matched rewards (in gray). E-F) Same as B-C for neural speed. Shaded areas and error bars ± SEM.*

Exploration may reflect a critical transition in brain state dynamics

Neural systems, like other complex networks, can undergo tipping points—irreversible "critical transitions" between stable operating regimes (O'Byrne and Jerbi, 2022; Scheffer, 2020; Scheffer et al., 2009; Wang et al., 2012). Because exploration occurs as the brain passes from exploiting one target to exploiting another, it is worth considering the possibility that exploration may represent a critical transition in brain states. Indeed, during exploration, we previously reported (Ebitz et al., 2018) several phenomena in the FEF and in behavior that are hallmarks of critical transitions, including a rapid flickering back and forth between choices (Wang et al.,

2012), an increase in the variance in neural activity (Scheffer et al., 2009), and a disruption of
long-term neuronal autocorrelations that suggests that passing through exploration causes time-
irreversible changes in the FEF network (Scheffer, 2020). However, there is another classic
feature of critical transitions that we did not consider: an early warning signal known as "critical
slowing". As the system nears the tipping point, the dynamics within the system begin to flatten
out in preparation for the change. As a result, the systems' processes slow down and take
longer to trace the same paths (Scheffer et al., 2009). Therefore, we next asked if there was any
evidence that decision-making slowed down in advance exploration in this dataset.

To test for critical slowing, we examined two measures of decision speed: one behavioral and
one neural. First, we looked at response time, a measure of how long it takes the brain to
generate saccadic decisions. Response time was not only slower in the trials before exploration,
compared to matched-reward control trials (**Figure 5A-C**; GLM offset = 0.39, p < 0.0001, n =
28), but it slowed down over trials before the onset of exploration (interaction = 0.05, p < 0.001).
Second, we looked at the mean rate of change in neural population choice signals during the
decision process ("neural speed", see **Methods**). Neural speed was only weakly correlated with
response time across sessions (mean = -0.07, min = -0.36, max = 0.09, Pearson's correlation),
suggesting that the measures were complementary, rather than redundant. Like response time,
neural speed was also significantly slower on average in the trials before exploration, compared
to matched-reward controls (**Figure 5D–F**; GLM offset = -0.17, p < 0.0001, n = 28). However,
unlike response time, neural speed did not show a significant slowing trend over trials
(interaction = -0.01, p = 0.08). Although the notion that the brain may be subject to critical
tipping points is controversial (O'Byrne and Jerbi, 2022), these results are consistent with the
idea that exploration could reflect a critical transition between exploiting one option and
exploiting another.

We first asked whether slowing effects could be better explained by the typical reward histories
that precede exploration, rather than internal states like arousal. However, reward history alone
did not have a significant effect on either neural or behavioral slowing (response time: slope of
matched-reward trials = 0.0002, p > 0.5; neural speed: slope = -0.018, p > 0.1). This suggests
that some internal variable, like arousal, could be driving increased slowing and, perhaps, also
the systems' proximity to a tipping point. Indeed, increasing pupil size predicted slower
response times (**Figure 5B**; GLM beta = 0.08, p < 0.0001, n = 28 sessions), even within periods
of exploitation (beta = 0.05, p < 0.0001). The same was true of neural slowing (**Figure 5E**; all

trials: beta = -0.03, p < 0.0005; exploit only: beta = -0.09, p < 0.0001). Further, structural

equation modeling revealed that both measures of slowing mediated the relationship between

pupil size and the onset of exploration (**Figure 5C and F**; Table S2-3). In sum, the pupil-linked

mechanisms that anticipated exploration included both a disorganization of neural activity and a

slowing of decision-related computations in brain and behavior—hallmarks of a system

approaching a critical transition.

Discussion

Random decision-making is a powerful strategy for exploration (Dayan and Daw, 2008; Ebitz et

al., 2018; Gershman, 2019; Wilson et al., 2021, 2014) that is linked to disorganized patterns of

neural activity in the prefrontal cortex (Ebitz et al., 2018; Muller et al., 2019; Wilson et al., 2021).

Here, we sought to identify some of the neurobiological mechanisms that drive random

exploration and its neural signatures. We found that pupil size, a peripheral correlate of

autonomic arousal, predicted exploration and certain measures of neural population activity

previously linked to exploration. Consistent with previous studies (Jepma and Nieuwenhuis,

2011a), pupil size was generally larger during exploration, compared to exploitation. However,

there was also a complex temporal relationship, where pupil size ramped up between periods of

exploration and decreased during exploration. As a result, pupil size was largest at the

beginning or "onset" of exploration and explained variance in the onset of exploration that could

not be explained by other variables. Together, these results suggest that pupil-linked

mechanisms may play a role in driving the brain into an exploratory state.

Our behavioral results largely replicate previous findings linking exploration to increased pupil

size (Jepma and Nieuwenhuis, 2011a). However, where we found gradual ramping before

exploration and sudden constriction after, Jepma and Nieuwenhuis (2011) reported an abrupt (if

modest) increase of pupil size at the onset of exploration and then a gradual decrease at the

return to exploitation. The discrepancy may be due to differences in the operational definition of

exploration. Jepma and Nieuwenhuis (2011) fit an RL model to behavior and defined "explore

choices" as the choices that were not reward-maximizing according to the model. This definition

conflates exploration with errors of reward maximization. A strategy that is non-reward-

maximizing would produce choices that are orthogonal to value, not consistently bad. Here, we

used an HMM to identify latent explore and exploit states on the basis of the temporal profiles of

choices alone, with no assumptions on the underlying value computations. This allowed us to

dissociate the effects of reward history from the explore/exploit choice labels. We reported here

(**Figure 1C**), and in previous studies (Chen et al., 2021; Ebitz et al., 2018), that HMM labels
outperform RL labels in explaining behavioral and neural measures, suggesting that the HMM
may more accurately separate distinct neural and behavioral states. If the HMM allows for more
precise identification of exploratory and exploitative choices, it would follow that it also allows for
more precise reconstruction of the temporal relationship between the pupil and exploration.

The precision of our explore/exploit labels revealed that the U-shaped relationship between
pupil size and exploration was caused by a refractory constriction in the pupil. When exploration
was plotted as a function of pupil size, the relationship appeared non-linear: both small- and
large-pupil choices were more likely to be exploratory. This superficially resonated with the idea
of a U-shaped relationship between arousal and task performance (i.e. the "Yerkes-Dodson
curve"; (Aston-Jones and Cohen, 2005; Yerkes and Dodson, 1908): perhaps reliable exploitation
is only possible at intermediate levels of arousal. However, when we examined the temporal
relationship between exploration and pupil size, we found that pupil size only predicted the
onset of exploration, the first explore choice in a sequence. Small-pupil explore choices
happened because starting to explore seemed to "reset" the level of pupil-linked arousal,
causing it to quickly fall below baseline. If increased pupil size promotes a transition to
exploration, then it is possible that post-exploration constriction represents a refractory period
for exploration. Given that uncertainty grows with time in this task (and in all dynamic
environments), it may not be smart to start to explore again immediately after you have just
explored. A refractory period could ensure that non-reward-maximizing explore choices are
deployed only when needed. Future work is needed to test this hypothesis and to determine the
cognitive and/or neurobiological mechanisms at play.

Before exploration, we observed an oscillatory dynamic that was about twice as fast as the 10
trials it took the pupil to recover after exploration. This 0.06-0.09 Hz oscillation entrained the
onset of exploration: onsets tended to occur during the rising phase of pupil size, whereas
identical trial sequences that did not result in exploration were on the opposite phase. This
implies that it is the confluence of pupil size, pupil phase, and trial history that best predicts the
onset of exploration. This result reinforces the idea that arousal or arousal-linked mechanisms
help trigger random exploration (Ebitz and Moore, 2019; Gilzenrat et al., 2010; Reimer et al.,
2016), rather than just tracking the reward-linked variables that make exploration more
probable. It is also notable that the period of the pupil oscillation was close to the frequency of
the Mayer wave: an oscillation in blood pressure that entrains other autonomic measures,

including respiration and heart rate (Borjon et al., 2016a; Japundzic et al., 1990; Julien, 2020,
2006; Kamiya et al., 2005). There is precedent for the idea that behavior can be entrained by
the Mayer wave: in marmosets, fluctuations in arousal predict the spontaneous onset of a call
(Borjon et al., 2016a). This paper argued that the Mayer wave may function to organize vocal
communication by bringing the system closer to the threshold for transitioning from inaction to
action. It is possible that oscillations in the pupil and pupil-linked mechanisms function the same
way here, organizing important state changes in time. In parallel, pupil-linked mechanisms seem
to anticipate other state transitions, including belief updating (Filipowicz et al., 2020; O'Reilly et
al., 2013), task disengagement (Kane et al., 2017), and other behavioral state changes (Bouret
and Sara, 2005). Together, these results suggest an important role for pupil-linked mechanisms
in driving successful transitions between certain neural and behavioral states.

Critically, pupil size and pupil oscillations did not predict all state transitions here, but only the
transition into exploration. What kinds of state transitions might be entrained by pupil- linked
arousal? It is possible that the pupil may have a special relationship with certain "critical" kinds
of transitions. Critical transitions are abrupt, large-scale, and irreversible changes in the
dynamics and behavior of complex systems, like the brain. As these systems go from being in
one conformation (i.e. always choosing the left option) into another conformation (i.e. always
choosing the right), the system dynamics that support the old state have to disappear and the
new dynamics have to emerge. During this brief transitory period, when both dynamics co-exist
in the system, certain signatures can be observed in the system. We previously reported that
the exploration was accompanied by abrupt changes in neural population activity, certain
patterns of noise in brain and behavior, and disruptions in long-term neuronal autocorrelations:
all observations that could be interpreted as suggesting that exploration is a critical transition in
the brain (Ebitz et al., 2018). Here, we found that pupil size predicts these features of neural
activity and also an prominent "early warning sign" of critical transitions: a slowing, in brain and
behavior, of the decision process. While there are certain patterns of activity in FEF that predict
response speed (Hauser et al., 2018; Yao and Vanduffel, 2023), here we identified *independent*
neural and behavioral measures of decision speed that both mediated the relationship between
pupil size and exploration. Notably, pupil size also predicted slower neural and behavioral
responses within exploit-only trials, suggesting that these effects are not an artifact of
differences between explore and exploit states. This suggests that these effects may reflect a
domain-general influence of arousal on cognitive dynamics, consistent with the idea that
fluctuations in pupil-linked neuromodulation shape the temporal structure of decision-making in

general, beyond any role in state transitions. In this view, the transition into exploration in FEF

may reflect an extremum of these domain-general arousal effects—a tipping point—rather than

signals that are highly specific to state transitions. Together these results suggest that pupil-

linked arousal pushes neural and behavioral states to a critical tipping point and highlights the

crucial role of pupil-linked mechanisms in changing the dynamics of the brain.

What underlying, pupil-linked mechanisms could support critical transitions? Changes in pupil

diameter coincide with neuromodulator system activity, especially norepinephrine (NE) and

acetylcholine (Breton-Provencher and Sur, 2019; de Gee et al., 2020; Joshi et al., 2016; Joshi

and Gold, 2020; Murphy et al., 2014; Reimer et al., 2016). At the neuronal level, central NE

flattens tuning curves, at least in the auditory cortex (Martins and Froemke, 2015), though it may

have different effects in non-cortical structures (Manella et al., 2017). Here, we made a parallel

observation: as pupil size increases, neuronal turning curves flattened and choice-predictive

neural population activity became disorganized. These results resonate with a particularly

influential theory of NE function: the idea that NE release may facilitate "resets" in cortical

networks in order to effect long-lasting changes in brain and behavior (Aston-Jones and Cohen,

2005; Bouret and Sara, 2005). More recent studies seem to consistently report that elevated

levels of NE predict an increase in behavioral variability, while pharmacological blockade of NE

receptors reduces variability (Chen et al., 2023; Kane et al., 2017; Sadacca et al., 2017; Tervo

et al., 2014). In combination with the present study, these results could suggest that phasic NE

signaling functions to push the brain towards a critical tipping point where it is better able to

transition from one regime to another. In this view, behavioral variability would be linked to NE

not because NE increases variability directly, but because the brain is more likely to transition

into a high variability regime after it is released. Of course, pupil size is also associated with

other neuromodulatory systems, cognitive factors, and other measures of arousal. Thus, future

work is needed to identify the neurobiological mechanisms that underpin the relationship

between pupil size and critical transitions that we report here.

Materials and Methods

Surgical and electrophysiological procedures. All procedures were approved by the Stanford

University Institutional Animal Care and Use Committee. Subjects were two male rhesus

macaques, surgically-prepared with head restraint prostheses, craniotomies, and recording

chambers under isoflurane anesthesia via techniques described previously (Ebitz et al., 2018).

Following surgery, analgesics were used to minimize discomfort, and antibiotics were delivered
prophylactically. After recovery, subjects were acclimated to the laboratory and head restraint,
then placed on controlled access to fluids and trained to perform the task.

In order to train the animals on the explore/exploit task a gradual procedure was used in which
the two animals were first trained to make saccadic eye movements in exchange for liquid
rewards. Once the animals reliably made controlled eye movements to a single target (generally
within 1–2 days), a second target was introduced, and the animals were free to choose between
them. At the outset, each target was associated with a probability of reward (initially 10% and
90%), which was reversed in blocks at the experimenter's discretion. Over a period of 2–4
months, the difference in reward probabilities between the targets was gradually reduced, the
blocks transitioned into gradual reward probability shifts (reward walks), and a third target was
introduced. The speed and order of these changes depended on each animal's performance
and engagement with the task. One animal (monkey O) was naïve to laboratory tasks prior to
this experiment, whereas the second (monkey B) had been previously trained on covert and
overt attention tasks, but not on any prior value-based tasks.

Recording sites were located within the FEF, which was identified via a combination of
anatomical and functional criteria. The location of recording sites in the anterior bank of the
arcuate sulcus was verified histologically in one subject and via microstimulation in both
subjects (Ebitz et al., 2018). Recordings were conducted with 16-channel U-probes (Plexon),
located such that each contact was within gray matter at an FEF site. An average of 20 units
were recorded in each session (131 single units, 443 multi units; 576 total units across 28
sessions).

General behavioral procedures. Eye position and pupil size were monitored at 1000 Hz via an
infrared eye tracking system (SR Research; Eyelink). The manufacturer's standard methods for
calculating pupil area were used. MATLAB (Psychtoolbox-3; (Kleiner et al., 2007)) was used to
display stimuli and record behavioral responses and pupil size measurements. Task stimuli
were presented against a dark gray background (7 cd/m2) on a 47.5 cm wide LCD monitor
(Samsung; 120 Hz refresh rate, 1680 x 1050 resolution), located 34 cm in front of the subject.

Three-armed bandit task. The subjects performed a sequential decision-making task in which
they chose between 3 targets whose values changed over time. The subject first fixated a
central fixation square (0.5° stimulus, +/- 1.5-2° of error) for a variable interval (450-750ms). At

any point within 2s after the onset of the targets, subjects indicated their choice by making a saccade to one of the targets and fixating it (+/- 3°) for 150 ms. Reward magnitude was fixed within session (0.2-0.4 µL). Reward probability was determined by the current reward probability of the chosen target, which changed independently over trials for each of the three targets. On every correct trial, each target had a 10% chance of the reward parameter changing either up or down by a fixed step of 0.1, bounded at 0.1 and 0.9. Because rewards were variable, independent, and probabilistic, the subjects could only infer the values of the different targets by sampling them and integrating noisy experienced rewards over multiple trials.

General analysis procedures. Data were analyzed with custom software in MATLAB. Unless otherwise noted, all t-tests were paired, two-sided t-tests, and generalized linear models were run on raw data, with session number coded as a dummy variable to account for session-to-session variability. Model comparison was based on standard methods that involve calculating the likelihood of the data and Akaike information criteria (AIC) of each model, then using AIC weights to identify (1) the model that is most likely to minimize information loss, and (2) the relative likelihood of competing models to do the same (Burnham and Anderson, 2004). For analyses of any behavioral or neural variables on the trials before or after exploration, continuous runs of exploit trials were required. The values of behavioral and neural variables were *z*-scored within a session to facilitate comparisons across sessions. In the results section, we refer to a z-score of 0 as "baseline". The 200 ms window immediately preceding target onset was chosen as the analysis epoch for all choice-predictive neural measures. A longer, whole-trial epoch was chosen for neural speed analyses (0 to 500 ms) following target presentation. Firing rates were computed per trial.

Pupil size. Pupil size was measured during the first 200 ms of fixation, a time at which the eye was fixed at a known point on the screen, illumination was identical across trials, and anticipatory changes in the pupil were minimal. To remove any blinks or movement artifacts, trials where pupil size or the change in pupil size from the first time bin of this epoch to the last was +/- 6 standard deviations from average were eliminated from further analyses. A total of 178 trials (out of 21,793, approximately 0.8% of observations) were outliers.

Hidden Markov Model. To identify when subjects were exploring versus exploiting, we employed a hidden Markov model (Chen et al., 2021; Ebitz et al., 2018). In this framework, choices Y_t are treated as emissions from a latent decision-making state z_t, which can either be an explore or one of the multiple exploit states.

746   The emission model for exploration assumed a uniform probability of selecting any option:

$$p(y_t = k \mid z_t = explore) = \frac{1}{N_k}$$

748   Where N_k is the total number of options. In contrast, exploit states deterministically emitted
749   choices to the exploited option i:

$$p(y_t = k \mid z_t = exploit_i, k = i) = 1$$

$$p(y_t = k \mid z_t = exploit_i, k \neq i) = 0$$

751   Latent state transitions followed a Markov process, such that the probability of the current state
752   depended only on the previous state:

$$p(z_t \mid z_{t-1}, y_{t-1} \ldots, z_1, y_1) = p(z_t \mid z_{t-1})$$

To reduce model complexity, parameters were shared across exploit states, and subjects were
assumed to begin in the explore state. The final HMM included only two free parameters: the
probability of persisting in exploration and the probability of persisting in exploitation. The model
was fit using expectation-maximization with 20 random restarts, and the solution maximizing the
observed data log-likelihood was selected. The most probable sequence of latent states was
recovered using the Viterbi algorithm.

Reinforcement learning model. To compare goal state labels derived from an RL and HMM
model, we employed a Rescorla-Wagner model. This was fit using maximum likelihood
estimation. The value of each option is iteratively updated according to:

$$V_{i,t+1} = V_{i,t} + \alpha(r_t - V_{i,t})$$

764   Where $V_{i,t}$ is the value of option *i* at time *t*, r_t is the reward at time *t*, and α represents the fitted
765   learning rate, which determines how much the difference between the predicted and actual

766   reward (the prediction error) influences value. To make a decision, the values are passed
767   through a softmax decision rule:

$$p_{i,t} = \frac{\exp(\beta V_{i,t})}{\Sigma_{j=1:n} \exp(\beta V_{j,t})}$$

Where n is the total number of available options, and β is the inverse temperature, which
controls the level of random noise in decision-making. After (Daw et al., 2006; Jepma and
Nieuwenhuis, 2011a; Pearson et al., 2009), decisions that were not reward maximizing were
labeled as exploratory (i.e. any decision where $V_{chosen,t}$ was not the maximum V at time t).

Generalized Linear Model (GLM). To examine the relationship between behavioral and neural
variables, we employed generalized linear models (GLMs). These models were fit using
maximum likelihood estimation. Each dependent variable   (i.e., scatter index, response time,
or choice probability) was modeled as a linear combination of predictors:

$$Y_i = \beta_0 + \beta_1 \cdot explore\ state_i + \beta_2 \cdot pupile\ size_i + \beta_3 \cdot (explore\ state_i \times pupile\ size_i) + \varepsilon_i$$

Where Y_i is the dependent variable on trial i, β_0 is the intercept, and β_1, β_2, β_3 are regression
coefficients quantifying the influence of the corresponding predictors (e.g., explore state, pupil
size, and their interactions). ε_i is the residual error term.

In analyses where categorical variables (e.g., explore state: 0 = exploit, 1 = explore) were used
as predictors, these were coded as binary dummy variables. Models were fit using the identity
link function and assumed normally distributed errors.

Learning Index. To investigate whether learning differed with pupil size within the exploratory
choices, we calculated a learning index that captured the effect of rewards experienced during
exploration on future choices. Because reward effects decay exponentially quickly (Lau and
Glimcher, 2008), a 1-trial-ahead index should capture most of the variability in how much is
learned between trial types. The equation was:

$$\text{learning index}_t = \frac{p(\text{switch}_{t+1} \mid \text{reward}_t) - p(\text{switch}_{t+1} \mid \text{no reward}_t)}{p(\text{switch}_{t+1})}$$

Lagged change in pupil size. To determine whether exploration impacted pupil size, we
measured the change in pupil size (Δ pupil) between pairs of trials that either were or were not
separated by at least 1 explore trial. Segments of twenty-five consecutive trials were identified
that either included a single bout of exploration or did not include exploration. For each pair of
trials within these sequences, we then measured the change in pupil size between the first
exploit trial of the sequence (t1) and the remaining exploit trials in the sequence (t2:25). This
was repeated for all unique pairs of trials that met our selection criteria.

Matched reward trials. To test whether the rising trend in pupil size before exploration is best
explained by reward history, we identified trial sequences with identical reward and state
histories that did not end in exploration ("matched rewards"). For each onset of exploration
preceded by at least 6 exploit trials, we searched for identical sequences of exploit trials, with
identical reward histories, that did not end in exploration. We chose 6 previous trials because
this was the longest sequence of reward history we could regularly match within the majority of
sessions (we could find at least 10 matched sequences in 96% [27/28] of sessions for 6 trials
sequences; that dropped to 75% [21/28] at 7 trials). Identical results were obtained with other
sequence lengths, though these analyses included fewer sessions.

Mediation analysis. To determine if the predictive relationship between pupil size and
exploration was mediated by other variables, we used structural equation modeling to test for
mediation. Mediation analyses involve fitting three regression models. The first model measures
the total effect (c) of the independent variable (here, pupil size) on the independent variable
(here, onset of exploration):

$$\text{Explore}_t = \gamma_1 + c(\text{Pupil}_{t\text{-}1}) + \epsilon_1$$

814 In these equations, represents the intercept for each equation, while □ represents the error of
815 the model. Note that the estimated parameter c will include both direct effects of pupil size on
816 exploration, but also indirect effects that may be mediated by other variables. Therefore, we

817   also fit a second model that tests if the independent variable also predicts a potential mediator
818   variable (here, neural network scatter):

$$\text{Scatter}_{t\text{-}1} = \gamma_2 + a(\text{Pupil}_{t\text{-}1}) + \epsilon_2$$

820   Model parameter a thus captures the effect of pupil size on the mediator. Finally, a third model
821   estimates the unique contributions of both the potential mediator (scatter, b) and the
822   independent variable (pupil size, c'), now controlling for the mediator:

$$\text{Explore}_t = \gamma_3 + b(\text{Scatter}_{t\text{-}1}) + c'(\text{Pupil}_{t\text{-}1}) + \epsilon_3$$

A drop between c and c' indicates that the effect of the independent variable (pupil) on the
dependent variable (exploration) is reduced when the mediating variable is considered. The
mediation effect (the indirect effect of the pupil size on the onset of exploration via the mediating
factor) can also be estimated directly, via taking the product of the coefficients a and b. Sobel's
test is used to determine the significance of the mediation path (Sobel, 1986).

Phase analysis. To determine if the onset of exploration happened at a specific phase of pupil
size over trials, we performed a wavelet analysis. Because this method only assumes local
stationarity, it is more suitable than other methods for analyzing pupil size, which tended to
ramp over trials. A wavelet was constructed by multiplying a complex sine wave (frequency = 5
trials) with a Gaussian envelope ($\mu = 0$, σ = cycles / (2π*frequency), cycles = 5; (Cohen, 2014)).
The wavelet was convolved with the baseline pupil size time series and the phase of the signal
was calculated on each trial (Matlab; angle). Standard circular statistics were used to measure
the differences between phase distributions for explore onsets and reward-matched controls
(Zar, 1999) and the phase alignment within these trial types (Berens, 2009). The latter was also
verified via comparison with bootstrapped null distributions (1000 samples).

Targeted dimensionality reduction. Neural state spaces have as many dimensions as there are
recorded neurons, but converging evidence suggests (1) that the neural states that are
observed in practice are generally confined to a lower-dimensional "manifold", and (2) that task-
relevant information is encoded by a small number of dimensions in that manifold. Because we
wanted to isolate the effects of arousal on choice-related activity from well-known effects of
arousal on neural activity (Ebitz and Platt, 2015; McGinley et al., 2015; Pfeffer et al., 2022;
Podvalny et al., 2021; Reimer et al., 2016, 2014; van Kempen et al., 2019; Waschke et al.,

2019), we focused all our neural population analyses on activity within the choice-predictive
subspace, rather than on neural activity more broadly.

To do this, we used targeted dimensionality reduction to identify the choice-predictive
dimensions of the neural state space (Cohen and Maunsell, 2010; Cunningham and Yu, 2014;
Ebitz et al., 2018; Peixoto et al., 2021). Specifically, we used multinomial logistic regression
(Matlab; mnrfit, mnrval, (Hastie et al., 2009)) to identify the separating hyperplanes that best
discriminated each choice from the alternative choices. This is equivalent to fitting a system of
binary classifiers of the form:

$$p(\text{choice} = \mathrm{k} \mid \mathrm{X}) = \frac{1}{1 - \exp^{-\beta_k \mathrm{X}}}$$

Where one classifier discriminates target 1 choices from targets 2 and 3 and a second
discriminates target 2 choices from targets 1 and 3. The classifier that discriminates target 3
from targets 1 and 2 is then just the negative of target 1 and target 2. These axes span the
subspace in which neural activity best predicts choice. Classifiers were trained on firing rates
from an epoch that began when the targets appeared and ended at the time of the saccade.
Mean imputation was used for the small number of occasions where a unit was not held for the
whole duration of the session (~3% of trials, ~12% of units) and a small fraction of units were
omitted from these analyses because their mean firing rates were less than 2 spikes/s, which
makes their weights difficult to identify (~8% of units).

Choice Probability Decoding. Within the choice-predictive subspace, the distance from the
separating hyperplanes (the vectors illustrated in **Figure 4E**) are the decoding vectors: the
vectors along which we can project neural activity in order to decode the log odds of choice.
This projection is equivalent to the decoded choice probability from the multinomial logistic
regression model and this is the figure we took as the decoded choice probability in **Figures 3F**
and **3H**. We evaluated decoding accuracy by measuring how often the most-likely choice
predicted by the model coincided with the choice the subject made.

Scatter index. The scatter index measures how much choice-predictive population neural
activity is clustered between trials with the same choice (Ebitz et al., 2018). It is calculated by
measuring the average Euclidean distance of each trial from all other trials where the same

choice was made and dividing it by the average Euclidean distance to all other trials where a
different choice was made:

$$\text{scatter} = \frac{\text{d}_{\text{within}}}{\text{d}_{\text{between}}}$$

Each trial thus has its own scatter index value, with a value of 1 indicating no difference in
clustering between same-choice and different-choice trials, and a value less than 1 indicating
greater clustering with same-choice trials compared to different-choice trials.

Neural speed. To determine how the speed of the decision-making process changed before and
during exploration, we calculated the rate of change in neural states within the choice-predictive
subspace during the first 400 ms following target presentation. Each trial's neural activity was
sampled in non-overlapping 20 ms bins and then projected into the choice-predictive subspace.
The change in neural activity within the subspace was then calculated between each pair of
samples. Finally, the changes were averaged together across the trial and normalized to the bin
width to produce an average rate of change in choice-predictive activity for that trial.

Acknowledgments

891 This work was supported by the Natural Sciences and Engineering Research Council of Canada
892 (Discovery Grant RGPIN-2020-05577), the Fonds de Recherche du Québec–Santé (Junior 1
893 Chercheur-Boursier 284309 to R.B.E.), the Jacobs Foundation (Research Fellowship, seed
894 grant to R.B.E.), and CIFAR Azrieli Global Scholars (seed grant to R.B.E.), the National Eye
895 Institute (R01-EY014924 to T.M.), and l'Institut de valorisation des données (fellowship to K.J.).

References

Aston-Jones G, Cohen JD. 2005. AN INTEGRATIVE THEORY OF LOCUS COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal Performance. *Annu Rev Neurosci* **28**:403–450. doi:10.1146/annurev.neuro.28.061604.135709

Berens P. 2009. CircStat: A MATLAB Toolbox for Circular Statistics. *J Stat Softw* **31**:1–21. doi:10.18637/jss.v031.i10

Bijleveld E, Custers R, Aarts H. 2009. The Unconscious Eye Opener: Pupil Dilation Reveals Strategic Recruitment of Resources Upon Presentation of Subliminal Reward Cues. *Psychol Sci* **20**:1313–1315. doi:10.1111/j.1467-9280.2009.02443.x

Borjon JI, Takahashi DY, Cervantes DC, Ghazanfar AA. 2016a. Arousal dynamics drive vocal production in marmoset monkeys. *J Neurophysiol* **116**:753–764. doi:10.1152/jn.00136.2016

Borjon JI, Takahashi DY, Cervantes DC, Ghazanfar AA. 2016b. Arousal dynamics drive vocal production in marmoset monkeys. *J Neurophysiol* **116**:753–764. doi:10.1152/jn.00136.2016

Bouret S, Sara SJ. 2005. Network reset: a simplified overarching theory of locus coeruleus noradrenaline function. *Trends Neurosci* **28**:574–582. doi:10.1016/j.tins.2005.09.002

Bradley MM, Miccoli L, Escrig MA, Lang PJ. 2008. The pupil as a measure of emotional arousal and autonomic activation. *Psychophysiology* **45**:602–607. doi:10.1111/j.1469-8986.2008.00654.x

Breton-Provencher V, Sur M. 2019. Active control of arousal by a locus coeruleus GABAergic circuit. *Nat Neurosci* **22**:218–228. doi:10.1038/s41593-018-0305-z

Bruce CJ, Goldberg ME. 1985. Primate frontal eye fields. I. Single neurons discharging before saccades. *J Neurophysiol* **53**:603–635. doi:10.1152/jn.1985.53.3.603

Burnham KP, Anderson DR, editors. 2004. Model Selection and Multimodel Inference. New York, NY: Springer. doi:10.1007/b97636

Chen CS, Knep E, Han A, Ebitz RB, Grissom NM. 2021. Sex differences in learning from exploration. *eLife* **10**:e69748. doi:10.7554/eLife.69748

Chen CS, Mueller D, Knep E, Ebitz RB, Grissom NM. 2023. Dopamine and norepinephrine differentially mediate the exploration-exploitation tradeoff. doi:10.1101/2023.01.09.523322

Clewett D, Gasser C, Davachi L. 2020. Pupil-linked arousal signals track the temporal organization of events in memory. *Nat Commun* **11**:4007. doi:10.1038/s41467-020-17851-9

Cohen MR, Maunsell JHR. 2010. A Neuronal Population Measure of Attention Predicts Behavioral Performance on Individual Trials. *J Neurosci* **30**:15241–15253. doi:10.1523/JNEUROSCI.2171-10.2010

Cohen MX. 2014. Analyzing neural time series data: theory and practice. MIT press.

Costa VD, Averbeck BB. 2020. Primate Orbitofrontal Cortex Codes Information Relevant for Managing Explore–Exploit Tradeoffs. *J Neurosci* **40**:2553–2561. doi:10.1523/JNEUROSCI.2355-19.2020

Costa VD, Rudebeck PH. 2016. More than Meets the Eye: the Relationship between Pupil Size and Locus Coeruleus Activity. *Neuron* **89**:8–10. doi:10.1016/j.neuron.2015.12.031

Cunningham JP, Yu BM. 2014. Dimensionality reduction for large-scale neural recordings. *Nat Neurosci* **17**:1500–1509. doi:10.1038/nn.3776

Daw ND, O'Doherty JP, Dayan P, Seymour B, Dolan RJ. 2006. Cortical substrates for exploratory decisions in humans. *Nature* **441**:876–879. doi:10.1038/nature04766

Dayan P, Daw ND. 2008. Decision theory, reinforcement learning, and the brain. *Cogn Affect*

Behav Neurosci **8**:429–453. doi:10.3758/CABN.8.4.429

de Gee JW, Tsetsos K, Schwabe L, Urai AE, McCormick D, McGinley MJ, Donner TH. 2020. Pupil-linked phasic arousal predicts a reduction of choice bias across species and decision domains. *eLife* **9**:e54014. doi:10.7554/eLife.54014

Ding L, Hikosaka O. 2006. Comparison of Reward Modulation in the Frontal Eye Field and Caudate of the Macaque. *J Neurosci* **26**:6695–6703. doi:10.1523/JNEUROSCI.0836-06.2006

Ebitz RB, Albarran E, Moore T. 2018. Exploration Disrupts Choice-predictive Signals and Alters Dynamics in Prefrontal Cortex. *Neuron* **97**:450-461.e9. doi:10.1016/j.neuron.2017.12.007

Ebitz RB, Hayden BY. 2021. The population doctrine in cognitive neuroscience. *Neuron* **109**:3055–3068. doi:10.1016/j.neuron.2021.07.011

Ebitz RB, Moore T. 2019. Both a Gauge and a Filter: Cognitive Modulations of Pupil Size. *Front Neurol* **9**.

Ebitz RB, Pearson JM, Platt ML. 2014. Pupil size and social vigilance in rhesus macaques. *Front Neurosci* **8**.

Ebitz RB, Platt ML. 2015. Neuronal Activity in Primate Dorsal Anterior Cingulate Cortex Signals Task Conflict and Predicts Adjustments in Pupil-Linked Arousal. *Neuron* **85**:628–640. doi:10.1016/j.neuron.2014.12.053

Ebitz RB, Sleezer BJ, Jedema HP, Bradberry CW, Hayden BY. 2019. Tonic exploration governs both flexibility and lapses. *PLOS Comput Biol* **15**:e1007475. doi:10.1371/journal.pcbi.1007475

Ebitz RB, Tu JC, Hayden BY. 2020. Rules warp feature encoding in decision-making circuits. *PLOS Biol* **18**:e3000951. doi:10.1371/journal.pbio.3000951

Eldar E, Cohen JD, Niv Y. 2013. The effects of neural gain on attention and learning. *Nat Neurosci* **16**:1146–1153. doi:10.1038/nn.3428

Fiete IR, Fee MS, Seung HS. 2007. Model of Birdsong Learning Based on Gradient Estimation by Dynamic Perturbation of Neural Conductances. *J Neurophysiol* **98**:2038–2057. doi:10.1152/jn.01311.2006

Filipowicz AL, Glaze CM, Kable JW, Gold JI. 2020. Pupil diameter encodes the idiosyncratic, cognitive complexity of belief updating. *eLife* **9**:e57872. doi:10.7554/eLife.57872

Gershman SJ. 2019. Uncertainty and Exploration. *Decis Wash DC* **6**:277–286. doi:10.1037/dec0000101

Gilzenrat MS, Nieuwenhuis S, Jepma M, Cohen JD. 2010. Pupil diameter tracks changes in control state predicted by the adaptive gain theory of locus coeruleus function. *Cogn Affect Behav Neurosci* **10**:252–269. doi:10.3758/CABN.10.2.252

Graves JE, Egré P, Pressnitzer D, de Gardelle V. 2021. An implicit representation of stimulus ambiguity in pupil size. *Proc Natl Acad Sci* **118**:e2107997118. doi:10.1073/pnas.2107997118

Hastie T, Tibshirani R, Friedman J. 2009. The Elements of Statistical Learning, Springer Series in Statistics. New York, NY: Springer. doi:10.1007/978-0-387-84858-7

Hauser CK, Zhu D, Stanford TR, Salinas E. 2018. Motor selection dynamics in FEF explain the reaction time variance of saccades to single targets. *eLife* **7**:e33456. doi:10.7554/eLife.33456

Japundzic N, Grichois M-L, Zitoun P, Laude D, Elghozi J-L. 1990. Spectral analysis of blood pressure and heart rate in conscious rats: effects of autonomic blockers. *J Auton Nerv Syst* **30**:91–100. doi:10.1016/0165-1838(90)90132-3

Jepma M, Nieuwenhuis S. 2011a. Pupil Diameter Predicts Changes in the Exploration–Exploitation Trade-off: Evidence for the Adaptive Gain Theory. *J Cogn Neurosci* **23**:1587–1596. doi:10.1162/jocn.2010.21548

Jepma M, Nieuwenhuis S. 2011b. Pupil diameter predicts changes in the exploration–exploitation trade-off: Evidence for the adaptive gain theory. *J Cogn Neurosci* **23**:1587–1596.

Joshi S, Gold JI. 2020. Pupil Size as a Window on Neural Substrates of Cognition. *Trends Cogn Sci* **24**:466–480. doi:10.1016/j.tics.2020.03.005

Joshi S, Li Y, Kalwani R, Gold JI. 2016. Relationships between pupil diameter and neuronal activity in the locus coeruleus, colliculi, and cingulate cortex. *Neuron* **89**:221–234. doi:10.1016/j.neuron.2015.11.028

Julien C. 2020. An update on the enigma of Mayer waves. *Cardiovasc Res* **116**:e210–e211. doi:10.1093/cvr/cvz327

Julien C. 2006. The enigma of Mayer waves: Facts and models. *Cardiovasc Res* **70**:12–21. doi:10.1016/j.cardiores.2005.11.008

Jurewicz K, Sleezer BJ, Mehta PS, Hayden BY, Ebitz RB. 2022. Irrational choices via a curvilinear representational geometry for value. doi:10.1101/2022.03.31.486635

Kamiya A, Hayano J, Kawada T, Michikami D, Yamamoto K, Ariumi H, Shimizu S, Uemura K, Miyamoto T, Aiba T, Sunagawa K, Sugimachi M. 2005. Low-frequency oscillation of sympathetic nerve activity decreases during development of tilt-induced syncope preceding sympathetic withdrawal and bradycardia. *Am J Physiol-Heart Circ Physiol* **289**:H1758–H1769. doi:10.1152/ajpheart.01027.2004

Kane GA, Vazey EM, Wilson RC, Shenhav A, Daw ND, Aston-Jones G, Cohen JD. 2017. Increased locus coeruleus tonic activity causes disengagement from a patch-foraging task. *Cogn Affect Behav Neurosci* **17**:1073–1083. doi:10.3758/s13415-017-0531-y

Kaske EA, Chen CS, Meyer C, Yang F, Ebitz B, Grissom N, Kapoor A, Darrow DP, Herman AB. 2022. Prolonged physiological stress is associated with a lower rate of exploratory learning that is compounded by depression. *Biol Psychiatry Cogn Neurosci Neuroimaging*. doi:10.1016/j.bpsc.2022.12.004

Kleiner M, Brainard DH, Pelli D. 2007. What's new in Psychtoolbox-3? *Perception* **36**:1–16.

Koss MC. 1986. Pupillary dilation as an index of central nervous system α2-adrenoceptor activation. *J Pharmacol Methods* **15**:1–19. doi:10.1016/0160-5402(86)90002-1

Lau B, Glimcher PW. 2008. Value Representations in the Primate Striatum during Matching Behavior. *Neuron* **58**:451–463. doi:10.1016/j.neuron.2008.02.021

Laurie V-J, Shourkeshti A, Chen CS, Herman A, Grissom N, Ebitz RB. 2025. Persistent Decision-Making in Mice, Monkeys, and Humans. *bioRxiv* 2024.05.07.592970. doi:10.1101/2024.05.07.592970

Loewenfeld IE. 1999. The Pupil: Anatomy, Physiology, and Clinical Applications. Butterworth-Heinemann.

Manella LC, Petersen N, Linster C. 2017. Stimulation of the Locus Ceruleus Modulates Signal-to-Noise Ratio in the Olfactory Bulb. *J Neurosci* **37**:11605–11615. doi:10.1523/JNEUROSCI.2026-17.2017

Martins ARO, Froemke RC. 2015. Coordinated forms of noradrenergic plasticity in the locus coeruleus and primary auditory cortex. *Nat Neurosci* **18**:1483–1492. doi:10.1038/nn.4090

McGinley MJ, David SV, McCormick DA. 2015. Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. *Neuron* **87**:179–192. doi:10.1016/j.neuron.2015.05.038

Moore T, Armstrong KM. 2003. Selective gating of visual signals by microstimulation of frontal cortex. *Nature* **421**:370–373. doi:10.1038/nature01341

Moore T, Fallah M. 2001. Control of eye movements and spatial attention. *Proc Natl Acad Sci U S A* **98**:1273–1276. doi:10.1073/pnas.98.3.1273

Muller TH, Mars RB, Behrens TE, O'Reilly JX. 2019. Control of entropy in neural models of

environmental state. *eLife* **8**:e39404. doi:10.7554/eLife.39404

Murphy PR, Vandekerckhove J, Nieuwenhuis S. 2014. Pupil-Linked Arousal Determines Variability in Perceptual Decision Making. *PLOS Comput Biol* **10**:e1003854. doi:10.1371/journal.pcbi.1003854

Nassar MR, Rumsey KM, Wilson RC, Parikh K, Heasly B, Gold JI. 2012. Rational regulation of learning dynamics by pupil-linked arousal systems. *Nat Neurosci* **15**:1040–1046. doi:10.1038/nn.3130

O'Byrne J, Jerbi K. 2022. How critical is brain criticality? *Trends Neurosci*. doi:10.1016/j.tins.2022.08.007

O'Reilly JX, Schüffelgen U, Cuell SF, Behrens TEJ, Mars RB, Rushworth MFS. 2013. Dissociable effects of surprise and model update in parietal and anterior cingulate cortex. *Proc Natl Acad Sci* **110**:E3660–E3669. doi:10.1073/pnas.1305373110

Pearson JM, Hayden BY, Raghavachari S, Platt ML. 2009. Neurons in posterior cingulate cortex signal exploratory decisions in a dynamic multi-option choice task. *Curr Biol CB* **19**:1532–1537. doi:10.1016/j.cub.2009.07.048

Peixoto D, Verhein JR, Kiani R, Kao JC, Nuyujukian P, Chandrasekaran C, Brown J, Fong S, Ryu SI, Shenoy KV, Newsome WT. 2021. Decoding and perturbing decision states in real time. *Nature* **591**:604–609. doi:10.1038/s41586-020-03181-9

Pfeffer T, Keitel C, Kluger DS, Keitel A, Russmann A, Thut G, Donner TH, Gross J. 2022. Coupling of pupil- and neuronal population dynamics reveals diverse influences of arousal on cortical processing. *eLife* **11**:e71890. doi:10.7554/eLife.71890

Pisupati S, Chartarifsky-Lynn L, Khanal A, Churchland AK. 2021. Lapses in perceptual decisions reflect exploration. *eLife* **10**:e55490. doi:10.7554/eLife.55490

Platt ML, Glimcher PW. 1999. Neural correlates of decision variables in parietal cortex. *Nature* **400**:233–238. doi:10.1038/22268

Podvalny E, King LE, He BJ. 2021. Spectral signature and behavioral consequence of spontaneous shifts of pupil-linked arousal in human. *eLife* **10**:e68265. doi:10.7554/eLife.68265

Preacher KJ, Rucker DD, Hayes AF. 2007. Addressing Moderated Mediation Hypotheses: Theory, Methods, and Prescriptions. *Multivar Behav Res* **42**:185–227. doi:10.1080/00273170701341316

Preuschoff K, 't Hart B, Einhauser W. 2011. Pupil Dilation Signals Surprise: Evidence for Noradrenaline's Role in Decision Making. *Front Neurosci* **5**.

Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS. 2014. Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. *Neuron* **84**:355–362. doi:10.1016/j.neuron.2014.09.033

Reimer J, McGinley MJ, Liu Y, Rodenkirch C, Wang Q, McCormick DA, Tolias AS. 2016. Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. *Nat Commun* **7**:13289. doi:10.1038/ncomms13289

Roesch MR, Olson CR. 2007. Neuronal Activity Related to Anticipated Reward in Frontal Cortex. *Ann N Y Acad Sci* **1121**:431–446. doi:10.1196/annals.1401.004

Sadacca B, Wikenheiser AM, Schoenbaum G. 2017. Towards a theoretical role for tonic norepinephrine in the orbitofrontal cortex in facilitating flexible learning. *Neuroscience* **345**:124–129. doi:10.1016/j.neuroscience.2016.04.017

Schall JD, Hanes DP. 1993. Neural basis of saccade target selection in frontal eye field during visual search. *Nature* **366**:467–469. doi:10.1038/366467a0

Scheffer M. 2020. Critical Transitions in Nature and Society, Princeton Studies in Complexity. Princeton University Press.

Scheffer M, Bascompte J, Brock WA, Brovkin V, Carpenter SR, Dakos V, Held H, van Nes EH, Rietkerk M, Sugihara G. 2009. Early-warning signals for critical transitions. *Nature*

461:53–59. doi:10.1038/nature08227

Schultz W, Preuschoff K, Camerer C, Hsu M, Fiorillo CD, Tobler PN, Bossaerts P. 2008. Explicit neural signals reflecting reward uncertainty. *Philos Trans R Soc B Biol Sci* **363**:3801–3811. doi:10.1098/rstb.2008.0152

Slooten JCV, Jahfari S, Knapen T, Theeuwes J. 2018. How pupil responses track value-based decision-making during and after reinforcement learning. *PLOS Comput Biol* **14**:e1006632. doi:10.1371/journal.pcbi.1006632

Sobel ME. 1986. Some New Results on Indirect Effects and Their Standard Errors in Covariance Structure Models. *Sociol Methodol* **16**:159–186. doi:10.2307/270922

Tervo DGR, Proskurin M, Manakov M, Kabra M, Vollmer A, Branson K, Karpova AY. 2014. Behavioral Variability through Stochastic Choice and Its Gating by Anterior Cingulate Cortex. *Cell* **159**:21–32. doi:10.1016/j.cell.2014.08.037

Urai AE, Braun A, Donner TH. 2017. Pupil-linked arousal is driven by decision uncertainty and alters serial choice bias. *Nat Commun* **8**:14637. doi:10.1038/ncomms14637

van Kempen J, Loughnane GM, Newman DP, Kelly SP, Thiele A, O'Connell RG, Bellgrove MA. 2019. Behavioural and neural signatures of perceptual decision-making are modulated by pupil-linked arousal. *eLife* **8**:e42541. doi:10.7554/eLife.42541

Wang R, Dearing JA, Langdon PG, Zhang E, Yang X, Dakos V, Scheffer M. 2012. Flickering gives early warning signals of a critical transition to a eutrophic lake state. *Nature* **492**:419–422. doi:10.1038/nature11655

Waschke L, Tune S, Obleser J. 2019. Local cortical desynchronization and pupil-linked arousal differentially shape brain states for optimal sensory performance. *eLife* **8**:e51501. doi:10.7554/eLife.51501

Wilson RC, Bonawitz E, Costa VD, Ebitz RB. 2021. Balancing exploration and exploitation with information and randomization. *Curr Opin Behav Sci*, Computational cognitive neuroscience **38**:49–56. doi:10.1016/j.cobeha.2020.10.001

Wilson RC, Geana A, White JM, Ludwig EA, Cohen JD. 2014. Humans use directed and random exploration to solve the explore-exploit dilemma. *J Exp Psychol Gen* **143**:2074–2081. doi:10.1037/a0038199

Wu HG, Miyamoto YR, Castro LNG, Ölveczky BP, Smith MA. 2014. Temporal structure of motor variability is dynamically regulated and predicts motor learning ability. *Nat Neurosci* **17**:312–321. doi:10.1038/nn.3616

Yao T, Vanduffel W. 2023. Spike rates of frontal eye field neurons predict reaction times in a spatial attention task. *Cell Rep* **42**:112384. doi:10.1016/j.celrep.2023.112384

Yerkes RM, Dodson JD. 1908. The relation of strength of stimulus to rapidity of habit-formation. *J Comp Neurol Psychol* **18**:459–482. doi:10.1002/cne.920180503

Yokoi A, Weiler J. 2022. Pupil diameter tracked during motor adaptation in humans. *J Neurophysiol* **128**:1224–1243. doi:10.1152/jn.00021.2022

Zar JH. 1999. Biostatistical analysis. Pearson Education India.

896 **Table S1**

897 Regression coefficients and p values for the mediation analysis testing whether the scatter
898 index mediates the relationship between pupil size and the onset of exploration. Related to
899 **Figure 4J**.

		$\text{pupil}_{t-1} \rightarrow \text{scatter}_{t-1} \rightarrow \text{explore}_t$	
		est. coefficient	p value <
Total effect	c	0.090	0.005
Effect on mediator	a	0.036	0.0005
Unique mediator effect	b	0.092	0.005
Indirect effect	ab	0.003 (z = 2.70*)	0.005
Direct effect	c'	0.086	0.005

900 *Sobel's test

902 **Table S2**

903 Regression coefficients and p values for the mediation analysis testing whether response time
904 slowing mediates the relationship between pupil size and the onset of exploration on the next
905 trial. Related to **Figure 5C**.

		$\text{pupil}_{t-1} \rightarrow \text{RT slowing}_{t-1} \rightarrow \text{explore}_t$	
		est. coefficient	p value <
Total effect	c	0.090	0.005
Effect on mediator	a	0.078	0.0001
Unique mediator effect	b	0.106	0.0005
Indirect effect	ab	0.008 (z = 3.48*)	0.0005
Direct effect	c'	0.080	0.01

906 *Sobel's test

907 **Table S3**

908 Regression coefficients and p values for the mediation analysis testing whether neural slowing
909 mediates the relationship between pupil size and the onset of exploration on the next trial.
910 Related to **Figure 5F**.

		$pupil_{t-1} \rightarrow$ neural $slowing_{t-1} \rightarrow explore_t$	
		est. coefficient	p value <
Total effect	c	0.095	0.005
Effect on mediator	a	-0.025	0.0005
Unique mediator effect	b	-0.059	0.06
Indirect effect	ab	0.001 (z = 1.69*)	0.05
Direct effect	c'	0.093	0.005

911 *Sobel's test
912
913

 Figure S1
915

916
917　**Figure S1.** Same as **Figure 1E**, but without first explore trials.

918 **Figure S2**

919
Figure S2. Decoded choice probability and scatter index across behavioral states and pupil
size. (A) Decoded choice probability (projection onto the correct coding dimension) for exploit
and explore states. Dots represent individual sessions, with lines connecting values from the
same session across states. (B) Decoded choice probability plotted as a function of pupil size
quantile for explore trials alone. (C) Scatter index, a measure of variance in choice-predictive
population activity, for exploit and explore states, with lines connecting values from the same
session across states. (D) The scatter index plotted as a function of pupil size quantile for
explore trials.
928