Title:

Pupil size predicts the onset of exploration in brain and behavior.

Authors:

Akram Shourkeshti[1], Gabriel Marrocco[1], Katarzyna Jurewicz[1,2], Tirin Moore[3,4], R. Becket Ebitz*[1]

Affiliations:

[1]Department of Neurosciences, Université de Montréal, Montréal, QC, Canada

[2]Department of Physiology, McGill University, Montréal, QC, Canada

[3]Department of Neurobiology, Stanford University School of Medicine, Stanford, CA, USA

[4]Howard Hughes Medical Institute, Chevy Chase, MD, USA

*Corresponding author and lead contact:

R. Becket Ebitz

Department of Neurosciences

Université de Montréal

Montréal, QC CANADA H3T 1J4

Email: becket@ebitzlab.com

Abstract

In uncertain environments, intelligent decision-makers exploit actions that have been rewarding in the past, but also explore actions that could be even better. Several neuromodulatory systems are implicated in exploration, based, in part, on work linking exploration to pupil size–a peripheral correlate of neuromodulatory tone and index of arousal. However, pupil size could instead track variables that make exploration more likely, like volatility or reward, without directly predicting either exploration or its neural bases. Here, we simultaneously measured pupil size, exploration, and neural population activity in the prefrontal cortex while two rhesus macaques explored and exploited in a dynamic environment. We found that pupil size under constant luminance specifically predicted the onset of exploration, beyond what could be explained by reward history. Pupil size also predicted disorganized patterns of prefrontal neural activity at both the single neuron and population levels, even within periods of exploitation. Ultimately, our results support a model in which pupil-linked mechanisms promote the onset of exploration via driving the prefrontal cortex through a critical tipping point where prefrontal control dynamics become disorganized and exploratory decisions are possible.

Introduction

Many decisions maximize immediate rewards. However, in uncertain or changing environments, it is important to occasionally sacrifice some immediate rewards in the service of long-term goals. This gives us an opportunity to learn about the value of alternative options and discover new, more valuable strategies for interacting with the world. In short, in complex environments, intelligent decision-makers exploit rewarding strategies, but also explore alternative strategies that could be even better.

Because exploitation maximizes immediate reward, it can rely on the same value-based decision-making processes that have been the subject of neurobiological studies for decades (Ding and Hikosaka, 2006; Jurewicz et al., 2022; Platt and Glimcher, 1999; Roesch and Olson, 2007; Schultz et al., 2008). However, we are only just beginning to understand the neural bases of exploration (Costa and Averbeck, 2020; Daw et al., 2006; Pearson et al., 2009; Wilson et al., 2021, 2014). One clue is that many organisms seem to explore via random sampling (Gershman, 2019; Wilson et al., 2021, 2014). Randomness is a critical component of exploratory discovery in bird song and motor learning (Fiete et al., 2007; Wu et al., 2014), can perform about as well as more sophisticated exploratory strategies in many environments (Dayan and Daw, 2008), and humans and other primates tend to explore randomly even when more sophisticated strategies are available (Ebitz et al., 2018; Wilson et al., 2014). There is some evidence linking random exploration to disorganized patterns of activity in the prefrontal cortex (Ebitz et al., 2019, 2018; Muller et al., 2019; Wilson et al., 2021), but more work is needed to understand how these disorganized patterns of activity emerge and what neurobiological processes drive their emergence.

One promising hypothesis is that random exploration and its neural correlates could be under the control of some process(es) linked to pupil size. Pupil size under constant luminance is a

peripheral index of autonomic arousal (Bradley et al., 2008; Ebitz and Moore, 2019; Loewenfeld,
1999) that also predicts widespread changes in neural population activity (McGinley et al., 2015;
Reimer et al., 2014)–including in regions implicated in noisy decision-making (Ebitz and Platt,
2015; Tervo et al., 2014). Among other neuromodulators (Gilzenrat et al., 2010; Koss, 1986;
Reimer et al., 2016), pupil size is correlated with central norepinephrine (Costa and Rudebeck,
2016; Joshi et al., 2016): a catecholamine that flattens neuronal tuning functions (Martins and
Froemke, 2015) and predicts abrupt "resets" in cortical networks (Aston-Jones and Cohen,
2005; Bouret and Sara, 2005). Behaviorally, pupil size predicts noisy decision-making (Aston-
Jones and Cohen, 2005; Ebitz et al., 2014; Eldar et al., 2013; Gilzenrat et al., 2010; O'Reilly et
al., 2013; Wilson et al., 2021), including errors of reward-maximization (Jepma and
Nieuwenhuis, 2011) and task performance (Ebitz et al., 2014; Ebitz and Platt, 2015), at least
some of which are likely to be caused by exploratory processes (Ebitz et al., 2019; Jepma and
Nieuwenhuis, 2011; Pisupati et al., 2021).

While a growing body of circumstantial evidence implicates pupil-linked mechanisms in
exploration, there is also a plausible alternative interpretation of these results: perhaps pupil
size simply tracks the variables that make exploration more likely without actually predicting
exploration. Pupil size under constant luminance increases with some variables that should
make exploration more likely, including the volatility of the environment, the surprise of reward
outcomes, novelty, uncertainty, and context changes (Clewett et al., 2020; Filipowicz et al.,
2020; Graves et al., 2021; Preuschoff et al., 2011; Slooten et al., 2018; Yokoi and Weiler, 2022).
Critically, in circumstances where these variables change behavior (i.e. increasing learning,
increasing decision noise, inducing exploration), it is not clear whether the pupil is signaling
these variables or instead directly predicting behavioral change (Nassar et al., 2012; O'Reilly et
al., 2013; Urai et al., 2017). Fortunately, recent results suggest that at least some exploration
appears to occur tonically, regardless of these variables (Ebitz et al., 2019; Pisupati et al., 2021;
Wilson et al., 2021). In parallel, a new computational approach now allows us to determine
when exploration is occurring independently of the reward-based computations thought to drive
it (Chen et al., 2021; Ebitz et al., 2020, 2019, 2018). These advances mean that it is now
possible to determine whether pupil size predicts exploration itself or instead simply tracks the
variables that make exploration more likely.

Here, we measured pupil size and recorded from populations of prefrontal neurons while two
rhesus macaques performed a task that encouraged both exploration and exploitation. To infer
whether the subjects were exploring or exploiting, we modeled exploration and exploitation as
the latent goal states underlying decision-making (Chen et al., 2021; Ebitz et al., 2020, 2019,
2018). Pupil size under constant luminance was larger during explore choices than exploit
choices, consistent with both the idea that pupil-linked processes are the proximate cause of
exploration and the idea that pupil size tracks variables that make exploration more likely.
However, the temporal relationship between pupil size and exploration was both precise and
complex, with pupil size explaining variability in brain and behavior that could not be explained
by rewards or time. Together, these results support the hypothesis that pupil-linked processes
drive the prefrontal cortex through a critical tipping point that permits exploratory decisions.

Results

Two male rhesus macaques performed a total of 28 sessions of a classic explore/exploit task: a restless three-armed bandit (subject B: 10 sessions, subject O: 18 sessions; a total of 21,793 trials). We have previously analyzed parts of this dataset (Ebitz et al., 2018), but not the pupil size data, and all analyses presented here are new. In this task, the reward probability (value) of three targets walks randomly and independently over time (**Figure 1A**). This means that the subjects have to take advantage of valuable options when they are available (exploit), but also occasionally sample alternative options to determine if they have become more valuable (explore).

Rather than instructing subjects to explore and exploit, this task takes advantage of the subjects' natural tendency to alternate between exploration and exploitation in a changing environment. We have previously shown that both monkeys and mice exhibit 2 behavioral modes in this task: one exploitative mode in which they repeatedly choose the same option—learning little but maximizing reward—and one exploratory mode in which they alternate rapidly between the options—choosing randomly with respect to rewards and learning rapidly (Chen et al., 2021; Ebitz et al., 2018). We infer which of these modes is driving behavior with a hidden Markov model (HMM; **Figure 1B**). This approach models the exploratory and exploitative modes as latent goal states and the maximum *a posteriori* goal is taken as the state label for each choice. We have previously shown that this method identifies explore/exploit state labels that match normative definitions (Chen et al., 2021; Ebitz et al., 2018) and explain variance in prefrontal neural activity that cannot be explained by reward value, reward history, and switch/stay decisions (Ebitz et al., 2018).

Some previous studies used a different method to identify exploratory choices (Daw et al., 2006; Jepma and Nieuwenhuis, 2011; Pearson et al., 2009). This method fits a reinforcement learning (RL) model to the behavior and identifies the choices that are not consistent with the model's subjective values as exploratory. However, this approach (1) equates exploration with errors of reward maximization, not a goal that is orthogonal to reward maximization, and (2) its accuracy depends on precise knowledge of the computations involved in the choice, which are highly variable, both across individuals and over time (Chen et al., 2021, 2021; Kaske et al., 2022). The HMM approach, conversely, makes no assumptions about the computations involved in the choice and identifies choices that are orthogonal to reward value, not anti-correlated with it (Chen et al., 2021; Ebitz et al., 2018). Here, we found that state labels from the HMM method explained more variance in behavior and neural activity than choice labels from the previous, RL method (**Figure 1C;** response time: both subjects, paired t-test: $p < 0.005$, $t(27) = 3.41$, the mean difference of beta weights = 0.004, 95% CI = 0.002 to 0.007; scatter index [(Ebitz et al., 2018)]: both subjects, paired t-test: $p < 0.001$, $t(27) = 3.84$, the mean difference of beta weights = 0.15, 95% CI = 0.07 to 0.24: see **Methods**). In short, we find that the HMM approach is a more robust and accurate method, with better face validity, than the RL-based method for identifying explore choices.

Figure 1. Task design and pupil. *A) Top: Subjects made saccadic choices between three identical options. One of the options was in the receptive field of an FEF neuron (dotted circle). Bottom: Reward probabilities for the 3 options (lines), with choices overlaid (dots) for 200 example trials. Gray bars = explore-labels. B) The HMM models exploration and exploitation as latent goal states underlying choice sequences. C) Comparing the regression coefficients HMM-inferred and RL-inferred explore choices for a measure of the disorganization of neural population responses ("scatter index"; see Methods; (Ebitz et al., 2018)) and response time. D) Average pupil size on explore and exploit choices. Right: Same for individual subjects. E) The probability of explore choices as a function of pupil size quantile. Dotted line: linear GLM fit. Solid line: quadratic fit. Right: Same for individual subjects. F) Several behavior measures compared across median-split large- and small-pupil-size explore choices. Left to right: reward probability, a one-trial-back learning index (see Methods), saccadic peak velocity of saccades, and the scatter index. No significant differences between pupil bins. The blue line is the mean ± SEM for exploit choices. Error bars depict ± SEM throughout.*

Using the previous method, Jepma & Nieuwenhuis (2011) reported that pupil size under
constant luminance is larger during explore choices compared to exploit ones. Therefore, we
first asked if this is also true using the HMM method. Indeed, we found that pupil size at fixation
(see **Methods**) was larger on explore-labeled trials than exploit-labeled trials in both subjects
(**Figure 1D**; both subjects, paired t-test: $p < 0.0001$, $t(27) = 4.95$, mean offset = 0.23, 95% CI =
0.13 to 0.32; subject B: $p < 0.001$, $t(9) = 5.50$, mean offset = 0.4, 95% CI = 0.24 to 0.57; subject
O: $p < 0.02$, $t(17) = 2.85$, mean offset = 0.13, 95% CI = 0.03 to 0.23). However, the probability of
exploration did not increase linearly as a function of pupil size (**Figure 1E**). Although larger pupil
size generally predicted more explore choices (both subjects, 1st order GLM: beta = 0.2, $p <$
0.0001, n = 28 sessions; subject B: 1st order GLM, beta = 0.34, $p < 0.0001$, n = 10; subject O:
1st order GLM, beta = 0.13, $p < 0.0001$, n = 18), the relationship was obviously nonlinear in both
subjects (2nd order, quadratic model was a better fit in both animals, 2nd order GLM, beta 1 =
0.16, $p < 0.0001$, beta 2 = 0.04, $p < 0.005$, n = 28 sessions; subject B: 2nd order GLM, beta 1 =
0.27, $p < 0.0001$, beta 2 = 0.05, $p < 0.05$, n = 10; subject O: 2nd order GLM, beta 1 = 0.11, $p <$
0.0001, beta 2 = 0.02, $p > 0.1$, n = 18; linear model AIC = 17319.5, quadratic model AIC =
17311.7, AIC weight [relative likelihood] of linear model = 0.02; see **Methods**). This was not an
artifact of some systematic error in the HMM fits: there was also a U-shaped relationship
between pupil size and the probability of switching (both subjects, 1st order GLM: beta = 0.2 $p <$
0.0001, n = 28 sessions; subject B: 1st order GLM, beta = 0.31, $p < 0.0001$; subject O: 1st order
GLM, beta = 0.18, $p < 0.0001$, 2nd order, quadratic model was a better fit in both animals, 2nd
order GLM, beta 1 = 0.16 $p < 0.0001$, beta 2 = 0.05, $p < 0.0001$; subject B: 2nd order GLM, beta
1 = 0.18, $p < 0.0005$, beta 2 = 0.09, $p < 0.0005$; subject O: 2nd order GLM, beta 1 = 0.15, $p <$
0.0001, beta 2 = 0.03, $p < 0.01$; linear model AIC = 16407.9, quadratic model AIC = 16390.6,
AIC weight of linear model = 0.0001). Thus, although pupil size was generally larger during
exploration, compared to exploitation, the relationship between pupil size and exploration was
U-shaped.

A U-shaped relationship would be expected if some "explore" choices (or switch choices) were
not real explore choices, but instead the result of disengagement at low levels of arousal.
However, if this were the case, then the valid, large-pupil explore choices would systematically
differ from the false, small-pupil "explore" choices. They did not. Small- and large-pupil explore
choices (median split) were indistinguishable along several of the key dimensions that
differentiate explore choices from exploit choices (**Figure 1F**). For example, both were equally
likely to be rewarded (mean difference = 0.03 ± 0.24 STD) between large- and small pupil-
explore choices ($p > 0.4$, $t(1,27) = 0.75$, paired t-test; AUC for discriminating explore and exploit
= 0.65 ± 0.05 STD across sessions). Both had similar peak saccadic velocities (mean difference
= -0.05 ± 0.23 STD, $p > 0.2$, $t(27) = -1.08$; explore/exploit AUC = 0.61 ± 0.10 STD) and both had
more variability in neural population choice information ("scatter index", mean difference = 0.03
± 0.33 STD, $p > 0.6$, $t(27) = 0.45$; explore/exploit AUC = 0.60 ± 0.07 STD). Both had similar
levels of reward learning (see **Methods**; the mean difference = -0.03 ± 0.57 STD, $p > 0.7$, $t(27)$
= 0.27): in both cases, learning was substantially enhanced relative to the exploit choices
(small-pupil, the mean difference from exploit = 0.24 ± 0.48 STD, $p < 0.02$, $t(27) = 2.69$; large-
pupil, the mean difference from exploit = 0.21 ± 0.39 STD, $p < 0.01$, $t(27) = 2.91$). These results
are incompatible with the idea that either type of explore choice reflects disengagement in the

task or that small- and large-pupil explore choices have different causes. Instead, we will see
that the U-shape was due to the complex temporal relationship between pupil size and
exploration.

Pupil size ramped up across trials before explore choices in both subjects and shrank down to
below-baseline levels when exploitation began (**Figure 2A**). This ramping meant that pupil size
was larger not just during exploration, but also during the exploit choices immediately before
exploration (both subjects, GLM slope = 0.01, $p < 0.005$, n = 28; subject B: beta = 0.02, $p <$
0.02, n = 10; subject O: beta = 0.01, $p < 0.05$, n = 18; average pupil size compared to the exploit
choices, post-hoc paired t-tests, 1 trial before exploration mean = 0.12, $p < 0.005$, $t(27) = 3.42$;
2 trials mean = 0.09, $p < 0.03$, $t(27) = 2.41$; 3 trials mean = 0.03, $p > 0.1$, $t(27) = 1.42$; 4 trials
mean = 0.05, $p < 0.05$, $t(27) = 2.07$). By the first exploit choice after exploration, pupil size had
already begun shrinking to below-baseline levels (post-hoc paired t-tests, 1 trial after exploration
mean = 0.03, $p = 0.09$, $t(27) = 1.73$; 2 trials after mean = -0.11, $p < 0.02$, $t(27) = -2.67$; 3 trials
after mean = -0.16, $p < 0.02$, $t(27) = -2.72$; 4 trials after mean = -0.08, $p > 0.2$, $t(27) = -1.27$; 5
trials after mean = -0.16, $p < 0.001$, $t(27) = -3.94$; p-values are significant with a Holm-
Bonferroni correction). The shrinking to below-baseline levels could suggest a refractory
mechanism that would prevent exploration from re-occurring immediately after it happened.

Neither the ramping in pupil size before exploration nor the shrinking after were artifacts of
some misalignment in the model's labels. We saw no evidence of ramping in peak saccadic
velocity, another behavioral measure that differentiated explore trials and exploit trials (**Figure
2B**; no significant decrease from baseline 1 trial before, paired t-test: $p > 0.7$, $t(27) = 0.42$; a
GLM was nonsignificant with the trend pointing in the opposite direction: 10 trials preceding
exploration, beta = 0.008, $p > 0.1$) and no significant change from baseline afterward (not
greater than the baseline during the 5 trials after exploration, when pupil shrinking was maximal,
mean = -0.47 ± 0.15 STD, $p > 0.9$, one-sided $t(27) = -1.67$). We previously reported similar
results for certain neural measures (Ebitz et al., 2018). Thus, while pupil size ramped before the
onset of exploration and shrank afterward, the same was not true of other behavioral and neural
variables, suggesting that these dynamics were not some artifact of misalignment.
227
Next, we examined the extent to which these across-trial pupil dynamics were affected by
exploration (see **Methods**). When the subjects did not explore the pupil size increased steadily
across trials (**Figure 2C**; both subjects, GLM: beta = 0.004, $p < 0.0001$; subject B: beta = 0.003,
$p < 0.0001$; subject O: beta = 0.005, $p < 0.0001$, n = 25 lags over 28 sessions). This implies that
the ramping in pupil size before explore choices may be a general dynamic of how pupil size
evolves in the absence of exploration. However, a different pattern emerged when we looked at
how the pupil changed between exploit trials that were separated by exploration. When 2 exploit
trials were separated by at least 1 explore choice, pupil size was smaller on the second exploit
trial (both subjects, GLM: beta = -0.09, $p < 0.0001$; subject B: beta = -0.14, $p < 0.0001$; subject
O: beta = -0.07, $p < 0.0001$). Critically, passing through exploration only produced a baseline
decrease in pupil size but did not alter the rate at which pupil size grew over trials (no significant
interaction between slope and condition in both subjects, GLM: beta < -0.0001, $p > 0.9$; subject
B: beta = 0.003, $p < 0.05$; subject O: beta = -0.002, $p > 0.1$; also nonsignificant on trials 5-25:

both subjects: beta < 0.0005, p > 0.5). Therefore, pupil size tended to ramp across trials but
exploit choices temporarily decreased pupil size without disrupting this ramping in the long term.
243

Figure 2. Pupil size ramps up before exploration and shrinks down after.
A) Average pupil size for 10 trials before and 10 trials after explore choices. Purple line: GLM fit. Right: Same for each subject separately. B) Same as Figures 1D and 2A but for peak velocity rather than pupil size. C) Change in pupil size between exploit trials that are either in a single bout of exploitation (gray) or separated by explore trials (purple). Right: Same for each subject separately. D) Change in pupil size over certain pairs of trials: starting (exploit to explore), during (explore to explore), and leaving (explore to exploit) exploration. *p < 0.001 E) The probability of starting to explore as a function of pupil size quantile. Solid line: Linear GLM fit. Error bars and shaded regions depict mean ± SEM.

Pupil size was generally smaller after explore choices, but was this driven by the onset of
exploration or the onset of exploitation? If the pupil starts to shrink only after exploration ends, it
would support models that suggest that pupil size decreases with commitment to a new option
or belief state (O'Reilly et al., 2013). Conversely, if the pupil shrinks immediately after
exploration begins, it might suggest that pupil-linked mechanisms are important for initiating
exploration, but not sustaining it. Our results were consistent with the latter hypothesis: the pupil
immediately began shrinking as soon as exploration began, not when it ended (**Figure 2D**;
mean change in pupil size between neighboring explore choices = -0.17, t-test, t(27) = -2.69, p
< 0.02; 95% CI = -0.30 to -0.04). This was essentially identical to the magnitude with which the
pupil shrank on exploit trials that followed explore trials (mean change = -0.17, t-test, t(27) = -
2.56, p < 0.02; 95% CI = -0.31 to -0.03). Validating the ramping we observed with other
methods, we also found that pupil size tended to grow on explore trials that followed exploit
trials here (mean change = 0.14, t-test, t(27) = 2.96, p < 0.01; 95% CI = 0.04 to 0.24). Together,
these results suggest that pupil size and pupil-linked mechanisms specifically predict the onset
of exploration and may not be important for sustaining exploration after the first explore choice.
260
The fact that pupil size increased before exploration and shrank after the first explore trial in a
sequence could explain the U-shaped relationship between pupil size and exploration. Perhaps
the small-pupil-size explore choices are the later explore choices in a sequence and larger pupil
size only predicts the first explore choice, the "onset" of exploration. Indeed, pupil size had a
purely linear relationship with the onset of exploration (**Figure 2E**; 1st order GLM: beta = 0.26, p
< 0.0001, n = 28 sessions; 1st order, linear model AIC = 8443.1, 2nd order, quadratic model
AIC = 8444.7, AIC weight of quadratic model = 0.45; see **Methods**). Conversely, there was no
special relationship between pupil size and probability of starting to exploit (1st order GLM: beta
= 0.05, p > 0.05). Thus, pupil size specifically predicted the onset of exploration, rather than
explore choices or state switches more generally.
271
It remained possible that the relationship between pupil size and the onset of exploration was a
by-product of the effect of reward history on both variables. Exploration (Daw et al., 2006; Ebitz
et al., 2018; Wilson et al., 2014) and pupil size (Bijleveld et al., 2009; Jepma and Nieuwenhuis,
2011) both tend to increase when rewards are omitted. To determine if there was a direct effect
of pupil-related processes on exploration, we compared pupil size across exploit trials before
exploration with pupil size from matched trial sequences where exploration did not happen (see
Methods). There was a significant increase in pupil size during the trials before exploration
compared to "matched rewards" control trials (**Figure 3A**; GLM, beta = 0.025, p < 0.01, n = 28),
suggesting that pupil size predicted the onset of exploration beyond what could be explained by
reward. Again, pupil size ramped up over time (GLM, beta = 0.119, p < 0.02, n = 28), but this
ramping did not differ between the traces (GLM, beta = 0.007, p > 0.5, n = 28). This implies that
either reward history or time (i.e. the number of trials) may explain the pupil ramping before
exploration, although there is still an offset in pupil size that predicts the onset of exploration
above and beyond the effect of reward history.
286
Critically, the trials where pupil size best predicted the onset of exploration were not those
immediately before exploration (trial t-1, t-2, etc.). Instead, they were earlier trials (**Figure 3B**;

trial t-4, mean difference = 0.117, p < 0.05, t(27) = 2.09; trial t-5 = 0.170, p < 0.01, t(27) = 2.84).
Visual inspection suggested that this could be due to pupil size being out of phase in the trials
before exploration compared to matched-reward controls. To test this hypothesis, we asked if
the onset of exploration was phase-locked to the pupil size (see **Methods**). Median trial length
in this task was ~3 seconds (range = [2.2, 3.2]), meaning that the ~5 trial period we observed in
Figure 3B would correspond to an oscillation in the 0.06-0.09 Hz range. This is close to the
reported frequency of the Mayer wave (0.05-1 Hz; (Borjon et al., 2016; Julien, 2006)): a well-
known oscillation of the autonomic system that appears in heart rate, blood pressure, and
sympathetic efferents (Borjon et al., 2016; Japundzic et al., 1990; Julien, 2020, 2006; Kamiya et
al., 2005). We found that pupil phase at the onset of exploration was concentrated at the rising
phase (**Figure 3C**; mean phase = 47.18°, Hodges-Ajne test for non-uniformity, p < 0.01, vector
length = 0.075, same for bootstrapped null distribution = 0.026, 95% CI = 0.004 to 0.057, p <
0.0001). In contrast, the distribution of phases in reward-matched trials pointed in the opposite
direction (mean phase = 207.58°; significantly different from onsets: p < 0.02, Watsons U^2 =
0.25, n = 2170 phases including 1135 onsets). Thus, the onset of exploration was phase-locked
to a slow oscillation in pupil size across trials, suggesting a model in which reward information
interacts with oscillations in arousal to trigger exploration (**Figure 3D**).
306

Figure 3. The onset of exploration is phase-locked to pupil size. *A) Average pupil size over*
sequences of exploit trials before the onset of exploration (black line) and sequences with matched
rewards, but no exploration at the end (gray line). Lines: GLM fit. B) Difference in pupil size between the
traces in A. C) Phase distribution of pupil size at the onset of exploration (blue) and bootstrapped null
distribution (black). The vectors at the center indicate the mean vector direction and length for the trials
before exploration (blue) and the matched reward trials (gray). Shaded areas ± SEM throughout. D)
Cartoon illustrating how oscillations in arousal (top) could interact with reward history (middle), to regulate
exploration. The bottom panel illustrates a hypothetical pupil trace that has an additive effect of reward
omissions and by oscillating arousal. Exploration (diamond shapes) begins when pupil size reaches a
threshold (dotted line). Note that identical patterns of reward delivery and omission have different

outcomes, depending on how they align with the phase of arousal (gray = no exploration, blue =
exploration).
320
321
To understand the pupil-linked neural mechanisms that might contribute to the onset of
exploration, we next looked at the relationship between pupil size and certain neural signatures
of exploration in the frontal eye fields (FEF), a part of the prefrontal cortex implicated in directing
gaze and attention (Bruce and Goldberg, 1985; Moore and Armstrong, 2003; Moore and Fallah,
2001; Schall and Hanes, 1993) (**Figure 4A**). We previously reported that exploration is
associated with flattened neuronal turning curves in FEF (Ebitz et al., 2018). While FEF neurons
tend to predict the choice the subject will make during exploration, many neurons do not
differentiate between choices to different options during exploration. Here, we found that pupil
size also predicted flattened tuning in FEF neurons. A total of 88 out of 155 single neurons were
tuned for choice (**Figure 4B**; 57%, one sample proportion test: $p < 0.001$). Of those neurons, 21
were also modulated by pupil size (24%; $p < 0.05$), and 16 had a significant interaction between
choices and pupil size (18%; $p < 0.5$). On average, we found that neuronal tuning curves tended
to flatten as pupil size increased (**Figure 4C-D**).
335

Figure 4. Pupil size predicts choice tuning curves and population disorganization. A)
Recordings were made in the FEF. Right: The cartoon illustrates the relative positions of the receptive
field target (T_{in}, red) and the ipsilateral and contralateral targets (T_{out}, blue and green). B) Percent of
neurons with significant tuning for choice target, pupil size, and the interaction. C) Tuning curve for an
example neuron across target locations, separated by pupil size. Lighter = larger pupil. D) Same for all
tuned neurons. E) Cartoon illustrating how neural population measures consider patterns of firing rates
across neurons as vectors in neural state space. Targeted dimensionality reduction is used to find the
hyperplane where the distribution of neural activity across trials best predicts choice. Vectors here are the
coding dimensions that separate the choices. F) The decoded choice probability (projection onto the
correct coding dimension) plotted as a function of pupil size quantile. Inset: Same for exploit trials alone.
G) The scatter index, a measure of the variance in choice-predictive population activity, plotted as a
function of pupil size quantile. Inset: Same for exploit trials. H) Mediation analysis between pupil size,
scatter index, and the onset of exploration. Top: Direct model. Bottom: Indirect, mediated model.
Asterisks marked significant paths (*p < 0.01 **p < 0.001). I) Decoded choice probability for trials before
the onset of exploration (in blue) and trials with matched rewards (in gray). J) Scatter index for trials
before the onset of exploration (in blue) and trials with matched rewards (in gray). Error bars and shaded
regions ± SEM.

Neuronal tuning functions are too noisy to partial out the contributions of exploration and pupil
size, but changes in single-neuron tuning functions also imply changes in the organization of the
neural population (Ebitz and Hayden, 2021; **Figure 4E**). Indeed, we found that pupil size also
predicted changes in how accurately we could decode choice information from populations of
simultaneously-recorded FEF neurons. Consistent with the results of our previous study (Ebitz
et al., 2018), there was a decrease in decoded choice probability during explore choices
compared to exploit choices (paired t-test: both subjects, $p < 0.0001$, $t(27) = 8.65$; subject B, $p <$
0.005, $t(9) = -4.72$; subject O, $p < 0.0001$, $t(17) = -7.97$). Critically, larger pupil size predicted
less choice information both across all trials (**Figure 4F**; GLM: beta = -0.032, $p < 0.0001$), and
within exploit trials alone (GLM: beta = -0.037, $p < 0.005$). Therefore, changes in pupil size
predicted variations in the amount of choice-predictive information in FEF neurons, even within
periods of exploitation.

Our previous study reported that the decrease in choice-predictive activity in FEF was related to
an increase in what we called the "scatter index" of FEF populations (Ebitz et al., 2018): the
spread within clusters of same-choice population activity (see **Methods**). A high scatter index
indicates that neural activity on a given trial was dissimilar to other trials where the same choice
was made, whereas a low scatter index indicates that neural activity was tightly clustered. In the
present study, we observed that increasing pupil size predicted an increase in the scatter index
in both subjects (**Figure 4G**; GLM: beta = 0.04, $p < 0.0001$). Again, this was true and of a similar
magnitude even within exploit trials (beta = 0.03, $p < 0.0005$). Thus, pupil size predicted
disorganization of choice-predictive signals in the FEF, at both the level of single neurons and in
the population.

To determine if neural population measures also predicted transition into exploration, we looked
at how they evolved across trials before exploration, compared to matched-reward control trials.
While sudden changes in the decoded choice probability and scatter index were largely aligned
with the onset of exploration (as reported previously), these neural measures were at a different
average level on the trials preceding exploration, compared to reward-matched controls (**Figure
4H-I**; choice probability, offset = -0.611, $p < 0.001$, n = 28; scatter index = 0.258, $p < 0.001$).
Reward information did not cause a change in either variable (choice probability, slope = -0.004,
$p > 0.5$; scatter index = 0.002, $p > 0.5$), while small, but significant interaction terms suggested
that both variables anticipated the onset of exploration (choice probability interaction = -0.058, p
< 0.01; scatter index interaction: beta = 0.040, $p < 0.001$). To determine if these neural
measures might explain or mediate some of the relationship between pupil size and exploration,
we turned to structural equation modeling (Preacher et al., 2007; Sobel, 1986). We found that
the scatter index was a significant mediator of the relationship between pupil size and the onset
of exploration (**Figure 4J**; effect of mediation, ab = 0.003, $p < 0.005$; full report in Table S1).
Together, these results suggest that pupil size predicts disruptions in the organization of
prefrontal neural activity that mediate its relationship with the onset of exploration.
392

Figure 5. Pupil size predicts behavioral and neural slowing. *A) Response time on exploit trials before the onset of exploration (blue) and trials with matched rewards but no exploration (gray). B) Response time plotted as a function of pupil size quantile. Inset: Same for exploit trials alone. C) Mediation analysis between pupil size, response time, and the onset of exploration. Top: Direct model. Asterisks marked significant paths (*p < 0.01). Bottom: Indirect, mediated model. D) Neural speed on exploit trials before the onset of exploration (in blue) and trials with matched rewards (in gray). E-F) Same as B-C for neural speed. Shaded areas and error bars ± SEM.*

Complex systems like neural networks can experience tipping points: irreversible "critical transitions" that mark the passage from one operating regime into another (O'Byrne and Jerbi, 2022; Scheffer, 2020; Scheffer et al., 2009; Wang et al., 2012). Because exploration occurs as the brain passes from exploiting one target to exploiting another, it is worth considering the possibility that exploration may represent a critical transition in brain states. Indeed, during exploration, we previously reported (Ebitz et al., 2018) several phenomena in the FEF and in behavior that are hallmarks of critical transitions, including a rapid flickering back and forth between choices (Wang et al., 2012), an increase in the variance in neural activity (Scheffer et al., 2009), and a disruption of long-term neuronal autocorrelations that suggests that passing through exploration causes time-irreversible changes in the FEF network (Scheffer, 2020). However, there is another classic feature of critical transitions that we did not consider: an early warning signal known as "critical slowing". As the system nears the tipping point, the dynamics within the system begin to flatten out in preparation for the change. As a result, the systems' processes slow down and take longer to trace the same paths (Scheffer et al., 2009). Therefore,

415 we next asked if there was any evidence that decision-making slowed down in advance
416 exploration in this dataset.
417
We considered two measures of the speed of decision-making: one behavioral and one neural.
First, we looked at response time, a measure of how long it takes the brain to generate saccadic
decisions. Response time was not only slower in the trials before exploration, compared to
matched-reward control trials (Figure 5A-C; GLM offset = 0.39, $p < 0.0001$, n = 28), but it
slowed down over trials before the onset of exploration (interaction = 0.05, $p < 0.001$). Second,
we looked at the mean rate of change in neural population choice signals during the decision
process ("neural speed", see **Methods**). Neural speed was only weakly correlated with
response time across sessions (mean = -0.07, min = -0.36, max = 0.09, Pearson's correlation),
suggesting that the measures were complementary, rather than redundant. Like response time,
neural speed was also significantly slowed in the trials before exploration, compared to
matched-reward controls (**Figure 5D-F**; GLM offset = -0.17, $p < 0.0001$, n = 28; interaction = -
0.01, $p = 0.08$). Although the notion that the brain may be subject to critical tipping points is
controversial (O'Byrne and Jerbi, 2022), these results are consistent with the idea that
exploration could reflect a critical transition between exploiting one option and exploiting
another.
433
In the last analysis, we found that typical explore-triggered reward histories did not, by
themselves, have a significant effect on neural or behavioral slowing (response time: slope of
matched-reward trials = 0.0002, $p > 0.5$; neural speed: slope = -0.018, $p > 0.1$). This suggests
that some internal variable, like arousal, could be driving slowing and, perhaps, also the
systems' proximity to a tipping point. Indeed, increasing pupil size predicted slower response
times (**Figure 5B**; GLM beta = 0.08, $p < 0.0001$, n = 28 sessions), even within periods of
exploitation (beta = 0.05, $p < 0.0001$). The same was true of neural slowing (**Figure 5E**; all
trials: beta = -0.03, $p < 0.0005$; exploit only: beta = -0.09, $p < 0.0001$). Further, structural
equation modeling revealed that both measures of slowing mediated the relationship between
pupil size and the onset of exploration (**Figure 5C and F**; Table S2-3). In sum, the pupil-linked
mechanisms that anticipated the onset of exploration included an increase in the disorganization
and trial-to-trial variability of neural activity, and a slowing of decision-making computations in
brain and behavior.

Discussion

Random decision-making is a powerful strategy for exploration (Dayan and Daw, 2008; Ebitz et
al., 2018; Gershman, 2019; Wilson et al., 2021, 2014) that is linked to disorganized patterns of
neural activity in the prefrontal cortex (Ebitz et al., 2018; Muller et al., 2019; Wilson et al., 2021).
Here, we sought to identify some of the neurobiological mechanisms that drive random
exploration and its neural signatures. We found that pupil size, a peripheral correlate of
autonomic arousal, predicted exploration and certain measures of neural population activity
previously linked to exploration. Consistent with previous studies (Jepma and Nieuwenhuis,
2011), pupil size was generally larger during exploration, compared to exploitation. However,
there was also a complex temporal relationship, where pupil size ramped up between periods of

exploration and decreased during exploration. As a result, pupil size was largest at the
beginning or "onset" of exploration and explained variance in the onset of exploration that could
not be explained by other variables. Together, these results suggest that pupil-linked
mechanisms may play a role in driving the brain into an exploratory state.
461
Our behavioral results largely replicate previous findings linking exploration to increased pupil
size (Jepma and Nieuwenhuis, 2011). However, where we found gradual ramping before
exploration and sudden constriction after, Jepma and Nieuwenhuis (2011) reported an abrupt (if
modest) increase of pupil size at the onset of exploration and then a gradual decrease at the
return to exploitation. The discrepancy may be due to differences in the operational definition of
exploration. Jepma and Nieuwenhuis (2011) fit an RL model to behavior and defined "explore
choices" as the choices that were not reward-maximizing according to the model. This definition
conflates exploration with errors of reward maximization. A strategy that is non-reward-
maximizing would produce choices that are orthogonal to value, not consistently bad. Here, we
used an HMM to identify latent explore and explore states on the basis of the temporal profiles
of choices alone, with no assumptions on the underlying value computations. This allowed us to
dissociate the effects of reward history from the explore/exploit choice labels. We reported here
(**Figure 1C**), and in previous studies (Chen et al., 2021; Ebitz et al., 2018), that HMM labels
outperform RL labels in explaining behavioral and neural measures, suggesting that the HMM
may more accurately separate distinct neural and behavioral states. If the HMM allows for more
precise identification of exploratory and exploitative choices, it would follow that it also allows for
more precise reconstruction of the temporal relationship between the pupil and exploration.
479
The precision of our explore/exploit labels revealed that the U-shaped relationship between
pupil size and exploration was caused by a refractory constriction in the pupil. When exploration
was plotted as a function of pupil size, the relationship appeared non-linear: both small- and
large-pupil choices were more likely to be exploratory. This superficially resonated with the idea
of a U-shaped relationship between arousal and task performance (i.e. the "Yerkes-Dodson
curve"; (Aston-Jones and Cohen, 2005; Yerkes and Dodson, 1908)): perhaps reliable
exploitation is only possible at intermediate levels of arousal. However, when we examined the
temporal relationship between exploration and pupil size, we found that pupil size only predicted
the onset of exploration, the first explore choice in a sequence. Small-pupil explore choices
happened because starting to explore seemed to "reset" the level of pupil-linked arousal,
causing it to quickly fall below baseline. If increased pupil size promotes a transition to
exploration, then it is possible that post-exploration constriction represents a refractory period
for exploration. Given that uncertainty grows with time in this task (and in all dynamic
environments), it may not be smart to start to explore again immediately after you have just
explored. A refractory period could ensure that non-reward-maximizing explore choices are
deployed only when needed. Future work is needed to test this hypothesis and to determine the
cognitive and/or neurobiological mechanisms at play.
497
Before exploration, we observed an oscillatory dynamic that was about twice as fast as the 10
trials it took the pupil to recover after exploration. This 5 trial, 0.06-0.09 Hz oscillation entrained
the onset of exploration: onsets tended to occur during the rising phase of pupil size, whereas

identical trial sequences that did not result in exploration were on the opposite phase. This
implies that it is the confluence of pupil size, pupil phase, and trial history that best predicts the
onset of exploration. This result reinforces the idea that arousal or arousal-linked mechanisms
help trigger random exploration (Ebitz and Moore, 2019; Gilzenrat et al., 2010; Reimer et al.,
2016), rather than just tracking the reward-linked variables that make exploration more
probable. It is also notable that the period of the pupil oscillation was close to the frequency of
the Mayer wave: an oscillation in blood pressure that entrains other autonomic measures,
including respiration and heart rate (Borjon et al., 2016; Japundzic et al., 1990; Julien, 2020,
2006; Kamiya et al., 2005). There is precedent for the idea that behavior can be entrained by
the Mayer wave: in marmosets, fluctuations in arousal predict the spontaneous onset of a call
(Borjon et al., 2016). This paper argued that the Mayer wave may function to organize vocal
communication by bringing the system closer to the threshold for transitioning from inaction to
action. It is possible that oscillations in the pupil and pupil-linked mechanisms function the same
way here, organizing important state changes in time. In parallel, pupil-linked mechanisms seem
to anticipate other state transitions, including belief updating (Filipowicz et al., 2020; O'Reilly et
al., 2013), task disengagement (Kane et al., 2017), and other behavioral state changes (Bouret
and Sara, 2005). Together, these results suggest an important role for pupil-linked mechanisms
in driving successful transitions between certain neural and behavioral states.
519
Critically, pupil size and pupil oscillations did not predict all state transitions here, but only the
transition into exploration. What kinds of state transitions might be entrained by pupil- linked
arousal? It is possible that the pupil may have a special relationship with certain "critical" kinds
of transitions. Critical transitions are abrupt, large-scale, and irreversible changes in the
dynamics and behavior of complex systems, like the brain. As these systems go from being in
one conformation (i.e. always choosing the left option) into another conformation (i.e. always
choosing the right), the system dynamics that support the old state have to disappear and the
new dynamics have to emerge. During this brief transitory period, when both dynamics co-exist
in the system, certain signatures can be observed in the system. We previously reported that
the exploration was accompanied by abrupt changes in neural population activity, certain
patterns of noise in brain and behavior, and disruptions in long-term neuronal autocorrelations:
all observations that could be interpreted as suggesting that exploration is a critical transition in
the brain (Ebitz et al., 2018). Here, we found that pupil size predicts these features of neural
activity and also an prominent "early warning sign" of critical transitions: a slowing, in brain and
behavior, of the decision process. While there are certain patterns of activity in FEF that predict
response speed (Hauser et al., 2018; Yao and Vanduffel, 2023), here we identified *independent*
neural and behavioral measures of decision speed that both mediated the relationship between
pupil size and exploration. Together these results suggest that pupil-linked arousal pushes
neural and behavioral states to a critical tipping point and highlights the crucial role of pupil-
linked mechanisms in changing the dynamics of the brain.
540
What underlying, pupil-linked mechanisms could support critical transitions? Changes in pupil
diameter coincide with neuromodulator system activity, especially norepinephrine (NE) and
acetylcholine (Breton-Provencher and Sur, 2019; de Gee et al., 2020; Joshi et al., 2016; Joshi
and Gold, 2020; Murphy et al., 2014; Reimer et al., 2016). At the neuronal level, central NE

flattens tuning curves, at least in the auditory cortex (Martins and Froemke, 2015), though it may
have different effects in non-cortical structures (Manella et al., 2017). Here, we made a parallel
observation: as pupil size increases, neuronal turning curves flattened and choice-predictive
neural population activity became disorganized. These results resonate with a particularly
influential theory of NE function: the idea that NE release may facilitate "resets" in cortical
networks in order to effect long-lasting changes in brain and behavior (Aston-Jones and Cohen,
2005; Bouret and Sara, 2005). More recent studies seem to consistently report that elevated
levels of NE predict an increase in behavioral variability, while pharmacological blockade of NE
receptors reduces variability (Chen et al., 2023; Kane et al., 2017; Sadacca et al., 2017; Tervo
et al., 2014). In combination with the present study, these results could suggest that phasic NE
signaling functions to push the brain towards a critical tipping point where it is better able to
transition from one regime to another. In this view, behavioral variability would be linked to NE
not because NE increases variability directly, but because the brain is more likely to transition
into a high variability regime after it is released. Of course, pupil size is also associated with
other neuromodulatory systems, cognitive factors, and other measures of arousal. Thus, future
work is needed to identify the neurobiological mechanisms that underpin the relationship
between pupil size and critical transitions that we report here.
562

Acknowledgments

564 This work was supported by the Natural Sciences and Engineering Research Council of Canada
565 (Discovery Grant RGPIN-2020-05577), the Fonds de Recherche du Québec–Santé (Junior 1
566 Chercheur-Boursier 284309 to R.B.E.), the Jacobs Foundation (Research Fellowship, seed
567 grant to R.B.E.), and CIFAR Azrieli Global Scholars (seed grant to R.B.E.), the National Eye
568 Institute (R01-EY014924 to T.M.), and l'Institut de valorisation des données (fellowship to K.J.).

Author Contributions

570 R.B.E. collected the data. K.J., T.M., and R.B.E., secured funding. R.B.E., G.M., and A.S.
571 formulated the hypotheses. A.S., G.M., K.J., and R.B.E. analyzed the data. A.S., G.M., and
572 R.B.E. drafted the manuscript. A.S., G.M., K.J., T.M., and R.B.E. reviewed and edited the
573 manuscript.

Declaration of Interests

575 The authors declare no competing interests.
576

Resource Availability

Lead Contact

Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Becket Ebitz (becket@ebitzlab.com).

Materials Availability

This study did not involve generating new material.

Data and Code Availability

Data and software are available upon request to the Lead Contact (Becket Ebitz, becket@ebitzlab.com)

Methods

Surgical and electrophysiological procedures. All procedures were approved by the Stanford University Institutional Animal Care and Use Committee. Subjects were two male rhesus macaques, surgically-prepared with head restraint prostheses, craniotomies, and recording chambers under isoflurane anesthesia via techniques described previously (Ebitz et al., 2018). Following surgery, analgesics were used to minimize discomfort, and antibiotics were delivered prophylactically. After recovery, subjects were acclimated to the laboratory and head restraint, then placed on controlled access to fluids and trained to perform the task.

Recording sites were located within the FEF, which was identified via a combination of anatomical and functional criteria. The location of recording sites in the anterior bank of the arcuate sulcus was verified histologically in one subject and via microstimulation in both subjects (Ebitz et al., 2018). Recordings were conducted with 16-channel U-probes (Plexon), located such that each contact was within gray matter at an FEF site. An average of 20 units were recorded in each session (131 single units, 443 multi units; 576 total units across 28 sessions).

General behavioral procedures. Eye position and pupil size were monitored at 1000 Hz via an infrared eye tracking system (SR Research; Eyelink). The manufacturer's standard methods for calculating pupil area were used. MATLAB (Psychtoolbox-3; (Kleiner et al., 2007)) was used to display stimuli and record behavioral responses and pupil size measurements. Task stimuli were presented against a dark gray background (7 cd/m2) on a 47.5 cm wide LCD monitor (Samsung; 120 Hz refresh rate, 1680 x 1050 resolution), located 34 cm in front of the subject.

Three-armed bandit task. The subjects performed a sequential decision-making task in which they chose between 3 targets whose values changed over time. The subject first fixated a central fixation square (0.5° stimulus, +/- 1.5-2° of error) for a variable interval (450-750ms). At any point within 2s after the onset of the targets, subjects indicated their choice by making a

saccade to one of the targets and fixating it (+/- 3°) for 150 ms. Reward magnitude was fixed
within session (0.2-0.4 μL). Reward probability was determined by the current reward probability
of the chosen target, which changed independently over trials for each of the three targets. On
every correct trial, each target had a 10% chance of the reward parameter changing either up or
down by a fixed step of 0.1, bounded at 0.1 and 0.9. Because rewards were variable,
independent, and probabilistic, the subjects could only infer the values of the different targets by
sampling them and integrating noisy experienced rewards over multiple trials.

General analysis procedures. Data were analyzed with custom software in MATLAB. Unless
otherwise noted, all t-tests were paired, two-sided t-tests, and generalized linear models were
run on raw data, with session number coded as a dummy variable to account for session-to-
session variability. Model comparison was based on standard methods that involve calculating
the likelihood of the data and Akaike information criteria (AIC) of each model, then using AIC
weights to identify (1) the model that is most likely to minimize information loss, and (2) the
relative likelihood of competing models to do the same (Burnham and Anderson, 2004). For
analyses of any behavioral or neural variables on the trials before or after exploration,
continuous runs of exploit trials were required. The values of behavioral and neural variables
were *z*-scored within a session to facilitate comparisons across sessions.

Pupil size. Pupil size was measured during the first 200 ms of fixation, a time at which the eye
was fixed at a known point on the screen, illumination was identical across trials, and
anticipatory changes in the pupil were minimal. To remove any blinks or movement artifacts,
trials where pupil size or the change in pupil size from the first time bin of this epoch to the last
was +/- 6 standard deviations from average were eliminated from further analyses. A total of
178 trials (out of 21,793, approximately 0.8% of observations) were outliers.

Reinforcement learning model. To compare goal state labels derived from an RL and HMM
model, we employed a Rescorla-Wagner model (Rescorla and Wagner, 1972). This was fit
using maximum likelihood estimation. The value of each option is iteratively updated according
to:

$$V_{i,t+1} = V_{i,t} + \alpha(r_t - V_{i,t})$$

639   Where $V_{i,t}$ is the value of option *i* at time *t*, r_t is the reward at time *t*, and *α* represents the fitted
640   learning rate, which determines how much the difference between the predicted and actual
641   reward (the prediction error) influences value. To make a decision, the values are passed
642   through a softmax decision rule:

$$p_{i,t} = \frac{\exp(\beta V_{i,t})}{\Sigma_{j=1:n} \exp(\beta V_{j,t})}$$

Where n is the total number of available options, and β is the inverse temperature, which
controls the level of random noise in decision-making. After (Daw et al., 2006; Jepma and
Nieuwenhuis, 2011; Pearson et al., 2009), decisions that were not reward maximizing were
labeled as exploratory (i.e. any decision where $V_{chosen,t}$ was not the maximum V at time t).

Learning Index. To investigate whether learning differed with pupil size within the exploratory
choices, we calculated a learning index that captured the effect of rewards experienced during
exploration on future choices. Because reward effects decay exponentially quickly (Lau and
Glimcher, 2008), a 1-trial-ahead index should capture most of the variability in how much is
learned between trial types. The equation was:

$$\text{learning index}_t = \frac{p(\text{switch}_{t+1} \mid \text{reward}_t) - p(\text{switch}_{t+1} \mid \text{no reward}_t)}{p(\text{switch}_{t+1})}$$

Lagged change in pupil size. To determine whether exploration impacted pupil size, we
measured the change in pupil size (Δ pupil) between pairs of trials that either were or were not
separated by at least 1 explore trial. Segments of twenty-five consecutive trials were identified
that either included a single bout of exploration or did not include exploration. For each pair of
trials within these sequences, we then measured the change in pupil size between the first
exploit trial of the sequence (t1) and the remaining exploit trials in the sequence (t2:25). This
was repeated for all unique pairs of trials that met our selection criteria.
661
Matched reward trials. To test whether the rising trend in pupil size before exploration is best
explained by reward history, we identified trial sequences with identical reward and state
histories that did not end in exploration ("matched rewards"). For each onset of exploration
preceded by at least 6 exploit trials, we searched for identical sequences of exploit trials, with
identical reward histories, that did not end in exploration. We chose 6 previous trials because
this was the longest sequence of reward history we could regularly match within the majority of
sessions (we could find at least 10 matched sequences in 96% [27/28] of sessions for 6 trials
sequences; that dropped to 75% [21/28] at 7 trials). Identical results were obtained with other
sequence lengths, though these analyses included fewer sessions.
671
672   *Mediation analysis.* To determine if the predictive relationship between pupil size and
673   exploration was mediated by other variables, we used structural equation modeling to test for
674   mediation. Mediation analyses involve fitting three regression models. The first model measures

675 the total effect (c) of the independent variable (here, pupil size) on the independent variable
676 (here, onset of exploration):

$$\text{Explore}_t = \gamma_1 + c(\text{Pupil}_{t-1}) + \epsilon_1$$

677
678 In these equations, γ represents the intercept for each equation, while ϵ represents the error of
679 the model. Note that the estimated parameter c will include both direct effects of pupil size on
680 exploration, but also indirect effects that may be mediated by other variables. Therefore, we
681 also fit a second model that tests if the independent variable also predicts a potential mediator
682 variable (here, neural network scatter):

$$\text{Scatter}_{t-1} = \gamma_2 + a(\text{Pupil}_{t-1}) + \epsilon_2$$

683
684 Model parameter a thus captures the effect of pupil size on the mediator. Finally, a third model
685 estimates the unique contributions of both the potential mediator (scatter, b) and the
686 independent variable (pupil size, c'), now controlling for the mediator:

$$\text{Explore}_t = \gamma_3 + b(\text{Scatter}_{t-1}) + c'(\text{Pupil}_{t-1}) + \epsilon_3$$

687
688 A drop between c and c' indicates that the effect of the independent variable (pupil) on the
689 dependent variable (exploration) is reduced when the mediating variable is considered. The
690 mediation effect (the indirect effect of the pupil size on the onset of exploration via the mediating
691 factor) can also be estimated directly, via taking the product of the coefficients a and b. Sobel's
692 test is used to determine the significance of the mediation path (Sobel, 1986).
693
Phase analysis. To determine if the onset of exploration happened at a specific phase of pupil
size over trials, we performed a wavelet analysis. Because this method only assumes local
stationarity, it is more suitable than other methods for analyzing pupil size, which tended to
ramp over trials. A wavelet was constructed by multiplying a complex sine wave (frequency = 5
trials) with a Gaussian envelope (μ = 0, σ = cycles / (2π*frequency), cycles = 5; (Cohen, 2014)).
The wavelet was convolved with the baseline pupil size time series and the phase of the signal
was calculated on each trial (Matlab; angle). Standard circular statistics were used to measure
the differences between phase distributions for explore onsets and reward-matched controls
(Zar, 1999) and the phase alignment within these trial types (Berens, 2009). The latter was also
verified via comparison with bootstrapped null distributions (1000 samples).
Targeted dimensionality reduction. Neural state spaces have as many dimensions as there are
recorded neurons, but converging evidence suggests (1) that the neural states that are
observed in practice are generally confined to a lower-dimensional "manifold", and (2) that task-
relevant information is encoded by a small number of dimensions in that manifold. Because we
wanted to isolate the effects of arousal on choice-related activity from well-known effects of

arousal on neural activity (Ebitz and Platt, 2015; McGinley et al., 2015; Pfeffer et al., 2022;
Podvalny et al., 2021; Reimer et al., 2016, 2014; van Kempen et al., 2019; Waschke et al.,
2019), we focused all our neural population analyses on activity within the choice-predictive
subspace, rather than on neural activity more broadly.
713
To do this, we used targeted dimensionality reduction to identify the choice-predictive
dimensions of the neural state space (Cohen and Maunsell, 2010; Cunningham and Yu, 2014;
Ebitz et al., 2018; Peixoto et al., 2021). Specifically, we used multinomial logistic regression
(Matlab; mnrfit, mnrval, (Hastie et al., 2009)) to identify the separating hyperplanes that best
discriminated each choice from the alternative choices. This is equivalent to fitting a system of
binary classifiers of the form:

$$p(\text{choice} = \text{k} \mid \text{X}) = \frac{1}{1 - \exp^{-\beta_k \text{X}}}$$

720
Where one classifier discriminates target 1 choices from targets 2 and 3 and a second
discriminates target 2 choices from targets 1 and 3. The classifier that discriminates target 3
from targets 1 and 2 is then just the negative of target 1 and target 2. These axes span the
subspace in which neural activity best predicts choice. Classifiers were trained on firing rates
from an epoch that began when the targets appeared and ended at the time of the saccade.
Mean imputation was used for the small number of occasions where a unit was not held for the
whole duration of the session (~3% of trials, ~12% of units) and a small fraction of units were
omitted from these analyses because their mean firing rates were less than 2 spikes/s, which
makes their weights difficult to identify (~8% of units).
730
Choice Probability Decoding. Within the choice-predictive subspace, the distance from the
separating hyperplanes (the vectors illustrated in **Figure 4E**) are the decoding vectors: the
vectors along which we can project neural activity in order to decode the log odds of choice.
This projection is equivalent to the decoded choice probability from the multinomial logistic
regression model and this is the figure we took as the decoded choice probability in **Figures 3F**
and **3H**. We evaluated decoding accuracy by measuring how often the most-likely choice
predicted by the model coincided with the choice the subject made.
738
739   *Scatter index.* The scatter index measures how much choice-predictive population neural
739   activity is clustered between trials with the same choice (Ebitz et al., 2018). It is calculated by
740   measuring the average Euclidean distance of each trial from all other trials where the same
741   choice was made and dividing it by the average Euclidean distance to all other trials where a
742   different choice was made:

$$\text{scatter} = \frac{\text{d}_{\text{within}}}{\text{d}_{\text{between}}}$$

Each trial thus has its own scatter index value, with a value of 1 indicating no difference in
clustering between same-choice and different-choice trials, and a value less than 1 indicating
greater clustering with same-choice trials compared to different-choice trials.

Neural speed. To determine how the speed of the decision-making process changed before and
during exploration, we calculated the rate of change in neural states within the choice-predictive
subspace during the first 400 ms following target presentation. Each trial's neural activity was
sampled in non-overlapping 20 ms bins and then projected into the choice-predictive subspace.
The change in neural activity within the subspace was then calculated between each pair of
samples. Finally, the changes were averaged together across the trial and normalized to the bin
width to produce an average rate of change in choice-predictive activity for that trial.

Table S1

Regression coefficients and p values for the mediation analysis testing whether the scatter
index mediates the relationship between pupil size and the onset of exploration. Related to
Figure 4J.

		pupil$_{t-1}$ → scatter$_{t-1}$ → explore$_t$	
		est. coefficient	p value <
Total effect	c	0.090	0.005
Effect on mediator	a	0.036	0.0005
Unique mediator effect	b	0.092	0.005
Indirect effect	ab	0.003 (z = 2.70*)	0.005
Direct effect	c'	0.086	0.005

758 *Sobel's test

759 **Table S2**

760 Regression coefficients and p values for the mediation analysis testing whether response time
761 slowing mediates the relationship between pupil size and the onset of exploration on the next
762 trial. Related to **Figure 5C**.

		pupil$_{t-1}$ → RT slowing$_{t-1}$ → explore$_t$	
		est. coefficient	p value <
Total effect	c	0.090	0.005
Effect on mediator	a	0.078	0.0001
Unique mediator effect	b	0.106	0.0005
Indirect effect	ab	0.008 (z = 3.48*)	0.0005
Direct effect	c'	0.080	0.01

763 *Sobel's test

764 **Table S3**

765 Regression coefficients and p values for the mediation analysis testing whether neural slowing
766 mediates the relationship between pupil size and the onset of exploration on the next trial.
767 Related to **Figure 5F**.

		pupil$_{t-1}$ → neural slowing$_{t-1}$ → explore$_t$	
		est. coefficient	p value <
Total effect	c	0.095	0.005
Effect on mediator	a	-0.025	0.0005
Unique mediator effect	b	-0.059	0.06
Indirect effect	ab	0.001 (z = 1.69*)	0.05
Direct effect	c'	0.093	0.005

*Sobel's test

References

Aston-Jones G, Cohen JD. 2005. AN INTEGRATIVE THEORY OF LOCUS COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal Performance. *Annu Rev Neurosci* **28**:403–450. doi:10.1146/annurev.neuro.28.061604.135709

Berens P. 2009. CircStat: A MATLAB Toolbox for Circular Statistics. *J Stat Softw* **31**:1–21. doi:10.18637/jss.v031.i10

Bijleveld E, Custers R, Aarts H. 2009. The unconscious eye opener: pupil dilation reveals strategic recruitment of resources upon presentation of subliminal reward cues. *Psychol Sci* **20**:1313–1315. doi:10.1111/j.1467-9280.2009.02443.x

Borjon JI, Takahashi DY, Cervantes DC, Ghazanfar AA. 2016. Arousal dynamics drive vocal production in marmoset monkeys. *J Neurophysiol* **116**:753–764. doi:10.1152/jn.00136.2016

Bouret S, Sara SJ. 2005. Network reset: a simplified overarching theory of locus coeruleus noradrenaline function. *Trends Neurosci* **28**:574–582. doi:10.1016/j.tins.2005.09.002

Bradley MM, Miccoli L, Escrig MA, Lang PJ. 2008. The pupil as a measure of emotional arousal and autonomic activation. *Psychophysiology* **45**:602–607. doi:10.1111/j.1469-8986.2008.00654.x

Breton-Provencher V, Sur M. 2019. Active control of arousal by a locus coeruleus GABAergic circuit. *Nat Neurosci* **22**:218–228. doi:10.1038/s41593-018-0305-z

Bruce CJ, Goldberg ME. 1985. Primate frontal eye fields. I. Single neurons discharging before saccades. *J Neurophysiol* **53**:603–635. doi:10.1152/jn.1985.53.3.603

Burnham KP, Anderson DR, editors. 2004. Model Selection and Multimodel Inference. New York, NY: Springer. doi:10.1007/b97636

Chen CS, Knep E, Han A, Ebitz RB, Grissom NM. 2021. Sex differences in learning from exploration. *eLife* **10**:e69748. doi:10.7554/eLife.69748

Chen CS, Mueller D, Knep E, Ebitz RB, Grissom NM. 2023. Dopamine and norepinephrine differentially mediate the exploration-exploitation tradeoff. doi:10.1101/2023.01.09.523322

Clewett D, Gasser C, Davachi L. 2020. Pupil-linked arousal signals track the temporal organization of events in memory. *Nat Commun* **11**:4007. doi:10.1038/s41467-020-17851-9

Cohen MR, Maunsell JHR. 2010. A Neuronal Population Measure of Attention Predicts Behavioral Performance on Individual Trials. *J Neurosci* **30**:15241–15253. doi:10.1523/JNEUROSCI.2171-10.2010

Cohen MX. 2014. Analyzing neural time series data: theory and practice. MIT press.

Costa VD, Averbeck BB. 2020. Primate Orbitofrontal Cortex Codes Information Relevant for Managing Explore–Exploit Tradeoffs. *J Neurosci* **40**:2553–2561. doi:10.1523/JNEUROSCI.2355-19.2020

Costa VD, Rudebeck PH. 2016. More than Meets the Eye: the Relationship between Pupil Size and Locus Coeruleus Activity. *Neuron* **89**:8–10. doi:10.1016/j.neuron.2015.12.031

Cunningham JP, Yu BM. 2014. Dimensionality reduction for large-scale neural recordings. *Nat Neurosci* **17**:1500–1509. doi:10.1038/nn.3776

Daw ND, O'Doherty JP, Dayan P, Seymour B, Dolan RJ. 2006. Cortical substrates for exploratory decisions in humans. *Nature* **441**:876–879. doi:10.1038/nature04766

Dayan P, Daw ND. 2008. Decision theory, reinforcement learning, and the brain. *Cogn Affect Behav Neurosci* **8**:429–453. doi:10.3758/CABN.8.4.429

de Gee JW, Tsetsos K, Schwabe L, Urai AE, McCormick D, McGinley MJ, Donner TH. 2020. Pupil-linked phasic arousal predicts a reduction of choice bias across species and decision domains. *eLife* **9**:e54014. doi:10.7554/eLife.54014

Ding L, Hikosaka O. 2006. Comparison of Reward Modulation in the Frontal Eye Field and Caudate of the Macaque. *J Neurosci* **26**:6695–6703. doi:10.1523/JNEUROSCI.0836-06.2006

Ebitz RB, Albarran E, Moore T. 2018. Exploration Disrupts Choice-predictive Signals and Alters Dynamics in Prefrontal Cortex. *Neuron* **97**:450-461.e9. doi:10.1016/j.neuron.2017.12.007

Ebitz RB, Hayden BY. 2021. The population doctrine in cognitive neuroscience. *Neuron* **109**:3055–3068. doi:10.1016/j.neuron.2021.07.011

Ebitz RB, Moore T. 2019. Both a Gauge and a Filter: Cognitive Modulations of Pupil Size. *Front Neurol* **9**.

Ebitz RB, Pearson JM, Platt ML. 2014. Pupil size and social vigilance in rhesus macaques. *Front Neurosci* **8**.

Ebitz RB, Platt ML. 2015. Neuronal Activity in Primate Dorsal Anterior Cingulate Cortex Signals Task Conflict and Predicts Adjustments in Pupil-Linked Arousal. *Neuron* **85**:628–640. doi:10.1016/j.neuron.2014.12.053

Ebitz RB, Sleezer BJ, Jedema HP, Bradberry CW, Hayden BY. 2019. Tonic exploration governs both flexibility and lapses. *PLOS Comput Biol* **15**:e1007475. doi:10.1371/journal.pcbi.1007475

Ebitz RB, Tu JC, Hayden BY. 2020. Rules warp feature encoding in decision-making circuits. *PLOS Biol* **18**:e3000951. doi:10.1371/journal.pbio.3000951

Eldar E, Cohen JD, Niv Y. 2013. The effects of neural gain on attention and learning. *Nat Neurosci* **16**:1146–1153. doi:10.1038/nn.3428

Fiete IR, Fee MS, Seung HS. 2007. Model of Birdsong Learning Based on Gradient Estimation by Dynamic Perturbation of Neural Conductances. *J Neurophysiol* **98**:2038–2057. doi:10.1152/jn.01311.2006

Filipowicz AL, Glaze CM, Kable JW, Gold JI. 2020. Pupil diameter encodes the idiosyncratic, cognitive complexity of belief updating. *eLife* **9**:e57872. doi:10.7554/eLife.57872

Gershman SJ. 2019. Uncertainty and Exploration. *Decis Wash DC* **6**:277–286. doi:10.1037/dec0000101

Gilzenrat MS, Nieuwenhuis S, Jepma M, Cohen JD. 2010. Pupil diameter tracks changes in control state predicted by the adaptive gain theory of locus coeruleus function. *Cogn Affect Behav Neurosci* **10**:252–269. doi:10.3758/CABN.10.2.252

Graves JE, Egré P, Pressnitzer D, de Gardelle V. 2021. An implicit representation of stimulus

ambiguity in pupil size. *Proc Natl Acad Sci* **118**:e2107997118.
doi:10.1073/pnas.2107997118

Hastie T, Tibshirani R, Friedman J. 2009. The Elements of Statistical Learning, Springer Series
in Statistics. New York, NY: Springer. doi:10.1007/978-0-387-84858-7

Hauser CK, Zhu D, Stanford TR, Salinas E. 2018. Motor selection dynamics in FEF explain the
reaction time variance of saccades to single targets. *eLife* **7**:e33456.
doi:10.7554/eLife.33456

Japundzic N, Grichois M-L, Zitoun P, Laude D, Elghozi J-L. 1990. Spectral analysis of blood
pressure and heart rate in conscious rats: effects of autonomic blockers. *J Auton Nerv*
Syst **30**:91–100. doi:10.1016/0165-1838(90)90132-3

Jepma M, Nieuwenhuis S. 2011. Pupil Diameter Predicts Changes in the Exploration–
Exploitation Trade-off: Evidence for the Adaptive Gain Theory. *J Cogn Neurosci*
23:1587–1596. doi:10.1162/jocn.2010.21548

Joshi S, Gold JI. 2020. Pupil Size as a Window on Neural Substrates of Cognition. *Trends Cogn*
Sci **24**:466–480. doi:10.1016/j.tics.2020.03.005

Joshi S, Li Y, Kalwani R, Gold JI. 2016. Relationships between pupil diameter and neuronal
activity in the locus coeruleus, colliculi, and cingulate cortex. *Neuron* **89**:221–234.
doi:10.1016/j.neuron.2015.11.028

Julien C. 2020. An update on the enigma of Mayer waves. *Cardiovasc Res* **116**:e210–e211.
doi:10.1093/cvr/cvz327

Julien C. 2006. The enigma of Mayer waves: Facts and models. *Cardiovasc Res* **70**:12–21.
doi:10.1016/j.cardiores.2005.11.008

Jurewicz K, Sleezer BJ, Mehta PS, Hayden BY, Ebitz RB. 2022. Irrational choices via a
curvilinear representational geometry for value. doi:10.1101/2022.03.31.486635

Kamiya A, Hayano J, Kawada T, Michikami D, Yamamoto K, Ariumi H, Shimizu S, Uemura K,
Miyamoto T, Aiba T, Sunagawa K, Sugimachi M. 2005. Low-frequency oscillation of
sympathetic nerve activity decreases during development of tilt-induced syncope
preceding sympathetic withdrawal and bradycardia. *Am J Physiol-Heart Circ Physiol*
289:H1758–H1769. doi:10.1152/ajpheart.01027.2004

Kane GA, Vazey EM, Wilson RC, Shenhav A, Daw ND, Aston-Jones G, Cohen JD. 2017.
Increased locus coeruleus tonic activity causes disengagement from a patch-foraging
task. *Cogn Affect Behav Neurosci* **17**:1073–1083. doi:10.3758/s13415-017-0531-y

Kaske EA, Chen CS, Meyer C, Yang F, Ebitz B, Grissom N, Kapoor A, Darrow DP, Herman AB.
2022. Prolonged physiological stress is associated with a lower rate of exploratory
learning that is compounded by depression. *Biol Psychiatry Cogn Neurosci*
Neuroimaging. doi:10.1016/j.bpsc.2022.12.004

Kleiner M, Brainard DH, Pelli D. 2007. What's new in Psychtoolbox-3? *Perception* **36**:1–16.

Koss MC. 1986. Pupillary dilation as an index of central nervous system α2-adrenoceptor
activation. *J Pharmacol Methods* **15**:1–19. doi:10.1016/0160-5402(86)90002-1

Lau B, Glimcher PW. 2008. Value Representations in the Primate Striatum during Matching
Behavior. *Neuron* **58**:451–463. doi:10.1016/j.neuron.2008.02.021

Loewenfeld IE. 1999. The Pupil: Anatomy, Physiology, and Clinical Applications. Butterworth-
Heinemann.

Manella LC, Petersen N, Linster C. 2017. Stimulation of the Locus Ceruleus Modulates Signal-
to-Noise Ratio in the Olfactory Bulb. *J Neurosci* **37**:11605–11615.
doi:10.1523/JNEUROSCI.2026-17.2017

Martins ARO, Froemke RC. 2015. Coordinated forms of noradrenergic plasticity in the locus
coeruleus and primary auditory cortex. *Nat Neurosci* **18**:1483–1492.
doi:10.1038/nn.4090

McGinley MJ, David SV, McCormick DA. 2015. Cortical Membrane Potential Signature of
Optimal States for Sensory Signal Detection. *Neuron* **87**:179–192.

902    doi:10.1016/j.neuron.2015.05.038

903 Moore T, Armstrong KM. 2003. Selective gating of visual signals by microstimulation of frontal
904    cortex. *Nature* **421**:370–373. doi:10.1038/nature01341

905 Moore T, Fallah M. 2001. Control of eye movements and spatial attention. *Proc Natl Acad Sci U*
906    *S A* **98**:1273–1276. doi:10.1073/pnas.98.3.1273

907 Muller TH, Mars RB, Behrens TE, O'Reilly JX. 2019. Control of entropy in neural models of
908    environmental state. *eLife* **8**:e39404. doi:10.7554/eLife.39404

909 Murphy PR, Vandekerckhove J, Nieuwenhuis S. 2014. Pupil-Linked Arousal Determines
910    Variability in Perceptual Decision Making. *PLOS Comput Biol* **10**:e1003854.
911    doi:10.1371/journal.pcbi.1003854

912 Nassar MR, Rumsey KM, Wilson RC, Parikh K, Heasly B, Gold JI. 2012. Rational regulation of
913    learning dynamics by pupil-linked arousal systems. *Nat Neurosci* **15**:1040–1046.
914    doi:10.1038/nn.3130

915 O'Byrne J, Jerbi K. 2022. How critical is brain criticality? *Trends Neurosci*.
916    doi:10.1016/j.tins.2022.08.007

917 O'Reilly JX, Schüffelgen U, Cuell SF, Behrens TEJ, Mars RB, Rushworth MFS. 2013.
918    Dissociable effects of surprise and model update in parietal and anterior cingulate
919    cortex. *Proc Natl Acad Sci* **110**:E3660–E3669. doi:10.1073/pnas.1305373110

920 Pearson JM, Hayden BY, Raghavachari S, Platt ML. 2009. Neurons in posterior cingulate cortex
921    signal exploratory decisions in a dynamic multi-option choice task. *Curr Biol CB*
922    **19**:1532–1537. doi:10.1016/j.cub.2009.07.048

923 Peixoto D, Verhein JR, Kiani R, Kao JC, Nuyujukian P, Chandrasekaran C, Brown J, Fong S,
924    Ryu SI, Shenoy KV, Newsome WT. 2021. Decoding and perturbing decision states in
925    real time. *Nature* **591**:604–609. doi:10.1038/s41586-020-03181-9

926 Pfeffer T, Keitel C, Kluger DS, Keitel A, Russmann A, Thut G, Donner TH, Gross J. 2022.
927    Coupling of pupil- and neuronal population dynamics reveals diverse influences of
928    arousal on cortical processing. *eLife* **11**:e71890. doi:10.7554/eLife.71890

929 Pisupati S, Chartarifsky-Lynn L, Khanal A, Churchland AK. 2021. Lapses in perceptual
930    decisions reflect exploration. *eLife* **10**:e55490. doi:10.7554/eLife.55490

931 Platt ML, Glimcher PW. 1999. Neural correlates of decision variables in parietal cortex. *Nature*
932    **400**:233–238. doi:10.1038/22268

933 Podvalny E, King LE, He BJ. 2021. Spectral signature and behavioral consequence of
934    spontaneous shifts of pupil-linked arousal in human. *eLife* **10**:e68265.
935    doi:10.7554/eLife.68265

936 Preacher KJ, Rucker DD, Hayes AF. 2007. Addressing Moderated Mediation Hypotheses:
937    Theory, Methods, and Prescriptions. *Multivar Behav Res* **42**:185–227.
938    doi:10.1080/00273170701341316

939 Preuschoff K, 't Hart B, Einhauser W. 2011. Pupil Dilation Signals Surprise: Evidence for
940    Noradrenaline's Role in Decision Making. *Front Neurosci* **5**.

941 Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS. 2014. Pupil
942    Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. *Neuron*
943    **84**:355–362. doi:10.1016/j.neuron.2014.09.033

944 Reimer J, McGinley MJ, Liu Y, Rodenkirch C, Wang Q, McCormick DA, Tolias AS. 2016. Pupil
945    fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. *Nat*
946    *Commun* **7**:13289. doi:10.1038/ncomms13289

947 Rescorla R, Wagner A. 1972. A theory of Pavlovian conditioning: Variations in the effectiveness
948    of reinforcement and nonreinforcementClassical Conditioning II: Current Research and
949    Theory.

950 Roesch MR, Olson CR. 2007. Neuronal Activity Related to Anticipated Reward in Frontal
951    Cortex. *Ann N Y Acad Sci* **1121**:431–446. doi:10.1196/annals.1401.004

952 Sadacca B, Wikenheiser AM, Schoenbaum G. 2017. Towards a theoretical role for tonic

norepinephrine in the orbitofrontal cortex in facilitating flexible learning. *Neuroscience* **345**:124–129. doi:10.1016/j.neuroscience.2016.04.017

Schall JD, Hanes DP. 1993. Neural basis of saccade target selection in frontal eye field during visual search. *Nature* **366**:467–469. doi:10.1038/366467a0

Scheffer M. 2020. Critical Transitions in Nature and Society, Princeton Studies in Complexity. Princeton University Press.

Scheffer M, Bascompte J, Brock WA, Brovkin V, Carpenter SR, Dakos V, Held H, van Nes EH, Rietkerk M, Sugihara G. 2009. Early-warning signals for critical transitions. *Nature* **461**:53–59. doi:10.1038/nature08227

Schultz W, Preuschoff K, Camerer C, Hsu M, Fiorillo CD, Tobler PN, Bossaerts P. 2008. Explicit neural signals reflecting reward uncertainty. *Philos Trans R Soc B Biol Sci* **363**:3801–3811. doi:10.1098/rstb.2008.0152

Slooten JCV, Jahfari S, Knapen T, Theeuwes J. 2018. How pupil responses track value-based decision-making during and after reinforcement learning. *PLOS Comput Biol* **14**:e1006632. doi:10.1371/journal.pcbi.1006632

Sobel ME. 1986. Some New Results on Indirect Effects and Their Standard Errors in Covariance Structure Models. *Sociol Methodol* **16**:159–186. doi:10.2307/270922

Tervo DGR, Proskurin M, Manakov M, Kabra M, Vollmer A, Branson K, Karpova AY. 2014. Behavioral Variability through Stochastic Choice and Its Gating by Anterior Cingulate Cortex. *Cell* **159**:21–32. doi:10.1016/j.cell.2014.08.037

Urai AE, Braun A, Donner TH. 2017. Pupil-linked arousal is driven by decision uncertainty and alters serial choice bias. *Nat Commun* **8**:14637. doi:10.1038/ncomms14637

van Kempen J, Loughnane GM, Newman DP, Kelly SP, Thiele A, O'Connell RG, Bellgrove MA. 2019. Behavioural and neural signatures of perceptual decision-making are modulated by pupil-linked arousal. *eLife* **8**:e42541. doi:10.7554/eLife.42541

Wang R, Dearing JA, Langdon PG, Zhang E, Yang X, Dakos V, Scheffer M. 2012. Flickering gives early warning signals of a critical transition to a eutrophic lake state. *Nature* **492**:419–422. doi:10.1038/nature11655

Waschke L, Tune S, Obleser J. 2019. Local cortical desynchronization and pupil-linked arousal differentially shape brain states for optimal sensory performance. *eLife* **8**:e51501. doi:10.7554/eLife.51501

Wilson RC, Bonawitz E, Costa VD, Ebitz RB. 2021. Balancing exploration and exploitation with information and randomization. *Curr Opin Behav Sci*, Computational cognitive neuroscience **38**:49–56. doi:10.1016/j.cobeha.2020.10.001

Wilson RC, Geana A, White JM, Ludvig EA, Cohen JD. 2014. Humans use directed and random exploration to solve the explore-exploit dilemma. *J Exp Psychol Gen* **143**:2074–2081. doi:10.1037/a0038199

Wu HG, Miyamoto YR, Castro LNG, Ölveczky BP, Smith MA. 2014. Temporal structure of motor variability is dynamically regulated and predicts motor learning ability. *Nat Neurosci* **17**:312–321. doi:10.1038/nn.3616

Yao T, Vanduffel W. 2023. Spike rates of frontal eye field neurons predict reaction times in a spatial attention task. *Cell Rep* **42**:112384. doi:10.1016/j.celrep.2023.112384

Yerkes RM, Dodson JD. 1908. The relation of strength of stimulus to rapidity of habit-formation. *J Comp Neurol Psychol* **18**:459–482. doi:10.1002/cne.920180503

Yokoi A, Weiler J. 2022. Pupil diameter tracked during motor adaptation in humans. *J Neurophysiol* **128**:1224–1243. doi:10.1152/jn.00021.2022

Zar JH. 1999. Biostatistical analysis. Pearson Education India.