Structure of mitoribosome reveals mechanism of mRNA binding, tRNA interactions with L1 stalk, roles of cofactors and rRNA modifications

Vivek Singh[1,14], Yuzuru Itoh[1,2,14], Samuel Del'Olio[3,14], Asem Hassan[4,5,14], Andreas Naschberger[1,6], Rasmus Kock Flygaard[7], Yuko Nobe[8], Keiichi Izumikawa[9], Shintaro Aibara[10], Juni Andréll[11], Paul C Whitford[4,5], Antoni Barrientos[3,12,13], Masato Taoka[8], Alexey Amunts[1*]

[1] Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, 17165 Solna, Sweden.
[2] Department of Biological Sciences, Graduate School of Science, University of Tokyo, 113-0033 Tokyo, Japan.
[3] Department of Molecular and Cellular Pharmacology, University of Miami Miller School of Medicine, Miami, FL 33136, United States.
[4] Department of Physics, Northeastern University, Boston, MA, 02115, United States.
[5] Center for Theoretical Biological Physics, Northeastern University, Boston, MA, 02115, United States.
[6] King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia.
[7] Department of Molecular Biology and Genetics, Danish Research Institute of Translational Neuroscience - DANDRITE, Nordic EMBL Partnership for Molecular Medicine, Aarhus University, 8000 Aarhus C, Denmark.
[8] Department of Chemistry, Graduate School of Science, Tokyo Metropolitan University, Minami-osawa 1-1, Hachioji-shi, Tokyo 192-0397, Japan.
[9] Department of Biotechnology, Global innovation research institute, Tokyo University of Agriculture and Technology, Saiwai-cho 3-5-8, Fuchu-shi, Tokyo 183-8509, Japan.
[10] BioPharmaceuticals Research and Development, AstraZeneca, UK.
[11] Department of Medical Biochemistry and Biophysics, Karolinska Institutet, 17177, Stockholm, Sweden.
[12] Department of Neurology, University of Miami Miller School of Medicine, Miami, FL 33136, United States.
[13] Department of Biochemistry and Molecular Biology, University of Miami Miller School of Medicine, Miami, FL 33136, United States.
[14] These authors contributed equally

* correspondence: amunts@scilifelab.se

The mitoribosome translates mitochondrial mRNAs and regulates energy conversion that

is a signature of aerobic life forms. We present a 2.2 Å resolution structure of human

mitoribosome together with validated mitoribosomal RNA (rRNA) modifications, including

aminoacylated CP-tRNA^{Val}. The structure shows how mitoribosomal proteins stabilise

binding of mRNA and tRNA helping to align it in the decoding center, whereas the GDP-

bound mS29 stabilizes intersubunit communication. Comparison between different states,

with respect to tRNA position, allowed to characterize a non-canonical L1 stalk, and

molecular dynamics simulations revealed how it facilitates tRNA transition in a way that

does not require interactions with rRNA. We also report functionally important

polyamines that are depleted when cells are subjected to an antibiotic treatment. The

structural, biochemical, and computational data illuminate the principal functional

components of the translation mechanism in mitochondria and provide the most complete

description so far of the structure and function of the human mitoribosome.

The human mitoribosome translates at least 13 respiratory chain protein-coding mRNAs from the mitochondrial genome [1]. It consists of three rRNAs and at least 82 mitoribosomal proteins, 36 of which have no corresponding homologs in bacterial and cytosolic ribosomes. The mitochondrial rRNA is reduced compared to its bacterial counterpart (2,512 vs 4,568 nucleotides in *Escherichia coli*), tRNA^{Val} has been incorporated as an additional constituent, and protein components are extended (~18,075 vs 7,536 residues in *E. coli*) [2]. The mitoribosomal proteins might have functional roles in mitochondria-specific aspects of translation, and therefore mutations in the corresponding genes result in cardio- and encephalomyopathies [3, 4]. However, due to the limited resolution and heterogeneity of the characterized complexes, the available models are currently incomplete, and many mitoribosomal protein key elements are described as unassigned densities [5, 6, 7, 8, 9, 10]. The existing models suggest that mitoribosomal proteins might play a role in binding mRNA, regulatory elements such as the L1 stalk are probably remodeled, and the specific mitoribosomal protein mS29, a putative GTPase is associated with inter-subunit communication. Yet, the lack of experimental data leaves open the questions of a functional involvement.

The mitochondrial rRNA is modified, and dedicated post-transcriptionally activated enzymes provide means for the regulation of gene expression and mitoribosome assembly [11]. Due to the central role of the human mitoribosome in cellular energy production, defects in rRNA modification result in fatal clinical syndromes from birth [12, 13]. Some of the bacterial counterpart modifying enzymes are missing from mitochondria, for example methyltransferases RlmA, RlmB, RmsE and pseudouridine synthase RluA, while other mitochondria-specific ones have evolved [14]. Therefore, rRNA modifications are different, and they have been identified by various experimental approaches [11, 15, 16, 17, 18, 19, 20, 21, 22, 23], furthermore recent structural studies visualized individual rRNA modifications on the mitoribosomal large (LSU) [24] and small (SSU) subunits [25, 26]. However, the entire set of modifications is yet to be quantitatively identified and visualized in the presence of bound ligands in order to enable a detailed examination of their roles. To date, the molecular mechanisms compensating for the lack of bacteria-like modifications remain unknown, as well as the modifications and the amino-acylation state of the structural tRNA^{Val}.

Mitoribosomes are also linked to age-related diseases that are associated with decline in mitochondrial function and biogenesis [27, 28, 29, 30]. Aging-related pathological changes can be inhibited by polyamine consumption [31, 32], which also has anti-inflammatory effects in mouse models *in vivo* [33]. Polyamines have been shown to be essential for mammalian cell differentiation and proliferation through their impact on translation dynamics [34, 35]. However, no polyamine biosynthetic pathways within mitochondria have been found, and their role in mitochondrial physiology remained unknown. Whether mitoribosomes might be related to these phenomena and associated with cellular senescence via polyamines needs to be explored.

To address these issues, we combined high resolution structural studies of mitoribosomal complexes with mass spectrometry based quantitative RNA analysis, biochemistry, and molecular dynamics simulations.

Structure determination

To produce a 2.2 Å overall resolution map of the mitoribosome, we collected five separate cryo-EM datasets. Selected monosome particles were subjected to per-particle defocus, beam-tilt and per-particle astigmatism correction, followed by Bayesian polishing [36, 37]. The data was then sorted into 86 optics groups based on acquisition areas followed by beam-tilt, magnification anisotropy and higher order aberration correction [38]. A final set of 509,691 particles yielded a 2.2 Å map. The local resolution was further improved by performing masked refinements around different regions of the monosome. In addition, we obtained a 2.6 Å resolution map in the presence of mRNA and three tRNAs in the classical state, and 3.0 Å resolution map with two tRNAs in the hybrid state (Supplementary Figs. 1 and 2). The improved resolution allowed building the most complete available model of the mitoribosome, including several key protein extensions involved in tRNA binding (Fig. 1a). The presence of mRNA on the mitoribosome further enabled modeling 32 of its residues, which we then used to trace mitochondria-specific elements involved in mRNA binding and put the rRNA modifications in a functional context (Fig. 1b).

To provide a description of the conformational changes associated with tRNA movement, we next focused efforts on modelling the L1 stalk. Thus, we merged particles containing tRNA in the E-site and through partial signal subtraction [39] and obtained a 2.9 Å resolution map of the region, enabling model building of the L1 stalk (Supplementary Fig. 1). We then used this model to perform molecular simulations of the complete ribosome with tRNAs.

rRNA and protein modifications, aminoacylated tRNA^{Val}, polyamines

On the rRNA level, the quality of the map allowed the modeling of 13 rRNA modifications, of which 6 are mitochondria specific compared to *E. coli* [21]. All the rRNA modifications were validated by a tailored quantitative method of Stable Isotope-Labeled riboNucleic Acid as an internal Standard (SILNAS) [40] (Supplementary Figs. 3,4). The mitochondria specific modifications include base methylations m^{5}U1076, m^{5}C1488 in the SSU; m^{1}A2617 in the LSU; and m^{1}A9, m^{2}G10 and ψ39 in the tRNA^{Val} incorporated in the central protuberance (CP) [41, 42, 43] (Fig. 1b). In the map, methylations of rRNA were identified by systematically examining the density map for the presence of unmodeled densities continuous with the base or sugar moieties of rRNA residues (Fig. 2). Identity of the two high-occupancy pseudo-uridines was established based on the presence of a structured water molecule within hydrogen-bonding distance from the N5 atom.

The tRNA^{Val} modifications are consistent with the previous mass-spectrometry data on isolated mitochondrial tRNAs from a general pool [44] (Fig. 3). The density further revealed that the tRNA^{Val} incorporated in the CP is amino acylated (Fig. 2a). In the structure, ψ39 of tRNA^{Val} contributes to the conformation of the anticodon arm, which contacts uL18m, mL38, mL40 and mL48 (Fig. 3a). In addition, two methylated residues, m^{1}A and m^{2}G, are observed at the positions 9 and 10, respectively in the D arm (Fig. 3a). m^{1}A9 is stacked between G45 and A46

151	and forms a base triple with the second base pair (C11:G24) of the D stem. On the other hand,
152	m^2G10 is stacked between C11 and A26 and forms the first base pair (m^2G10:C25) of the stem,
153	which further interacts with G45 to be a base triple. These methylations facilitate the stacking
154	and hydrogen bonds among bases, and thereby contribute to the conformation of the D arm,
155	which contacts uL18m and mL38 (Fig. 3a-c). The terminal A76 has a base-specific interaction
156	with Q175 of mL46 and together with Valine is stacked with F165 and F215 (Fig. 3d).
157	A notable feature of the cryo-EM map is the presence of continuous tubular densities associated
158	with rRNA that we assigned as polyamines, justified by the electrostatic and hydrogen-bonding
159	interactions at the amine groups (Supplementary Fig. 5). Thus, in addition to a previously
160	assigned spermine (SPM) [45], we detected one triamine spermidine (SPD) in the SSU, and a
161	diamine putrescine (PUT) and three SPDs in the LSU. The identified polyamines are bound to
162	negatively charged rRNA where mitochondria specific changes occur, providing required
163	structural support (Fig. 4; Supplementary Fig. 5). For example, SPD3301 in the LSU fills the
164	formed space of a shortened mitoribosomal protein uL2m, which has undergone reduction in the
165	human mitoribosome. SPD3303 is tethered to uL3m, where a bacterial counterpart has a
166	signature Q150 methylation, therefore replacing the post-translational modification. The PUT is
167	found in the exit tunnel upper region of uL22m hairpin.
168	Finally, the model contains seven protein modifications; three iron-sulfur clusters, one of which
169	is coordinated by uL10m and mL66 and reported here for the first time, and a molecule of
170	nicotinamide adenosine dinucleotide (NAD) reported previously [26, 45, 46] (Fig. 1b, Supplementary
171	Table 1). We also identified a disulfide bond in one of the two redox active cysteine pairs in
172	mS37 (CHCHD1) (Fig. 2) that is oxidized in the intermembrane space during mitochondrial
173	import, representing a quality control mechanism [47].
174	

Fig. 1: Structure of the human mitoribosome. a, Overview of the 2.2 Å resolution model with

mRNA and tRNAs. Proteins are shown as cartoons colored in blue for the LSU and yellow for

the SSU. The rRNA is shown as grey surface. Newly identified features are shown as spheres. **b,**

The newly identified cofactors and modifications are indicated against background of translucent

rRNA and proteins with tRNAs and mRNA shown in cartoon. We observe 13 rRNA

modifications (green), 7 protein modifications (pink), 3 iron-sulfur clusters (red-yellow),

guanosine diphosphate (GDP), adenosine triphosphate (ATP), NAD, SPM, 4 SPDs and PUT.

Fig. 2: High resolution features. Modified rRNA residues and disulfide-linked or reduced cysteine residues of mS37 in the SSU (yellow background). Modified rRNA and protein residues, and 2Fe-2S cluster in the LSU (cyan background). Their corresponding densities in the 2.2 Å map are shown as mesh at indicated threshold (bottom left) and local resolution (bottom right) levels.

Fig. 3: Interactions and post-transcriptional modifications in CP-tRNA of the mitoribosome. a, CP-tRNA[Val] (left to right) secondary structure diagram with modified nucleotides (green) and valine (cyan) highlighted, protein-contact map, B-factor representation. b,c, Interactions of the modified m^1A9 and m^2G10 nucleotides of CP-tRNA[Val]. d, Interactions of A76 and V77 of CP-tRNA[Val] with mL46. The cryo-EM density map is displayed for panels b-d (consensus map) as blue mesh. Panels (b) and (c) are related to panel (a) on the left by colored boxes.

Fig. 4: Polyamines functionally compensate for rRNA and protein alterations. Central inset shows positions of polyamines and interacting rRNA and proteins in the LSU. For each polyamine, its environment and interacting partners are shown in zoom-in panels with *E. coli* counterparts [48] (white) for comparison.

Initial binding of mRNA involves Y-N-C-Y motif coevolved with mS39

The modulation of a stable mitoribosomal complex allowed us to resolve the mRNA path and investigate mitoribosome-specific components involved in mRNA recognition (Fig. 5, Supplementary Fig. 6, Supplementary Video 1). It engages seven helical repeats of mS39 with specific contacts within a prominent positively charged groove (Fig. 5a). Particularly, a pyrimidine ring of the mRNA nucleotide in position 25 is stacked between H489 and R524 and interacts with S490 of mS39 (Fig. 5a,b). Hydrogen bonding between the side chain of R377 of mS39 and nucleotide 27 favors cytosine over uracil because the protonated N3 of a uracil would partially clash with the protonated Nη of R377. R343 of mS39 forms a hydrogen bond with the pyrimidine ring of nucleotide 28. Thus, the structure reveals a binding motif: 25 pyrimidine (Y), 27 cytosine (C), 28 pyrimidine (Y) (Fig. 5a,b). Bioinformatic analysis showed that the Y-N-C-Y motif is enriched in mitochondria and distributed over the entire length of each of the 12 open reading frames in the guanine-rich heavy DNA strand, suggesting a co-evolution of the mitochondrial DNA and the mitoribosome (Fig. 5c, Supplementary Fig. 7). The only mammalian protein-coding gene with a reduced frequency of the Y-N-C-Y motif is the *MT-ND6* that is on the light strand and codes for an 18-kDa subunit of complex I. The expression of the ND6 mRNA is unique, as it requires an auxiliary factor FASTK to generate the mature form, suggesting a dedicated regulatory mechanism [49].

The most 5' contact between mRNA and mS39 is ~50 Å from the channel entrance, and by following the mRNA density toward the decoding center, we identified six specific mitoribosomal proteins that are important for mRNA recruitment, channel opening, and precise alignment of the message for translation in the decoding center. In our structure, the Y-N-C-Y motif forms a ~100° turn (kink-1) in the mRNA midway through mS39 (Fig. 5a and Supplementary Video 1). After the kink, the N-terminal mitochondria-specific extension of uS5m supports the path of mRNA over a distance of ~60 Å, involving a unique N-terminal polybasic stretch of 14 residues 111-124 (KKGRGKRTKKKKRK) together with R237 and K239 (Fig. 5d, Supplementary Video 1, Supplementary Fig. 6). From the side of the SSU head, mS35 provides potential stabilizing electrostatic interactions to mRNA with N-terminal residues K46 and K49 (Fig. 5e,f). This mitochondria-specific extension of uS5m structurally replaces bacterial bS4 and uS3-CTD that interact with mRNA (Supplementary Fig. 6) and confer RNA helicase activity on the bacterial ribosome [50]. Just before the channel entry, the binding of mRNA induces a conformational change of the N-terminal extension of uS9m that contacts the mRNA nucleotide at position 15 via the backbone of V55 (Fig. 5e,g; Supplementary Video 1), which in the absence of mRNA blocks the entrance to the decoding center [51]. Thus, uS9m exists in a closed conformation in the absence of mRNA, and upon mRNA threading, a conformational change in uS9m allows the passage (Fig 5h).

Fig. 5: Specific protein elements involved in mRNA binding.

a, Electrostatic coulomb potential surface representation of the model shows a positively charged groove (blue) on mS39 that accommodates the mRNA. N-terminal loop of mS31 interacts with mS39. Y-N-C-Y motif and its interactions with mS39 residues are depicted as a schematic. Position of kink-2 is marked. **b,** Interactions of the Y-N-C-Y motif with mS39 and density map (mesh). Protein mS31 has been removed for clarity. **c,** Average percentage frequency of Y-N-C-Y motif (left) with +/- standard deviation for mitochondrial mRNAs (excluding ND6; dark grey bar) and nuclear mRNA (brown bar). P-value (<0.0001) calculated by Welch's unpaired t-test. Positions of the Y-N-C-Y motif in the mt-mRNAs are shown as discs (right) to depict its distribution along the mt-mRNA sequences. The total number of residues for each mt-mRNA is indicated. **d,** Overview of uS5m showing a region of contact with mRNA. The polybasic stretch 111-124 along the mRNA is colored by electrostatic potential. Interacting residues between uS5m and mRNA are shown with the local density map. **e,** Interactions of mS35 and uS9m with mRNA at the channel entry and the P-site. **f,** R46 and R48 of mS35 potentially interact with the mRNA backbone, and the density suggests flexibility of the side-chains. **g,** uS9m forms stacking and hydrogen bonding interactions with mRNA position 15 via F54 and V55, respectively. **h,** uS9m N-terminus (purple cartoon) adopts alternative conformations that result in mRNA channel blocked or open states regulating mRNA access to the SSU (gold surface).

mL40 N-terminal tail stabilizes the A- and P-site tRNAs and decoding interactions

We then examined the role of mitoribosomal proteins in the mRNA alignment. The structure shows that to enter the SSU channel, the mRNA must be kinked again (kink-2), and in the A-site the nucleotide at position 9 is stabilized by uS12m (Supplementary Fig. 8). Here, a conserved K72 side chain forms a salt bridge with the phosphate of nucleotide 9 (last nucleotide in the A-site codon). At the same time, *cis*-proline P73 (Supplementary Fig. 8) coordinates two potassium ions, one of which directly coordinates 2′O of nucleotide 9 (Supplementary Fig. 8, Supplementary Video 1) and stabilizes the decoding center. The densities identified as two potassium ions have been observed previously, but modeled as Mg^{2+} (PDB 4V51) [52] or waters (PDB 7K00) [48, 52]. However, they have been experimentally identified as K^+ by long wavelength X-ray diffraction analysis (PDB 6QNR) [53], which agrees with our assignment (Supplementary Fig. 8).

Next, between the A- and P-site, the map reveals a third kink in the mRNA (kink-3), which is co-localized with a 35-Å long density extending from the CP in between the two tRNAs (Fig. 6a, Supplementary Fig. 9a) that could not be previously interpreted. We identified it as a positively charged N-terminus of mL40 with the terminal backbone amino group of S47 forming a salt bridge with the backbone phosphate of the mRNA at position 7 in the A-site (Fig. 6a, Supplementary Fig. 9b), and the interaction is further stabilized via a magnesium ion coordinated by the phosphate of G1485. In addition, the hydroxyl group of S47 interacts with the mRNA nucleotide base at position 6 in the P-site (Fig. 6a, Supplementary Video 1), although the contacts are likely to be flexible. Moreover, the base orientation allows any nucleotide to interact with S47 (Supplementary Fig. 9c). The concept is analogous to the cytosolic ribosome that

engages uS19 for tRNA stabilization and mRNA interactions in the decoding center [54] (Fig. 6b).
The presence of four lysine residues 52-55 (Fig. 6a, Supplementary Fig. 9b), might suggest that
mL40 contributes to balancing the charge between the A- and P-tRNAs. Together with R51,
which forms salt bridges with A1560 and C1561 in h44, the positively charged patch
compensates for the negative RNA charges accumulated in that region (Fig. 6a, Supplementary
Fig. 9b, Supplementary Video 1). Further in the P-site, we observe a well-resolved density
showing the conserved C-terminal R396 of uS9m interacting with the anticodon loop of the P-
site tRNA (Supplementary Fig. 9d). Finally, in the E-site, uS7m residues G164-165 further
contribute to mRNA stabilization (Supplementary Fig. 9e). Thus, mitoribosomal proteins play a
role in mRNA binding.

292
Fig. 6: Analysis of mRNA binding in tRNA sites. a, Interactions of mL40 (shown with
electrostatic potential of the model) with mRNA between A/A- and P/P-tRNAs. Hydrogen bonds
G6 base and G7 phosphate are shown as dashed lines. **b,** Comparison of mL40-tRNA interaction
(left) with those of uS19 (middle) and uS13 (right) in human cytosolic [54] and *T. thermophilus*
ribosome [52], respectively.

mRNA alignment in the decoding center and rRNA modifications

In the decoding center, the universally conserved G902 is in *anti* conformation, while A1557 and
A1558 are flipped out from h44 to interact with the A-site codon:anti-codon helix in the
canonical manner [55, 56] (Supplementary Fig. 10). A well-resolved density, combined with the
mass-spectrometry SILNAS [40, 57, 58] approach further allowed us to validate the complete set of
the rRNA modifications, including six mitochondria specific, and propose a role in stabilizing
the decoding center (Fig. 5, Supplementary Figs. 4 and 5). Close to the decoding center, the
residue G899 was recently shown to interact with streptomycin in human mtSSU [45]. Its

methylated to m^7G527 in *E.coli* (Supplementary Fig. 10) and this modification partly contributes to streptomycin sensitivity [59]. In the P-site, two base-methylated nucleotides m^4C1486 and m^5C1488 flank the base triple C1487:A1564:U1563 by stabilizing the stacking interactions (Fig. 5a,b). The orientation of N^4-methylated m^4C1486 towards mRNA is conserved with respect to *E. coli*, where the corresponding residue is also 2'-O methylated ($m^4Cm1402$) (Supplementary Fig. 10). m^4C1486 together with C1487 interacts with the phosphate group of the last nucleotide in the P-site mRNA codon via hydrogen bonds (Fig. 5a,b). Further, phosphate group of m^4C1486 coordinates a magnesium ion, which stabilizes a conserved kink of the mRNA between A- and P-sites (Fig. 7a). In contrast to the *E. coli* P-site, U1563 does not have an N^3-methylation (m^3U1498 in *E. coli*) (Supplementary Fig. 10). This lack of N^3-methylation allows U1563 together with C1487 to form a water mediated interaction with the P-site codon (Fig. 7a,b). Further, the conserved dimethylated residues, m^6_2A1583 and m^6_2A1584 orient the nucleotide U1563 such that the ribose of U1563 forms a hydrogen bond to a phosphate group of the second nucleotide in the P-site mRNA codon, while the phosphate of U1563 forms a hydrogen bond with the ribose on the first codon nucleotide (Fig. 7b). The two dimethylated nucleotides facilitate A1080 positioning for the 2'-OH to hydrogen bond with N3 of m^5U1076. This together with methylation of m^5U1076 contributes to the stability of h24 loop that interacts with the P-tRNA anticodon stem at position 38 via A1078 (Fig. 7a,b).

In the A-loop of H92, there are two 2'-O-methyl modified nucleotides Um3039 and Gm3040. The 2'-O-methyl of Um3039 sterically positions the bases of Gm3040 and U3041 to form interactions with the nucleotides C75 and C74 of the A-tRNA, respectively (Fig. 7c,d). The 2'-O-methyl of Gm3040 present in the cytosolic but not bacterial ribosome and interacts with the base of A3069 to stabilize the conformation of Gm3040. Furthermore, the bacterial counterpart rRNA that interacts with A-tRNA (C75) is G:C, whereas in our structure it's A:U (3069:2994) that is less stable, and thus stabilized by the mitochondria-specific methylation of Gm3040 and base-pairing interactions with ψ3067 (Fig. 7d). In the LSU, mitochondria-specific m^1A2617 positions U2614 into a wedged interaction (Fig. 7e). Surprisingly, there is only weak sugar puckering for m^1A2617 (Fig. 2a). It bears a partial positive charge at N1 and interacts with U2649 of H61. The interactions mediated by m^1A2617 potentially contribute to the structuring of H71 loop that participates in an intersubunit bridge shares an interface with the A-loop (Fig. 7e). The universally conserved Gm2815 in the P-loop forms Watson-Crick base pairing with C75 of P/P-tRNA, similar to the interactions of Gm3040 of the A-loop (Fig. 7f).

Fig. 7: Post-transcriptional modifications in functional centers of the mitoribosome.
a, Base-methylated m^{4}C1486 and m^{5}C1488 and the interactions at the P-site mRNA codon. **b,**
The dimethylated m$^6_2$A1583, m$^6_2$A1584 and base-methylated m^{5}1076 facilitate interactions with
mRNA codon and P-tRNA. **c,** 2′-*O*-methyl modified Um3039 and Gm3040 lead to interactions
with A-tRNA. **d,** Gm3040 and Ψ3067 contribute to the interface with the A-tRNA. **e,** m^{1}A2617
contributes to a stabilization of inter-subunit bridge and the A-loop. **f,** Gm2815 base-pairs with
P-tRNA. Cryo-EM density map (A/A P/P E/E state) is displayed as a blue mesh.

Structure of the mitoribosomal L1 stalk reveals its unique protein features
To investigate whether mitoribosomal proteins might also have influence on tRNA
rearrangement in the E-site, we aimed to perform molecular simulations of the mitochondrial L1
stalk. Therefore, we modeled the complete structure of the L1 stalk, which relies on
mitochondria specific protein elements. In the structure, the E-tRNA is stabilized via
mitochondria-specific protein elements of the C-terminal domain of uL9m and 33-residue N-
terminal extension of uL1m (Fig. 8a,b). The orientation of the uL9m C-terminal domain adopts a

previously unseen folded conformation involving a ~75° rotation towards the E-tRNA (Fig. 8c).
This unexpected conformational difference is facilitated by the C-terminal helix in the extension
that reaches and contacts the non-base-pairing part and the distal short hairpin of the rRNA H76
(Fig. 8d). The resulting projection of uL9m towards the ribosomal core reduces the distance
between the central helix and the C-terminus of uL1m from ~70 to only ~30 Å. This distance is
spanned by the uL1m mitochondria-specific extension that forms direct contacts with the central
helix of uL9m (Fig. 8d-f). Therefore, our model of the L1 stalk establishes a tertiary
mitoribosomal complex uL9m:uL1m:rRNA-H76, where uL9m acquires a new functional role.
From the opposite side, the mitochondria-specific mL64 approaches the T-arm through its long
C-terminal helix extending to the tRNA binding sites.

365
366 **Fig. 8: Structural features of the L1 stalk and interactions with tRNA.**

a, The L1 stalk components uL9m (green), uL1m (purple) rRNA helix H76-77 (residues 2757-2790, light blue), as well as mL64 (magenta) interact with E/E-tRNA. The structure and interactions are different from *T. thermophilus* (PDB ID: 4V51) [52] **b,** Components of the L1 stalk modeled into their respective densities (shown as mesh). **c,** Comparison of uL9m conformation with that of *T. thermophilus* uL9 (white PDB ID 4V51). **d,** Model of uL1m stalk with uL9m in green and uL1m in purple, highlighting mitochondria-specific protein elements. **e,** 2D diagram of uL9m showing interactions with E/E-tRNA (brown bars), rRNA (light-blue bars), uL1m (dark-blue bars). Conserved regions are shown in light and mitochondria-specific in dark colors. **f,** 2D diagram of uL1m color-coded as in panel d.

Recognition of tRNA by the L1 stalk and intersubunit communication

We next used an all-atom structure-based model [60, 61] to simulate hundreds of transitions of the tRNA from the P site to P/E state (Fig. 9a,b; Supplementary Video 2). To isolate the influence of uL1m and mL64, we compared the dynamics with four variants of the model (Fig. 9c; see Methods): 1) baseline model; 2) truncated mL64 (ΔmL64) with C-terminal region starting with Q112 being removed to prevent steric interactions between mL64 and EE-tRNA without disturbing the rest of the protein.; 3) modified uL1m (ΔuL1m) where stabilizing interactions between uL1m and P-site-tRNA are scaled down compared to the baseline model (values available in methods); 4) both ΔmL64 and ΔuL1m modifications are included (see methods section for details of the models). Using these models, we simulated 1066 spontaneous (non-targeted) transitions between the classical (A/A P/P) and hybrid (A/P P/E) states, which include a ~40 Å displacement of the P-tRNA and spontaneous rotation of the SSU (Supplementary Video 2). A mechanistic insight from these calculations is that mL64 imposes strict limits on the pathways that are accessible to tRNA, which thereby favors a specific sequence of conformational substeps during P/E formation (Fig. 9d-g). As a result, mL64 ensures that the tRNA is properly positioned and may reliably engage uL1m as the tRNA approaches the E site. The structure also shows that due to the different type of tRNA binding, no base pairs are formed between the E/E-tRNA and mRNA, unlike those found in bacteria [62] (Supplementary Fig. 9e). Instead, the protein uS7m undergoes a conformational change in which residues G164-165 directly contact the mRNA and stabilize its binding (Supplementary Fig. 9e, Supplementary Video 1). Thus, upon each translation step, a tRNA is recognized in the E-site and escorted from the mitoribosome without involving rRNA, which is strikingly different from bacterial and cytosolic counterparts.

Fig. 9: Simulations of P/E formation reveal the influence of mitochondria-specific proteins.
a, Distances R_{elbow} and R_{CCA} are used to describe the position of the tRNA during P/E
formation. **b,** Probability distribution as a function of P-site tRNA elbow position (R_{elbow}) and the
fraction of uL1m-tRNA contacts ($Q_{L1\text{-}tRNA}$) relative to those formed in the E/E state, calculated
from simulations that used the baseline model (left). When uL1m-P-tRNA contacts are
attenuated (ΔuL1m model; right), $Q_{L1\text{-}tRNA}$ remains low, consistent with a lack of uL1m
binding. **c,** In simulations of P/E formation, perturbations to uL1m and/or mL64 (models
ΔuL1m, ΔmL64, and ΔmL64-ΔuL1m models) lead to a wider range of tRNA elbow positions
and longer timescales for P/E formation. The narrower range of pathways accessed with the
complete/baseline model leads to a lower average value of R_{elbow} as a function of the tRNA CCA
position R_{CCA}, showing how the corridor favors a process where the elbow motion precedes CCA
movement. **d,** Probability distribution (327 simulated events) as a function of elbow and 3′-CCA
position reveals distinct intermediates during P/E formation. Representative structural snapshots
of I1-I3 are shown in panels e-g. **e,** Intermediate I1 is associated with a small displacement of the
tRNA elbow and CCA tail toward the E site. **f,** In order to reach I2, the tRNA elbow is displaced
by an additional ~10Å, where it it confined by contacts with mL64. **g,** mL64 then guides the
tRNA elbow towards the E site, allowing it to adopt I3. After reaching I3, a final ~40Å
displacement of the CCA tail results in adoption of the P/E conformation.

**Mitoribosome function is independent of mS29 GTPase activity, but GDP stabilizes
intersubunit communication**

For each translation step, the movement of mRNA:tRNA is also accompanied by the SSU head
swiveling, but it has remained an open question whether intrinsic GTPase activity of mS29 (also
known as death-associated protein 3, DAP3) is coupled to the process [63, 64]. We now resolve it by
analyzing the structure in two different conformations combined with structure-based
mutagenesis and biochemical analysis (Fig. 10, Supplementary Fig. 11a-c). Particularly, we
performed site-directed mutagenesis targeting the previously reported nucleotide-binding site [2, 63]

and found that the mutant K134A, which based on the structure would affect the nucleotide
binding, appears to be dispensable for mitoribosomal function (Supplementary Fig. 11a,b).
Recently, another putative binding site has been proposed, formed with residues 173, 177, 208,
209, 210, 238, 242, 245, 291, 292, 295 [45, 46] (Fig. 10a,b, Supplementary Fig. 11d), and our map at
the local resolution of 2.65 Å suggests the presence of GDP in that site. To address the functional
relevance of the GDP, we compared it with the SSU structure in the presence of a
nonhydrolyzable analog GMPPNP [45] (Fig. 10a,b). It showed a very similar global architecture of
mS29, implying that no net conformational change is involved upon GTP hydrolysis (Fig.
10a,b). Furthermore, there is no loss of interactions of mS29 with its binding partners in the SSU
head (Fig. 10b). Together, the data suggest that the GTPase activity is not coupled to the
translation cycle and the related SSU head swiveling.
We next used structural and biochemical analysis to identify the role of the GDP. We compared
the interactions of mS29, which is the signature protein of the SSU head with the CP between
our classical (2.6 Å resolution) and hybrid (3.0 Å resolution) structures. In the classical state,
they are extensive and manifest through three contact sites. Site-1 and -2 are formed between
GDP-associated region and the respective mL46 helices α2 and α3 extending from the CP.
Contact site-3 is formed with mL48 β-turn protruding into the intersubunit space. Upon subunit
ratcheting, in the hybrid state mS29 shifts resulting in disruption of contact site-3. Further, the
mS29 β-hairpin 208-216 (stabilized by GDP) that formed contact site-2 in the classical state
switches in the hybrid state and remains the only inter-subunit bridge in the dimeter of 50 Å (Fig.
10c, Supplementary Fig. 11e). To assess the role of GDP binding in translation, we generated
cell lines stably expressing mS29 mutants to disrupt the GDP binding site (Fig. 10d-g,
Supplementary Fig. 11a-c).
Expression of the double mutants R177A/K245A or R177A/K295A, which target residues
interacting with the phosphate groups of GDP, led to reduced levels of mitochondrion-encoded
proteins and cellular respiration (Fig. 10d-g, Supplementary Fig. 11c), suggesting a stabilizing
role for GDP in this pocket. The R177A/K245A mutation shows only a tendency toward a more
stringent effect on mitochondrial protein synthesis than the R177A/K295A mutation in our short
15 min pulses. However, the effect is significant when looking at the steady-state levels of
mtDNA-encoded proteins. (Fig. 10d,e). Interestingly, the double mutant R177A/H291A
displayed a rate of mitochondrial protein synthesis comparable to WT (Fig. 10d-g). Although
K245 and K295 stabilize the same beta phosphate on GDP, it is possible that H291 can
compensate for the loss of K295, explaining the decreased stability of the R177A/K245A mutant
compared to the R177A/K295A mutant. Change of nearby hydrophobic residues Y173A/Y208A
yields a decrease in translation similar to the R177A/K245A implying their role in GDP
stabilization (Fig. 10d-g). Lastly, double mutant pairs V209P/D238A and Y208A/D238A, which
directly disrupt GDP base binding to the mS29 β-hairpin, showed drastic decreases in the
synthesis and stability of mitochondrion-encoded proteins (Fig. 10d-g). Thus, the biochemical
assays based on mutagenesis show that GDP binding to mS29 is required for efficient
mitochondrial translation. Our structure-function correlation analyses suggest that GDP acts as a

stabilizing agent for mS29 β-hairpin 208-216 to maintain the intersubunit communication as
tRNA moves to the hybrid state.

470
Fig. 10: Binding of GDP to mS29 and role in mitoribosome function.

a, Superposition of GDP-bound (yellow) and GMPPNP-bound (white) states. **b,** Close-up view
illustrating side chains that coordinate nucleotide binding. **c,** Intersubunit contacts in the classical
and hybrid states mediated by mS29 (cryo-EM density map shown as colored surface). **d,**
Immunoblot analysis comparing steady-state levels of proteins using whole cell lysates from
HEK293T (WT), *mS29*-KO (KO), and *mS29*-KO cells stably expressing either WT mS29 or
double mutants targeting the GDP-mS29β-hairpin binding site under the control of an attenuated
CMV6 promoter (Δ5pCMV6). COX1, COX2, and CTYB were used as markers of mitoribosome
function, and β-Actin was used as a loading control. **e,** Images in panel (d) were digitalized, and
densitometry was conducted using the histogram function in Adobe Photoshop. The bar graphs
represent the mean ± SD from three independent repetitions. Statistical differences were
estimated using one-way ANOVA in GraphPad Prism. *$p<0.05$, **$p<0.01$; ***$p<0.001$,
****$p<0.0001$. **f,** Metabolic labeling of mitochondrially translated peptides in whole cells from
the indicated cell lines using ^{35}S-methionine for 15 minutes in the presence of emetine to inhibit
cytosolic protein synthesis. Immunoblotting for β-ACTIN was used as a loading control. Newly
synthesized peptides are identified on the left. **g**, Images in panel (d) were digitalized, and
densitometry was conducted using the histogram function in Adobe Photoshop. The bar graphs
represent the mean ± SD from four independent repetitions. Statistical differences were
estimated using one-way ANOVA in GraphPad Prism. *$p<0.05$, **$p<0.01$; ***$p<0.001$,
****$p<0.0001$. The source data file contains all original uncropped and unprocessed
immunoblots as well as densitometry values used to perform statistics and generate
quantification graphs in panels (d-g).
493

Discussion

This study illustrates the modifications and cofactors of the human mitoribosome (Fig. 1-4,
Supplementary Fig. 3-5) and explains how mRNA and tRNAs are stabilized (Fig. 5-9,
Supplementary movie 1). While the basic mechanism of translation and the rRNA in the
decoding center are conserved, our structures attribute additional roles to mitoribosomal proteins
in mRNA and tRNA binding.

For mRNA, previous work showed that the delivery of some mitochondrial transcripts for
translation is facilitated by the factor LRPPRC [65, 66]. We now demonstrate that once an mRNA is
received by the mitoribosome, it involves six mitochondria-specific proteins that sequentially
help to direct the mRNA to the decoding center, where it's aligned with a tRNA (Fig. 5,6,
Supplementary Fig. 6,8-9). Most of these proteins compensate for the reduced rRNA through the
mechanisms of structural patching [67] and protein ratcheting [68] that drive neutral evolution of
protein complexity. The first protein to receive an mRNA is mS39, and we detected a co-
evolution of mS39 with an mRNA motif Y-N-C-Y that is enriched in all 12 transcripts encoded
in the heavy strand of the mitochondrial DNA. The only mRNA that is not enriched with the
motif is ND6 (Fig. 5, Supplementary Fig. 7). Interestingly, *ND6* possesses a long 3′UTR and is
the only transcript in human mitochondria lacking a polyA tail. This gene is also the only one in
mitochondria of vertebrates that was found to be translocated to the control region through gene

duplication [69]. This occurs in Antarctic fish species, and while the selectively advantageous
duplicate is functionally neutral on the protein level [70], we found that there is a change in the Y-
N-C-Y frequency in the mRNA sequence, which is increased by 2-fold (from 14 to 30
occurrences in 525 residues). It suggests a possible adaptive correlation between the mRNA
stability on the mitoribosome and mt-DNA reorganization.

For tRNA, the structure demonstrates two cases with an involvement of mitoribosomal proteins.
First, between the A- and P-site tRNAs, the N-terminus of mL40 extends from the CP to the
phosphate backbone of mRNA. Second, the recognition of a tRNA in the E-site is achieved by a
protein-made L1 stalk region and mL64. Both architectures are unique to mitochondria. Our
molecular dynamics simulations further illustrate how the process of E-tRNA movement
fundamentally relies on specific protein extensions and insertions of uL1m and uL9m without
any involvement of rRNA (Fig. 8,9, Supplementary movie 2). These adaptations of
mitoribosomal proteins are particularly important in the context of different functional states, as
previous structural studies of elongation [8], termination [10], and rescue [6] complexes reported
unidentified densities associated with translational factors, which we now assign specific
proteins. Since our data resolves those densities as mammalian protein extensions (Fig. 5,
Supplementary Fig. 6), the structural information can now be used for reinterpretation of the
functional complexes, thus completing previous works.

In addition to proteins, newly revealed structural details include polyamines as stabilizing
intramolecular agents in human mitochondria. The visualization of polyamines in the human
mitoribosome and the reduced occupancy in the presence of antibiotics further supports their
stimulating effect on mitochondrial translation, which is essential for cell proliferation. In
mitoribosomes from cell culture treated with antibiotics [45, 71], the occupancy of the polyamines is
reduced [45], and they are generally replaced with Mg^{2+} or K^+, accompanied by some residue
rearrangement (Supplementary Fig. 5). Those changes are associated with attenuation of
translation, and thus our structure rationalizes previous report that the positive effect of
polyamines could not be functionally replaced by the addition of Mg^{2+} in a cell-free system [72].
Together, it links mitochondrial translation to reports assigning key regulatory effects to
polyamines in mitochondria. This includes anti-inflammatory effects and inhibition of aging-
associated pathologies upon consumption of polyamine in mice [31, 32, 73], the ability to boost
mitochondrial respiratory capacity in a *Drosophila* aging model [32], restorative effects on
oxidative functions of aged mitochondria [74, 75], and antitumor immunity in aged mice [76].
Polyamines were also proposed to regulate mitochondrial metabolism via Src kinase signaling to
preserve bioenergetics in the spinal cord [77]. Considering our results on the involvement of iron-
sulphur clusters as protein-protein mediating agents [45], as well as in the mitoribosomal assembly
[78], the new structural data provides a wealth of additional information on cofactors that can be
further assessed in disease models and with respect to tissue and cell specificity, as well as
ageing processes. The detection of disulfide bond in mS37 that was proposed to occur during the
mitochondrial protein import pathway [47] further links translation with metabolic state and

NADH/NAD$^+$ ratio in the intermembrane space and the matrix due to the special role of mS37
that links assembly to the initiation of translation [25].

On the rRNA level, through tailored quantitative mass-spectrometry, we describe a fully
modified mitoribosome, including in the decoding center. While the critical modifications are
conserved with bacteria, an unexpected feature of the structure is that the rRNA component CP-
tRNA^{Val} is modified, as well as aminoacylated. This suggests that this tRNA is recognized in the
matrix by the elongation factor mtEF-Tu prior the assembly onto the CP. The involvement
mtEF-Tu in the mitoribosome assembly has been previously reported, and direct interaction with
the assembly factor GTPBP5 was visualized by cryo-EM of an LSU intermediate [79]. Thus, the
regulation of the assembly of the mammalian mitoribosome relies on the availability of mtEF-Tu
in active developmental stages of the organelle, which might be a potential early checkpoint for
translation. A possible model for these observations is that accumulation of deacylated tRNAs
might trigger starvation responses, and since tRNA^{Val} is likely to be abundant at the onset of
mitoribosomal assembly, its aminoacylation is essential, which requires the presence of mtEF-
Tu. The feature is likely to be mammalian-specific, as representative from other lineages, such as
fungi [46], ciliates [80], and algae [81, 82] do not possess the CP-tRNA component in their
mitoribosomes.

Finally, the data addresses a long-standing question of whether a putative GTPase and death-
associated protein mS29 might provide the human mitoribosome with an intrinsic activity, while
undergoing structural rearrangements. This protein is of a particular clinical interest, because
several cancer types are associated with its aberrant expression [83]. We show that mS29-GDP
does not change its conformation during translation but is a stable structural element that
contributes to inter-subunit communication during transitions between classical and hybrid states
(Fig. 10, Supplementary Fig. 11).

Overall, these data explain how the newly identified components modulate the functional and
structurally distinct mitoribosome. With the continuous progress in improvement of resolution of
cryo-EM reconstructions, and availability of newly developed methods for base editing of
protein-coding genes in mammalian mitochondrial genome [84, 85], the field of mitochondrial
translation now steps into a phase, where it becomes possible to analyze fine regulatory
mechanisms specific transcripts, and study how it's affected by modifications, cofactors, and
mutants. Therefore, the structural approach has a potential to lead towards a more detailed
description of the mitoribosome-associated pathologies, including mitochondrial disease and
ageing, and the current study will provide a reference model of the human mitoribosome for
future investigation.

References
1.   Kienzle L*, et al.* A small protein coded within the mitochondrial canonical gene nd4
regulates mitochondrial bioenergetics. *BMC Biol* **21**, 111 (2023).
589

590   2.   Amunts A, Brown A, Toots J, Scheres SHW, Ramakrishnan V. Ribosome. The structure
591   of the human mitochondrial ribosome. *Science* **348**, 95-98 (2015).
592

593   3.   Ferrari A, Del'Olio S, Barrientos A. The Diseased Mitoribosome. *FEBS Lett* **595**, 1025-
594   1061 (2021).
595

596   4.   Amarasekera SSC*, et al.* Multi-omics identifies large mitoribosomal subunit instability
597   caused by pathogenic MRPL39 variants as a cause of pediatric onset mitochondrial
598   disease. *Hum Mol Genet*, (2023).
599

600   5.   Aibara S, Singh V, Modelska A, Amunts A. Structural basis of mitochondrial translation.
601   *Elife* **9**, (2020).
602

603   6.   Desai N, Yang H, Chandrasekaran V, Kazi R, Minczuk M, Ramakrishnan V.
604   Elongational stalling activates mitoribosome-associated quality control. *Science* **370**,
605   1105-1110 (2020).
606

607   7.   Koripella RK*, et al.* Structures of the human mitochondrial ribosome bound to EF-G1
608   reveal distinct features of mitochondrial translation elongation. *Nat Commun* **11**, 3830
609   (2020).
610

611   8.   Kummer E, Ban N. Structural insights into mammalian mitochondrial translation
612   elongation catalyzed by mtEFG1. *EMBO J* **39**, e104820 (2020).
613

614   9.   Kummer E*, et al.* Unique features of mammalian mitochondrial translation initiation
615   revealed by cryo-EM. *Nature* **560**, 263-267 (2018).
616

617   10.   Kummer E, Schubert KN, Schoenhut T, Scaiola A, Ban N. Structural basis of translation
618   termination, rescue, and recycling in mammalian mitochondria. *Mol Cell* **81**, 2566-2582
619   e2566 (2021).
620

621   11.   Rebelo-Guiomar P, Powell CA, Van Haute L, Minczuk M. The mammalian
622   mitochondrial epitranscriptome. *Biochim Biophys Acta Gene Regul Mech* **1862**, 429-446
623   (2019).
624

625   12.   D'Souza AR, Minczuk M. Mitochondrial transcription and translation: overview. *Essays*
626   *Biochem* **62**, 309-320 (2018).
627

628   13.   Garone C*, et al.* Defective mitochondrial rRNA methyltransferase MRM2 causes
629   MELAS-like clinical syndrome. *Hum Mol Genet* **26**, 4257-4266 (2017).
630

631   14.   Lopez Sanchez MIG, Cipullo M, Gopalakrishna S, Khawaja A, Rorbach J. Methylation
632   of Ribosomal RNA: A Mitochondrial Perspective. *Front Genet* **11**, 761 (2020).
633

15. Antonicka H, Choquet K, Lin ZY, Gingras AC, Kleinman CL, Shoubridge EA. A pseudouridine synthase module is essential for mitochondrial protein synthesis and cell viability. *EMBO Rep* **18**, 28-38 (2017).

16. Baer RJ, Dubin DT. Methylated regions of hamster mitochondrial ribosomal RNA: structural and functional correlates. *Nucleic Acids Res* **9**, 323-337 (1981).

17. McCulloch V, Seidel-Rogol BL, Shadel GS. A human mitochondrial transcription factor is related to RNA adenine methyltransferases and binds S-adenosylmethionine. *Mol Cell Biol* **22**, 1116-1125 (2002).

18. Metodiev MD*, et al.* Methylation of 12S rRNA is necessary for in vivo stability of the small subunit of the mammalian mitochondrial ribosome. *Cell Metab* **9**, 386-397 (2009).

19. Metodiev MD*, et al.* NSUN4 is a dual function mitochondrial protein required for both methylation of 12S rRNA and coordination of mitoribosomal assembly. *PLoS Genet* **10**, e1004110 (2014).

20. Seidel-Rogol BL, McCulloch V, Shadel GS. Human mitochondrial transcription factor B1 methylates ribosomal RNA at a conserved stem-loop. *Nat Genet* **33**, 23-24 (2003).

21. Sergiev PV, Aleksashin NA, Chugunova AA, Polikanov YS, Dontsova OA. Structural and evolutionary insights into ribosomal RNA methylation. *Nat Chem Biol* **14**, 226-235 (2018).

22. Van Haute L*, et al.* METTL15 introduces N4-methylcytidine into human mitochondrial 12S rRNA and is required for mitoribosome biogenesis. *Nucleic Acids Res* **47**, 10267-10281 (2019).

23. Zaganelli S*, et al.* The Pseudouridine Synthase RPUSD4 Is an Essential Component of Mitochondrial RNA Granules. *J Biol Chem* **292**, 4519-4532 (2017).

24. Hillen HS*, et al.* Structural basis of GTPase-mediated mitochondrial ribosome biogenesis and recycling. *Nat Commun* **12**, 3672 (2021).

25. Itoh Y*, et al.* Mechanism of mitoribosomal small subunit biogenesis and preinitiation. *Nature* **606**, 603-608 (2022).

26. Harper NJ, Burnside C, Klinge S. Principles of mitoribosomal small subunit assembly in eukaryotes. *Nature* **614**, 175-181 (2023).

27. Bobba-Alves N*, et al.*, (2022).

28. Haas RH. Mitochondrial Dysfunction in Aging and Diseases of Aging. *Biology (Basel)* **8**, (2019).

680   29.   Hong HJ, *et al.* Mitoribosome insufficiency in beta cells is associated with type 2
681   diabetes-like islet failure. *Exp Mol Med* **54**, 932-945 (2022).
682

683   30.   Richman TR, *et al.* Mitochondrial mistranslation modulated by metabolic stress causes
684   cardiovascular disease and reduced lifespan. *Aging Cell* **20**, e13408 (2021).
685

686   31.   Soda K, Dobashi Y, Kano Y, Tsujinaka S, Konishi F. Polyamine-rich food decreases age-
687   associated pathology and mortality in aged mice. *Exp Gerontol* **44**, 727-732 (2009).
688

689   32.   Soda K, Kano Y, Chiba F, Koizumi K, Miyaki Y. Increased polyamine intake inhibits
690   age-associated alteration in global DNA methylation and 1,2-dimethylhydrazine-induced
691   tumorigenesis. *PLoS One* **8**, e64357 (2013).
692

693   33.   Medina CB, *et al.* Metabolites released from apoptotic cells act as tissue messengers.
694   *Nature* **580**, 130-135 (2020).
695

696   34.   Igarashi K, Kashiwagi K. The functional role of polyamines in eukaryotic cells. *Int J*
697   *Biochem Cell Biol* **107**, 104-115 (2019).
698

699   35.   Mandal S, Mandal A, Johansson HE, Orjalo AV, Park MH. Depletion of cellular
700   polyamines, spermidine and spermine, causes a total arrest in translation and growth in
701   mammalian cells. *Proc Natl Acad Sci U S A* **110**, 2169-2174 (2013).
702

703   36.   Zivanov J, *et al.* New tools for automated high-resolution cryo-EM structure
704   determination in RELION-3. *Elife* **7**, (2018).
705

706   37.   Zivanov J, Nakane T, Scheres SHW. A Bayesian approach to beam-induced motion
707   correction in cryo-EM single-particle analysis. *IUCrJ* **6**, 5-17 (2019).
708

709   38.   Zivanov J, Nakane T, Scheres SHW. Estimation of high-order aberrations and anisotropic
710   magnification from cryo-EM data sets in RELION-3.1. *IUCrJ* **7**, 253-267 (2020).
711

712   39.   Scheres SH. Processing of Structurally Heterogeneous Cryo-EM Data in RELION.
713   *Methods Enzymol* **579**, 125-157 (2016).
714

715   40.   Taoka M, *et al.* A mass spectrometry-based method for comprehensive quantitative
716   determination of post-transcriptional RNA modifications: the complete chemical
717   structure of Schizosaccharomyces pombe ribosomal RNAs. *Nucleic Acids Res* **43**, e115
718   (2015).
719

720   41.   Brown A, *et al.* Structure of the large ribosomal subunit from human mitochondria.
721   *Science* **346**, 718-722 (2014).
722

723   42.   Brown A, *et al.* Structures of the human mitochondrial ribosome in native states of
724   assembly. *Nat Struct Mol Biol* **24**, 866-869 (2017).
725

726   43.   Greber BJ*, et al.* The complete structure of the large subunit of the mammalian
727   mitochondrial ribosome. *Nature* **515**, 283-286 (2014).

729   44.   Suzuki T*, et al.* Complete chemical structures of human mitochondrial tRNAs. *Nat*
730   *Commun* **11**, 4269 (2020).

732   45.   Itoh Y*, et al.* Structure of the mitoribosomal small subunit with streptomycin reveals Fe-S
733   clusters and physiological molecules. *Elife* **11**, (2022).

735   46.   Itoh Y, Naschberger A, Mortezaei N, Herrmann JM, Amunts A. Analysis of translating
736   mitoribosome reveals functional characteristics of translation in mitochondria of fungi.
737   *Nat Commun* **11**, 5187 (2020).

739   47.   Peker E*, et al.* A two-step mitochondrial import pathway couples the disulfide relay with
740   matrix complex I biogenesis. *J Cell Biol* **222**, (2023).

742   48.   Watson ZL*, et al.* Structure of the bacterial ribosome at 2 A resolution. *Elife* **9**, (2020).

744   49.   Jourdain AA*, et al.* A mitochondria-specific isoform of FASTK is present in
745   mitochondrial RNA granules and regulates gene expression and function. *Cell Rep* **10**,
746   1110-1121 (2015).

748   50.   Takyar S, Hickerson RP, Noller HF. mRNA helicase activity of the ribosome. *Cell* **120**,
749   49-58 (2005).

751   51.   Khawaja A*, et al.* Distinct pre-initiation steps in human mitochondrial translation. *Nat*
752   *Commun* **11**, 2932 (2020).

754   52.   Selmer M*, et al.* Structure of the 70S ribosome complexed with mRNA and tRNA.
755   *Science* **313**, 1935-1942 (2006).

757   53.   Rozov A*, et al.* Importance of potassium ions for ribosome structure and function
758   revealed by long-wavelength X-ray diffraction. *Nat Commun* **10**, 2519 (2019).

760   54.   Bhaskar V*, et al.* Dynamics of uS19 C-Terminal Tail during the Translation Elongation
761   Cycle in Human Ribosomes. *Cell Rep* **31**, 107473 (2020).

763   55.   Ogle JM, Brodersen DE, Clemons WM, Jr., Tarry MJ, Carter AP, Ramakrishnan V.
764   Recognition of cognate transfer RNA by the 30S ribosomal subunit. *Science* **292**, 897-
765   902 (2001).

767   56.   Youngman EM, Cochella L, Brunelle JL, He S, Green R. Two distinct conformations of
768   the conserved RNA-rich decoding center of the small ribosomal subunit are recognized
769   by tRNAs and release factors. *Cold Spring Harb Symp Quant Biol* **71**, 545-549 (2006).

771   57.   Nakayama H, *et al.* Ariadne: a database search engine for identification and chemical
772   analysis of RNA using tandem mass spectrometry data. *Nucleic Acids Res* **37**, e47 (2009).
773

774   58.   Yamauchi Y*, et al.* A mass spectrometry-based method for direct determination of
775   pseudouridine in RNA. *Nucleic Acids Res* **44**, e59 (2016).
776

777   59.   Nishimura K*, et al.* Identification of the RsmG methyltransferase target as 16S rRNA
778   nucleotide G527 and characterization of Bacillus subtilis rsmG mutants. *J Bacteriol* **189**,
779   6068-6073 (2007).
780

781   60.   Noel JK, *et al.* SMOG 2: A Versatile Software Package for Generating Structure-Based
782   Models. *PLoS Comput Biol* **12**, e1004794 (2016).
783

784   61.   Whitford PC, Noel JK, Gosavi S, Schug A, Sanbonmatsu KY, Onuchic JN. An all-atom
785   structure-based potential for proteins: bridging minimal models with all-atom empirical
786   forcefields. *Proteins* **75**, 430-441 (2009).
787

788   62.   Polikanov YS, Melnikov SV, Soll D, Steitz TA. Structural insights into the role of rRNA
789   modifications in protein synthesis and ribosome assembly. *Nat Struct Mol Biol* **22**, 342-
790   344 (2015).
791

792   63.   Greber BJ*, et al.* Ribosome. The complete structure of the 55S mammalian mitochondrial
793   ribosome. *Science* **348**, 303-308 (2015).
794

795   64.   Ott M, Amunts A, Brown A. Organization and Regulation of Mitochondrial Protein
796   Synthesis. *Annu Rev Biochem* **85**, 77-101 (2016).
797

798   65.   Singh V*, et al.* Structural basis of LRPPRC-SLIRP-dependent translation by the
799   mitoribosome. *bioRxiv*, 2022.2006.2020.496763 (2022).
800

801   66.   Webster MW*, et al.* Structural basis of transcription-translation coupling and collision in
802   bacteria. *Science* **369**, 1355-1359 (2020).
803

804   67.   Petrov AS, Wood EC, Bernier CR, Norris AM, Brown A, Amunts A. Structural Patching
805   Fosters Divergence of Mitochondrial Ribosomes. *Mol Biol Evol* **36**, 207-219 (2019).
806

807   68.   Hochberg GKA, Liu Y, Marklund EG, Metzger BPH, Laganowsky A, Thornton JW. A
808   hydrophobic ratchet entrenches molecular complexes. *Nature* **588**, 503-508 (2020).
809

810   69.   Zhuang X, Cheng CH. ND6 gene "lost" and found: evolution of mitochondrial gene
811   rearrangement in Antarctic notothenioids. *Mol Biol Evol* **27**, 1391-1403 (2010).
812

813   70.   Mark FC*, et al.* Mitochondrial function in Antarctic nototheniids with ND6 translocation.
814   *PLoS One* **7**, e31860 (2012).
815

71.   Itoh Y*, et al.* Mechanism of membrane-tethered mitochondrial protein synthesis. *Science* **371**, 846-849 (2021).

72.   Ogasawara T, Ito K, Igarashi K. Effect of polyamines on globin synthesis in a rabbit reticulocyte polyamine-free protein synthetic system. *J Biochem* **105**, 164-167 (1989).

73.   Schroeder S*, et al.* Dietary spermidine improves cognitive function. *Cell Rep* **35**, 108985 (2021).

74.   Jing YH*, et al.* Spermidine ameliorates the neuronal aging by improving the mitochondrial function in vitro. *Exp Gerontol* **108**, 77-86 (2018).

75.   Phillips JE, Chaffee RR. Restorative effects of spermine on oxidative phosphorylation and respiration in heat-aged mitochondria. *Biochem Biophys Res Commun* **108**, 174-181 (1982).

76.   Al-Habsi M*, et al.* Spermidine activates mitochondrial trifunctional protein and improves antitumor immunity in mice. *Science* **378**, eabj3510 (2022).

77.   Zhang R, Ma XN, Liu K, Zhang L, Yao M. Exogenous spermine preserves mitochondrial bioenergetics via regulating Src kinase signaling in the spinal cord. *Mol Med Rep* **16**, 3619-3626 (2017).

78.   Ast T*, et al.* METTL17 is an Fe-S cluster checkpoint for mitochondrial translation. *bioRxiv*, 2022.2011.2024.517765 (2022).

79.   Cipullo M, Gese GV, Khawaja A, Hallberg BM, Rorbach J. Structural basis for late maturation steps of the human mitoribosomal large subunit. *Nat Commun* **12**, 3673 (2021).

80.   Tobiasson V, Amunts A. Ciliate mitoribosome illuminates evolutionary steps of mitochondrial translation. *Elife* **9**, (2020).

81.   Tobiasson V, Gahura O, Aibara S, Baradaran R, Zikova A, Amunts A. Interconnected assembly factors regulate the biogenesis of mitoribosomal large subunit. *EMBO J* **40**, e106292 (2021).

82.   Waltz F, Soufari H, Bochler A, Giege P, Hashem Y. Cryo-EM structure of the RNA-rich plant mitochondrial ribosome. *Nat Plants* **6**, 377-383 (2020).

83.   Sui L*, et al.* Death associated protein-3 (DAP3) and DAP3 binding cell death enhancer-1 (DELE1) in human colorectal cancer, and their impacts on clinical outcome and chemoresistance. *Int J Oncol* **62**, (2023).

84.   Silva-Pinheiro P*, et al.* A library of base editors for the precise ablation of all protein-coding genes in the mouse mitochondrial genome. *Nat Biomed Eng*, (2022).

862
863   85.   Willis JCW, Silva-Pinheiro P, Widdup L, Minczuk M, Liu DR. Compact zinc finger base
864   editors that edit mitochondrial or nuclear DNA in vitro and in vivo. *Nat Commun* **13**,
865   7204 (2022).
866

Methods

Experimental model and culturing

HEK293S-derived cells (T501) were grown in Freestyle 293 Expression Medium containing 5% tetracycline-free fetal bovine serum (FBS) in vented shaking flasks at 37°C, 5% CO_2 and 120 rpm (550 x g). Culture was scaled up sequentially, by inoculating at 1.5×10^6 cells/mL and subsequently splitting at a cell density of 3.0×10^6 cells/mL. Finally, a final volume of 2 L of cell culture at a cell density of 4.5×10^6 cells/mL was used for mitochondria isolation, as previously described [86].

mS29-KO cells were derived from HEK293T (CRL-3216) and cultured in complete DMEM supplemented with 20% FBS at 37 °C and 5% CO_2. HEK293T (CRL-3216) were cultured in high-glucose, sodium pyruvate, L-glutamine Dulbecco's modified Eagle medium (DMEM, Life Technologies) supplemented with 10% FBS, 100 µg/mL uridine, 3 mM sodium formate, and 1x MEM non-essential amino acids complete DMEM medium). Analysis of mycoplasma contamination was routinely performed.

For the labeling of cells with deuterated Uridine required for SILNAS analysis, TK6 cells deficient in uridine monophosphate synthase (*UMPS-/-*) were cultured at 37°C in RPMI-1640 medium containing 10% horse serum, 2 mM L-glutamine, penicillin (50 U/ml), streptomycin (50 µg/ml) and 0.1 mM 5,6-D2-uridine.

Generation of mS29 edited cell lines and plasmid transfections

To generate a stable human *mS29-KO* cell line in HEK293T cells (ATCC: CRL-3216), we used a pool of CRISPR/Cas9-mediated knockout cells generated by the genome editing company, Synthego (Synthego Corporation, Redwood City, CA). HEK293T cells were first tested to be negative for mycoplasma. Three guide RNAs targeting exon 5 of the *mS29* gene (DAP3-202 transcript ID ENST00000368336.10) were selected for high specificity and activity to create premature stop codons through frameshift mutations in the coding region via insertions and/or deletions (Indels). Targeting exon 5 ensured that all *mS29* transcript variants would be affected. Based on additional off-target analysis, the three specific guide RNAs selected (g1, g2, and g3) were: g1 5′-GUGAAGCUUGCCUGAUGGUA-3′ [AGG]-PAM, g2 5′-UAUAGCUGGAUAAGCAAAAC-3′ [TGG]-PAM, and g3 5′-AUGCUUUCUCCCCUCAAAUA-3′ [AGG]-PAM. The pool of knockout cells was then generated by electroporation of ribonucleoproteins (RNPs) containing the Cas9 protein and synthetic chemically modified sgRNA (Synthego) into the cells using Synthego's optimized protocol. The editing efficiency was assessed upon recovery, 48 h post-electroporation. Genomic

DNA is extracted from a portion of the cells, PCR-amplified, and sequenced using Sanger sequencing. The resulting chromatograms are processed using Synthego Inference of CRISPR Edits software (ice.synthego.com). The pooled *mS29-KO* was then plated into 96-well plates to screen for single clones in-house. Single clone candidates were screened by immunoblotting to determine the steady-state levels of mS29 and the mtDNA-encoded COX2 protein as a surrogate of mitochondrial protein synthesis capacity. Clones that had undetectable mS29 and attenuated COX2 levels were further analyzed by genotyping. For genotyping, the edited region of mS29 was amplified using oligonucleotides mS29 exon 5 seq (-164) Forward: 5′-GGATAGATTTTCAAACTCAGTACCA-3′ and mS29 exon 5 seq (+166) Reverse: 5′-TCCTGACTTCAGGCGATACG-3′, and subsequently cloned into the TOPO-TA vector (Thermo Fisher) for sequencing.

To establish a stable cell line reconstituted with mS29, we obtained the WT *mS29* gene as a Myc/DDK-tagged ORF from Origene (Cat# RC223182). Using restriction sites *Asi*SI and *Pme*I, WT mS29-Myc/DDK was cloned into a mammalian vector with hygromycin as the selection marker. The vector used was pCMV6, in which mS29 gene expression was placed under the control of an attenuated version of the human cytomegalovirus (CMV) enhancer/promoter (Δ5) in which a deletion in the promoter sequence eliminates most transcription factor binding sites [87]. Furthermore, to ensure functionality of the recombinant mS29, the Myc/DDK tag was removed via the Q5 site-directed mutagenesis kit (New England Biolabs) using the oligonucleotides mS29 ΔMyc-DDK Forward: 5′-TAAACGGCCGGCCGCGGT-3′ and mS29 Δmyc-DDK Reverse: 5′-GAGGTAGGCACAGTGCCGCTC-3′. Two μg of the construct containing *mS29* was transfected to the HEK293T *mS29*-KO cell line using 5 μL of EndoFectin™ Max (GeneCopoeia) pre-incubated in Opti-MEM (ThermoFisher). 72 h post-transfection, the medium was supplemented with 200μg/mL hygromycin B for three weeks. We use the same protocol to establish stable cell lines reconstituted with mutant variants of mS29.

Mutant variants of the *mS29*-containing construct were generated using the Q5 site-directed mutagenesis kit (New England Biolabs). Mutagenesis primers were designed using the NEBaseChanger tool. Oligonucleotides used for mutagenesis include: mS29 K134A Forward: 5′-GGGAACAGGAGCAACCCTAAGTC-3′; mS29 K134A Reverse: 5′-TTCTCTCCATACAGAAGATATC-3′; mS29 Y173A Forward: 5′-GCAGTCCAGCGCCAACAAACAGC-3′; mS29 Y173A Reverse: 5′-AGAAGATCCCGACAATTTTTC-3′; mS29 R177A Forward: 5′-CAACAAACAGGCCTTTGATCAACCTTTAGAG-3′; mS29 R177A Reverse: 5′-TAGCTGGACTGCAGAAGA-3′; mS29 K245A Forward: 5′-AATTGTGCTGGCAGAGCTAAAGAGGC-3′; mS29 K245A Reverse: 5′-CCAACTGCATCTGTGGCG-3′; mS29 H291A Forward: 5′-AGCACTTGTTGCCAACTTGAGGAAAATGATG-3′; mS29 H291A Reverse: 5′-AATTCCTCGGGGGCAATC-3′; mS29 K295A Forward: 5′-CAACTTGAGGGCAATGATGAAAAATGATTG-3′; mS29 K295A Reverse: 5′-TGAACAAGTGCTAATTCC-3′; mS29 Y208A Forward: 5'-

TCAAGAGAAGGCTGTCTGGAATAAGAG -3'; mS29 Y208A Reverse: 5'-
ACTTTTATCTGGTTCAGG -3'; mS29 V209P Forward: 5'-
AGAGAAGTATCCCTGGAATAAGAGAG -3'; mS29 V209P Reverse: 5'-
TGAACTTTTATCTGGTTCAG -3'; mS29 D238A Forward: 5'-
GAACGCCACAGCTGCAGTTGGAA -3'; mS29 D238A Reverse: 5'-
CTCACCCGTGTTATGCCC -3. All mutant sequences were validated by Sanger sequencing
using oligonucleotides Δ5pCMV6 Forward: 5′-CCTCTTCGCTATTACGCCAG-3′, mS29
cDNA (308) Forward: 5′-AACCAGCCCTAGAACTTCTGC-3′, and mS29 cDNA (829)
Forward: 5′-GAAGATAAAAGCCCGATTGC-3′.
950

<u>Mitoribosome purification</u>

Mitoribosome purification was carried out based on the previously developed protocol for rapid
isolation [86]. HEK293S-derived cells were harvested from the 2 L culture when the cell density
was 4.2 x 10^6 cells/mL by centrifugation at 1,000 g for 7 min, 4 °C. The pellet was washed and
resuspended in 200 mL Phosphate Buffered Saline (PBS). The washed cells were pelleted at
1,000 g for 10 min at 4 °C. The resulting pellet was resuspended in 120 mL of MIB buffer (50
mM HEPES-KOH, pH 7.5, 10 mM KCl, 1.5 mM $MgCl_2$, 1 mM EDTA, 1 mM EGTA, 1 mM
dithiothreitol, cOmplete EDTA-free protease inhibitor cocktail (Roche) and allowed to swell in
the buffer for 15 min in the cold room by gentle stirring. About 45 mL of SM4 buffer (840 mM
mannitol, 280 mM sucrose, 50 mM HEPES-KOH, pH 7.5, 10 mM KCl, 1.5 mM $MgCl_2$, 1 mM
EDTA, 1 mM EGTA, 1 mM DTT, 1X cOmplete EDTA-free protease inhibitor cocktail (Roche)
was added to the cells in being stirred in MIB buffer and poured into a nitrogen cavitation device
kept on ice. The cells were subjected to a pressure of 500 psi for 20 min before releasing the
nitrogen from the chamber and collecting the lysate. The lysate was clarified by centrifugation at
800 x g and 4 °C, for 15 min, to separate the cell debris and nuclei. The supernatant was passed
through a cheesecloth into a beaker kept on ice. The pellet was resuspended in half the previous
volume of MIBSM buffer (3 volumes MIB buffer + 1 volume SM4 buffer) and homogenized
with a Teflon/glass Dounce homogenizer. After clarification as described before, the resulting
lysate was pooled with the previous batch of the lysate and subjected to centrifugation at 1,000 x
g, 4 °C for 15 minutes to ensure complete removal of cell debris. The clarified and filtered
supernatant was centrifuged at 10,000 x g and 4 °C for 15 min to pellet crude mitochondria.
Crude mitochondria were resuspended in 10 mL MIBSM buffer and treated with 200 units of
Rnase-free Dnase (Sigma-Aldrich) for 20 min in the cold room to remove contaminating
genomic DNA. Crude mitochondria were again recovered by centrifugation at 10,000 g, 4 °C for
15 min and gently resuspended in 2 mL SEM buffer (250 mM sucrose, 20 mM HEPES-KOH,
pH 7.5, 1 mM EDTA). Resuspended mitochondria were subjected to a sucrose density step-
gradient (1.5 mL of 60% sucrose; 4 mL of 32% sucrose; 1.5 mL of 23% sucrose and 1.5 mL of
15% sucrose in 20 mM HEPES-KOH, pH 7.5, 1 mM EDTA) centrifugation in a Beckmann
Coulter SW40 rotor at 28,000 rpm (139,000 x g) for 60 min. Mitochondria seen as a brown band

at the interface of 32% and 60% sucrose layers were collected and snap-frozen using liquid
nitrogen and transferred to −80 °C.

Frozen mitochondria were transferred on ice and allowed to thaw slowly. Lysis buffer (25 mM
HEPES-KOH, pH 7.5, 50 mM KCl, 10 mM Mg(OAc)$_2$, 2% polyethylene glycol octylphenyl
ether, 2 mM DTT, 1 mg/mL EDTA-free protease inhibitors (Sigma-Aldrich) was added to
mitochondria and the tube was inverted several times to ensure mixing. A small Teflon/glass
Dounce homogenizer was used to homogenize mitochondria for efficient lysis. After incubation
on ice for 5-10 min, the lysate was clarified by centrifugation at 30,000 x g for 20 min, 4 °C. The
clarified lysate was carefully collected. Centrifugation was repeated to ensure complete
clarification. A volume of 1 mL of the mitochondrial lysate was applied on top of 0.4 mL of 1 M
sucrose (v/v ratio of 2.5:1) in thick-walled TLS55 tubes. Centrifugation was carried out at
231,500 x g for 45 min in a TLA120.2 rotor at 4 °C. The pellets thus obtained were washed and
sequentially resuspended in a total volume of 100 µl resuspension buffer (20 mM HEPES-KOH,
pH 7.5, 50 mM KCl, 10 mM Mg(OAc)$_2$, 1% Triton X-100, 2 mM DTT). The sample was
clarified twice by centrifugation at 18,000 g for 10 min at 4 °C. The sample was applied on to a
linear 15-30% sucrose (20 mM HEPES-KOH, pH 7.5, 50 mM KCl, 10 mM Mg(OAc)$_2$, 0.05% n-
dodecyl-β-D-maltopyranoside, 2 mM DTT) gradient and centrifuged in a TLS55 rotor at 213,600
x g for 120 min at 4 °C. The gradient was fractionated into 50 µL volume aliquots. The
absorption for each aliquot at 260 nm was measured and fractions corresponding to the
monosome peak were collected. The pooled fractions were subjected to buffer exchange with the
resuspension buffer to dilute away sucrose.

1001

<u>rRNA mass spectrometry</u>

Purification of mitochondrial rRNAs

Mitochondrial rRNAs of the HEK293 cells were extracted from the purified ribosome using
Isogen (Nippon Gene) and stored at -80 °C until use. TK6 cell culture (~1.0×10^9 cells) were
homogenized by Dounce homogenizer with 250 mM sucrose, 1 mM EDTA and 10 mM
Tris−HCl (pH 8.0). The lysate was centrifuged twice at 1500 x g for 3 min at 4 °C, and the
resulting supernatant was further centrifuged at 5000 x g for 10 min to obtain mitochondrial
precipitate. From the precipitate, total mitochondrial RNA (~100 µg) was extracted using TRIzol
Reagent (Thermo Fisher Scientific).

Each mitochondrial rRNA was purified by using reversed-phase LC through a PLRP-S 4000Å
column (4.6 × 150 mm, 10 µm, Agilent Technologies). After applying the mixture of rRNAs
from the purified ribosome or total mitochondrial RNA to the column, the rRNAs were eluted
with a 60-minute linear gradient of 11.5–13.5% (v/v) acetonitrile in 100 mM TEAA (pH 7.0) and
0.1 mM diammonium phosphate at a flow rate of 200 µl/min at 60 °C while monitoring the
eluate at A260 [88].

1017

LC-MS, MS/MS and MS/MS/MS analysis and database search of RNA fragments

RNA was digested as described below. Nucleolytic RNA fragments were analyzed with a direct nanoflow LC-MS system as described [89]. The LC eluate was sprayed online at –1.3 kV with the aid of a spray-assisting device [89] to a Q Exactive Plus mass spectrometer (Thermo Fisher Scientific) in negative ion mode. Other settings for MS, MS/MS and MS/MS/MS were as described [58].

Ariadne [57] was used for database searches and assignment of MS/MS RNA spectra. The composite of human cytosolic and mitochondrial rRNA and tRNA sequences was used as a database. The following default search parameters for Ariadne were used: maximum number of missed cleavages, 1; variable modification parameters, two methylations per RNA fragment for any residue; RNA mass tolerance, ±5 ppm, and MS/MS tolerance, ±20 ppm. For assignment of Ψ residues using 5,6-D2-uridine labeled RNAs, the mass table and the variable modification parameters were altered from default values to "5,6D_CU" and "Ψ", respectively, because both C and U were labeled with the medium and the pseudouridylation reaction results in the exchange of the position 5 deuterium of 5,6-D2-uridine to the proton of solvent, providing a −1 Da mass shift [52].

Internal standard RNAs and SILNAS-based quantitation of the stoichiometry of post-transcriptional modification

To construct the plasmids for i*n vitro* transcription of internal standard RNAs, DNAs encoding human 12S and 16S rRNAs were amplified by PCR from the mitochondrial DNA from TK6 cells. The PCR primers used are as follows: HindIII-12S-Forward TATAAAGCTTAATAGGTTTGGTCCTAGCCTTTCTATTAGC, 12S-*Bam*HI-reverse ATATGGATCCGTTCGTCCAAGTGCACTTTCCAGTACACTT, *Hind*III-16S-forward TATAAAGCTTGCTAAACCTAGCCCCAAACCCACTCCACCT and 16S-*Xho*I-reverse ATATCTCGAGAAACCCTGTTCTTGGGTGGGTGTGGGTATA). The amplified DNAs were inserted into the multiple cloning sites of plasmid pCDNA3.1 (+) (Thermo Fisher Scientific). To synthesize RNA, 2 µg of template DNA was incubated and transcribed using Megascript T7 kit (Thermo Fisher Scientific). When RNA was synthesized, guanosine-$^{13}C_{10}$ 5′-triphosphate, cytidine-$^{13}C_9$ 5′-triphosphate, or uridine-$^{13}C_9$ 5′-triphosphate solution was used instead of the respective 5′-triphosphate reagent that contained carbons with a natural isotope distribution. The RNA was precipitated in ethanol, solubilized in nuclease-free water, and purified further by reversed-phase LC as described above.

SILNAS-based quantitation was performed as described [90]. In brief, RNA (~100 fmol) from natural sources or cells grown in guanosine with natural isotope distribution was mixed with an equal amount of synthetic RNA transcribed *in vitro* with ^{13}C-labeled guanosine and digested with RNAse T1 (2 ng/µl). For the RNA transcribed with $^{13}C_9$-labeled cytidine and uridine, Rnase A (0.5 ng/µl) was used as the digestion enzyme. Digestion was carried out in 100 mM TEAA (pH 7.0) at 37°C. The 1:1 RNA mixing was performed based on the measurement of the

1057 absorbance at 260 nm and ensured later by a correction factor obtained experimentally. After
1058 obtaining the LC-MS spectrum of the digested RNA mixture, the stoichiometry of RNA
1059 modification at each site was estimated by Ariadne program designed for SILNAS [90]. The results
1060 were confirmed by manual inspection of the original MS spectrum to examine whether the
1061 estimates are based on 'uncontaminated' MS signals.

1062

Other procedures for RNA analysis

The masses of RNA fragments and a-, c-, w-, and y-series ions were calculated with Ariadne
(http://ariadne.riken.jp/). Sequence-specific Rnase H cleavage of rRNAs was performed as
described [91], using O-methylated RNA/DNA hybrid oligonucleotide complementary to the 12S
and 16S rRNAs,
GmUmUmCmGmUmCm(CAAG)UmGmCmAmCmUmUmUmCmCmAmGmUmAmCmAmC
m and (AmAmCmCmCmUmGm(TTCT)UmGmGmGmUmGmGmGmUm, respectively (where
Nm refers to 2'-O-methyl ribonucleotide and deoxyribonucleotides are indicated in parentheses).

1071

SDS-PAGE and immunoblotting analyses

Whole-cell protein extracts were obtained by solubilization in RIPA buffer (25mM Tris-HCl pH
7.6, 150 mM NaCl, 1% NP-40, 1% sodium deoxycholate, and 0.1% SDS) with 1 mM PMSF and
1x mammalian protease inhibitor. Extracts were cleared by 5 minutes of centrifugation at 10,000
g at 4 °C. Protein concentrations were determined with the Folin phenol reagent. 40-60 μg of
whole-cell protein extracts were separated by SDS-PAGE in the Laemmli buffer system.
Proteins were then transferred to nitrocellulose membranes, subsequently blocked with 5% skim
milk, and then incubated with antibodies against the indicated proteins followed by a second
reaction with anti-mouse or anti-rabbit IgG conjugated to horseradish peroxidase. Signals were
detected by chemiluminescence and exposure to X-ray films. For quantification, the signals in
digitalized images were quantified by densitometry using the histogram function of Adobe
Photoshop. Values were normalized to the signal of β-Actin and plotted as the percentage of WT
using the Prism 8 software.

1085

Pulse labeling of mitochondrial translation products

Mitochondrial protein synthesis was determined by pulse-labeling of 60-70% confluent human
wild-type (WT) HEK293T cells, *MRPS29*-KO HEK293T cells, and knockout cells stably
reconstituted with mutant *MRPS29* variants. Cultures were grown in a medium without
methionine and in the presence of 100 μg/mL emetine to inhibit cytoplasmic translation as
described [92, 93]. Cells were labeled for 15 min at 37 °C with 100 μCi of ^{35}S-methionine
(PerkinElmer Life Sciences, Boston, MA). After incubation, the medium containing ^{35}S-
methionine was removed, the cells were washed once with PBS, and then the cells were
incubated for 5 minutes in complete DMEM to allow recovery. After the short incubation, cells

were washed once more with PBS, collected by trypsinization, and whole-cell protein extracts

were obtained by solubilization with RIPA buffer. 40-50 µg of each sample was separated by

SDS-PAGE on a 17.5% polyacrylamide gel, transferred to a nitrocellulose membrane, and

exposed to X-ray film. Membranes were then probed by immunoblotting to assess the steady-

state levels of β-ACTIN as a loading control. Values were normalized to the signal of β-ACTIN

and plotted as the percentage of WT using the Prism 8 software. Potassium cyanide (KCN)-

sensitive endogenous cell respiration was measured polarographically using a Clark electrode in

the indicated cell lines. Cell respiration rates are presented as nmol of O_2 consumed per one

million cells per minute.

1104

<u>Cryo-EM data acquisition</u>

Five cryo-EM datasets of mitoribosomal complexes were collected in total using EPU 1.9

(Extended Data Table 1). First, 3 µL of ~120 nM mitoribosome was applied onto a glow-

discharged (20 mA for 30 sec) holey-carbon grid (Quantifoil R2/2, copper, mesh 300) coated

with continuous carbon (of ~3 nm thickness) and incubated for 30 sec in a controlled

environment of 100% humidity and 4 ºC. The grids were blotted for 3 sec, followed by plunge-

freezing in liquid ethane, using a Vitrobot MKIV (FEI/Thermofischer). For high resolution

mitoribosome analysis, five separate datasets were collected on FEI Titan Krios

(FEI/Thermofischer) transmission electron microscope operated at 300 keV, using C2 aperture of

70 µm and a slit width of 20 eV on a GIF quantum energy filter (Gatan). A K2 Summit detector

(Gatan) was used at a pixel size of 0.83 Å (magnification of 165,000X) with a dose of 29-32

electrons/Å^2 fractionated over 20 frames. A defocus range of -0.6 to -2.8 µm was used. More

detailed parameters for each one of the datasets are listed in Supplementary Table 1.

1118

<u>Cryo-EM data processing</u>

The complete workflows of the mitoribosomal complexes A/A-P/P-E/E, A/P-P/E, L1 stalk,

monosome consensus map, SSU and their 8 masked regions are given in Extended Data Figs. 1

and 2. For all the datasets, beam-induced motion correction and per-frame B-factor weighting

were performed for all data sets using RELION-3.0.2 [36, 37]. Motion-corrected micrographs were

used for contrast transfer function (CTF) estimation with gctf [94]. Unusable micrographs were

removed by manual inspection of the micrographs and their respective calculated CTF

parameters. Particles were picked in RELION-3.0.2 [36, 37], using reference-free followed by

reference-aided particle picking procedures. Reference-free 2D classification was carried out to

sort useful particles from falsely picked objects, which were then subjected to 3D classification.

3D classes corresponding to unaligned particles and LSU were discarded and monosome

particles were pooled and used for 3D auto-refinement yielding a map with an overall resolution

of 2.9-3.4 Å for the five datasets. Resolution was estimated using a Fourier Shell Correlation cut-

off of 0.143 between the two reconstructed half maps. Finally, the selected particles were

subjected to per-particle defocus estimation, beam-tilt correction and per-particle astigmatism

1134	correction followed by Bayesian polishing. Bayesian polished particles were subjected to a
1135	second round per-particle defocus correction. A total of 994,919 particles from all datasets were
1136	then pooled and separated into 86 optics groups in RELION-3.1 [38], based on acquisition areas
1137	and date of data collection. Beam-tilt, magnification anisotropy and higher-order (trefoil and
1138	fourth-order) aberrations were corrected in RELION-3.1 [38]. Subsequently, 509,691 particles from
1139	datasets 2-4 were pooled together and subjected to 3D auto-refinement giving a final nominal
1140	resolution of 2.21 Å. Masked refinement was performed on the small subunit head, body, mS39
1141	and tail to produce maps of 2.36 Å, 2.31 Å, 2.44 Å and 2.45 Å resolution, respectively and
1142	likewise for the SLU body, L10-L7/L12 stalk, CP and L1 stalk to yield maps of 2.08 Å, 2.38 Å,
1143	2.36 Å and 2.89 Å resolution, respectively. To improve the quality of the L1-stalk map, 220,906
1144	monosome particles with high occupancy of E/E-tRNA were pooled and subjected to partial
1145	signal subtraction using the mask that retains the E/E-tRNA and L1-stalk, followed by 3D auto-
1146	refinement to have a nominal resolution of 2.87 Å (Supplementary Fig. 1 and Supplementary
1147	Table 1).

1148	For separating the A/A-P/P-E/E and A/P-P/E states, all 994,919 monosome particles were
1149	subjected to signal subtraction using the mask that retains the tRNA binding region. The tRNA
1150	binding sites displayed weak densities corresponding to a heterogeneous mixture of tRNAs
1151	bound with partial occupancy. Signal subtracted particles were 3D classified without alignment
1152	using a mask on the A-site. This yielded two main classes, one with density in the A-site
1153	(332,187 particles) and the other lacking it (629,597 particles). Particles with density in the A-
1154	site were selected and subjected to a second round of 3D-classification using a mask around the
1155	P- and E-sites. Two classes were selected, one with P/P- and E/E-tRNA density and other in
1156	hybrid state with P/E-tRNA density. All 3D classifications with signal subtracted particles were
1157	performed using a T-value of 400. The subtracted signal was reverted for these two classes,
1158	corresponding to the A/A-P/P-E/E state (82,522 particles) and A/P-P/E state (20,143 particles).
1159	Following 3D auto-refinement, the nominal resolution of the cosmplexes was 2.63 Å and 2.98 Å,
1160	respectively. Masked refinement for the classical state was performed using local masks on SSU
1161	head, body, tail, and mS39 to produce maps at 2.72 Å, 2.75 Å, 2.91 Å and 2.93 Å resolution,
1162	respectively. For the LSU body, L10-L7/L12, CP, and the L1 stalk regions, the resolution was
1163	2.44 Å, 2.84 Å, 2.78 Å and 3.20 Å, respectively. Masked refinement for the hybrid state was
1164	performed using the same local masks as for the classical state but transformed to compensate
1165	for the structural differences. This produced maps for the SSU head, body, tail, and mS39 to
1166	produce maps of 3.03 Å, 3.07 Å, 3.39 Å and 3.47 Å resolution, respectively. For the LSU body,
1167	L10-L7/L12 region, CP, and the L1 stalk masks, the resolution was 2.84 Å, 3.29 Å, 3.25 Å and
1168	3.53 Å, respectively. The maps were then subjected to modulation transfer function correction,
1169	automatic *B*-factor sharpening and local resolution filtering using RELION-3.1 (Supplementary
1170	Fig. 2, Supplementary Table 1).

1171

1172

 Model building and refinement

We started by building a model into the 2.21 Å resolution consensus map. The starting model for
the LSU was taken from PDB ID: 6ZSG, and for the SSU from PDB ID: 6RW4. These models
were rigid body fitted using UCSF Chimera [95]. Individual local-masked refined maps with local-
resolution filtering and *B*-factor sharpening (RELION 3.1) superposed to the overall map were
combined into a single composite map using Phenix.combine_focused_maps [96] for model
building and refinement. *Coot* v0.9 [97] with Ramachandran and torsion restraints was used for
model building.

The L1-stalk model was built using the 2.87 Å resolution masked-refined map of L1 stalk from
E/E-tRNA-containing particles. Initial model for uL1m was generated by homology-modeling
with uL1 structure of *T. thermophilus* (PDB ID: 3U4M) as template using the Swiss-Model
webserver [98]. The N-terminal extension of uL1m (residues 61–93), the C-terminal domain
(residues 158–254) and the preceding loop (residues 148–157) of uL9m, and H76-77 (residues
2761–2786) of 16S rRNA were modeled manually into the density. Mitochondria-specific C-
terminal extension of uL9m was modeled as a helix based on the density and secondary structure
prediction by PSIPRED [99]. The model was rigid body fitted into the respective local-masked
refined maps for L1 stalk region in the A/A P/P E/E and A/P/E states followed by self-restrained
real space refinement in *Coot* v0.9.

In the CP, the tRNA^{Val} could be completely modeled, owing to the improved resolution of the
local masked-refined map (2.36 Å). The 3′-CCA terminus (C74–A76) and a valine residue linked
to A76 could be placed on the density map. The N-terminus of mL40 was extended by 13-
residues (47–59) modeled in a tubular wedged between the A- and P-site tRNAs to the phosphate
backbone of mRNA residue 7. Similarly, the C-terminus of bL31m was extended by 31 residues
(95–125) from the C-terminal helix onto SSU head. The local resolution was sufficient for
accurate side-chain modeling for most residues and secondary-structure assignments. We
modeled a disulfide linkage between C45-C76 of mS37. However, they might exist in both
disulfide-linked, and reduced states as indicated by the density and a water mediated hydrogen
bond with a neighbouring Arginine side chain (R240 from uS7m). The other pair C55-C66
appears to be reduced. However, as we had added 1mM DTT to our ribosome purification
buffers, this is not truly indicative of the native state of C45-C76 or C55-C66 pairs.

At the mRNA channel entrance, a more accurate and complete model of mS39 could be built
with 29 residues added to the structure. Improved local resolution enabled unambiguous
assignment of residues to the density which allowed us to address errors in the previous model.
Finally, a total of 28 α-helices could be modeled in their correct register and orientation. Further,
a 28-residue long N-terminal loop of mS31 (residues 247–275) along mS39 and mitochondria-
specific N-terminal extension of uS9m (residues 53–70) approaching mRNA were modeled
manually by fitting the loops into the density maps.

Polyamines were identified upon a further manual inspection of the density map for the presence
of continuous tubular densities which corresponded to spermine, spermidine or putrescine in

length. A total of one spermine, four spermidines and one putrescine were placed in the model,
justified by the electrostatic and hydrogen-bonding interactions that each polyamine exhibits at
its amine groups with its neighbors.

Metal ions, Mg^{2+}, K^+ and Zn^{2+}, were modeled into densities based on a manual inspection of
small density blobs and the chemical environment, specifically, coordination geometry and
strength of density in both unsharpened and local resolution filtered maps. The coordination
distance of Mg^{2+} is around 2 Å with an octahedral geometry, whereas that of K^+ is around 3 Å
with more relaxed constraints on coordination geometry. Notably, a density that was modeled
previously as Zn^{2+} coordinated by residues D224, D240, D241 and H93 of uS2m was assigned as
a magnesium ion, indicated by the octahedral geometry of coordination with the uS2m residues
mentioned above and two water molecules.

A-, P- and E-site tRNAs, and mRNA were modeled manually in the density. The density for
tRNAs and mRNAs comes from a natively-bound heterogeneous population. Hence, individual
residues were modeled as one of the nucleotides, A, U, G, or C, based on the density and/or
conservation. While the 70 residues of P-site tRNA could be modeled, fewer residues could be
modeled for the A-site (22 residues) and E-site tRNAs (43 residues), including the acceptor arm
and anticodon loop due to weaker respective densities. Along the mRNA channel, 34 residues of
mRNA were built. In the tRNA binding region, codon-anticodon pairs were clearly resolved.
Additionally, extra residues near the A-site (10-12, 14-16) and at the mRNA channel entrance
(25, 27-29) were modeled accurately into density. For the remaining mRNA chain at a relatively
lower resolution, correct overall chain trace and nucleotide orientation could be assigned. This
allowed specific protein-nucleotide interactions to be identified for uS5m, uS7m, uS9m, uS12m,
mS35, mS39 and mL40.

For modeling the classical A/A P/P E/E- and hybrid A/P P/E-states, the LSU and SSU from the
consensus model were fitted into their composite maps, followed by manual revision. As
described above, tRNAs and mRNA were modeled manually into the density.

The water molecules were automatically picked by *Coot* v0.9 [62], followed by manual revision.
Geometrical restraints of modified residues and ligands used for the refinement were calculated
by Grade Web Server (http://grade.globalphasing.org) or obtained from CCP4 library [100].
Hydrogens were added to the models except for water molecules by REFMAC5 [101] using the
prepared geometrical restraint files. Charged α-amino-group hydrogens (H2 and H3) and a free-
ribose hydrogen HO3′ were removed from non-terminal protein and RNA residues, respectively,
while those at terminal residues were kept. Protonation of Histidine residues were judged based
on the chemical environment and removed or kept their N-H hydrogens (HD1 and HE2)
accordingly. Protonation/deprotonation of ligands and modified residues were also adjusted.

Final models were further subjected to refinement with Phenix.real_space_refine v1.13_2998 [96],
wherein three macro-cycles of global energy minimization with reference restraints (using the
input model as the reference, sigma 5-7) and secondary structure restraints, rotamer restraints but
without Ramachandran restraints were carried out for each model. The composite maps were

used as the targets for the refinement. An 'edit' file that defines metal-coordination bonds was
generated by ReadySet in the Phenix suite and used. Non-canonical covalent bonds, which are
those between tRNA^{Val} and Valine and between 1-methyl-isoaspartate 184 and G185 in uS11m,
were also defined in the 'edit' file. Model refinement data are listed in Supplementary Table 1.

1255

<u>Bioinformatic analysis of the Y-N-C-Y motif</u>

The density and base-specific interactions of mRNA residues 25-28 with mS39 imply a sequence
of pyrimidine, any nucleotide, cytosine, pyrimidine, leading to a consensus motif: Y-N-C-Y. We
explored the possibility of increased frequency of this motif in mRNA sequences. The sequences
for all 13 mRNAs were obtained from GenBank. The consensus motif was searched in each
sequence and all occurrences (including ones that overlap) were recorded and used to calculate
the percentage frequency of occurrence for each transcript and the average percentage frequency.
To determine if there is an enrichment of this motif in mRNAs, we calculated its occurrence in a
set of nuclear-encoded mRNAs. To create this set, we acquired sequences randomly from
GenBank and subjected them to a 90% redundancy cut-off, resulting in a total of 1369
sequences. The percentage frequency of occurrence of Y-N-C-Y motif was determined for each
of the 1370 sequences and used to calculate an average percentage frequency for the entire set.
Next to determine the distribution of the Y-N-C-Y motif across the length of mRNAs, each
occurrence was plotted against its position number in the mRNA sequence. This distribution for
all 13 mRNA sequences was plotted together for comparison.

1271

<u>Molecular Dynamics simulations: Modified force fields for simulating a truncated mL64 protein</u>
<u>and modified interactions with protein uL1m</u>

Baseline force field used for simulations of P/E formation

An all-atom multi-basin structure-based "SMOG" model [60, 61] of the mitoribosome was used to
simulate P/E formation. For this, we first constructed single-basin structure-based models for the
pre- and post-translocation configurations (A/A-P/P and P/P-E/E). We then combined the two
models, such that the composite model stabilizes both the pre- and post-translocation
configurations. Here we describe the baseline model, which is described in the main text. In
subsequent sections, we provide detailed descriptions of all modified force fields.

In a single-basin all-atom structure-based model, every non-hydrogen atom is represented by a
bead of unit mass, and an experimentally-obtained structure is used to define the potential energy
minimum. The functional form of the potential is given by:

$$U = \sum_{\text{bonds}} \frac{\epsilon_r}{2}(r_i - r_{i,0})^2 + \sum_{\text{angles}} \frac{\epsilon_\theta}{2}(\theta_i - \theta_{i,0})^2$$

$$+ \sum_{\text{impropers}} \frac{\epsilon_{\chi_{imp}}}{2}(\chi_i - \chi_{i,0})^2 + \sum_{\text{planars}} \epsilon_{\chi_{planar}}[1 - \cos(2\chi_i)]$$

$$+ \sum_{\substack{\text{backbone} \\ \text{dihedrals}}} \epsilon_{bb}F(\phi_i - \phi_{i,0}) + \sum_{\substack{\text{sidechain} \\ \text{dihedrals}}} \epsilon_{sc}F(\phi_i - \phi_{i,0})$$

$$+ \sum_{\text{contacts}} \epsilon_c \left[\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{12} - 2\left(\frac{\sigma_{ij}}{r_{ij}}\right)^6 \right] + \sum_{\text{non-contacts}} \epsilon_{nc}\left(\frac{\sigma_{nc}}{r_{ij}}\right)^{12}$$

where

$$F(\phi) = [1 - \cos(\phi)] + \frac{1}{2}[1 - \cos(3\phi)]$$

r_0 and θ_0 parameters are assigned values found in the Amber ff03 force field [102]. The dihedral
parameters ϕ_0 are defined by the experimental structure. χ_0 non-planar improper dihedrals
(around chiral centers) are also defined to have the values adopted in the experimental structures.
Planar dihedral angles are given a cosine term with periodicity 2 and minima at 0 and 180
degrees. The energy scale is set to $\epsilon_r = 100\frac{\epsilon}{A^2}, \epsilon_\theta = 80\frac{\epsilon}{rad^2}, \epsilon_{\chi_{imp}} = 10\frac{\epsilon}{rad^2}, \epsilon_{\chi_{planar}} =$
$40\frac{\epsilon}{rad^2}$, where ϵ is the reduced energy scale. The dihedral and contact energy scales $\epsilon_{bb}, \epsilon_{sc}, \epsilon_C$
are normalized as in Ref. [60]. This combination of structure-based non-bonded terms and AMBER
bonded terms is identical to the implementation of a previous SMOG-amber variant (called
SBM_AA-amber-bonds [103] in the SMOG 2 force field repository (smog-server.org), though post-
transcriptionally-modified residues were added for the current model. Upon publication, the
SMOG 2 force field templates used in the current study will be available for download on the
SMOG 2 force field repository (ID: AA_PTM_Hassan21.v1).

For the non-bonded interactions, each atomic contact that is found in the experimental structure
(i.e. "native" contact) is given an attractive Lennard-Jones-like interaction that stabilizes a
preassigned structure. Native contacts were defined based on the Shadow Contact Map algorithm
with default parameters [104]. Σ_{ij} are the interatomic distances between contacts in the native
structure, multiplied by 0.96. As employed previously in simulations of the ribosome [105, 106] and
viral capsids [103], this scaling of the contact distance was employed to avoid artificial expansion
of the ribosome that results from thermal energy. That is, while the potential energy is defined
such that the experimental structure is the global minimum, thermal energy leads to a free-energy
minimum for which the ribosome is slightly expanded. By shortening the stabilizing interactions,
the free-energy minimum is consistent with the experimental structure. For this comparison, the
radius of gyration (R_g) was used as a measure of expansion. With this scaling of the contact

distances, R_g was found to match that of the cryo-EM model. All non-native contacts are given a
repulsive term that ensures excluded-volume steric interactions are present. The parameter σ_{nc} is
given the value 2.5 Å, while $\epsilon_{nc} = 0.1\epsilon$.

Stabilizing intra-ribosome contacts and dihedrals were assigned the values found in the A/P-P/E
(SSU rotated) hybrid structure, which is described in the main text. This ensures that the SSU
rotates forward in each simulation. Each simulation is initiated from the A/A-P/P conformation.
All intra-mt-tRNA and intra-mt-mRNA contacts were defined based on the A/A-P/P structure.
Since stabilizing contacts in a structure-based model describe effective interactions [107, 108], we
rescaled the strengths of subsets of contacts within the model. Consistent with recent simulations
of P/E formation in bacteria [105], interactions that stabilize the structure of the ribosome (i.e. intra-
ribosomal) are given larger weights than contacts that are only transiently-formed (e.g. tRNA-
ribosome contacts). Relative to intra-ribosome contacts, the energetic interactions between the
mt-mRNA-tRNA complex and the SSU (in A/A-P/P and P/P-E/E structures) were rescaled by a
factor of 0.3. While these interactions were weaker than intra-ribosomal contacts, since the
simulations were not performed under conditions in which translocation will occur (*e.g.*, binding
of elongation factors or back-rotation, etc), the mt-mRNA maintained its initial position on the
mRNA track of the SSU. P/P-E/E contacts between the mt-tRNA and the uL1m protein were
rescaled by a factor of 0.8 and all other interactions between mt-mRNA-tRNA and the LSU were
rescaled by a factor of 0.1. This rescaling was applied to account for the transient nature of mt-
mRNA-tRNA binding to the ribosome. The strength of the uL1m-tRNA contacts was set such
that the P/E configuration would stably form contacts with the uL1m protein. When contacts
were scaled below a factor of ~0.5, we found that uL1m could not form stable interactions with
the mt-tRNA.

Interactions between the 3′-CCA ends of A-site and P-site mt-tRNAs with the A-site and the P-
site were scaled by a factor of 0.2, while their interactions with P-site and E-site were scaled by a
factor of 0.8. This leads to the mt-tRNA molecules favoring binding of their target sites (*i.e.*,
sites bound in the P/E configuration) on the ribosome. While the interactions with the subsequent
binding site (*i.e.*, A-site mt-tRNA with the P-site and P-site mt-tRNA with the E-site) were
stronger, these contacts are short range (~6 Å). Accordingly, the stronger weights do not affect
the dynamics of the mt-tRNA until it is very close to the binding site. Thus, the details of the
final repositioning of the 3′-CCA tail of the P-site mt-tRNA should not be overinterpreted. While
the contacts are short-range, the structure of the mitoribosome may introduce steric barriers that
the mt-tRNA must navigate in order to adopt the P/E configuration [108]. With these considerations
in mind, the current model is aimed at identifying structural elements that can impose steric
limitations on mt-tRNA during P/E formation.

There were also two sets of harmonic potentials that were introduced. To mimic the presence of
a nascent peptide chain, which would restrain the 3′-CCA end of the A-site tRNA to the PTC,
we introduced weak harmonic restraints (weight of $1/nm^2$) between the 3′-CCA end of the A-site
tRNA and the P site of the LSU. Distances were assigned based on the P/P conformation. These
weak interactions ensured that the 3′-CCA end remains localized to the PTC. We also

1347	introduced harmonic potentials between base-pairing contacts in the mt-tRNA acceptor arm
1348	(weight $1000/nm^2$). Due to the details of the contact map, initial simulations exhibited partial
1349	unfolding/refolding of the acceptor arm during P/E formation. Since that was likely an artifact
1350	due to the simplicity of the model and the details of the contact map definition, these additional
1351	harmonic terms were introduced to ensure the tRNA acceptor arm remained folded throughout
1352	the transition.
1353	
1354	*Modified force fields*
1355	Multiple variants of the model (*i.e.*, force field) were considered, in order to investigate the
1356	influence of steric interactions between the P-site mt-tRNA and C-terminal residues of mL64 and
1357	to investigate the influence of attractive interactions between uL1m and P-site mt-tRNA on P/E
1358	formation.
1359	
1360	*Truncated-mL64 model: ΔmL64 force field*
1361	The C-terminal region of mL64 is involved in steric interactions with the mt-tRNA. For ΔmL64
1362	model, the C-terminal region of mL64 (beginning with Q112) was removed from the system.
1363	Q112 was chosen as truncating here guarantees no steric interactions between mL64 and the mt-
1364	tRNA and keeps the rest of the protein intact. Thus, all other interactions were unperturbed. This
1365	model is intended to mimic an ideal experiment, where one could delete the terminal region of
1366	mL64 without introducing indirect effects on the dynamics. In our application, this model allows
1367	us to ask what effect steric interactions between P-site mt-tRNA elbow and mL64 have on the
1368	dynamics of P/E formation. Deletion of this region from the model was achieved using the
1369	smog_extract tool of the SMOG 2 software package [60].
1370	
1371	*Modified-uL1m model: ΔuL1m force field*
1372	All stabilizing contacts between uL1m and the P-site mt-tRNA that are formed in the P/P-E/E
1373	state were scaled by a relative factor of 0.1, rather than 0.8 (baseline model, above). Thus, by
1374	weakening the attractive interactions, there is insufficient stabilizing energy between uL1m and
1375	the P/E tRNA, such that the tRNA elbow does not fully reach the P/E configuration.
1376	
1377	*Combined truncated mL64 and modified uL1m model: ΔmL64-ΔuL1m force field*
1378	All modifications applied in the truncated mL64 model and the modified-uL1m model were
1379	included in a single combined force field.
1380	
1381	*Simulation details*
1382	All simulations were initiated from the pre-translocation A/A-P/P configuration. All force field
1383	files were generated using the SMOG 2 software package [60]. Molecular dynamics simulations

1384	were performed using Gromacs (v5.1.4) [109, 110]. Reduced units were used in all calculations.
1385	Energy minimization was first performed using steepest descent minimization. The system was
1386	then coupled to a heat bath using Langevin Dynamics protocols, with reference temperature of
1387	$0.5\ \varepsilon/k_B$ reduced temperature units (60 Gromacs units). This temperature was chosen since it has
1388	been shown to produce structural fluctuations that are consistent with those inferred from
1389	anisotropic B-factors and explicit-solvent simulation [111]. The timestep was 0.002 reduced units
1390	and each simulation was continued until the distance of the 3'-CCA end from the LSU E-site
1391	R_{CCA} reached a value less than 4 Å. Energy minimization was initially performed using steepest
1392	descent minimization. Each simulation was performed for a maximum of 10^8 timesteps. For the
1393	primary/baseline model described in the main text (327 simulations) we obtained an aggregate
1394	simulated time of $2.27 * 10^6$ reduced units. For the truncated mL64 model (ΔmL64 model) (321
1395	simulations), the aggregate simulation time was $1.87 * 10^6$ reduced units. For the attenuated uL1m
1396	contacts model (ΔuL1m model) (237 simulations), the aggregate simulation time was $1.76 * 10^6$
1397	reduced units. For the ΔmL64-ΔuL1m model (181 simulations), the aggregate simulation time
1398	was $1.04 * 10^6$. Previous comparisons of diffusion coefficients of tRNA molecules in the
1399	ribosome (explicit-solvent simulations vs. SMOG models), estimate that 1 reduced unit
1400	corresponds to an effective timescale of approximately 1 nanosecond [112]. Using this conversion
1401	factor, the presented simulations represent an aggregate effective simulate time of ~ 6.94
1402	milliseconds.

Structural metrics for describing simulated dynamics

- A-site mt-tRNA, R_{CCA}^{P}: distance between the geometric centers of the side chains of C75 in the A-site mt-tRNA and Gm2815 in the LSU rRNA. These residues base pair in the A/P-P/E and P/P-E/E configurations.

- P-site mt-tRNA, R_{CCA}^{P}: distance between the geometric centers of the side chains of C75 in the P-site mt-tRNA and Gm2815 in the LSU rRNA. These residues base pair in the A/P-P/E configuration.

- R_{CCA}: distance between the geometric centers of the side chains of A76 in the P-site mt-tRNA and G2909 in the LSU rRNA. G2909 in the E-site forms a base stacking interaction with A76 of P-site mt-tRNA in the A/P-P/E configuration.

- R_{elbow}: distance between geometric centers of U54 of P-site mt-tRNA and its reference position in the A/P-P/E configuration after alignment of the LSU rRNA.

- $Q_{L1-tRNA}$: the fraction of P/P-E/E contacts between uL1m and P-site mt-tRNA. A contact is defined as formed if it is within a factor of 1.2 of the distance found in the P/P-E/E configuration.

SSU rotation and tilt angles

To follow the rotary motions of the SSU body with respect to the LSU, we calculated rotation and tilt angles for both the SSU body. For this, we first determined the corresponding *E. coli* numbering of the mitoribosome rRNA residues via STAMP alignment [113] of the mitoribosome to

1422   a reference *E. coli* structure (PDB ID: 4V9D). Alignment was performed separately for the LSU,
1423   SSU head and SSU body. The angles were then calculated, using the assigned *E. coli* numbering.

1424

Structural model preparation for simulations

Since the cryo-EM models did not resolve the exact same set of atoms, there was a need to
perform some additional structural models prior to performing simulations. For example, to
define the multi-basin structure-based model, it was necessary for each mt-tRNA molecule to
have identical atoms in the pre- and post-translocation configurations. Additionally, to define
the intra-ribosome interactions to stabilize the A/P-P/E conformation, it was necessary that all
mt-rRNA and mitoribosomal proteins had identical atomic composition. While most of the
resolved atoms are common to all reported models, we employed restraint-based modeling to
construct models of the mitoribosome and mt-tRNA that have identical compositions. Below, we
describe the steps required for modeling the mitoribosome and mt-tRNA.

1435

Structural modeling pre-processing steps for rRNA and proteins for simulations

Since the structure-based model only included intra-ribosome interactions from the A/P-P/E
structure, we first removed all mt-tRNA and mt-mRNA from the A/A-P/P and A/P-P/E models.
We also removed ligands and water molecules (HOH, SPD, PUT, NAD, SPM). For the
remaining atoms, we identified all atoms that were common to the A/A-P/P and A/P-P/E
structures. We then generated a single-basin structure-based model of the A/A-P/P structure.
Next, position restraints (harmonic terms, weight of 100 for each atom) were introduced for all
common atoms, with positions corresponding to the A/P-P/E structure. The system was then
energy minimized via steepest descent minimization. Then, the system was subject to simulated
annealing. The system was initially simulated at 0.15 reduced temperature units for 25,000
timesteps. The temperature was then linearly decreased to 0 over the next 50,000 timesteps. An
additional 25,000 steps were simulated at temperature 0. The final, annealed structural model
was used for force field generation steps, as described above.

1449

Structural modeling pre-processing steps for mt-tRNA and mt-mRNA

Since the cryo-EM reconstructions reported consensus/average sequences for each mt-tRNA, the
composition of the A-, P- and E-site mt-tRNAs were not identical. To perform simulations, we
homogenized the composition of the mt-tRNA molecules in the A/A-P/P-E/E structure and the
mt-tRNA codons. For this, we started with the structural model of the A-site mt-tRNA (and the
associated codon) from the A/A-P/P structure, and then used it to generate a structure-based
model. We then introduced position restraints based on the positions of the A-site mt-tRNA in
the A/A-P/P-E/E structure. All common atoms (*i.e.*, backbone atoms, or atoms in identical
residues) were restrained to the A-site mt-tRNA/mRNA positions. For the A-site mt-tRNA,
restraints were not imposed on residues 46-58 and 16-17. Cycles of minimization and annealing

1460	were performed, where the position restraints were given weights of 1, 10, 100 and then 1000.
1461	This process was repeated for the P-site and E-site mt-tRNAs, as well as for the upstream and
1462	downstream mt-mRNA. This process produced a model of the A/A-P/P-E/E configuration that
1463	has repeating codons and identical mt-tRNA composition. This structural model was used to
1464	define the pre-translocation and post-translocation structure-based models, which were used to
1465	construct the multi-basin model.

Describing the sequence of rearrangements in simulations

To describe the dynamics of the P-site mt-tRNA during P/E state formation, we have defined three states based on the 2D probability distribution of R^E_{elbow} and R^E_{CCA} distances (Fig. 8). These states are defined as all configurations of the P-site mt-tRNA in the baseline model that sample the following values of R^E_{elbow} & R^E_{CCA}:

state I_1: disk centered at (R^E_{elbow}, R^E_{CCA})=(46,35) Å with radius r=6 Å

state I_2: disk centered at (R^E_{elbow}, R^E_{CCA})=(63,23) Å with radius r=8 Å

state I_3: disk centered at (R^E_{elbow}, R^E_{CCA})=(42,13) Å with radius r=7 Å

Using this definition of the intermediate states, the most common sequence of transitions followed the order P/P => I_1 => I_2 => I_3 => P/E.

Structure analysis and Fig.s

Visualization and analysis of the models and maps was carried out using UCSF Chimera and ChimeraX [95, 114]. For model and map comparisons, models were superposed in *Coot* v0.9 [62] using the Secondary Structure Matching algorithm and maps downloaded from EMDB were resampled on our maps in UCSF Chimera. Simulated snapshots were visualized using Visual Molecular Dynamics [115].

Quantitation and statistical analysis

All experiments were done in at least biological triplicate unless otherwise stated. GraphPad Prism 8 software was used for statistical analyses of the data. One-way analysis of variance (ANOVA) was performed, comparing the mean of each cell line with the mean of the mS29-KO cell line reconstituted with WT mS29, followed by Dunnett's multiple comparisons test. The data is presented as the mean ± SD; * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$. For immunoblotting, band density is quantified and normalized to ACTIN. For metabolic labeling of mitochondrially translated peptides in whole cells, the data presented represents the mean of each sample's ^{35}S signals.

Additional References

86.   Aibara S, Andrell J, Singh V, Amunts A. Rapid Isolation of the Mitoribosome from HEK Cells. *J Vis Exp*, (2018).

1498

1499   87.   Morita E, Arii J, Christensen D, Votteler J, Sundquist WI. Attenuated protein expression
1500   vectors for use in siRNA rescue experiments. *Biotechniques* **0**, 1-5 (2012).

1501

1502   88.   Yamauchi Y*, et al.* Denaturing reversed phase liquid chromatographic separation of non-
1503   coding ribonucleic acids on macro-porous polystyrene-divinylbenzene resins. *J*
1504   *Chromatogr A* **1312**, 87-92 (2013).

1505

1506   89.   Nakayama H, Yamauchi Y, Taoka M, Isobe T. Direct identification of human cellular
1507   microRNAs by nanoflow liquid chromatography-high-resolution tandem mass
1508   spectrometry and database searching. *Anal Chem* **87**, 2884-2891 (2015).

1509

1510   90.   Taoka M*, et al.* Landscape of the complete RNA chemical modifications in the human
1511   80S ribosome. *Nucleic Acids Res* **46**, 9289-9298 (2018).

1512

1513   91.   Taoka M*, et al.* The complete chemical structure of Saccharomyces cerevisiae rRNA:
1514   partial pseudouridylation of U2345 in 25S rRNA by snoRNA snR9. *Nucleic Acids Res*
1515   **44**, 8951-8961 (2016).

1516

1517   92.   Bourens M, Barrientos A. A CMC1-knockout reveals translation-independent control of
1518   human mitochondrial complex IV biogenesis. *EMBO Rep* **18**, 477-494 (2017).

1519

1520   93.   Leary SC, Sasarman F. Oxidative phosphorylation: synthesis of mitochondrially encoded
1521   proteins and assembly of individual structural subunits into functional holoenzyme
1522   complexes. *Methods Mol Biol* **554**, 143-162 (2009).

1523

1524   94.   Zhang K. Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1-12
1525   (2016).

1526

1527   95.   Pettersen EF*, et al.* UCSF Chimera--a visualization system for exploratory research and
1528   analysis. *J Comput Chem* **25**, 1605-1612 (2004).

1529

1530   96.   Afonine PV*, et al.* Real-space refinement in PHENIX for cryo-EM and crystallography.
1531   *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).

1532

1533   97.   Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. *Acta*
1534   *Crystallogr D Biol Crystallogr* **60**, 2126-2132 (2004).

1535

1536   98.   Waterhouse A*, et al.* SWISS-MODEL: homology modelling of protein structures and
1537   complexes. *Nucleic Acids Res* **46**, W296-W303 (2018).

1538

1539   99.   Buchan DWA, Jones DT. The PSIPRED Protein Analysis Workbench: 20 years on.
1540   *Nucleic Acids Res* **47**, W402-W407 (2019).

1541

1542 100. Krissinel EB, *et al.* The new CCP4 Coordinate Library as a toolkit for the design of
1543   coordinate-related applications in protein crystallography. *Acta Crystallogr D Biol*
1544   *Crystallogr* **60**, 2250-2255 (2004).
1545
1546 101. Murshudov GN, *et al.* REFMAC5 for the refinement of macromolecular crystal
1547   structures. *Acta Crystallogr D Biol Crystallogr* **67**, 355-367 (2011).
1548
1549 102. Duan Y, *et al.* A point-charge force field for molecular mechanics simulations of proteins
1550   based on condensed-phase quantum mechanical calculations. *J Comput Chem* **24**, 1999-
1551   2012 (2003).
1552
1553 103. Whitford PC, Jiang W, Serwer P. Simulations of Phage T7 Capsid Expansion Reveal the
1554   Role of Molecular Sterics on Dynamics. *Viruses* **12**, (2020).
1555
1556 104. Noel JK, Whitford PC, Onuchic JN. The shadow map: a general contact definition for
1557   capturing the dynamics of biomolecular folding and function. *J Phys Chem B* **116**, 8692-
1558   8702 (2012).
1559
1560 105. Levi M, Walak K, Wang A, Mohanty U, Whitford PC. A steric gate controls P/E hybrid-
1561   state formation of tRNA on the ribosome. *Nat Commun* **11**, 5706 (2020).
1562
1563 106. Nguyen K, Whitford PC. Steric interactions lead to collective tilting motion in the
1564   ribosome during mRNA-tRNA translocation. *Nat Commun* **7**, 10586 (2016).
1565
1566 107. Chan HS, Zhang Z, Wallin S, Liu Z. Cooperativity, local-nonlocal coupling, and
1567   nonnative interactions: principles of protein folding from coarse-grained models. *Annu*
1568   *Rev Phys Chem* **62**, 301-326 (2011).
1569
1570 108. Levi M, Noel JK, Whitford PC. Studying ribosome dynamics with simplified models.
1571   *Methods* **162-163**, 128-140 (2019).
1572
1573 109. Hess B, Kutzner C, van der Spoel D, Lindahl E. GROMACS 4: Algorithms for Highly
1574   Efficient, Load-Balanced, and Scalable Molecular Simulation. *J Chem Theory Comput* **4**,
1575   435-447 (2008).
1576
1577 110. Lindahl E, Hess B, van der Spoel D. GROMACS 3.0: a package for molecular simulation
1578   and trajectory analysis. *Journal of Molecular Modeling* **7**, 306-317 (2001).
1579
1580 111. Jackson J, Nguyen K, Whitford PC. Exploring the balance between folding and
1581   functional dynamics in proteins and RNA. *Int J Mol Sci* **16**, 6868-6889 (2015).
1582
1583 112. Yang H, Bandarkar P, Horne R, Leite VBP, Chahine J, Whitford PC. Diffusion of tRNA
1584   inside the ribosome is position-dependent. *J Chem Phys* **151**, 085102 (2019).
1585

1586 113. Russell RB, Barton GJ. Multiple protein sequence alignment from tertiary structure
1587   comparison: assignment of global and residue confidence levels. *Proteins* **14**, 309-323
1588   (1992).

1589
1590 114. Pettersen EF*, et al.* UCSF ChimeraX: Structure visualization for researchers, educators,
1591   and developers. *Protein Sci* **30**, 70-82 (2021).

1592
1593 115. Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. *J Mol Graph* **14**,
1594   33-38, 27-38 (1996).

1595
1596

Data availability statement

The atomic coordinates were deposited in the RCSB Protein Data Bank, and EM maps have been deposited in the Electron Microscopy Data bank under accession numbers: 7QI4 and EMD-13980 (consensus monosome), 7QI5 and EMD-13981(classical state), 7QI6 and EMD-13982 (hybrid state).

The atomic coordinates that were used in this study: 3U4M (uL1 from *T. thermophilus*), 4V51 (*T. thermophilus* ribosome with mRNA, tRNA), 4V9D (*E. coli* ribosome), 4Y4P (*T. thermophilus* ribosome with A-, P-, E-site tRNAs), 6QNR (*T. thermophilus* ribosome with experimentally assigned K^+ ions) 6RW4 (mtSSU with mtIF3), 6ZM5 (mitoribosome from actinonin-treated cells) 6ZSG (mitoribosome with mRNA, tRNA), 6ZTJ (*E.coli* expressome), 6ZTN (*E.coli* expressome), 7K00 (*E. coli* ribosome at 2 Å resolution).

Acknowledgments

We thank the EMBO Young Investigator Program, the Northeastern University Discovery cluster and Northeastern University Research Computing staff. Swedish Foundation for Strategic Research (FFL15:0325), Ragnar Söderberg Foundation (M44/16), European Research Council (ERC-2018-StG-805230), Knut and Alice Wallenberg Foundation (2018.0080), Japan Science and Technology Agency (19198049), NIH-R35 grant (GM118141 to A.B.), and MDA Research Grant (MDA-381828 to A.B.). V.S. is supported by the Horizon 2020 - Marie Sklodowska-Curie Innovative Training Network (721757). Y.I. was supported by H2020-MSCA-IF-2017 (799399-Itohribo). The SciLifeLab cryo-EM facility is funded by the Knut and Alice Wallenberg, Family Erling Persson, and Kempe foundations. P.C.W. was supported by NSF grant MCB-1915843. Work at the Center for Theoretical Biological Physics was also supported by the NSF (Grant PHY-2019745). We would like to thank AMD for the donation of critical hardware and support resources from its HPC Fund that made this work possible.

Author contributions

1625 V.S., Y.I., S.A., and J.A. collected cryo-EM data. V.S. and Y.I. processed the data and built the
1626 models. V.S., Y.I., R.K.F., A.N., S.A. and A.A. carried out structural analysis. Y.N., K.I. and
1627 M.T. performed mass-spec characterization. S.L.D. and A.B. performed biochemical
1628 characterization. A.H. and P.C.W. performed molecular simulations. All authors contributed to
1629 data interpretation and manuscript writing.

1630

1631 **Competing Interests Statement**

1632 The authors declare no competing interests.

SUPPLEMENTARY INFORMATION

Table of contents:

Supplementary Fig. 1: Cryo-EM processing overview.
Data processing scheme for classical (A/A P/P E/E) and hybrid (A/P P/E) states. Bottom panel,
local-masked refinements, resolution and model validation for the L1 stalk. A total of 43686 movies
were recorded and analyzed, particles that were not mitoribosomes were discarded by classification,
since they cannot contribute to reconstruction. Consensus map was produced from 656846 particles.
The classical A/A P/P E/E and hybrid A/P P/E maps were obtained from 82522 and 20134 particles,
respectively. Cryo-EM structures were successfully obtained from 5 preliminary datasets.

Supplementary Fig. 2: Resolution and model validation.

a, Fourier Shell Correlation curves of the half maps and local-masked refinements. **b,** Binary masks used for local-masked refinements. **c,** Local-masked refinements colored by local resolution. **d,** Overall maps for classical and hybrid states and their corresponding local-masked refined maps fitted and colored by local resolution. Cryo-EM map resolution estimates by Fourier Shell Correlation were performed using half- maps from random half-sets.

Supplementary Fig. 3: SILNAS of human mitoribosomal rRNA.
a. Reversed phase-LC isolation of the rRNAs extracted from the HEK 293 mitoribosome (10 µg) applied to a PLRP-S 4000A column (4.6 × 150 mm, 10 µm, Agilent Technologies). **b.** Selective detection and identification of mono-methylated nucleoside isomers in human mitochondrial

rRNAs. The HEK293 12S rRNA, 16S rRNA and CP-tRNA^{Val} (each 200 fmol) were digested with RNase T1 or RNase A, and the products subjected to LC-MS with an in-source fragmentation energy of 70 eV. The signals initially extracted for the second step were at m/z 148.06, 164.06, 125.04, and 124.05 (mass tolerance, +/-5 ppm) for B(mA), B(mG), B(mU) and B(mC), respectively. The extracted chromatograms of the oligonucleotides containing the mono-methylated nucleobase dissociated from RNase A and RNAse T1 digestion of the mitochondrial 12S rRNA (top left and top right panels) and 16S rRNA (bottom left and bottom right panels) are shown, respectively. Pseudo-MS3 spectra of the mono-methylated nucleobases are shown below each. The sequences assigned by Ariadne are shown next to their signals (arrow). The assigned mono-methylated isomers are shown in red. The spectrum was obtained from the m^{2}G anion of the oligonucleotide because the ratio of the signal heights at m/z 133.016 and 149.034 are 0.75. **c.** Table lists all the HEK 293 mt-rRNA modifications identified with their respective occupancies.

Supplementary Fig. 4: Complete rRNA sequence showing rRNA modifications determined by SILNAS.

a, The complete sequence (numbered according to NC_012920.1) and modifications of the human mitochondrial rRNAs and the fragments used for the structural analysis. Digestion fragments produced by RNase T1 (Black solid); RNase A (black shaded bars) and RNase H (Blue solid bars) are indicated. All fragments were identified by Ariadne software (see Methods). Modified

nucleotides (red letters) and their abbreviations are summarized as follows: P, pseudouridine; Π, 1-methyladenosine; ζ, N6,N6-dimethyladenosine; ν, N4-methylcytidine; Σ, 5-methylcytidine; #, 2′-O-methylguanosine; L, N2-methylguanosine; J, 2′-O-methyluridine; T, 5-methyluridine. **b,** 3' end of 12S and 16S mt-rRNAs show hetero sequences. Mitochondrial genome sequence (NC_012920.1) around 3' end of 12S and 16S are indicated as green text. The transcription termination points are indicated with the residue number and arrow. The hetero 3' end of 12S and 16S mitochondrial rRNAs detected in this study are shown in plain text.

Supplementary Fig. 5: Comparison of polyamine densities with antibiotic-treated cells (PDB ID 6ZM5). Densities (colored by chain) corresponding to spermidine and putrescine are absent upon antibiotic treatment [71]. Absence of putrescine is further associated with a conformational rearrangement of the residue A2354. Interactions are shown with dashed lines.

Supplementary Fig. 6: Comparison of protein elements involved in mRNA binding with *E. coli*.

Proteins uS5m, uS7m, uS9m and uS12m (cartoon), mRNA (light-blue surface) from the mitoribosome (left) compared with their bacterial counterparts from *E. coli* [66] (PDB ID 6ZTJ; right). The residues K111 and K126 mark the poly-basic stretch of uS5m. Mitochondria-specific extensions of uS5m and uS9m are positioned along the mRNA channel.

Supplementary Fig. 7: Frequency of Y-N-C-Y motif in the human mitochondrial mRNA transcripts.
Percentage frequency of Y-N-C-Y motif in each of the 13 mitochondrial mRNAs (light grey bars).

Supplementary Fig. 8: Interactions of uS12m with codon in the A-site.

a, In the mitoribosome at 2.2 Å resolution, P73 is in *cis*, and it coordinates a K^+ ion. K72 side-chain interacts with U9 of the A-site codon. **b,** In *T. thermophilus* ribosome (PDB ID 4V51) at 2.8 Å resolution [52], the densities are assigned as Mg^{2+}, P73 is in *trans*, and K72 is pointing away from the A-site codon. **c,** In *E. coli* ribosome (PDB ID 7K00) at 2.0 Å resolution [48], two water molecules have been modeled, and P73 is in *cis*. **d,** In *T. thermophilus* ribosome (PDB ID 6QNR) at 3.1 Å resolution [53], the densities were assigned as K^+ based on long wavelength X-ray diffraction.

Supplementary Fig. 9: Mitoribosomal proteins involved in mRNA binding.
a, N-terminal region of mL40 lodged between A- and P-tRNA. **b,** basic residues 51-55 of mL40 interact with tRNA phosphate backbone. R51 forms salt bridges with A1560 and C1561 phosphates in h44. **c,** The mRNA residue at position 6 is mutated in Coot v0.9 to depict potential H-bond between S47 side chain and base regardless of base identity. **d,** uS9m C-terminal residue R396 interact with P/P-tRNA anticodon stem-loop. **e,** E-tRNA does not directly interact with mRNA in the mitoribosome. Instead, Gly164 and Gly165 of uS7m stack against mRNA nucleotide base. This is in contrast to the bacterial ribosome [62] (PDB ID 4Y4P) where E-tRNA nucleotide at position 34 stacks with mRNA nucleotide base.

Supplementary Fig. 10: mRNA-rRNA interactions in the decoding center and roles of modifications.

a, Mitoribosome in the classical state: rRNA yellow, mRNA blue, tRNA red, modified nucleotides are labeled and colored lime, cryo-EM density in blue mesh. **b,** *E. coli* ribosome [48] (PDB ID 7K00), the same color scheme as in (a). Differences are observed along the mRNA codons: m4C1486 (methylated m4Cm1402 in *E. coli*); m5C1488 (unmodified C1404 in *E. coli*), U1563 (N3-methylated m3U1498 in *E. coli*). Conserved nucleotides at the decoding center G902, A1557, A1558 interact directly with A-site tRNA anti-codon, equivalent to the corresponding residues G530, A1492 and A1493 in *E. coli*.

Supplementary Fig. 11: Analysis of nucleotide binding by mS29 and its role in mitoribosome function.

a, Immunoblot analysis comparing steady-state levels of proteins using whole cell protein lysates from HEK293T (WT), *mS29*-KO (KO), and *mS29*-KO cells stably expressing either WT mS29 or single mutants in the ATP binding site (K134A) or GDP binding site (Y173A, R177A, K245A, H291A, K295A) under an attenuated CMV6 promoter (Δ5pCMV6). COX1 and COX2 were used as surrogates of mitoribosome function, uS3m as a control for the SSU, and β-ACTIN as a loading control. Single mutations of mS29 did not have an effect on the steady-state levels of mitochondrial proteins. **b,** Metabolic labeling of mitochondrially translated peptides in whole cells from the indicated cell lines using ^{35}S-methionine for 15 min. Immunoblotting for β-ACTIN was used as a loading control. Newly synthesized peptides are identified on the left. **c,** Cyanide-sensitive endogenous cell respiration rate measured polarographically using a Clark electrode in the indicated cell lines. Cell respiration rates are presented as nmol of O_2 consumed per minute per one million cells. Data represents the mean ± SD of three independent repetitions. **d,** 2D diagram of GDP interactions. **e,** Schematic depiction of intersubunit contacts mediated by mS29.

	Monosome (consensus) PDB: 7QI4 EMD-13980	Monosome (A/A P/P E/E) PDB: 7QI5 EMD-13981	Monosome (A/P P/E) PDB: 7QI6 EMD-13982
Data collection and processing			
Electron microscope	Titan Krios	Titan Krios	Titan Krios
Camera	K2 Summit (counting mode)	K2 Summit (counting mode)	K2 Summit (counting mode)
Magnification	165,000	165,000	165,000
Voltage (kV)	300	300	300
Electron exposure ($e^-/Å^2$)	29–32	29–32	29–32
No. of frames	20	20	20
Defocus range (μm)	−0.6 to −2.8	−0.6 to −2.8	−0.6 to −2.8
Pixel size (Å)	0.83	0.83	0.83
Symmetry imposed	C_1	C_1	C_1
Final particle number (no.)	656,846	82,522	20,134
Map resolution (Å) (Overall/ LSU-body/ CP/ L10-L12-stalk/ L1-stalk/ SSU-body/ SSU-head/ mS39/ SSU-tail/ mS39-LRPPRC-SLIRP)	2.21/ 2.08/ 2.36/ 2.38/2.89/ 2.31/ 2.36/ 2.45/ 2.44/ −	2.63/ 2.44/ 2.78/ 2.84/3.20/ 2.75/2.72/ 2.93/ 2.91/ −	2.98/ 2.84/ 3.25/3.29/3.53/3.07/3.03/3.47/3.39/ −
FSC threshold	0.143	0.143	0.143
Map resolution range (Å)	1.9-8.0	2.1-8.0	2.4-8.0
Refinement			
Initial model used (PDB code)	6ZSG, 6RW4	6ZSG, 6RW4	6ZSG, 6RW4
Model resolution (Å)	2.10	2.50	2.80
Model to map CC (CC_{volume})	0.84	0.84	0.81
FSC threshold	0.5	0.5	0.5
Map-sharpening B factor ($Å^2$) (Overall/ LSU-body/ CP/ L10-L12-stalk/ L1-stalk/ SSU-body/ SSU-head/ mS39/ Model composition	-38/ -34/ -53/-55/-74/ -43/ -45/ -52/ -52/ −	-36/ -28/ -52/-53/-60/ -39/ -42/ -51/ -51/ −	-42/ -21/ -50/-59/-40/ -35/ -49/ -64/ -57/ −
Non-hydrogen atoms	187689	189567	183520
Hydrogen atoms	152459	154149	153182
Protein chains	88	88	88
RNA chains	7	7	6
Protein residues (non-modified/ *N*-acetylAla/ *N*-acetylSer/ *N*-acetylThr O^1-methylisoAsp)	14790/3/1/1/1	14891/3/1/1/1	14879/3/1/1/1
RNA residues (non-modified/ mG/ mU/ m^1A/ m^2G/ψ /m^4C/ m^5C/ m^5U/ m^6_2A)	2750/2/ 1/ 2/ 1/ 2/ 1/ 1/ 1/ 2	2814/2/ 1/ 2/ 1/ 2/ 1/ 1/ 1/ 2	2744/ 2/1/ 2/ 1/ 2/ 1/ 1/ 1/ 2
Ligands (ATP/ GDP/ NAD/ 2Fe-2S/ spermine/ spermidine/ putrescine)	1/ 1/ 1/ 3 /1 /4/ 1	1/ 1/ 1/ 3 /1 /4/ 1/	1/ 1/ 1/ 3 /1 /4/ 1
Ions (Zn^{2+}/ K^+/ Mg^{2+})	3/ 54/ 206	3/ 54/ 208	3/ 54/ 202
Waters	8,001	7,714	3,304
Mean atomic B-factor ($Å^2$)			
Protein	42.146	49.50	33.04
RNA	41.81	45.12	32.94
Ligand	27.34	26.81	16.35
Water	25.96	24.50	11.57
Validation			
Ramanchandran plot (%)			
Outliers	0.03	0.03	0.02
Allowed	1.45	1.50	1.54
Favored	98.52	98.47	98.44
Clash score	1.48	2.20	2.03
RMSD			
Bonds (Å)	0.002	0.004	0.002
Angles (°)	0.4531	0.457	0.434
Rotamer outliers (%)	0.00	0.00	0.00
C_β outliers (%)	0.00	0.00	0.00
CaBLAM outliers (%)	0.85	0.77	0.81

Supplementary Table 1: Data collection and model statistics.