Mitochondrial temperature homeostasis

resists external metabolic stresses

Mügen Terzioglu[1,#], Kristo Veeroja[1], Toni Montonen[1], Teemu O. Ihalainen[1],

Tiina S. Salminen[1], Paule Bénit[2], Pierre Rustin[2], Young-Tae Chang[3], Takeharu Nagai[4]

& Howard T. Jacobs[1,5,#,§]

[1]Faculty of Medicine and Health Technology, FI-33014 Tampere University, Finland; [2]Université Paris Cité, Inserm, Maladies neurodéveloppementales et neurovasculaires, F-75019 Paris, France; [3]Department of Chemistry, POSTECH, 37673 Pohang, Republic of Korea; [4]SANKEN (The Institute of Scientific and Industrial Research), Osaka University, Ibaraki, Japan; [5]Department of Environment and Genetics, La Trobe University, Melbourne, Victoria 3086, Australia

ORCID IDs: Terzioglu = 0000-0003-0872-3046; Rustin = 0000-0002-5500-7280; Montonen = 0000-0003-3784-0890; Ihalainen = 0000-0003-4351-8697; Salminen = 0000-0002-7232-0754; Bénit 0000-0002-2321-8972; Chang = 0000-0002-1927-3688; Nagai = 0000-0003-2650-9895; Jacobs = 0000-0003-1895-6003

[§]Corresponding author: Howard T. Jacobs, Faculty of Medicine and Health Technology, FI-33014 Tampere University, Finland. Phone: +358503412894, E-mail: howard.jacobs@tuni.fi, howy@jacobs.net

[#]who shares corresponding and senior authorship. E-mail: mugen.terzioglu@tuni.fi

Running title: mitochondrial temperature

ABSTRACT

Based on studies with a fluorescent reporter dye, Mito Thermo Yellow, and the genetically encoded gTEMP ratiometric fluorescent temperature indicator targeted to mitochondria, the temperature of active mitochondria in four mammalian and one insect cell-line was estimated to be up to 15 °C above that of the external environment to which the cells were exposed. High mitochondrial temperature was maintained in the face of a variety of metabolic stresses, including substrate starvation or modification, decreased ATP demand due to inhibition of cytosolic protein synthesis, inhibition of the mitochondrial adenine nucleotide transporter and, if an auxiliary pathway for electron transfer was available via the alternative oxidase, even respiratory poisons acting downstream of complex I. We propose that the high temperature of active mitochondria is an inescapable consequence of the biochemistry of oxidative phosphorylation and is homeostatically maintained as a primary feature of mitochondrial metabolism.

IMPACT STATEMENT

Mitochondria are at least 15 ºC hotter than their external environment in living cells. In response to diverse metabolic stresses, mitochondrial temperature re-adjusts to this value whenever possible.

INTRODUCTION

The mitochondrial system of oxidative phosphorylation (OXPHOS) is considered the most efficient energy conversion platform of non-photosynthetic eukaryotes. However, it does not operate at or anywhere near 100% thermodynamic efficiency. Much of the free energy released by the reoxidation of primary electron carriers such as NADH is converted to heat. Note that this applies under normal physiological conditions[1,2,3], not only in cells expressing uncoupler proteins or treated with a chemical uncoupler. Furthermore, this heat production is not mere 'waste', since it is a major and regulatable energy source for maintaining body temperature in homeotherms such as mammals and birds[4], and possibly many other organisms, even those living at a wide range of temperatures in varying environments. Uncoupler proteins merely serve an auxiliary function under cold stress, as shown by the fact that, in mice, ablation of the major uncoupling protein UCP1 is nonlethal[5] and does not affect basal body temperature.

Using the mitochondrially targeted fluorescent dye Mito Thermo Yellow (MTY), it was recently shown that intramitochondrial temperature in respiring, human HEK293-derived (immortalized embryonic kidney) cells is at least 10-12 °C higher than the ambient temperature at which the cells are maintained[6]. In essence, this inference was based on the fact that the MTY fluorescence change, after cells were treated with OXPHOS inhibitors such as rotenone (targeted on OXPHOS complex I, cI), antimycin (targeted on OXPHOS complex III, cIII), oligomycin (targeted on OXPHOS complex V, cV) or subjected to prolonged anaerobiosis indicated a temperature drop of this magnitude, based on internal calibration at the end of each experiment. Inferences from the use of protein-based fluorescence probes targeted to mitochondria are similar[7]. In addition, similar methods have indicated that the endoplasmic reticulum (ER)[8], cell nucleus[9,10], plasma membrane[11] and different cellular compartments such as the cell body and neurites of neurons[12], are at

temperatures distinct from the extracellular environment and from each other.

These findings have far-reaching implications for biochemistry and physiology, but need further validation and elaboration before they can be generally accepted. In particular, since the cells of many organisms, notably homeotherms, are unable to tolerate prolonged high or low temperatures when applied externally, the question arises as to how much the temperature of mitochondria is permitted to vary *in vivo*, and whether and how it is regulated according to physiological conditions[13].

In this study we undertook to assess mitochondrial temperature in a number of cultured cell-lines from different organisms, and subjected to different metabolic stresses involving well-characterized inhibitors and the transgenic expression of the thermogenic alternative oxidase from a marine invertebrate. Broadly, the outcome of these studies verifies that mitochondrial temperature is physiologically maintained at up to ~15 ℃ above ambient temperature, and is subject to homeostatic regulation in response to metabolic stress.

RESULTS

Based on MTY fluorescence mitochondrial temperature is up to 18 °C above ambient

In previous studies[6], MTY fluorescence was tracked in continuously oxygenated human HEK293-derived cells maintained at a constant external temperature, in which mitochondria had been prelabelled with MTY and then treated with one of several OXPHOS inhibitors under controlled oxygen concentration. Internal calibration in each experiment indicated a mitochondrial temperature decrease of 10-12 °C (Ref. 6) following the transition from normoxia to anaerobiosis, or inhibition of the OXPHOS system by classic inhibitors (rotenone, antimycin, cyanide or oligomycin – see Fig.

1). We conducted a systematic follow-up study with different inhibitors and cell-lines, and sufficient repeats to generate statistical reliability (Fig. 2: see Supplementary Figs. S1 and S2 for details of the instrumentation, its validation and calibration), once the MTY fluorescence intensity had reached a stable value following initial oxygenation. The findings may be summarized as follows.

In most cell-lines tested there was a clear gradation of mitochondrial temperature decrease using inhibitors that act further along the OXPHOS chain: the effect was always greatest after oligomycin (inhibitor of ATP synthase or OXPHOS complex V, cV) and generally least for rotenone (cI inhibitor), with potassium cyanide (inhibitor of OXPHOS complex IV, cIV) and antimycin (cIII inhibitor) intermediate (Fig. 2). In general, findings were similar for different cell-lines, with a temperature decrease of the order of 17-19 °C produced by oligomycin. This implies a mitochondrial temperature of at least 54 °C, assuming that cV inhibition abolishes all mitochondrial heat production. Two different isolates of immortalized mouse embryonic fibroblasts (iMEFs) gave similar results, with no significant difference in the extent of mitochondrial temperature decrease produced by rotenone to inhibit cI, by antimycin to inhibit cIII (Fig. 2C. 2D) or a combination of the two drugs (Fig. 2D). Unlike rotenone plus antimycin (Supplementary Fig. S3A, panel i) other combinations of OXPHOS inhibitors did not give stable fluorescence values (Supplementary Fig. S3A, panels ii-vi) and were not considered further. We were also unable to estimate directly the individual contribution to mitochondrial temperature maintenance of the ALR (Evr1) pathway (of mitochondrial protein import to the inter-membrane space), in which electrons are fed directly to cytochrome c (Fig.1): preparations of the ALR inhibitor MitoBloCK-6 (Ref. 14) that we were able to source had a distinct yellow colour that rendered measurements of MTY fluorescence uninterpretable (Supplementary Fig. S3B).

We investigated the effect of culturing iMEF cells in low-glucose medium, with or without galactose supplementation, aiming to enhance dependence on mitochondrial respiration[15]. This culture-medium alteration had no significant effect on the effect of cI inhibition on mitochondrial temperature (Supplementary Fig. S3C).

The inferred temperature of mitochondria in S2 cells from the fruit-fly *Drosophila*, a poikilotherm, when suspended and oxygenated at 25 °C, showed similar responses to OXPHOS inhibition as mammalian cells suspended at 37 °C, except that we were unable to derive a stable reading after oligomycin treatment (see Supplementary Fig. S3D, panel i). MTY fluorescence was still increasing 2 hours after the addition of oligomycin, whereas other inhibitors, such as KCN, when applied to S2 cells, gave a stable reading over the same time period (e.g. Supplementary Fig. S3D panel ii). Fluorescence reached a stable value after the addition of oligomycin in all of the other cell-lines tested, e.g. O2OS human osteosarcoma cells as shown (Supplementary Fig. S3D panel iii).

Protein ratiometric fluorescence confirms mitochondrial temperature measured by MTY
Although the previous study excluded the most obvious possible artifacts from the use of MTY fluorescence as a mitochondrial thermometer[6], such as an effect of membrane potential, the redox state of the respiratory chain or reactive oxygen species (ROS) production, artifacts of unknown provenance could always be in play. We therefore set about applying a different method to measure mitochondrial temperature in these cell lines, based on the ratiometric fluorescence of mito-gTEMP[7]. This technique is based on the fluorescence ratio between two previously developed fluorescent proteins, mT-Sapphire and Sirius, co-expressed as a single, self-cleaved polypeptide, with each cleavage product targeted to mitochondria using a reiterated cytochrome oxidase subunit 8 (COX8) presequence (Fig. 3A). The basis of the method is the fact that the fluorescence of Sirius is highly temperature sensitive, declining (like MTY) as temperature rises, whereas that of mT-

Sapphire is not[7]. We stably transfected HEK293T, U2OS and iMEF(P) cells with mito-gTEMP and confirmed mitochondrial targeting, as exemplified in Fig. 3B and Supplementary Fig. S4A. Since the two proteins are expressed at a fixed stoichiometry, the peak fluorescence ratio is independent of expression level, which may vary even within a cloned mammalian cell population[16], as seen, for example, in Fig. 3B.

To check for possible artifacts introduced by mito-gTEMP in the presence of OXPHOS inhibitors we checked whether the drugs alone in PBS affect background fluorescence in the absence of cells. Background fluorescence at the wavelengths used to track mT-Sapphire and Sirius were both more than an order of magnitude less than that of the fluorescent proteins expressed in iMEFs, and were unaffected by rotenone (Supplementary Fig. S4B). Background fluorescence at both wavelengths was slightly increased by oligomycin (Supplementary Fig. S4C), but the fluorescence ratio was unchanged (Supplementary Fig. S4E). In contrast, antimycin produced a substantial change in background fluorescence at the Sirius wavelength (Supplementary Fig. S4D), which affected the fluorescence ratio (Supplementary Fig. S4F), making it unsuitable for use in estimating mitochondrial temperature shifts.

Since, based on MTY fluorescence, we had inferred a maximal amount of mitochondrial temperature decrease in the presence of oligomycin, we used oligomycin-treated cells to calibrate the mito-gTEMP fluorescence ratio against externally imposed temperature. This assumes that any residual mitochondrial heat production in oligomycin-treated cells is negligible and invariant with respect to externally applied temperature changes.

After oligomycin treatment, iMEF mitochondria showed a decline in the mito-gTEMP fluorescence ratio over ~10 min, thereafter holding a constant value for at least a further 25 min that is assumed

to represent the ambient temperature (Fig. 3C). This enabled us to construct an *in vivo* calibration curve for the mito-gTEMP fluorescence ratio against externally imposed temperature (Fig. 3D), which fitted a linear function over the temperature range at which we were able to conduct the measurement: higher temperatures result in loss of cell integrity and viability[17], so were not used. Extrapolation to higher mitochondrial temperatures is therefore based on the assumption that the relationship remains linear, as was determined previously for the isolated gTEMP proteins[7].

Oligomycin treatment of mito-gTEMP expressing cell clones derived from three different mammalian cell-lines gave extrapolated values for the mitochondrial temperature decrease in the same range as those inferred by MTY, although slightly (1-2 °C) less in each case, implying an internal mitochondrial matrix temperature of ~52-54 °C (Fig. 3E). Rotenone treatment of iMEF(P) cells also produced a similar amount of mitochondrial temperature decrease when estimated by mito-gTEMP as by MTY (Fig. 3E).

Individual cells and organelles show mitochondrial temperature differences and fluctuations

As studied by either method, mitochondrial temperature represents an average over many cells, each containing many mitochondria. We used confocal imagining to look at the profile of mitochondrial temperature in individual iMEF(P) cells and organelles stained with MTY (Fig. 4A, panel i). This revealed much greater heterogeneity between and even within cells not seen, for example, with the ubiquitous mitochondrial stain MitoTracker™ Deep Red FM (Fig. 4A, panel ii). Within individual cells stained with MTY we observed some more brightly staining puncta (Fig. 4B; Supplementary Movie S1). Time-lapse imaging revealed that mitochondria in individual cells were cooled to variable extents by oligomycin treatment (Fig. 4C).

Metabolic adaptation can maintain or restore mitochondrial temperature

When mito-gTEMP expressing iMEF cells were incubated in PBS at 37 ºC without the addition of any inhibitor we observed that, after reaching a steady fluorescence ratio indicative of a temperature of 53-54 ºC, there was a modest but progressive decline in the fluorescence ratio, such that by 35 min of incubation the inferred mitochondrial temperature was around 4 ºC lower than the peak value reached at approximately 10 min (Fig. 5A, 5B). Although remaining much warmer than the surrounding medium, this suggests that, in the complete absence of an externally supplied, metabolizable substrate, mitochondrial metabolism adjusts to a slightly lower heat output and steady-state temperature. To explore this further we added the main substrates normally provided to mammalian cells in culture, namely glucose and glutamine. As expected, this enabled mitochondria to maintain a higher temperature of ~53 ºC at the end of the experiment, although this was still slightly less than the initial temperature peak of ~55 ºC between 5-10 min of incubation (Fig. 5B, 5C). However, the addition of glucose and glutamine at 10 min did produce a small rise in mitochondrial temperature (Fig. 5B, 5D). In sum, these data indicate that mitochondrial temperature fluctuates within a narrow range according to substrate supply.

In order to explore the issue further, we treated cells with bongkrekic acid (BKA), a potent inhibitor of the mitochondrial adenine nucleotide translocase, ANT (Ref. 18). The ANT exchanges ADP and ATP across the mitochondrial inner membrane (IMM), thus enabling OXPHOS to supply the rest of the cell with ATP. Previous investigators have shown that cultured mouse fibroblasts respond to BKA by inducing glycolysis and downregulating mitochondrial respiration, whilst maintaining a high mitochondrial membrane potential[19]. In initial trials we found that BKA quenches the fluorescence of MTY, even in the absence of cells (Supplementary Fig. S4G). Therefore, we used the mito-gTEMP fluorescence ratio, which was less disturbed by the drug, to estimate its effects on mitochondrial temperature. We reasoned that the metabolic switching induced by BKA should result in a substantial mitochondrial temperature decrease, due to the expected lowering of flux

through the respiratory chain. However, the mitochondrial temperature decrease that we observed was minimal, following an apparent slight increase upon addition of the drug (Fig. 5E, F), qualitatively consistent with the MTY signal after discounting the quenching effect (Supplementary Fig. S4G).

AOX maintains or restores high mitochondrial temperature in the presence of OXPHOS inhibitors

Next we looked at iMEF cells constitutively expressing the non-protonmotive alternative oxidase (AOX) from *Ciona intestinalis*, which provides an alternative pathway for respiratory electron flow from ubiquinol to oxygen, if cIII and/or cIV are inhibited[20]. We first verified by respirometry that iMEF AOX cells were resistant to inhibitors of cIII (antimycin, Supplementary Fig. S5A) and cIV (KCN, Supplementary Fig. S5B).

In cells that principally metabolize cI-linked substrates, AOX makes only a minor contribution to respiratory electron flow under non-inhibited conditions[21]. However, if cI is blocked, forcing cells to switch to other ubiquinone-linked substrates, AOX becomes active[21]. Once again, the cell must switch substantially to glycolysis to sustain ATP levels, and to cytosolic NADH reoxidation via lactate dehydrogenase.

Consistent with these expectations, the initial mitochondrial temperature decrease produced by the cI-inhibitor rotenone in AOX-expressing iMEFs was very similar to that of control iMEFs, with a mitochondrial temperature decrease of ~10 °C based both on MTY fluorescence (Fig. 6A, 6B) and the mito-gTEMP fluorescence ratio measured in iMEF cells stably transfected with AOX (Fig. 6B, Supplementary Fig. S6A). However, this was followed by a gradual mitochondrial rewarming (Fig. 6A panel i, 6B, 6C). AOX-expressing iMEFs treated with other OXPHOS inhibitors showed a

similar behavior (Fig. 6A panels ii-iv, 6C), except that the initial mitochondrial temperature decrease was substantially less than that seen in control iMEFs (Fig. 6A panels ii-iv, 6B), whilst the rate of rewarming differed according to the inhibitor used (Fig. 6C). We were not able to estimate directly the contribution of AOX to temperature maintenance, due to fluorescence quenching by the AOX inhibitor n-propyl gallate, nPG (Supplementary Fig. S6B).

Mitochondrial temperature is maintained despite varying ATP demand

To model the effects on mitochondrial temperature of a more physiologically relevant perturbation, we downregulated cytosolic protein synthesis, the major cellular consumer of ATP, by treating iMEFs with anisomycin (Fig. 7). Using our earlier calibrations, we initially estimated the mitochondrial temperature in cells prior to anisomycin treatment, then followed the effect of anisomycin treatment for up to 18 hours (Fig. 7). In contrast to our expectation that decreased ATP demand would decrease electron flux in the OXPHOS system and lead to a drop in mitochondrial temperature, the latter instead showed a transient slight increase (Fig. 7A). Thereafter, high mitochondrial temperature was sustained for many hours, although by 18 hours of exposure to the drug many of the cells were no longer attached and probably dying (the cells analyzed in Fig. 7A were those that had remained attached). AOX-expressing iMEF cells (Fig. 7B) showed an almost identical behavior, except that the cells remained healthy-looking even after 18 hours in anisomycin. To evaluate the contribution of cI and cV to the maintenance of mitochondrial temperature under these conditions of decreased ATP demand, we stained anisomycin pretreated iMEFs with MTY, then tracked fluorescence changes after treatment with rotenone (Fig. 7C) or oligomycin (Fig. 7D). The mitochondrial temperature of anisomycin-treated control iMEFs showed a significantly lesser effect of rotenone than untreated cells (Fig. 7C, blue trace), and this effect was increasingly pronounced after longer times of anisomycin exposure. Mitochondrial temperature in the surviving cells after 18 h in anisomycin was almost unaffected by rotenone. In contrast, and

despite some experiment-to-experiment variation, anisomycin-exposed control iMEFs were as affected by oligomycin as prior to treatment, even up to 18 h (Fig. 7D, blue trace). AOX-expressing iMEFs behaved differently. After anisomycin treatment, they showed a consistently greater mitochondrial temperature decrease produced by rotenone than did control iMEFs (Fig. 7C, red trace). Although the effect of oligomycin declined with increasing times of anisomycin exposure, it did so more gradually than for control iMEFs. The mitochondria of anisomycin-treated iMEF(AOX) cells showed a temperature decrease by oligomycin significantly less than anisomycin-treated control iMEF(P) cells, regardless of the length of time of anisomycin exposure, but also showed a possibly increased effect of oligomycin, with increasing anisomycin exposure time (Fig. 7D, red trace), at least for the first hours. Anisomycin-exposed iMEF(AOX) cells showed the same slow rewarming after rotenone treatment (Fig. S7A panel i, ii) as did iMEF(AOX) cells not exposed to anisomycin (Fig. 5A panel i, 5C), whilst control iMEFs did not show any such effect from rotenone (Fig. S7A, panel iii, iv). Conversely, the rewarming effect after oligomycin treatment was slower and of much lower magnitude for anisomycin-treated than untreated iMEF(AOX) cells (Fig. S7B panels i, ii), whilst a modest rewarming was also seen for control iMEFs (Fig. S7B panel iii, iv).

DISCUSSION

In this study, using the mitochondrially targeted, temperature-sensitive dye MTY applied to five cell-lines, we confirmed that physiologically active mitochondria *in vivo* exhibit high temperature, some 15-18 °C above that of the rest of the cell, i. e., in the 52-55 °C range in mammalian cells. We validated the measurements using the mito-gTEMP system, based on the ratiometric fluorescence of two mitochondrially localized reporter proteins. We visualized mitochondrial temperature fluctuations at the single-cell level, and documented the maintenance of high mitochondrial

temperature despite variations in growth conditions, substrate availability, ATP demand, and even OXPHOS poisons, if an alternative respiratory pathway is available.

No chemical reaction is 100% thermally efficient, and those of OXPHOS are no exception, with up to 60% of the energy yield from the biological oxidations of respiration and ATP synthesis released as heat[2,3]. It has long been assumed that this heat was simply radiated to the rest of the cell or the external environment, maintaining intracellular temperature at a uniform 37-42 ºC in mammals and birds, or at a variable, environmentally determined temperature in poikilotherms such as insects, generally considered as ectothermic. Our findings are inconsistent with this model, indicating instead that much of the heat generated by OXPHOS is retained within the organelle, keeping mitochondrial temperature far above that of other cell components and, whenever possible, within a narrow range. Heat conductance from the mitochondria might also be modulated to maintain mitochondrial temperature.

Validation and calibration of high mitochondrial temperature

In previous studies using MTY in human HEK293 and HeLa cells we documented fluorescence changes after treatment with OXPHOS inhibitors that were indicative of high mitochondrial temperature[6,22]. In the present study we extended these findings to other cell lines, including one from *Drosophila*, and two isolates of iMEFs, immortalized using a different cocktail of factors[23,24]. We systematically conducted multiple repeats with each of five drugs impacting the OXPHOS system. In each case, we implemented an internal calibration step after MTY fluorescence had reached a plateau level in the presence of the inhibitor, by abruptly shifting the external temperature up, then down by 3 or 5 ºC, at minimally 4 minute intervals. In each case, it is assumed that the observed fluorescence shift is due only to the change in the externally applied temperature (i.e., that any 'corrective' response of the cell in the presence of the inhibitor is negligible). This is supported

by the fact that the new fluorescence values reached during the calibration steps are essentially constant over the time-scales used (see Fig. S2C for examples). Internal calibration in each experiment provides a control for variation in the exact number of cells or the amount of label taken up. The use of a Peltier jacket surrounding the cuvette in the apparatus ensures that temperature changes in the cuvette rapidly and accurately reflect those applied, as verified (Fig. S2A, S2B). Importantly, the mito-gTEMP fluorescence ratio, and its calibration against externally applied temperature, provides a second check on the findings obtained using MTY. Although the principle of the mito-gTEMP method is different, being based on fluorescent reporter proteins rather than an externally applied dye, the calibration steps rely on the same, validated assumptions, and produced temperature readouts very similar to those derived using MTY, although typically several ºC lower in the case of each of the mammalian cell-lines tested (Fig. 3E).

Although the two methods gave essentially concordant results, both have limitations. Apart from our failure to obtain a stable fluorescence reading for S2 cells treated with oligomycin, we did not experience problems working with MTY on any of the cell-lines tested in the presence of single OXPHOS inhibitors, such as were reported earlier for a human primary fibroblast line[22]. However, using combinations of OXPHOS inhibitors, other than rotenone plus antimycin, we were not able to infer mitochondrial temperature by the MTY method, due to unstable fluorescence readings (Fig. S3A). These are likely due to dye leakage resulting from membrane-potential anomalies of the type reported earlier[22].

The excitation wavelength for mito-gTEMP (360 nm) falls within the ultra-violet range, and the resulting cellular damage under continuous illumination for more than 60 min renders calibration unreliable[25]. An alternative probe set, based on the recently developed B-gTEMP[25], should overcome the problem in future studies. In the current report, we used mito-gTEMP only up to 35

min, which generated a reliable calibration curve (Fig. 3D). However, this is not long enough to profile very slow temperature changes such as the rewarming we observed using MTY on AOX-expressing cells (Fig. 6). In addition, we found that some inhibitors, notably antimycin (Fig. S4F), though not rotenone (Fig. S4B) or oligomycin (Fig. S4E), affected the mito-gTEMP fluorescence ratio, compromising its use. Similarly, the effects of BKA (Fig. S4G), nPG (Fig. S6B) and MitoBloCK-6 on MTY fluorescence readings preclude their use with the probe. One important caveat regarding the application of mito-gTEMP is that it was necessary to extrapolate the calibration curve beyond 42 ºC, assuming the linearity seen earlier for the isolated proteins[7]. Above this temperature, *in vivo* calibration is not possible, due to the loss of cellular viability and integrity, accompanied by major changes in protein composition and distribution[17].

The finding that mitochondrial temperature is of the order of 50 ºC or more and distinct from that of the rest of the cell has been challenged on both theoretical[26] and experimental grounds[27]. However, recent data indicate that the slow thermal diffusivity of mammalian cells[25] is sufficient to account for the maintenance of intracellular temperature differences inferred under physiological conditions and further documented here. Furthermore, the maintenance of high mitochondrial temperature as a source of intracellular heat is supported by theoretical considerations[1]. It should also be noted that previous studies using different probes have shown that the temperature across the cell is not uniform. Some organelles or active regions manifest distinct temperature inhomogeneities: notably, the nucleus is 1-2 ºC warmer than the cytoplasm[9,10], the nucleolus even hotter[28] and perinuclear mitochondria hotter than those at the cellular periphery[29].

We previously reported[6] that isolated OXPHOS complexes I and V are heat labile, though OXPHOS complex II (cII), cIII and cIV activities have temperature optima in the 50-55 ºC range when studied in intact mitochondrial membranes. Others have recently reported that the imposition

of an external temperature of 43 ºC or above to cells or to isolated mitochondria damages respiratory function and disturbs OXPHOS complex and supercomplex organization[27]. Thus, externally applied heat delivered to mitochondria, which our assays indicate are already at elevated temperatures, may push them beyond their operational limit, implying that mitochondrial heat production *in vivo* must be finely attuned to cellular needs, to avoid catastrophic overheating. On the other hand, a recent meltome analysis of human proteins found that those constituting the respiratory chain complexes have median Tm values of around 60 ºC (Ref. 30), consistent with our inference of high mitochondrial temperature.

Inhibition of each OXPHOS complex produces specific temperature effects on mitochondria

As the OXPHOS system is traversed from cI through cV, we observed significant increases in the temperature effect of specific inhibitors, with essentially the same pattern observed in two iMEF lines, U2OS, HEK293T and even in *Drosophila* S2 cells, as far as methods allowed, i.e., oligomycin> KCN > antimycin ≥ rotenone. Oligomycin inhibits all protonmotive components of the OXPHOS system, including those that feed electrons via ubiquinol to cIII, even though they are not directly protonmotive. The lesser amount of mitochondrial temperature decrease produced by blocks on cI, cIII or cIV (Fig. 2) is not the result of incomplete inhibition: the inhibitor doses were pre-checked on all cell-lines by respirometry and are similar to those used in many other studies. Instead, we posit that it reflects the proportion of electron flow entering the respiratory chain at each step. Since rotenone and antimycin inhibition had the same mitochondrial temperature effect in iMEFs, both individually and in combination, we infer that electron flow entering the chain independently of cI in this cell background is negligible. In contrast, antimycin produced a significantly greater mitochondrial temperature decrease in U2OS cells than did rotenone (Fig. 2B), from which we conclude that around 20-30% of electron flow into cIII in this cell line must come from other ubiquinone-linked enzymes. In iMEFs (Fig. 2C, 2D) and S2 cells (Fig. 2E), though not

in HEK293T or U2OS cells, KCN had a significantly greater mitochondrial temperature effect than antimycin, which implies that, in these cell lines, a measurable amount of electron flow to oxygen enters the OXPHOS system via cytochrome c independently of cIII, i.e. via the Mia40-Evr1/ALR protein import pathway, albeit that we were not able to verify this directly using the Evr1/ALR inhibitor MitoBloCK-6. There may also be other, as yet uncharacterized, electron donors to cytochrome c. In all cell lines that we were able to test, oligomycin produced a significantly greater mitochondrial temperature decrease than KCN (Fig. 2A-2D). Since KCN should block all respiratory proton pumping across the IMM, we infer that cV itself must be able to generate some heat in the complete absence of cellular respiration, presumably by the hydrolysis of mitochondrially imported ATP. Since this residual thermogenesis is inhibited by oligomycin, it must nevertheless involve cV, in a reaction that is itself protonmotive, and hence limited by the rate of dissipation of the IMM proton gradient by other processes, such as protein, metabolite and ion transport.

In cells expressing AOX, the initial effects of OXPHOS inhibition at these various steps was different (Fig. 6A, 6B). Rotenone inhibition of electron flow through cI initially produced the same temperature decrease in AOX-expressing iMEF mitochondria as in control iMEF mitochondria (Fig. 6A, 6B). This is consistent with the fact that AOX by-passes cIII+cIV but not cI[20,31]. In contrast, the mitochondrial temperature decrease produced by antimycin inhibition of cIII was substantially less than in control cells (Fig. 6B). Electron flow through the inherently thermogenic AOX is expected substantially to replaces the normal heating effect of electron flow through cIII+cIV. Note that, when cIII and cIV are able to function normally, electrons flowing through cI should not be diverted via AOX[21]. However, the mitochondrial temperature decrease brought about by antimycin in AOX-expressing MEFs was not zero. Part of the explanation is that the capacity of AOX to support electron flow from cI through to oxygen in these cells remains below that of

cIII+cIV, as inferred previously by respirometry[24]. Furthermore, the full engagement of AOX for

cI-linked substrates[21] requires the accumulation of ubiquinol, the reduced form of coenzyme Q.

Nevertheless, the initial temperature decrease produced by antimycin in the mitochondria of AOX-

expressing cells is maintained only for a relatively short time, approximately 10 min (Fig. 6C),

before rewarming commences. The fact that the initial effect of KCN in AOX-expressing cells was

greater than that caused by antimycin supports the idea that some of the electron flow through cIV

in iMEFs is introduced via cytochrome *c* independently of cIII, for which AOX does not provide a

by-pass.

High mitochondrial temperature is maintained under diverse metabolic stress conditions

Despite the fact that. under the rather extreme conditions of toxic inhibition of the OXPHOS

complexes, high mitochondrial temperature is compromised, we uncovered several contexts

wherein it appeared to be homeostatically maintained, hinting at one or more metabolic adaptation

processes. First, it appears to decline only slowly and minimally in response to nutrient starvation,

and to be restored by renewed substrate availability (Fig. 5), consistent with previous estimates by

an independent method, using so-called Upconversion nanoparticles[32]. This implies the existence of

a metabolic reserve even in cells such as fibroblasts that are not normally considered repositories

for storage molecules such as triglycerides or glycogen. Second, in cells expressing AOX we

recorded a gradual restoration of high mitochondrial temperature after the initial decrease provoked

by inhibition of the OXPHOS complexes (Fig. 6A, 6C). Although AOX is not normally present in

vertebrate cells, the metabolic flexibility that it confers may be important in maintaining

mitochondrial temperature and organismal viability in the vast array of species that do possess

AOX but are subject to wide external temperature fluctuations, namely bacteria, fungi, plants,

eukaryotic protists and many animals as well[31]. The exact nature of the metabolic changes that

enable AOX-endowed cells to recover normal mitochondrial temperature over periods of 40-70 min

remains to be elucidated and may be complex. Logically it should involve a remodeling of metabolism towards the oxidation of flavin-linked substrates that directly reduce ubiquinone. Rewarming was evident in the case of cells in which cI was inhibited by rotenone (Fig. 6A, 6C), indicting that it does not involve the acquisition of additional capacity for AOX to receive electrons via cI. Rather, it must involve a metabolic switch to other pathways, even though via AOX these do not support ATP production, only heat generation.

The rather minimal effect of BKA, i.e. a possible transient, small increase in mitochondrial temperature, followed by an equally modest decrease, was also surprising. BKA treatment of immortalized fibroblasts has previously been reported to induce a switch to glycolytic ATP production and decreased flux through the respiratory chain, whilst maintaining high mitochondrial membrane potential[19]. The modest mitochondrial temperature decrease that was inferred may simply correspond with that seen in cells deprived of substrate over the time-period of observation (compare Fig. 5A and 5E or 5B and 5F). Thus, the efficient exchange of adenine nucleotides between mitochondria and the cytosol is not required for the maintenance of high mitochondrial temperature. Instead, the usually minor contribution of intramitochondrial ATP turnover can be boosted sufficiently to drive mitochondrial heat production to near normal levels, possibly involving, once again, a contribution from cV-linked ATP hydrolysis.

The most surprising result that we obtained was the maintenance of high mitochondrial temperature in cells subjected to prolonged treatment with anisomycin at a dose in excess of that which profoundly inhibits cytosolic protein synthesis[33,34]. Protein synthesis uses between 35-70% of the ATP produced in cells[35,36,37,38]. We therefore expected that decreased ATP demand would lead to decreased flux through the respiratory chain and to a sustained drop in mitochondrial temperature. Instead, anisomycin treatment led to a modest but transient increase (~4ºC) in mitochondrial

temperature (Fig. 7A), with gradual reversion to the original mitochondrial temperature after 2-3 h in the presence of the drug. Even after prolonged (18 h) anisomycin treatment, leading to widespread cell death, high mitochondrial temperature was maintained in the surviving cells. The same effects, but without cell death, were also seen in AOX-expressing cells (Fig. 7B).

Similarly, overnight growth in media that should stimulate mitochondrial as opposed to cytosolic ATP production had no effect on mitochondrial temperature (Fig. S3B). Once again, the precise metabolic changes that underlie these phenomena remain uncertain and may not be easy to dissect. Nevertheless, our observations imply a homeostatic mechanism adjusting mitochondrial heat production to metabolic supply and demand, but keeping mitochondrial temperature within a narrow range.

Based on the inhibitor studies shown in Fig. 7C and 7D, mitochondrial temperature maintenance in anisomycin-exposed iMEF cells is less and less dependent on cI with time, yet remains fully dependent on cV. This suggests that, as cellular ATP usage dwindles, cV may start to operate thermogenically as an ATPase, although this does not explain how the resulting proton gradient across the IMM is dissipated. A more complex mechanism, such as the futile creatine cycle recently documented in brown fat[39], may need to be invoked. The adenine nucleotide carrier (ATP-Mg/Pi), encoded by a small multigene family, which ensures the net delivery of adenine nucleotides to the mitochondrial matrix under specific physiological conditions[40], may also be involved in sustaining a futile cycle of intramitochondrial ATP hydrolysis, as well as possible reverse electron transport through cI. In anisomycin-exposed, AOX-expressing cells, where AOX itself provides an alternative mechanism of thermogenesis, cells remained more dependent on cI for the maintenance of high mitochondrial temperature over time, but less dependent on cV.

Biological implications of high mitochondrial temperature

Although the amount of mitochondrial temperature decrease produced after OXPHOS inhibition by oligomycin, when measured by the MTY and mito-gTEMP methods, was similar, it was not identical, and the rather small difference was judged to be significant for at least two of the cell-lines studied, U2OS osteosarcoma cells and iMEFs. Although both probes are efficiently targeted to mitochondria, they are most likely localized in different submitochondrial compartments. As derivatives of GFP, both of the mito-gTEMP reporter polypeptides should reside in the mitochondrial matrix[41]. Conversely, MTY is a rhodamine-related lipophilic dye whose uptake requires a mitochondrial membrane potential, and which is rapidly lost from mitochondria when the membrane potential is abrogated[6]. Thus, it should be mostly localized at or very close to the inner face of the IMM, despite reports that at least some of it interacts with the matrix protein aldehyde dehydrogenase 2, ALDH2 (Ref. 42). Since the IMM is also the location of mitochondrial heat generation, it should logically be warmer than the matrix that it surrounds, accounting for the small temperature differential implied by the two methods.

Since high mitochondrial temperature was here documented in five different cell-lines from three species of animal, including a poikilotherm, it is clearly not just an exceptional feature of one cell-type. Rather it would appear to be a common property of mitochondria and of the chemistry of the OXPHOS system. Moreover, it suggests a novel role for the universal double membrane system of mitochondria, as an insulating layer enabling mitochondria to maintain a high internal temperature, whilst allowing the remainder of the cell to operate at a lower temperature. If this is correct, a corollary is that the ratio of mitochondrial volume to surface area may change in response to metabolic needs and physiological signals that regulate mitochondrial heat output. Put more simply, mitochondria might be expected to fragment into smaller entities when heat output rises, so as to maintain internal mitochondrial temperature, and to fuse into filamentous structures when heat

output is low. They may also fragment if mitochondrial heat production is simply abolished and mitochondria need to absorb as much heat as possible from the rest of the cell, in order to maintain a minimal metabolism.

This proposition could account for some of the apparently paradoxical dynamics of mitochondria which, when subject to various stresses can either undergo widespread fission or hyperfusion[43]. The mitochondrial fusion/fission cycle has hitherto been associated primarily with mitochondrial quality control[44], which is disturbed in patholological states and during aging[45]. However, mitochondrial temperature homeostasis could also be a determinant. Accordingly, it has recently been reported that inhibition of cV, but not of the other OXPHOS complexes, triggers mitochondrial fragmentation[46]. Treatment with rotenone or antimycin were reported to result in rapid depletion of intramitochondrial ATP[46], consistent with our inference that these inhibitors allow some residual mitochondrial heat output driven by ATP hydrolysis, whilst oligomycin does not. Using mito-gTEMP, it was previously shown that treatment of cells with an uncoupler, which is known to provoke mitochondrial fragmentation[47], also resulted in a mitochondrial temperature increase of ~6 ºC (see Fig. 4 of Ref. 7). Using a different spectrophotometric method, other authors estimated the effect of a chemical uncoupler in the 4-5 ºC range[29]. Consistent with the idea that mitochondrial fusion is associated with low mitochondrial heat output, the cell line with the greatest inferred temperature difference between the two methods used in the present study to assess mitochondrial temperature, and also the one where this difference was the most variable experimentally, was U2OS (Fig. 3E), a cell line in which mitochondria are mostly filamentous, e.g. see Ref. 48. In other cell-lines, such as iMEFs, mitochondria are highly fragmented under standard culture conditions, e.g. see Ref. 49.

Staining of cells with MTY revealed bright puncta within mitochondria (Fig. 4B), the nature of

which is unknown. One possibility is that these may correspond with nucleoids. Their bright staining could indicate that they are cooler than their mitochondrial surroundings, or could be due to binding of the dye to one or more components of the nucleoid. In *Saccharomyces cerevisiae* the orthologue of ALDH2 (Ald4) is a nucleoid protein, but this has not been reported to be the case in mammals[50].

Since the OXPHOS system is of bacterial origin, its heat output in bacteria should be similar to that of mitochondria, and we would expect that the thermal insulation provided in gram-negatives by the double membrane and lipopolysaccharide (LPS) cell wall, or by the thicker cell wall in gram-positives, should be at least as effective as that of the double membrane of mitochondria. It will therefore be interesting to implement thermosensitive reporters such as gTEMP or B-gTEMP in bacteria, to assess intracellular temperature, which we would predict to be once again some 15-20 ºC warmer than ambient temperature. This has important potential implications for understanding the metabolic biochemistry of bacteria, including antibiotic action.

Similarly, considering that heat generation is an inescapable cellular function of metabolically active mitochondria, a loss of this activity, as in cases of mitochondrial disease, may compromise the maintenance of intracellular temperature or induce futile cycles of cytosolic ATP generation and usage that may have profound metabolic effects. This could be a crucial aspect of mitochondrial pathology[51] and may also influence progression in other metabolic diseases, including cancer[52].

Conclusion

On the basis of our findings, we propose that the maintenance of high mitochondrial temperature should be regarded as an important homeostatic process of the organelle, along with more traditionally considered parameters such as membrane potential, pH, ATP, metabolite levels and

ROS.

MATERIALS AND METHODS

Cell culture

All cell-lines used are freely available and are not subject to any ethical restrictions on use. Human embryonic kidney-derived cells expressing a mutant version of the SV40 large T antigen, HEK293T[53], U2OS (2T) osteosarcoma cells[54], two isolates of immortalized mouse embryonic fibroblasts, iMEF(L7)[24] and iMEF(P)[23] and immortalized mouse embryonic fibroblasts expressing *C. intestinalis* alternative oxidase, iMEF(AOX)[23] were cultured in DMEM medium containing 4.5 g/L glucose, 10% heat inactivated FBS, and 100 U/mL each penicillin and streptomycin, at 37 °C in 5% CO_2 with 95% humidity. Cells were also grown in low-glucose medium (Sigma, D5546) containing 1 g/L glucose, supplemented with 25 mM galactose where indicated. *Drosophila melanogaster* S2 (Schneider 2) cells (Invitrogen), originally derived from a primary culture of late stage (20-24 h old) embryos[55], were cultured in suspension in Schneider's medium (Sigma S9895-1L) supplemented with 10% heat-inactivated FBS, 100 U/mL each penicillin and streptomycin at 25 °C and diluted 1:6 every 3-4 days.

Transfection of mammalian cells with mito-gTEMP

To avoid the use of neomycin/geneticin selection, HEK293T cells already being resistant, the mito-gTEMP ratiometric reporter pair (see Fig. 3A) was recloned from mito-gTEMP_pcDNA3 (Addgene, plasmid #109117) into the pTriEx™-1.1 Hygro plasmid (Novagen 70928). Stably expressing cell lines were generated using FuGENE® HD Transfection Reagent (Promega), according to manufacturer's instructions. Based on preliminary trials, hygromycin selection was

implemented at the following doses: iMEF(P) and U2OS – 145 µg/mL, iMEF(AOX) 240 µg/mL, and HEK293T 450 µg/mL.

Spectrofluorometry

For spectrofluorometric measurement of mitochondrial temperature using Mito Thermo Yellow, MTY[42], sub-confluent cells were treated with 100 nM MTY supplemented with warmed fresh medium, for 15 min. Cells (5×10^6 HEK293T and U2OS cells, 7×10^6 iMEFs, and 10×10^6 S2 cells) were trypsinized and collected by centrifugation at 250 g_{max} for 3 min. The pellet was washed once in 10 mL PBS warmed to 37 °C, then maintained as a concentrated pellet at 37 ℃ for 10 min, for anaerobiosis. Cells were resuspended in PBS in a magnetically stirred 3.5 mL quartz cuvette with 10 mm optical path (Hellma Analytics, Germany), placed in a Peltier temperature-controlled chamber set at 38 °C (25 °C for S2 cells). The time-based fluorescence signal at constant emission (excitation 542 nm, emission 562 nm) was generated using a QuantaMaster™-6/2003 LPS-220B spectrofluorometer (PTI-Horiba, Japan, Supplementary Fig. S1). For ratiometric spectrofluorometry of cell lines genetically expressing mito-gTEMP, cell preparation and collection were carried out similarly, with the emission of both fluorophores and their ratio recorded over time, with excitation at 360 nm and emission for Sirius at 425 nm and for mT-Sapphire at 509 nm.

Ratiometric temperature calibration for mito-gTEMP *in vivo*

Cells were prepared as described above and, once resuspended in the cuvette, were treated with 5 µM oligomycin to block ATP synthesis, hydrolysis and respiratory electron transfer. As soon as the fluorescence ratio reached a steady-state, the temperature of the Peltier jacket holding the cuvette was shifted to 30 °C and then in 3 °C steps to 42 °C, in each case recording the fluorescence ratio during the third minute. The experiment was repeated (n=9) and the ratios of mT-Sapphire/Sirius fluorescence were plotted against temperature to generate a linear calibration function.

Respirometry

In order to determine the exact concentrations of inhibitors required for complete inhibition of each OXPHOS complex, we used the same amount of cells as used for spectrofluorometry to measure oxygen consumption with a Clark-type electrode (Oxytherm system, Hansatech, UK). Intact cell respiration was recorded from cells suspended in 500 µL of Respiraton Buffer A (225 mM Sucrose, 75 mM mannitol, 10 mM Tris/HCl, 10 mM KCl, 10 mM KH_2PO_4, 5 mM $MgCl_2$, pH 7.4) plus 10 mg/mL freshly added BSA, permeabilized by the addition of 80 µg/mL digitonin at 37 °C. Substrate concentrations were as follows; 10 mM ADP, 5 mM pyruvate and malate (cI+cIII), 10 mM succinate (cII+cIII), 50 mM TMPD/1 mM ascorbate (cIV). The required concentrations for full inhibition were determined to be 3 µM rotenone (cI inhibitor), 1 µM antimycin (cIII inhibitor), 0.8 mM potassium cyanide (KCN, cIV inhibitor), 5 µM oligomycin (cV inhibitor), and 100 µM n-propyl gallate (AOX inhibitor).

Thermo-profiling of mitochondria in cells treated with OXPHOS inhibitors

Cells were prepared for spectrofluorometry, and inhibitors were added to the above concentrations, once fluorescence or fluorescence ratio had reached a steady state, generally 10 min after initial oxygenation. Fluorescence values were then studied over time, until a new equilibrium value was reached, but only until 35 min for mito-gTEMP. MTY fluorescence was calibrated against temperature in each experiment, by raising and lowering the Peltier by 3 ºC at the end of the experiment, and recording the fluorescence values reached. The change in temperature was then estimated using this internal calibration. For estimating temperature changes using mito-gTEMP, the calibration function described above was applied to the actual values obtained for the mT-Sapphire/Sirius ratio. The adenine nucleotide transporter (ANT) inhibitor Bongkrekic acid (Sigma,

B-6179) and ALR/Evr1 inhibitor MitoBloCK-6 (Calbiochem, Sigma-5.05759) were used similarly at 100 µM.

Cytoplasmic protein synthesis inhibition

Cells were seeded and, after 24 h, the medium was replaced with otherwise complete medium without penicillin or streptomycin, but with 150 µM anisomycin (A-9789, Sigma). Cells were incubated for various times up to 18 h. For measuring the mito-gTEMP fluorescence ratio, cells were then treated as for control (untreated) cells expressing mito-gTEMP, except that for all time-points except that at t=0, the cells were resuspended in PBS containing 150 µM anisomycin. For measuring MTY fluorescence in anisomycin-treated cells, 100 nM MTY was added 15 min prior to each time point, followed by resuspension in PBS containing 150 µM anisomycin and the addition of OXPHOS inhibitors as for control (untreated cells)

Microscopy

Cells were seeded on glass bottom dishes (35 mm MatTek, USA) and grown for 24 h in standard growth media at 37 °C, 5% CO_2. The culture medium was replaced with prewarmed medium containing fluorescent dyes, namely 100 nM MTY and/or 100 nM MitoTracker™ Deep Red FM (ThermoFisher Scientific). After 15 min incubation, the staining medium was removed, and cells washed once with fresh, prewarmed PBS then kept in PBS at 37 °C until analyses were complete. The expression and localization of mT-Sapphire in transgenic cell lines was imaged by spinning disc (X-light V2, CrestOptics) confocal fluorescence microscopy (Nikon FN1), with excitation at 360 nm (Cool-Led pE-400 light source) and emission filter 525/50 nm, with Nikon 60X/1.0 dip objective (CFI Apochromat NIR 60X W). Laser excitation light intensity was adjusted to minimize photobleaching. Fluorescence images for MitoTracker™ Deep Red FM (excitation 633 nm, emission 700/75 nm) were generated similarly. Live-cell and time-lapse images of cells stained

with MitoTracker^{TM} Deep Red FM or MTY (excitation 542 nm, emission 562 nm) were captured by laser-scanning microscopy (Nikon A1R), using a Nikon 60x/1.27 water-immersion (CFI SR Plan Apo IR 60XC WI) objective. The detector sensitivity was adjusted for each sample to optimize the image brightness and to avoid saturation. The Nikon A1R with N-SIM laser scanning confocal microscopy system includes a live-cell supporting chamber where CO_2, temperature of the culture dish (surrounded by heated-water supply), and temperature of objective in use can be controlled. Time-lapse imaging was performed on cells maintained at 37 °C with one image /10sec for 10 mins, with a 2 min pause for addition of oligomycin to 5 µM, followed by a further 30 min of image capture. Signal analyses of time-lapse images were performed with ImageJ (Fuji), by surface-plot analysis.

Statistics

All data are presented as means ± SD. Statistical significance was calculated by standard unpaired one-way ANOVA with Tukey HSD test (www.astatsa.com) or, where appropriate, by Student's t test (LibreOffice Calc). Values of $p < 0.05$ were considered significant.

Image presentation

Micrographic images and fluorescence traces were cropped, rescaled and optimized for brightness and contrast, with labels added or removed, but with no other modifications.

FUNDING

This work was supported by Academy of Finland (award 324730 to HTJ and TSS, awards 322732 and 328969 to TSS), Basic Science Research Institute Fund (NRF grant 2021R1A6A1A10042944 to Y-TC), the Sigrid Juselius Foundation (grant 3122800849 to TSS), and by a grant from Core

Research for Evolutionary Science and Technology, Japan Science and Technology Agency (JPMJCR15N3 to TN).

ACKNOWLEDGMENTS

We thank Malgorzata Rak and Eric Dufour for useful discussions, Tea Tuomela for technical assistance, Maria Carretero-Junquera for artwork and the Biocenter Finland-supported Tampere Imaging Facility for access to its infrastructure.

AUTHOR CONTRIBUTIONS

M. Terzioglu = Conceptualization; Methodology; Validation; Formal analysis; Investigation; Resources; Data curation; Writing – original draft preparation; Writing – review & editing; Visualisation; Supervision

K. Veeroja = Conceptualization; Methodology; Validation; Investigation; Writing – review & editing

T. Montonen = Methodology; Investigation; Resources; Writing – review & editing; Visualisation

Teemu O. Ihalainen = Conceptualization; Methodology; Resources; Writing – review & editing; Visualisation; Supervision

T. S. Salminen = Conceptualization; Resources; Writing – review & editing; Supervision; Project administration; Funding acquisition

P. Bénit = Conceptualization; Methodology; Writing – review & editing

P. Rustin = Conceptualization; Methodology; Writing – review & editing

Y.-T. Chang = Conceptualization; Methodology; Resources; Writing – review & editing; Project administration; Funding acquisition

T. Nagai = Conceptualization; Methodology; Resources; Writing – review & editing; Project administration; Funding acquisition

H. T. Jacobs = Conceptualization; Methodology; Validation; Formal analysis; Resources; Writing –

original draft preparation; Visualisation; Supervision; Project administration; Funding acquisition

LIST OF SUPPLEMENTARY FILES

temp SI.pdf – 7 supplementary figures and their legends

Suppl movie S1.avi – supplementary movie

REFERENCES

1. Kang, J. S. Theoretical model and characteristics of mitochondrial thermogenesis. *Biophys. Rep.* 4, 63–67 (2018).

2. Brand, M. D. The efficiency and plasticity of mitochondrial energy transduction. *Biochem Soc. Trans.* 33, 897–904 (2005).

3. Nath, S. The thermodynamic efficiency of ATP synthesis in oxidative phosphorylation. *Biophys Chem.* 219, 69–74 (2016).

4. Brand, M. D., Chien, L. F., Ainscow, E. K., Rolfe, D. F. & Porter, R. K. The causes and functions of mitochondrial proton leak. *Biochim. Biophys. Acta* 1187, 132–139 (1994).

5. Enerbäck, S., Jacobsson, A., Simpson, E. M., Guerra, C., Yamashita, H., Harper, M. E. & Kozak, L. P. Mice lacking mitochondrial uncoupling protein are cold-sensitive but not obese. *Nature* 387, 90–94 (1997).

6. Chrétien, D., Bénit, P., Ha, H. H., Keipert, S., El-Khoury, R., Chang, Y. T., Jastroch, M., Jacobs, H. T., Rustin, P. & Rak, M. Mitochondria are physiologically maintained at close to 50 °C. *PLoS Biol.* 16, e2003992 (2018)

7. Nakano, M., Arai, Y., Kotera, I., Okabe, K., Kamei, Y. & Nagai, T. Genetically encoded ratiometric fluorescent thermometer with wide range and rapid response. *PLoS One* 12, e0172344 (2017).

8. Kriszt, R., Arai, S., Itoh, H., Lee, M. H., Goralczyk, A. G., Ang, X. M., Cypess, A. M., White, A. P., Shamsi, F., Xue, R., Lee, J. Y., Lee, S. C., Hou, Y., Kitaguchi, T., Sudhaharan, T., Ishiwata, S., Lane, E. B., Chang, Y. T., Tseng, Y. H., Suzuki, M. & Raghunath, M. Optical visualisation of thermogenesis in stimulated single-cell brown adipocytes. *Sci. Rep.* 7, 1383 (2017).

9. Okabe, K., Inada, N., Gota, C., Harada, Y., Funatsu, T. & Uchiyama, S. Intracellular

temperature mapping with a fluorescent polymeric thermometer and fluorescence lifetime imaging microscopy. *Nat. Commun.* 3, 705 (2012).

10. Hayashi, T., Fukuda, N., Uchiyama, S. & Inada, N. A cell-permeable fluorescent polymeric thermometer for intracellular temperature mapping in mammalian cell lines. *PLoS One* 10, e0117677 (2015).

11. Liang, H., Yang, K., Yang, Y., Hong, Z., Li, S., Chen, Q., Li, J., Song, X. & Yang, H.A Lanthanide Upconversion Nanothermometer for Precise Temperature Mapping on Immune Cell Membrane. *Nano Lett*. 22, 9045–9053 (2022).

12. Tanimoto, R., Hiraiwa, T., Nakai, Y., Shindo, Y., Oka, K., Hiroi, N. & Funahashi, A. Detection of Temperature Difference in Neuronal Cells. *Sci. Rep.* 6, 22071 (2016).

13. Lane, N. Hot mitochondria? *PLoS Biol.* 16, e2005113 (2018).

14. Dabir, D. V., Hasson, S. A., Setoguchi, K., Johnson, M. E., Wongkongkathep, P., Douglas, C. J., Zimmerman, J., Damoiseaux, R., Teitell, M. A. & Koehler, C. M. A small molecule inhibitor of redox-regulated protein translocation into mitochondria. *Dev. Cell* 25, 81–92 (2013).

15. Marroquin, L. D., Hynes, J., Dykens, J. A., Jamieson, J. D. & Will, Y. Circumventing the Crabtree effect: replacing media glucose with galactose increases susceptibility of HepG2 cells to mitochondrial toxicants. *Toxicological Sci.* 97, 539–547 (2007)

16. Pilbrough, W., Munro, T. P. & Gray, P. Intraclonal protein expression heterogeneity in recombinant CHO cells. *PLoS One* 4, e8432 (2009).

17. Roti Roti, J. L. Cellular responses to hyperthermia (40-46 degrees C): cell killing and molecular events. *Int. J. Hyperthermia* 24, 3–15.

18. Henderson, P. J. & Lardy, H. A. Bongkrekic acid. An inhibitor of the adenine nucleotide translocase of mitochondria. *J. Biol. Chem.* 245, 1319–1326 (1970).

19. Kano, A., Iwasaki, T. & Shindo, M. Bongkrekic acid facilitates glycolysis in cultured cells

and induces cell death under low glucose conditions. *Biochem. Biophys. Rep.* 20, 100683 (2019).

20. Dassa, E. P., Dufour, E., Gonçalves, S., Paupe, V., Hakkaart, G. A., Jacobs, H. T, & Rustin, P. Expression of the alternative oxidase complements cytochrome c oxidase deficiency in human cells. *EMBO Mol. Med.* 1, 30–36 (2009).

21. Szibor, M., Gainutdinov, T., Fernandez-Vizarra, E., Dufour, E., Gizatullina, Z., Debska-Vielhaber, G., Heidler, J., Wittig, I., Viscomi, C., Gellerich, F. & Moore, A. L. Bioenergetic consequences from xenotopic expression of a tunicate AOX in mouse mitochondria: Switch from RET and ROS to FET. *Biochim. Biophys. Acta Bioenerg.* 1861, 148137 (2020).

22. Chrétien, D., Bénit, P., Leroy, C., El-Khoury, R., Park, S., Lee, J. Y., Chang. Y.-T., Lenaers, G., Rustin, P. & Rak, M. Pitfalls in Monitoring Mitochondrial Temperature Using Charged Thermosensitive Fluorophores. *Chemosensors* 8, 124 (2020).

23. Andjelković, A., Mordas, A., Bruinsma, L., Ketola, A., Cannino, G., Giordano, L., Dhandapani, P. K., Szibor, M., Dufour, E. & Jacobs, H. T. Expression of the Alternative Oxidase Influences Jun N-Terminal Kinase Signaling and Cell Migration. *Mol. Cell. Biol.* 38, e00110-18 (2018).

24. Giordano, L., Farnham, A., Dhandapani, P. K., Salminen, L., Bhaskaran, J., Voswinckel, R., Rauschkolb, P., Scheibe, S., Sommer, N., Beisswenger, C., Weissmann, N., Braun, T., Jacobs, H. T., Bals, R., Herr, C. & Szibor, M. Alternative Oxidase Attenuates Cigarette Smoke-induced Lung Dysfunction and Tissue Damage. *Am. J. Respir. Cell Mol. Biol.* 60, 515–522 (2019).

25. Lu, K., Wazawa, T., Sakamoto, J., Vu, C. Q., Nakano, M., Kamei, Y. & Nagai, T. Intracellular Heat Transfer and Thermal Property Revealed by Kilohertz Temperature Imaging with a Genetically Encoded Nanothermometer. *Nano Lett.* 22, 5698–5707 (2022).

26. Baffou, G., Rigneault, H., Marguet, D. & Jullien, L. A critique of methods for temperature

imaging in single cells. *Nat. Methods* 11, 899–901 (2014).

27. Moreno-Loshuertos, R., Marco-Brualla, J., Meade, P., Soler-Agesta, R., Enriquez, J. A. & Fernández-Silva, P. How hot can mitochondria be? Incubation at temperatures above 43 °C induces the degradation of respiratory complexes and supercomplexes in intact cells and isolated mitochondria. *Mitochondrion* 69, 83–94 (2023).

28. Piñol, R., Zeler, J., Brites, C. D. S., Gu, Y., Téllez, P., Carneiro Neto, A. N., da Silva, T. E., Moreno-Loshuertos, R., Fernandez-Silva, P., Gallego, A. I., Martinez-Lostao, L., Martínez, A., Carlos, L. D. & Millán, A. Real-Time Intracellular Temperature Imaging Using Lanthanide-Bearing Polymeric Micelles. *Nano Lett.* 20, 6466–6472 (2020).

29. Savchuk, O. A., Silvestre, O. F., Adão, R. M. R. & Nieder, J. B. GFP fluorescence peak fraction analysis based nanothermometer for the assessment of exothermal mitochondria activity in live cells. *Sci. Rep.* 9, 7535 (2019).

30. Jarzab, A., Kurzawa, N., Hopf, T., Moerch, M., Zecha, J., Leijten, N., Bian, Y., Musiol, E., Maschberger, M., Stoehr, G., Becher, I., Daly, C., Samaras, P., Mergner, J., Spanier, B., Angelov, A., Werner, T., Bantscheff, M., Wilhelm, M., Klingenspor, M., Lemeer, S., Liebl, W., Hahne, H., Savitski. M. M. & Kuster, B. Meltome atlas-thermal proteome stability across the tree of life. *Nat. Methods.* 17, 495–503 (2020).

31. McDonald, A. E. & Gospodaryov, D. V. Alternative NAD(P)H dehydrogenase and alternative oxidase: Proposed physiological roles in animals. *Mitochondrion* 45, 7–17 (2019).

32. Di, X., Wang, D., Zhou, J., Zhang, L., Stenzel, M. H., Su, Q. P. & Jin, D. Quantitatively Monitoring *In Situ* Mitochondrial Thermal Dynamics by Upconversion Nanoparticles. *Nano Lett.* 21, 1651–1658 (2021).

33. Grollman, A. P. Inhibitors of protein biosynthesis. II. Mode of action of anisomycin. *J. Biol. Chem.* 242, 3226–3233 (1967).

34. Sidhu, J. S. & Omiecinski, C. J. Protein synthesis inhibitors exhibit a nonspecific effect on

phenobarbital-inducible cytochome P450 gene expression in primary rat hepatocytes. *J. Biol. Chem.* 273, 4769–4775 (1998).

35. Stouthamer, A. H. A theoretical study on the amount of ATP required for synthesis of microbial cell material. *Antonie Van Leeuwenhoek* 39, 545–565 (1973).

36. Pontes, M. H., Sevostyanova, A. & Groisman, E. A. When Too Much ATP Is Bad for Protein Synthesis. *J. Mol. Biol.* 427, 2586–2594 (2015).

37. Siems, W. G., Schmidt, H., Gruner, S. & Jakstadt, M. Balancing of energy-consuming processes of K 562 cells. *Cell Biochem. Funct.* 10, 61–66 (1992).

38. Rolfe, D.F. & Brown, G. C. Cellular energy utilization and molecular origin of standard metabolic rate in mammals. *Physiol. Rev.* 77, 731–758 (1997).

39. Rahbani, J. F., Roesler, A., Hussain, M. F., Samborska, B., Dykstra, C. B., Tsai, L., Jedrychowski, M. P., Vergnes, L., Reue, K., Spiegelman, B. M. & Kazak, L. Creatine kinase B controls futile creatine cycling in thermogenic fat. *Nature* 590, 480–485 (2021).

40. Fiermonte, G., De Leonardis, F., Todisco, S., Palmieri, L., Lasorsa, F. M, & Palmieri, F. Identification of the mitochondrial ATP-Mg/Pi transporter. Bacterial expression, reconstitution, functional characterization, and tissue distribution. *J. Biol. Chem.* 279, 30722–30730 (2004).

41. Partikian, A., Olveczky, B., Swaminathan, R., Li, Y. & Verkman, A. S. Rapid diffusion of green fluorescent protein in the mitochondrial matrix. *J. Cell. Biol.* 140, 821–829 (1998).

42. Arai, S., Suzuki, M., Park, S. J., Yoo, J. S., Wang, L., Kang, N. Y., Ha, H. H. & Chang, Y. T. Mitochondria-targeted fluorescent thermometer monitors intracellular temperature gradient. *Chem. Commun.* 51, 8044–8047 (2015).

43. Das, R. & Chakrabarti, O. Mitochondrial hyperfusion: a friend or a foe. *Biochem. Soc. Trans.* 48, 631–644 (2020).

44. Gottlieb, R. A., Piplani, H., Sin, J., Sawaged, S., Hamid, S. M., Taylor, D. J. & de Freitas

Germano, J. At the heart of mitochondrial quality control: many roads to the top. *Cell Mol. Life. Sci.*78, 3791–3801 (2021).

45. Seo, A. Y., Joseph, A. M., Dutta, D., Hwang, J. C., Aris, J. P. & Leeuwenburgh, C. New insights into the role of mitochondria in aging: mitochondrial dynamics and more. *J. Cell Sci.* 123, 2533–2542 (2010).

46. van der Stel, W., Yang, H., le Dévédec, S. E., van de Water, B., Beltman, J. B. & Danen, E. H. J. High-content high-throughput imaging reveals distinct connections between mitochondrial morphology and functionality for OXPHOS complex I, III, and V inhibitors. *Cell Biol. Toxicol.* https://doi.org/10.1007/s10565-022-09712-6 (2022).

47. Legros, F., Lombès, A., Frachon, P. & Rojo, M. Mitochondrial fusion in human cells is efficient, requires the inner membrane potential, and is mediated by mitofusins. *Mol. Biol. Cell* 13, 4343–4354 (2002).

48. Westrate, L. M., Drocco, J. A., Martin, K. R., Hlavacek, W. S. & MacKeigan, J. P. Mitochondrial morphological features are associated with fission and fusion events. *PLoS One* 9, e95265 (2014).

49. Abdullah, M. O., Zeng, R. X., Margerum, C. L., Papadopoli, D., Monnin, C., Punter, K. B., Chu, C., Al-Rofaidi, M., Al-Tannak, N. F., Berardi, D., Rattray, Z., Rattray, N. J. W., Abraham, S. A., Eskelinen, E. L., Watson, D. G., Avizonis, D., Topisirovic, I. & Chan, E. Y. W. Mitochondrial hyperfusion via metabolic sensing of regulatory amino acids. *Cell Rep.* 40, 111198 (2022).

50. Kucej, M. & Butow, R. A. Evolutionary tinkering with mitochondrial nucleoids. *Trends Cell Biol.* 17, 586–592 (2007).

51. Rango, M., Arighi, A., Bonifati, C., Del Bo, R., Comi, G. & Bresolin, N. The brain is hypothermic in patients with mitochondrial diseases. *J. Cereb. Blood Flow Metab.* 34, 915–920 (2014).

52. El-Gammal, Z., Nasr, M. A., Elmehrath, A. O., Salah, R. A., Saad, S. M. & El-Badri, N. Regulation of mitochondrial temperature in health and disease. *Pflugers Arch.* 474, 1043–1051 (2022).

53. Lebkowski, J. S., Clancy, S. & Calos, M. P. Simian virus 40 replication in adenovirus-transformed human cells antagonizes gene expression. *Nature* 317, 169–171 (1985).

54. Pontén, J. & Saksela, E. Two established *in vitro* cell lines from human mesenchymal tumours. *Int. J. Cancer* 2, 434–447 (1967).

55. Schneider, I. Cell lines derived from late embryonic stages of *Drosophila melanogaster*. *J. Embryol. Exp. Morphol.* 27, 353–365 (1972).

56. Donnelly, M. L. L., Hughes, L. E., Luke, G., Mendoza, H., Ten Dam, E., Gani, D. & Ryan, M. D. The 'cleavage' activities of foot-and-mouth disease virus 2A site-directed mutants and naturally occurring '2A-like' sequences. *J. Gen. Virol.* 82, 1027-1041 (2001).

57. Donnelly, M. L. L., Luke, G., Mehrotra, A., Li, X., Hughes, L. E., Gani, D. & Ryan, M. D. Analysis of the aphthovirus 2A/2B polyprotein 'cleavage' mechanism indicates not a proteolytic reaction, but a novel translational effect: a putative ribosomal 'skip'. *J. Gen. Virol.* 82, 1013–1025 (2001).

Figure 1

The mitochondrial OXPHOS system and inhibitors

Summary of the major components of the OXPHOS system and classic inhibitors. Protonmotive

OXPHOS enzyme complexes (cI, cIII, cIV, cV) shown in green, the non-protonmotive

transgenically introduced alternative oxidase (AOX) from *C. intestinalis* in orange, the inner

mitochondrial membrane in grey. Ions and small molecules indicated in purple, inhibitors in brick-

red. BKA – bongkrekic acid, an inhibitor of the adenine nucleotide translocase (ANT). nPG – *n*-

propyl gallate, an inhibitor of AOX, Q – ubiquinone (oxidized coenzyme Q), QH_2 – ubiquinol

(reduced coenzyme Q), c_{red}, c_{ox} – reduced and oxidized forms of cytochrome *c*, respectively.

Complex I (cI), along with a number of flavin-linked dehydrogenases, one of which is succinate

dehydrogenase, also described as complex II, each reduce ubiquinone, thus contributing input

electrons to complex III (cIII). ALR – 'augmenter of liver regeneration', Evr1 in yeast, acts as an

additional feeder of electrons to cytochrome *c*, from the oxidation of sulfhydryl groups in proteins

destined for the mitochondrial intermembrane space, in the Mia40 pathway and is inhibited by

MitoBLloCK-6.

Figure 2

OXPHOS inhibitors decrease mitochondrial temperature to different extents

Extrapolated mitochondrial temperature shifts (means $\pm$ SD for at least 5 independent experiments

in each case), based on MTY fluorescence in different cell-lines treated with the indicated

OXPHOS inhibitors, with internal temperature calibration at the end of each experiment. *, **, ***

denote significantly different data sets (one-way ANOVA with Tukey HSD *post-hoc* test, $p < 0.05$,

0.01 and 0.001, respectively). Note that a stable reading could not be obtained for S2 cells treated

with oligomycin (Fig. S3B), hence reliable calibration was not possible. iMEF(L7) and iMEF(P) are two isolates of immortalized, embryonic fibroblasts derived from wild-type C57Bl/6J mice.

Figure 3

Ratiometric fluorescence probes confirm mitochondrial temperature measurements by MTY

(A) Diagrammatic map (not to scale) of the self-cleaving polyprotein encoded by plasmid mito-gTEMP (coxVIIIx2-gTEMP)/pcDNA3, indicating the reiterated COX8 targeting presequences, the Sirius and mT-Sapphire coding regions and the T2A element[56], which promotes self-cleavage via ribosome frameshifting[57]. (B) Illustrative mT-Sapphire fluorescence image of iMEF cell clone expressing mito-gTEMP, showing variable expression even within cells of the clone. Scale bar 5 μm. See Fig. S4A for confirmation of mitochondrial localization. (C) Illustrative trace of mT-Sapphire/Sirius fluorescence ratio in HEK293T cell clone expressing mito-gTEMP, following treatment with oligomycin. Note that, as for MTY, Sirius fluorescence decreases with increasing temperature, so that the decreased mT-Sapphire/Sirius fluorescence ratio after oligomycin treatment represents a decrease in temperature. (D) Calibration curve for mT-Sapphire/Sirius fluorescence ratio (means $\pm$ SD, 9 independent experiments in each case) in mito-gTEMP expressing iMEFs treated with oligomycin, and shifted to various temperatures up to 42 ºC. Solid red line represents linear best fit (as per equation shown) extrapolated to higher temperatures (dotted red line) that were not directly tested due to the effects of prolonged high temperatures on cell integrity and viability[17]. Note that, during these trials, we observed that continuous illumination of mito-gTEMP expressing cells at the excitation wavelength for more than 40 min caused damage affecting the fluorescence signal. All calibration measurements were therefore conducted over < 35 min. We also noted that the internal temperature of the cuvette took ~2 minutes to equilibrate with that of the surrounding Peltier (Fig. S2), for which reason fluorescence values were measured over just the final minute at each calibration step. (E) Extrapolated mitochondrial temperature shifts (means $\pm$

SD for at least 5 independent experiments in each case) in the indicated cell-lines and mito-gTEMP transfected cell clones following treatment with oligomycin, based on on (red bars) MTY fluorescence, reproduced from corresponding panels of Fig. 2, and (blue bars) mito-gTEMP fluorescence ratio, applying calibration curves such as shown in panel (C), derived separately for each cell line studied. ** denotes significant differences between the two methods (Student's t test within each cell-line, $p < 0.01$).

Figure 4

Mitochondrial temperature variation between and within cells

(A, B) Micrographic images of iMEF cells stained with MTY (A, panel i and B), alongside a parallel culture of cells stained with Mitotracker™ Deep Red FM (A, panel ii), which shows much more uniform staining. The intensity of MTY staining appears to vary between and even within cells. Scale bars – 10 μm. (B) Still image from Supplementary Movie S1, brightness and contrast optimized, showing bright (i.e., potentially cooler) foci (arrowed) within the less intensely stained mitochondrial network. Many of these foci appear mobile within the mitochondrial network, over the 5 min time-scale of the movie. Scale bar – 2 μm. (C) Fluorescence Intensity histograms of a field of cells immediately before and 33 min after oligomycin treatment. Note that, in addition to the general pattern of increased brightness (e.g., peaks represented by red dots), mitochondria in some cells showed only minor changes in fluorescence intensity (e.g., peaks denoted by blue dots). Scale bars – 50 μm.

Figure 5

Mitochondrial temperature fluctuates in response to substrate availability

(A, C, D, E) Representative fluorescence ratio traces of mito-gTEMP expressing iMEF(P) cells resuspended in various media without OXPHOS inhibitors. PBS – phosphate-buffered saline, glc –

glucose and gln – glutamine at standard cell-culture concentrations, BKA – bongkrekic acid.

Arrows indicate time of addition of substrates or of BKA. Mitochondrial temperatures (°C)

extrapolated from calibration curve (Fig. 3D) shown in red boxes. (B) Mitochondrial temperatures

(°C, means $\pm$ SD of 5 independent experiments), averaged over the time intervals since cell

resuspension as shown, extrapolated from mito-gTEMP fluorescence ratios using calibration curve

(Fig. 3D), for cells with and without substrate additions as shown. *, **, *** denote statistical

significance (p < 0.05, 0.01, 0.001, respectively, one-way ANOVA with Tukey *post-hoc* HSD test).

(F) Mitochondrial temperatures (°C, means $\pm$ SD, n = 3 independent experiments), averaged over

the time intervals since cell resuspension as shown, extrapolated from mito-gTEMP fluorescence

ratios using calibration curve (Fig. 3D), for cells treated with BKA at 11 min (values averaged over

11-16 min) and then tracked until 35 min (final values averaged over 30-35 min). *, **, *** denote

statistical significance (p < 0.05, 0.01, 0.001, respectively, one-way ANOVA with Tukey *post-hoc*

HSD test).

Figure 6

Restoration of high mitochondrial temperature in AOX-expressing cells after OXPHOS

inhibition

(A) Representative traces of MTY fluorescence over time, in control and AOX-expressing iMEFs

treated with the OXPHOS inhibitors shown. Vertical axes are arbitrary values, horizontal axes

scaled as indicated. Note that increased fluorescence indicates cooler mitochondria. Control

iMEF(P) cells reached and maintained plateau levels of fluorescence, indicating a sustained

decrease in mitochondrial temperature, whilst iMEF(AOX) cells showed rewarming after reaching

a peak fluorescence value. (B) Peak mitochondrial temperature decrease (means $\pm$ SD, n $\geq$ 4

independent experiments in the indicated cell-lines following treatment with OXPHOS inhibitors,

inferred from MTY fluorescence or mito-gTEMP fluorescence ratio as shown. Horizontal bars

indicate significant differences for pairwise comparisons between iMEF(P) and iMEF(AOX) cells treated with the indicated inhibitors (MTY data, Student's t test, $p < 0.05$, 0.01 and 0.001, denoted, respectively, by *, ** and ***). There was no significant difference between the methods, applied to rotenone-treated iMEF(AOX) cells (Student's t test, $p > 0.05$). (C) Mitochondrial rewarming (means $\pm$ SD, $n \geq 5$ independent experiments except at latest time-points, where $n = 3$ or 4) in iMEF(AOX) cells treated with the inhibitors shown, inferred from MTY fluorescence traces.

Figure 7

Cells maintain high mitochondrial temperature despite varying ATP demand

(A, B) Intramitochondrial temperatures (means $\pm$ SD) estimated by mito-gTEMP fluorescence ratio in iMEF(P) and iMEF(AOX) cells as indicated, treated with 150 μM anisomycin for the times shown. Temperatures extrapolated from calibration curve shown in Fig. 3D. Horizontal bars denote data points that were significantly different (one-way ANOVA with Tukey HSD test, $p < 0.05$). (C) Extrapolated mitochondrial temperature shifts (means $\pm$ SD for at least 4 independent experiments in each case) based on MTY fluorescence, as indicated, in iMEF(P) and iMEF(AOX) cells pre-treated with 150 μM anisomycin for the times shown, followed by treatment with rotenone. Zero-time values (i.e. cells not treated with anisomycin) are reproduced from Figs. 2D and 5B for iMEF(P) and iMEF(AOX) cells, respectively. Values for the two cell lines are significantly different at each time-point except 0 and 4 h (Student's t test, $p < 0.05$). Within each cell-line, in addition to the clear downward trend with time (i.e., lesser amount of mitochondrial temperature decrease), the values for the extreme time points (0 and 18 h) are significantly different from each other and from all other time-points for the iMEF(P) line (denoted by asterisks), whilst for the iMEF(AOX) line significant pairwise differences are indicated by horizontal lines above the graphed values (one-way ANOVA with Tukey *post-hoc* HSD test, $p > 0.05$). (D) Extrapolated mitochondrial temperature shifts (means $\pm$ SD for at least 4 independent experiments in each case)

based on MTY fluorescence, as indicated, in iMEF(P) and iMEF(AOX) cells pre-treated with 150 μM anisomycin for the times shown, followed by treatment with oligomycin. Zero-time values (i.e. cells not treated with anisomycin) are reproduced from Figs. 2D and 5B for iMEF(P) and iMEF(AOX) cells, respectively. Values for the two cell lines are significantly different at each time-point (Student's t test, $p < 0.05$). Within each cell-line, and despite the apparent trend of the means for iMEF(AOX) cells, the values for different time points are, in general, not significantly different from each other (one-way ANOVA with Tukey *post-hoc* HSD test, $p > 0.05$). Exceptions for iMEF(P) cells are indicated by horizontal lines ($p < 0.01$).

Terzioglu et al, Figure 1

Terzioglu et al, Figure 2

A

Bam H1   EcoR1

☐ COX8 presequence

■ Sirius

■ T2A

■ mT-Sapphire

B

iMEF (P)
mito-gTEMP clone 1

C

HEK293T
mito-gTEMP clone 1

fluorescence ratio

oligomycin

time (min)

D

fluorescence ratio

y = 0.0609x + 6.611

temperature (ºC)

E

oligomycin   rotenone

−ΔTemperature (ºC)

HEK293T   U2OS

iMEF (P)

■ MTY

■ mito-gTEMP

Terzioglu et al, Figure 3

A

MTY

MitoTracker™ Deep Red

B

C

4095.0

4095.0

0.0

0.0

151.8 microns

151.8 microns

151.8 microns

151.8 microns

before oligomycin

after oligomycin

Terzioglu et al, Figure 4

Terzioglu et al, Figure 5

Terzioglu et al, Figure 6

Terzioglu et al, Figure 7

Mitochondrial temperature homeostasis

resists external metabolic stresses

Mügen Terzioglu *et al*

SUPPLEMENTARY INFORMATION

- Supplementary Figure Legends

- Supplementary Movie Legend (movie in separate file, Suppl Movie S1.avi)

- Supplementary Figures

LEGENDS TO SUPPLEMENTARY FIGURES

Supplementary Figure S1

Instrumentation for mitochondrial temperature estimation using fluorescent probes

Schematic illustration of spectrophotometer, QuantaMaster™ QM-6/2003 LPS-220B with a Peltier temperature-controlled sample cell chamber. Waterbath (Julabo® CORIO™ CD, Germany) with refrigerated/heated circulator (-20 to 150 ℃) provides accurate temperature range via circulation around the sample cell chamber. The xenon lamp (Xenon short arc UXL 75XE, USHIO, Japan) used as the light excitation source. The excitation monochromator selects the desired wavelength of excitation light, which is focused at the sample position. The emission monochromator is scanned across the desired emission range and fluorescence intensity is recorded in the detector as a function of emission wavelength. While data is recorded, the cell sample is stirred continuously, maintaining oxygen concentration far above the level required for fully active cellular respiration. For instrument validation see Fig. S2. Illustration by Maria Carretero-Junquera.

Supplementary Figure S2

Instrument validation steps for estimation of mitochondrial temperature

(A) In an initial step, an external temperature probe was used to validate the temperature readings of components of the apparatus (water-bath, Peltier, cuvette) as shown on the digital displays. After a 3 ℃ temperature shift, the Peltier took ~2 min to stabilize at the new temperature, and held steady despite minor fluctuations in the external room temperature. (B) The cuvette temperature was not affected by water-bath temperature changes or differences from the set temperature of the Peltier. The water bath temperature (not shown) itself took ~5-6 min to restabilize. Note that the cuvette temperature in steady-state was 1 ℃ lower than that of the Peltier, which was taken account of in the calibration shown in Fig. 3D. (C) Sample traces showing calibration steps after fluorescence of

the indicated inhibitor-treated cells had reached a steady value. Vertical axes are arbitrary values, horizontal axes (time) rescaled for uniformity. In each case fluorescence reached a new equilibrium value rapidly after the externally applied shifts in temperature, as indicated (ºC).

Supplementary Figure S3

Supplementary data on effects of OXPHOS inhibitors on mitochondrial temperature

(A) Representative fluorescence traces, uniformly rescaled, for MTY-labelled iMEF(P) cells treated with the indicated combinations of OXPHOS inhibitors. Vertical scale (not shown): arbitrary values to similar scale. Rotenone + antimycin (panel i) gave the same stable fluorescence reading as all individual inhibitors, whereas other combinations (panels ii-vi) showed prolonged and/or continuous decline of the fluorescence reading, which is likely be due to dye leakage upon loss of membrane potential, as shown previously[6]. For this reason we did not attempt to interpret findings from combinations of inhibitors other than rotenone *plus* antimycin (Fig. 2D). (B) Fluorescence trace for addition of 100 μM MitoBloCK-6 to MTY-stained cells. Note the immediate >20-fold increase in fluorescence intensity upon addition of the drug. (C) Extrapolated mitochondrial temperature shifts for iMEFs grown in different media and treated with rotenone. Data for lines iMEF(L7) and iMEF(P) grown in normal glucose (glc) medium are from the experiments shown in Fig. 2C and 2D, respectively. Gal – galactose. Lo glc – low-glucose medium (see Materials and Methods). Data are shown as scatter plots to indicate the distributions. There are no significant differences between the groups (one-way ANOVA with Tukey HSD *post-hoc* test). (C) Representative, uniformly scaled traces of MTY fluorescence in S2 (panel i) and U2OS (panel iii) cells treated with oligomycin and S2 cells treated with KCN (panel ii), as indicated. For clarity, vertical scale (arbitrary values) and temperature calibration are omitted. Fluorescence in oligomycin-treated S2 cells had not reached a stable value even 2 h after addition of the drug; therefore, no meaningful calibration could be conducted. Treatment of other cell-lines with

oligomycin or of S2 cells with other OXPHOS inhibitors did not produce this problem, as per the examples shown.

Supplementary Figure S4

Supplementary data on mitochondrially targeted temperature-sensitive fluorescent probes

(A) Fluorescence microscopy images of iMEF(P) cells transfected with mito-gTEMP and imaged as indicated, as described in Materials and Methods. Scale bars 10 μm. (B, C, D) Fluorescence traces (AU – arbitrary units) and (E, F) fluorescence ratios at the wavelengths used to measure mT-Sapphire and Sirius, in PBS at 38 °C, to which were added the indicated drugs (arrowed). (G) MTY fluorescence traces for cell-free PBS or iMEF(P) cells treated with bongkrekic acid (BKA) as indicated. The drug produced an immediate quenching of fluorescence in both cases. In cells, this was followed by a gradual regain of fluorescence, possibly indicative of mitochondrial temperature decrease by several ºC, based on the internal calibration (red boxes). See also Fig. 5E, 5F.

Supplementary Figure S5

AOX-expressing iMEFs show antimycin- and KCN-resistant respiration

Respirometric traces from digitonin-permeabilized cells of the indicated lines, treated successively with digitonin – dig, pyruvate+glutamate+malate substrate mix – S and ADP – A, followed by either (A) antimycin (ant) or (B) either one or two successive doses of KCN. For details see Methods. In each part, panels (i) and (ii) represent two independent trials. Relative rates of inhibitor-resistant oxygen consumption of iMEF (AOX) cells, based on the indicated slopes, and following subtraction of inhibitor-resistant oxygen consumption of control iMEF (P) cells, were as follows: (A) 70% and 67% antimycin resistance, (B) 50% and 47% KCN resistance.

Supplementary Figure S6

Supplementary data on effects of inhibitors on fluorescence in AOX-expressing cells

(A) Representative traces of mito-gTEMP fluorescence ratio for (I) control iMEF(P) and (ii) iMEF(AOX) cells, after rotenone delivery at 15 min until 35 min, with continuous recording (two replicate experiments in each case). Text in red boxes indicates inferred mitochondrial temperature (to nearest ºC) from calibration curve of Fig. 3D. AOX-expressing cells show a consistent warming of ~2 ºC by 20 min post-drug treatment, consistent with the MTY data of Fig. 5C. (B) Representative traces of MTY fluorescence for (i) control iMEF(P) cells and (ii) iMEF(AOX) cells treated with nPG (red arrows). Note that, prior to nPG addition, fluorescence in both cases had reached a stable level – in control cells after antimycin treatment and in AOX-expressing cells after rewarming, following KCN treatment. In both cases, nPG addition produced an abrupt quenching of the fluorescence, making its effect on mitochondrial temperature unquantifiable.

Supplementary Figure S7

Supplementary data on mitochondrial temperature decrease in anisomycin-treated cells

Representative fluorescence traces, uniformly rescaled, for (panels i, iii) MTY-labelled iMEF(AOX) and (panels ii, iv) iMEF(P)cells pretreated for 2 h with anisomycin, then incubated with (A) rotenone and (B) oligomycin.

LEGEND TO SUPPLEMENTARY MOVIE

Supplementary Movie S1

Mitochondrial temperature variation between and within cells

Time-lapse micrographic images (1 frame per 10 s) of iMEF cells stained with MTY, brightness and contrast optimized, showing bright (i.e., potentially cooler) foci within the less intensely stained mitochondrial network. Many of these foci appear mobile within the mitochondrial network, over the 5 min time-scale of the movie. See Fig. 4B for a still image taken from the movie.

Cuvette socket and Peltier jacket

Terzioglu et al, Figure S1

Terzioglu et al, Figure S2, page 1 of 2

C

HEK293

38   38   antimycin
41

25   25   rotenone, S2
30

38   38   KCN
41

38   38   rotenone, U2OS
41

38   38   oligomycin
41

38   38   rotenone, iMEF(P)
41

A

i   rotenone + antimycin

ii   rotenone + KCN

iii   rotenone + oligomycin

iv   antimycin + KCN

v   antimycin + oligomycin

vi   KCN + oligomycin

50 min

B

iMEF(P), MitoBloCK-6

fluorescence (a.u.)

C

iMEF, rotenone

−ΔTemperature (°C)

iMEF(P)
- normal glc
- lo glc + gal
- lo glc

iMEF(L7)
- normal glc
- lo glc + gal

from Fig. 2D   from Fig. 2C

D

i   S2, oligomycin

ii   S2, KCN

iii   U2OS, oligomycin

50 min

Terzioglu et al, Figure S3

A

mT-Sapphire

MitoTracker™ Deep Red

Merge

Terzioglu et al, Supplementary Figure S5

Terzioglu et al, Figure S6

A

iMEF(AOX)

iMEF(P)

i

↓ rotenone

ii

50 min

2 h anisomycin

B

iii

↓ oligomycin

iv

iMEF(AOX)

iMEF(P)

Terzioglu et al, Figure S7