Sex-biased regulatory changes in the placenta of native highlanders contribute to adaptive fetal development

Tian Yue[1,4#], Yongbo Guo[1,4#], Xuebin Qi[1,2,3#], Wangshan Zheng[1#], Hui Zhang[1,3#], Bin Wang[2], Kai Liu[1], Bin Zhou[1,4], Xuerui Zeng[1,4], Ouzhuluobu[2], Yaoxi He[1]*, Bing Su[1,5]*

[1]State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China;

[2]Fukang Obstetrics, Gynecology and Children Branch Hospital, Tibetan Fukang Hospital, Lhasa 850000, China;

[3]State Key Laboratory of Primate Biomedical Research, Institute of Primate Translational Medicine, Kunming University of Science and Technology, Kunming, China.

[4]Kunming College of Life Science, University of Chinese Academy of Sciences, Beijing 100101, China;

[5]Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming 650223, China;

[#]These authors contributed equally to this work.

Corresponding authors: Yaoxi He and Bing Su,

Email: heyaoxi@mail.kiz.ac.cn; sub@mail.kiz.ac.cn

Summary

Compared with lowlander migrants, native Tibetans have a higher reproductive

success at high altitude though the underlying mechanism remains unclear. Here, we

compared the transcriptome and histology of full-term placentas between native

Tibetans and Han migrants. We found that the placental trophoblast shows the largest

expression divergence between Tibetans and Han, and Tibetans show decreased

immune response and endoplasmic reticulum stress. Remarkably, we detected a

sex-biased expression divergence, where the male-infant placentas show a greater

between-population difference than the female-infant placentas. The umbilical cord

plays a key role in the sex-biased expression divergence, which is associated with the

higher birth weight of the male newborns of Tibetans. We also identified adaptive

histological changes in the male-infant placentas of Tibetans, including larger

umbilical artery wall and umbilical artery intima and media, and fewer syncytial knots.

These findings provide valuable insights into the sex-biased adaptation of human

populations, with significant implications for medical and genetic studies of human

reproduction.

Introduction

Hypobaric hypoxia at high altitude can restrict fetal growth, resulting in reduced neonatal birth weight (BW) and increased infant mortality (Moore et al., 2001b). Due to long-term natural selection at high altitude, native highlanders such as Tibetans have acquired a higher reproductive success than the lowlander migrants (e.g. Han Chinese moving to high altitude), reflected by the higher newborn BW and the lower prenatal and postnatal mortality (Beall et al., 2004; Moore et al., 2011). As expected, Tibetans performed better than Han migrants during fetal development (He et al., 2023; Julian et al., 2009; Moore et al., 2011; Moore et al., 2001a). Previous genomic studies have identified a group of genes potentially contributing to the genetic adaptation of Tibetans to high-altitude environments, and these studies were mostly focused on the blood and cardiopulmonary systems of adult Tibetans (Beall et al., 2010a; Bigham et al., 2010; Deng et al., 2019; Peng et al., 2017; Peng et al., 2011; Simonson et al., 2010; Song et al., 2020; Xiang et al., 2013; Xu et al., 2011; Yang et al., 2017; Yi et al., 2010). Although a recent study reported the association of more pregnancies and a higher survival birth rate based on GWAS analysis of 1,008 Nepali Tibetans, no adaptive genetic variant was identified (Jeong et al., 2018). Therefore, whether and how the Tibetan adaptive variants affect their fetal development and eventually improve the reproductive fitness are yet to be explored.

The placenta is a temporary organ that connects fetus to the mother's uterus during pregnancy, and it begins to develop around 6-7 days after gestation (Knofler et al., 2019). As an important organ of maternal-fetal exchange of nutrient, oxygen and waste, the placenta plays a critical role in fetal growth and development. Previous studies have reported that placental dysfunction can lead to premature birth, abnormal fetal growth and defects of neurological development (Guttmacher et al., 2014). At the same time, the genetic study reported that placental transcription in utero plays a key role in determining postnatal body size (Peng et al., 2018), and placenta weight is positively correlated with BW, an important trait related to the postnatal survival rate of newborns (Haeussner et al., 2013). Therefore, gene expression profiling of placenta

is essential to dissecting the molecular mechanism of the successful reproduction.

Recently, a transcriptomic study comparing 91 placental samples (including 47

Tibetans giving birth at high altitudes) found that the genes related to autophagy and

tricarboxylic acid cycle showed significant up-regulation in Tibetans. However, no

lowlander migrants living at high altitude (>2500m) were included in this study (Wu

et al., 2022).

In addition, it is known that the intrauterine growth rate of the male fetus is faster

than that of the female fetus, and this growth rate difference leads to a higher

proportion of preeclampsia, premature delivery and intrauterine growth restriction of

the male fetus (Clifton, 2010). The between-sex difference of fetal development was

explained by the higher transport efficiency, but weaker reserve capacity of the male

fetal placenta (Eriksson et al., 2010). Consistently, we recently reported a male-biased

pattern of BW seasonality in Tibetans and Han migrants living at high altitude.

Compared to the female fetuses, the male fetuses are more sensitive to hypobaric

hypoxia, reflected by their more serious BW reduction and relatively lower survival

rates (He et al., 2022). However, whether the placenta plays a key role in the observed

sex-biased sensitivity to hypoxia during fetal development at high altitude remains

elusive.

In this study, we sampled 69 full-term placental tissues, including 35 native

Tibetans and 34 Han migrants living at the same altitude (an elevation of 3,650m).

Seven tissue layers of the placenta were dissected and analyzed in detail. With the use

of RNA-seq, we generated a systematic map of placental transcriptomes of indigenous

and migrant populations living at high altitude. Markedly, we found a male-biased

transcriptomic divergence between Tibetans and Han migrants, and the gene

expression pattern of the fetal umbilical cord may affect the BW of the male

newborns. We also observed adaptive histological changes in the Tibetan male-infant

placentas, providing further evidence on the sex-biased adaptation of fetal

development.

Results

Gene expression profiles of placentas of native Tibetans and Han migrants

To understand the transcriptomic patterns of placentas of Tibetans and Han migrants at high altitude, we collected full-term (>37 weeks of gestation) placental samples from 35 Tibetans (16 male newborns and 19 female newborns) and 34 Han migrants (21 male newborns and 13 female newborns) with singleton birth. These Tibetan and Han migrant mothers live at the same high altitude (Lhasa, Tibet Autonomous Region, China; elevation = 3,650 m), and had experienced their entire pregnancy at this altitude. The population ancestry of the subjects was confirmed by self-claim, and further validated by the genome-sequencing data (**Methods, Figure S1**).

We first compared the newborn traits between the Tibetan group and the Han group. As expected, the average BW of the Tibetan newborns is significantly higher than that of the Han newborns (p=0.0121, unpaired Student's t-test), a clear indication of a better fetal development in Tibetans, while there are no significant differences in placental weight and placental volume (**Figure S2**). Given the placenta is a highly heterogeneous organs underlying functional and histological specializations in discrete anatomical parts (Sood et al., 2006), we dissected each placenta into seven tissue layers according to its anatomic structure (from fetal side to maternal side), including umbilical cord (UC), amnion (AN), chorion (CN), chorionic plate (CP), villus of fetal (VF), villus of intermediate (VI) and villus of maternal (VM), and we conducted RNA-seq of 483 placental samples (69 individuals × 7 layers) (**Figure 1A**), In total, ~12 billion short reads were generated. After applying a stringent quality control (**Methods**), we kept 448 placental transcriptome data, and each sample contains ~27 million reads on average (sample details in **Table S1**). In addition, we obtained the genome-wide variants of these mothers and their newborns from our previous studies (He et al., 2023; Zheng et al., 2023).

As the placenta is a maternal-fetal mosaic organ, we firstly determined the maternal-fetal compositions of each dissected layer using the genomic and

transcriptomic data (see **Methods** for details). The results show that UC and VF are

100% fetal origin, so are VM, VI, CP and AN (>95%) with tiny proportions of

maternal origin. By contrast, CN is mostly maternal origin (70%) (**Figure 1B**). This

pattern is consistent with the anatomic structure of the placenta (Sood et al., 2006).

Consequently, the RNA-seq data of placenta mostly represent the transcriptomes of

the newborns, except for the CN reflecting the mixed transcriptomes of mothers and

newborns.

The principal component analysis (PCA) indicates that the placental samples

cluster by layers instead of population groups, suggesting that the transcriptomic

difference between placental structures is greater than the between-population

divergence. This pattern is in line with previous transcriptome of three-layer placentas

(chorion, amnion and decidua) of Nepali Tibetans, in which the samples were grouped

by tissue type instead of ethnicity (Wu et al., 2022). Notably, the four trophoblast

layers (CP, VF, VI and VM) cannot be clearly separated in the PCA map and they

form a large cluster. These four layers belong to the chorionic villus, the essential

structure in placenta (Wapner and Jackson, 1988). The UC, AN and CN layers form

three separate clusters (**Figure 1C**). Accordingly, the gene expression heat-map

analysis generates the same clustering pattern (**Figure 1D**). These results suggest that

different layers of the placenta have distinct gene expression profiles, which can

provide detailed regulatory information in understanding the role of placenta in fetal

development.

VF and VI show the largest expression divergence between Tibetans and Han

To gain insights into the differences of gene expression in placenta between

Tibetans and Han, we performed differential gene expression analysis of the seven

placental layers (see **Methods** for technical details). In total, we identified 579

differentially expressed genes (DEGs) between Tibetans and Han, accounting for 3.4%

of the total number of expressed genes. Our results show that VF and VI have the

largest numbers of DEGs between Tibetans and Han (305 DEGs in VI and 173 DEGs

in VF), accounting for 82.56% (478/579) of the total DEGs in the placenta. By

contrast, the numbers of DEGs in the other five layers are much less (8-46 DEGs) (**Figure 2A, Table S2**). Notably, most of the DEGs (399/579, 68.91%) are detected in only one layer (**Figure 2A**), an indication of a layer-specific differential expression pattern. The results of GO enrichment analyses show that for VF and VI, and the top functional terms reflect an enhancement of mRNA regulation and a decrease of immune response (indicated by the terms of defense response to virus) in VF of Tibetans compared to Han (**Figure 2B, Table S3**). In VI, the significant functional terms (decreased expression in Tibetans) are mostly related to proteins targeting endoplasmic reticulum (ER) (**Figure 2B, Table S3**). Taken together, these results indicate that the differences between Tibetans and Han are mostly unfolded in the VF and VI layers, and Tibetans have less hypoxia-induced immune response and ER stress in the placenta (Elvekrog and Walter, 2015; Plumb et al., 2015; Tenzing et al., 2021). We did not see significant enrichment terms in the other layers due to the small numbers of DEGs.

There are 85 shared DEGs among the placental layers. As expected, most of them are two-layer shared DEGs between VF and VI (76/85, 89.41%), and the other 9 are multi-layer-shared DEGs (**Figure 2C**). In particular, there are two four-layer shared DEGs, and the two involved genes are *KCNE1* (potassium voltage-gated channel subfamily E regulatory subunit 1) and *AC004057.1*. *KCNE1* is significantly up-regulated in all four trophoblast layers (CP, VF, VI and VM) of Tibetans compared to Han (**Figure 2D**), and it is also the top up-regulated DEG in both VF and VI (**Figure 2E**). It encodes a potassium ion voltage-gated channel protein. The expression of *KCNE1* is reportedly down-regulated in preeclampsia placentas (Mistry et al., 2011), and the *in vitro* experiment also showed its down-regulation in the cultured cells under hypoxia (Luo et al., 2013). Therefore, the relative up-regulation of *KCNE1* in Tibetans is presumably beneficial to the reproductive outcomes at high altitude. The function of *AC004057.1* is currently unknown.

For the down-regulated genes in Tibetans, the top genes in VF and VI are *PADI1* (peptidyl arginine deiminase 1) and *TAC3* (tachykinin precursor 3), respectively

(**Figure 2E**). *PADI1* is a VF-specific DEG with 3.4-fold lower expression in Tibetans compared to Han ($p = 0.0001$) (**Figure S3**). *PADI1* encodes a member of the peptidyl arginine deiminase family enzymes. The expression of *PADI1* is triggered by hypoxia and it stimulates *PKM2* activity and contributes to the increased glycolysis under hypoxic condition (Coassolo et al., 2021). *TAC3* is a VI-VF-shared DEG with both > 3-fold changes in the two layers (Tibetans lower than Han, $p = 0.04$ for VI, and $p = 0.04$ for VF) (**Figure S3**). *TAC3* encodes a member of the tachykinin family of secreted neuropeptides, and its high expression can lead to preeclampsia (Page et al., 2006) and severe fetal growth restriction (Whitehead et al., 2013). Hence, the observed down-regulation of these two genes suggests that compared to the Han migrants, the placenta of Tibetans is more adapted to hypoxic environment, presumably with a low level of hypoxia-induced glycolysis and a reduced risk of fetal growth restriction.

Sex-biased expression divergence in the placenta between Tibetans and Han

Given the male fetuses are more sensitive to high-altitude hypoxia than the female fetuses as described in our previous report (He et al., 2022), we next analyzed the placental transcriptomes by taking into account of the infant gender (**Methods**). When the gender identity is superimposed onto the PCA map in **Figure 1D**, we did not see any clustering patterns by genders, implying that the gender-related expression changes do not cause a dramatic profile shift (**Figure S4**). However, surprisingly, we found a striking difference in view of the numbers of DEGs (Tibetans vs. Han), where the male-fetus placentas have many more DEGs than the female-fetus placentas (759 genes vs. 31 genes), and there is no intersection between them (**Figure 3A**). This between-sex difference is also reflected by the shared DEGs between the gender-combined DEG set and the gender-separated DEG sets, where the male set has 128 shared DEGs, but only 5 shared DEGs in the female set (**Figure S4**), suggesting that the male-fetus placentas contribute more to the detected between-population expression divergence.

Of the 759 male-fetus DEGs, 728 DEGs are detected in one-layer. There are 25

two-layer-shared DEGs, 4 three-layer-shared DEGs and 2 four-layer-shared DEGs (**Figure S5**). By contrast, there is no multi-layer shared DEGs in the female-fetus placentas. Among the 31 layer-shared DEGs in the male-fetus placentas, 22 DEGs are up-regulated and 9 DEGs are down-regulated in Tibetans. Similar to the above results, there are a large portion of overlap between the layer-shared DGEs of the male-fetus placentas and the gender-combined placentas (**Figure S5**).

Layer-wise, in the male-fetus placentas, UC, AN and VF have the largest numbers of DEGs (396, 177 and 189, respectively, **Figure 3A, Table S4**), accounting for 95.49% (762/798) of all DEGs, while the DEGs in the other layers are much less (< 15). By contrast, the great majority of the DEGs in the female-fetus placentas is from UC (23 genes, **Table S5**), while the other layers only have a few DEGs (< 5). This DEG pattern is highly different from that seen in the gender-combined result (**Figure 2A**), again, an indication of a sex-biased expression divergence between Tibetans and Han migrants at high altitude. Similar with the gender-combined result, the great majority (>95% in both males and females) of the DEGs are one-layer DEGs. Given the previous studies have been mostly focused on the functional role of the placental villi (Lorca et al., 2021; Tana et al., 2021; Vaughan et al., 2020), our gender-separated analyses illustrate the importance of UC and AN in fetal development, especially for the male fetus.

Markedly, in the gender-combined result, we only detected 31 DEGs in the UC layer, contrasting the large number of DEGs (396 genes) in the male-fetus placentas. We found that this might be caused by the large portion of DEGs (248/396, 62.62%) with an opposite direction of between-population expression divergence in the males and females, respectively (**Figure S6**), which canceled out the differences in the gender-combined analysis. Consistently, the between-sex expression comparison (DEGs between genders in a population: Tibetan male vs. female; Han male vs. female) indicates that the UC shows the largest DEG count difference between native Tibetans and Han migrants (**Figure 3B** and **Figure S6**). Consequently, the UC layer shows the most pronounced gender-dependent expression divergence.

The sex-biased expression divergence in placenta is likely caused by the
differential responses of male and female fetuses to hypoxic stress at altitude, as
proposed in our previous study (He et al., 2022). In UC, AN and VF, where the largest
numbers of DEGs were detected in the male placentas, the top-5 DEGs are indicated
in **Figure 3C**. These genes are primarily involved in immune responses, embryonic
development, cell migration and cholesterol metabolism, all of which are closely
related to fetal development (Barlan et al., 2017; Choi et al., 2019; Sifakis et al., 2018;
Yang et al., 2021). In UC, the enriched GO terms include up-regulation of
cytoplasmic translational initiation and DNA-binding transcription factor activity, and
down-regulation of presynaptic endocytosis in Tibetans (**Figure 3D, Table S3**). In AN,
Tibetans show a down-regulation in alcohol metabolic related process, phospholipid
metabolic process and cholesterol biosynthetic process (**Figure 3D, Table S3**). These
enriched functional categories reflect a more active protein synthesis and a reduced
risk of hypoxia-induced metabolic disturbance, which are presumably beneficial to
fetal development of the Tibetan male newborns (Guzel et al., 2017; Yung et al., 2008;
Zhang et al., 2017). We did not see significant functional terms in VF (**Table S3**).

It is known that hypoxia can lead to inhibition of global protein synthesis and
activation of ER stress (Koumenis, 2006). Our results suggest that compared to native
Tibetans, the Han migrants are more responsive to hypoxic stress, reflected by their
relative down-regulation of translational initiation and transcription factor activity,
and this trend is mostly exemplified in the UC layer of the male placentas since UC is
the key channel of maternal-fetal exchange of oxygen and nutrition. On the other hand,
cholesterol plays an important role in embryonic development (Cortes et al., 2014).
We detected the down-regulation of a cholesterol regulating gene (*DHCR7*) in
Tibetans (**Figure 3C**). The *DHCR7*-encoded protein can covalently link to *Shh* (sonic
hedgehog) to participate in brain, limb and genital development (Roux et al., 2000).
Accumulation of *DHCR7* and deficiency in cholesterol production will cause a
devastating developmental disorder (Prabhu et al., 2016).
272

Gene expression changes in the umbilical cord are associated with birth weight of the male newborns

As a critical organ for maternal-fetal nutrition and oxygen exchanges, the physiological status of the placenta is closely related to the neonatal status. We performed Weighted Gene Co-expression Network Analysis (WGCNA) to investigate the association between gene expression modules and the newborn-related traits. The investigated traits include birth weight (BW), biparietal diameter (BPD), femur length (FL), gestation time (GT), placental weight (PW), placental volume (PLV), abdominal girth (AG), amniotic fluid maximcon depth (AFMD), amniotic fluid (AFI), fetal heart rate (FH) and fundal height (FUH).

A total of 136 and 161 molecular modules were obtained from the placentas of the male infants and the female infants, respectively. We define a module as a significantly differential module (SDM) when the gene expression of a module is significantly correlated to population divergence (Tibetans vs. Han, $p < 0.05$) and the newborn-related traits ($p < 0.05$) (see **Methods** for details). Totally, there are 10 SDMs in males, but only 4 in females (**Figure 4A, Tables S6-9, Methods**). Markedly, the UC layer of the male infants has 3 SDMs, including Module 4 ($R = 0.39$, $p = 0.03$), Module 8 ($R = -0.45$, $p = 8.6 \times 10^{-3}$) and Module 9 ($R = -0.47$, $p = 6.4 \times 10^{-3}$) (**Figure 4A**), which are significantly associated with two newborn traits (BW and PLV), supporting a close involvement of UC in fetal development, especially for the male fetuses. There are four AN SDMs, two VF SDMs and one VI SDM that are associated with five newborn traits, including AFI (Module 16, Module 31 and Module 36), AFMI (Module 16 and Module 31), FUH (Module 18), AG (Module 95 and Module 96), and FH (Module 109), respectively (**Figure 4A**). For the female placentas, four SDMs are associated with AG (Module 107), AFMD (Module 45), FH (Module 45) and FUH (Module 34), respectively (**Figure 4A**). These identified SDMs suggest that the difference of transcriptional regulation of the placental layers between Tibetans and Han migrants do have impact on the newborn status, especially on the male infants.

We next looked at the DEGs in the SDMs. For the male placentas, there are a lot of DEGs involved in the ten SDMs. By contrast, among the four SMDs in the female placentas, there is only one DEG (**Figure 4B**). This pattern is consistent with the male-biased gene expression divergence between Tibetans and Han. In particular, in the UC layer of the male placentas, there are 196 DEGs in Module 8, and 104 DEGs in Module 4. Most of the DEGs in Module 8 are down-regulated in the UC layer of Tibetans (**Figure 4C**), and this module is negatively correlated with BW (**Figure 4B** and **Figure S7**), indicating that the down-regulated gene expression module in the UC layer of the Tibetan male placentas may contribute to the higher BW of the Tibetan newborns compared to Han. There are 7 hub genes in Module 8, and all of them are down-regulated in the Tibetan male placentas (**Figure 4C, 4D**). For example, *SNRNP70* (small nuclear ribonucleoprotein U1 subunit 70) enables U1 snRNA binding activity, and it is one of the known hypoxia-inducible target genes (Kurihara et al., 2016). *TSC2* (TSC complex subunit 2) is a tumor suppressor gene that encodes the growth inhibitory protein tuberin. Inhibition of the mTOR pathway by hypoxia requires *TSC2*, leading to cell proliferation under hypoxia (Brugarolas et al., 2004). Hence, the down-regulation of the great majority of DEGs in Module 8 suggests that the male fetuses are likely more sensitive to hypoxic stress than the female fetuses, and compared to native Tibetans, they are more responsive in the Han migrants, which is associated with the higher BW in the adapted natives (Tibetans) compared to the acclimatized migrants (Han). Taken together, the above results indicate that the UC layer plays a key role in the development of the male fetus and ultimately affects BW.

GHR (growth hormone receptor) is the only DEG in the SDMs of the female newborns (**Figure 4B**). The gene expression level of *GHR* in the VF layer is positively correlated with FUH ($R = 0.44$, $p = 0.017$) of the female newborns, an indicator of fetal growth (Morse et al., 2009), but not the male newborns (**Figure S7**). Consistently, it is up-regulated in Tibetans, corresponding to the higher FUH in Tibetans (He et al., 2023). Interestingly, the expression pattern of *GHR* mirrors *GH*

(growth hormone) in placenta early development and embryonic growth (Harvey and Baudet, 2014).

Histological outcomes of the sex-biased expression divergence

To further test the impact of the observed sex-biased expression divergence, we performed histological examination of the UC, VF and VI where the most prominent sex-biased expression and the largest numbers of DEGs were detected (**Figure 5A**). We analyzed 20 individual placenta samples (male-infants: 5 Tibetans vs. 5 Han; female-infants: 5 Tibetans vs. 5 Han). For the UC, five parameters of UC were evaluated using cross-sections, including umbilical vein lumen (UVL), umbilical vein wall (UVW), umbilical artery lumen (UAL), umbilical artery wall (UAW) and umbilical artery intima and media (UAIM) (**Methods**).

In line with the expression data, we observed an obvious sex-biased histological differences in UC, and the values of UVW and UAIM are significantly larger in Tibetan male infants than Han ($p = 0.003$ and $p = 0.03$), but no differences seen in the female infants ($p > 0.05$) (**Figure 5B, 5C**). The previous study reported a significant association between preeclampsia and a smaller thickness and wall area of the umbilical vein and artery, independent of gestation time and birth weight (Herzog et al., 2017), which explains the presumably adaptive histological changes of UC (increased UVM and UAIM) in Tibetans. No significant differences between Tibetans and Han were observed in the other UC parameters (**Figure S8**).

For placental trophoblast (VF and VI), we counted the number of villi and syncytial knots in the placenta (**Methods**). In the male placentas, Tibetans have more villi in the VI layer than Han ($p < 0.0001$, **Figure 5D**), and the same trend was seen in the VF layer though not significant ($p = 0.13$, **Figure S8**). Moreover, when we counted the syncytial knots in the villi, we saw a significantly lower ratio of syncytial knots/villi in male placenta of Tibetans ($p < 0.0001$, **Figure 5E**), an indication of healthier villi because the appearance of syncytial knots are the degenerating feature of placenta with remarkable accumulations of nuclei and degenerating organelles

(Heazell et al., 2007). By contrast, no difference was detected in the placentas of the

female newborns in view of the number of villi and the ratio of syncytial knots/villi in

either the VI layer or the VF layer (**Figure 5D, 5E** and **Figure S8**).

These sex-biased histological results are consistent with the patterns seen in gene

expression. The male infants show remarkably more DEGs between native Tibetans

and Han migrants than female infants, especially those genes involved in immune

responses. Also, our results prove that such male-biased expression divergence is

closely associated with birth weight. Therefore, as a distinctive remodeling of the UC

in male infants, the male-biased histological differences between Tibetans and Han

Chinese is most likely induced by the different inflammatory responses and

intrauterine hemodynamics. For placenta trophoblast, the difference of the syncytial

knots/villi ratios between VF and VI may be caused by their differences in the

distribution of placental villi. Branches of umbilical artery pass from the CP layer

before producing stem villi inferiorly (Castellucci et al., 1990). The VF layer is next

to the CP layer and the increased villus number in Tibetans suggests that the stem villi

extending out from umbilical artery produce more villous branches, which is

presumably beneficial to the maternal-fetal material exchange in Tibetans.

Consistently, the reduced number of syncytial knots in the VI layer of Tibetans might

be conducive to a better material exchange efficiency (Fogarty et al., 2013).

Natural selection acts on the placental DEGs with expression divergence between

Tibetans and Han

Given the overall gene expression divergence in the placenta between Tibetans

and Han migrants can explain the better newborn traits of Tibetans, the involved

genes could be the target of natural selection. At the same time, most of the adaptive

variants identified in Tibetan population are located in the noncoding regions of the

genome, suggesting the functional outcomes of these genetic variations are likely

achieved by gene expression regulation.

To see whether the detected DEGs of placenta tissues are subject to Darwinian

positive selection, therefore contributing to the genetic adaptation of reproductive

fitness in Tibetans, we checked all the identified DEGs and see whether they were listed in the 192 reported Tibetan selection-nominated genes (TSNGs), i.e. the genes showing signatures of Darwinian positive selection in Tibetans (Zheng et al., 2023). In total, we found that 13 DEGs are TSNGs, including 4 DEGs from the gender-combined gene list and 9 DEGs from the male-only gene list. No DEGs from the female-only gene list is overlapped with the TSNGs (**Figure 6A**). Permutation test shows that the overlapped DEGs with TSNGs are over-represented in the genome, especially for the male-only DEGs (p = 9e-4) (**Figure S9**). Among the four DEGs with selection signals in the gender-combined gene list, we saw *EPAS1*, the gene with the strongest signal of selection in Tibetans, and it shows a significant down-regulation in the UC layer of Tibetans compared to Han migrants (p = 0.04) (**Figure 6B**). Previously, there were multiple SNPs of *EPAS1* with reported GWAS signals for BW in lowland populations with a large sample size (N > 300,000) (Warrington et al., 2019). Therefore, the down-regulation of *EPAS1* in the Tibetan placentas likely reflects a blunted hypoxic response that may improve vasodilation of UC for better blood flow, and eventually leading to the higher BW in Tibetans.

FOXJ3 and *HSD11B1* are the other two genes overlapped between the placenta DEGs and the TSNG set (**Figure 6B**). *FOXJ3* (Forkhead Box J3) is a transcriptional activator, and reportedly plays an important role in spermatogenesis (Ni et al., 2016). We observed a significant correlation of *FOXJ3* expression of the CN layer with placenta weight and BW (p = 0.02 and p = 0.004, respectively) (**Figure S10**), suggesting a potential role of *FOXJ3* in transcription regulation underlying early fetal development. *HSD11B1* (Hydroxysteroid 11-Beta Dehydrogenase 1) is a microsomal enzyme that catalyzes the conversion of the inactive hormone cortisol to the active metabolite cortisone. The Tibetan placentas show a lower *HSD11B1* expression than Han migrants (p = 0.01), which may bring a lower cortisol stress hormone level. Given the hypoxia-induced glucocorticoid elevation can have impact on placental vascular development and nutrient transport (Ozmen et al., 2017), the lower cortisol stress hormone level in the Tibetan placentas might be advantageous under hypobaric

hypoxia at high altitude.

Among the nine DEGs with selection signals from the male-only gene list, four

are DEGs of the UC layer, including *EGLN1*, *L3MBTL2*, *MKL1* and *TRABD* (**Figures**

6C and S11 and **Table S10**). *EGLN1* is one of the TSNGs with strong signals of

selection in Tibetans, and it is a member of the HIF (hypoxia induced factor) pathway

and works together with *EPAS1* (Peng et al. 2011). There are two Tibetan-enriched

missense mutations in the *EGLN1* gene, and the mutations are reportedly related to

the lower erythropoiesis and augmented hypoxic ventilatory response in Tibetans

(Lorenzo et al., 2014; Song et al., 2020). *EGLN1* is temporally expressed in an

oxygen-dependent fashion, with the greatest mRNA expression at 10-12 week of

gestation (Ietta et al., 2006). In the UC layer, we detected a significant

down-regulation of *EGLN1* in the Tibetan male placentas ($p = 0.01$), but not in the

Tibetan female placentas ($p = 0.91$) (**Figure 6C**). Among the other three genes, *MKL1*

(myocardin related transcription factor A, also named *MRTFA*) was reportedly

involved in vasoconstriction mediated by human vein endothelial cells under hypoxia

(Yang et al., 2013), and it possesses a series Tibetan-enriched SNPs and structural

variants (Ouzhuluobu et al., 2020; Zheng et al., 2023). The rat gene knockout model

indicated that a *MKL1* deletion could improve hypoxia induced pulmonary

hypertension (Yuan et al., 2014), and the Tibetan-enriched 163-bp deletion at *MKL1* is

associated with the lower pulmonary arterial pressure of Tibetans (Ouzhuluobu et al.,

2020). Therefore, the down-regulation of *MKL1* in the male placentas likely works in

the same way as *EGLN1* and *EPAS1* do, and may help reduce the risk of hypoxia

induced hypertension in Tibetans. In addition, *L3MBTL2* (L3MBTL histone

methyl-lysine binding protein 2) is involved in DNA damage response (Nowsheen et

al., 2018), and cell proliferation and differentiation (Markus et al., 2003). The

function of *TRABD* (TraB domain containing) is not known. More functional data are

needed to understand the adaptive roles of these genes in fetal development at high

altitude. Except for the DEGs of the UC layer, two DEGs of the AN layer (*ACSS2* and

STXBP3) and three DEGs of the VF layer (*ZNF160*, *FOXP1* and *ENDOU*) are also

overlapped with the TSNG set (**Figure S11**). These genes are mainly involved in the

functions about embryonic development, cardiac valve morphology and immune

response (Hu et al., 2006; Kanda et al., 2005).

Furthermore, we tested the speculated polygenic effect on the adaptation of gene

expression using eQTL (expression quantitative trait loci) in Tibetans (**Methods**). The

PolyGraph (Racimo et al., 2018) detected a significant signature of polygenic

selection on gene expression in Tibetans (adjusted $p = 0.003$) (**Figure S12**).

Discussion

The placenta is the key organ for fetal development, and the transcriptional

profiles of the placenta are highly informative in revealing the gene expression

patterns and the regulatory networks, which eventually contribute to the newborn

health. In this study, with a placental transcriptomic and histological profiling of

native Tibetans and Han migrants living at the same high altitude, we intend to dissect

the molecular mechanism of the observed higher reproductive fitness of Tibetans and

to understand how the gene expression regulation in the placenta explains the genetic

adaptation of Tibetans in view of fetal development, the determinant outcome of

natural selection.

We show that among the 7 anatomic layers of the placenta, VF and VI display

the largest expression divergence between Tibetans and Han migrants, which is

expected given the known crucial role of the placental trophoblast in fetal

development (Kingdom et al., 2000). The down-regulated DEGs in VF and VI of

Tibetans (compared to Han) are mainly involved in immune response and ER stress. It

is known that besides of the maternal-fetal exchanges of nutrient, oxygen and waste,

the placenta is also an important immune organ, and the immune response is one of

the physiological reactions under chronic hypoxia (Facco et al., 2005). Placental

hypoxia/ischemia can lead to the release of a variety of placental factors, the

activation of circulating immune cells and autoantibodies, and have a profound impact

on blood flow and arterial pressure regulation (Zarate et al., 2014). Hence, our results

suggest that Tibetans might acquire the ability of overcoming the hypoxia-induced

inflammation and ER stress protein translation inhibition at high altitude.

It is known that the sex of the fetus has a detectable impact on the developmental

status (Clifton, 2010), and maternal hypoxia could alter placental formation in a sex

specific manner in rodents (Cuffe et al., 2014). In addition, sex differences in

placental traits are reportedly associated with high elevation adaptation in Andeans

(Jackson et al., 1987). However, previous studies on human placenta did not take the

fetus's gender into account in gene expression analyses. In this study, by looking at

the transcriptomic profiles of the male and female placentas separately, we discovered

a striking pattern of a male-biased expression divergence between Tibetans and Han

migrants. By contrast, there is almost no difference in placental gene expression of the

female newborns between Tibetans and Han migrants. This result suggests that the

male placenta is more sensitive to environmental stresses such as hypobaric hypoxia

at altitude, and consequently, the expression divergence between Tibetans and Han

migrants is more pronounced in males.

In particular, the UC layer possesses the largest number of DEGs (396 genes) in

the male placentas, and the enriched functional categories of these DEGs imply a

more active protein synthesis and a reduced risk of hypoxia-induced metabolic

disorder in Tibetans. UC is the main channel of maternal-fetal exchange. Previous

studies have shown that UC is important for blood vitamin D (Alp et al., 2016) and

oxygen transportation (Postigo et al., 2009; Yancey et al., 1992). We speculate that

UC may play a key role in the differential development of the male fetus between

Tibetans and Han migrants, reflected by the three detected male-specific molecular

modules that ultimately affect BW. In addition, in the UC layer, the between-sex

expression comparison indicates that the expression differences between the male

placentas and the female placentas are more pronounced in Han migrants (117 DEGs)

compared to Tibetans (33 DEGs) (**Figure S6**). Hence, under hypoxic stress, besides of

the known important functions of the trophoblast layers of the placenta, UC and the

other non-trophoblast layers are also crucial, especially for the development of the

male fetus. This observation serves as an important guiding information for future placental research.

Furthermore, at the histological level, we also see the male-biased differences between Tibetans and Han migrants (**Figure 5**). The UC and VI/VF show distinctive histological changes only in the male infants, the phenotypic outcome of the observed sex-biased expression divergence between Tibetans and Han migrants. The UC and fetal vasculature share the same embryonic origin, and vessels are often used as an important index reflecting newborn vascular health (Jin and Patterson, 2009). We observed that the umbilical artery wall and artery intima and media are significantly larger in the Tibetan male infants than in Han, consistent with the previous report that native Tibetans can prevent hypoxia-associated IUGR accompanied with the higher uteroplacental blood flow in pregnancy (Moore et al., 2004). It is known that the UC is highly influenced by systemic and local hemodynamic conditions of pregnancy, such as blood flow, oxygen tension and oxidative stress (Blanco et al., 2011). Preeclampsia is a complex disease characterized by an increased maternal blood pressure during pregnancy and increased risks of fetal growth restriction and preterm birth (Duley, 2009). It was reported that environmental hypoxia of high altitude impairs fetal growth, increases the incidence of preeclampsia, and, as a result, significantly increases the risk of perinatal and/or maternal morbidity and mortality (Julian, 2011). Additionally, the histological and morphological studies reported that the preeclampsia is associated with a smaller UC vein area and wall thickness (Herzog et al., 2017; Inan et al., 2002). Therefore, our observation of the larger umbilical artery wall and umbilical artery intima and media suggest that native Tibetans might benefit from a remolding of the UC vessels, due to the decreased expression of inflammation-related genes in the Tibetan placentas. Eventually, Tibetans have achieved a higher uteroplacental blood flow, a lower prevalence of preeclampsia, a higher newborn birth weight and a lower mortality at high altitude. Notably, all these adjustments are more significant in the male infants than in the female infants.

The adaptive changes are also reflected in the placental trophoblast. The VF and VI layers are the substantial parts of placental villi, which are composed of trophoblastic villi and are responsible for the exchange of nutrients and active transportation. Syncytial knot is a specialized structure of trophoblastic villi, which increases with fetal age and is a feature of placenta maturity (Loukeris et al., 2010). It is reportedly increased in the placenta of eclampsia and fetal growth restriction. In vitro experiments have proved that syncytial knots can be induced by hypoxia and other stressful conditions (Heazell et al., 2007). In the Tibetan male placentas, there are fewer syncytial knots compared to Han migrants, likely resulting from the observed male-biased expression changes. More functional data is needed to establish the mechanistic connection between the expression profiles and the histological outcomes.

As the gene with the strongest signal of natural selection in Tibetans, *EPAS1* has been reported in numerus studies on its contribution to high altitude adaptation. In this study, we detected a significant expression reduction of *EPAS1* in the Tibetan UC compared to the high-altitude Han. It was reported that the selected-for *EPAS1* variants/haplotype were associated with lower hemoglobin levels in the Tibetan highlanders with a major effect (Beall et al., 2010; Peng et al., 2017), and the low hemoglobin concentration of Tibetans is causally associated with a better reproductive success (Cho et al., 2017). Therefore, we speculate that the selective pressure on *EPAS1* is likely through its effect on hemoglobin, rather than directly on the reproductive traits. The down-regulation of *EPAS1* in placenta likely reflects a blunted hypoxic response that may improve vasodilation of UC for better blood flow, and eventually leading to the higher BW in Tibetans (He et al., 2023). For *EGLN1*, another well-known gene in Tibetans, we detected between-population expression difference in the male UC layer, but not in other placental layers. Considering the known adaptation mechanism of *EGLN1* is attributed to the two Tibetan-enriched missense mutations, the contribution of *EGLN1* to the gene expression change in the Tibetan UC was unexpected and worth to be explored in the future.

562  It should be noted that there are limitations in our study. We only analyzed the
563 full-term placentas. At this stage, the placenta has fully matured and we may not be
564 able to capture the dynamic changes during earlier placental and fetal development.
565 Also, we did include placenta data of a low-elevation reference, and we could not
566 completely rule out the possible effect of random genetic drift on gene expression
567 divergence between populations, which is not due to adaptation in Tibetans.
568 Additionally, although we have identified four DEGs (Tibetans vs. Han migrants)
569 with reported signals of Darwinian positive selection (Zheng et al., 2023), the
570 transcriptome data is insufficient to explain the underlying molecular mechanisms of
571 genetic adaptation in Tibetans. Future single-cell transcriptome analysis and
572 functional validations of the candidate genes are warranted to reveal the responsible
573 cell types and the molecular pathways.

574  In summary, we present a comprehensive analysis of transcriptional profiles and
575 histological differences of placentas of Tibetans and Han migrants. Our data suggest
576 that the male fetuses are more sensitive to high-altitude hypoxia, and the UC layer
577 plays a key role in the male-biased transcriptional divergence of the placenta between
578 Tibetans and Han migrants, suggesting its important role in the higher reproductive
579 fitness of Tibetans as the outcome of intense natural selection at high altitude.

580 **Materials and methods**

581 **Ethical Approval**

582  All participants signed the written informed consent. To make sure the local
583 native Tibetans can fully understand the content of the consent, the informed consent
584 was prepared in two language versions (Chinese and Tibetan), and we provided oral
585 interpretation for those who were not able to read. The protocol of this study was
586 reviewed and approved by the Internal Review Board of Kunming Institute of
587 Zoology, Chinese Academy of Sciences (Approval ID: SMKX-20160311–45) and the
588 research scheme is in accordance with the Regulations of the People's Republic of
589 China on the Administration of Human Genetic Resources.

Sample collection and processing

We collected 69 healthy full-term placental samples at a hospital in Lhasa (elevation = 3,650 m), the capital of Tibet Autonomous Region, China. The detailed information of the investigated samples and statistic results were provided in Tables S11 and S12. Based on the anatomic structure and the published protocol (Sood et al., 2006), each placenta was divided into seven tissue layers, from fetal side to maternal side, including umbilical cord (UC), amnion (AN), chorion (CN), chorionic plate (CP), villus of fetal (VF), villus of intermediate (VI) and villus of maternal (VM). The UC layer was dissected by cutting approximately 4 cm from the placental insertion point. The AN and CN layers were obtained by stripping. We sampled AN from the area approximately 5 cm from the placental insertion point and CN from the area 3 cm from the placental junction. We obtained the placental parenchyma ~2.5 cm thick at ~ 3 cm from the placental insertion point. After separation of CP, the remaining part was dissected into triplicates of equal thickness: the maternal layer (including the thin basal plate), the intermediate layer, and the fetal layer. Tissues were snap frozen in liquid nitrogen and stored at -80℃. Part of the placenta tissue samples were also subject to OCT (optimal cutting temperature compound) embedding for tissue section analysis. All placental samples were obtained and stored within two hours of delivery. The information of population ancestry of the sampled subjects was collected by self-claim with no recorded admixture in the past three generations, which was further validated by the genome-sequencing data (Zheng et al. 2023).

RNA extraction and sequencing

Total RNA of the placental samples was extracted using TRIzol reagent, and RNA purity was determined using agarose gel electrophoresis and NanoDrop 2000 (Thermo Scientific). After RNA extraction, we assessed the RNA integrity and purity using agarose gel electrophoresis. The RIN value of the extracted RNA was 7.56 ± 0.71. The extracted RNA was stored at -80℃. RNA library construction and sequencing were conducted through commercial service (Novogene). The Illumina HiSeq 2500 platform was utilized to sequencing with paired-end 150 bp reads. We

619   obtained >6G raw sequencing data for each sample.

Read alignment and quality control

In total, we generated 483 placental RNA-seq data from 69 full-term deliveries
of Tibetans (35 placentas) and Han migrants (34 placentas). The quality controls (QCs)
including following criteria: 1) filter out low-quality reads; 2) remove the sequence
with low quality ends (the applied threshold was 30); 3). Trim reads embracing the
joint sequences and the sequences containing the ambiguous base N; 4) remove the
transcripts of each sample with reads length < 60. After QCs, we obtained a
high-quality clean data for subsequent analyses.

The reference human genome (GRCh38 version) and annotation file were
downloaded from the *ensemble* database (http://www.ensembl.org/index.html), and
the clean reads were then mapped to the reference genome using STAR release 2.6.1a
(Dobin et al., 2013) with the following default parameters "*--runThreadN 8*
--readFilesCommand gunzip –c –outSAMtype BAM SortedByCoordinate –
outSAMmapqUnique 60 --outSAMattrRGline ID:sample SM:sample
PL:ILLUMINA --outSAMmultNmax 1 --outSJfilterReads Unique
--outFilterMismatchNmax 2 --alignIntronMin 20 --alignIntronMax 50000
--sjdbOverhang 149 --quantMode TranscriptomeSAM GeneCounts --twopassMode
Basic". The position information of the sequencing reads on the reference genome
and the feature information of each sample were obtained to create bam files.
Subsequently, the bam files were fed to the *resm* software
(https://github.com/deweylab/RSEM) (Li and Dewey, 2011) to generate read counts
and TPM (transcripts per million reads). In total, 10.6 billion reads were mapped to
the annotated regions, and 17,283 genes express in all the investigated placenta.

Determination of fetal and maternal origin of placenta

To trace the maternal-fetal origin of the placenta, we calculated the maternal and
fetal ratio by using the maternal genomic data and the fetal genome-array data. An
informative SNP is defined as fetal-specific when it is heterozygous in fetus (A/B)

and meanwhile is homozygous in mother (A/A) (Tsang et al., 2017).

Maternal-specific informative SNP definition is opposite. We randomly selected fetal

specific and mother specific tag SNP (at least 5 informative SNPs for each sample),

respectively. B is the specific allele and A is defined as common allele. For the

RNA-seq data of each layer of placental tissue, we calculate allelic ratio (*R*) according

to the following formula:

$$R = \frac{B}{B + A}$$

653   We calculated five times and got five *R* values each sample, and calculated the

654   average value as maternal-specific allelic ratio (R_m) and fetal-specific allelic ratio (R_f)

655   of the sample.

Differential expression analysis

Differential expression analysis was performed by R package DESeq2 (Love et

al., 2014) on the seven layers of placenta. We added two covariates (fetal sex and

maternal age) to correct for gene expression: design = ~ fetal sex + maternal age. A

nominal significance threshold of the adjusted *p* < 0.05 was used to identify

differentially expressed genes (DEGs). The Benjamini-Hochberg FDR (false

discovery rate) procedure was used in multiple testing correction of *p*-values.

Functional enrichment analysis of DEGs was evaluated using the clusterProfiler (Yu

et al., 2012), we calculated the FDR-adjusted *q*-values from the overall *p*-values to

correct for multiple testing. For the gender-separated analysis of differential

expression, only maternal age was included in covariates.

Construction of gene co-expression module

We used R package WGCNA (Langfelder and Horvath, 2008) to construct gene

expression module for the male-infant placentas and the female-infant placentas

separately. The purpose of this analysis is to capture the interactions between gene

co-expression modules and reproductive phenotypes. The 11 reproductive phenotypes

were included: birth weight (BW), biparietal diameter (BPD), femur length (FL),

gestation (GT), placental weight (PW), placental volume (PLV), abdominal girth

(AG), amniotic fluid maximcon depth (AFMD), amniotic fluid (AFI), fetal heart (FH)

and fundal height (FUH).

We used *varianceStabilizingTransformation* data from DESeq2 as input

expression data. Use the *pickSoftThreshold* function to perform the analysis of

network topology and choose a proper soft-thresholding power. Call

blockwiseModules function to build gene network and identify modules. Then we

correlated phenotypic characteristics with summary profile and look for the most

significant associations. Considering there are 12 traits, we calculated a Bonferroni

threshold (p-value = 0.05/number of independent traits) using the correlation matrix

of the traits to evaluate the significant modules. The estimated the number of

independent traits among the 12 investigated traits was 4. Therefore, we used a more

stringent significant threshold of p-value = 0.0125 (0.05/4) as the final threshold to

correct multiple testing brought by multiple traits in the WGCNA analysis.

Identification of key differential modules and hub genes

The module is defined differential when the p-value < 0.05 in population-based

WGCNA analysis and p-value < 0.05 for phenotype-based WGCNA analysis. We

defined a module as key differential modules when a differential module with the

most significant p value, at the same time, with the most significant overlaps with

DEGs. Co-expression gene networks were visualized using Cytoscape (v3.7.0)

(Shannon et al., 2003). Functional enrichment analysis of genes in the modules of

interest was performed using clusterProfiler.

We defined hub genes with three criteria: 1) gene module correlation > 0.2; 2)

gene phenotype correlation > 0.8; 3) interaction degree ranked in the top 3.

Simulation analysis

We adopted the permutation approach to evaluate the enrichment of genes

overlapped between DEGs and TSNGs. For each permutation, we randomly extracted

192 genes from the human genome, then overlapped them with the three DEG sets

(female and male; female only; male only) and counted the gene number. After 10,000

permutations, we constructed a null distribution for each set, and found that the overlapped genes between DEGs and TSNGs were significantly enriched in the "female and male" set ($p = 0.048$) and the "male" set ($p = 9e-4$), but not in the "female" set ($p = 0.1158$) (**Figure S9**). This result suggests that the observed DEGs were significantly enriched in TSNGs when compared to random sampling, especially for the male DEGs.

Analysis of polygenic adaptation

We employed the PolyGraph (Racimo et al., 2018) tool to identify the signatures of polygenic selection in Tibetans using eQTL information. We conducted eQTL analysis for seven layers, and collected a set of 5,251 eQTLs, conversing the SNPs associated with gene expression with a significant p-value < 5e-8. To obtain a list of independent eQTLs, we removed those SNPs in linkage disequilibrium (r2 > 0.2 in 1000 Genome Project). Finally, we obtained 176 independent eQTLs. At the same time, we generated a set of 1,308,436 independent SNPs of Tibetans as the control panel.

Histological analysis

We performed histological analysis for three parts of placenta: UC, VF and VI, and compared the differences between native Tibetans and Han migrants. In total, 20 individual placenta samples from random selection were analyzed, including 10 from male infants (5 Tibetans vs. 5 Han) and 10 from female infants (5 Tibetans vs. 5 Han). For the UC, we firstly fixed the UC tissue in 4% paraformaldehyde 48h after picked out from -80°C freezer, then cut the middle part of the UC (~0.5cm long) and fixed overnight. Next, the OCT embedding was performed after sucrose gradient dehydration (15% sucrose dewatering bottling, 30% sucrose dewatering bottling, 4°C), with the umbilical cord facing upward. Finally, we performed frozen section (10 μm thickness) and H&E staining. Five parameters of UC were evaluated based on published methods (Lan et al., 2018), including area size of umbilical vein lumen (UVL), umbilical vein wall (UVW), umbilical artery lumen (UAL), umbilical artery

wall (UAW) and umbilical artery intima and media (UAIM). All these parameters

were analyzed by ImageJ (Collins, 2007). Considering lack of paraffin embedded

material, we placed OCT embedded tissue in 4% paraformaldehyde for fixation and

paraffin embedding. Then, the samples were cut into 0.3 μm thick sections and

counterstain to H&E staining. Five fields with equal area per section were randomly

selected for counting the numbers of villi and the numbers of syncytial knots, and the

average numbers was taken in the Tibetan-Han comparison.

Acknowledgements

We are grateful to all participants in this study. We would like to acknowledge Chunxia Li, Rongrong Bai and all other Doctors and Nurses from Obstetrical Department and Delivery Room of Fukang Obstetrics and Gynecology Children's Branch Hospital for their assistance in data collection of the pregnant cohorts.

Funding

This study was funded by grants from the National Natural Science Foundation of China (NSFC) (32288101 and 91631306 to B.S; 3217040584 and 32000390 to Y.H.; 32070578 and U22A20340 to X.Q and 32170629 to H.Z.), the Youth Innovation Promotion Association of CAS (to Y.H.), the Science and Technology General Program of Yunnan Province (202301AW070010 and 202001AT070110 to Y.H.), the Provincial Key Research, Development and Translational Program of Tibetan Autonomous Region of China (XZ202201ZY0035G to X.Q.), and the State Key Laboratory of Genetic Resources and Evolution (GREKF22-15 to H.Z.).

Author contribution

B.S., Y.H. designed the study. T.Y., Y.G., X.Q., W.Z., H.Z, Ouzhuluobu, B. W. collected the placenta samples and phenotypic data. T.Y., Y.G. performed the experiments. T.Y., Y.G., W.Z., Y.H., K.L., B.Z., X.Z. analyzed the data. B.S. supervised the project; T.Y., Y.H., B.S. wrote the manuscript with input from all authors.

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

Transcriptome data and phenotypic data generated by this study are stored on GSA under the project ID of PRJCA014064. Additional material supporting this study is available from the corresponding authors upon request.

References

Alp, H., Tekgunduz, K.S., and Akkar, M.K. (2016). Maternal and cord blood vitamin D status in high-altitude pregnancy. J Matern Fetal Neonatal Med *29*, 571-575.

Barlan, K., Cetera, M., and Horne-Badovinac, S. (2017). Fat2 and Lar Define a Basally Localized Planar Signaling System Controlling Collective Cell Migration. Dev Cell *40*, 467-477 e465.

Beall, C.M., Cavalleri, G.L., Deng, L., Elston, R.C., Gao, Y., Knight, J., Li, C., Li, J.C., Liang, Y., McCormack, M.*, et al.* (2010a). Natural selection on EPAS1 (HIF2alpha) associated with low hemoglobin concentration in Tibetan highlanders. Proc Natl Acad Sci U S A *107*, 11459-11464.

Beall, C.M., Cavalleri, G.L., Deng, L.B., Elston, R.C., Gao, Y., Knight, J., Li, C.H., Li, J.C., Liang, Y., McCormack, M.*, et al.* (2010b). Natural selection on EPAS1 (HIF2 alpha) associated with low hemoglobin concentration in Tibetan highlanders. P Natl Acad Sci USA *107*, 11459-11464.

Beall, C.M., Song, K.J., Elston, R.C., and Goldstein, M.C. (2004). Higher offspring survival among Tibetan women with high oxygen saturation genotypes residing at 4,000 m. P Natl Acad Sci USA *101*, 14300-14304.

Bigham, A., Bauchet, M., Pinto, D., Mao, X., Akey, J.M., Mei, R., Scherer, S.W., Julian, C.G., Wilson, M.J., Lopez Herraez, D.*, et al.* (2010). Identifying signatures of natural selection in Tibetan and Andean populations using dense genome scan data. PLoS Genet *6*, e1001116.

Blanco, M.V., Vega, H.R., Giuliano, R., Grana, D.R., Azzato, F., Lerman, J., and Milei, J. (2011). Histomorphometry of umbilical cord blood vessels in preeclampsia. J Clin Hypertens (Greenwich) *13*, 30-34.

Brugarolas, J., Lei, K., Hurley, R.L., Manning, B.D., Reiling, J.H., Hafen, E., Witters, L.A., Ellisen, L.W., and Kaelin, W.G., Jr. (2004). Regulation of mTOR function in response to hypoxia by REDD1 and the TSC1/TSC2 tumor suppressor complex. Genes Dev *18*, 2893-2904.

Castellucci, M., Scheper, M., Scheffen, I., Celona, A., and Kaufmann, P. (1990). The development of the human placental villous tree. Anat Embryol (Berl) *181*, 117-128.

Cho, J.I., Basnyat, B., Jeong, C., Di Rienzo, A., Childs, G., Craig, S.R., Sun, J., and Beall, C.M. (2017). Ethnically Tibetan women in Nepal with low hemoglobin concentration have better reproductive outcomes. Evol Med Public Health *2017*, 82-96.

Choi, W.S., Lee, G., Song, W.H., Koh, J.T., Yang, J., Kwak, J.S., Kim, H.E., Kim, S.K., Son, Y.O., Nam, H.*, et al.* (2019). The CH25H-CYP7B1-RORalpha axis of cholesterol metabolism regulates osteoarthritis. Nature *566*, 254-258.

Clifton, V.L. (2010). Review: Sex and the human placenta: mediating differential strategies of fetal growth and survival. Placenta *31 Suppl*, S33-39.

Coassolo, S., Davidson, G., Negroni, L., Gambi, G., Daujat, S., Romier, C., and Davidson, I. (2021). Citrullination of pyruvate kinase M2 by PADI1 and PADI3 regulates glycolysis and cancer cell proliferation. Nat Commun *12*, 1718.

Collins, T.J. (2007). ImageJ for microscopy. Biotechniques *43*, 25-30.

Cortes, V.A., Busso, D., Maiz, A., Arteaga, A., Nervi, F., and Rigotti, A. (2014). Physiological and pathological implications of cholesterol. Front Biosci (Landmark Ed) *19*, 416-428.

Cuffe, J.S., Walton, S.L., Singh, R.R., Spiers, J.G., Bielefeldt-Ohmann, H., Wilkinson, L., Little, M.H., and Moritz, K.M. (2014). Mid- to late term hypoxia in the mouse alters placental morphology, glucocorticoid regulatory pathways and nutrient transporters in a sex-specific manner. J Physiol *592*, 3127-3141.

Deng, L., Zhang, C., Yuan, K., Gao, Y., Pan, Y., Ge, X., He, Y., Yuan, Y., Lu, Y., Zhang, X.*, et al.*
(2019). Prioritizing natural-selection signals from the deep-sequencing genomic data suggests
multi-variant adaptation in Tibetan highlanders. Natl Sci Rev *6*, 1201-1222.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and
Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Duley, L. (2009). The global impact of pre-eclampsia and eclampsia. Semin Perinatol *33*, 130-137.

Elvekrog, M.M., and Walter, P. (2015). Dynamics of co-translational protein targeting. Curr Opin Chem
Biol *29*, 79-86.

Eriksson, J.G., Kajantie, E., Osmond, C., Thornburg, K., and Barker, D.J. (2010). Boys live
dangerously in the womb. Am J Hum Biol *22*, 330-335.

Facco, M., Zilli, C., Siviero, M., Ermolao, A., Travain, G., Baesso, I., Bonamico, S., Cabrelle, A.,
Zaccaria, M., and Agostini, C. (2005). Modulation of immune response by the acute and chronic
exposure to high altitude. Med Sci Sports Exerc *37*, 768-774.

Fogarty, N.M., Ferguson-Smith, A.C., and Burton, G.J. (2013). Syncytial knots (Tenney-Parker
changes) in the human placenta: evidence of loss of transcriptional activity and oxidative damage. Am
J Pathol *183*, 144-152.

Guttmacher, A.E., Maddox, Y.T., and Spong, C.Y. (2014). The Human Placenta Project: placental
structure, development, and function in real time. Placenta *35*, 303-304.

Guzel, E., Arlier, S., Guzeloglu-Kayisli, O., Tabak, M.S., Ekiz, T., Semerci, N., Larsen, K., Schatz, F.,
Lockwood, C.J., and Kayisli, U.A. (2017). Endoplasmic Reticulum Stress and Homeostasis in
Reproductive Physiology and Pathology. Int J Mol Sci *18*.

Haeussner, E., Schmitz, C., von Koch, F., and Frank, H.G. (2013). Birth weight correlates with size but
not shape of the normal human placenta. Placenta *34*, 574-582.

Harvey, S., and Baudet, M.L. (2014). Extrapituitary growth hormone and growth? Gen Comp
Endocrinol *205*, 55-61.

He, Y., Guo, Y., Zheng, W., Yue, T., Zhang, H., Wang, B., Feng, Z., Ouzhuluobu, Cui, C., Liu, K.*, et al.*
(2023). Polygenic adaptation leads to a higher reproductive fitness of native Tibetans at high altitude.
Curr Biol.

He, Y., Li, J., Yue, T., Zheng, W., Guo, Y., Zhang, H., Chen, L., Li, C., Li, H., Cui, C.*, et al.* (2022).
Seasonality and Sex-Biased Fluctuation of Birth Weight in Tibetan Populations. Phenomics *2*, 64-71.

Heazell, A.E., Moll, S.J., Jones, C.J., Baker, P.N., and Crocker, I.P. (2007). Formation of syncytial
knots is increased by hyperoxia, hypoxia and reactive oxygen species. Placenta *28 Suppl A*, S33-40.

Herzog, E.M., Eggink, A.J., Reijnierse, A., Kerkhof, M.A., de Krijger, R.R., Roks, A.J., Reiss, I.K.,
Nigg, A.L., Eilers, P.H., Steegers, E.A.*, et al.* (2017). Impact of early- and late-onset preeclampsia on
features of placental and newborn vascular health. Placenta *49*, 72-79.

Hu, H., Wang, B., Borde, M., Nardone, J., Maika, S., Allred, L., Tucker, P.W., and Rao, A. (2006).
Foxp1 is an essential transcriptional regulator of B cell development. Nat Immunol *7*, 819-826.

Ietta, F., Wu, Y., Winter, J., Xu, J., Wang, J., Post, M., and Caniggia, I. (2006). Dynamic HIF1A
regulation during human placental development. Biol Reprod *75*, 112-121.

Inan, S., Sanci, M., Can, D., Vatansever, S., Oztekin, O., and Tinar, S. (2002). Comparative
morphological differences between umbilical cords from chronic hypertensive and preeclamptic
pregnancies. Acta Med Okayama *56*, 177-186.

Jackson, M.R., Mayhew, T.M., and Haas, J.D. (1987). Morphometric studies on villi in human term
placentae and the effects of altitude, ethnic grouping and sex of newborn. Placenta *8*, 487-495.

Jeong, C., Witonsky, D.B., Basnyat, B., Neupane, M., Beall, C.M., Childs, G., Craig, S.R., Novembre,

J., and Di Rienzo, A. (2018). Detecting past and ongoing natural selection among ethnically Tibetan

women at high altitude in Nepal. PLoS genetics *14*, e1007650.

Jin, S.W., and Patterson, C. (2009). The opening act: vasculogenesis and the origins of circulation.

Arterioscler Thromb Vasc Biol *29*, 623-629.

Julian, C.G. (2011). High altitude during pregnancy. Clin Chest Med *32*, 21-31.

Julian, C.G., Wilson, M.J., and Moore, L.G. (2009). Evolutionary Adaptation to High Altitude: A View

From In Utero. Am J Hum Biol *21*, 614-622.

Kanda, H., Tamori, Y., Shinoda, H., Yoshikawa, M., Sakaue, M., Udagawa, J., Otani, H., Tashiro, F.,

Miyazaki, J., and Kasuga, M. (2005). Adipocytes from Munc18c-null mice show increased sensitivity

to insulin-stimulated GLUT4 externalization. J Clin Invest *115*, 291-301.

Kingdom, J., Huppertz, B., Seaward, G., and Kaufmann, P. (2000). Development of the placental

villous tree and its consequences for fetal growth. Eur J Obstet Gynecol Reprod Biol *92*, 35-43.

Knofler, M., Haider, S., Saleh, L., Pollheimer, J., Gamage, T., and James, J. (2019). Human placenta

and trophoblast development: key molecular mechanisms and model systems. Cell Mol Life Sci *76*,

3479-3496.

Koumenis, C. (2006). ER stress, hypoxia tolerance and tumor progression. Curr Mol Med *6*, 55-69.

Kurihara, T., Westenskow, P.D., Gantner, M.L., Usui, Y., Schultz, A., Bravo, S., Aguilar, E., Wittgrove,

C., Friedlander, M., Paris, L.P.*, et al.* (2016). Hypoxia-induced metabolic stress in retinal pigment

epithelial cells is sufficient to induce photoreceptor degeneration. Elife *5*.

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation network

analysis. BMC Bioinformatics *9*, 559.

Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. BMC Bioinformatics *12*, 323.

Lorca, R.A., Houck, J.A., Laurent, L.C., Matarazzo, C.J., Baker, K., Horii, M., Nelson, K.K., Bales,

E.S., Euser, A.G., Parast, M.M.*, et al.* (2021). High altitude regulates the expression of AMPK

pathways in human placenta. Placenta *104*, 267-276.

Lorenzo, F.R., Huff, C., Myllymaki, M., Olenchock, B., Swierczek, S., Tashi, T., Gordeuk, V., Wuren,

T., Ri-Li, G., McClain, D.A.*, et al.* (2014). A genetic mechanism for Tibetan high-altitude adaptation.

Nat Genet *46*, 951-956.

Loukeris, K., Sela, R., and Baergen, R.N. (2010). Syncytial knots as a reflection of placental maturity:

reference values for 20 to 40 weeks' gestational age. Pediatr Dev Pathol *13*, 305-309.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol *15*, 550.

Luo, Y., Kumar, P., and Mendelson, C.R. (2013). Estrogen-related receptor gamma (ERRgamma)

regulates oxygen-dependent expression of voltage-gated potassium (K+) channels and tissue kallikrein

during human trophoblast differentiation. Mol Endocrinol *27*, 940-952.

Markus, J., Feikova, S., Sramko, M., Wolff, L., and Bies, J. (2003). Proliferation-linked expression of

the novel murine gene m4mbt encoding a nuclear zinc finger protein with four mbt domains. Gene *319*,

117-126.

Mistry, H.D., McCallum, L.A., Kurlak, L.O., Greenwood, I.A., Broughton Pipkin, F., and Tribe, R.M.

(2011). Novel expression and regulation of voltage-dependent potassium channels in placentas from

women with preeclampsia. Hypertension *58*, 497-504.

Moore, L.G., Charles, S.M., and Julian, C.G. (2011). Humans at high altitude: hypoxia and fetal growth.

Respir Physiol Neurobiol *178*, 181-190.

Moore, L.G., Shriver, M., Bemis, L., Hickler, B., Wilson, M., Brutsaert, T., Parra, E., and Vargas, E.

(2004). Maternal adaptation to high-altitude pregnancy: an experiment of nature--a review. Placenta *25*

Suppl A, S60-71.

Moore, L.G., Young, D., McCullough, R.E., Droma, T., and Zamudio, S. (2001a). Tibetan protection

from intrauterine growth restriction (IUGR) and reproductive loss at high altitude. Am J Hum Biol *13*,

635-644.

Moore, L.G., Young, D., McCullough, R.E., Droma, T., and Zamudio, S. (2001b). Tibetan protection

from intrauterine growth restriction (IUGR) and reproductive loss at high altitude. Am J Hum Biol *13*,

635-644.

Morse, K., Williams, A., and Gardosi, J. (2009). Fetal growth screening by fundal height measurement.

Best Pract Res Clin Obstet Gynaecol *23*, 809-818.

Ni, L., Xie, H., and Tan, L. (2016). Multiple roles of FOXJ3 in spermatogenesis: A lesson from Foxj3

conditional knockout mouse models. Mol Reprod Dev *83*, 1060-1069.

Nowsheen, S., Aziz, K., Aziz, A., Deng, M., Qin, B., Luo, K., Jeganathan, K.B., Zhang, H., Liu, T., Yu,

J.*, et al.* (2018). L3MBTL2 orchestrates ubiquitin signalling by dictating the sequential recruitment of

RNF8 and RNF168 after DNA damage. Nat Cell Biol *20*, 455-464.

Ouzhuluobu, He, Y., Lou, H., Cui, C., Deng, L., Gao, Y., Zheng, W., Guo, Y., Wang, X., Ning, Z.*, et al.*

(2020). De novo assembly of a Tibetan genome and identification of novel structural variants

associated with high-altitude adaptation. Natl Sci Rev *7*, 391-402.

Ozmen, A., Unek, G., and Korgun, E.T. (2017). Effect of glucocorticoids on mechanisms of placental

angiogenesis. Placenta *52*, 41-48.

Page, N.M., Dakour, J., and Morrish, D.W. (2006). Gene regulation of neurokinin B and its receptor

NK3 in late pregnancy and pre-eclampsia. Mol Hum Reprod *12*, 427-433.

Peng, S., Deyssenroth, M.A., Di Narzo, A.F., Cheng, H., Zhang, Z., Lambertini, L., Ruusalepp, A.,

Kovacic, J.C., Bjorkegren, J.L.M., Marsit, C.J.*, et al.* (2018). Genetic regulation of the placental

transcriptome underlies birth weight and risk of childhood obesity. PLoS genetics *14*, e1007799.

Peng, Y., Cui, C., He, Y., Ouzhuluobu, Zhang, H., Yang, D., Zhang, Q., Bianbazhuoma, Yang, L., He, Y.*,*

et al. (2017). Down-Regulation of EPAS1 Transcription and Genetic Adaptation of Tibetans to

High-Altitude Hypoxia. Mol Biol Evol *34*, 818-830.

Peng, Y., Yang, Z., Zhang, H., Cui, C., Qi, X., Luo, X., Tao, X., Wu, T., Ouzhuluobu, Basang, *et al.*

(2011). Genetic variations in Tibetan populations and high-altitude adaptation at the Himalayas. Mol

Biol Evol *28*, 1075-1081.

Plumb, R., Zhang, Z.R., Appathurai, S., and Mariappan, M. (2015). A functional link between the

co-translational protein translocation pathway and the UPR. Elife *4*.

Postigo, L., Heredia, G., Illsley, N.P., Torricos, T., Dolan, C., Echalar, L., Tellez, W., Maldonado, I.,

Brimacombe, M., Balanza, E.*, et al.* (2009). Where the O2 goes to: preservation of human fetal oxygen

delivery and consumption at high altitude. J Physiol *587*, 693-708.

Prabhu, A.V., Luu, W., Sharpe, L.J., and Brown, A.J. (2016). Cholesterol-mediated Degradation of

7-Dehydrocholesterol Reductase Switches the Balance from Cholesterol to Vitamin D Synthesis. J Biol

Chem *291*, 8363-8373.

Racimo, F., Berg, J.J., and Pickrell, J.K. (2018). Detecting Polygenic Adaptation in Admixture Graphs.

Genetics *208*, 1565-1584.

Roux, C., Wolf, C., Mulliez, N., Gaoua, W., Cormier, V., Chevy, F., and Citadelle, D. (2000). Role of

cholesterol in embryonic development. Am J Clin Nutr *71*, 1270S-1279S.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B.,
and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular
interaction networks. Genome Res *13*, 2498-2504.

Sifakis, S., Androutsopoulos, V.P., Pontikaki, A., Velegrakis, A., Papaioannou, G.I., Koukoura, O.,
Spandidos, D.A., and Papantoniou, N. (2018). Placental expression of PAPPA, PAPPA-2 and PLAC-1
in pregnacies is associated with FGR. Mol Med Rep *17*, 6435-6440.

Simonson, T.S., Yang, Y., Huff, C.D., Yun, H., Qin, G., Witherspoon, D.J., Bai, Z., Lorenzo, F.R., Xing,
J., Jorde, L.B.*, et al.* (2010). Genetic evidence for high-altitude adaptation in Tibet. Science *329*, 72-75.

Song, D., Navalsky, B.E., Guan, W., Ingersoll, C., Wang, T., Loro, E., Eeles, L., Matchett, K.B., Percy,
M.J., Walsby-Tickle, J.*, et al.* (2020). Tibetan PHD2, an allele with loss-of-function properties.
Proceedings of the National Academy of Sciences of the United States of America *117*, 12230-12238.

Sood, R., Zehnder, J.L., Druzin, M.L., and Brown, P.O. (2006). Gene expression patterns in human
placenta. Proc Natl Acad Sci U S A *103*, 5478-5483.

Tana, W., Noryung, T., Burton, G.J., van Patot, M.T., and Ri-Li, G. (2021). Protective Effects from the
Ischemic/Hypoxic Stress Induced by Labor in the High-Altitude Tibetan Placenta. Reprod Sci *28*,
659-664.

Tenzing, N., van Patot, M.T., Liu, H., Xu, Q., Liu, J., Wang, Z., Wang, Y., Wuren, T., and Ge, R.L.
(2021). Identification of a miRNA-mRNA Regulatory Networks in Placental Tissue Associated With
Tibetan High Altitude Adaptation. Front Genet *12*, 671119.

Tsang, J.C.H., Vong, J.S.L., Ji, L., Poon, L.C.Y., Jiang, P., Lui, K.O., Ni, Y.B., To, K.F., Cheng, Y.K.Y.,
Chiu, R.W.K.*, et al.* (2017). Integrative single-cell and cell-free plasma RNA transcriptomics elucidates
placental cellular dynamics. Proc Natl Acad Sci U S A *114*, E7786-E7795.

Vaughan, O.R., Thompson, F., Lorca, R.A., Julian, C.G., Powell, T.L., Moore, L.G., and Jansson, T.
(2020). Effect of high altitude on human placental amino acid transport. J Appl Physiol (1985) *128*,
127-133.

Wapner, R.J., and Jackson, L. (1988). Chorionic villus sampling. Clinical obstetrics and gynecology *31*,
328-344.

Warrington, N.M., Beaumont, R.N., Horikoshi, M., Day, F.R., Helgeland, O., Laurin, C., Bacelis, J.,
Peng, S., Hao, K., Feenstra, B.*, et al.* (2019). Maternal and fetal genetic effects on birth weight and
their relevance to cardio-metabolic risk factors. Nat Genet *51*, 804-814.

Whitehead, C.L., Walker, S.P., Ye, L., Mendis, S., Kaitu'u-Lino, T.J., Lappas, M., and Tong, S. (2013).
Placental specific mRNA in the maternal circulation are globally dysregulated in pregnancies
complicated by fetal growth restriction. J Clin Endocrinol Metab *98*, E429-436.

Wu, D., Liu, Y., Chen, W., Shao, J., Zhuoma, P., Zhao, D., Yu, Y., Liu, T., Yu, R., Gan, Y.*, et al.* (2022).
How Placenta Promotes the Successful Reproduction in High-Altitude Populations: A Transcriptome
Comparison between Adaptation and Acclimatization. Molecular biology and evolution *39*.

Xiang, K., Ouzhuluobu, Peng, Y., Yang, Z., Zhang, X., Cui, C., Zhang, H., Li, M., Zhang, Y., Bianba*, et*
al. (2013). Identification of a Tibetan-specific mutation in the hypoxic gene EGLN1 and its
contribution to high-altitude adaptation. Molecular biology and evolution *30*, 1889-1898.

Xu, S., Li, S., Yang, Y., Tan, J., Lou, H., Jin, W., Yang, L., Pan, X., Wang, J., Shen, Y.*, et al.* (2011). A
genome-wide search for signals of high-altitude adaptation in Tibetans. Mol Biol Evol *28*, 1003-1011.

Yancey, M.K., Moore, J., Brady, K., Milligan, D., and Strampel, W. (1992). The effect of altitude on
umbilical cord blood gases. Obstet Gynecol *79*, 571-574.

Yang, F., Whelan, E.C., Guan, X., Deng, B., Wang, S., Sun, J., Avarbock, M.R., Wu, X., and Brinster, R.L. (2021). FGF9 promotes mouse spermatogonial stem cell proliferation mediated by p38 MAPK signalling. Cell Prolif *54*, e12933.

Yang, J., Jin, Z.B., Chen, J., Huang, X.F., Li, X.M., Liang, Y.B., Mao, J.Y., Chen, X., Zheng, Z., Bakshi, A.*, et al.* (2017). Genetic signatures of high-altitude adaptation in Tibetans. Proc Natl Acad Sci U S A *114*, 4189-4194.

Yang, Y., Chen, D., Yuan, Z., Fang, F., Cheng, X., Xia, J., Fang, M., Xu, Y., and Gao, Y. (2013). Megakaryocytic leukemia 1 (MKL1) ties the epigenetic machinery to hypoxia-induced transactivation of endothelin-1. Nucleic Acids Res *41*, 6005-6017.

Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z.X.P., Pool, J.E., Xu, X., Jiang, H., Vinckenbosch, N., Korneliussen, T.S.*, et al.* (2010). Sequencing of 50 Human Exomes Reveals Adaptation to High Altitude. Science *329*, 75-78.

Yu, G.C., Wang, L.G., Han, Y.Y., and He, Q.Y. (2012). clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. Omics *16*, 284-287.

Yuan, Z., Chen, J., Chen, D., Xu, G., Xia, M., Xu, Y., and Gao, Y. (2014). Megakaryocytic leukemia 1 (MKL1) regulates hypoxia induced pulmonary hypertension in rats. Plos One *9*, e83895.

Yung, H.W., Calabrese, S., Hynx, D., Hemmings, B.A., Cetin, I., Charnock-Jones, D.S., and Burton, G.J. (2008). Evidence of placental translation inhibition and endoplasmic reticulum stress in the etiology of human intrauterine growth restriction. Am J Pathol *173*, 451-462.

Zarate, A., Saucedo, R., Valencia, J., Manuel, L., and Hernandez, M. (2014). Early disturbed placental ischemia and hypoxia creates immune alteration and vascular disorder causing preeclampsia. Arch Med Res *45*, 519-524.

Zhang, R., Dong, S., Ma, W.W., Cai, X.P., Le, Z.Y., Xiao, R., Zhou, Q., and Yu, H.L. (2017). Modulation of cholesterol transport by maternal hypercholesterolemia in human full-term placenta. Plos One *12*, e0171934.

Zheng, W., He, Y., Guo, Y., Yue, T., Zhang, H., Li, J., Zhou, B., Zeng, X., Li, L., Wang, B.*, et al.* (2023). Large-scale genome sequencing redefines the genetic footprints of high-altitude adaptation in Tibetans. Genome Biol *24*, 73.

Figure legends

Figure 1. Sampling strategy and gene expression patterns of the placental layers of Tibetans and Han migrants at high altitude. (A) The strategy of sampling the full-term placentas. The placenta was dissected into seven layers, as shown from the fetal side to the maternal side, are umbilical cord (UC), amnion (AN), chorion (CN), chorionic plate (CP), villus of fetal (VF), villus of intermediate (VI) and villus of maternal (VM). The seven layers are labeled with seven different colors. **(B)** Analysis of the maternal-fetal origins of the placental layers. **(C)** The map of principal component analysis (PCA) showing the clustering pattern of placental tissue layers. **(D)** The heat map of gene expression in the seven placental layers of Tibetans and Han migrants reveals the same clustering pattern seen in the PCA map. The layers are color-coded (see panel-A).

Figure 2. The gene expression differences of the placental layers between Tibetans and Han migrants. **(A)** The numbers of differentially expressed genes (DEGs) between Tibetans and Han in each placenta layer; Purple: the number of up-regulated DEGs in Tibetans; Green: the number of down-regulated DEGs in Tibetans; The pie chart indicates the shared DEGs among two and more placental layers. **(B)** The enriched functional categories (GO terms) of the up-regulated (top) and the down-regulated (bottom) DEGs in VF and VI, respectively. The dashed line denotes the threshold of significant test (adjusted p-value < 0.05). **(C)** The heat map of the 85 shared DEGs among two and more placental layers. **(D)** Comparison of the expression levels of *KCNE1* between Tibetans and Han in the seven layers of placenta. The significant between-population differences are indicated. adjusted p-value (p): *-$p < 0.05$; **-$p < 0.01$; ***-$p < 0.001$. **(E)** The volcano plots of the DEGs in VF and VI, respectively. The top genes are indicated.

Figure 3. Sex-biased placental expression divergence between Tibetans and Han migrants. **(A)** The DEG numbers of the seven placental layers in the male infants and the female infants, respectively. Left: female (pink); Right: male (blue); Venn chart: the overlapped DEGs between male-infant placentas and female-infant placentas. **(B)** Bar plot show the difference of gene counts between cross-sex DEGs in native Tibetans and Han migrants. **(C)** The up- and down-regulated genes in the male infants of Tibetans compared to Han, shown in the volcano plots of UC, AN and VF; The top-5 DEGs are indicated. The example gene *DHCR7* is highlighted in red. **(D)** The

enriched functional categories (GO terms) of the up-regulated (purple) and the down-regulated (green) DEGs in UC and AN, respectively. The dashed line denotes the threshold of significant test (adjusted *p*-value < 0.05).

Figure 4. Gene expression modules in UC of the male infants correlate with neonatal phenotypes. (A) The heat maps of *p*-values showing the correlations between gene expression and the newborn traits of the male-infant placentas (left) and the female-infant placentas (right). Inner ring: module name; Outer ring: layers; Red: significantly differential modules (SDMs). **(B)** The counts of the module-associated DEGs in the seven placental layers of the male infants and the female infants (the upper panels), and their correlations with the newborn traits (the lower panels). **(C)** The gene co-expression network of Module 8 of the male infants. **(D)** Gene expression comparisons of two hub genes in UC of the male infants between Tibetans and Han.

Figure 5. Sex-biased histological changes in the male-infant placentas of Tibetans. **(A)** Schematic diagram of the sampling strategy in histological analysis. **(B)(C)** The H&E stained cross-sections of the UC vessels (umbilical vein (UV); panel B; and umbilical artery (UA); panel C). The umbilical vein wall (UVW) and umbilical artery intima and media (UAIM) are highlighted in red. The boxplots show the comparisons of area size of UVW and UAIM between Tibetan and Han male infants (left) or female infants (right). The two umbilical arteries of each umbilical cord were subject to analysis in Panel-C. **(D)(E)** The H&E stained sections of the placental VI and VF. The arrows denote the syncytial knots. The boxplots show the comparison of the villi numbers or the syncytial knot/villi ratios between Tibetan and Han male infants (left) or female infants (right). Unpaired Student's t-tests are used to evaluate the significance of difference. *, *p*-value < 0.05; **, *p*-value < 0.01; ns, not significant.

Figure 6. The differentially expressed genes in the placenta that underwent positive selection in Tibetans. (A) The Venn plots show the intersections between the placental DEGs and the 192 Tibetan selection-nominated genes (TSNGs), covering the gender-combined and the gender-separated DEGs. **(B)** The three DEGs (gender-combined) under positive selection in Tibetans. The upper panels show the allele frequencies of variants with the strongest signals of selection within the gene. Besides of Tibetans and Han Chinese, the data from three other reference populations are also presented, including Japanese (JPT), Europeans (CEU) and Africans (YRI)

from the 1000 Genome Project. The solid and hollow bars in red denote the allele frequencies in the published 1001 Tibetan individuals and 35 Tibetan individuals, respectively, and the solid and hollow bars in blue denote the allele frequencies in the published 103 Han individuals and 34 Han individuals, respectively. The bottom panels show the comparison of the expression levels between Tibetans and Han in the seven layers of placenta. Only the significant between-population differences are indicated. adjusted *p*-value: *, *p*-value < 0.05; **, *p*-value < 0.01; ***, *p*-value < 0.001. **(C)** The three DEGs (gender-separated analysis) with signals of positive selection in Tibetans.