Title

Orai mediated Calcium entry sets the excitability threshold of central dopaminergic neurons by regulation of gene expression

Authors

Rishav Mitra[1], Shlesha Richhariya[1,2], Gaiti Hasan[1]

[1] National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, India
[2] Department of Biology, Brandeis University, Waltham, USA

Correspondence: gaiti@ncbs.res.in

Keywords

Neuromodulation, Drosophila flight, PPL1 DANs, histone modifications, Trithorax-like

Highlights

1.   Store-operated Ca^{2+} entry (SOCE) through Orai is required in a set of flight-promoting central dopaminergic neurons (fpDANs) during late pupae and early adults to establish their gene expression profile.

2.   SOCE activates a homeobox transcription factor, '*Trithorax-like*' and thus regulates expression of histone modifiers Set2 and *E(z)* to generate a balance between opposing epigenetic signatures of H3K36me3 and H3K27me3 on downstream genes.

3.   SOCE drives a transcriptional feedback loop to ensure expression of key genes required for neuronal function including the muscarinic acetylcholine receptor (*mAChR*) and the inositol 1,4,5-trisphosphate receptor (*itpr)*.

4.   The transcriptional program downstream of SOCE is key to functional maturation of the dopaminergic neurons, enabling their neuronal excitability and synaptic transmission required for adult flight.

1 **Graphical Abstract**

Abstract

Maturation and fine-tuning of neural circuits frequently requires

neuromodulatory signals that set the excitability threshold, neuronal connectivity and

synaptic strength. Here we present a mechanistic study of how neuromodulator

stimulated intracellular Ca^{2+} signals, through the store - operated Ca^{2+} channel Orai,

regulate intrinsic neuronal properties by control of developmental gene expression in

flight promoting central dopaminergic neurons (fpDANs). The fpDANs receive

cholinergic inputs for release of dopamine at a central brain tripartite synapse that

sustains flight (Sharma and Hasan, 2020). Cholinergic inputs act on the muscarinic

acetylcholine receptor to stimulate intracellular Ca^{2+} release through the endoplasmic

reticulum (ER) localised inositol 1,4,5-trisphosphate receptor followed by ER-store

depletion and Orai mediated store-operated Ca^{2+} entry (SOCE). Analysis of gene

expression in fpDANs followed by genetic, cellular and molecular studies identified

Orai-mediated Ca^{2+} entry as a key regulator of excitability in fpDANs during circuit

maturation. SOCE activates the transcription factor Trithorax-like (Trl) which in turn

drives expression of a set of genes including *Set2*, that encodes a histone 3 Lysine 36

methyltransferase (H3K36me3). Set2 function establishes a positive feedback loop,

essential for receiving neuromodulatory cholinergic inputs and sustaining SOCE. Chromatin modifying activity of Set2 changes the epigenetic status of fpDANs and drives expression of key ion channel and signaling genes thus setting the excitability threshold that determines dopamine release for maintenance of long flight.

Introduction

Neural circuitry underlying mature adult behaviours emerges from a combination of developmental gene expression programs and experience-dependent neuronal activity. Holometabolous insects such as *Drosophila*, reconfigure their nervous system during metamorphosis to generate circuitry capable of supporting adult behaviours (Levine, 1984). During pupal development, neurons undergo maturation of their electrical properties with a gradual increase in depolarizing responses and consequent synaptic transmission (Hardie et al., 1993; Jarvilehto and Finell, 1983). Apart from voltage-gated Ca^{2+} channels and fast-acting neurotransmitters specific to ionotropic receptors, the developing nervous system also uses neuromodulators, that target metabotropic receptors, show slower response kinetics and are capable of affecting a larger subset of neurons by means of diffusion aided volumetric transmission (Taber and Hurley, 2014). Although neuromodulators alter intrinsic neuronal properties (Marder, 2012) by changes in gene expression, the molecular mechanisms through which this is achieved and maintained over developmental timescales needs further understanding.

Ca^{2+} signals generated by neuronal activity can determine neurotransmitter specification (Spitzer, 2012), synaptic plasticity (O'Hare et al., 2022; Takechi et al., 1998), patterns of neurite growth (Gu and Spitzer, 1995) and gene expression programs (Ciceri et al., 2022; Rosenberg and Spitzer, 2011) over developmental timescales (McKinney et al., 2022) where output specificity is defined by signal dynamics. Neuromodulators generate intracellular Ca^{2+} signals by stimulation of cognate G-Protein Coupled Receptors (GPCRs) linked to the inositol 1,4,5-trisphosphate (IP_3) and Ca^{2+} signalling pathway. IP_3 transduces intracellular Ca^{2+} release through the endoplasmic reticulum (ER) localised ligand-gated ion channel, the inositol 1,4,5-trisphosphate receptor (IP_3R; Streb et al., 1984), followed by store-operated Ca^{2+} entry (SOCE) through the plasma membrane localised Orai channel (Prakriya and Lewis, 2015; Thillaiappan et al., 2019). Cellular and physiological

consequences of Ca^{2+} signals generated through SOCE exhibit different time-scales and sub-cellular localisation from activity induced signals, suggesting that they alter neuronal properties through novel mechanisms.

Expression studies, genetics and physiological analysis support a role for IP_3/Ca^{2+} and SOCE in neuronal development and the regulation of adult neuronal physiology across organisms (Hasan and Sharma, 2020; Mitra and Hasan, 2022; Somasundaram et al., 2014). In *Drosophila*, neuromodulatory signals of neurotransmitters and neuropeptides stimulate cognate GPCRs on specific neurons to generate IP_3 followed by intracellular Ca^{2+} release and SOCE (Agrawal et al., 2013; Megha and Hasan, 2017; Shakiryanova et al., 2011). Genetic and cellular studies identified neuromodulatory inputs and SOCE as an essential component for maturation of neural circuits for flight (Pathak et al., 2015; Venkiteswaran and Hasan, 2009). A focus of this flight deficit lies in a group of central dopaminergic neurons (or DANs) that include the PPL1 and PPM3 DANs (Liu et al. 2012) marked by *THD' GAL4,* alternately referred to henceforth as flight promoting DANs (fpDANs). Among the fpDANs some PPL1 DANs project to the γ2α'1 lobe of the Mushroom Body (MB; Mao and Davis 2009) a dense region of neuropil in the insect central brain, that forms a relay centre for flight (Sharma and Hasan, 2020). Axonal projections from Kenyon cells (KC) carry sensory inputs (Cervantes-Sandoval et al., 2017; Tsao et al., 2018; Yagi et al., 2016) and those from DANs carry internal state information (Mao and Davis, 2009; Riemensperger et al., 2005; Zolin et al., 2021) to functional compartments of the MB, where dopamine release modulates the KC and MB output neuron synapse (MBONs; Aso et al. 2014). The KC-DAN-MBON tripartite synapse, carries a dynamically updating representation of the motivational and behavioural state of the animal (Aso and Rubin, 2016; Berry et al., 2015; Owald et al., 2015; Waddell, 2016). Here we have investigated the molecular mechanisms that underlie the ability of neuromodulatory acetylcholine signals, acting through SOCE during circuit maturation, to determine fpDAN function required for *Drosophila* flight.

Results

A spatio-temporal requirement for SOCE determines flight

To understand how neuromodulator stimulated SOCE alters neuronal

properties over developmental time scales we began by refining further the existing

spatio-temporal coordinates of SOCE requirement for flight. Flight promoting

dopaminergic neurons (DANs) with a requirement for SOCE have been identified

earlier by expression of a dominant negative *Orai* transgene, which renders the

channel Ca^{2+} impermeable, thus abrogating all SOCE (*Orai^{E180A}*; Figure 1A; Pathak et

al. 2015; Yeromin et al. 2006). These include a smaller subset of *THD'GAL4* marked

DANs, where acetylcholine promotes dopamine release through the muscarinic

acetylcholine receptor (mAchR) by stimulating IP_3/Ca^{2+} signaling (Sharma and Hasan,

2020). *Orai^{E180A}* expression in a subset of 21-23 dopaminergic neurons marked by

THD' GAL4 (Figure 1B) led to complete loss of flight (Figure 1C) while overexpression

of the wildtype *Orai* transgene had a relatively minor effect (Figure S1A).

THD' marks DANs in the PPL1, PPL2, and PPM3 clusters of the adult central

brain (Figure 1B) of which the PPL1 (10-12) and PPM3 (6-7) cell clusters constitute

the fpDANs (Pathak et al., 2015). Among these 16-19 DANs, two pairs of PPL1 DANs

projecting to the $\gamma 2\alpha$'1 lobes of the mushroom body (marked by the MB296B split

GAL4 driver and schematised in Figure S1B; Aso et al. 2014; Aso and Rubin 2016)

were identified as contributing to the flight deficit to a significant extent (Figures 1C;

S1C). The MB296B DANs form part of a central flight relay centre identified earlier as

requiring IP_3/Ca^{2+} signaling (Sharma and Hasan, 2020). That the *THD'* neurons are

required for flight was further confirmed by inhibiting their function. Acute activation of

either a temperature sensitive *Dynamin* mutant (*Shibire^{ts}*; Kitamoto 2001), which

blocks exocytosis at an elevated temperature, or acute expression of the tetanus toxin

light chain fragment (*TeTxLC* ; Sweeney et al. 1995) which cleaves *Synaptobrevin*, an

essential component of the synaptic release machinery in *THD'* neurons led to severe

flight defects (Figure S1D).

Previous work has demonstrated that intracellular Ca^{2+} release through the

IP_3R, that precedes SOCE (Figure 1A), is required during late pupal development in

THD' marked DANs for flight (Pathak et al., 2015; Sharma and Hasan, 2020). The

precise temporal requirement for SOCE in *THD'* neurons was investigated by inducing

Orai^{E180A} transgene expression for specific periods of pupal development and in adults

with the *tubulinGAL80^{ts}* based temperature sensitive expression system, (TARGET-

Temporal And Regional Gene Expression Targeting; McGuire et al., 2004).

Approximately 100 hours of pupal development were binned into 24 hour windows,

wherein SOCE was abrogated by $Orai^{E180A}$ expression, following which, normal

development and growth was permitted. Flies were assayed for flight 5 days post

eclosion. Abrogation of SOCE by expression of the $Orai^{E180A}$ dominant negative

transgene at 72 to 96 hrs after puparium formation (APF), resulted in complete loss of

flight as compared to minor flight deficits observed upon expression of $Orai^{E180A}$ during

earlier developmental windows (Figure 1D). Abrogation of flight also occurred upon

$Orai^{E180A}$ expression during the first 2 days post-eclosion (see discussion) whereas

abrogation of SOCE 5 days after eclosion as adults resulted in only modest flight

deficits (Figure 1D), indicating that SOCE is required for maturation of *THD'* marked

flight dopaminergic neurons. Earlier work has shown that loss of SOCE by expression

of $Orai^{E180A}$ does not alter the number of dopaminergic neurons or affect neurite

projection patterns of PPL1 and PPM3 neurons (Pathak et al., 2015). Of these two

clusters, PPM3 DANs project to the ellipsoid body in the central complex (Kong et al.,

2010), and PPL1 DANs project to the mushroom body and the lateral horn (Mao and

Davis, 2009) and aid in the maintenance of extended flight bouts (Sharma and Hasan,

2020).

Loss of SOCE in late pupae leads to a re-organisation of neuronal gene

expression

The molecular consequences of loss of SOCE in *THD'* neurons were

investigated next by undertaking a comprehensive transcriptomic analysis from

fluorescence activated and sorted (FACS) *THD'* neurons with or without $Orai^{E180A}$

expression and marked with eGFP (Figure 1E; S1E). *THD'* neurons were obtained

from pupae at 72 ± 6h APF (essential SOCE requirement for flight; Figure 1E).

Expression analysis of RNAseq libraries generated from *THD'* neurons revealed a

reorganisation of the transcriptome upon loss of SOCE (Figure 1F). Expression of 822

genes was downregulated (with a fold change cut-off of < -1) as assessed using 3

different methods of Differentially Expressed Genes (DEG) analysis (Figure 1G)

whereas 137 genes were upregulated (Figure S1F). To understand if loss of SOCE

affects genes expressed in neurons from the pupal stage, we used the modEncode

dataset (Celniker et al., 2009; modENCODE Consortium et al., 2010) to reconstruct

developmental trajectories of all genes expressed in *THD'* neurons (Figure S1G).

Using an unsupervised clustering algorithm, we classified genes expressed in *THD'*

neurons into 3 clusters where 'Cluster 0' is low expression throughout development;

'Cluster 1' exhibits a larval peak in expression and 'Cluster 2' exhibits a pupal peak in

expression (Figure 1H). The majority of expressed genes (>95%) classify as low

expression throughout (Cluster 0) and were removed from further analysis. Genes

belonging to either the downregulated gene set or the upregulated gene set in SOCE-

deficient *THD'* neurons and a random set of genes were analysed further. A

comparison of the proportion of DEGs classified in the larval and pupal clusters

revealed that >75% of downregulated genes belonged to the pupal peak (Cluster 2)

whereas <20% of either the upregulated genes or a random set of genes classified as

belonging to the pupal peak (Figure 1I). Moreover, downregulated genes exhibit higher

baseline expression in wildtype pupal *THD'* neurons (Figures S1H, I). These analyses

suggest that SOCE induces expression of a set of genes in *THD'* neurons during the

late pupal phase which are subsequently required for flight. These are henceforth

referred to as SOCE-responsive genes.

SOCE regulates gene expression through a balance of Histone 3 Lysine 36

trimethylation and Histone 3 Lysine 27 trimethylation

Gene ontology (GO) analysis of SOCE-responsive genes revealed

'Transcription', 'Ion transporters', 'Ca^{2+} dependent exocytosis', 'GPCRs', 'Synaptic

components', and 'Kinases' as top GO categories (Figure 2A). Among these,

'Transcription' and 'Ion transporters' represented categories with the highest

enrichment. Further analysis of genes that classified under 'Transcription' revealed

SET domain containing genes that encode histone lysine methyltransferases

implicated in chromatin regulation and gene expression (Dillon et al., 2005) and

among the SET domain containing genes, *Set1* and *Set2* showed distinct

downregulation (Figure 2B). One of these genes, the H3K36me3 methyltransferase

Set2 (Figure 2C), was identified in a previous RNAseq from larval neurons with loss

of IP$_3$R-mediated Ca^{2+} signaling (Mitra et al., 2020). RT-PCRs from sorted *THD'*

neurons confirmed that *Set2* levels are indeed downregulated (by 76%) in *THD'*

neurons expressing $Orai^{E180A}$ and to the same extent as seen upon expression of $Set2^{RNAi}$ (79% downregulation; Figure 2C). $Set1$ and $Set2$ perform methyltransferase activity at H3K4 and H3K36 histone residues respectively (Ardehali et al., 2011; Stabell et al., 2007). Both these markers are epigenetic signatures for transcriptional activation, where H3K4me3 is enriched at the 5' end of the gene body and H3K36me3 is enriched in the gene bodies of actively transcribed genes ; schematised in Figure S2A). In the adult CNS, $Set2$, is expressed at 4-fold higher levels compared to $Set1$ (Figure S2B; modENCODE Consortium et al., 2010). $Set2$ also shows a steep 5.5 fold increase in expression levels during the pupal to adult transition, compared to a 2 fold increase in $Set1$ (Figure S2B). These observations coupled with the previous report of Set2 function downstream of IP_3R/Ca^{2+} signaling in larval glutamatergic neurons(Mitra et al., 2020), led us to pursue the role of Set2 in THD' neurons.

Set2 encodes the only $Drosophila$ methyltransferase which is specific for H3K36 trimethylation (Stabell et al., 2007). To understand the functional significance, if any, of down-regulation of $Set2$ in THD' neurons upon loss of SOCE, four independent $Set2$ RNAi constructs were obtained. Knockdown of $Set2$ in THD' neurons with all four RNAi lines tested resulted in significant flight defects (Figure 2D; S2C). $Set2$ function for flight was additionally verified in the PPL1 DANs projecting to the $\gamma2\alpha'1$ lobe of the mushroom body ($MB296BGAL4$; Figure S2D). To test the hypothesis that $Set2$ expression and function for flight is regulated by SOCE, we overexpressed $Set2$ in the $Orai^{E180A}$ background, using GAL4-UAS driven heterologous $Set2^{WT}$ expression or CRISPR-dCas9::VPR driven overexpression (Gilbert et al., 2013) from the endogenous locus, and measured flight (Figure 2D data in purple and brown). Flies expressing $Orai^{E180A}$ in the THD' neurons, which are flightless, exhibit significant rescue in flight bout durations upon $Set2$ overexpression, through either method (Figure 2D, compare red with purple and brown data), but not upon expression of a control transgene, $GCamP6m$ (Figure S2E, compare data in green and red). Additionally, overexpression of a key SOCE component, the ER Ca^{2+} sensor $STIM$ in a $Set2^{RNAi}$ background led to a rescue (Figure S2E, compare pink and grey data). Excess STIM presumably drives higher SOCE sufficient to rescue flight bout durations caused by deficient $Set2$ levels. These two sets of data taken together with downregulation of $Set2$ upon expression of $Orai^{E180A}$ (Figure 2C) support the

hypothesis that SOCE driven expression of *Set2* in fpDANs is required for flight. That Set2 function downstream of SOCE, occurs specifically through its methyltransferase activity was supported through experiments where knockdown of the demethylase *Kdm4B*, the primary histone demethylase expressed in these neurons (Figure S2F), rescued flight in *THD'>Orai^{E180A}* flies (Figure 2D; data in pink). Knockdown of the lower expressed *Drosophila* H3K36 demethylase isoform, *Kdm4A,* did not show a similar rescue (Figure S2G; data in pink). Moreover, both *Set2* depletion and loss of SOCE (*THD'>Orai^{E180A}*) lead to a decrease in the normalised H3K36me3 signal in *THD'* neurons (Figure 2E).

To understand if deficient H3K36me3 is found in the SOCE-responsive genes, we quantified the enrichment of H3K36me3 at SOCE-responsive loci from an H3K36me3 ChIP-Seq dataset (modENCODE Consortium et al., 2010) from wild-type adult heads (Figure 2F). We observed an enrichment of the H3K36me3 signal over the gene bodies of the SOCE-responsive genes (Figure 2F). Additionally, genes that were more affected by loss of SOCE (greater extent of downregulation), had a correspondingly higher H3K36me3 signal (Figure 2G) as compared with a random set of genes (Figure 2G; data in grey), further indicating that one of the ways SOCE regulates gene expression is through Set2 mediated deposition of H3K36me3.

The gene *E(z)* that encodes a component of the PRC2 complex (*EZH2*), which deposits H3K27me3 is up-regulated upon loss of SOCE (Figure 2B) suggesting that SOCE could affect additional histone modifications. While H3K36me3 is a marker for transcriptional activation (Krogan et al., 2003), H3K27me3 is a repressive signature which silences transcription (Cai et al., 2021). These two marks are antagonistic as deduced from studies which have shown that deposition of H3K36me3 allosterically inhibits the Polycomb Repressive Complex (PRC2), thereby preventing it from depositing the H3K27me3 signature at the same genomic loci (Finogenova et al., 2020). A comparison of H3K36me3 and H3K27me3 peaks over the 822 SOCE-responsive genes revealed a higher enrichment of H3K36me3 versus H3K27me3 in wild type fly heads (Figure 2H; modENCODE Consortium et al., 2010), suggesting that robust expression of these genes in adult neurons occurs through a SOCE-dependent mechanism initiated in late pupae (schematised in Figure 2I). To test this hypothesis, we fed SOCE-deficient flies (*THD'GAL4>Orai^{E180A}*) GSK343, a pharmacological

inhibitor (Verma et al., 2012) of the EZH2 component of PRC2 that is known to reduce

H3K27me3 chromatin marks and performed flight assays. Feeding of GSK343 lead to

partial rescue of flight in $THD'> Orai^{E180A}$ flies in a dose dependent manner with

maximal flight bout durations of up to 400 sec (Figure 2J; S2H).

Normal cellular responses of PPL1 DANs require Orai mediated SOCE and

H3K36 trimethylation

Next we investigated Ca^{2+} responses in SOCE and H3K36 trimethylation

deficient *THD'* PPL1 DANs upon stimulation with a neuromodulatory signal for the

muscarinic acetylcholine receptor (mAChR). It is known that *THD'* marked PPL1 DANs

receive cholinergic inputs that stimulate store Ca^{2+} release through the IP_3R, leading

to SOCE through Orai (Figure 3A; Ebihara et al., 2006; Sharma and Hasan, 2020). In

support of this idea, either expression of $Orai^{E180A}$ or knockdown of the IP_3R attenuated

the response to carbachol (CCh), a mAChR agonist, in PPL1 DANs (Figure 3B-D;

Figure S3A-C). Importantly, knockdown of *Set2* also abrogated the Ca^{2+} response to

CCh (Figure 3B-D), whereas overexpression of a transgene encoding *Set2* in *THD'*

neurons either with loss of SOCE ($Orai^{E180A}$) or with knockdown of the IP_3R ($itpr^{RNAi}$),

lead to significant rescue of the Ca^{2+} response, indicating that *Set2* is required

downstream of SOCE (Figure 3B-D) and IP_3/Ca^{2+} signalling (Figure S3A-C).

An understanding of how Set2 might rescue cellular IP_3/Ca^{2+} signaling was

obtained in an earlier study where it was demonstrated that *Set2* participates in a

transcriptional feedback loop to control the expression of key upstream components

such as the *mAChR* and the *IP_3R* in a set of larval glutamatergic neurons (Mitra et al.,

2020). To test whether *Set2* acts through a similar transcriptional feedback loop in the

current context, *THD'* DANs were isolated using FACS from appropriate genetic

backgrounds and tested for expression of key genes required for IP_3/Ca^{2+} signaling

(*mAChR* and *itpr)* and SOCE (*Stim* and *Orai*). Loss of SOCE upon expression of

$Orai^{E180A}$ lead to a significant decrease in the expression of *itpr (72%* downregulation;

Figure 3E*), mAChR (56%* downregulation; Figure 3E*), Stim (63%* downregulation;

Figure 3E) in *THD'* DANs. As expected *Orai* expression (Figure 3E) increased 2-fold

presumably due to overexpression of the $Orai^{E180A}$ transgene. Importantly, knockdown

of *Set2* in *THD'* neurons also led to downregulation of *itpr* (43%), *mAChR* (78%) and

Stim (61%) (Figure 3E). Overexpression of *Set2* in the background of *Orai^{E180A}* led to

a rescue in the levels of *itpr* (69% increase), *mAChR* (43% increase), and *Stim* (40%

increase) (Figure 3E). Overexpression of *Set2* in wildtype *THD'* neurons resulted in

upregulation of *mAChR* and *Orai* (Figure 3E). These findings indicate that while SOCE

regulates *Set2* expression (Figure 2C), *Set2* in turn functions in a feedback loop to

regulate expression of key components of IP$_3$/Ca^{2+} and SOCE such as *mAChR, IP$_3$R,*

STIM and *Orai* (Figure 3E). Thus, ectopic *Set2* overexpression in SOCE deficient *THD'*

neurons, leads to increase in expression of key genes that facilitate intracellular Ca^{2+}

signalling and SOCE (schematised in Figure 3F).

Trithorax-like or Trl is an SOCE responsive transcription factor in *THD'* DANs

The data so far identify SOCE as a key developmental regulator of the neuronal

transcriptome in *THD'* marked DANs, where it upregulates Set2 expression and thus

enhances the activation mark of H3K36 trimethylation on specific chromatin regions.

However, the mechanism by which SOCE regulates expression of *Set2,* and other

relevant effector genes remained unresolved (Figure 3F). In mammalian T cells,

SOCE leads to de-phosphorylation of the NFAT (Nuclear Factor of Activated T-cells)

family of transcription factors by Ca^{2+}/Calmodulin sensitive phosphatase Calcineurin,

followed by their nuclear translocation and ultimately transcription of relevant target

genes (Hogan et al., 2003). Unlike mammals, the *Drosophila* genome encodes a

single member of the NFAT gene family, which does not possess calcineurin binding

sites, and is therefore insensitive to intracellular Ca^{2+} and SOCE (Keyser et al., 2007).

In order to identify transcription factors (TFs) that regulate SOCE-mediated

gene expression in *Drosophila* neurons, we examined the upstream regions (up to 10

kb) of SOCE-regulated genes using motif enrichment analysis (Zambelli et al., 2009)

for TF binding sites (Figure 4A). This analysis helped identify several putative TFs with

enriched binding sites in the regulatory regions of SOCE-regulated genes (Figure 4B,

S4A) which were then analyzed for developmental expression trajectories (Hu et al.,

2017) in the *Drosophila* CNS (Figure 4B; lower panel). Among the top 10 identified

TFs, the homeobox transcription factor Trl/GAF had the highest enrichment value,

lowest p-value (Figure 4B) and was consistently expressed in the CNS through all

developmental stages, with a distinct peak during pupal development (Figure 4B;

lower panel). To test the functional significance of Trl/GAF, we tested flight in flies with *THD'* specific knockdown of *Trl* using two independent RNAi lines (Figure 4C) as well as in existing *Trl* mutant combinations that were viable as adults (Figure S4B). In all cases significant flight deficits were observed (Figure 4C, S4B). Moreover, the flight deficits could be rescued by raising SOCE either directly through overexpression of *STIM* or indirectly by *Set2* overexpression (Figure 4C; data in brown and pink respectively).

To further test Trl function, as an effector of intracellular Ca^{2+} signaling and SOCE, we investigated genetic interactions of Trl^{13C}, a hypomorphic mutant *Trl* allele, with an existing deficiency for the ER-Ca^{2+} sensor STIM ($STIM^{KO}$), an essential activator of the Orai channel (Wang et al., 2010) as well as existing mutants for an intracellular ER-Ca^{2+} release channel, the IP_3R (Joshi et al., 2004). While $STIM^{KO}/+$ flies demonstrate normal flight bout durations, adding a single copy of the Trl^{13C} allele to the $STIM^{KO}$ heterozygotes (i.e., $STIM^{KO}/Trl^{13C}$ trans-heterozygotes) resulted in significant flight deficits, which could be rescued by overexpression of *STIM* or *Set2* (Figure 4D). A single copy of Trl^{13C} placed in combination with a single copy of various *itpr* mutant alleles, also showed a significant reduction in the duration of flight bouts (Figure S4C). Together these genetic data provide strong support for Trl as a SOCE-responsive TF. Due to the strong flight deficit observed by expression of $Orai^{E180A}$ we were unable to test genetic interactions of *Trl* with Orai directly. *Orai RNAi* strains exhibit off-target effects and *Orai* hypomorphs are unavailable.

The requirement of *Trl* for SOCE-dependent gene expression was tested directly by measuring *Set2* transcripts in *THD'* neurons with knockdown of *Trl* ($THD'>Trl^{RNAi}$). *Set2* has 20 Trl binding sites in a 2 Kb region upstream of the transcription start site (Figure S4D) and knockdown of *Trl* resulted in downregulation of *Set2* (32%) and the Set2 regulated genes *itpr* (56%) and *mAchR* (73%; Figure 4E). Moreover, a trans-heterozygotic combination of hypomorphic Trl alleles ($Trl^{13C/62}$) had markedly reduced levels of brain H3K36me3 (Figure 4F).

To test if Trl drives cellular function in the fpDANs, we stimulated the mAChR with CCh and measured Ca^{2+} responses in fpDANs with knockdown of *Trl*. Compared to *WT* fpDANs, which respond robustly to cholinergic stimulation, Trl^{RNAi} expressing DANs, exhibit strongly attenuated responses (Figure 4G-I). Overexpression of *Set2* in

352 the Trl^{RNAi} background rescued the cholinergic response (Figure 4G-I). The rescue of

353 both flight (Figure 4C, D) and the cholinergic response (Figure 4G-I) by overexpression

354 of *Set2* in flies with knockdown of *Trl* in *THD'* neurons (*THD'>Trl^{RNAi}*), confirms that Trl

355 acts upstream of *Set2* to ensure optimal neuronal function and flight.

Trl function downstream of SOCE targets neuronal excitability through changes
in ion channel gene expression including VGCCs

The results obtained so far indicate that Trl could function as an intermediary

transcription factor between SOCE and its downstream effector gene *Set2*.

Overexpression of *Trl*, however, is unable to rescue the loss of flight caused by loss

of SOCE (*THD'>Orai^{E180A}*; Figure 5A). Moreover, *Trl* transcript levels are unchanged

in the *Orai^{E180A}* condition (Figure 5B) suggesting that Trl requires Ca^{2+} influx through

SOCE for its function to go from an 'inactive' form to an 'active' form (schematized in

Figure 5C). Although Trl does not possess a defined Ca^{2+} binding domain, it has been

shown to interact with a diverse set of proteins (Lomaev et al., 2017) including Ca^{2+}

sensitive kinases, and is known to undergo phosphorylation (Zhai et al., 2008). These

data support a model (schematized in Figure 5C) wherein Ca^{2+} influx through SOCE

sustains Trl activation, leading to expression of *Set2*.

Next we investigated the extent to which the identified SOCE-Trl-Set2

mechanism impacts fpDAN function. 'Ion transport' is among the top GO terms

identified in SOCE-responsive genes (Figure 2A). Indeed, multiple classes of ion

channel genes including several Na^+, K^+ and Cl^- channels are downregulated in

fpDANs upon expression of *Orai^{E180A}* (Fig 5D, E). Regulation of ion channel gene

expression by Trl was indicated from the enrichment of Trl binding sites in regulatory

regions of some ion channel loci (Figure 5F). We purified fpDANs with knock down of

Trl (*THD'>Trl^{RNAi-1}*) and measured expression of a few key voltage-gated ion channel

genes (Fig 5F). *NaCP60E*, *para* (Na^+ channels), *cac*, *ca-α1D* (VGCC subunits), *hk*,

and *eag* (outward rectifying K^+ channels) are all downregulated between 0.6-0.9 fold

upon *Trl* knockdown (Fig 5G) as well as upon loss of SOCE (Fig 5B). Taken together,

these results indicate that the expression of key voltage-gated ion channel genes

which are required for the depolarization-mediated response, maintenance of

electrical excitability and neurotransmitter release is a focus of the transcriptional program set in place by SOCE-Trl-Set2 .

To test the functional consequences of altered ion channel gene expression downstream of Trl, neuronal activity of the fpDANs was tested by means of KCl-evoked depolarisation. PPL1 DANs exhibited a robust Ca^{2+} response that was lost upon knock down of *Trl* (Figure 5H-J; data in orange). In consonance with the earlier behavioural experiments (Figure 5A), overexpression of Trl^+ in *THD'* DANs did not rescue the KCl response in PPL1 DANs with loss of SOCE ($OraiE^{180A}$; Trl^{OE}, Figure 5I, J), further reinforcing the hypothesis that, Trl activity in fpDANs is dependent of Ca^{2+} entry through Orai.

Response to KCl is also lost in PPL1 neurons by knockdown of *Set2* and restored to a significant extent by overexpression of *Set2* in *THD'* DANs lacking SOCE ($Orai^{E180A}$; $Set2^{OE}$; Figure 6A-C). We confirmed that the KCl response of PPL1 neurons requires VGCC function, either by treatment with Nimodopine (Xu and Lipscombe, 2001), an L-type VGCC inhibitor or by knockdown of a conserved VGCC subunit *cac* (cac^{RNAi}). Both forms of perturbation abrogated the depolarisation response to KCl in PPL1 neurons (Figures 6D-F; data in purple or orange respectively). Moreover, the Ca^{2+} entry upon KCl-mediated depolarization in PPL1 neurons is a cell-autonomous property as evident by measuring the response after treatment with the Na^+ channel inhibitor, Tetrodotoxin (TTX; Figures 6D-F; data in magenta).

Having confirmed that the Ca^{2+} response towards KCl depolarisation requires VGCCs, we tested the expression level of key components of *Drosophila* VGCCs (Figure 6G) in *THD'* DANs, including *cacophony* (*cac*), *ca-α1D*, *ca-α1T*, and *ca-β*. All four subunits of *Drosophila* VGCC are downregulated upon loss of SOCE ($Orai^{E180A}$; Figure 6H) in *THD'* DANs whereas overexpression of *Set2* in the background of $Orai^{E180A}$ restored expression of the four VGCC subunits tested (Figure 6H), thus explaining recovery of the KCl response upon *Set2* overexpression in PPL1 neurons lacking SOCE ($Orai^{E180A}$; Figures 6A-C).

The functional significance of downregulation of VGCC subunits by the SOCE-Trl-Set2 pathway was tested directly by knockdown of the 4 VGCC subunits independently in PPL1 neurons followed by measurement of flight bout durations. Significant loss of flight was observed with knockdown of each subunit (Figure 6I)

though in no case was the phenotype as strong as what is observed with loss of SOCE ($Orai^{E180A}$; Figure 1D). These data suggest that in addition to VGCC subunit expression, appropriate expression of other ion channels in the fpDANs is of functional significance. This idea is further supported by the observation that loss of flight upon loss of SOCE cannot be restored by overexpression of *cac* alone (Figure 6J) indicating that optimal neuronal activity requires an ensemble of genes, including ion channels, whose expression is regulated by SOCE-Trl-Set2.

THD' DANs require SOCE for developmental maturation of the neuronal excitability threshold

The functional relevance of changes in neuronal excitability during pupal maturation of the PPL1-dependent flight circuit was tested next. Inhibition of *THD'* DANs from 72 to 96 hours APF using acute induction of the inward rectifying K^+ channel *Kir2.1*(Johns et al., 1999) or optogenetic inhibition through the hyperpolarizing Cl^- channel *GtACR2* (Govorunova et al., 2015) resulted in significant flight deficits (Figure 7A; data in dark or light green). Similar flight defects were recapitulated upon hyperactivation of *THD'* neurons through either optogenetic (*CsChrimson*; Klapoetke et al., 2014) or thermogenetic (*TrpA1*; Viswanath et al., 2003) stimulation (Figure 7A; data in orange or red), indicating that these neurons require balanced neuronal activity during a critical window in late pupal development, failing which the flight circuit malfunctions. To test if restoring excitability in fpDANs lacking SOCE *(THD'>Orai^{E180A})* is sufficient to restore flight, we induced hyperexcitability by overexpressing *NachBac* (Nitabach, 2006) a bacterial Na^+ channel. *NachBac* expression rescued both flight (*THD'>Orai^{E180A}*; Figure 7B) and excitability (Figures 7C-E). A partial rescue of flight was also obtained by optogenetic stimulation of neuronal activity through activation of *THD'>Orai^{E180A}* DANs using *CsChrimson* either 72-79 hrs APF or 0-2 days post eclosion (Figure 7E). In this case, flight rescues were accompanied by a corresponding rescue of *CsChrimson* activation induced Ca^{2+} entry (Figures 7F, G). Moreover inducing neuronal hyperactivation with indirect methods such as excess K^+ supplementation (Figure S7A) or impeding glial K^+ uptake, by genetic depletion of the glial K^+ channel *sandman* (Weiss et al., 2019), partially rescued flight in animals lacking SOCE in the *THD'* DANs (Figure S7B). Together, these results indicate that

SOCE, through Trl and Set2 activity, determines the threshold of fpDAN activation
during circuit maturation by regulating expression of voltage-gated ion channel genes,
like *cac.*

fpDANs also require a balance between H3K36me3/H3K27me3 mediated
epigenetic regulation (Figure 2K). To understand if this epigenetic balance affects
flight through modulation of neuronal excitability, we measured KCl-induced
depolarizing responses of PPL1 DANs in Orai deficient animals (*THD'>Orai^{E180A}*) fed
on the H3K27me3 antagonist GSK343 (0.5 mM). GSK343 fed animals were sorted
into 2 groups (Fliers/Non-Fliers) on the basis of flight bout durations of greater or lesser
than 30 s (Figure S7C), and tested for responses to KCl (Figures S7D, E).
THD'>Orai^{E180A} flies fed on GSK343 showed a rescue in KCl responses, with a clearly
enhanced response in the fliers compared to the non-fliers (Figures S7D, E), thereby
reiterating the hypothesis that SOCE-mediated regulation of activity in these neurons
acts through epigenetic regulation between opposing histone modifications.

***THD'* DANs require SOCE-mediated gene expression for neuromodulatory**
dopamine release in the mushroom body γ lobe

Dopaminergic neurotransmission requires presynaptic voltage gated Ca^{2+}
channels (VGCCs; Brimblecombe et al., 2015), and has been implicated in the
maturation of intrinsic neuronal excitability during circuit formation (Lieberman et al.,
2018). A number of fpDANs send projections to the γ2α'1 lobe of the MB (Figure 8A).
We examined the consequences of loss of SOCE on CCh-evoked DA release at the
γ2α'1 region of the mushroom body (Figure 8B; red arrow). For this purpose we used
the dopamine sensor *GRAB-DA* (Sun et al., 2018). Loss of SOCE (*THD'>Orai^{E180A}*)
attenuated DA release (Figures 8C, D), which could be partially rescued by
overexpression of *Set2*. Taken together these data identify an SOCE-dependent gene
regulation mechanism acting through the transcription factor Trl, and the histone
modifier Set2 for timely expression of genes that impact diverse aspects of neuronal
function including excitability and sustained neurotransmitter release (schematised in
Figure 8E).

Discussion

Over the course of development, neurons define an excitability set point within
a dynamic range (Truszkowski and Aizenman, 2015), which stabilizes existing
connections (Mayseless et al., 2023) and enables circuit maturation (Johnson-
Venkatesh et al., 2015). In this study, we identify an essential role for the store-
operated Ca^{2+} channel *Orai* in setting the excitability threshold of a subset of
dopaminergic neurons central to a MB circuit for *Drosophila* flight. Orai-mediated Ca^{2+}
entry, initiated by neuromodulatory acetylcholine signals, is required for amplification
of a signalling cascade, that begins by activation of the homeo-box transcription factor
Trl, followed by upregulation of several genes, including *Set2*, a gene that encodes an
enzyme for an activating epigenetic mark, H3K36me3. Set2 in turn establishes a
transcriptional feedback loop to drive expression of key neuronal signaling genes,
including a repertoire of voltage-gated ion channel genes required for neuronal activity
in mature PPL1 dopaminergic neurons.

SOCE supports gene expression transitions during critical developmental
windows

Studies in the *Drosophila* mushroom body have identified spontaneous bouts
of voltage-gated Ca^{2+} channel mediated neuronal activity, through as yet unknown
mechanisms, which occur during early adulthood driving refinement and maturation of
behaviours such as associative learning (Leinwand and Scott, 2021). Our studies
here, on a subset of 21-23 central dopaminergic neurons (Figures 1B, C) of which
some send projections to the mushroom body and regulate flight behaviour, provide a
mechanistic explanation for neuromodulation-dependent intracellular signaling
culminating in transcriptional maturation of a circuit supporting adult flight. In the critical
developmental window of 72-96 hrs APF, a cohort of SOCE-responsive genes, that
include a range of ion channels, undergo a wave of induction. Loss of SOCE at this
stage thus renders these neurons functionally incompetent (Figures 3B-D, 6G-I, 7D-
F) and abrogates flight (Figure 1D). Developmentally assembled neuronal circuits
require experience/activity-dependent maturation (Akin and Zipursky, 2020). The
requirement of SOCE for flight circuit function during the early post eclosion phase (0-
2 days) suggests that maturation of the flight circuit extends into early adulthood,
where presumably feedback from circuit activity may further refine synaptic strengths

(Sugie et al., 2018). Expression of $Orai^{E180A}$ in 5 day old adults had no effect on flight

(Figure 1E), indicating that either SOCE is not required for maintaining gene

expression in adults or that wildtype Orai channel proteins from late pupal and early

adults perdure for long periods. The absence of any visible changes to neurite

patterning suggest that SOCE in the MB-DANs works primarily to refine synaptic

function. In summary, our findings here identify Orai-driven Ca^{2+} entry as a key

signaling step for driving coordinated gene expression enabling the maturation of

nascent neuronal circuits and sustenance of adult behaviour.

SOCE regulates a balance between competing epigenetic signatures

Chromatin structure and function is dynamic over the course of brain

development (Kishi and Gotoh, 2018). One way neurons achieve dynamic spatio-

temporal control over developmental gene expression is via epigenetic mechanisms

such as post translational histone modifications (Geng et al., 2021). Cells possess an

extensive toolkit of histone modifiers, with characterised effects on transcriptional

output by either activating or repressing gene expression (Bannister and Kouzarides,

2011). However, relatively little, is known about how modifiers of histone marks are in

turn regulated to bring about the requisite changes in neuronal gene expression over

developmental timescales.

The SET-domain containing family of histone modifiers (Figure 2B), including

H3K36me3 methyltransferase Set2 (Figure 2C), identified here appear to function as

key transcriptional effectors induced downstream of neuromodulatory inputs that

stimulate Store-operated Ca^{2+} entry through Orai during pupal and adult maturation of

flight promoting dopaminergic neurons. SOCE-driven *Set2*-mediated H3K36me3

enhances the expression of key GPCRS, components of intracellular Ca^{2+} signaling

(Figure 2A) and ion channels (Figures 6A, B) for optimal dopaminergic neuron function

(Figure 7D-F) and flight (Figure 1C). Rescue of flight by genetic depletion of an

epigenetic 'eraser' *Kdm4B*, an H3K36me3 demethylase (Figure 2D) suggests that a

balance between perdurance of H3K36me3 and its removal is actively maintained on

the expressed genes. Interestingly, another member of the SET domain family, *E(z)*

is up-regulated upon loss of SOCE (Figure 2B). *E(z)* is a component of the *Drosophila*

PRC2 complex (Polycomb repressive complex), which represses gene expression

through H2K27me3 (Margueron and Reinberg, 2011). Structural studies on *Drosophila* nucleosomes have elucidated that the H3K36me3 modification allosterically inhibits the PRC2 complex activity (Finogenova et al., 2020). Our findings support a model wherein at key developmental stages, SOCE determines the level of these competing epigenetic signatures thus allowing gene expression by enhancing Set2 mediated H3K36me3 (Fig 2H). Loss of SOCE leads to deficient Set2-mediated H3K36me3 (Figure 2E), a likely shifting of the balance in favour of PRC2 mediated H3K27me3 and overall repression in gene expression (as observed in Fig S1H, I). Feeding SOCE-deficient animals a pharmacological inhibitor of PRC2 (GSK434) led to a significant rescue in behaviour (Figure 2J) presumably by disinhibition of H3K27me3 mediated gene repression. Though flight initiation is rescued in a dose dependent manner, the duration of flight bouts is not sustained (Figure S2H) indicating that suppression of H3K27me3 marks in the SOCE-deficient flies is partial. Our findings indicate additional roles for other histone modifications such as Set1 mediated H3K4me3, which may work in concert to regulate developmental gene expression programs stimulated by SOCE. Future studies directed at a comprehensive cataloguing of multiple epigenetic signatures in defined neuronal subsets using cell-specific techniques should help elucidate these mechanisms.

Trl as an SOCE responsive transcription factor

SOCE-mediated regulation of gene expression has been reported in mammalian neural progenitor cells (Somasundaram et al., 2014) and *Drosophila* pupal neurons (Pathak et al., 2015; Richhariya et al., 2017), through as yet unknown mechanisms. A combination of motif enrichment analysis over regulatory regions of SOCE-responsive genes and expression enrichment analysis in cells of interest, helped generate a list of putative SOCE-responsive transcription factors (Figure 4A, B). Among these we experimentally validated Trl/Trithorax-like/GAGA factor as a transcription factor required for maturation of excitability in flight promoting dopaminergic neurons (Figure 4). Although Trl function has been reported in the context of early embryonic development, where it has been implicated in zygotic genome activation (Gaskill et al., 2021) (ZGA), expression of *Hox* genes (Shimojima, 2003), dosage compensation (Greenberg et al., 2004), and expression of a *Drosophila*

voltage-gated calcium channel subunit in germ cell development (Dorogova et al., 2014), his is the first ever report of its role in regulating neuronal gene expression in the context of SOCE and circuit maturation.

In vitro studies on *Drosophila* Trl reveal that it utilizes a Q-rich intrinsically disordered domain (IDD) to self-multimerize or interact with a wide range of accessory proteins (Wilkins and Lis, 1999). Multimeric complexes of Trl reside on chromatin with an exceptionally long residence time and maintain chromatin in an 'open' state (Tang et al., 2022). Trl has also been reported to directly interact with components of the Transcription initiation and elongation complex (Chopra et al., 2008; Li et al., 2013), support long distance promoter-enhancer interactions (Mahmoudi, 2003), mediate active ATP-dependent chromatin remodeling by maintaining nucleosome free stretches (Tsukiyama et al., 1994), and regulating global gene expression by controlling transcriptional stalling and pausing (Tsai et al., 2016). SOCE-dependent gene regulation may rely upon multiple transcription factors acting in concert in a cell type and developmental stage specific contexts, of which Trl may be just one. Future studies directed at other possible transcription factors downstream of SOCE would be of interest.

SOCE and the control of neuronal excitability and cholinergic neuromodulation of a tripartite synapse in the mushroom body

Neurons undergo maturation of their electrical properties with a gradual increase in depolarizing responses and synaptic transmission over the course of pupal development (Jarvilehto and Finell, 1983). Here we show that a subset of dopaminergic neurons require a balance of ion channels for optimal excitation and inhibition essential for adult function. Neuromodulatory signals, such as acetylcholine are essential for acquiring this balance and act via an SOCE-Trl-Set2 mechanism during late pupal and early adult stages. The ability of *Set2* overexpression to rescue SOCE-deficient phenotypes (Figures 6A-C) indicates that anomalous gene expression is the underlying basis of this phenotype. Indeed, SOCE deficient dopaminergic neurons show reduced expression of an ensemble of voltage-gated ion channel genes (Figures 5E, F). including genes encoding the outward rectifying K^+ channel *shaw*, slow-inactivating voltage gated K^+ channel *shaker*, a Ca^{2+} gated K^+ channel *slowpoke*

and *cacophony* a subunit of the voltage gated Ca^{2+} channel (Figure 6G). Ion channels like *Shaker* and *slowpoke* play an important role in establishing resting membrane potential (Singh and Wu, 1990), neuronal repolarisation (Lichtinghagen et al., 1990) and mediating afterhyperpolarisation (AHP) after repeated bouts of activity (Ping et al., 2011). Though dopaminergic neurons require *Set2* mediated expression of *cacophony* for excitability, *cacophony* is in itself insufficient to rescue the loss in excitability and flight deficits upon loss of SOCE (Figure 6J). Both flight and KCl-depolarisation in SOCE-deficient neurons could, however, be rescued by expression of a heterologous depolarisation activated Na^+ channel NachBac (Nitabach, 2006) (Figures 7B-D), which increases neuronal excitability by stimulating a low threshold positive feedback loop and to a lesser extent by CsChrimson-mediated optogenetic induction of neuronal activity (Figure 7E). The poorer rescue by optogenetic activation may stem from the fact that sustained bursts of activity are non-physiological and can deplete the readily releasable pool of neurotransmitters (Arrigoni and Saper, 2014; Kaeser and Regehr, 2017). Other indirect forms of inducing neuronal hyperactivation (Figures S7A, B) in SOCE-deficient neurons also achieved minor rescues in flight. These results identify SOCE as a key driver downstream of developmentally salient neuromodulatory signals for expression of an ion channels suite that enables generation of intrinsic electrical properties and functional maturation of the PPL1 DANs, and the flight circuit.

Methods

Fly maintenance

Drosophila strains were grown on standard cornmeal medium consisting of 80 g corn flour, 20 g glucose, 40 g sugar, 15 g yeast extract, 4 ml propionic acid, 5 ml p-hydroxybenzoic acid methyl ester in ethanol and 5 ml ortho-butyric acid in a total volume of 1 l (ND) at 25°C under a light-dark cycle of 12 h and 12 h. Canton S was used as wild type throughout. Mixed sex populations were used for all experiments. Several fly stocks used in this study were sourced using FlyBase (https://flybase.org), which is supported by a grant from the National Human Genome Research Institute at the US National Institutes of Health (#U41 HG000739), the British Medical Research Council (#MR/N030117/1) and FlyBase users from across the world. The stocks were

obtained from the Bloomington Drosophila Stock Centre (BDSC) supported by NIH
P40OD018537. All fly stocks used and their sources are listed in Supplementary Table
1.

Flight Assays
Flight assays were performed as previously described (Manjila and Hasan, 2018).
Briefly, flies aged 3–5 days of either sex were tested in batches of 8–10 flies, and a
minimum of 30 flies were tested for each genotype. Adult flies were anaesthetized on
ice for 2–3 min and then tethered between at the head-thorax junction using a thin
metal wire and nail polish. Post recovery at room temperature for 2-3 min, an air puff
was provided as stimulus to initiate flight. Flight duration was recorded for each fly for
15 min. For all control genotypes, GAL4 or UAS strains were crossed to the Wild Type
strain, *Canton S*. Flight assays are represented in the form of a swarm plot, wherein
each dot represents a flight bout duration for a single fly. The colours indicate different
genotypes. The Δ Flight parameter refers to the mean difference for comparisons
against the shared CS control which is shown as a Cumming estimation plot(Ho et al.,
2018). On the lower axes, mean differences are plotted as bootstrap sampling
distributions. Each mean difference is depicted as a dot. Each 95% confidence interval
is indicated by the ends of the vertical error bars. Letters beneath each distribution
refer to statistically indistinguishable groups after performing a Kruskall-Wallis test
followed by a post hoc Mann-Whitney U-test (p<0.05).

FACS
Fluorescence-activated cell sorting (FACS) was used to enrich eGFP labelled DANs
from pupal/adult. The following genotypes were used for sorting: wild type
(THD'GAL4> UAS-eGFP), Orai^{E180A} (THD'GAL4> Orai^{E180A}), Set2^{IR-1} *(THD'GAL4>*
Set2^{IR-1}), Orai^{E180A}; Set2^{OE} (THD'GAL4> Orai^{E180A}; Set2^{OE}), Set2^{OE} (THD'GAL4>
Set2^{OE}), Trl^{IR-1} (THD'GAL4> Trl^{IR-1}). Approximately 100 pupae per sample were
washed in 1×PBS and 70% ethanol. Pupal/adult CNSs were dissected in Schneider's
medium (ThermoFisher Scientific) supplemented with 10% fetal bovine serum, 2%
PenStrep, 0.02 mM insulin, 20 mM glutamine and 0.04 mg/ml glutathione. Post
dissection, the CNSs were treated with an enzyme solution [0.75 g/l Collagenase and
0.4 g/l Dispase in Rinaldini's solution (8 mg/ml NaCl, 0.2 mg/ml KCl, 0.05 mg/ml

NaH2PO4, 1 mg/ml NaHCO3 and 0.1 mg/ml glucose)] at room temperature for 30 min.

They were then washed and resuspended in ice-cold Schneider's medium and gently

triturated several times using a pipette tip to obtain a single-cell suspension. This

suspension was then passed through a 40 mm mesh filter to remove clumps and kept

on ice until sorting (less than 1 h). Flow cytometry was performed on a FACS Aria

Fusion cell sorter (BD Biosciences) with a 100 mm nozzle at 60 psi. The threshold for

GFP-positive cells was set using dissociated neurons from a non GFP-expressing

wild-type strain (*Canton S*). The same gating parameters were used to sort other

genotypes in the experiment. GFP-positive cells were collected directly in TRIzol and

then frozen immediately in dry ice until further processing. Details for all reagents used

are listed in Supplementary Table 2.

RNA-Seq

RNA from at least 600 sorted *THD'* DANs from the relevant genotypes, expressing

UAS-eGFP was subjected to 14 cycles of PCR amplification (SMARTer Seq V4 Ultra

Low Input RNA Kit; Takara Bio). 1 ng of amplified RNA was used to prepare cDNA

libraries (Nextera XT DNA library preparation kit; Illumina). cDNA libraries for 4

biological replicates for both control (*THD'GAL4/+*) and experimental

(*THD'GAL4>UAS-Orai^{E180A}*) genotypes were run on a Hiseq2500 platform. 50-70

million unpaired sequencing reads per sample were aligned to the dm6 release of the

Drosophila genome using HISAT2(Kim et al., 2015, 2019) and an overall alignment

rate of 95.2-96.8% was obtained for all samples. Featurecounts(Liao et al., 2014) was

used to assign the mapped sequence reads to the genome and obtain read counts.

Differential expression analysis was performed using three independent methods:

DESeq2(Love et al., 2014), limma-voom(Ritchie et al., 2015), and edgeR(Robinson et

al., 2009). A fold change cutoff of a minimum twofold change was used. Significance

cutoff was set at an FDR-corrected P value of 0.05 for DESeq2 and edgeR.

Volcanoplots were generated using VolcaNoseR (https://huygens.science.uva.nl/).

Comparison of gene lists and generation of Venn diagrams was performed using

Whitehead BaRC public tools (http://jura.wi.mit.edu/bioc/tools/). Gene ontology

analysis for molecular function was performed using DAVID(Huang et al., 2008, 2009).

Developmental gene expression levels were measured for downregulated genes

using FlyBase(Larkin et al., 2020) and DGET(Hu et al., 2017) (www.flyrnai.org/tools/dget/web/), and were plotted as a heatmap using ClustVis (Metsalu and Vilo, 2015) (https://biit.cs.ut.ee/clustvis/).

qRT-PCRs

Central nervous systems (CNSs) of the appropriate genotype and age were dissected in 1× phosphate-buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4 and 1.8 mM KH_2PO_4) prepared in double distilled water treated with DEPC. Each sample consisted of five CNS homogenized in 500 µl of TRIzol (Ambion, ThermoFisher Scientific) per sample. At least three biological replicate samples were made for each genotype. After homogenization the sample was kept on ice and either processed further within 30 min or stored at −80°C for up to 4 weeks before processing. RNA was isolated following the manufacturer's protocol. Purity of the isolated RNA was estimated by NanoDrop spectrophotometer (ThermoFisher Scientific) and integrity was checked by running it on a 1% Tris-EDTA agarose gel. Approximately 100 ng of total RNA was used per sample for cDNA synthesis. DNAse treatment and first-strand synthesis were performed as described previously(Pathak et al., 2015). Quantitative real time PCRs (qPCRs) were performed in a total volume of 10 µl with Kapa SYBR Fast qPCR kit (KAPA Biosystems) on an ABI 7500 fast machine operated with ABI 7500 software (Applied Biosystems). Technical duplicates were performed for each qPCR reaction. The fold change of gene expression in any experimental condition relative to wild type was calculated as 2−ΔΔCt, where $\Delta\Delta C_t$=[C_t (target gene) –C_t (rp49)] Expt. – [C_t (target gene) – C_t (rp49)]. Primers specific for rp49 and ac5c were used as internal controls. Sequences of all primers used are provided in Supplementary Table 3.

Functional imaging

Adult brains were dissected in adult hemolymph-like saline (AHL: 108 mM NaCl, 5 mM KCl, 2 mM CaCl2, 8.2 mM $MgCl_2$, 4 mM $NaHCO_3$, 1 mM NaH_2PO_4, 5 mM Trehalose, 10 mM Sucrose, 5 mM Tris, pH 7.5) after embedded in a drop of 0.1% low-melting agarose (Invitrogen). Embedded brains were bathed in AHL and then subjected to functional imaging. The genetically encoded calcium sensor *GCaMP6m*(Chen et al., 2013) was used to record changes in intracellular cytosolic

Ca^{2+} in response to stimulation with carbachol or KCl. The GRAB-DA sensor(Sun et

al., 2018) was used to measure evoked dopamine release. Images were taken as a

time series on a xy plane at an interval of 1 s using a 20× objective with an NA of 0.7

on an Olympus FV3000 inverted confocal microscope. A 488 nm laser line was used

to record GCaMP6m / GRAB-DA measurements. All live-imaging experiments were

performed with at least 10 independent brain preparations. For measuring evoked

responses, KCl/Carbachol, was added on top of the samples. For optogenetics

experiments, flies were reared on medium supplemented with 200 μM ATR (all trans

retinal), following which neuronal activation was achieved using a 633-nm laser line

to stimulate *CsChrimson*(Klapoetke et al., 2014) while simultaneously acquiring

images with the 488 nm laser line, and the images were acquired every 1 s. The raw

images were extracted using FIJI(Schindelin et al., 2012) (based on ImageJ version

2.1.0/1.53c). ΔF/F was calculated from selected regions of interest (ROIs) using the

formula ΔF/F=(Ft-F0)/F0, where Ft is the fluorescence at time t and F0 is baseline

fluorescence corresponding to the average fluorescence over the first 40-time

frames. Mean ΔF/F time-lapses were plotted using PlotTwist(Goedhart, 2020)

(https://huygens.science.uva.nl/PlotTwist/). A shaded error bar around the mean

indicates the 95% confidence interval for CCh (50 μM) or KCl (70 mM) responses.

Peaks for individual cellular responses for each genotype was calculated from before

or after the point of stimulation, using Microsoft Excel and plotted as a paired plot

using SuperPlotsOfData(Goedhart, 2021)

(https://huygens.science.uva.nl/SuperPlotsOfData/).

Boxplots were plotted using PlotsOfData (Postma and Goedhart, 2019)

(https://huygens.science.uva.nl/PlotsOfData/).

Immunohistochemistry

Immunostaining of larval *Drosophila* brains was performed as described

previously(Daniels et al., 2008). Briefly, adult brains were dissected in 1x phosphate

buffered saline (PBS) and fixed with 4% Paraformaldehyde. The dissected brains were

washed 3-4 times with 0.2% phosphate buffer, pH 7.2 containing 0.2% Triton-X 100

(PTX) and blocked with 0.2% PTX containing 5% normal goat serum (NGS) for 2 hours

at room temperature. Respective primary antibodies were incubated overnight (14–16

hr) at 4°C. After washing 3-4 times with 0.2% PTX at room temperature, they were

incubated in the respective secondary antibodies for 2 hr at room temperature. The following primary antibodies were used: chick anti-GFP antibody (1:10,000; A6455, Life Technologies), mouse anti-nc82 (anti-brp) antibody (1:50), rabbit anti-H3K36me3, mouse anti-RFP. Secondary antibodies were used at a dilution of 1:400 as follows: anti-rabbit Alexa Fluor 488 (#A11008, Life Technologies), anti-mouse Alexa Fluor 488 (#A11001, Life Technologies). Confocal images were obtained on the Olympus Confocal FV1000 microscope (Olympus) with a 40x, 1.3 NA objective or with a 60x, 1.4 NA objective. Images were visualized using either the FV10-ASW 4.0 viewer (Olympus) or Fiji(Schindelin et al., 2012). Details for all reagents used are listed in Supplementary Table 2.

Western Blotting

Adult CNSs of appropriate genotypes were dissected in ice-cold PBS. Between 5 and 10 brains were homogenized in 50 μl of NETN buffer [100 mM NaCl, 20 mM Tris-Cl (pH 8.0), 0.5 mM EDTA, 0.5% Triton-X-100, 1× Protease inhibitor cocktail (Roche)]. The homogenate (10-15 μl) was run on an 8% SDS-polyacrylamide gel. The protein was transferred to a PVDF membrane by standard semi-dry transfer protocols (10 V for 10 min). The membrane was incubated in the primary antibody overnight at 4°C. Primary antibodies were used at the following dilutions: rabbit anti-H3K36me3 (1:5000, Abcam, ab9050) and rabbit anti-H3 (1:5000, Abcam, ab12079). Secondary antibodies conjugated with horseradish peroxidase were used at dilution of 1:3000 (anti-rabbit HRP; 32260, Thermo Scientific). Protein was detected by a chemiluminescent reaction (WesternBright ECL, Advansta K12045-20). Blots were first probed for H3K36me3, stripped with a solution of 3% glacial acetic acid for 10 min, followed by re-probing with the anti-H3 antibody.

In silico ChIP-Seq Analysis

H3K36me3 enrichment data was obtained from a ChIP-chip dataset (ID_301) generated in *Drosophila* ML-DmBG3-c2 cells submitted to modEncode(Celniker et al., 2009; modENCODE Consortium et al., 2010). Enrichment scores for genomic regions were calculated using 'computematrix' and plotted as a tag density plot using 'plotHeatmap' from deeptools2(Ramírez et al., 2016). All genes were scaled to 2 kb with a flanking region of 250 bp on either end. A 50 bp length of non-overlapping bins

804 was used for averaging the score over each region length. Genes were sorted based
805 on mean enrichment scores and displayed on the heatmap in descending order.
806

Statistical tests
Non-parametric tests were employed to test significance for data that did not follow a
normal distribution. Significant differences between experimental genotypes and
relevant controls were tested either with the Kruskal-Wallis test followed by Dunn's
multiple comparison test (for multiple comparisons) or with Mann-Whitney U tests (for
pairwise comparisons). Data with normal distribution were tested by the Student's T-
test. Statistical tests and p-values are mentioned in each figure legend. For calculation
and representation of effect size, estimationstats.com was used.
815

Acknowledgements

We acknowledge the Central Imaging and Flow Cytometry Facility (CIFF, NCBS) for
maintenance of microscopes and the *Drosophila* facility (Flyfacility, NCBS) for Fly
stock maintenance and development of transgenics.

Competing Interests

The authors declare no competing interests

Funding

This work was funded by grant No. BT/PR28450/MED/122/166/2018 from the
Department of Biotechnology, Govt. of India and core support by NCBS, TIFR. RM
and SR received graduate student fellowships from NCBS, TIFR.

Data availability

The RNA Sequencing data has been submitted to GEO under accession number
GSE230134.

References

Agrawal, T., Sadaf, S., & Hasan, G. (2013). A Genetic RNAi Screen for IP3/Ca2+ Coupled GPCRs in Drosophila Identifies the PdfR as a Regulator of Insect Flight. *PLoS Genetics*, *9*, e1003849. https://doi.org/10.1371/journal.pgen.1003849

Akin, O., & Zipursky, S. L. (2020). Activity regulates brain development in the fly. *Current Opinion in Genetics & Development*, *65*, 8–13. https://doi.org/10.1016/j.gde.2020.04.005

Ardehali, M. B., Mei, A., Zobeck, K. L., Caron, M., Lis, J. T., & Kusch, T. (2011). *Drosophila*Set1 is the major histone H3 lysine 4 trimethyltransferase with role in transcription. *The EMBO Journal*, *30*, 2817–2828. https://doi.org/10.1038/emboj.2011.194

Arrigoni, E., & Saper, C. B. (2014). What optogenetic stimulation is telling us (and failing to tell us) about fast neurotransmitters and neuromodulators in brain circuits for wake–sleep regulation. *Current Opinion in Neurobiology, 29*, 165–171. https://doi.org/10.1016/j.conb.2014.07.016

Aso, Y., Hattori, D., Yu, Y., Johnston, R. M., Iyer, N. A., Ngo, T.-T. B., Dionne, H., Abbott, L. F., Axel, R., Tanimoto, H., & Rubin, G. M. (2014). The neuronal architecture of the mushroom body provides a logic for associative learning. *ELife, 3*, e04577. https://doi.org/10.7554/eLife.04577

Aso, Y., & Rubin, G. M. (2016). Dopaminergic neurons write and update memories with cell-type-specific rules. *ELife, 5*. https://doi.org/10.7554/eLife.16135

Bannister, A. J., & Kouzarides, T. (2011). Regulation of Chromatin by Histone Modifications. *Cell Research, 21*, 381–395. https://doi.org/10.1038/cr.2011.22

Berry, J. A., Cervantes-Sandoval, I., Chakraborty, M., & Davis, R. L. (2015). Sleep Facilitates Memory by Blocking Dopamine Neuron-Mediated Forgetting. *Cell, 161*, 1656–1667. https://doi.org/10.1016/j.cell.2015.05.027

Brimblecombe, K. R., Gracie, C. J., Platt, N. J., & Cragg, S. J. (2015). Gating of dopamine transmission by calcium and axonal N-, Q-, T- and L-type voltage-gated calcium channels differs between striatal domains. *The Journal of Physiology, 593*, 929–946. https://doi.org/10.1113/jphysiol.2014.285890

Cai, Y., Zhang, Y., Loh, Y. P., Tng, J. Q., Lim, M. C., Cao, Z., Raju, A., Lieberman

Aiden, E., Li, S., Manikandan, L., Tergaonkar, V., Tucker-Kellogg, G., &
Fullwood, M. J. (2021). H3K27me3-rich genomic regions can function as
silencers to repress gene expression via chromatin interactions. *Nature*
Communications, 12. https://doi.org/10.1038/s41467-021-20940-y

Celniker, S. E., Dillon, L. A. L., Gerstein, M. B., Gunsalus, K. C., Henikoff, S.,
Karpen, G. H., Kellis, M., Lai, E. C., Lieb, J. D., MacAlpine, D. M., Micklem, G.,
Piano, F., Snyder, M., Stein, L., White, K. P., & Waterston, R. H. (2009).
Unlocking the secrets of the genome. *Nature, 459*(7249), 927–930.
https://doi.org/10.1038/459927a

Cervantes-Sandoval, I., Phan, A., Chakraborty, M., & Davis, R. L. (2017). Reciprocal
synapses between mushroom body and dopamine neurons form a positive
feedback loop required for learning. *ELife, 6*. https://doi.org/10.7554/elife.23789

Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A.,
Schreiter, E. R., Kerr, R. A., Orger, M. B., Jayaraman, V., Looger, L. L.,
Svoboda, K., & Kim, D. S. (2013). Ultrasensitive fluorescent proteins for imaging
neuronal activity. *Nature, 499*, 295–300. https://doi.org/10.1038/nature12354

Chopra, V. S., Srinivasan, A., Kumar, R. P., Mishra, K., Basquin, D., Docquier, M.,
Seum, C., Pauli, D., & Mishra, R. K. (2008). Transcriptional activation by GAGA
factor is through its direct interaction with dmTAF3. *Developmental Biology, 317*,
660–670. https://doi.org/10.1016/j.ydbio.2008.02.008

Ciceri, G., Cho, H., Kshirsagar, M., Baggiolini, A., Aromolaran, K. A., Walsh, R. M.,
Goldstein, P. A., Koche, R. P., Leslie, C. S., & Studer, L. (2022). An epigenetic
barrier sets the timing of human neuronal maturation. *BioArxiv*.
https://doi.org/10.1101/2022.06.02.490114

Daniels, R. W., Gelfand, M. V, Collins, C. A., & DiAntonio, A. (2008). Visualizing
glutamatergic cell bodies and synapses inDrosophila larval and adult CNS. *The*
Journal of Comparative Neurology, 508, 131–152.
https://doi.org/10.1002/cne.21670

Dillon, S. C., Zhang, X., Trievel, R. C., & Cheng, X. (2005). The SET-domain protein
superfamily: protein lysine methyltransferases. *Genome Biology, 6*, 227.
https://doi.org/10.1186/gb-2005-6-8-227

Dorogova, N. V, Fedorova, E. V, Bolobolova, E. U., Ogienko, A. A., & Baricheva, E.

M. (2014). GAGA protein is essential for male germ cell development

inDrosophila. *Genesis*, *52*, 738–751. https://doi.org/10.1002/dvg.22789

Ebihara, T., Guo, F., Zhang, L., Kim, J. Y., & Saffen, D. (2006). Muscarinic

Acetylcholine Receptors Stimulate Ca2+ Influx in PC12D Cells Predominantly

via Activation of Ca2+ Store–Operated Channels. *The Journal of Biochemistry*,

139, 449–458. https://doi.org/10.1093/jb/mvj064

Finogenova, K., Bonnet, J., Poepsel, S., Schäfer, I. B., Finkl, K., Schmid, K., Litz, C.,

Strauss, M., Benda, C., & Müller, J. (2020). Structural basis for PRC2 decoding

of active histone methylation marks H3K36me2/3. *ELife*, *9*.

https://doi.org/10.7554/elife.61964

Gaskill, M. M., Gibson, T. J., Larson, E. D., & Harrison, M. M. (2021). GAF is

essential for zygotic genome activation and chromatin accessibility in the early

Drosophila embryo. *ELife*, *10*. https://doi.org/10.7554/elife.66668

Geng, H., Chen, H., Wang, H., & Wang, L. (2021). The Histone Modifications of

Neuronal Plasticity. *Neural Plasticity*, *2021*, 1–7.

https://doi.org/10.1155/2021/6690523

Gilbert, L. A., Larson, M. H., Morsut, L., Liu, Z., Brar, G. A., Torres, S. E., Stern-

Ginossar, N., Brandman, O., Whitehead, E. H., Doudna, J. A., Lim, W. A.,

Weissman, J. S., & Qi, L. S. (2013). CRISPR-Mediated Modular RNA-Guided

Regulation of Transcription in Eukaryotes. *Cell*, *154*, 442–451.

https://doi.org/10.1016/j.cell.2013.06.044

Goedhart, J. (2020). PlotTwist: A web app for plotting and annotating continuous

data. *PLOS Biology*, *18*, e3000581. https://doi.org/10.1371/journal.pbio.3000581

Goedhart, J. (2021). SuperPlotsOfData—a web app for the transparent display and

quantitative comparison of continuous data from different conditions. *Molecular*

Biology of the Cell, *32*, 470–474. https://doi.org/10.1091/mbc.e20-09-0583

Govorunova, E. G., Sineshchekov, O. A., Janz, R., Liu, X., & Spudich, J. L. (2015).

Natural light-gated anion channels: A family of microbial rhodopsins for

advanced optogenetics. *Science*, *349*, 647–650.

https://doi.org/10.1126/science.aaa7484

Greenberg, A. J., Yanowitz, J. L., & Schedl, P. (2004). The Drosophila GAGA Factor

Is Required for Dosage Compensation in Males and for the Formation of the

Male-Specific-Lethal Complex Chromatin Entry Site at 12DE. *Genetics*, *166*, 279–289. https://doi.org/10.1534/genetics.166.1.279

Gu, X., & Spitzer, N. C. (1995). Distinct aspects of neuronal differentiation encoded by frequency of spontaneous Ca2+ transients. *Nature*, *375*, 784–787. https://doi.org/10.1038/375784a0

Hardie, R. C., Peretz, A., Suss-Toby, E., Rom-Glas, A., Bishop, S. A., Selinger, Z., & Minke, B. (1993). Protein kinase C is required for light adaptation in Drosophila photoreceptors. *Nature*, *363*, 634–637. https://doi.org/10.1038/363634a0

Hasan, G., & Sharma, A. (2020). Regulation of neuronal physiology by Ca2+ release through the IP3R. *Current Opinion in Physiology*, *17*, 1–8. https://doi.org/10.1016/j.cophys.2020.06.001

Ho, J., Tumkaya, T., Aryal, S., Choi, H., & Claridge-Chang, A. (2018). Moving beyond *P* values: Everyday data analysis with estimation plots. *BioArxiv*. https://doi.org/10.1101/377978

Hogan, P. G., Chen, L., Nardone, J., & Rao, A. (2003). Transcriptional regulation by calcium, calcineurin, and NFAT. *Genes & Development*, *17*, 2205–2232. https://doi.org/10.1101/gad.1102703

Hu, Y., Comjean, A., Perrimon, N., & Mohr, S. E. (2017). The Drosophila Gene Expression Tool (DGET) for expression analyses. *BMC Bioinformatics*, *18*(1), 1–9. https://doi.org/10.1186/s12859-017-1509-z

Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2008). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*, *4*, 44–57. https://doi.org/10.1038/nprot.2008.211

Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*, *4*(1), 44–57. https://doi.org/10.1038/nprot.2008.211

Jarvilehto, M., & Finell, N. (1983). Development of the function of visual receptor cells during the pupal life of the flyCalliphora. *Journal of Comparative Physiology ? A*, *150*, 529–536. https://doi.org/10.1007/bf00609579

Johns, D. C., Marx, R., Mains, R. E., O'Rourke, B., & Marbán, E. (1999). Inducible Genetic Suppression of Neuronal Excitability. *The Journal of Neuroscience*, *19*, 1691–1697. https://doi.org/10.1523/jneurosci.19-05-01691.1999

Johnson-Venkatesh, E. M., Khan, M. N., Murphy, G. G., Sutton, M. A., & Umemori,

H. (2015). Excitability governs neural development in a hippocampal region

specific manner. *Development.* https://doi.org/10.1242/dev.121202

Joshi, R., Venkatesh, K., Srinivas, R., Nair, S., & Hasan, G. (2004). Genetic

Dissection of itpr Gene Function Reveals a Vital Requirement in Aminergic Cells

of Drosophila Larvae. *Genetics, 166*(1), 225–236.

https://doi.org/10.1534/genetics.166.1.225

Kaeser, P. S., & Regehr, W. G. (2017). The readily releasable pool of synaptic

vesicles. *Current Opinion in Neurobiology, 43*, 63–70.

https://doi.org/10.1016/j.conb.2016.12.012

Keyser, P., Borge-Renberg, K., & Hultmark, D. (2007). The Drosophila NFAT

homolog is involved in salt stress tolerance. *Insect Biochemistry and Molecular*

Biology, 37, 356–362. https://doi.org/10.1016/j.ibmb.2006.12.009

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with

low memory requirements. *Nature Methods, 12*(4), 357–360.

https://doi.org/10.1038/nmeth.3317

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based

genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature*

Biotechnology, 37(8), 907–915. https://doi.org/10.1038/s41587-019-0201-4

Kishi, Y., & Gotoh, Y. (2018). Regulation of Chromatin Structure During Neural

Development. *Frontiers in Neuroscience, 12*.

https://doi.org/10.3389/fnins.2018.00874

Kitamoto, T. (2001). Conditional modification of behavior inDrosophila by targeted

expression of a temperature-sensitiveshibire allele in defined neurons. *Journal*

of Neurobiology, 47, 81–92. https://doi.org/10.1002/neu.1018

Klapoetke, N. C., Murata, Y., Kim, S. S., Pulver, S. R., Birdsey-Benson, A., Cho, Y.

K., Morimoto, T. K., Chuong, A. S., Carpenter, E. J., Tian, Z., Wang, J., Xie, Y.,

Yan, Z., Zhang, Y., Chow, B. Y., Surek, B., Melkonian, M., Jayaraman, V.,

Constantine-Paton, M., … Boyden, E. S. (2014). Independent optical excitation

of distinct neural populations. *Nature Methods, 11*, 338–346.

https://doi.org/10.1038/nmeth.2836

Kong, E. C., Woo, K., Li, H., Lebestky, T., Mayer, N., Sniffen, M. R., Heberlein, U.,

Bainton, R. J., Hirsh, J., & Wolf, F. W. (2010). A Pair of Dopamine Neurons Target the D1-Like Dopamine Receptor DopR in the Central Complex to Promote Ethanol-Stimulated Locomotion in Drosophila. *PLoS ONE*, *5*. https://doi.org/10.1371/journal.pone.0009954

Krogan, N. J., Kim, M., Tong, A., Golshani, A., Cagney, G., Canadien, V., Richards, D. P., Beattie, B. K., Emili, A., Boone, C., Shilatifard, A., Buratowski, S., & Greenblatt, J. (2003). Methylation of Histone H3 by Set2 in *Saccharomyces cerevisiae* Is Linked to Transcriptional Elongation by RNA Polymerase II. *Molecular and Cellular Biology*, *23*, 4207–4218. https://doi.org/10.1128/mcb.23.12.4207-4218.2003

Larkin, A., Marygold, S. J., Antonazzo, G., Attrill, H., dos Santos, G., Garapati, P. V, Goodman, J. L., Gramates, L. S., Millburn, G., Strelets, V. B., Tabone, C. J., Thurmond, J., Perrimon, N., Gelbart, S. R., Agapite, J., Broll, K., Crosby, M., dos Santos, G., Falls, K., … Lovato, T. (2020). FlyBase: updates to the Drosophila melanogaster knowledge base. *Nucleic Acids Research*, *49*, D899–D907. https://doi.org/10.1093/nar/gkaa1026

Leinwand, S. G., & Scott, K. (2021). Juvenile hormone drives the maturation of spontaneous mushroom body neural activity and learned behavior. *Neuron*, *109*, 1836-1847.e5. https://doi.org/10.1016/j.neuron.2021.04.006

Levine, R. B. (1984). Changes in Neuronal Circuits During Insect Metamorphosis. *Journal of Experimental Biology*, *112*, 27–44. https://doi.org/10.1242/jeb.112.1.27

Li, J., Liu, Y., Rhee, H. S., Ghosh, S. K. B., Bai, L., Pugh, B. F., & Gilmour, D. S. (2013). Kinetic Competition between Elongation Rate and Binding of NELF Controls Promoter-Proximal Pausing. *Molecular Cell*, *50*(5), 711–722. https://doi.org/10.1016/j.molcel.2013.05.016

Liao, Y., Smyth, G. K., & Shi, W. (2014). FeatureCounts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930. https://doi.org/10.1093/bioinformatics/btt656

Lichtinghagen, R., Stocker, M., Wittka, R., Boheim, G., Stühmer, W., Ferrus, A., & Pongs, O. (1990). Molecular basis of altered excitability in Shaker mutants of Drosophila melanogaster. *The EMBO Journal*, *9*, 4399–4407.

https://doi.org/10.1002/j.1460-2075.1990.tb07890.x

Lieberman, O. J., McGuirt, A. F., Mosharov, E. V, Pigulevskiy, I., Hobson, B. D.,
Choi, S., Frier, M. D., Santini, E., Borgkvist, A., & Sulzer, D. (2018). Dopamine
Triggers the Maturation of Striatal Spiny Projection Neuron Excitability during a
Critical Period. *Neuron*, *99*, 540-554.e4.
https://doi.org/10.1016/j.neuron.2018.06.044

Liu, C., Plaçais, P.-Y., Yamagata, N., Pfeiffer, B. D., Aso, Y., Friedrich, A. B.,
Siwanowicz, I., Rubin, G. M., Preat, T., & Tanimoto, H. (2012). A subset of
dopamine neurons signals reward for odour memory in Drosophila. *Nature*, *488*,
512–516. https://doi.org/10.1038/nature11304

Lomaev, D., Mikhailova, A., Erokhin, M., Shaposhnikov, A. V, Moresco, J. J.,
Blokhina, T., Wolle, D., Aoki, T., Ryabykh, V., Yates, J. R., Shidlovskii, Y. V,
Georgiev, P., Schedl, P., & Chetverina, D. (2017). The GAGA factor regulatory
network: Identification of GAGA factor associated proteins. *PLOS ONE*, *12*,
e0173602. https://doi.org/10.1371/journal.pone.0173602

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21.
https://doi.org/10.1186/s13059-014-0550-8

Mahmoudi, T. (2003). GAGA facilitates binding of Pleiohomeotic to a chromatinized
Polycomb response element. *Nucleic Acids Research*, *31*(14), 4147–4156.
https://doi.org/10.1093/nar/gkg479

Manjila, S., & Hasan, G. (2018). Flight and Climbing Assay for Assessing Motor
Functions in Drosophila. *BIO-PROTOCOL*, *8*.
https://doi.org/10.21769/bioprotoc.2742

Mao, Z., & Davis, R. (2009). Eight different types of dopaminergic neurons innervate
the Drosophila mushroom body neuropil: anatomical and physiological
heterogeneity. *Frontiers in Neural Circuits*, *3*.
https://doi.org/10.3389/neuro.04.005.2009

Marder, E. (2012). Neuromodulation of Neuronal Circuits: Back to the Future.
Neuron, *76*, 1–11. https://doi.org/10.1016/j.neuron.2012.09.010

Margueron, R., & Reinberg, D. (2011). The Polycomb complex PRC2 and its mark in
life. *Nature*, *469*, 343–349. https://doi.org/10.1038/nature09784

Mayseless, O., Shapira, G., Rachad, E. Y., Fiala, A., & Schuldiner, O. (2023). Neuronal excitability as a regulator of circuit remodeling. *Current Biology*, *33*, 981-989.e3. https://doi.org/10.1016/j.cub.2023.01.032

McGuire, S. E., Mao, Z., & Davis, R. L. (2004). Spatiotemporal Gene Expression Targeting with the TARGET and Gene-Switch Systems in Drosophila. *Science Signaling*, *2004*, pl6–pl6. https://doi.org/10.1126/stke.2202004pl6

McKinney, A., Petrova, R., & Panagiotakos, G. (2022). Calcium and activity-dependent signaling in the developing cerebral cortex. *Development*, *149*. https://doi.org/10.1242/dev.198853

Megha, & Hasan, G. (2017). IP3R-mediated Ca2+ release regulates protein metabolism in drosophila neuroendocrine cells: Implications for development under nutrient stress. *Development (Cambridge)*, *144*(8), 1484–1489. https://doi.org/10.1242/dev.145235

Metsalu, T., & Vilo, J. (2015). ClustVis: A web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Research*, *43*(W1), W566–W570. https://doi.org/10.1093/nar/gkv468

Mitra, R., & Hasan, G. (2022). Store-operated Ca2+ entry regulates neuronal gene expression and function. *Current Opinion in Neurobiology*, *73*, 102520. https://doi.org/10.1016/j.conb.2022.01.005

Mitra, R., Richhariya, S., Jayakumar, S., Notani, D., & Hasan, G. (2020). IP3/Ca2+ signals regulate larval to pupal transition under nutrient stress through the H3K36 methyltransferase dSET2. *BioRxiv*. https://doi.org/10.1101/2020.11.25.399329

modENCODE Consortium, Roy, S., Ernst, J., Kharchenko, P. V, Kheradpour, P., Negre, N., Eaton, M. L., Landolin, J. M., Bristow, C. A., Ma, L., Lin, M. F., Washietl, S., Arshinoff, B. I., Ay, F., Meyer, P. E., Robine, N., Washington, N. L., Di Stefano, L., Berezikov, E., … Kellis, M. (2010). Identification of functional elements and regulatory circuits by Drosophila modENCODE. *Science (New York, N.Y.)*, *330*, 1787–1797. https://doi.org/10.1126/science.1198374

Nitabach, M. N. (2006). Electrical Hyperexcitation of Lateral Ventral Pacemaker Neurons Desynchronizes Downstream Circadian Oscillators in the Fly Circadian Circuit and Induces Multiple Behavioral Periods. *Journal of Neuroscience*, *26*,

479–489. https://doi.org/10.1523/jneurosci.3915-05.2006

O'Hare, J. K., Gonzalez, K. C., Herrlinger, S. A., Hirabayashi, Y., Hewitt, V. L.,

Blockus, H., Szoboszlay, M., Rolotti, S. V, Geiller, T. C., Negrean, A., Chelur, V.,

Polleux, F., & Losonczy, A. (2022). Compartment-specific tuning of dendritic

feature selectivity by intracellular Ca2+ release. *Science, 375*.

https://doi.org/10.1126/science.abm1670

Owald, D., Felsenberg, J., Talbot, C. B., Das, G., Perisse, E., Huetteroth, W., &

Waddell, S. (2015). Activity of Defined Mushroom Body Output Neurons

Underlies Learned Olfactory Behavior in Drosophila. *Neuron, 86*, 417–427.

https://doi.org/10.1016/j.neuron.2015.03.025

Pathak, T., Agrawal, T., Richhariya, S., Sadaf, S., & Hasan, G. (2015). Store-

Operated Calcium Entry through Orai Is Required for Transcriptional Maturation

of the Flight Circuit in *Drosophila*. *The Journal of Neuroscience, 35*, 13784–

13799. https://doi.org/10.1523/jneurosci.1680-15.2015

Ping, Y., Waro, G., Licursi, A., Smith, S., Vo-Ba, D.-A., & Tsunoda, S. (2011).

Shal/Kv4 Channels Are Required for Maintaining Excitability during Repetitive

Firing and Normal Locomotion in Drosophila. *PLoS ONE, 6*, e16043.

https://doi.org/10.1371/journal.pone.0016043

Postma, M., & Goedhart, J. (2019). PlotsOfData—A web app for visualizing data

together with their summaries. *PLOS Biology, 17*, e3000202.

https://doi.org/10.1371/journal.pbio.3000202

Prakriya, M., & Lewis, R. S. (2015). STORE-OPERATED CALCIUM CHANNELS.

Physiol Rev, 307, 1383–1436. https://doi.org/10.1152/physrev.00020.2014

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S.,

Heyne, S., Dündar, F., & Manke, T. (2016). deepTools2: a next generation web

server for deep-sequencing data analysis. *Nucleic Acids Research, 44*(W1),

W160–W165. https://doi.org/10.1093/nar/gkw257

Richhariya, S., Jayakumar, S., Abruzzi, K., Rosbash, M., & Hasan, G. (2017). A

pupal transcriptomic screen identifies Ral as a target of store-operated calcium

entry in Drosophila neurons. *Scientific Reports, 7*(1), 1–12.

https://doi.org/10.1038/srep42586

Riemensperger, T., Völler, T., Stock, P., Buchner, E., & Fiala, A. (2005). Punishment

Prediction by Dopaminergic Neurons in Drosophila. *Current Biology*, *15*, 1953–1960. https://doi.org/10.1016/j.cub.2005.09.042

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, *43*, e47–e47. https://doi.org/10.1093/nar/gkv007

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2009). edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1), 139–140. https://doi.org/10.1093/bioinformatics/btp616

Rosenberg, S. S., & Spitzer, N. C. (2011). Calcium Signaling in Neuronal Development. *Cold Spring Harbor Perspectives in Biology*, *3*, a004259–a004259. https://doi.org/10.1101/cshperspect.a004259

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Shakiryanova, D., Morimoto, T., Zhou, C., Chouhan, A. K., Sigrist, S. J., Nose, A., Macleod, G. T., Deitcher, D. L., & Levitan, E. S. (2011). Differential Control of Presynaptic CaMKII Activation and Translocation to Active Zones. *Journal of Neuroscience*, *31*, 9093–9100. https://doi.org/10.1523/jneurosci.0550-11.2011

Sharma, A., & Hasan, G. (2020). Modulation of flight and feeding behaviours requires presynaptic IP3Rs in dopaminergic neurons. *ELife*, *9*. https://doi.org/10.7554/elife.62297

Shimojima, T. (2003). Drosophila FACT contributes to Hox gene expression through physical and functional interactions with GAGA factor. *Genes & Development*, *17*(13), 1605–1616. https://doi.org/10.1101/gad.1086803

Singh, S., & Wu, C. F. (1990). Properties of potassium currents and their role in membrane excitability in Drosophila larval muscle fibers. *Journal of Experimental Biology*, *152*, 59–76. https://doi.org/10.1242/jeb.152.1.59

Somasundaram, A., Shum, A. K., McBride, H. J., Kessler, J. A., Feske, S., Miller, R. J., & Prakriya, M. (2014). Store-Operated CRAC Channels Regulate Gene

Expression and Proliferation in Neural Progenitor Cells. *Journal of Neuroscience*, *34*, 9107–9123. https://doi.org/10.1523/jneurosci.0263-14.2014

Spitzer, N. C. (2012). Activity-dependent neurotransmitter respecification. *Nature Reviews Neuroscience*, *13*, 94–106. https://doi.org/10.1038/nrn3154

Stabell, M., Larsson, J., Aalen, R. B., & Lambertsson, A. (2007). Drosophila dSet2 functions in H3-K36 methylation and is required for development. *Biochemical and Biophysical Research Communications*, *359*, 784–789. https://doi.org/10.1016/j.bbrc.2007.05.189

Streb, H., Bayerdörffer, E., Haase, W., Irvine, R. F., & Schulz, I. (1984). Effect of inositol-1,4,5-trisphosphate on isolated subcellular fractions of rat pancreas. *The Journal of Membrane Biology*, *81*, 241–253. https://doi.org/10.1007/bf01868717

Sugie, A., Marchetti, G., & Tavosanis, G. (2018). Structural aspects of plasticity in the nervous system of Drosophila. *Neural Development*, *13*. https://doi.org/10.1186/s13064-018-0111-z

Sun, F., Zeng, J., Jing, M., Zhou, J., Feng, J., Owen, S. F., Luo, Y., Li, F., Wang, H., Yamaguchi, T., Yong, Z., Gao, Y., Peng, W., Wang, L., Zhang, S., Du, J., Lin, D., Xu, M., Kreitzer, A. C., … Li, Y. (2018). A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific Detection of Dopamine in Flies, Fish, and Mice. *Cell*, *174*, 481-496.e19. https://doi.org/10.1016/j.cell.2018.06.042

Sweeney, S. T., Broadie, K., Keane, J., Niemann, H., & O'Kane, C. J. (1995). Targeted expression of tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and causes behavioral defects. *Neuron*, *14*, 341–351. https://doi.org/10.1016/0896-6273(95)90290-2

Taber, K. H., & Hurley, R. A. (2014). Volume Transmission in the Brain: Beyond the Synapse. *The Journal of Neuropsychiatry and Clinical Neurosciences*, *26*, iv–4. https://doi.org/10.1176/appi.neuropsych.13110351

Takechi, H., Eilers, J., & Konnerth, A. (1998). A new class of synaptic response involving calcium release in dendritic spines. *Nature*, *396*, 757–760. https://doi.org/10.1038/25547

Tang, X., Li, T., Liu, S., Wisniewski, J., Zheng, Q., Rong, Y., Lavis, L. D., & Wu, C. (2022). Kinetic principles underlying pioneer function of GAGA transcription factor in live cells. *Nature Structural & Molecular Biology*, *29*, 665–676.

https://doi.org/10.1038/s41594-022-00800-z

Thillaiappan, N. B., Chakraborty, P., Hasan, G., & Taylor, C. W. (2019). IP3 receptors and Ca2+ entry. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, *1866*, 1092–1100. https://doi.org/10.1016/j.bbamcr.2018.11.007

Truszkowski, T. L. S., & Aizenman, C. D. (2015). Neurobiology: Setting the Set Point for Neural Homeostasis. *Current Biology*, *25*, R1132–R1133. https://doi.org/10.1016/j.cub.2015.10.021

Tsai, S.-Y., Chang, Y.-L., Swamy, K. B. S., Chiang, R.-L., & Huang, D.-H. (2016). GAGA factor, a positive regulator of global gene expression, modulates transcriptional pausing and organization of upstream nucleosomes. *Epigenetics & Chromatin*, *9*. https://doi.org/10.1186/s13072-016-0082-4

Tsao, C.-H., Chen, C.-C., Lin, C.-H., Yang, H.-Y., & Lin, S. (2018). Drosophila mushroom bodies integrate hunger and satiety signals to control innate food-seeking behavior. *ELife*, *7*. https://doi.org/10.7554/elife.35264

Tsukiyama, T., Becker, P. B., & Wu, C. (1994). ATP-dependent nucleosome disruption at a heat-shock promoter mediated by binding of GAGA transcription factor. *Nature*, *367*(6463), 525–532. https://doi.org/10.1038/367525a0

Venkiteswaran, G., & Hasan, G. (2009). Intracellular Ca2+ signalling and store operated Ca2+ entry are required in Drosophila neurons for flight. *Proceedings of the National Academy of Sciences*, *2009*.

Verma, S. K., Tian, X., LaFrance, L. V, Duquenne, C., Suarez, D. P., Newlander, K. A., Romeril, S. P., Burgess, J. L., Grant, S. W., Brackley, J. A., Graves, A. P., Scherzer, D. A., Shu, A., Thompson, C., Ott, H. M., Aller, G. S. Van, Machutta, C. A., Diaz, E., Jiang, Y., … Miller, W. H. (2012). Identification of Potent, Selective, Cell-Active Inhibitors of the Histone Lysine Methyltransferase EZH2. *ACS Medicinal Chemistry Letters*, *3*, 1091–1096. https://doi.org/10.1021/ml3003346

Viswanath, V., Story, G. M., Peier, A. M., Petrus, M. J., Lee, V. M., Hwang, S. W., Patapoutian, A., & Jegla, T. (2003). Opposite thermosensor in fruitfly and mouse. *Nature*, *423*, 822–823. https://doi.org/10.1038/423822a

Waddell, S. (2016). Neural Plasticity: Dopamine Tunes the Mushroom Body Output Network. *Current Biology*, *26*, R109–R112.

https://doi.org/10.1016/j.cub.2015.12.023

Wang, Y., Deng, X., & Gill, D. L. (2010). Calcium Signaling by STIM and Orai:

Intimate Coupling Details Revealed. *Science Signaling*, *3*, pe42–pe42.

https://doi.org/10.1126/scisignal.3148pe42

Weiss, S., Melom, J. E., Ormerod, K. G., Zhang, Y. V, & Littleton, J. T. (2019). Glial

Ca2+signaling links endocytosis to K+ buffering around neuronal somas to

regulate excitability. *ELife*, *8*. https://doi.org/10.7554/elife.44186

Wilkins, R. C., & Lis, J. T. (1999). DNA distortion and multimerization: novel

functions of the glutamine-rich domain of GAGA factor 1 1Edited by M. Yaniv.

Journal of Molecular Biology, *285*(2), 515–525.

https://doi.org/10.1006/jmbi.1998.2356

Xu, W., & Lipscombe, D. (2001). Neuronal CaV1.3α1 L-Type Channels Activate at

Relatively Hyperpolarized Membrane Potentials and Are Incompletely Inhibited

by Dihydropyridines. *The Journal of Neuroscience*, *21*, 5944–5951.

https://doi.org/10.1523/JNEUROSCI.21-16-05944.2001

Yagi, R., Mabuchi, Y., Mizunami, M., & Tanaka, N. K. (2016). Convergence of

multimodal sensory pathways to the mushroom body calyx in Drosophila

melanogaster. *Scientific Reports*, *6*. https://doi.org/10.1038/srep29481

Yeromin, A. V, Zhang, S. L., Jiang, W., Yu, Y., Safrina, O., & Cahalan, M. D. (2006).

Molecular identification of the CRAC channel by altered ion selectivity in a

mutant of Orai. *Nature*, *443*, 226–229. https://doi.org/10.1038/nature05108

Zambelli, F., Pesole, G., & Pavesi, G. (2009). Pscan: finding over-represented

transcription factor binding site motifs in sequences from co-regulated or co-

expressed genes. *Nucleic Acids Research*, *37*, W247–W252.

https://doi.org/10.1093/nar/gkp464

Zhai, B., Villén, J., Beausoleil, S. A., Mintseris, J., & Gygi, S. P. (2008).

Phosphoproteome Analysis of Drosophila melanogaster Embryos. *Journal of*

Proteome Research, *7*, 1675–1682. https://doi.org/10.1021/pr700696a

Zolin, A., Cohn, R., Pang, R., Siliciano, A. F., Fairhall, A. L., & Ruta, V. (2021).

Context-dependent representations of movement in Drosophila dopaminergic

reinforcement pathways. *Nature Neuroscience*, *24*, 1555–1566.

https://doi.org/10.1038/s41593-021-00929-y

Figure 1

Figure 1. Orai-mediated Ca²⁺ entry sets the gene expression profile of flight-promoting dopaminergic neurons in late development and early adulthood.

A. A schematic of Ca²⁺ release through the IP₃R and SOCE through STIM/Orai (upper panel) followed by representation of the wildtype (Ca²⁺ permeable) and mutant (Ca²⁺ impermeable) Orai channels (lower panel). **B**. Anatomical location of *THD'* DANs in the fly central brain immunolabelled for mCD8GFP (upper panel) followed by a cartoon of central brain DAN clusters. Scale bar indicates 20 μm. PPL1 and PPM3 clusters are labelled in red and green respectively. **C**. Measurement of flight bout durations demonstrate a requirement for Orai-mediated Ca²⁺ entry in *THD'* DANs and in 2 pairs of PPL1 DANs marked by the MB296B driver. **D**. *THD'* DANs require Ca²⁺ entry through Orai at 72-96 hours APF and 0-2 days post eclosion to promote flight. In **C** and **D** flight bout durations in seconds (s) are represented as a swarm plot where each genotype is represented by a different color, and each fly as a single data point. The Δ Flight parameter shown below indicates the mean difference for comparisons against the shared Canton S control and is shown as a Cumming estimation plot.

1263 Mean differences are plotted as bootstrap sampling distributions. Each 95%
1264 confidence interval is indicated by the ends of the vertical error bars. The same letter
1265 beneath each distribution refers to statistically indistinguishable groups after
1266 performing a Kruskal-Wallis test followed by a post hoc Mann-Whitney U-test
1267 ($p<0.005$). At least 30 flies were tested for each genotype. **E**. *THD'* DANs were labelled
1268 with cytosolic eGFP (10 μm scale), isolated using FACS and validated for enrichment
1269 of *GFP* mRNA by qRT-pCR (lower panel). The qRT-pCR results are from 4 biological
1270 replicates with different letters representing statistically distinguishable groups after
1271 performing a two-tailed t test ($p<0.05$). **F.** RNA-seq comparison of FAC sorted
1272 populations of GFP labelled THD' DANs from *THD'>GFP* and *THD'>GFP;Orai^{E180A}*
1273 pupal dissected CNSs. The RNA-Seq data is represented in the form of a volcano plot
1274 of fold change vs FDR. Individual dots represent genes, coloured in red (upregulated)
1275 or blue (downregulated) by >1 fold. **G.** Downregulated genes were identified by 3
1276 different methods of DEG analysis, quantified and compared as a Venn diagram. **H.**
1277 Gene expression trajectories of SOCE-induced DEGs plotted as a function of
1278 developmental time (modENCODE Consortium et al., 2010) and clustered into 3
1279 groups using k-means analysis. **I**. The relative proportion of downregulated,
1280 upregulated genes, and a random set of genes found in the clusters described in (**H**)
1281 indicate that 75% of downregulated genes exhibit a pupal peak of expression.

Figure 2

Figure 2. **Ca²⁺ entry through Orai regulates gene expression by Set2-mediated histone modification**. **A**. Scatterplot of GO categories enriched in SOCE-responsive genes. Individual GO terms are represented as differently colored circles, with radius size indicating number of genes enriched in that category. **B**. Fold change of SET domain containing genes as indicated. Individual circles on the Y-axis for each gene represent transcript variants pertaining to that gene. **C**. Transcripts of the H3K36 methyltransferase (left) are significantly diminished (right) in *THD'* DANs either upon loss of Orai function (*THD'>Orai^E180A*) or by knockdown of *Set2* (*THD'>Set2^RNAi*). qRT-PCRs were performed from FAC sorted *THD'* DANs with 3 biological replicates. Individual $2^{-\Delta\Delta CT}$ values are shown as points. Letters represent statistically distinguishable groups after performing an ANOVA and post hoc Tukey test (p<0.05). **D**. Significant rescue of flight bout durations seen in *THD' > Orai^E180A* flies by

overexpression of *Set2* and by knockdown of the *Kdm4B* demethylase indicating a net
requirement for H3K36me3. Flight durations of single flies are depicted as swarm plots
and the Δ flight parameter is shown below. Both were measured as described in the
legend to Figure 1. N = 30 or more flies for each genotype. Letters represent
statistically distinguishable groups after performing an ANOVA and post hoc Tukey
test (p<0.005) **E**. Representative images (upper panel) and quantification (lower
panel) from immunostaining of H3K36me3 and H2BmRFP in nuclei of *THD'* DANs
from at least 10 brains. Scale bar represents 5 μm. The boxplot represents individual
H3K36me3/H2BmRFP ratios from each *THD'* DAN for each genotype. Letters
represent statistically distinguishable groups after performing an ANOVA and post hoc
Tukey test (p<0.05) **F**. H3K36me3 enrichment over the gene bodies of SOCE-
responsive genes in wildtype fly heads represented in the form of a tag density plot.
G. H3K36me3 signal is enriched on WT SOCE-responsive genes with greater
downregulation upon loss of Orai function. Individual data points represent *WT*
H3K36me3 ChIP-Seq signals from adult fly heads represented as a boxplot (left) and
a regression plot (right; Pearson's correlation coefficient = 0.11) indicating a
correlation between extent of downregulation upon loss of SOCE and greater
enrichment of H3K36me3. **H**. SOCE-responsive genes are enriched in H3K36me3
signal as compared to H3K27me3 signal as measured from relevant ChIP-Seq
datasets. Adult fly head ChIP-Seq datasets for measurements in **F** to **H** were obtained
from modEncode (modENCODE Consortium et al., 2010). **I**. Schematic
representation of how Orai mediated Ca^{2+} entry regulates a balance of two opposing
epigenetic signatures in developing DANs. **J**. Pharmacological inhibition of H3K27me3
using GSKS343 (left) in *THD'>Orai^{E180A}* flies results in a dose-dependent rescue of
flight bout durations (right). Flight assay measurements from N> 30 flies, are
represented as described earlier. Letters represent statistically distinguishable groups
after performing an ANOVA and post hoc Tukey test (p<0.005).

Figure 3

Figure 3. Orai-mediated Ca^{2+} entry potentiates cellular Ca^{2+} responses to
cholinergic inputs through Set2 and a transcriptional feedback loop
A. A schematic of intracellular Ca^{2+} signaling downstream of neuromodulatory
signaling where activation of mAChR stimulates intracellular Ca^{2+} release through the
IP$_3$R followed by SOCE through STIM/Orai. **B.** Cholinergic inputs by addition of
carbachol (CCh) evoke Ca^{2+} signals as measured by change in the fluorescence of
GCaMP6m in *THD'* DANs of ex-vivo brains. Representative GCaMP6m images of
THD' DANs are shown with baseline and peak evoked responses in the indicated
genotypes. Scale bar = 10 μM. **C**. Median GCaMP6m responses plotted as a function
of time. A shaded region around the solid line represents the 95% confidence interval
from 4-5 cells imaged per brain from 10 or more brains per genotype. **D.** Individual
cellular responses depicted as a paired plot where different letters above indicate
statistically distinguishable groups after performing a Kruskal Wallis Test and a Mann-
Whitney U-test (p<0.05). **E**. qRT-PCR measurements of *itpr*, *mAChR*, *Stim* and *Orai*
from FAC sorted *THD'* DANs obtained from three biological replicates of appropriate
genetic backgrounds. The genotypes include DANs with loss of cellular Ca^{2+}
responses (*THD'>Orai^{E180A}*; and *THD'>Set2^{RNAi-1}*), rescue of cellular Ca^{2+} response by
overexpression of *Set2* (*THD'>Orai^{E180A}*; *Set2^{OE}*) and *Set2* overexpression. Bar plots
indicate mean expression levels in comparison to *rp49*, with individual data points
represented as hollow circles. The letters above indicate statistically indistinguishable
groups after performing a Kruskal Wallis Test and a Mann-Whitney U-test (p<0.05). **F.**

Schematic representation of a transcriptional feedback loop downstream of cholinergic
stimulation in *THD'* DANs.

1364

Figure 4

Figure 4. Identification of Trl as an SOCE responsive transcription factor
A. Schematic of motif enrichment analysis for identification of putative SOCE
dependent transcription factors (TFs). **B.** Candidate TFs identified (upper panel) and
their expression through development (lower panel; modEncode(modENCODE
Consortium et al., 2010)) suggests Trl as a top SOCE-responsive candidate TF. **C**.
Genetic depletion of *Trl* in *THD'* DANs results in significant flight defects, that can be
rescued by overexpression of either *STIM* or *Set2*. **D.** Heteroallelic combination of the
Trl hypomorphic allele (*Trl^13C*) with a *STIM* deficiency causes significant flight deficits,
that can be rescued by overexpression of *STIM* or *Set2*. **E.** qRT-PCRs measured
relative to *rp49* show reduced *Set2, itpr, and mAChR* levels in *THD'* DANs upon *Trl*
knockdown with *Trl^RNAi-1*. Individual data points of 4 biological replicates are shown as
circles and mean expression level as a bar plot (± SEM). The letters above indicate
statistically indistinguishable groups after performing a Kruskal Wallis Test and a
Mann-Whitney U-test (p<0.05). **F**. A representative western blot (left) showing reduced
H3K36me3 levels in a *Trl* mutant combination. Quantification of H3K36me3 from 3
biological replicates of *WT* (control) and *Trl* mutant brain lysates (panel on the right).
Letters indicate statistically indistinguishable groups after performing a two-tailed t-
test (p<0.05). **G**. Knockdown of *Trl* in *THD'* neurons attenuate Ca^{2+} response to CCh
as shown in representative images of GCaMP6m fluorescence quantified in (**H-I**).
Scale bar = 10 μM. *Set2* overexpression in the background of *THD'>Trl^RNAi* rescues
the cholinergic response (**G-I**). Quantification of Ca^{2+} responses are from 10 or more
brains per genotype and were performed as described in the legend to Figure 3. The

 letters above indicate statistically indistinguishable groups after performing a Kruskal
Wallis Test and a Mann-Whitney U-test (p<0.05).

Figure 5

Figure 5. Trl activity downstream of SOCE is required for *THD'* DAN activity.
A. Overexpression of *WT Trl* is insufficient to rescue flight deficits in *THD'>Orai^{E180A}*
flies as evident from flight bout measurements in the indicated genotypes. Flight assay
measurements are as described earlier. **B**. *Trl* transcript levels are not altered in
THD'>Orai^{E180A} neurons with loss of SOCE. qRT-PCR data are measured relative to
rp49. The bar plot indicates mean expression levels, with individual data points
represented as circles. The letters above indicate statistically indistinguishable groups
from 3 independent biological replicates after performing a two-tailed t-test (p<0.05).
C. Schematic with possible Ca^{2+} - mediated activation of Trl downstream of Orai-
mediated Ca^{2+} entry (SOCE). **D.** Genes encoding ion channels are enriched among
SOCE responsive genes in *THD'* DANs as determined by GO analysis. Circles of
varying radii are scaled according to the number of genes enriched in that category.
E. Downregulation of individual ion channel genes depicted as a heatmap. **F**. Number
of Trl binding sites (GAGA repeats) shown as a bar plot in the regulatory regions (2 kb

Figure 6

Figure 6. fpDAN excitability requires Orai-mediated Ca^{2+} entry acting through *Set2* -mediated VGCC gene expression. KCl-induced depolarizing responses in *fp*DANs of the indicated genotypes measured using GCaMP6m indicate a requirement for Orai and *Set2*. Representative images **A. (**scale bar = 10 μM), quantified in **B.** and **C**. KCl evoked responses are modulated upon treatment with 10 μM TTX (magenta), 10 μM Nimodipine (orange), and upon *cac^RNAi* (purple)- representative images (scale bar = 10 μM), (**D**), quantified in **E**. Median KCl-evoked GCaMP6m responses plotted as a function of time. The solid line indicates the time point of addition of 70 mM KCl. KCl responses are quantified as a paired plot of peak responses and **F**. with letters

representing statistically indistinguishable groups as measured using a Kruskall Wallis
test and post hoc Mann-Whitney U test ($p < 0.05$). Quantification of Ca^{2+} responses are
from 10 or more brains per genotype and were performed as described in the legend
to Figure 3. **G.** Schematic of a typical VGCC and its constituent subunits. **H.** Heatmap
of $2^{\wedge}\text{-}\Delta\Delta CT$ values measured using qRT-PCRs from sorted *THD'* DANs show a
reduction in the expression of VGCC subunit genes upon loss of Orai function and a
rescue by *Set2* overexpression. **I**. Flight assays representing the effect of various
VGCC subunit gene RNAis showing flight defects to varying extents. Overexpression
of the key VGCC subunit gene-*cac* is required (**I**) but not sufficient (**J**) for restoring
flight defects caused by loss of *Orai* function. Flight assay measurements are as
described earlier.

Figure 7

 Figure 7. SOCE-mediated gene expression sets the excitability threshold during
pupal development   A. Altering excitability in *THD'* DANs using *Kir2.1* or *GtACR2*
mediated inhibition or *CsChrimson* or *TrpA1* during the critical 72-96 hour APF
developmental window results in significant flight defects. Orai loss of function
phenotypes can be rescued by overexpression of *NachBac* or *CsChrimson* in terms
of flight bout durations (**B, E**) and depolarizing KCl responses (**C, D, D, G**). **H**.
Schematic for Orai-mediated Ca^{2+} entry regulating expression of key genes that
regulate the excitability threshold of dopaminergic neurons regulating flight during a
critical developmental window. Flight assays are represented as described earlier,
(n>30). Ca^{2+} responses were quantified as described in the legend to Figure 3 and
were from 10 or more brains per genotype. Letters above each genotype represent
statistically indistinguishable groups as measured using a Kruskall-Wallis test and post
hoc Tukey test (p<0.05).

Figure 8

Figure 8. *THD'* **DANs require SOCE-mediated gene expression for neuromodulatory dopamine release in the mushroom body γ lobe. A.** Schematic of a KC-DAN-MBON tripartite synapse at the $\gamma 2\alpha'1$ MB lobe (upper), innervated by fpDANs (lower). The $\gamma 2\alpha'1$ MB lobe is marked by a red circle and arrow and the fpDAN clusters are marked with a green arrow (PPL1) and blue arrow (PPM3) . Scale bar=50 μM. **B**. CCh-evoked DA release measured at the $\gamma 2\alpha'1$ MB lobe using the GRAB-DA sensor expressed in *THD'* DANs. Representative GRAB-DA images of *THD'* DANs are shown with baseline and peak evoked responses in the indicated genotypes. Scale bar = 10 μM. **C**. Median GRAB-DA responses plotted as a function of time. A shaded region around the solid line represents the 95% confidence interval from 10 or more brains per genotype. **D.** Individual cellular responses depicted as a paired plot of peak responses where different letters above indicate statistically distinguishable groups after performing a Kruskal Wallis Test and a Mann-Whitney U-test. **E**. Schematic describing how Orai-mediated Ca^{2+} entry regulates expression of key genes that control neuronal activity in fpDANs.