**RAPSYN-Mediated Neddylation of BCR-ABL Alternatively Determines the Fate of Philadelphia Chromosome-positive Leukemia**

Mengya Zhao[1†], Beiying Dai[2†], Xiaodong Li[1†], Yixin Zhang[2†], Chun Qiao[2, 3], Yaru Qin[2], Zhao Li[1], Qingmei Li[1], Shuzhen Wang[1]\*, Yong Yang[2]\*, Yijun Chen[1, 2, 4]\*

[1]Laboratory of Chemical Biology, School of Life Science and Technology, China Pharmaceutical University, Nanjing, 211198, China

[2]State Key Laboratory of Natural Medicines, China Pharmaceutical University, Nanjing, 211198, China

[3]Department of Hematology, The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, Nanjing, 210029, China

[4]Chongqing Innovation Institute of China Pharmaceutical University, Chongqing 401135, China

[†] These authors contributed equally

\***Correspondence**: Yijun Chen (yjchen@cpu.edu.cn), Yong Yang (yy@cpu.edu.cn) and Shuzhen Wang (shuzhenwang@cpu.edu.cn)

**Abstract**

Philadelphia chromosome-positive (Ph$^+$) leukemia is a fatal hematological malignancy. Although standard treatments with tyrosine kinase inhibitors (TKI) have achieved remarkable success in prolonging patient survival, intolerance, relapse and TKI resistance remain serious issues for patients with Ph$^+$ leukemia. Here, we report a new leukemogenic process in which RAPSYN and BCR-ABL co-occur in Ph$^+$ leukemia, and RAPSYN mediates the neddylation of BCR-ABL. Consequently, neddylated BCR-ABL enhances the stability by competing its c-CBL-mediated degradation. Furthermore, SRC phosphorylates RAPSYN to activate its NEDD8 E3 ligase activity, promoting BCR-ABL stabilization and disease progression. Moreover, in contrast to *in vivo* ineffectiveness of PROTAC-based degraders, depletion of RAPSYN expression or its ligase activity decreased BCR-ABL stability and, in turn, inhibited tumor formation and growth. Collectively, these findings represent an alternative to tyrosine kinase activity for the oncoprotein and leukemogenic cells and generate a rationale of targeting RAPSYN-mediated BCR-ABL neddylation for the treatment of Ph$^+$ leukemia.

**Introduction**

Philadelphia chromosome-positive (Ph[+]) leukemia is a myeloproliferative neoplasm characterized by the reciprocal translocation between the long arms of chromosome 9 and 22, t (9;22) (q34.1; q11.2) ( de Klein et al., 1982; Deininger et al., 2000). This cytogenetic abnormality results in a *BCR-ABL* fusion gene, which encodes the chimeric protein BCR-ABL with enhanced tyrosine kinase activity (Cortes et al., 2021). Based on its oncogenic role in Ph[+] leukemia, BCR-ABL has been regarded as the most pivotal target for Ph[+] leukemia therapy, especially for chronic myeloid leukemia (CML). Tyrosine kinase inhibitors (TKI) have been the main treatment option for Ph[+] leukemia, remarkably prolonging the patients' life span and improving their quality of life (Hochhaus et al., 2020; Jabbour and Kantarjian, 2020). However, most patients develop TKI resistance and relapse after long-term treatment (Braun et al., 2020). It is worth noting that the increase of BCR-ABL expression can affect the sensitivity to TKIs and eventually determine the rate of TKI resistance in patients with Ph[+] leukemia in addition to the mutations in the kinase domain (Jabbour et al., 2007). Mutations in the kinase domain can change the conformation of BCR-ABL, thus interfering with the binding between TKIs and BCR-ABL and resulting in decreased therapeutic efficacy (Lussana et al., 2018). In parallel, the increase of BCR-ABL expression can affect the sensitivity to TKIs and eventually determine the rate of disease progression and TKI resistance in patients with Ph[+] leukemia (Barnes et al., 2005). Therefore, effective degradation of BCR-ABL could address the issues on TKI resistance and leukemia-initiating cells (LICs), and PROTAC-based protein degradation strategy may represent a new therapeutic approach (Bekes et al., 2022). Currently, based on different ubiquitin E3 ligases, including VHL, CRBN and IAP, PROTAC-based degraders at nM level have shown significant degradation of BCR-ABL in CML cell lines, cell lines carrying mutations in BCR-ABL as well as patient-derived primary cells containing multiple BCR-ABL mutations ( Demizu et al., 2016; Lai et al., 2016; Shimokawa et al., 2017; Zhao et al., 2019; Burslem et al., 2019; Liu et al., 2022) . Unfortunately, the excellent cellular activity by the PROTAC-based degraders

has not been able to translate to *in vivo* efficacy, even in rare examples of xenografted

mouse models ( Zhao, Ren et al., 2019; Jiang et al., 2021a), resulting in uncertain

usefulness of these degraders. Nonetheless, the unsatisfactory results are not really

surprising because the underlying mechanism of elevated BCR-ABL expression

remains largely unclear.

Receptor-associated protein of the synapse (RAPSYN) has been identified as a

classic synaptic adaptor protein that binds to the acetylcholine receptor (AChR) and

several cytoskeleton-associated proteins, contributing to AChR clustering and

neuromuscular junction (NMJ) formation (Huh and Fuhrer, 2002; Witzemann et al.,

2013). Later, RAPSYN was found to exert NEDD8 E3 ligase activity to catalyze the

neddylation for AChR aggregation.(Li et al., 2016) Despite the extensive studies of

RAPSYN in muscular and neuronal cells and tissues (Legay and Mei, 2017; Li et al.,

2018),  with regard to its involvement in leukemogenesis, there is no available

information thus far except for our previous finding. Previously, RAPSYN was found to

be located in the cytosol of the typical Ph[+] leukemia cell line K562 when a small

molecule was used to probe its binding proteins using a proteomics approach (Wang

et al., 2015). Because of its newly identified E3 ligase activity for neddylation and its

occurrence in the Ph[+] leukemia cell line, we speculated that RAPSYN might contribute

to Ph[+] leukemia development through its enzymatic activity instead of only serving as

a scaffolding protein.

As a type of post-translational modification (PTM), neddylation is sequentially

catalyzed by the neuronal precursor cell-expressed developmentally downregulated

protein 8 (NEDD8)-activating enzyme E1 (NAE1), NEDD8-conjugating enzyme E2

(UBE2M/UBC12 or UBE2F), and a substrate-specific NEDD8 E3 ligase to complete

the covalent conjugation of NEDD8 to a lysine residue of its substrates ( van der Veen

and Ploegh, 2012; Enchev et al., 2015). The neddylation of proteins can be reversed

by deneddylases such as NEDP1. In the last two decades, accumulating evidence

indicated the strong involvement of dysregulated neddylation in tumor progression,

neurodegenerative and cardiac diseases, aberrant immunoregulation and others ( Ying

et al., 2018; Zhou et al., 2019; Li et al., 2020; Yao et al., 2020), which rationalizes the

modulation of neddylation as a feasible therapeutic strategy.

In this study, we report that RAPSYN is highly expressed along with BCR-ABL in

patients with Ph[+] leukemia and promotes disease progression, presumably by

stabilizing the BCR-ABL fusion protein *via* neddylation. The neddylation of BCR-ABL

by RAPSYN subsequently competes its ubiquitination-dependent degradation to

increase the stability of BCR-ABL. Additionally, the NEDD8 E3 ligase activity of

RAPSYN can be substantially increased by SRC-mediated phosphorylation, leading

to enhanced stability and activity of RAPSYN.

**Results**

**High protein levels of RAPSYN promoted Ph[+] leukemia progression**

Prior to investigating the biological roles of RAPSYN in the pathogenesis of Ph[+]

leukemia, its expression at both mRNA and protein levels was analyzed. Neither a

publicly available database nor our collection of patient samples and cell lines showed

a significant increase in mRNA levels (Figure1-figure supplement 1A-C). The protein

levels of RAPSYN were substantially elevated in the peripheral blood mononuclear

cells (PBMC) of Ph[+] CML (#8-11) and the bone marrow of ALL (#7) patient samples in

comparison to that of healthy donors (#1-6), which was in a direct accordance with the

expression of BCR-ABL (Figure 1A). This co-expression of RAPSYN and BCR-ABL

was also found in Ph[+] cell lines (Figure 1B), suggesting that the function of RAPSYN

in Ph[+] leukemia could be closely related to BCR-ABL.

To examine the relationship between RAPSYN and Ph[+] leukemia progression, we

first performed knockdown of its expression by using shRNAs. Marked reduction of

RAPSYN levels and notable cytotoxicity were observed in all tested Ph[+] leukemia cells

transduced with RAPSN shRNAs (Figure 1C, Figure1-figure supplement 1D-F),

indicating the dependence on the presence and expression level of BCR-ABL.

Conversely, exogenous expression of RAPSN rescued Ph[+] leukemia cells from

shRNA-generated toxicity (Figure 1D). Knockdown of RAPSYN also changed the

phenotypes of Ph[+] leukemia cells, including proliferation, G0/G1 cell cycle arrest and

apoptosis (Figure 1E-G, Figure1-figure supplement 1G, H**)**.

Next, we subcutaneously implanted empty body transduced sh*RAPSN*#3 or K562

cells into NCG mice to establish a cell line derived xenotransplantation mouse model

(Figure 1H). Tumor growth was significantly inhibited by RAPSYN silencing (Figure 1I,

J, Figure1-figure supplement 1I). Meanwhile, immunoblotting of tumor samples

showed a notable downregulation of RAPSYN expression, along with the reduction of

BCR-ABL levels (Figure 1K). After knockout of RAPSYN for remarkable depletion of

BCR-ABL expression in K562 cells (Figure 1L, Figure1-figure supplement 1K), these

cells along with the empty vector-transduced K562 cells were intravenously injected

into NCG mice to establish the leukemogenic mouse model (Figure 1H). Consequently,

the survival of tumor-bearing mice was profoundly prolonged by the knockout of

RAPSYN compared to the controls (Figure 1M). Altogether, our findings indicate that

RAPSYN is highly expressed at protein level with the accordance to BCR-ABL in Ph$^+$

leukemia and its depletion results in inhibiting the progression of Ph$^+$ leukemia.

**RAPSYN directly neddylated BCR-ABL**

Previous reports determined that both nicotinic AChR subunit $\alpha_7$ and muscarinic AChR

subtypes $M_2$, $M_3$, and $M_4$ were involved in the the cell proliferation of K562 cells

(Cabadak et al., 2011; Önder Narin et al., 2021). Furthermore, RAPSYN was found to

exert its NEDD8 E3 ligase activity toward AChR in neuronal systems (Li, Cao et al.,

2016). To determine whether RAPSYN functioned in a similar manner in leukemogenic

cells, we investigated whether RAPSYN promoted Ph$^+$ leukemia progression through

neddylation of AChRs. Despite AChR subunits $\alpha_7$, $M_2$, $M_3$, and $M_4$ were expressed in

all tested cells, no change in their neddylation was observed upon RAPSYN ablation

(Figure2-figure supplement 2A, B). On the basis of the co-expression of RAPSYN and

BCR-ABL, we postulated that RAPSYN could specifically mediate neddylation of BCR-

ABL to promote Ph$^+$ leukemia development.

To test this hypothesis, reciprocal immunoprecipitation was performed to reveal a

strong interaction between RAPSYN and BCR-ABL in Ph$^+$ leukemia cells (Figure 2A,

Figure2-figure supplement 2C). Similar results were obtained with exogenous

expression in HEK293T cells (Figure 2B), further confirming the specific interaction of

RAPSYN with BCR-ABL. Furthermore, GST pull-down assay with purified proteins

displayed specific binding of GST-tagged RAPSYN to His-tagged BCR-ABL (Figure

2C), indicating that BCR-ABL is the primary target of RAPSYN-mediated neddylation.

Domain mapping revealed that the Δ1 domain (1-927 aa) of BCR-ABL was responsible

for the interaction with RAPSYN (Figure 2D).

Next, we studied whether RAPSYN could directly mediate BCR-ABL neddylation.

Preeminent BCR-ABL neddylation was detected in all $Ph^+$ leukemia cell lines (Figure

2E, Figure2-figure supplement 2D). Treatment with the NAE1 inhibitor, MLN4924,

significantly dampened the neddylation of BCR-ABL (Figure 2F, Figure2-figure

supplement 2E). In addition, the mutation of two glycine residues at the C-terminus of

NEDD8 required for its covalent conjugating ability, or the co-expression of NEDP1

(NEDD8-specific protease 1) essentially diminished the neddylation of BCR-ABL

(Figure 2G, H). As shown in Figure 2I, we co-expressed either WT-RAPSYN or its

C366A mutant along with BCR-ABL and NEDD8, revealing that mutation of Cys to Ala

at the catalytic residue C366 significantly decreased the neddylation level of BCR-ABL.

Additionally, knockout of RAPSYN abrogated BCR-ABL neddylation in the cells, and

this effect was restored by transduction of *RAPSN* cDNA (Figure 2J). These results

were further corroborated by *in vitro* experiments, which showed that BCR-ABL could

hardly be neddylated in the absence of RAPSYN (Figure 2K). Consistently, the amount

of neddylated BCR-ABL was markedly reduced in tumors generated by K562 cells

transfected with sh*RAPSN*#3 (Figure 2L), indicating an essential role of the ligase

activity of RAPSYN in BCR-ABL neddylation.

In addition, neither overall BCR-ABL expression nor its neddylation levels were

affected after the knockdown of AChRs (Figure2-figure supplement 2F). Similarly,

modulation of AChR activities and their downstream PKC-RAS-ERK and JAK2-AKT

signaling pathways ( Kawamata et al., 2011; Aydin et al., 2013) by either an agonist

(carbachol) (Jakubik et al., 2008) or antagonists (benzethonium and homatropine)

(Durieux and Nietgen, 1997) did not alter the expression or neddylation status of BCR-

ABL (Figure2-figure supplement 2G, H).

Subsequently, we tried to identified specific modification sites on BCR-ABL. The

purified proteins were used for *in vitro* neddylation reactions, and the target bands were

digested with trypsin for LC-MS/MS analyses. Eight lysine residues were found to be

potential NEDD8 accepting sites in BCR-ABL (Figure2-figure supplement 3). To

confirm these modification sites, a series of individual Lys-to-Arg mutants were

generated. Except for K257, neddylation levels of BCR-ABL at other candidate sites

were all significantly reduced, confirming the modification sites of these Lys residues

(Figure 2M).

**RAPSYN attenuated c-CBL-mediated BCR-ABL ubiquitination and degradation**

As decreased neddylation of BCR-ABL following either MLN4924 treatment or

RAPSYN/KO was accompanied by a strong decline in its overall protein expression

level (Figure 2F, 3A-B, Figure2-figure supplement 2E, 4), we asked whether RAPSYN-

mediated BCR-ABL neddylation affects protein stability. Subsequently, the protein

synthesis inhibitor CHX was applied to K562 cells transduced with vectors encoding

doxycycline-inducible *RAPSN* shRNA #3. Indeed, the expression levels of BCR-ABL

declined much faster in cells with the induction of shRNA expression (Figure 3C).

Meanwhile, we found that a clear inverse correlation between the neddylation and

ubiquitination levels of BCR-ABL was observed (Figure 3D, E). BCR-ABL ubiquitination

was remarkably reduced in the cells transfected with NEDD8 (Figure 3F). Consistent

with these results, treatment of the cells with the proteasome inhibitor MG132

significantly increased the amount of ubiquitinated BCR-ABL accompanied by the

decrease of BCR-ABL neddylation (Figure 3G).

To clarify the molecular basis of the battle between BCR-ABL neddylation and its

ubiquitination, we detected that whether RAPSYN competes for binding to BCR-ABL

with c-CBL, a reported E3 ligase mediating BCR-ABL ubiquitin-proteasome

degradation (Mao et al., 2010). As a result, exogenous expression of RAPSYN

interfered with the interactions between BCR-ABL and c-CBL, whereas RAPSYN

ablation in K562 cells promoted c-CBL binding to BCR-ABL (Figure 3H, I). These data

indicate that RAPSYN competes with c-CBL for binding to BCR-ABL, leading to

subsequent BCR-ABL neddylation to enhance BCR-ABL stability by counteracting its

proteasomal degradation.

**SRC-mediated phosphorylation stabilized RAPSYN by repressing its**

**proteasomal degradation**

SRC-family protein tyrosine kinases are capable of phosphorylating RAPSYN in

neuronal system, among which SRC exerts the strongest function (Mohamed and

Swope, 1999). In addition, SRC has been shown to be highly expressed in primary

CML cells (Yang et al., 2017). We then studied whether SRC acts as an upstream

regulator to mediate RAPSYN. SRC inhibition with saracatinib or shRNA not only

significantly downregulated phosphorylated (Tyr418) SRC, but also inhibited the

phosphorylation of endogenous RAPSYN, resulting in a substantial decline in its

protein level, whereas heterologous expression of *SRC* increased RAPSYN

phosphorylation (Figure 4A-C, Figure4-figure supplement 5A). Furthermore, *in vitro*

incubation with recombinant RAPSYN, SRC, and ATP resulted in strong

phosphorylation of RAPSYN, which could be fully abrogated by saracatinib treatment

(Figure 4D). LC-MS/MS analyses indicated that Tyr residues at positions 59, 152, and

336 in RAPSYN are potential phosphorylation sites by SRC (Figure4-figure

supplement 5B). Then, after mutagenesis of these sites from Tyr to Phe, Y336, an

evolutionarily conserved Tyr residue, was confirmed to be the primary site of RAPSYN

phosphorylation (Figure 4E, Figure4-figure supplement 5C). As SRC has no effect on

*RAPSN* mRNA levels (Figure4-figure supplement 5D, E), implying that SRC-mediated

phosphorylation also affects RAPSYN stability. In fact, Ph[+] leukemia cells were treated

with saracatinib, SRC silencing or mutation of the key phosphorylation site significantly

accelerated the diminishment of RAPSYN expression following CHX treatment,

conversely, expressing exogenous SRC cDNA prolonged the half-life of RAPSYN

(Figure 4F-I).

To explore the molecular mechanisms responsible for the increased stability of

phosphorylated RAPSYN, Ph[+] leukemia cells were treated with saracatinib or

transduction of sh*SRC* followed by incubation with MG132. In all circumstances,

MG132 could rescue the decrease of RAPSYN induced by saracatinib treatment or

sh*SRC* knockdown (Figure 4J, K). Clearly, the specific phosphorylation of RAPSYN at

Y336 by SRC led to its increased stability by preventing the proteasomal degradation,

thereby maintaining the high levels of RAPSYN in Ph$^+$ leukemia.

**261 Phosphorylated RAPSYN potentiated its NEDD8 E3 ligase activity and promoted**

**262 BCR-ABL stabilization**

To dissect the role of SRC-mediated phosphorylation of RAPSYN, we tested whether

phosphorylation of RAPSYN at Y336 affects its ligase activity. Immunoblotting revealed

saracatinib treatment or SRC silencing reduced BCR-ABL neddylation and its protein

expression, while exogenous expression of *SRC* cDNA strongly increased it (Figure

5A-C). Additionally, co-expression in HEK293T cells showed that Y336F mutation had

no impact on BCR-ABL neddylation compared to co-transfection with SRC (Figure 5D).

These results were supported by stronger neddylation of endogenous BCR-ABL in

cells overexpressing WT RAPSYN, but not in the Y336F mutant (Figure 5E).

Furthermore, protein turnover rates of BCR-ABL were determined in Ph$^+$ leukemia cells

and the cells with exogenous *RAPSN*$^{Y336F}$ expression displayed larger decrease in

BCR-ABL level than those expressing *RAPSN*$^{WT}$ (Figure 5F). Therefore, RAPSYN

phosphorylation at Y336 by SRC was a major contributing factor to its NEDD8 E3

ligase activity and BCR-ABL stability in Ph$^+$ leukemia cells.

**276 Phosphorylation of RAPSYN at Y336 promoted Ph$^+$ leukemia progression**

To assess the extent to which SRC-mediated phosphorylation of RAPSYN at Y336

contributes to the enhanced viability of RAPSYN-dependent Ph$^+$ leukemia cells, we

first identified a specific sh*SRC* by screening five candidates using toxicity tests and

then performing rescue experiments with *SRC* cDNA. Toxicity tests revealed that,

albeit to varying degrees, sh*SRC*#2, #4, and #5 induced cytotoxicity in all Ph$^+$ leukemia

cell lines (Figure 6A, Figure6-figure supplement 6A). However, exogenous *SRC* cDNA

expression only restored the growth of Ph$^+$ leukemia cells transduced with sh*SRC*#2

(Figure 6B, Figure6-figure supplement 6B, C).

Subsequently, we performed rescue experiments by introducing $RAPSN^{WT/Y336F}$

cDNA or an empty vector into shSRC#2-transducted Ph$^+$ leukemia cells and found that

virtually complete $RAPSN^{WT}$-induced rescue was detected in both cell lines, but

$RAPSN^{Y336F}$ exhibited no restoring effect (Figure 6C, D). In addition, transduction of

$RAPSN^{WT}$ cDNA conferred an increased resistance against saracatinib treatment,

whereas the expression of $RAPSN^{Y336F}$ cDNA did not affect the drug sensitivity of the

cells (Figure 6E, F). Furthermore, knockdown of RAPSYN substantially sensitized both

cell lines to saracatinib (Figure 6G).

In animal models, while the overall survival of mice intravenously injected with

K562 cells expressing control empty vector was significantly improved by either

saracatinib administration or shRNA-mediated SRC inhibition, overexpression of WT

RAPSYN fully counteracted these effects and shortened the lifespan of mice to the

levels comparable to those of mice injected with K562 cells without SRC inhibition. In

contrast, expression of exogenous $RAPSN^{Y336F}$ attenuated the protective effects of

SRC inhibition to a much lesser extent (Figure 6H-J). Taken together, these results

suggest that phosphorylation at Y336 by SRC is a major event in the pro-leukemogenic

functions of RAPSYN in Ph$^+$ leukemia development.

**Discussion**

In this study, we identified a novel role of RAPSYN in hematology. Indeed,

RAPSYN inhibition significantly suppressed the survival of Ph$^+$ leukemia. This

phenotype was found to be linked to the NEDD8 E3 ligase activity of RAPSYN, which

mediated BCR-ABL neddylation to enhance its stability for promoting leukemogenesis

(Figure 7).

Balanced protein synthesis and degradation are pivotal for maintaining protein

homeostasis and normal cellular function. Currently, a variety of degraders for BCR-

ABL by PROTAC strategy have been reported ( Demizu, Shibata et al., 2016; Lai,

Toure et al., 2016; Shimokawa, Shibata et al., 2017; Zhao, Ren et al., 2019; Burslem,

Schultz et al., 2019; Liu, Mi et al., 2022;), and this approach for the degradation of

BCR-ABL has appeared to be effective on overcoming drug-resistance in various cell

lines (Demizu, Shibata et al., 2016; Jiang, Wang et al., 2021a). However, very few reports with *in vivo* data in animal models have shown that the efficacy is way far from satisfactory ( Zhao et al., 2019; Jiang et al., 2021a), due likely to the inherited limitations of the degraders, such as unfavored physiochemical properties, significant off-target effects, various safety issues and others (Li and Song, 2020; Bekes, Langley et al., 2022). On the other hand, neddylation is a crucial type of PTM that modifies multiple Lys residues in BCR-ABL, competing with and shielding this oncoprotein from ubiquitination-mediated degradation, which provides a reasonable explanation on the poor *in vivo* efficacy of PROTAC-based degraders for BCR-ABL. Meanwhile, accumulating evidence has shown that targeting the neddylation process could be an appealing strategy for anticancer therapy (Bhalla and Fiskus., 2016; McGrail et al., 2020; Norton et al., 2021; Xie et al., 2021).  MLN4924, a NEDD8-activating E1 enzyme inhibitor, has been shown to inhibit the survival of both wild-type (WT) and T315I-BCR-ABL leukemia cells as well as leukemia-initiating cells ( Liu et al., 2018; Bahjat et al., 2019; Guo et al., 2019). Moreover, clinical trials of MLN4924 in combination with anticancer agents in acute myeloid leukemia (AML) have progressed to phase II (NCT03745352 and PEVENAZA [NCT04266795]) and III (PANTHER[NCT03268954] and PEVOLAM[NCT04090736]). However, the neddylation system works in a substrate- and context-dependent manner, which essentially defines its role in tumorigenesis as anti or pro, particularly relying on the substrate specificity of the NEDD8 E3 ligase. Neddylation can either facilitate ubiquitination-dependent degradation of its substrates, such as EGFR (Oved et al., 2006) and c-SRC (Lee et al., 2018), or enhance protein stability in the cases of HuR (Embade et al., 2012) and TGF-β type II receptor (Zuo et al., 2013). Thus, the antitumor effects of MLN4924 are the integrative outcome of inhibiting more *pro-* than *anti-* tumorigenic neddylation activities in the reported tumor types. Recent studies uncovered that neddylation could also inhibit tumor progression and MLN4924 stimulates tumor sphere formation and wound healing as well as promotes glycolysis (Zhou et al., 2016; Zhou, Jiang et al., 2019; Zhou and Sun, 2019). Therefore, rather

than suppressing the entire neddylation system to affect a wide range of proteins, targeted inhibition of substrate-specific NEDD8 E3 ligase, such as RAPSYN, might offer a potential therapeutic opportunity for more elegant anticancer intervention with fewer side effects.

Functional studies on RAPSYN have focused on its contribution to neuromuscular transmission (Xing et al., 2019; Xing et al., 2020). To date, whether RAPSYN is involved in biological processes other than the formation and maintenance of NMJ remains poorly understood. In this study, we found that RAPSYN promoted disease progression by neddylating BCR-ABL for its resistance to c-CBL-mediated proteasomal degradation. Intriguingly, although AChRs, the primary substrates of RAPSYN E3 ligase activity in neural and muscle cells, were also found to be expressed in Ph[+] leukemia cells, they were not identified as substrates of RAPSYN for neddylation, suggesting highly orchestrated tissue-specificity of the RAPSYN enzymatic activity on its substrates.

It is well known that the generation of BCR-ABL fusion protein is a decisive characteristic of Ph[+] leukemia (de Klein et al., 1982). Aside from the occurrence of RAPSYN in patients with Ph[+] leukemia, the present study revealed a fascinating finding that BCR-ABL expression levels were correlated with those of RAPSYN, demonstrating the specificity of RAPSYN-mediated neddylation of BCR-ABL. Moreover, as a new type of PTM for BCR-ABL, RAPSYN-mediated neddylation was found to compete c-CBL-mediated ubiquitination, causing a reduction in BCR-ABL degradation. Given the fact that the increase of BCR-ABL expression can affect the sensitivity to TKIs and eventually determine the rate of TKI resistance and LIC population in patients with Ph[+] leukemia ( Issaad et al., 2000; Barnes, Palaiologou et al., 2005), effective degradation of BCR-ABL is an alternative opportunity for the treatment. Although recent studies have greatly advanced our understanding of the regulation of BCR-ABL degradation (Burslem, Schultz et al., 2019; Shibata et al., 2020; Jiang et al., 2021b), most reported modulatory proteins are not ideal for translation to clinical settings because of their pivotal roles in sustaining normal hematological

functions. In contrast, RAPSYN was nearly unexpressed in the blood of healthy donors.

Thus, it is reasonable to expect that the inhibition of RAPSYN expression could lead

to cytotoxicity in Ph$^+$ leukemia with high specificity and marginal side effects.

Furthermore, our present results showed that knockdown of RAPSYN significantly

increased the sensitivity of leukemia cells to saracatinib, implying that a combination

of RAPSYN inhibition and TKI treatment can effectively control mutations and LIC-

derived TKI resistance in Ph$^+$ leukemia.

In summary, our work has uncovered the pivotal role that RAPSYN exerts its

NEDD8 E3 ligase activity in neddylating and stabilizing BCR-ABL in the pathogenesis

of Ph$^+$ leukemia and thus delineate it as a potential novel therapeutic target for the

treatment of Ph$^+$ leukemia. More importantly, our results shed a light on future

investigations that may help to extend to other cancer types for broadening our

understanding of RAPSYN's involvement in hematology and oncology.

**Materials and Methods**

**Human clinical samples**

This study was approved by the ethics committee of the First Affiliated Hospital of Nanjing

Medical University (2019-SR-485.A1). Human peripheral blood samples were obtained from

the remaining material utilized for routine laboratory tests at the First Affiliated Hospital of

Nanjing Medical University (Nanjing, China) and derived from 21 patients with Ph$^+$ chronic

myeloid leukemia (CML), and six healthy volunteers. And one human bone marrow sample of

Ph$^+$ acute lymphoblastic leukemia patient was obtained from the remaining material utilized for

routine laboratory tests at the First Affiliated Hospital of Nanjing Medical University (Nanjing,

China). Peripheral blood and bone marrow mononuclear cells were isolated by density gradient

centrifugation using the Ficoll® Paque Plus solution (17-1440-02, GE Healthcare).

**Cell culture**

K562 (female; CBP60529), MEG-01 (male; BP61104), and KU812 (male; BP60732) cells were

purchased from COBIOER and cultured in Roswell Park Memorial Institute 1640 medium

(RPMI 1640; KGM31800, KeyGEN BioTECH) containing 10-20% fetal bovine serum (FBS;

FS301-02, TransGen Biotech) and 100 mg/mL streptomycin/penicillin (FG101-01, TransGen

Biotech). HS-5 (male; CRL11882, ATCC), purchased from the American Type Culture Collection (ATCC) and HEK293T (KG405) from KeyGEN BioTECH were cultured in Dulbecco's modified Eagle's medium (DMEM; KGM12800, KeyGEN BioTECH) containing 10% FBS and 100 mg/mL streptomycin/penicillin. All cells were cultured in a humidified incubator with 5% $CO_2$ at 37°C. All the cell lines were authenticated using short tandem repeat matching analysis and tested negative for mycoplasma contamination.

**Animal studies**

Female NOD/ShiLtJGpt-Prkdcem26Cd52Il2rgem26Cd22/Gpt (NCG) mice (6-8 weeks), purchased from GemPharmatech Co., Ltd., were used for all in vivo studies. Mice were housed under specific pathogen-free conditions at 24 ± 1°C and 55 ± 5% humidity in a barrier facility with 12-h light-dark cycles. All animal experiments were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals with the approval of the Center for New Drug Evaluation and Research, China Pharmaceutical University (approval number: B20190925-1; Nanjing, China).

**Apoptosis assay**

A total of 1-5 x $10^5$ cells were washed with phosphate-buffered saline (PBS; 02-024-1ACS, Biological Industries) and resuspended in 100 μL of 1 × Annexin V binding buffer (E-CK-A211, Elabscience). The cell suspension was incubated with 2.5 μL of Annexin V-AF647 (E-CK-A211, Elabscience) and 2.5 μL of propidium iodide (PI; E-CK-A211, Elabscience) for 20 min in the dark, followed by the addition of 400 μL of 1 × Annexin V binding buffer, and detection by flow cytometry (Thermo Attune NxT, MA, USA).

**Cell viability assay**

Cells were seeded at a density of 5,000 (K562) or 20,000 cells (MEG-01) per well in round-bottom 96-well plates and incubated with different concentrations of saracatinib (AZD0530, Selleck) or the corresponding amount of solvent for 72 h. Cells were then transferred to flat-bottom 96-well plates for the determination of cell viability using the CCK-8 Cell Counting Kit (A311-01, Vazyme) following the manufacturer's instructions. Each experiment was performed at least three times for each cell line.

**Cell proliferation assay**

Cells were washed twice with PBS, resuspended with a cell number of 2 x 106 in 1 mL PBS,

and incubated with 2.5 µM carboxyfluorescein succinimidyl ester (CFSE) solution (1948076,

Thermo Scientific) or 5 µM carboxylic acid, acetate, and succinimidyl ester (SNARF-1) solution

(S-22801, Invitrogen) for 20 min at 37°C in the dark, respectively. Subsequently, 1 mL of FBS

was added to stop the staining, and the cells were washed twice with complete media. Cell

division was monitored by measuring CFSE or SNARF-1 dilution using flow cytometry via

channels BL1 and YL1.

**Cell cycle analysis**

439    A total of 1×106 cells were washed once with pre-chilled PBS, fixed in ice-cold 70% ethanol,

440    vortexing and kept for at least 20 min at -20°C. The fixed cells were washed twice with PBS

441    and stained with 20 µg/mL propidium iodide (PI) containing 100 mg/mL RNAse A (740505,

442    MACHEREY-NAGEL) for 15 min at room temperature. Stained nuclei were analyzed by flow

443    cytometry and quantified using FlowJo software (BD Biosciences, NJ, USA).

**Cell transfection and viral transduction**

Transfection of the indicated DNA plasmids into HEK293T cells was performed using

Lipofectamine 2000 (11668500, Thermo Fisher Scientific), according to the manufacturer's

instructions. Briefly, transfection of HEK293T was performed when cell confluency reached 60-

70%. Plasmids and Lipofectamine 2000 reagent were diluted in Opti-MEM medium (31985-070,

Thermo Fisher Scientific) and incubated for 5 min at room temperature. They were then mixed

together, incubated for another 20 min at room temperature, and added to the target cells. The

transfected cells were collected after 48 h for further analysis.

**GST pull-down assay**

Purified GST or GST-RAPSYN (GenBank: NM_005055.5; 1236 bp ORF sequence) proteins

were incubated with glutathione beads 4FF (SA010010, Smart-Lifescience) overnight at 4°C in

binding buffer (0.14 M NaCl, 2.68 mM KCl, 2 mM KH2PO4, 0.01 M Na2HPO4, 10 mM DTT, pH

7.4), respectively, and incubated with purified His-BCR-ABL (p210 BCR-ABL (b3a2); 6126 bp

ORF sequence) protein for another 4 h at 4°C. Beads were washed three times with washing

buffer(0.14 M NaCl, 2.68 mM KCl, 2 mM KH2PO4, 0.01 M Na2HPO4, 0.5 mM reduced GSH,

10 mM DTT, pH 7.4), eluted with elution buffer(0.14 M NaCl, 2.68 mM KCl, 2mM KH2PO4,

0.01M Na2HPO4, 10 mM reduced GSH, 10mM DTT, pH 7.4), and subjected to immunoblotting

detection.

**Immunoblotting**

The cells were lysed on ice with Nonidet® P-40 (NP-40) lysis buffer (150 mM NaCl, 100 mM

NaF, 50 mM Tris-HCl (pH 7.6), and 0.5% NP-40) supplemented with a protease inhibitor

cocktail (78446, Thermo Fisher Scientific). Lysates were centrifuged, quantified, subjected to

sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and transferred to

polyvinylidene difluoride (PVDF) membranes using a Bio-Rad transfer apparatus. Membranes

were blocked with 5% non-fat milk in Tris-buffered saline buffer containing 0.1% Tween-20

(TBST) at 20-25°C for 2 h, followed by incubation with primary antibody overnight at 4°C. The

membranes were then washed three times in TBST buffer and incubated with species-specific

HRP-conjugated secondary antibodies for 2 h at room temperature. Afterwards, the

membranes were washed three times in TBST buffer, developed using the enhanced

chemiluminescence (ECL) reagent, and exposed to the ChemiDoc Imaging System (Tanon,

Shanghai, China).

**Immunoprecipitation**

Immunoprecipitation assays were performed in accordance with the manufacturer's instructions.

Briefly, the cells were lysed on ice with NP-40 lysis buffer supplemented with a protease

inhibitor cocktail. Cell lysates were centrifuged, quantified, and incubated with the appropriate

primary antibody overnight at 4°C, and subsequently with protein A agarose beads (16-125,

Millipore) for another 4 h at 4°C. Agarose was washed three times with lysis buffer and eluted

with SDS-PAGE loading buffer. The eluted immunocomplexes were separated by SDS-PAGE

and transferred to PVDF membranes, which were probed with primary and corresponding

secondary antibodies, washed three times in TBST buffer, developed using the ECL reagent,

and exposed to the ChemiDoc Imaging System.

***In vitro* neddylation assay**

486    A 30 µL reaction mixture containing 2 mM ATP-Mg2+ (B-20, R&D), 50 ng E1(APPBP1/UBA3;

487    E-313-25, R&D), 400 ng E2 (UBE2M; E2-656-100, R&D), 0.25 µg NEDD8 (UL-812-500, R&D),

488   0.35 μg His-BCR-ABL, with or without 4.77 μg RAPSYN was incubated at 37°C for 4 h. Reaction

489   was terminated with SDS-PAGE loading buffer and assayed using immunoblotting.

***In vitro* phosphorylation assay**

491   A 40 μL reaction mixture containing 2 mM ATP-Mg2+, 3 μg RAPSYN, and 1 μg SRC (GenBank:

492   NM_005417.5; 1608 bp ORF sequence) protein, with or without 2 μM saracatinib (AZD0530,

493   Selleck), was incubated at 30°C for 30 min. Reaction was terminated with SDS-PAGE loading

494   buffer and assayed using immunoblotting.

**Animal experiments with mouse models**

Female NCG mice aged 6-8 weeks were used in all the animal experiments. In the

subcutaneous tumor experiment shown in Figure 1G, 20 mice were randomly divided into two

groups, followed by subcutaneous injection of 1×106 K562-shNC or K562-shRAPSYN #3 cells

in 60 μL Matrigel (354234, Corning) into the right foreleg. Tumor size was measured every two

days using a digital caliper. The tumor volume was quantified using the following equation:

tumor volume = 0.5 × (long diameter) × (short diameter) 2. When the average volume of the

control group exceeded 2,000 mm3, the mice were sacrificed. The tumors were separated, and

their weights were measured. As shown in the survival experiment in Figure 1J, NCG mice

were inoculated with K562-RAPSYNWT or K562-RAPSYNKO (1×107 cells/mouse) via the tail

vein. The survival time was recorded until the mice died. As shown in the survival experiment

of Figure 6I, 40 mice were randomly divided into 4 groups and intravenously inoculated with

K562-OE-NC (20 mice), K562-OE-RAPSYNWT (10 mice), or K562-OE-RAPSYNY336F (10

mice). From day 6 to 26 after tumor inoculation, 10 mice inoculated with K562-OE-NC were

treated with vehicle orally, while the other 30 mice inoculated with K562-OE-NC, K562-OE-

RAPSYNWT, or K562-OE-RAPSYNY336F, 10 mice in each group, were administered

saracatinib orally (50 mg/kg/d). The survival time was recorded until the mice died. In the

survival experiment shown in Figure 6J, 40 mice were randomly divided into four groups and

intravenously inoculated with double-transfected K562 cells, as indicated. The survival time

was recorded until the mice died.

**Identification of modification sites**

To determine which lysines in BCR-ABL were neddylated using NEDD8, a neddylation reaction

(50 µL) was performed. After incubation at 37°C for 4 h, the reaction was run on an SDS-PAGE

gel, silver-stained bands were excised and sent to BiotechPack Scientific Co., Ltd (Beijing,

China) for liquid chromatography-mass spectrometry (LC-MS/MS) analysis. To determine

which tyrosines in RAPSYN were phosphorylated by SRC, a phosphorylation reaction (50 µL)

was performed. After a 30 min reaction was run on an SDS-PAGE, silver-stained bands were

excised and sent to Applied Protein Technology Co., Ltd. (Shanghai, China) for liquid

chromatography-mass spectrometry (LC-MS/MS) analysis.

**Plasmid construction**

Eukaryotic expression vectors encoding His-, GST-, HA-, Myc-, or Flag-tagged proteins were

generated by inserting PCR-amplified fragments into the pcDNA3.1(+) mammalian expression

vector (V79020, Invitrogen). Eukaryotic expression vectors encoding green fluorescent protein

(GFP)-tagged proteins were generated by inserting PCR-amplified fragments into the pd1-

EGFP-N1 vector (6073-1, Clontech). Prokaryotic plasmids encoding GST-fusion proteins were

constructed using the pGEX-4T-1 bacterial expression vector (27-4580-01, Addgene). Mutants

of His-, HA- GST-, or GFP-tagged proteins were generated using the QuickMutation Site-

Directed Mutagenesis Kit (D0206, Beyotime) according to the manufacturer's instructions.

Briefly, whole plasmid DNA was amplified by polymerase chain reaction (PCR) for 20 cycles

with specific mutant primers (Supplemental Table S1) using the QuickMutation site-directed

mutagenesis kit. Next, 1 µL DpnI was directly added to the PCR reaction mixture, followed by

incubation at 37°C for 30 min and transformation. To verify the mutation sites, single clones

were selected for DNA sequencing.

**Preparation of stable K562 RAPSYN KO cell line**

K562 RAPSYN KO cells were generated using the CRISPR/Cas9 system (Genloci

Biotechnologies Inc.). RAPSYN single guide RNAs (sgRNAs) were designed using the online

CRISPR design tool (http://crispr.mit.edu/). The sgRNA sequences were

ATGGGGCGCTTCCGCGTGCTGGG,          GTAGCGGCCCATCTCCGAGTGGG,          and

TCTGGTTGGACTGGTACAGCTGG, which were cloned into the pGK1.1/CRISPR/Cas9 vector

(Genloci Biotechnologies Inc.). To obtain single clones of RAPSYN KO cells, K562 cells were

transfected with the pGK1.1/CRISPR/Cas9 plasmid containing the aforementioned sgRNA

sequence, expanded, selected with puromycin (0120A21, LEAGENE), and isolated by single-

cell culture. Single clones obtained from RAPSYN KO cells were validated by DNA sequencing

and immunoblotting.

**Preparation of stable RAPSYN KD and SRC KD cell lines**

Lentivirus-producing shRNA targeting either human RAPSYN or SRC mRNA was used to

inhibit endogenous RAPSYN or SRC expression, respectively. All shRNAs (Supplemental

Table S1) were designed using online shRNA design tools

(https://rnaidesigner.thermofisher.com and https://portals.broadinstitute.org). The shRNA

primers were ordered from GenScript and annealed in a thermal cycler according to the

following procedure (95°C, 2 min; 85°C, 9 min; 75°C, 9 min; 65°C, 9 min; 55°C, 9 min; 45°C, 9

min; 35°C, 9 min; 25°C, 10 min; 4°C, hold) with the presence of NEBuffer 2.1 (B7202S, NEW

ENGLAND BioLabs) to form a double strand with EcoRI and AgeI sticky ends. Using T4 DNA

ligase (M0202L, NEW ENGLAND BioLabs), the double-stranded shRNAs were ligated into

either the lentiviral backbone plasmid vector pLKO-EGFP-puro or a tet pLKO-EGFP-puro,

which was digested with the restriction enzymes EcoRI-HF (R3101S, NEW ENGLAND BioLabs)

and AgeI-HF (R3552S, NEW ENGLAND BioLabs). Plasmids containing shRNA or the

corresponding empty vector were cotransfected with lentivirus packaging plasmids (pLP1,

pLP2, and pLP/VSVG) into HEK293T cells using the linear polyethylenimine (23966,

Polyscience) transfection method. After transfection for 6–8 h, the transfection reagent was

replaced with a fresh medium. After incubation at 37°C, 5% $CO_2$ for 48 and 72 h, the resulting

lentivirus supernatant was collected, respectively, and filtrated through a 0.22 μm disc filter.

Then, 15 mL of the filtered lentivirus supernatant was concentrated through a 100 kD

ultrafiltration tube at 1,500 × g and 4°C for 1 h. Ph+ leukemia cell lines were infected with

concentrated lentivirus supernatant containing 8 μg/mL polybrene (H9268, Sigma). The culture

plate or dish was centrifuged in a horizontal rotor centrifuge at 2,000 × g and 32°C for 1.5 h.

After 48 h, the viral particles were replaced with fresh medium and 3 μg/mL puromycin was

added for selection for another 48–72 h. Protein expression levels were analyzed by

immunoblotting with anti-RAPSYN (ab118491, Abcam) or SRC (11097-1-AP, Proteintech)

antibodies.

**Preparation of stable RAPSYN WT，RAPSYNY 336F, or SRC expression cell lines**

Lentiviruses overexpressing RAPSYN WT, RAPSYN Y336F, or SRC were purchased from

GenePharma. The volume of virus required was calculated using the following equation: Ph+

leukemia cell lines were infected using the spin-infection method described above. 48 h after

infection, the viral particles were replaced with fresh medium. Stable RAPSYN WT, RAPSYNY

336F, or SRC expressing cells were selected in the presence of 3 μg/mL puromycin for 48-72

h. Protein expression levels were analyzed by immunoblotting with anti-RAPSYN or SRC

antibodies.

583
$$\text{virus volume} = \frac{\text{MOI} \times \text{cell number}}{\text{virus titer}}$$

**Protein expression and purification**

Recombinant pGEX-4T-1-GST-RAPSYN plasmid were transformed into the ArcticExpress

(DE3) pRARE2 competent cells (AYBIO-G6023, ANGYUBIO) and treated with 0.4 mM

isopropyl-β-D-thiogalactoside (367-93-1, Sangon Biotech) to induce fusion protein expression

under 8°C culture condition. After 50 h, cells were harvested, resuspended in PBS (0.14 M

NaCl, 2.68 mM KCl, 2mM KH2PO4, 0.01M Na2HPO4, 10 mM DTT, pH 7.4), and sonicated on

ice. Precipitates were removed from the cell lysates by centrifugation. Recombinant GST-

RAPSYN was purified from the supernatant by GST-affinity chromatography (SA010010,

Smart-Lifescience) and size-exclusion chromatography (17-0060-01, GE Healthcare). Purified

GST-RAPSYN protein was digested with thrombin (T8021, Solarbio) for 6 h at 4°C to remove

the GST tag. Recombinant pcDNA3.1-His-BCR-ABL plasmids were transfected into HEK293T

cells, which were collected after 48 h and lysed on ice using Nonidet P-40 (NP-40) lysis buffer

with a protease inhibitor cocktail. Cell lysates were centrifuged, then the supernatant fraction

incubated with the anti-BCR-ABL antibody overnight at 4°C and subsequently with protein A

magnetic beads (73778, Cell Signaling Technology) for another 4 h at 4°C. The bead

complexes were washed three times with washing buffer (Tris 25 mM, NaCl 0.15 M, Tween-20

0.005%, pH 7.5), eluted with elution buffer (glycine 0.1 M, pH 2.0), and mixed with neutralization

buffer (Tris 1 M, pH 9.0) for neutralization.

**Protein stability assay**

RAPSYN$^{WT}$, RAPSYN$^{Y336F}$, or BCR-ABL-transfected Ph$^+$ leukemia cells were incubated with 100 mg/mL cycloheximide (CHX, A8244; Cell Signaling Technology) for the indicated time points. Cells were harvested and lysed on ice using NP-40 lysis buffer supplemented with a protease inhibitor cocktail. The supernatant was collected and subjected to immunoblotting using anti-RAPSYN or SRC antibodies.

**Quantitative reverse transcription-PCR (RT-PCR)**

High-quality RNA was isolated from cells or tissues using Trizol reagent (AJF1807A, Takara) according to the manufacturer's instructions. cDNA was synthesized from 1 µg of total RNA using the HiScriptIIRT SuperMix for qPCR (R233-01, Vazyme). The ChamQ SYBR qPCR Master Mix (Q331-02, Vazyme) was used for two-step RT-PCR analysis on an Applied Biosystems StepOnePlus Real-Time PCR instrument. The samples were analyzed in triplicate. The expression value of the target gene in a given sample was normalized to the corresponding expression of ACTB or GAPDH. The 2-ΔΔCt method was used to calculate the relative expression of target genes. The primers used are listed in Supplemental Table S1.

**Cytotoxicity assay**

Lentiviruses co-expressing GFP were used to assess the toxicity of shRNAs. Flow cytometry was performed two days after shRNA transduction to determine the initial GFP-positive proportion of live cells for each shRNA. Subsequently, the cells were sampled every two days over time. The GFP-positive proportion at each time point was normalized to that of the day two. Each shRNA experiment was performed at least three times for each cell line.

**Statistical analysis**

All in vitro experiments were repeated at least three times. Animals were randomly assigned to different groups for each in vivo study. Kaplan-Meier survival analysis was used for all survival studies, and the log-rank test was used to determine significant differences between groups. Differences with * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$ were considered significant. Prism 8 (GraphPad Software, CA, USA) was used for statistical analysis. Representative results from at least three independent replicates are shown. Data are

630 presented as the means ± SD, and significant differences were determined using the Student's

631 t test, unpaired Student's t test, or one-way analysis of variance test.

## Acknowledgement

This work was supported by grants from the National Key R&D Program of China (2018YFA0902000), the National Science Foundation of China (No. 81872924, 81973386 and 82002971), the "Double First-Class" University Project (CPU2022QZ014), the Project Program of the State Key Laboratory of Natural Medicines, China Pharmaceutical University (SKLNMZZ202201) and PAPD of Jiangsu Province.

## Additional information

### Funding

| Funder | Grant reference number | Author |
|---|---|---|
| National Natural Science Foundation of China | 81973386 | Yijun Chen |
| National Natural Science Foundation of China | 81872924 | Shuzhen Wang |
| National Natural Science Foundation of China | 82002971 | Beiying Dai |
| National Key R&D Program of China | 2018YFA0902000 | Yijun Chen |
| "Double First-Class" University Project | CPU2022QZ014 | Yijun Chen |
| Project Program of the State Key Laboratory of Natural Medicines | SKLNMZZ202201 | Yijun Chen |
| Key Research and Development Project of Guangdong Province | 2022B1111070004 | Yijun Chen |

The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

**Author contributions**

Mengya Zhao, Formal analysis, Investigation, Data curation, Methodology, Writing - original draft; Beiying Dai, Formal analysis, Investigation, Data curation, Methodology, Writing - original draft, Project administration, Funding acquisition; Xiaodong Li, Formal analysis, Investigation, Data curation, Methodology; Yixin Zhang, Formal analysis, Investigation, Data curation, Methodology; Chun Qiao, Data curation; Yaru Qin, Data curation; Qingmei Li, Data curation; Zhao Li, Data curation; Shuzhen Wang, Supervision, Project administration, Funding acquisition, Writing - review & editing; Yong Yang, Supervision, Resources; Yijun Chen, Conceptualization, Supervision, Resources, Writing - review & editing, Funding acquisition.

**Competing interests**

A Chinese patent application was filed with the number of 202210107464.7. China Pharmaceutical University owns the patent rights, and Y. C., Y. Y., M. Z., B. D., X. L., S. W. and X. Z. are the inventors of the patent.

**Data availability**

The data generated in this study are available upon request from the corresponding author.

**Ethics**

This study was approved by the ethics committee of the First Affiliated Hospital of Nanjing Medical University (2019-SR-485.A1). All animal experiments were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals with the approval of the Center for New Drug Evaluation and Research, China Pharmaceutical University (approval number: B20190925-1; Nanjing, China).

## Figures and figures legends

Figure 1

**Figure 1. High protein levels of RAPSYN promotes Ph⁺ leukemia progression.**

(A) Immunoblot of RAPSYN and BCR-ABL in the PBMCs of clinical samples.

(B) Immunoblots of RAPSYN and BCR-ABL in Ph⁺ leukemic cells and normal bone

marrow stromal cells (HS-5).

(C) Cytotoxicity induced by shRNA-mediated RAPSYN knockdown in leukemic cells.

(D) Rescue of leukemic cells from sh*RAPSN* #3-induced toxicity by exogenous

expression of RAPSN cDNA.

(E) Analysis of SNARF-1 labeling intensity in K562 cells transduced with shNC or

shRAPSN #3.

(F) Quantification of cell cycle analyses in K562 cells transduced with shNC or shRAPSN #3.

(G) Quantification of cell apoptosis analyses in K562 cells transduced with shNC or shRAPSN #3.

(H) An in vivo experimental design for testing the effects of RAPSYN on tumor growth and survival.

(I) The growth curve of subcutaneous xenograft tumors was measured every two days from the third day after tumor inoculation for 19 days (five mice in each group).

(J) Photograph and weight quantification of excised tumor xenografts from (I).

(K) Immunoblot of RAPSYN and BCR-ABL in mouse xenograft tumor biopsies from K562 cells transduced with shRAPSN #3 or shNC.

(L) Immunoblot of RAPSYN and BCR-ABL in K562-RAPSYN$^{WT}$ and K562-RAPSYN$^{KO}$ cells.

(M) Kaplan-Meier survival curve of NCG mice following intravenous injection of K562-RAPSYN$^{WT}$ or K562-RAPSYN$^{KO}$ cells, as shown in (H) (ten mice in each group).

All data represent the mean ± SD of at least three independent experiments. *P* values were calculated using unpaired Student's t-test or log-rank test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 2

**Figure 2. RAPSYN neddylates BCR-ABL.**

(A) Co-immunoprecipitation of BCR-ABL and RAPSYN in leukemic cells.

(B) Immunoblot of GST and His after immunoprecipitation of His or GST in HEK293T

cells transfected with His-tagged BCR-ABL and GST-tagged RAPSYN.

(C) Immunoblot of GST and His following GST pull-down after in vitro incubation of

purified His-tagged BCR-ABL and GST or GST-tagged RAPSYN.

(D) His-immunoblot of GST immunoprecipitates from HEK293T cells transfected with

GST-tagged RAPSYN alone or in combination with His-tagged full-length or truncated

BCR-ABL (Δ1: aa 1-927, Δ2: aa 928-2047).

(E) Analysis of BCR-ABL neddylation levels in leukemic cells.

(F) Analysis of BCR-ABL neddylation levels in leukemic cells treated with MLN4924 or

DMSO for 24 h.

(G) HA-immunoblot of His-immunoprecipitate from HEK293T cells transfected with

His-tagged BCR-ABL and HA-tagged NEDD8 or NEDD8 ΔGG.

(H) HA-immunoblot of the His-immunoprecipitate from HEK293T cells transfected with

the indicated constructs.

(I) Analysis of BCR-ABL neddylation levels in K562 WT, RAPSYN KO and RAPSYN

KO with exogenous expression of a RAPSN cDNA cells.

(J) HA-immunoblot after immunoprecipitation of His in HEK293T cells transfected with

His-tagged BCR-ABL, HA-tagged NEDD8, GFP-tagged WT RAPSYN or C366A

RAPSYN.

(K) Assessment of BCR-ABL neddylation by RAPSYN in vitro. Purified RAPSYN and

BCR-ABL were incubated with APPBP1/UBA3, UBE2M or NEDD8 in in vitro

neddylation assay.

(L) Analysis of BCR-ABL neddylation levels in excised tumor xenografts from Fig. 1J.

(M) Verification of BCR-ABL neddylation sites in HEK293T cells transfected with

indicated constructs.

Figure 3

**Figure 3. RAPSYN attenuates BCR-ABL ubiquitination and degradation.**

(A) Immunoblot of BCR-ABL in leukemic cells treated with MLN4924 or DMSO for 24 h and corresponding quantification of three independent replicates.

(B) Immunoblot of BCR-ABL in K562 WT and RAPSYN KO cells and corresponding quantification of three independent replicates.

(C) Assessment of BCR-ABL protein stability in K562 cells expressing DOX-inducible sh*RAPSN* #3 treated with CHX alone or in combination with DOX at indicated time points by immunoblotting.

(D) Analysis of BCR-ABL neddylation and ubiquitination levels in leukemic cells treated with MLN4924 or DMSO for 24 h.

(E) Analysis of BCR-ABL neddylation and ubiquitination levels in K562 WT and RAPSYN KO cells.

(F) Immunoblot of HA and Myc after His-immunoprecipitation in HEK293T cells transfected with His-tagged BCR-ABL, HA-tagged Ub or without Myc-tagged NEDD8.

(G) Analysis of BCR-ABL ubiquitination and neddylation in leukemic cells treated with MG132 or DMSO for 12 h.

(H) Co-immunoprecipitation of BCR-ABL, c-CBL and RAPSYN in leukemic cells expressing exogenous RAPSN cDNA or empty vector.

(I) Co-immunoprecipitation of BCR-ABL, c-CBL and RAPSYN in K562 WT and RAPSYN KO cells.

All data represent the mean ± SD of at least three independent experiments. P values were calculated using unpaired Student's t-test. ** $p < 0.01$, *** $p < 0.001$.

Figure 4

**Figure 4. SRC-mediated phosphorylation at Y336 promotes RAPSYN stability by repressing its proteasomal degradation.**

(A) Assessment of RAPSYN phosphorylation levels in leukemic cells treated with saracatinib or DMSO for 24 h.

(B) Assessment of RAPSYN phosphorylation levels in leukemic cells transduced with shSRC or shNC.

(C) Assessment of RAPSYN phosphorylation levels in leukemic cells expressing exogenous SRC cDNA or empty vector.

(D) Assessment of RAPSYN phosphorylation by SRC in vitro. Purified RAPSYN and SRC were incubated with ATP in the presence or absence of saracatinib for phosphorylation assay.

(E) Verification of RAPSYN phosphorylation sites. Purified SRC and RAPSYN WT or indicated mutants were incubated with ATP for phosphorylation assay.

(F) Assessment of RAPSYN protein stability in leukemic cells treated with CHX in combination with saracatinib or DMSO at indicated time points by immunoblotting.

(G) Assessment of RAPSYN protein stability in leukemic cells transduced with shSRC or shNC by immunoblotting.

(H) Assessment of RAPSYN protein stability in leukemic cells transduced with exogenous SRC cDNA or empty vector by immunoblotting.

(I) Assessment of RAPSYN protein stability in leukemic cells transduced with exogenous RAPSYN WT or Y336F cDNA by immunoblotting.

(J) Immunoblot of RAPSYN in leukemic cells treated with saracatinib or DMSO for 12 h, and subsequently with MG132 or DMSO for another 12 h.

(K) Immunoblot of RAPSYN in leukemic cells transduced with shNC or shSRC and treated with MG132 or DMSO for 12 h.

**Figure 5. RAPSYN phosphorylation at Y336 potentiates its E3 ligase activity and promotes BCR-ABL stabilization.**

(A) Immunoblot of BCR-ABL neddylation levels in leukemic cells treated with saracatinib or DMSO for 24 h.

(B) Immunoblot of BCR-ABL neddylation levels in leukemic cells transduced with sh*SRC* or shNC.

(C) Immunoblot of BCR-ABL neddylation levels in leukemic cells expressing exogenous SRC cDNA or empty vector.

(D) Effects of RAPSYN phosphorylation on BCR-ABL neddylation levels in HEK293T cells transfected with indicated constructs.

(E) Effects of RAPSYN phosphorylation at Y336 on BCR-ABL neddylation levels in leukemic cells expressing exogenous RAPSYN WT, Y336F cDNA, or empty vector.

(F) Assessment of BCR-ABL protein stability in leukemic cells transduced with exogenous RAPSYN WT, Y336F cDNA, or empty vector by immunoblotting.

Figure 6

**Figure 6. SRC-mediated phosphorylation of RAPSYN at Y336 promotes Ph⁺**

**leukemia progression.**

(A) Cytotoxicity induced by sh*SRC* #2-mediated SRC knockdown in leukemic cells.

(B) Rescue of leukemic cells from sh*SRC* #2-induced toxicity by exogenous expression

of SRC cDNA.

(C) Rescue of leukemic cells from sh*SRC* #2-induced toxicity by exogenous expression of RAPSN$^{WT}$ cDNA.

(D) Failed rescue of leukemic cells from sh*SRC* #2-induced toxicity by exogenous expression of RAPSN$^{Y336F}$ cDNA.

(E) Viability of leukemic cells transduced with either RAPSN$^{WT}$ cDNA or corresponding empty vector after 72 h of incubation with indicated concentrations of saracatinib.

(F) Viability of leukemic cells transduced with either RAPSN$^{Y336F}$ cDNA or corresponding empty vector after 72 h of incubation with indicated concentrations of saracatinib.

(G) Viability of leukemic cells transduced with either shNC or sh*RAPSN* #3 after 72 h of incubation with indicated concentrations of saracatinib.

(H) Experimental design used to test in vivo effects of RAPSYN phosphorylation at Y336 on Ph$^+$ leukemia progression and survival time.

(I) Kaplan-Meier survival curve of NCG mice following intravenous injection of K562-RAPSYN$^{WT}$ or K562-RAPSYN$^{Y336F}$ cells and intragastric administration of saracatinib or corresponding vehicle from days 6 to 26 as indicated (ten mice in each group).

(J) Kaplan-Meier survival curve of NCG mice following intravenous injection of double-transfected K562 cells (ten mice in each group). Representative results from at least three independent experiments are shown (A-G); error bars, mean ± SD; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; log-rank test (I-L).

**Figure.7. Schematic representation of the competition between neddylation and**

**ubiquitination on BCR-ABL in Ph⁺ leukemia.**

**Reference**

Aydin, B., Kan, B., and Cabadak, H. 2013. The role of intracellular pathways in the proliferation
of human K562 cells mediated by muscarinic receptors. *Leuk Res* **37**: 1144-1149. DOI:
10.1016/j.leukres.2013.05.018.

Bahjat, M., de Wilde, G., van Dam, T., Maas, C., Bloedjes, T., Bende, R.J., van Noesel, C.J.M.,
Luijks, D.M., Eldering, E., Kersten, M.J., and Guikema, J.E.J. 2019. The NEDD8-activating
enzyme inhibitor MLN4924 induces DNA damage in Ph+ leukemia and sensitizes for ABL
kinase inhibitors. *Cell Cycle* **18**: 2307-2322. DOI:10.1080/15384101.2019.1646068.

Barnes, D.J., Palaiologou, D., Panousopoulou, E., Schultheis, B., Yong, A.S., Wong, A.,
Pattacini, L., Goldman, J.M., and Melo, J.V. 2005. Bcr-Abl expression levels determine the rate
of development of resistance to imatinib mesylate in chronic myeloid leukemia. *Cancer Res* **65**:
8912-8919. DOI:10.1158/0008-5472.CAN-05-0076.

Bekes, M., Langley, D.R., and Crews, C.M. 2022. PROTAC targeted protein degraders: the
past is prologue. *Nat Rev Drug Discov* **21**: 181-200. DOI:10.1038/s41573-021-00371-6.

Braun, T.P., Eide, C.A., and Druker, B.J. 2020. Response and Resistance to BCR-ABL1-
Targeted Therapies. *Cancer Cell* **37**: 530-542. DOI:10.1016/j.ccell.2020.03.006.

Burslem, G.M., Schultz, A.R., Bondeson, D.P., Eide, C.A., Savage Stevens, S.L., Druker, B.J.,
and Crews, C.M. 2019. Targeting BCR-ABL1 in chronic myeloid leukemia by PROTAC-
mmediated targeted protein degradation. *Cancer Res* **79**: 4744-4753. DOI:10.1158/0008-
5472.CAN-19-1236.

Cabadak, H., Aydin, B., and Kan, B. 2011. Regulation of M2, M3, and M4 muscarinic receptor
expression in K562 chronic myelogenous leukemic cells by carbachol. *J Recept Signal*
*Transduct Res* **31**: 26-32. DOI:10.3109/10799893.2010.506484.

Cortes, J., Pavlovsky, C., and Saußele, S. 2021. Chronic myeloid leukaemia. *The Lancet* **398**:
1914-1926. DOI:10.1016/s0140-6736(21)01204-6.

Deininger, M.W., Goldman, J.M., and Melo, J.V. 2000. The molecular biology of chronic myeloid
leukemia. *Blood* **96**: 3343-3356.

Demizu, Y., Shibata, N., Hattori, T., Ohoka, N., Motoi, H., Misawa, T., Shoda, T., Naito, M., and
Kurihara, M. 2016. Development of BCR-ABL degradation inducers via the conjugation of an
imatinib derivative and a cIAP1 ligand. *Bioorg Med Chem Lett* **26**: 4865-4869.
DOI:10.1016/j.bmcl.2016.09.041.

Durieux, M.E., and Nietgen, G.W. 1997. Synergistic inhibition of muscarinic signaling by
ketamine stereoisomers and the preservative benzethonium chloride. *Anesthesiology* **86**: 1326-
1333. DOI:10.1097/00000542-199706000-00014.

Embade, N., Fernandez-Ramos, D., Varela-Rey, M., Beraza, N., Sini, M., Gutierrez de Juan, V.,
Woodhoo, A., Martinez-Lopez, N., Rodriguez-Iruretagoyena, B., Bustamante, F.J., de la Hoz,
A.B., Carracedo, A., Xirodimas, D.P., Rodriguez, M.S., Lu, S.C., Mato, J.M., and Martinez-
Chantar, M.L. 2012. Murine double minute 2 regulates Hu antigen R stability in human liver and
colon cancer through NEDDylation. *Hepatology* **55**: 1237-1248. DOI:10.1002/hep.24795.

Enchev, R.I., Schulman, B.A., and Peter, M. 2015. Protein neddylation: beyond cullin-RING
ligases. *Nat Rev Mol Cell Biol* **16**: 30-44. DOI:10.1038/nrm3919.

Guo, N., Azadniv, M., Coppage, M., Nemer, M., Mendler, J., Becker, M., and Liesveld, J. 2019.
Effects of neddylation and mTOR Inhibition in acute myelogenous leukemia. *Transl Oncol* **12**:

602-613. DOI:10.1016/j.tranon.2019.01.001.

Hochhaus, A., Baccarani, M., Silver, R.T., Schiffer, C., Apperley, J.F., Cervantes, F., Clark, R.E.,
Cortes, J.E., Deininger, M.W., Guilhot, F., Hjorth-Hansen, H., Hughes, T.P., Janssen, J.,
Kantarjian, H.M., Kim, D.W., Larson, R.A., Lipton, J.H., Mahon, F.X., Mayer, J., Nicolini, F.,
Niederwieser, D., Pane, F., Radich, J.P., Rea, D., Richter, J., Rosti, G., Rousselot, P., Saglio,
G., Saussele, S., Soverini, S., Steegmann, J.L., Turkina, A., Zaritskey, A., and Hehlmann, R.
2020. European LeukemiaNet 2020 recommendations for treating chronic myeloid leukemia.
*Leukemia* *34*: 966-984. DOI:10.1038/s41375-020-0776-2.

Huh, K.H., and Fuhrer, C. 2002. Clustering of nicotinic acetylcholine receptors: from the
neuromuscular junction to interneuronal synapses. *Mol Neurobiol* *25*: 79-112.
DOI:10.1385/MN:25:1:079.

Issaad, C., Ahmed, M., Novault, S., Bonnet, M.L., Bennardo, T., Varet, B., Vainchenker, W., and
Turhan, A.G. 2000. Biological effects induced by variable levels of BCR-ABL protein in the
pluripotent hematopoietic cell line UT-7. *Leukemia* *14*: 662-670. DOI:10.1038/sj.leu.2401730.

Jabbour, E., Cortes, J., O'Brien, S., Giles, F., and Kantarjian, H. 2007. New targeted therapies
for chronic myelogenous leukemia: opportunities to overcome imatinib resistance. *Semin*
*Hematol* *44*: S25-31. DOI:10.1053/j.seminhematol.2006.12.003.

Jabbour, E., and Kantarjian, H. 2020. Chronic myeloid leukemia: 2020 update on diagnosis,
therapy and monitoring. *Am J Hematol* *95*: 691-709. DOI:10.1002/ajh.25792.

Jakubik, J., Michal, P., Machova, E., and Dolezal, V. 2008. Importance and prospects for design
of selective muscarinic agonists. *Physiol Res* *57 Suppl 3*: S39-S47.
DOI:10.33549/physiolres.931449.

Jiang, L., Wang, Y., Li, Q., Tu, Z., Zhu, S., Tu, S., Zhang, Z., Ding, K., and Lu, X. 2021a. Design,
synthesis, and biological evaluation of Bcr-Abl PROTACs to overcome T315I mutation. *Acta*
*Pharm Sin B* *11*: 1315-1328. DOI:10.1016/j.apsb.2020.11.009.

Jiang, S., Wang, X., He, Y., Huang, H., Cao, B., Zhang, Z., Liu, J., Wang, Q., Huang, Z., and
Mao, X. 2021b. Suppression of USP7 induces BCR-ABL degradation and chronic myelogenous
leukemia cell apoptosis. *Cell Death Dis* *12*: 456. DOI:10.1038/s41419-021-03732-6.

Bhalla KN, Fiskus W. 2016. NEDD8 and HDACs: promising cotargets in AML. *Blood* *127*:
2167-2170. DOI:10.1182/blood-2016-02-699058.

Kawamata, J., Suzuki, S., and Shimohama, S. 2011. Enhancement of nicotinic receptors
alleviates cytotoxicity in neurological disease models. *Ther Adv Chronic Dis* *2*: 197-208.
DOI:10.1177/2040622310397691.

Klein, A.d., Kessel, A.G.v., Grosveld, G., Bartram, C.R., Hagemeijer, A., Bootsma, D., Spurr,
N.K., Heisterkamp, N., and J Groffen, J.R.S. 1982. A cellular oncogene is translocated to the
Philadelphia chromosome in chronic myelocytic leukaemia. *Nature* *300*: 765-767.
DOI:10.1038/300765a0.

Lai, A.C., Toure, M., Hellerschmied, D., Salami, J., Jaime-Figueroa, S., Ko, E., Hines, J., and
Crews, C.M. 2016. Modular PROTAC design for the degradation of oncogenic BCR-ABL.
*Angew Chem Int Ed Engl* *55*: 807-810. DOI:10.1002/anie.201507634.

Lee, G.W., Park, J.B., Park, S.Y., Seo, J., Shin, S.H., Park, J.W., Kim, S.J., Watanabe, M., and
Chun, Y.S. 2018. The E3 ligase C-CBL inhibits cancer cell migration by neddylating the proto-
oncogene c-Src. *Oncogene* *37*: 5552-5568. DOI:10.1038/s41388-018-0354-5.

Legay, C., and Mei, L. 2017. Moving forward with the neuromuscular junction. *J Neurochem*

**142 Suppl 2**: 59-63. DOI:10.1111/jnc.14028.

Li, J., Zou, J., Littlejohn, R., Liu, J., and Su, H. 2020. Neddylation, an emerging mechanism
regulating cardiac development and function. *Front Physiol* **11**: 612927.
DOI:10.3389/fphys.2020.612927.

Li, L., Cao, Y., Wu, H., Ye, X., Zhu, Z., Xing, G., Shen, C., Barik, A., Zhang, B., Xie, X., Zhi, W.,
Gan, L., Su, H., Xiong, W.C., and Mei, L. 2016. Enzymatic activity of the scaffold protein rapsyn
for synapse formation. *Neuron* **92**: DOI:1007-1019. 10.1016/j.neuron.2016.10.023.

Li, L., Xiong, W.C., and Mei, L. 2018. Neuromuscular junction formation, aging, and disorders.
*Annu Rev Physiol* **80**: 159-188. 10.1146/annurev-physiol-022516-034255.

Li, X., and Song, Y. 2020. Proteolysis-targeting chimera (PROTAC) for targeted protein
degradation and cancer therapy. *J Hematol Oncol* **13**: 50. DOI:10.1186/s13045-020-00885-3.

Liu, C., Nie, D., Li, J., Du, X., Lu, Y., Li, Y., Zhou, J., Jin, Y., and Pan, J. 2018. Antitumor effects
of blocking protein neddylation in T315I-BCR-ABL leukemia cells and leukemia stem cells.
*Cancer Res* **78**: 1522-1536. DOI:10.1158/0008-5472.CAN-17-1733.

Liu, H., Mi, Q., Ding, X., Lin, C., Liu, L., Ren, C., Shen, S., Shao, Y., Chen, J., Zhou, Y., Ji, L.,
Zhang, H., Bai, F., Yang, X., Yin, Q., and Jiang, B. 2022. Discovery and characterization of
novel potent BCR-ABL degraders by conjugating allosteric inhibitor. *Eur J Med Chem* **244**:
DOI:114810. 10.1016/j.ejmech.2022.114810.

Lussana, F., Intermesoli, T., Stefanoni, P., and Rambaldi, A. 2018. Mechanisms of resistance
to targeted therapies in chronic myeloid leukemia. *Handb Exp Pharmacol* **249**: 231-250.
DOI:10.1007/164_2017_81.

Mao, J.H., Sun, X.Y., Liu, J.X., Zhang, Q.Y., Liu, P., Huang, Q.H., Li, K.K., Chen, Q., Chen, Z.,
and Chen, S.J. 2010. As4S4 targets RING-type E3 ligase c-CBL to induce degradation of BCR-
ABL in chronic myelogenous leukemia. *Proc Natl Acad Sci U S A* **107**: 21683-21688.
DOI:10.1073/pnas.1016311108.

McGrail, D.J., Garnett, J., Yin, J., Dai, H., Shih, D.J.H., Lam, T.N.A., Li, Y., Sun, C., Li, Y.,
Schmandt, R., Wu, J.Y., Hu, L., Liang, Y., Peng, G., Jonasch, E., Menter, D., Yates, M.S., Kopetz,
S., Lu, K.H., Broaddus, R., Mills, G.B., Sahni, N., and Lin, S.Y. 2020. Proteome instability is a
therapeutic vulnerability in Mismatch Repair-Deficient cancer. *Cancer Cell* **37**: 371-386 e312.
DOI:10.1016/j.ccell.2020.01.011.

Mohamed, A.S., and Swope, S.L. 1999. Phosphorylation and cytoskeletal anchoring of the
acetylcholine receptor by Src class protein-tyrosine kinases. Activation by rapsyn. *J Biol Chem*
**274**: 20529-20539. DOI:10.1074/jbc.274.29.20529.

Norton, J.P., Augert, A., Eastwood, E., Basom, R., Rudin, C.M., and MacPherson, D. 2021.
Protein neddylation as a therapeutic target in pulmonary and extrapulmonary small cell
carcinomas. *Genes Dev* **35**: 870-887. DOI:10.1101/gad.348316.121.

Önder Narin G., Aydin, B., and Cabadak, H. 2021. Studies on the role of alpha 7 nicotinic
acetylcholine receptors in K562 cell proliferation and signaling. *Mol Biol Rep* **48**: 5045-5055.
DOI:10.1007/s11033-021-06498-4.

Oved, S., Mosesson, Y., Zwang, Y., Santonico, E., Shtiegman, K., Marmor, M.D.,
Kochupurakkal, B.S., Katz, M., Lavi, S., Cesareni, G., and Yarden, Y. 2006. Conjugation to
Nedd8 instigates ubiquitylation and down-regulation of activated receptor tyrosine kinases. *J*
*Biol Chem* **281**: 21640-21651. DOI:10.1074/jbc.M513034200.

Shibata, N., Ohoka, N., Tsuji, G., Demizu, Y., Miyawaza, K., Ui-Tei, K., Akiyama, T., and Naito,

M. 2020. Deubiquitylase USP25 prevents degradation of BCR-ABL protein and ensures proliferation of Ph-positive leukemia cells. *Oncogene 39*: 3867-3878. DOI:10.1038/s41388-020-1253-0.

Shimokawa, K., Shibata, N., Sameshima, T., Miyamoto, N., Ujikawa, O., Nara, H., Ohoka, N., Hattori, T., Cho, N., and Naito, M. 2017. Targeting the allosteric site of oncoprotein BCR-ABL as an alternative strategy for effective target protein degradation. *ACS Med Chem Lett 8*: 1042-1047. DOI:10.1021/acsmedchemlett.7b00247.

van der Veen, A.G., and Ploegh, H.L. 2012. Ubiquitin-like proteins. *Annu Rev Biochem 81*: 323-357. DOI:10.1146/annurev-biochem-093010-153308.

Wang, S., Xie, W., Wang, D., Peng, Z., Zheng, Y., Liu, N., Dai, W., Wang, Y., Wang, Z., Yang, Y., and Chen, Y. 2015. Discovery of a small molecule targeting SET-PP2A interaction to overcome BCR-ABL$^{T315I}$ mutation of chronic myeloid leukemia. *Oncotarget 6*: 12128-12140. DOI:10.18632/oncotarget.3665.

Witzemann, V., Chevessier, F., Pacifici, P.G., and Yampolsky, P. 2013. The neuromuscular junction: selective remodeling of synaptic regulators at the nerve/muscle interface. *Mech Dev 130*: 402-411. DOI:10.1016/j.mod.2012.09.004.

Xie, P., Peng, Z., Chen, Y., Li, H., Du, M., Tan, Y., Zhang, X., Lu, Z., Cui, C.P., Liu, C.H., He, F., and Zhang, L. 2021. Neddylation of PTEN regulates its nuclear import and promotes tumor development. *Cell Res 31*: 291-311. DOI:10.1038/s41422-020-00443-z.

Xing, G., Jing, H., Zhang, L., Cao, Y., Li, L., Zhao, K., Dong, Z., Chen, W., Wang, H., Cao, R., Xiong, W.C., and Mei, L. 2019. A mechanism in agrin signaling revealed by a prevalent Rapsyn mutation in congenital myasthenic syndrome. *eLlife 8*. DOI:10.7554/eLife.49180.

Xing, G., Xiong, W.C., and Mei, L. 2020. Rapsyn as a signaling and scaffolding molecule in neuromuscular junction formation and maintenance. *Neurosci Lett 731*: 135013. DOI:10.1016/j.neulet.2020.135013.

Yang, Y., Ding, L., Guo, Z.K., Zheng, X.L., Wang, L.S., Sun, H.Y., Jin, Z.G., and Wang, H.X. 2017. The epigenetically-regulated miR-34a targeting c-SRC suppresses RAF/MEK/ERK signaling pathway in K-562 cells. *Leuk Res 55*: 91-96. DOI:10.1016/j.leukres.2017.01.020.

Yao, J., Liang, X., Liu, Y., and Zheng, M. 2020. Neddylation: A Versatile Pathway Takes on Chronic Liver Diseases. *Front Med (Lausanne) 7*: 586881. DOI:10.3389/fmed.2020.586881.

Ying, J., Zhang, M., Qiu, X., and Lu, Y. 2018. Targeting the neddylation pathway in cells as a potential therapeutic approach for diseases. *Cancer Chemother Pharmacol 81*: 797-808. DOI:10.1007/s00280-018-3541-8.

Zhao, Q., Ren, C., Liu, L., Chen, J., Shao, Y., Sun, N., Sun, R., Kong, Y., Ding, X., Zhang, X., Xu, Y., Yang, B., Yin, Q., Yang, X., and Jiang, B. 2019. Discovery of SIAIS178 as an effective BCR-ABL degrader by recruiting Von Hippel-Lindau (VHL) E3 ubiquitin ligase. *J Med Chem 62*: 9281-9298. 10.1021/acs.jmedchem.9b01264.

Zhou, L., Chen, S., Zhang, Y., Kmieciak, M., Leng, Y., Li, L., Lin, H., Rizzo, K.A., Dumur, C.I., Ferreira-Gonzalez, A., Rahmani, M., Povirk, L., Chalasani, S., Berger, A.J., Dai, Y., and Grant, S. 2016. The NAE inhibitor pevonedistat interacts with the HDAC inhibitor belinostat to target AML cells by disrupting the DDR. *Blood 127*: 2219-2230. DOI:10.1182/blood-2015-06-653717.

Zhou, L., Jiang, Y., Luo, Q., Li, L., and Jia, L. 2019. Neddylation: a novel modulator of the tumor microenvironment. *Mol Cancer 18*: 77. DOI:10.1186/s12943-019-0979-1.

Zhou, Q., and Sun, Y. 2019. MLN4924: additional activities beyond neddylation inhibition. *Mol*

1015    *Cell Oncol 6*: e1618174. DOI:10.1080/23723556.2019.1618174.

1016    Zuo, W., Huang, F., Chiang, Y.J., Li, M., Du, J., Ding, Y., Zhang, T., Lee, H.W., Jeong, L.S.,

1017    Chen, Y., Deng, H., Feng, X.H., Luo, S., Gao, C., and Chen, Y.G. 2013. c-Cbl-mediated

1018    neddylation antagonizes ubiquitination and degradation of the TGF-beta type II receptor. *Mol*

1019    *Cell 49*: 499-510. DOI:10.1016/j.molcel.2012.12.002.

1020