Rapid intracellular acidification is a novel plant defense response countered by the brown
planthopper

Yanjuan Jiang[a,b,1*], Xiao-Ya Zhang[b,c,1], Shaoqin Li[d,e], Yu-Cheng Xie[c], Xu-Mei Luo[c], Yongping
Yang[d,e], Zhengyan Pu[a], Li Zhang[f,g], Jia-Bao Lu[h], Hai-Jian Huang[h], Chuan-Xi Zhang[c,h,*], Sheng
Yang He[b,f,g,*].

[a]State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan
University, Kunming 650091, China

[b]DOE Plant Research Laboratory, Michigan State University, East Lansing, MI, USA

[c]Institute of Insect Science, Zhejiang University, Hangzhou, 310058, China.

[d]CAS Key Laboratory of Tropical Plant Resources and Sustainable Use, Xishuangbanna Tropical
Botanical Garden, Chinese Academy of Sciences, Kunming, Yunnan 650223, China

[e]University of Chinese Academy of Sciences, Beijing 100049, China.

[f]Howard Hughes Medical Institute, Duke University, Durham, NC, USA.

[g]Department of Biology, Duke University, Durham, NC, USA.

[h]State Key Laboratory for Managing Biotic and Chemical Threats to the Quality and Safety of
Agro-products, Ningbo University, China.

[1]Yanjuan Jiang and Xiao-Ya Zhang contributed equally to this work.
*Corresponding authors
Email: yanjuanjiang@ynu.edu.cn, chxzhang@zju.edu.cn or Shengyang.he@duke.edu

Keywords: Brown planthopper | Carbonic anhydrase | Insect effector | Plant immunity | Callose
deposition|

Abstract

The brown planthopper (BPH; *Nilaparvata lugens*) is the most destructive insect pest in rice. Through a stylet, BPH secrets a plethora of salivary proteins into rice phloem cells as a crucial step of infestation. However, how various salivary proteins function in rice cells to promote insect infestation is poorly understood. Among them, one of the salivary proteins is predicted to be a carbonic anhydrase (NICA). The survival rate of the *NICA*-RNA interference (RNAi) BPH insects was extremely low on rice, indicating a vital role of this salivary protein in BPH infestation. We generated *NICA* transgenic rice plants and found that *NICA* expressed in rice plants could restore the ability of *NICA*-RNAi BPH to survive on rice. Next, we produced rice plants expressing the ratiometric pH sensor pHusion and found that *NICA*-RNAi BPH induced rapid intracellular acidification of rice cells during feeding. Further analysis revealed that both *NICA*-RNAi BPH feeding and artificial of lowering intracellular pH activated plant defense responses, and that NICA-mediated intracellular pH stabilization is linked to diminished defense responses, including reduced callose deposition at the phloem sieve plates and suppressed defense gene expression. Given the importance of pH homeostasis across the kingdoms of life, discovery of NICA-mediated intracellular pH modulation uncovered a new dimension in the interaction between plants and piecing/sucking insect pests. The crucial role of NICA for BPH infestation of rice suggests that NICA is a promising target for chemical or trans-kingdom RNAi-based inactivation for BPH control strategies in plants.

Introduction

The brown planthopper (BPH; *Nilaparvata lugens* Stål, Hemiptera, Delphacidae) is a monophagous insect pest of rice (*Oryza sativa* L.) found in all rice-growing Asian countries. The BPH sucks rice phloem sap via its stylet, causing leaf yellowing and wilting, stunted plant growth, reduced photosynthesis and ultimately death of rice plant [1]. During severe BPH outbreaks, tens of thousands of insects swarm on a rice field, resulting in the 'hopperburn' phenomenon, which is characterized by large-scale wilting, yellowing and lethal drying of rice plants [2]. Besides direct damages, the BPH may also indirectly damage rice plants by oviposition and transmitting viral disease agents [3,4].

Application of chemical insecticides has been a main strategy for controlling BPH. Although it has the advantages of having rapid effects on killing insects and low costs, use of insecticides leads to environmental pollution and resistance of BPH to pesticides. In recent decades, breeding rice resistant varieties to control BPH has attracted increasing attention. To date, more than 30 resistance genes have been found in the rice genome [5]. However, BPH often quickly evolves new

biological types that evade rice resistant genes. Therefore, additional BPH controlling methods
need to be developed to complement the current control measures toward long-term solutions of
achieving durable BPH resistance in rice. One method could be based on disruption of key steps
of BPH's natural infestation process. However, development of such methods will require a
comprehensive understanding of the basic biology of the BPH-rice interaction.

A critical step in BPH infestation of rice is secreting bioactive substances into the plant tissues
through the stylet [6-8]. Specifically, during feeding, BPH secretes both colloidal and watery saliva
[9,10]. The main function of the colloidal saliva is to form a saliva sheath around the piercing-sucking
mouthparts, stabilizing the overall feeding apparatus. The composition and function of watery
saliva is more complex, and it contains salivary proteins that are believed, in most cases, to
regulate various pathways in plant cells to enhance BPH feeding and survival in rice plants. For
example, NlEG1, a salivary endo-b-1,4-glucanase, degrades plant celluloses to help the BPH's
stylet reach to the phloem [11]. NlSEF1, an EF-hand Ca^{2+}-binding protein, interferes with calcium
signaling and H_2O_2 production during BPH feeding [12]. Salivary protein 7 is required for normal
feeding behavior and for countering accumulation of a defense compound, tricin [13]. On the other
hand, mucin-like protein (NlMLP) triggers defense responses in rice cells, including cell death,
callose deposition and up-regulation of pathogen-responsive genes [14,15].

Carbonic anhydrases (CAs) (EC 4.2.1.1) are zinc metalloenzymes that function as catalysts in the
bidirectional conversion of CO_2 and water into bicarbonate and protons [16]. There are at least five
distinct CA families (α-, β-, γ-, δ-, and ε-CAs) and three of them (α-, β-, and γ-CAs) are
ubiquitously distributed among animal, plant, and bacterial species. The widespread distribution
and adequate abundance of these CA families underline their evolutionary importance throughout
the kingdoms of life. CAs participate in a wide range of biological processes, such as pH
regulation, CO_2 homeostasis, stomatal aperture, and plant defense [17-21]. NlCA belongs to the α-
CA subfamily. Our previous study showed that *NlCA* expressed in BPH salivary glands [6].
Surprisingly, however, RNA interference (RNAi) of the *NlCA* transcript in BPH insects affects
neither pH maintenance within the salivary gland, watery saliva or gut, nor insect feeding
behavior or honeydew excretion, but greatly reduced survival of BPH on rice plants, suggesting a
critical function *in planta* via an unknown mechanism [6].

Here, we report that *NlCA*-RNAi BPH feeding results in rapid intracellular acidification of rice cells.
We found that NlCA is secreted into the rice tissues and functions as an effector that stabilizes
host cell intracellular pH, accompanied by suppression of defense responses, during BPH feeding.
Thus, we have uncovered intracellular pH homeostasis is a previously uncharacterized
battleground in plant-insect interactions.

Results

NICA is detected in rice sheath tissues during BPH feeding. NICA was previously found to be

highly expressed in salivary glands and present in the watery saliva of BPH fed on artificial diet [6].

We conducted a more detailed characterization of *NICA* expression in this study. RNA *in situ*

hybridization showed that the expression level of *NICA* was detectable throughout the principal

glands (PGs) and accessory glands (AGs), but not in A-follicle of the principal gland (APG) (Fig.

1a), which further raised the possibility that NICA may be one of the "effector proteins" secreted

into the rice tissue during BPH feeding on rice plants. To test this possibility, we compared protein

profiles in the leaf sheaths of Nipponbare rice plants before and after BPH feeding using liquid

chromatograph-mass spectrometer (LC-MS). We found 8 NICA-specific peptides in BPH-fed leaf

sheath tissue (Fig. 1b*;* Supplementary Fig. 1), confirming that NICA is secreted into the host

tissues during BPH feeding.

Transgenic expression of *NICA* in rice rescues the ability of *NICA*-silenced BPH to feed

and survive. To further clarify the site of function (i.e., in insect vs. in plant) of NICA in the BPH-

rice interaction, we produced transgenic Nipponbare plants expressing *NICA* (see Methods). A

total of 26 lines were produced and 6 lines were found to robustly express the *NICA* transcript.

NICA-expressing plants exhibited no noticeable changes in appearance compared to Nipponbare

plants (Supplementary Fig. 2). *NICA*-expressing plant lines were propagated to T3 generation,

and 3 lines were subjected to further characterization, including BPH feeding. For BPH feeding

assay, double-stranded RNA (dsRNA) of *NICA* (ds*NICA*) or the control green fluorescent protein

gene (ds*GFP*) was injected into 3rd instar BPH nymphs to initiate RNAi of the *NICA* transcript [22].

The sequence of *NICA* used for RNAi treatment is expected to be specific to BPH

(Supplementary Fig. 3). Quantitative real-time PCR analysis confirmed that the transcript levels of

the *NICA* gene were reduced by 99% and 97%, respectively, in tested individuals when compared

with non-RNAi control and ds*GFP*-treated insects (Figure 2a). There were no significant

differences in survival rates between ds*GFP* and ds*NICA* BPH when fed on the artificial diet,

indicating that silencing *NICA* expression has no obvious impact on the basic physiology of BPH

(Fig. 2b). However, we found that the survival rate of ds*NICA* BPH was sharply decreased,

starting at day 9 post-infestation, to ~40% at day 14, whereas the control ds*GFP* BPH survived

normally on wild-type Nipponbare plants (Fig. 2c). This result is consistent with previous results

conducted in the *japonica* cultivar Xiushui134 rice plants [6], suggesting that the requirement of

NICA for BPH survival is not specific to a specific rice genotype. Strikingly, *NICA* transgenic

plants almost fully restored the survival of *NICA*-silenced BPH insects (Fig. 2c; Supplementary

Fig. 4), demonstrating that *NI*CA expressed in the host tissue can complement the infestation

defect of *NICA*-silenced BPH insects.

The conserved catalytic site amino acids of *Nl*CA are critical to its function in rice. NlCA
contains several conserved amino acid residues predicted to be at the catalytic site of carbonic
anhydrases (Fig. 2d). We asked if some of these conserved active site residues are critical to the
function of *Nl*CA in the rice-BPH interaction. Accordingly, we expressed three different *Nl*CA
mutants in Nipponbare plants. "*NlCA-m1*" plants contain a *NlCA* transgene in which His^{159} and
His^{161} were mutated to Ala; "*NlCA-m2*" plants contain a *NlCA* transgene in which Glu^{182} and
His^{184} were replaced by Ala; and "*NlCA-m3*" plants carry a *NlCA* transgene in which all seven
conserved active site residues, His^{159}, His^{161}, Glu^{171}, Glu^{182}, His^{184}, Thr^{269} and Phe^{279}, were
changed to Ala. Like *NlCA* transgenic plants, transgenic plants expressing these *NlCA*-mutants
exhibited no noticeable changes in appearance (Supplementary Fig. 2). We conducted BPH
survival assay and found that, in contrast to plants expressing wild-type *NlCA*, the *NlCA-m2* and
NlCA-m3 plants could not fully rescue the infestation defect of *NlCA*-silenced BPH insects. The
NlCA-m1 plants, on the other hand, could recover the infestation defect of ds*NlCA* BPH (Fig. 2e
and f; Supplementary Fig. 5). Thus, several conservative residues at the predicted catalytic site
are indispensable for the function of *Nl*CA inside the plant cell.

After having demonstrated the requirement of catalytic site residues for the function of NlCA in
rice, we next investigated the subcellular localization of NlCA in both *Nicotiana benthamiana* leaf
cells and rice leaf protoplasts transiently expressing *NlCA:CFP*. In both cases, NlCA:CFP showed
a nonuniform distribution of CFP signal in the cell. Further colocalization studies with YFP signal
revealed that NlCA was colocalized in the cytoplasm but not in the nucleus (Fig. 1c*;*
Supplementary Fig. 6).

NlCA counters rapid intracellular acidification during BPH feeding of rice. BPH feeding on
rice plants can be divided into two phases based on the electropenetrography (EPG) waveforms
[23]. The first phase involves BPH's stylet penetrates the plant through the cell walls and cell
membranes of various epidermal and mesophyll cells until the stylet reaches the phloem sieve
cells inside the vascular system. In Nipponbare, the time to reach the phloem sieve cells can be
around 1.2 h before sustained phloem sap ingestion occurs at 3.8 h [23]. Our finding of the site of
NlCA action in the plant prompted us to test the hypothesis that NlCA might modulate rice cell pH
changes during BPH feeding. To directly test this hypothesis, we constructed a rice transgenic
line expressed a cytoplasmic ratiometric pH sensor, cyto-pHusion. Cyto-pHusion consists of the
tandem concatenation of enhanced green fluorescent protein (EGFP) and a monomeric red
fluorescent protein 1 (mRFP1) [24]. EGFP is highly sensitive to pH variation with the brightest
EGFP signal emitted at a pH of 7-8. EGFP fluorescence is gradually quenched at lower pH
values and totally quenched at pH values < 5. In contrast, mRFP1 is insensitive to pH changes in
the physiologically relevant range and serves as an internal reference. We placed 5th instar BPH

nymphs on the leaf sheath of cyto-pHusion-expressing rice plants and recorded GFP and RFP
signals of the feeding sites under microscopic observation at different time points with a 12 h test
period. As shown in the Fig. 3b-e, ds*GFP* BPH feeding did not elicit an obvious intracellular pH
change in rice cells during this period. In contrast, ds*NICA* BPH feeding induced a significant pH
decrease at 4 h and 8 h (Fig. 3c and d). The decreased ratio of EGFP: mRFP was caused by the
quenched EGFP signals after ds*NICA* BPH feeding, while the internal control, mRFP, kept at a
stable level (Supplementary Fig. 7). These results uncovered a previously uncharacterized plant
cellular response, rapid intracellular acidification, during the early stage of ds*NICA* BPH feeding
and demonstrated that BPH has evolved a critical virulence effector, NICA, to counter this novel
plant cellular response.

Because extracellular pH change is associated with plant responses to biotic stress [25,26], we
speculated that rapid intracellular acidification of ds*NICA* BPH feeding might induce defense gene
expression. To test this possibility, Nipponbare plants infested by ds*GFP* and ds*NICA* BPH were
examined for defense gene expression. We found that expression of defense response genes
(e.g., *OsNH1*, *OsNH2*, *OsWRKY45*, *OsWRKY13*) was significantly induced at a higher level by
ds*NICA* BPH infestation than by dsGFP BPH control at 4 h and 8 h (Fig. 3f-i), which is consistent
with the intracellular pH change at 4 h and 8 h (Fig. 3c-d). Taken together, these results indicate
that intercellular acidification is linked to rice defense activation.

Ectopic intracellular acidification causes defense gene expression in rice plants. Because
plant defense responses are important for limiting BPH survival in rice [27-30], we conducted
experiments to determine if intracellular acidification would trigger defense gene expression. Rice
is normally grown in Yoshida medium, pH of 4 [31]. To test if medium pH changes can modulate
defense responses in rice, we first grew rice seedlings in Yoshida medium until they reached the
5-leaf stage and then placed them to fresh Yoshida medium with pH adjusted to 2, 3, 4, 5, 6 and
7, respectively, for 48 hours. Interestingly, drastically increased expression of defense response
genes (e.g., *OsNH1*, *OsNH2*, *OsWRKY45*, *OsPBZ1*) was observed in Nipponbare plants at acidic
pH of 2 (Supplementary Fig. 8-9). The pHusion sensor plants exposed to external pH of 2
showed significant intracellular acidification (Fig. 3j-k). Survival rates of BPH insects were lower in
Nipponbare plants growing in media with pH of 2, compared to those growing in media with pH of
4 (Fig. 3l) These results suggest that cellular acidification is sufficient to induce defense gene
expression in rice and to reduce BPH's ability to survive on rice.

Rice defense responses are inhibited by NICA. As NICA stabilizes intracellular pH during WT
BPH feeding (Fig. 3a-d) and both ds*NICA* BPH feeding and ectopic intracellular acidification
causes activation of defense gene expression (Fig. 3f-i; Supplementary Fig. 8-9), we next tested

the hypothesis that a major function of NICA-mediated stabilization of intracellular pH may be to
prevent over-stimulation of downstream rice defense responses during feeding. Callose
deposition in the phloem sieve tubes is a classical defense response that is associated with
feeding of piercing-sucking insects [32]. We examined this response using aniline blue to stain
callose in the phloem sieve cells. Indeed, we found that fewer and smaller callose deposition was
found in the sieve plates of *NICA*-expressing leaf sheaths compared to those found in the sieve
plates of Nipponbare fed by BPH (Fig. 4a-f), demonstrating that *NICA* suppresses callose
deposition in sieve cell pates. Interestingly, the transcript levels of callose biosynthesis genes,
such as *OsGSL1*, *OsGSL3*, *OsGSL5* and *OsGns5,* were expressed at lower levels in *NICA*
transgenic plants compared to Nipponbare plants (Fig. 4g-j). We also measured the transcript
levels of defense marker genes (e.g., *OsNH1, OsNH2, OsWRKY45* and *OsWRKY13*) in
Nipponbare and *NICA*-expressing plants fed by ds*GFP* and ds*NICA* BPH. As shown in Fig. 4k-n,
induced expression of *OsNH1, OsNH2, OsWRKY13* and *OsWRKY45* by ds*NICA* BPH was
suppressed in *NICA* transgenic plants compared to those in Nipponbare plants. Collectively,
these results showed that NICA-mediated intracellular acidification is linked to downregulation of
callose deposition in phloem sieve cells as well as defense gene expression in rice plants.

Discussion

In this study, we provided evidence that intracellular acidification is a previously unrecognized
plant defense response that occurs during BPH feeding on rice. This finding was facilitated by our
attempt to understand the role of NICA in the rice-BPH interaction. We found that the *NICA*
transcript is detected mainly in the salivary glands (Fig. 1a) and that the NICA protein is found in
rice tissues fed by BPH (Fig. 1b). *NICA*-silenced (ds*NICA*) insect survived very poorly on at least
two independent cultivars of rice plants, Xiushui134 and Nipponbare ([6], Fig. 2), whereas *NICA*
expressing rice plants can restore the normal survival of *NICA*-silenced (ds*NICA*) insects (Fig. 2;
Supplementary Fig. 4), suggesting that NICA functions in plant cells. Using the cytoplasm pH
sensor, we found that NICA is required for BPH to maintain a normal plant cytoplasm pH during
BPH feeding (Fig. 3a-d). Pathogen/insect-derived effectors can be powerful molecular probes to
discovering novel plant regulators/responses to biotic attacks. Although other piercing/sucking
herbivores derived effectors, such as Btfer1, LsPDI1, LsSP1, Mp55, DNase II and BISP, have
been reported as defense-suppressive effectors [33-38], our discovery of intracellular acidification as
a plant defense response and our finding that brown planthopper (BPH) secretes NICA as a
counter-defense measure illustrates host cell pH homeostasis as a novel battlefield that has not
been revealed in any plant-biotic interactions.

Extracellular alkalinization of culture plant cells has long been recognized as a canonical plant
response to microbial elicitors as well as endogenous plant signals [26]. Extracellular alkalinization
caused by plant endogenous RAPID ALKALINIZATION FACTOR (RALF) peptides, for example,
are perceived by the FERONIA-family receptors [39,40]. There is evidence that this perception
causes phosphorylation of PM-localized H(+)-ATPase 2, resulting in the inhibition of proton
transport across the PM [41]. Notably, extracellular alkalinization has recently been shown to inhibit
or promote growth- or immunity-associated cell surface receptor functions through specific pH-
sensitive amino acid sensors [26]. In contrast, intracellular pH changes have so far escaped the
discovery by researchers until this study. Our demonstration of a link between intracellular
acidification and defense activation has laid a foundation for future discovery of potentially diverse
pH-sensitive intracellular regulators of defense responses, which could add a new dimension in
the study of plant-biotic interactions.

Because cellular pH alterations could potentially affect multiple biomolecules and, hence, multiple
cellular processes, future research should comprehensively define all cellular processes that are
affected by intracellular pH acidification. In this study, we found that this pH change is linked to
activation of callose deposition at phloem sieve cells and expression of defense response genes,
such as *OsNH1, OsNH2, OsPBZ1* and *OsWRKY45* (Fig. 3; and Supplementary Fig. 8-9). In
reverse, NICA-mediated intracellular pH stabilization dampens these defense responses (Fig. 4).
Callose deposition in phloem sieve cells, in particular, is a classical defense response to a variety
of sucking/piecing insects and is thought to limit nutrient flow during insect feeding [32,42-45].
Together, these results suggest that a major effect of NICA-mediated pH stabilization is to
prevent overactivation of defense responses during BPH feeding.

The mechanism by which NICA counters intracellular acidification is likely inherent to its
reversible inter-conversion of carbon dioxide and water into carbonic acid, protons and
bicarbonate ions. CAs are universally present in all organisms (Supplementary Fig. 10); other
piecing/sucking insects may use CAs or another mechanism to manipulate host intracellular pH
as part of their infestation strategy. Indeed, CA has been reported as a protein component of
saliva in rice green leafhopper, *Nephotettix cincticeps* [46] and aphid *Myzus persicae* [47]. In the case
of *M. persicae,* CA-II was shown to increase viral transmission via plant apoplastic acidification-
mediated acceleration of intracellular vesicular trafficking [47]. However, because NICA plays a
critical role in BPH's survival on rice plants *per se* (i.e., in the absence of viral infection), as shown
in this study, it is more likely that insect-secreted CAs constitute play a primary role in facilitating
insect survival by countering intracellular acidification-associated defense activation. In fact,
future research should examine if CA-mediated increase in viral transmission may be a

secondary consequence of defense suppression, which was not examined in the previous study
[47].

Discovering the role of pH regulation during plant response to biotic and abiotic stresses and
characterizing the impact of such pH alterations could be an important area for future research.
Because maintaining proper external and internal pH is critical for all forms of life, prokaryotic and
eukaryotic organisms, alike, have evolved mechanisms to achieve pH homeostasis. Facing the
fluctuating external pH, prokaryotes have evolved diverse mechanisms for sensing external pH.
For instance, the bimodal sensing of pH is employed by *Bacillus subtilis* and *Escherichia coli* [48].
Fungi employ a conserved pathway, mediated by Rim101 and PacC, to sense external pH [49]. It
has been reported that different subcellular compartments within the plant cell maintain different
pH values, presumably as part of carrying out their unique physiological functions [50]. The
demonstrated ability of NICA to counter stimulus-dependent pH changes in plant cells could
make NICA a useful molecular tool to modulate and broadly understand the effects of pH
stabilization on plant signaling and metabolic pathways in different cell types, organelles, and
tissues in plants by, for example, targeting NICA expression in specific tissues, cells, or
organelles. Additionally, the crucial role of NICA for BPH infestation of rice suggests that NICA is
an important target for chemical or trans-kingdom RNAi-based inactivation for the development of
novel BPH control strategies in plants.

Methods
Plant materials and insects. *Oryza sativa* ssp. Japonica cv. Nipponbare was used as the wild
type and for generating transgenic plants. The *Nilaparvata lugens* population used in this study
was originally derived from a rice field on the Huajiachi Campus of Zhejiang University, Hangzhou
(30° 16' N, 120° 11' E), China, in 2008, and artificially propagated for many generations until now.
BPH insects were reared in environmentally controlled growth chambers on a susceptible cultivar
rice variety Xiushui 134 at 27 ± 0.5°C and 50 ± 0.5% relative humidity under a 16-h light and 8-h
dark photoperiod in a modified Yoshida solution (in which ammonium nitrate was replaced by
1.06 M carbamide) with pH of 4.

Transgenic rice. The *NICA* coding sequence was amplified from BPH. Several *NICA* mutants
(*NICA-m1, NICA-m2, and NICA-m3*) were made by site-directed mutagenesis. The *NICA-m1*,
NICA-m2 and *NICA-m3* PCR products encoding proteins in which His159 and His161 were
mutated to Ala, Glu182 and His184 were replaced by Ala, and all seven conserved active site
residues, His159, His161, Glu171, Glu182, His184, Thr269 and Phe279, were changed to Ala,
respectively. The WT and mutant *NICA* PCR products were recombined into the intermediate
vector pDonor207 using the BP recombination kit (Invitrogen), and then recombined into the

destination vector pEarleygate104 using the LR recombination kit (Invitrogen) to generate

P$_{35S}$::NICA-YFP and *P$_{35S}$::mutant NICA-YFP* expression vectors. The transgenic rice plants were

generated by Agrobacterium-mediated transformation at BIOGLE GeneTech (Hangzhou Biogle

Co., LTD.) where rice plants were reared in a greenhouse at 27 ± 0.5°C and 60 ± 0.5% relative

humidity under a 16-h light and 8-h dark photoperiod.

***In situ* mRNA hybridization analysis.** Fresh salivary tissues were isolated from 100 5th instar

BPH and washed 3 times before being placed in the 4% paraformaldehyde (PFA) fix solution

(Servicebio, G1113) at 4°C overnight, dehydrated and embedded in paraffin. Ten µM-thick

sections were cut on glass slides, dewaxed and digested by Protease K (Servicebio, G1205) at

37°C for 15 min. After three washes in phosphate buffered saline (PBS) buffer (Servicebio,

G0020), the sections were incubated in pre-hybridization solution at 37°C for 1 h and then in

hybridization solution containing the *NICA* probe (1µM) at 42°C overnight. The PCR primer for

generating the probe was 5'-GGTTCTGTGATTGAGTGTTTGGATCTCCCTGCG-3' with a single-

end CY3 tag generated using the DIG RNA Labeling Kit (Roche Applied Science, Penzberg,

Germany) based on the manufacturer's instructions. The slides were stained with DAPI for 8 min

in the dark after being washed in saline sodium citrate (SSC) buffer (Servicebio, G3016-4) for

three times (2 × SSC, 10 min; 1 × SSC, 2×5 min; 0.5 × SSC, 10 min). Sections hybridized with

sense probe was used as a negative control and antisense probe was used to detect the target

mRNA. All steps were performed at 37°C and sections were examined under a Nikon

fluorescence microscope with a 40 × lens (330 nm – 380 nm for DAPI, 510 nm-560 nm for CY3).

Rice sheath sample preparation for LC-MS. Three independent sets of Nipponbare rice sheath

protein samples (three biological replicates) were collected after feeding by 5th instar nymphs as

follows. Ten 6-leaf-stage rice plants were individually placed into a breathable transparent plastic

bottle (5 cm in diameter, 30 cm in height) containing Yoshida medium, pH=4, and then fed by

~500 fifth instar BPH for 72 h. Next, the two outermost layers of leaf sheath surrounding the stem

were cut into small pieces, rinsed with ultrapure water for three times and ground to powder in

liquid nitrogen. Total protein was extracted in a buffer containing 60 mM Tris-HCl (pH=7.6), 2.5%

Glycerol, 6% SDS, 0.1 M DL-Dithiothreitol (DTT), 1x Protein inhibitor cocktail and 1 mM

phenylmethanesulfonyl fluoride. Protein samples were concentrated using 3-kDa filtration units

and shipped on dry ice to Shanghai Hoogen Biotech for shot-gun LC-MS analysis. Briefly, protein

samples digested by trypsin for 16 h at 37°C. The hydrolysates were then loaded into Zorbax

300SB-C18 peptide traps (Agilent Technologies, Wilmington, DE) and analyzed by a Q Exactive

mass spectrometer (Thermo Fisher). The raw LC-MS data were searched against the BPH

protein database (NCBI Accession No. PRJNA669454) using software MaxQuant 1.5.5.1.

Peptides with a high confidence score was extracted, and the database search was conducted
with false discovery rate (FDR) under 0.01.

RNA interference (RNAi). The nucleotide sequence (around 500 bp long) specific to the *NlCA*
coding region was cloned into the pMD 19-T vector (TAKARA). The double-stranded RNA was
synthesized through PCR amplification by using the Mega script T7 High Yield RNA Transcription
Kit (Vazyme, Nanjing, China), following the manufacturer's instructions. The procedure for gene
knockdown by RNAi was described previously [22]. Briefly, the 2nd and 5th instar BPH nymphs in
an insect rearing growth chamber were anaesthetized with carbon dioxide for ~10 s.
Approximately 50-250 ng of dsRNA was injected into BPH mesothorax using FemtoJet
(Eppendorf-Netheler-Hinz, Hamburg, Germany). The efficiency of RNA interference was
assessed 48 h after injection. RNA from 30 nymphs was extracted using the RNAiso Plus kit
(TaKaRa) according to the manufacturer's instructions as one biological replicate. The relative
level of the *NlCA* transcript was quantified using a CFX96TM Real-Time PCR Detection System
(Bio-Rad, Hercules, CA, USA) with primers shown in Supplementary Table 1. The Aequorea
Victoria green fluorescent protein (*GFP*) gene sequence was used as the control template in
RNAi experiments. Four biological replicates (30 individual insects pooled for each biological
replicate) were conducted.

BPH survival test on artificial diet. The 5th instar BPH nymphs injected with ds*GFP* or ds*NlCA*
were immediately placed on one-week-old rice seedlings (1-leaf-stage) for recovery in the small
breathable transparent plastic bottle for 3 days. Fifty normal-appearing BPH adults were gently
transferred into a shaded black plastic bottle of 5 cm in diameter and 15 cm in height, with
translucent double-layer parafilm loaded with fresh artificial diet. The first layer of parafilm closed
to BPH was needled 40 holes by a 1 ml syringe and fresh artificial diet AADM, pH of 6.0, was
filtered through a 0.22 micrometer needle-type bacterial filter (Sangon Biotech, F513142-0001).
Fresh artificial diet was replaced every 24 hours. The second layer of parafilm wrapped the
artificial diet to isolate the diet from bacteria in the air. Dead BPHs were separated from live BPHs
daily and mucus adhering to the inner wall of the black plastic bottle was cleaned daily to
eliminate possible impact on live BPHs. and the pH was changed from 6.8 to 6. There were three
biological replicates for each treatment, and each biological replicate includes 50 BPH adults.
Experiments were repeated three times with similar trends.

BPH survival test on rice. BPH nymphs with verified *NlCA* knockdown by RNAi were placed into
a small breathable transparent plastic bottle of 10 cm in diameter and 9 cm in height containing
one-week-old rice seedlings (1-leaf-stage) for 12 h to recover. About 20 healthy-appearing 2nd

instar nymphs were gently transferred into a longer breathable transparent plastic bottle of 5 cm
in diameter and 30 cm in height containing four-week-old rice plants (4- to 5-leaf-stage) and
observed for up to 15 days. The feeding space for BPH in each plastic bottle was around 200
cm3 divided by two sponges of 5 cm in diameter and 2 cm in thickness. The number of surviving
BPHs were counted daily on each plant. ds*GFP* BPH were used as control. The average of
survival rate on each seedling was calculated.

For BPH survival test on normal and acidic media growing rice plants, the early 5th instar BPH
nymphs inoculated ds*GFP* and ds*NICA* fed on one-week-old rice seedlings (1-leaf-stage) were
recovered in the small breathable transparent plastic bottle for 24 hours. The Nipponbare rice
plants (5-leaf stage) growing on normal liquid media (pH = 4) was transferred into fresh liquid
media with pH of 4 and 2 for 24 hours, respectively, before conducting BPH survival test. There
were five biological replicates for every treatment, and one biological replicate is one rice plant
fed by 20 BPHs. Experiments were repeated three times with similar trends.

pH assay. Transgenic rice plants expressing the cyto-pHusion pH sensor were grown in the
modified Yoshida medium as described above. Each 5-leaf-stage seedlings were fed by 30 5th
instar BPH, which were starved in the breathable plastic tubes for 3 h before feeding. No BPH
treatment was included as control. The microscopic images were taken from the phloem of rice
leaf sheaths that were fed by BPH for 1.5 h, 4 h, 8h and 12 h using an Olympus FV3000 with a
40.0 x objective. EGFP was imaged with λEm = 540 nm and the detection wavelength at 500-540
nm. mRFP was imaged with λEm = 570 nm and detection wavelength wat 570 to 620 nm. EGFP
fluorescence data and EGFP:mRFP ratio data were obtained using 64-bit professional edition Fiji
ImageJ2. Each treatment includes > 24 circular areas of leaf sheath with a diameter of 100 μm
each, including the feeding site at the center.

RNA isolation and quantitative real-time PCR (qRT-PCR). The total RNA was extracted using
the TRIzol reagent kit (Invitrogen), and then treated with Rnase-free Dnase I (Invitrogen)
according to the manufacturer's protocol. The RNA was reverse-transcribed using a cDNA
synthesis kit (TaKaRa) and one μg of each RNA samples was reverse transcribed. qRT-PCR was
performed from cDNA using SYBR Green qPCR mix (TaKaRa, RR601A) following the
manufacturer's instructions using the LightCycler 480 system (Roche Diagnostics). Fold changes
were calculated using the ΔCt method. Each assay was replicated at least three times, each with
three biological replicates (independent RNA preparations). The rice Actin gene
(*LOC_Os03g50885*) and the BPH 18S rRNA (*LOC_111047943*) gene were used as a reference.
Sequences of qRT-PCR primers are given in Supplementary Table 1.

Callose deposition in rice sheath. For counting callose deposition, four-leaf-stage rice plants
were infested with ~15 fifth instar BPH. The two outermost rice leaf sheaths were collected before
and after BPH feeding for 72 h and fixed in FAA solution as described by Guo et al. [29],
dehydrated and embedded in paraffin. The sections were cut to 5 um thickness on glass slides,
dewaxed and stained with 0.1 % aniline blue in 0.1 % 5 M K_2PHO4 for 10 min, and examined
under Zeiss confocal microscope with a 40× oil objective lens. The total area and number of
callose deposition in the rice sieve pores were counted. Each treatment has at least 400 cross
sections and 12 biological replicates.

Data Availability.
All data are presented within this paper.

Acknowledgments
We thank our lab members Richard Hilleary, Houping Wang, Yanru Hu and Xiao Han for their
comments and suggestions during this work. This study was supported by funding from the
Natural Science Foundation of China (Grants 31870259 to Y.J.), funds for The Young and Middle
Aged Academic and technical Leaders of Yunnan Province (202105AC160028 to Y.J.J., which
supported her research at Michigan State University), Zhejiang University (to support the
research of X.Y.Z. at Michigan State University), Howard Hughes Medical Institute to S.Y.H. and
the National Key Research and Development Plan in the 14th five year plan (2021YFD1401100)
to C.X.Z.

Author Contributions: S.Y.H., C.X.Z., Y.J.J. and X.Y.Z. conceptualized and designed this study;
Y.J.J., X.Y.Z., S.Q.L., Y.C.X., X.M.L., Y.P.Y., Z.Y.P., L.Z., J.B.L., and H.J.H. performed
experiments and analyzed data; C.X.Z. and S.Y.H. analyzed data. Y.J.J., X.Y.Z., C.X.Z. and
S.Y.H. wrote the paper with all authors approved the final article.

Competing Interest Statement: The authors declare no competing interest.

References

1. Jing, S. L. et al. Genomics of interaction between the brown planthopper and rice. *Curr. Opin.
   Insect Sci.* **19**, 82-87 (2017).
2. Sogawa, K. Feeding of the rice plant- and leafhoppers. *Rev. Plant Prot. Res.* **6**, 31-43 (1973).

3. Ling, K. C., Tiongco, E.R., & Aguiero, V.M. Rice ragged stunt, a new virus disease. *Plant Disease Report* **62**, 701-705 (1978).

4. Zhou, S. X. et al. OsMKK3, a Stress-Responsive Protein Kinase, Positively Regulates Rice Resistance to *Nilaparvata lugens* via Phytohormone Dynamics. *Int. J. Mol. Sci.* **20**, 3023-3038 (2019).

5. Du, B., Chen, R. Z., Guo, J. P. & He, G. C. Current understanding of the genomic, genetic, and molecular control of insect resistance in rice. *Mol. Breeding* **40**, 1103-3 (2020).

6. Huang, H. J. et al. Screening and Functional Analyses of *Nilaparvata lugens* Salivary Proteome. *J. Proteome Res.* **15**, 1883-1896 (2016).

7. Rao, W. et al. Secretome analysis and in planta expression of salivary proteins identify candidate effectors from the brown planthopper, *Nilaparvata lugens. Mol. Plant Microbe Interact.* **32**:227-239 (2019).

8. Liu, X. Q., Zhou, H. Y., Zhao, J., Hua, H. X. & He, Y. P. Identification of the secreted watery saliva proteins of the rice brown planthopper, *Nilaparvata lugens* (Stål) by transcriptome and Shotgun LC-MS/MS approach. *J. Insect Physiol.* **89**, 60-69 (2016).

9. Wang, Y. Y. et al. Responses of two contrasting genotypes of rice to brown planthopper. *Mol. Plant Microbe Interact.* **21**, 122-132 (2008).

10. Miles, P. W. Aphid saliva. *Biol. Reviews* **74**, 41-85 (1999).

11. Ji, R. et al. A Salivary Endo-beta-1,4-Glucanase Acts as an Effector That Enables the Brown Planthopper to Feed on Rice. *Plant Physiol.* **173**, 1920-1932 (2017).

12. Ye, W. F. et al. A salivary EF-hand calcium-binding protein of the brown planthopper *Nilaparvata lugens* functions as an effector for defense responses in rice. *Sci. Rep.* **7**, 40498 (2017).

13. Gong, G. et al. Salivary protein 7 of the brown planthopper functions as an effector for mediating tricin metabolism in rice plants. *Sci. Rep.* **12**, 3205 (2022).

14. Huang, H. J., Liu, C. W., Xu, H. J., Bao, Y. Y. & Zhang, C. X. Mucin-like protein, a saliva component involved in brown planthopper virulence and host adaptation. *J. Insect Physiol.* **98**, 223-230 (2017).

15. Shangguan, X. X. et al. A Mucin-Like Protein of Planthopper Is Required for Feeding and Induces Immunity Response in Plants. *Plant Physiol.* **176**, 552-565 (2018).

16. Meldrum NU, R. F. Carbonic anhydrase. Its preparation and properties. *J. Physiol.* **80**, 113-142 (1933).

17. Engineer, C. B. et al. Carbonic anhydrases, EPF2 and a novel protease mediate CO2 control of stomatal development. *Nature* **513**, 246-250 (2014).

18. Yu, X. J., McGourty, K., Liu, M., Unsworth, K. E. & Holden, D. W. pH sensing by intracellular Salmonella induces effector translocation. *Science* **328**, 1040-1043 (2010).

19. Smith, K. S. & Ferry, J. G. Prokaryotic carbonic anhydrases. *FEMS Microbiol. Rev.* **24**, 335-
366 (2000).

20. Henry, R. P. Multiple roles of carbonic anhydrase in cellular transport and metabolism. *Annu.*
Rev. Physiol. **58**, 523-538 (1996).

21. Zhou, Y. L. et al. Carbonic anhydrases CA1 and CA4 function in atmospheric CO2-modulated
disease resistance. *Planta* **251**, 75 (2020).

22. Xu, H. J. et al. Two insulin receptors determine alternative wing morphs in planthoppers.
Nature **519**, 464-467 (2015).

23. Ghaffar, M. B., Pritchard, J. & Ford-Lloyd, B. Brown planthopper (*N. lugens* Stål) feeding
behaviour on rice germplasm as an indicator of resistance. *PLoS One* **6**, e22137 (2011).

24. Gjetting, K. S. K., Ytting, C. K., Schulz, A. & Fuglsang, A. T. Live imaging of intra- and
extracellular pH in plants using pHusion, a novel genetically encoded biosensor. *J. Exp. Bot.* **63**,
12 (2012).

25. Kesten, C. et al. Pathogen-induced pH changes regulate the growth-defense balance in
plants. *EMBO J.* **38**, e101822 (2019).

26. Liu, L. et al. Extracellular pH sensing by plant cell-surface peptide-receptor complexes. *Cell*
185: 3341-3355 (2022).

27. Fu, Z. Q. & Dong, X. N. Systemic acquired resistance: turning local infection into global
defense. *Annu. Rev. Plant Biol.* **64**, 839-863 (2013).

28. Innes, R. The Positives and Negatives of NPR: A Unifying Model for Salicylic Acid Signaling
in Plants. *Cell* **173**, 1314-1315 (2018).

29. Guo, J. P. et al. *Bph6* encodes an exocyst-localized protein and confers broad resistance to
planthoppers in rice. *Nat. Genet.* **50**, 297-306 (2018).

30. He, J. et al. An R2R3 MYB transcription factor confers brown planthopper resistance by
regulating the phenylalanine ammonia-lyase pathway in rice. *Proc. Natl. Acad. Sci. USA* **117**,
271-277 (2020).

31. Yoshida S, F. D., Cock JH, Gomez KA. Laboratory manual for physiological studies of rice,
3rd edn. Manila, *The Philippines: International Rice Research Institute*, 61-64 (1976).

32. Hao, P. Y. et al. Herbivore-induced callose deposition on the sieve plates of rice: an important
mechanism for host resistance. *Plant Physiol.* **146**, 1810-1820 (2008).

33. Guo, J. et al. A tripartite rheostat controls self-regulated host plant resistance to insects.
Nature **618**:799-807 (2023).

34. Huang, H.-J. et al. Planthopper salivary sheath protein LsSP1 contributes to manipulation of
rice plant defenses. *Nat. Commun.* **14**, 737 (2023).

35. Huang, H.-J. et al. Salivary DNase II from *Laodelphax striatellus* acts as an effector that
suppresses plant defense. *New Phytol.* **224**, 860-874 (2019).

36. Elzinga, D. A., De Vos, M. & Jander, G. Suppression of plant defenses by a *Myzus persicae*
(green peach aphid) salivary effector protein. *Mol. Plant Microbe Interact.* **27**, 747-756 (2014).
37. Su, Q. et al. A salivary ferritin in the whitefly suppresses plant defenses and facilitates host
exploitation. *J. Exp. Bot.* **70**, 3343-3355 (2019).
38. Fu, J. et al. Planthopper-Secreted Salivary Disulfide Isomerase Activates Immune Responses
in Plants. *Front. Plant Sci.* **11**, 622513 (2020).
39. Murphy, E. & De Smet, I. Understanding the RALF family: a tale of many species. *Trends*
Plant Sci. **19**, 664-671 (2014).
40. Abarca, A., Franck, C. M. & Zipfel, C. Family-wide evaluation of RAPID ALKALINIZATION
FACTOR peptides. *Plant Physiol.* **187**, 996-1010 (2021).
41. Haruta, M., Sabat, G., Stecker, K., Minkoff, B. B. & Sussman, M. R. A peptide hormone and
its receptor protein kinase regulate plant cell expansion. *Science* **343**, 408-411 (2014).
42. Koh, E. J. et al. Callose deposition in the phloem plasmodesmata and inhibition of phloem
transport in citrus leaves infected with "*Candidatus Liberibacter asiaticus*". *Protoplasma* **249**,
687-697 (2012).
43. Ernst, A. M. et al. *Sieve element occlusion* (*SEO*) genes encode structural phloem proteins
involved in wound sealing of the phloem. *Proc. Natl. Acad. Sci. USA* **109**, E1980-1989 (2012).
44. Batailler, B. et al. Soluble and filamentous proteins in Arabidopsis sieve elements. *Plant Cell*
Environ. **35**, 1258-1273 (2012).
45. Jiang, Y. J., Zhang, C. X., Chen, R. Z. & He, S. Y. Challenging battles of plants with phloem-
feeding insects and prokaryotic pathogens. *Proc. Natl. Acad. Sci. USA* **116**: 23390-23397
(2019).
46. Hattori, M., Komatsu, S., Noda, H. & Matsumoto, Y. Proteome Analysis of Watery Saliva
Secreted by Green Rice Leafhopper, *Nephotettix cincticeps*. *PLoS One* **10**, e0123671 (2015).
47. Guo, H. et al. Salivary carbonic anhydrase II in winged aphid morph facilitates plant infection
by viruses. *Proc. Natl. Acad. Sci. USA* **120**, e2222040120 (2023).
48. Yang, Y. L. & Sourjik, V. Opposite responses by different chemoreceptors set a tunable
preference point in *Escherichia coli* pH taxis. *Mol. Microbiol.* **86**, 1482-1489 (2012).
49. Hayashi, M., Fukuzawa, T., Sorimachi, H. & Maeda, T. Constitutive activation of the pH-
responsive Rim101 pathway in yeast mutants defective in late steps of the MVB/ESCRT
pathway. *Mol. Cell. Biol.* **25**, 9478-9490 (2005).
50. Shen, J. et al. Organelle pH in the Arabidopsis Endomembrane System. *Mol. Plant* **6**, 1419-
1437 (2013).
558

Figures

Fig. 1| Initial characterization of *Nilaparvata lugens* carbonic anhydrase (NICA). a, *in situ* RNA hybridisation of salivary glands in 5-instar BPHs (lower row) using antisense NICA sequence as a probe. Sense *NICA* probe was used as a negative control (upper row). White arrows indicate glands. Nuclei are stained blue by 4',6-diamidino-2-phenylindole (DAPI). **b**, The amino acid sequence of NICA. The highlighted amino acid residues indicate the peptides detected in BPH-infested rice sheath tissue by LC-MS analysis. **c**, NICA is colocalized with the YFP signals in *N. benthamiana* (upper row) and rice cells (lower row). YFP and NICA-CFP fusion proteins were co-expressed in *N. benthamiana* leaf cells for 48 h using the *Agrobacterium*-mediated transient expression method. YFP and NICA-CFP fusion proteins were co-expressed in rice protoplasts 16 h after the corresponding DNA constructs were introduced into rice protoplasts via polyethylene glycol-mediated transformation. Experiments were repeated three times with similar trends.

Fig. 2| Effects of *NlCA* RNAi on BPH survival on rice cultivar Nipponbare. a, The *NlCA* transcript levels in BPH at 3 dpi post-injection of ds*GFP* or ds*NlCA* were determined by qRT-PCR (displayed as % of the *NlCA* transcript abundance in control BPH). Values are displayed as mean ± SEM of 4 experimental replicates (Two-way ANOVA; 3 biological replicates in each experiment and 30 individual insects pooled for each biological replicate). **b**, The daily survival rates of ds*NlCA*-injected BPH insects feeding on artificial diet. Values are displayed as mean ± SEM of 3 biological replicates (one biological replicate includes 50 individual BPH adults fed on artificial diet in a lucifugal plastic bottle). **c**, The daily survival rates of ds*NlCA*-injected BPH insects feeding on Nipponbare and NlCA-expressing line 1. Values are displayed as mean ± SEM of 8 biological replicates (Two-way ANOVA; 20 individual 2[nd] BPH nymphs per rice plant for each biological replicate). The black dashed line indicates significant differences (*P* < 0.05) in survival rates start day 9 between ds*NlCA* BPH on Nipponbare (Nip) and other treatments. **d**, Schematic display of

the NICA protein with the N-terminus displayed in bule, the C-terminus in green and the central

catalytic domain in yellow/orange. The amino acid sequence (158 - 280 aa) of the putative active

site is showed. Amino acid substitutions in NICA-m1, NICA-m2 and NICA-m3 are indicated in red.

e-f, The survival rates of 20 ds*GFP* and ds*NICA* BPH insects in Nipponbare and *NICA* transgenic

plants after 1- and 15-days post-infestation. Values are displayed as mean ± SEM (n ≥ 6

biological replicates; 20 individual 2nd BPH nymphs per rice plant for each biological replicate). ns

indicates no significant difference between treatments (Two-way ANOVA). Different letters

indicate statistically significant differences analyzed by two-way ANOVA (Tukey test, $P < 0.05$).

Experiments were repeated three times with similar trends.

Fig. 3| Role of NICA in maintaining intracellular pH of rice cells. a, Confocal microscopic

images at BPH feeding sites in Nipponbare plants expressing a ratiometric cytoplasmic pH

sensor (Cyto-pHusion) 4 h after placing 30 5[th]-instar BPH nymphs on each plant. Confocal

images from plants with no BPH feeding served as the control. **b-e**, EGFP:mRFP signal ratios at

1.5 h (a), 4 h (c), 8 h (d) and 12 h (e) of ds*GFP* or ds*NICA* BPH treatment compared with no BPH

control. EGFP was imaged at λ_{Ex} = 500 nm and λ_{Em} = 540 nm. mRFP was imaged at λ_{Ex} = 570

nm and λ_{Em} = 620 nm. Values are displayed as mean ± SEM (n ≥ 24 circular areas of leaf sheath

phloem with a diameter of 100 µm with the feeding site at the center). **f-i**, Expression of defense

response genes, *OsWRKY45*, *OsWRKY13*, *OsNH1* and *OsNH2*, in Nipponbare plants infested by

ds*GFP* or ds*NICA* BPH. Values are displayed as mean ± SEM of three biological replicates. Each

biological replicate represents pooled leaf sheaths from three individual rice plants fed by 20 5[th]

instar BPH nymphs per plant. **j**, Confocal microscopic images of Nipponbare plants expressing a

ratiometric cytoplasmic pH sensor (Cyto-pHusion) at 12 h after being transferred to Yoshida

medium at pH of 2. Confocal images from plants grown in Yoshida medium with pH of 4 served as the control. **k**, EGFP:mRFP signal ratios. EGFP was imaged at λ_{Ex} = 500 nm and λ_{Em} = 540 nm. mRFP was imaged at λ_{Ex} = 570 nm and λ_{Em} = 620 nm. Values are displayed as mean ± SEM (n ≥ 33 calculation area per condition). **l**, The 7[th]-day survival rates of ds*GFP*- and ds*NlCA*-injected BPH insects feeding on Nipponbare growing in media with different pHs. Values are displayed as mean ± SEM of 3 biological replicates (20 individual insects per each biological replicate). Different letters indicate statistically significant differences analyzed by two-way ANOVA (Tukey test, *P* < 0.05). Experiments were repeated at least three times with similar trends in this figure.

653 **Fig. 4| NICA dampens callose deposition and defense gene expression. a-d**, Callose
654 accumulation (bright blue fluorescence indicated by red arrows) on the sieve plates of
655 Nipponbare leaf sheaths (a and c) and *NICA-OE* leaf sheaths (b and d) plants before and 72 h

after BPH feeding. P, phloem; X, xylem. The pictures were taken under a Zeiss Microscope. (a-b) Cross-sections. (c-d) Longitudinal sections. **e**, Total areas of callose deposition in BPH-infested leaf sheaths of Nipponbare and *NICA*-OE. Each data point represents the total areas of callose deposition found in 300 cross-sections of each experiment. Values are displayed as mean ± SEM of four experiments. **f**, Total number of callose deposits in BPH-infested leaf sheaths of Nipponbare and *NICA*-OE. Each data point represents the total number of callose deposits found in 300 cross-sections of each experiment. Values are displayed as mean ± SEM of four experiments. **g-j**, Relative expression levels of the callose synthase gene *OsGSL1* (g), *OsGSL3* (h), *OsGSL5* (i) and *OsGns5* (j) in response to BPH feeding. Values are displayed as mean ± SEM of 3 biological replicates. Each biological replicate represents pooled leaf sheath form three individual rice plants fed by 20 5^{th} instar BPH nymphs per plant. **k-n**, Expression of defense marker gene *OsWRKY45* (k)*, OsWRKY13* (l)*, OsNH1* (m) and *OsNH1* (n) in response to BPH feeding. Values are displayed as mean ± SEM of 3 biological replicates. Each biological replicate represents pooled leaf sheaths from three individual rice plants fed by 20 5^{th} instar BPH nymphs per plant. ns indicates no significant difference between treatments (Two-way ANOVA). Different letters indicate statistically significant differences analyzed by two-way ANOVA (Tukey test, $P <$ 0.05). Experiments were repeated three times with similar trends.