Rapid intracellular acidification is a novel plant defense response countered by the brown
planthopper

Yanjuan Jiang[1,3,a*], Xiao-Ya Zhang[2,3,a], Shaoqin Li[4,5], Yu-Cheng Xie[2], Xu-Mei Luo[2], Yongping
Yang[4,5], Zhengyan Pu[1], Li Zhang[6,7], Jia-Bao Lu[8], Hai-Jian Huang[8], Chuan-Xi Zhang[2,8,*], Sheng
Yang He[3,6,7,9,*]

[1] State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan
University, Kunming 650091, China.

[2] Institute of Insect Science, Zhejiang University, Hangzhou, 310058, China.

[3] DOE Plant Research Laboratory, Michigan State University, East Lansing, MI, USA.

[4] CAS Key Laboratory of Tropical Plant Resources and Sustainable Use, Xishuangbanna Tropical
Botanical Garden, Chinese Academy of Sciences, Kunming, Yunnan 650223, China.

[5] University of Chinese Academy of Sciences, Beijing 100049, China.

[6] Howard Hughes Medical Institute, Duke University, Durham, NC, USA.

[7] Department of Biology, Duke University, Durham, NC, USA.

[8] State Key Laboratory for Managing Biotic and Chemical Threats to the Quality and Safety of
Agro products, Ningbo University, China.

[9] Lead contract

[a] Yanjuan Jiang and Xiao-Ya Zhang contributed equally to this work.
* Corresponding authors
Email: yanjuanjiang@ynu.edu.cn, chxzhang@zju.edu.cn or Shengyang.he@duke.edu

Highlights

● Rapid intracellular acidification is a novel plant defense response.
● Brown planthopper delivers into rice tissues a major virulence protein, carbonic anhydrase
(*Nl*CA), critical for survival.
● NlCA counters intracellular acidification to suppress plant defense.
● Results implicate existence of pH-responsive regulators of defense responses in the rice-brown
planthopper interaction.

In brief

The brown planthopper (BPH) is the most devasting insect pest in rice. Jiang *et al.* discovered that BPH secretes a salivary carbonic anhydrase (*NI*CA) to regulate the intracellular pH of the rice cell to facilitate its feeding and survival on rice plants and that NICA-counters host intracellular pH acidification to diminished defense responses. These findings uncovered that intracellular pH homeostasis is a previously uncharacterized battleground in plant-insect interactions, but also open a door to future discovery of pH-responsive intracellular regulators of defense responses, which could add a new research dimension in the study of plant-biotic interactions.

SUMMARY

The brown planthopper (BPH) is the most destructive insect pest in rice. Through a stylet, BPH secrets a plethora of salivary proteins into rice phloem cells as a crucial step of infestation. However, how various salivary proteins function in rice cells to promote insect infestation is poorly understood. Among them, one of the salivary proteins is predicted to be a carbonic anhydrase (NICA). The survival rate of the *NICA*-RNA interference (RNAi) BPH insects was extremely low on rice, indicating a vital role of this salivary protein in BPH infestation. We generated *NICA* transgenic rice plants and found that *NICA* expressed in rice plants could restore the ability of *NICA*-RNAi BPH to survive on rice. Next, we produced rice plants expressing the ratiometric pH sensor pHusion and found that *NICA*-RNAi BPH induced rapid intracellular acidification of rice cells during feeding. Further analysis revealed that both *NICA*-RNAi BPH feeding and artificial of lowering intracellular pH activated plant defense responses, and that NICA-mediated intracellular pH stabilization is linked to diminished defense responses, including reduced callose deposition at the phloem sieve plates and suppressed defense gene expression. Given the importance of pH homeostasis across the kingdoms of life, discovery of NICA-mediated intracellular pH modulation uncovered a new dimension in the interaction between plants and piecing/sucking insect pests. The crucial role of NICA for BPH infestation of rice suggests that NICA is a promising target for chemical or trans-kingdom RNAi-based inactivation for BPH control strategies in plants.

INTRODUCTION

The brown planthopper (BPH; *Nilaparvata lugens* Stål, Hemiptera, Delphacidae) is a monophagous insect pest of rice (*Oryza sativa* L.) found in all rice-growing Asian countries. The BPH sucks rice phloem sap via its stylet, causing leaf yellowing and wilting, stunted plant growth, reduced photosynthesis and ultimately death of rice plant[1]. During severe BPH outbreaks, tens of thousands of insects swarm on a rice field, resulting in the 'hopperburn' phenomenon, which is

characterized by large-scale wilting, yellowing and lethal drying of rice plants[2]. Besides direct

damages, the BPH may also indirectly damage rice plants by oviposition and transmitting viral

disease agents[3,4].

Application of chemical insecticides has been a main strategy for controlling BPH. Although it has

the advantages of having rapid effects on killing insects and low costs, use of insecticides leads

to environmental pollution and resistance of BPH to pesticides. In recent decades, breeding rice

resistant varieties to control BPH has attracted increasing attention. To date, more than 30

resistance genes have been found in the rice genome[5]. However, BPH often quickly evolves new

biological types that evade rice resistant genes. Therefore, additional BPH controlling methods

need to be developed to complement the current control measures toward long-term solutions of

achieving durable BPH resistance in rice. One method could be based on disruption of key steps

of BPH's natural infestation process. However, development of such methods will require a

comprehensive understanding of the basic biology of the BPH-rice interaction.

A critical step in BPH infestation of rice is secreting bioactive substances into the plant tissues

through the stylet[6-8]. Specifically, during feeding, BPH secretes both colloidal and watery

saliva[9,10]. The main function of the colloidal saliva is to form a saliva sheath around the piercing-

sucking mouthparts, stabilizing the overall feeding apparatus. The composition and function of

watery saliva is more complex, and it contains salivary proteins that are believed, in most cases,

to regulate various pathways in plant cells to enhance BPH feeding and survival in rice plants. For

example, NlEG1, a salivary endo-b-1,4-glucanase, degrades plant celluloses to help the BPH's

stylet reach to the phloem[11]. NlSEF1, an EF-hand Ca^{2+}-binding protein, interferes with calcium

signaling and H_2O_2 production during BPH feeding[12]. Salivary protein 7 is required for normal

feeding behavior and for countering accumulation of a defense compound, tricin[13]. On the other

hand, mucin-like protein (NlMLP) triggers defense responses in rice cells, including cell death,

callose deposition and up-regulation of pathogen-responsive genes[14,15].

Carbonic anhydrases (CAs) (EC 4.2.1.1) are zinc metalloenzymes that function as catalysts in the

bidirectional conversion of CO_2 and water into bicarbonate and protons[16]. There are at least five

distinct CA families (α-, β-, γ-, δ-, and ε-CAs) and three of them (α-, β-, and γ-CAs) are

ubiquitously distributed among animal, plant, and bacterial species. The widespread distribution

and adequate abundance of these CA families underline their evolutionary importance throughout

the kingdoms of life. CAs participate in a wide range of biological processes, such as pH

regulation, CO_2 homeostasis, stomatal aperture, and plant defense[17-21]. NlCA belongs to the α-

CA subfamily. Our previous study showed that *NlCA* expressed in BPH salivary glands[6].

Surprisingly, however, RNA interference (RNAi) of the *NlCA* transcript in BPH insects affects

neither pH maintenance within the salivary gland, watery saliva or gut, nor insect feeding
behavior or honeydew excretion, but greatly reduced survival of BPH on rice plants, suggesting a
critical function *in planta* via an unknown mechanism[6].

Here, we report that *NlCA*-RNAi BPH feeding results in rapid intracellular acidification of rice
cells. We found that NlCA is secreted into the rice tissues and functions as an effector that
stabilizes host cell intracellular pH, accompanied by suppression of defense responses, during
BPH feeding. Thus, we have uncovered intracellular pH homeostasis is a previously
uncharacterized battleground in plant-insect interactions.

Results

NlCA is detected in rice sheath tissues during BPH feeding. NlCA was previously found to be
highly expressed in salivary glands and present in the watery saliva of BPH fed on artificial diet[6].
We conducted a more detailed characterization of *NlCA* expression in this study. RNA *in situ*
hybridization showed that the expression level of *NlCA* was detectable throughout the principal
glands (PGs) and accessory glands (AGs), but not in A-follicle of the principal gland (APG)
(Figure 1A), which further raised the possibility that NlCA may be one of the "effector proteins"
secreted into the rice tissue during BPH feeding on rice plants. To test this possibility, we
compared protein profiles in the leaf sheaths of Nipponbare rice plants before and after BPH
feeding using liquid chromatograph-mass spectrometer (LC-MS). We found 8 NlCA-specific
peptides in BPH-fed leaf sheath tissue (Figure 1B), confirming that NlCA is secreted into the host
tissues during BPH feeding. As BPH is a phloem-feeding insect, its effector proteins, such as
NlCA, presumably act in the phloem. We therefore collected the phloem exudate from
Nipponbare rice leaf sheaths with or without BPH feeding for LC-MS analysis (Figure 1C). Six
NlCA-specific peptides were detected in the phloem exudate of Nipponbare plants fed by BPH
(Figure 1B), demonstrating that NlCA is delivered into phloem by BPH.

**Transgenic expression of *NlCA* in rice rescues the ability of *NlCA*-silenced BPH to feed
and survive.** To further clarify the site of function (i.e., in insect vs. in plant) of NlCA in the BPH-
rice interaction, we produced transgenic Nipponbare plants expressing *NlCA* (see Methods). A
total of 26 lines were produced and 6 lines were found to robustly express the *NlCA* transcript
(Supplemental Figure 1A). *NlCA*-expressing plants exhibited no noticeable changes in
appearance compared to Nipponbare plants (Supplemental Figure 1B–1D). *NlCA*-expressing
plant lines were propagated to T3 generation, and three lines were subjected to further
characterization, including BPH feeding. For BPH feeding assay, double-stranded RNA (dsRNA)
of *NlCA* (ds*NlCA*) or the control green fluorescent protein gene (ds*GFP*) was injected into 3[rd]

instar BPH nymphs to initiate RNAi of the *NICA* transcript[22]. Quantitative real-time PCR analysis

confirmed that the transcript levels of the *NICA* gene were reduced by 99% and 97%,

respectively, in tested individuals when compared with non-RNAi control and ds*GFP*-treated

insects (Figure 2A). There were no significant differences in the survival rates between ds*GFP*

and ds*NICA* BPH when fed on the artificial diet, indicating that silencing *NICA* expression has no

obvious impact on the basic physiology of BPH (Figure 2B). However, we found that the survival

rate of ds*NICA* BPH was sharply decreased, starting at day 9 post-infestation, to ~40% at day 14,

whereas the control ds*GFP* BPH survived normally on wild-type Nipponbare plants (Figure 2C).

This result is consistent with previous results conducted in the *japonica* cultivar Xiushui134 rice

plants[6], suggesting that the requirement of NICA for BPH survival is not specific to a specific rice

genotype. Strikingly, *NICA* transgenic plants almost fully restored the survival of *NICA*-silenced

BPH insects (Figure 2C; Supplemental Figure 1E–1F), demonstrating that *NI*CA expressed in the

host tissue (Figure 2D and 2E) can complement the infestation defect of *NICA*-silenced BPH

insects.

The conserved catalytic site amino acids of *NI*CA are critical to its function in rice. NlCA

contains several conserved amino acid residues predicted to be at the catalytic site of carbonic

anhydrases (Figure 2F). We asked if some of these conserved active site residues are critical to

the function of *NI*CA in the rice-BPH interaction. Accordingly, we expressed three different *NICA*

mutants in Nipponbare plants (Supplemental Figure 1A and 1G). "*NICA-m1*" plants contain a

NICA transgene in which His^{159} and His^{161} were mutated to Ala; "*NICA-m2*" plants contain a *NICA*

transgene in which Glu^{182} and His^{184} were replaced by Ala; and "*NICA-m3*" plants carry a *NICA*

transgene in which all seven conserved active site residues, His^{159}, His^{161}, Glu^{171}, Glu^{182}, His^{184},

Thr^{269} and Phe^{279}, were changed to Ala. Like *NICA* transgenic plants, transgenic plants

expressing these *NICA*-mutants exhibited no noticeable changes in appearance (Supplemental

Figure 1B-1D). We conducted BPH survival assay and found that, in contrast to plants expressing

wild-type *NICA*, the *NICA-m2* and *NICA-m3* plants could not fully rescue the infestation defect of

NICA-silenced BPH insects. The *NICA-m1* plants, on the other hand, could recover the infestation

defect of ds*NICA* BPH (Figure 2G and 2H; Supplemental Figure 1H). Thus, the conservative

residues Glu182 and His184 at the predicted catalytic site are indispensable for the function of

*NI*CA inside the plant cell.

After showing the requirement of catalytic site residues for the function of NICA in rice, we next

investigated the subcellular localization of NICA in both *Nicotiana benthamiana* leaf cells and rice

leaf protoplasts transiently expressing *NICA:CFP*. In both cases, NlCA:CFP showed a nonuniform

distribution of CFP signal in the cell. Further colocalization studies with YFP signal revealed that

NICA was predominantly localized in the cytoplasm (Figure 1D). There was some nuclear

localization of NlCA-CFP in rice protoplasts. Intact NlCA:CFP proteins were detected in these

localization experiments (Figure 1E).

NICA counters rapid intracellular acidification during BPH feeding of rice. BPH feeding on

rice plants can be divided into two phases based on the electropenetrography (EPG)

waveforms[23]. The first phase involves BPH's stylet penetrates the plant through the cell walls and

cell membranes of various epidermal and mesophyll cells until the stylet reaches the phloem

sieve cells inside the vascular system. In Nipponbare, the time to reach the phloem sieve cells

can be around 1.2 h before sustained phloem sap ingestion occurs at 3.8 h[23]. Our finding of the

site of NICA action in the plant prompted us to test the hypothesis that NICA might modulate rice

cell pH changes during BPH feeding. To directly test this hypothesis, we constructed a rice

transgenic line expressed a cytoplasmic ratiometric pH sensor driven by the 35S promoter, cyto-

pHusion. Cyto-pHusion consists of the tandem concatenation of enhanced green fluorescent

protein (EGFP) and a monomeric red fluorescent protein 1 (mRFP1)[24]. EGFP is highly sensitive

to pH variation with the brightest EGFP signal emitted at a pH of 7–8. EGFP fluorescence is

gradually quenched at lower pH values and totally quenched at pH values <5. In contrast, mRFP1

is insensitive to pH changes in the physiologically relevant range and serves as an internal

reference. We placed 5^{th} instar BPH nymphs on the leaf sheath of cyto-pHusion-expressing rice

plants and recorded GFP and RFP signals of the feeding sites under microscopic observation at

different time points with a 12 h test period. BPH can move around from one feeding site to

another during the 12-h feeding experiments. The average feeding time at a feeding site is

around 8 h. Therefore, the first 8 h of cellular responses best capture mostly synchronized rice-

BPH interactions at a typical feeding cycle. As shown in the Figure 3A–D, ds*NICA* BPH feeding

induced a significant pH decrease in the phloem sieve elements/companion cells at 4 h and 8 h

(Figure 3C and D). The decreased ratio of EGFP: mRFP was caused by the quenched EGFP

signals after ds*NICA* BPH feeding, while the internal control, mRFP, kept at a stable level

(Supplemental Figure 2A–2H). In contrast, ds*GFP* BPH feeding did not elicit an obvious

intracellular pH change in the phloem sieve elements/companion cells during this period. These

results uncovered a previously uncharacterized plant cellular response, rapid intracellular

acidification, during the early stage of ds*NICA* BPH feeding and demonstrated that BPH has

evolved a critical virulence effector, NICA, to counter this novel plant cellular response.

Interestingly, we noted that the control ds*GFP* wild-type BPH feeding induced slight intracellular

acidification at 12 h (Figure 3E). At this late time point of feeding, most BPHs would have left their

initial feeding sites and have relocated to new feeding sites. In our analysis, we could not

distinguish between old and new feeding sites. It is therefore likely that active BPH feeding (i.e.,

active NlCA injection) is needed to maintain intracellular pH homeostasis at all feeding sites.

Next, we conducted experiments to determine if intracellular acidification is a highly localized cell-
type specific response or a spreading local response. We imaged and calculated the intracellular
pH in the mesophyll cells and epidermal cells at the feeding site. We found that intracellular
acidification occurred in the mesophyll cells and epidermal cells (Supplemental Figure 2I–2L) in
response to feeding by ds*NICA* BPH, albeit to a lesser degree compared to that in the sieve
elements/companion cells. In contrast, rice plants fed by ds*GFP* wild-type BPH responded with
an initial slight intracellular acidification at 1.5 h, but then consistently maintained a more alkalized
intracellular pH in the mesophyll and epidermal cells compared to non-fed rice plants and ds*NICA*
BPH (Supplemental Figure 2I–2L).

Because extracellular pH change is associated with plant responses to biotic stress[25,26], we
speculated that rapid intracellular acidification of ds*NICA* BPH feeding might induce defense gene
expression. To test this possibility, Nipponbare plants infested by ds*GFP* and ds*NICA* BPH were
examined for defense gene expression. Commonly used marker genes (e.g., *OsNH1*, *OsNH2*,
OsWRKY45, *OsWRKY13*) in the studies of rice defense against BPH were selected[27-30]. We
found that expression of those genes was significantly induced at a higher level by ds*NICA* BPH
infestation than by ds*GFP* BPH control at 4 h and 8 h (Figure 3F–I), which is consistent with the
intracellular pH change at 4 h and 8 h (Figure 3C–D). These results indicate that intercellular
acidification is linked to rice defense activation.

Ectopic intracellular acidification causes defense gene expression in rice plants. Because
plant defense responses are important for limiting BPH survival in rice[28,31-33], we conducted
experiments to determine if intracellular acidification would trigger defense gene expression. Rice
is normally grown in Yoshida medium, pH of 4[34]. To test if medium pH changes can modulate
defense responses in rice, we first grew rice seedlings in Yoshida medium until they reached the
5-leaf stage and then placed them to fresh Yoshida medium with pH adjusted to 2, 3, 4, 5, 6 and
7, respectively, for 48 hours. Interestingly, drastically increased expression of defense response
genes (e.g., *OsNH1*, *OsNH2*, *OsWRKY45*, *OsPBZ1*) was observed in Nipponbare plants at acidic
pH of 2 (Supplemental Figure 3A–H). The pHusion sensor plants exposed to external pH of 2
showed intracellular acidification that simulates pH changes observed during BPH feeding (Figure
3J–K). Survival rates of BPH insects were lower in Nipponbare plants growing in media with pH of
2, compared to those growing in media with pH of 4 (Figure 3L). Rice growth medium of pH 2 was
used to attain an intracellular pH similar to that during BPH feeding (Figure 3K). These results
suggest that cellular acidification is sufficient to induce defense gene expression in rice and to
reduce BPH's ability to survive on rice.

Rice defense responses are inhibited by NICA. As NICA stabilizes intracellular pH during WT
BPH feeding (Figure 3A–D) and both ds*NICA* BPH feeding and ectopic intracellular acidification
causes activation of defense gene expression (Figure 3F–I; Supplemental Figure 3), we next
tested the hypothesis that a major function of NICA-mediated stabilization of intracellular pH may
be to prevent over-stimulation of downstream rice defense responses during feeding. Callose
deposition in the phloem sieve tubes is a classical defense response that is associated with
feeding of piercing-sucking insects[35]. We examined this response using aniline blue to stain
callose in the phloem sieve cells. Indeed, we found that fewer and smaller callose deposition was
found in the sieve plates of *NICA*-expressing leaf sheaths compared to those found in the sieve
plates of Nipponbare fed by BPH (Figure 4A–H) at 72 h after BPH feeding, demonstrating that
*NI*CA suppresses callose deposition in sieve cell pates. Furthermore, ds*NICA* BPH induced
higher expression of callose biosynthesis genes, such as *OsGSL1*, *OsGSL3*, *OsGLS5* and
OsGns5, than the ds*GFP* control BPH; however, the induction of these genes was greatly
compromised in *NICA* transgenic plants with or without BPH treatment (Figure 4I–L). We also
measured the transcript levels of defense marker genes (e.g., *OsNH1, OsNH2, OsWRKY45* and
OsWRKY13) in Nipponbare and *NICA*-expressing plants fed by ds*GFP* and ds*NICA* BPH. As
shown in Figure 4M–P, induced expression of *OsNH1, OsNH2, OsWRKY13* and *OsWRKY45* by
ds*NICA* BPH was suppressed in *NICA* transgenic plants compared to those in Nipponbare plants.
Collectively, these results showed that NICA-mediated intracellular pH homeostasis is linked to
downregulation of callose deposition in phloem sieve cells as well as defense gene expression in
rice plants.

Discussion
262
In this study, we provided evidence that intracellular acidification is a previously unrecognized
plant defense response that occurs during BPH feeding on rice. This finding was facilitated by our
attempt to understand the role of NICA in the rice-BPH interaction. We found that the *NICA*
transcript is detected mainly in the salivary glands (Figure 1A) and that the NICA protein is found
in rice tissues, including the phloem sap, fed by BPH (Figure 1B). *NICA*-silenced (ds*NICA*) insect
survived very poorly on at least two independent cultivars of rice plants, Xiushui134[6] and
Nipponbare (Figure 2), whereas *NICA* expressing rice plants can restore the normal survival of
NICA-silenced (ds*NICA*) insects (Figure 2; Supplemental Figure 1), suggesting that NICA
functions in plant cells. Using the cytoplasm pH sensor, we found that NICA is required for BPH to
maintain a normal plant cytoplasm pH during BPH feeding (Figure 3A–D; Supplemental Figure 2).
Pathogen/insect-derived effectors can be powerful molecular probes to discovering novel plant
regulators/responses to biotic attacks. Although other piercing/sucking herbivores derived
effectors, such as Btfer1, LsPDI1, LsSP1, Mp55, DNase II and BISP, have been reported as

defense-suppressive effectors[36-41], our discovery of intracellular acidification as a plant defense
response and our finding that brown planthopper (BPH) secretes NICA as a counter-defense
measure illustrates host cell pH homeostasis as a novel battlefield that has not been revealed in
any plant-biotic interactions.

Extracellular alkalinization of culture plant cells has long been recognized as a canonical plant
response to microbial elicitors as well as endogenous plant signals[26]. Extracellular alkalinization
caused by plant endogenous RAPID ALKALINIZATION FACTOR (RALF) peptides, for example,
are perceived by the FERONIA-family receptors[42,43]. There is evidence that this perception
causes phosphorylation of PM-localized H(+)-ATPase 2, resulting in the inhibition of proton
transport across the PM[44]. Notably, extracellular alkalinization has recently been shown to inhibit
or promote growth- or immunity-associated cell surface receptor functions through specific pH-
sensitive amino acid sensors[26]. In contrast, defense regulation by intracellular pH changes has so
far escaped the discovery by researchers until this study. Our demonstration of a link between
intracellular acidification and defense activation has laid a foundation for future discovery of
potentially diverse pH-sensitive intracellular regulators of defense responses, which could add a
new dimension in the study of plant-biotic interactions.

Because cellular pH alterations could potentially affect multiple biomolecules and, hence, multiple
cellular processes, future research should comprehensively define all cellular processes that are
affected by intracellular pH acidification. In this study, we found that this pH change is linked to
activation of callose deposition at phloem sieve cells and expression of defense response genes,
such as *OsNH1, OsNH2, OsPBZ1* and *OsWRKY45* (Figure 3; Supplemental Figure 3). In
reverse, NICA-mediated intracellular pH stabilization dampens these defense responses (Figure
4). Callose deposition in phloem sieve cells, in particular, is a classical defense response to a
variety of sucking/piecing insects and is thought to limit nutrient flow during insect feeding[35,45-48].
Together, these results suggest that a major effect of NICA-mediated pH stabilization is to
prevent overactivation of defense responses during BPH feeding.

The mechanism by which NICA counters intracellular acidification is likely inherent to its
reversible inter-conversion of carbon dioxide and water into carbonic acid, protons and
bicarbonate ions. CAs are universally present in all organisms (Supplemental Figure 4); other
piecing/sucking insects may use CAs or another mechanism to manipulate host intracellular pH
as part of their infestation strategy. Indeed, CA has been reported as a protein component of
saliva in rice green leafhopper, *Nephotettix cincticeps*[49], and aphid *Myzus persicae*[50]. In the case
of *M. persicae,* CA-II was shown to increase viral transmission via plant apoplastic acidification-
mediated acceleration of intracellular vesicular trafficking[50]. However, because NICA plays a

critical role in BPH's survival on rice plants *per se* (i.e., in the absence of viral infection), as shown

in this study, it is more likely that insect-secreted CAs constitute play a primary role in facilitating

insect survival by countering intracellular acidification-associated defense activation. In fact,

future research should examine if CA-mediated increase in viral transmission may be a

secondary consequence of defense suppression, which was not examined in the previous

study[50].

Discovering the role of pH regulation during plant response to biotic and abiotic stresses and

characterizing the impact of such pH alterations could be an important area for future research.

Because maintaining proper external and internal pH is critical for all forms of life, prokaryotic and

eukaryotic organisms, alike, have evolved mechanisms to achieve pH homeostasis. Facing the

fluctuating external pH, prokaryotes have evolved diverse mechanisms for sensing external pH.

For instance, the bimodal sensing of pH is employed by *Bacillus subtilis* and *Escherichia coli*[51].

Fungi employ a conserved pathway, mediated by Rim101 and PacC, to sense external pH[52]. It

has been reported that different subcellular compartments within the plant cell maintain different

pH values, presumably as part of carrying out their unique physiological functions[53]. The

demonstrated ability of NICA to counter stimulus-dependent pH changes in plant cells could

make NICA a useful molecular tool to modulate and broadly understand the effects of pH

stabilization on plant signaling and metabolic pathways in different cell types, organelles, and

tissues in plants by, for example, targeting NICA expression in specific tissues, cells, or

organelles. Additionally, the crucial role of NICA for BPH infestation of rice suggests that NICA is

an important target for chemical or trans-kingdom RNAi-based inactivation for the development of

novel BPH control strategies in plants.

Acknowledgments

We thank our lab member Richard Hilleary for his comments and suggestions during this work.

This study was supported by fundings from the Natural Science Foundation of China (32360082

and 31870259, which supported her research at Michigan State University) and the Yunnan

Fundamental Research Projects (202401AS070122 and 202105AC160028) to Y.J.J., Zhejiang

University (to support the research of X.Y.Z. at Michigan State University), Howard Hughes

Medical Institute to S.Y.H. and the National Key Research and Development Plan in the 14th five-

year plan (2021YFD1401100) to C.X.Z.

Author Contributions: S.Y.H., C.X.Z., Y.J.J. and X.Y.Z. conceptualized and designed this study;

Y.J.J., X.Y.Z., S.Q.L., Y.C.X., X.M.L., Y.P.Y., Z.Y.P., L.Z., J.B.L., and H.J.H. performed

experiments and analyzed data; C.X.Z. and S.Y.H. analyzed data. Y.J.J., X.Y.Z., C.X.Z. and

S.Y.H. wrote the paper with all authors approved the final article.

Competing Interest Statement: The authors declare no competing interest.

Figure legends

Figure 1. Initial characterization of *Nilaparvata lugens* carbonic anhydrase (NICA).

(A) *in situ* RNA hybridisation of salivary glands in 5-instar BPHs. Left, a schematic diagram of

BPH salivary glands, including the principal glands (PG), the accessory glands (AG) and the A-

follicle of the principle gland (APG). Right, *NICA* expression was detectable in the PG (red) using

antisense *NICA* sequence as a probe. Sense *NICA* probe was used as a negative control. Nuclei

are stained blue by 4',6-diamidino-2-phenylindole (DAPI). Scale bar = 50 μm. (B) The amino acid

sequence of NICA. The highlighted amino acid residues indicate the peptides detected in BPH-

infested rice sheath tissue by LC-MS analysis. The underlined amino acid residues indicate the

peptides detected in phloem exudate of BPH-infested rice by LC-MS analysis. (C) A schematic

diagram of the phloem exudate collection. (D) NICA is colocalized with the YFP signals in *N.*

benthamiana (upper row; Scale bar = 10 μm) and rice cells (lower row; Scale bar = 5 μm). YFP

and NICA-CFP fusion proteins were co-expressed in *N. benthamiana* leaf cells for 48 h using the

Agrobacterium-mediated transient expression method. YFP and NICA-CFP fusion proteins were

co-expressed in rice protoplasts 16 h after the corresponding DNA constructs were introduced

into rice protoplasts via polyethylene glycol-mediated transformation. Experiments were repeated

three times with similar trends. (E) NICA-CFP-HA fusion protein levels are detected with anti-HA

(Zenbio, 301113) in *N. benthamiana* leaves and rice protoplasts. Protein samples were extracted

from NICA-CFP-HA expressing *N. benthamiana* and rice protoplasts. Proteins from mock *N.*

benthamiana leaves and rice protoplasts were as negative controls (N). Ponceau S staining of

Rubisco shows protein loading control. See also Table S1.

Figure 2. Effects of *NICA* RNAi on BPH survival on rice cultivar Nipponbare.

(A) The *NICA* transcript levels in BPH at 3 dpi post-injection of ds*GFP* or ds*NICA* were

determined by qRT-PCR (displayed as % of the *NICA* transcript abundance in control BPH).

Values are displayed as mean ± SEM of 4 experimental replicates (Two-way ANOVA; 3 biological

replicates in each experiment and 30 individual insects pooled for each biological replicate). (B)

The daily survival rates of ds*NICA*-injected BPH insects feeding on artificial diet. Values are

displayed as mean ± SEM of 3 biological replicates (one biological replicate includes 50 individual

BPH adults fed on artificial diet in a lucifugal plastic bottle). (C) The daily survival rates of ds*NICA*-

injected BPH insects feeding on Nipponbare and *NICA*-expressing line 1. Values are displayed as

mean ± SEM of 8 biological replicates (Two-way ANOVA; 20 individual 2nd BPH nymphs per rice

plant for each biological replicate). The black dashed line indicates significant differences ($P <$

0.05) in survival rates start day 9 between ds*NICA* BPH on Nipponbare (Nip) and other

treatments. (D) The amino acid sequence of NICA. The highlighted amino acid residues indicate

the peptides detected in phloem exudate of *NICA*-OE transgenic rice plants by LC-MS analysis.

(E) NICA:YFP protein is detectable in the phloem exudate of *NICA*-OE transgenic plants by anti-

GFP antibody (Abmart, 20004). (F) Schematic display of the NICA protein with the N-terminus

displayed in bule, the C-terminus in green and the central catalytic domain in yellow/orange. The

amino acid sequence (158–280 aa) of the putative active site is showed. Amino acid substitutions

in NICA-m1, NICA-m2 and NICA-m3 are indicated in red. (G and H) The survival rates of 20

ds*GFP* and ds*NICA* BPH insects in Nipponbare and *NICA* transgenic plants after 1-day (G) and

15-days (H) post-infestation. Values are displayed as mean ± SEM ($n \geq 6$ biological replicates; 20

individual 2^{nd} BPH nymphs per rice plant for each biological replicate). ns indicates no significant

difference between treatments (Two-way ANOVA). Different letters indicate statistically significant

differences analyzed by two-way ANOVA (Tukey test, $P < 0.05$). Experiments were repeated

three times with similar trends. See also Figure S1.

Figure 3. Role of NICA in maintaining intracellular pH of rice cells.

(A) Confocal microscopic images at BPH feeding sites in Nipponbare plants expressing a

ratiometric cytoplasmic pH sensor (Cyto-pHusion) 4 h after placing 30 5^{th}-instar BPH nymphs on

each plant. A 40.0 x objective was used to capture every feeding site on the tiled 1 cm x 1 cm

section of rice leaf sheath. Confocal images from plants with no BPH feeding served as the

control. Scale bar = 20 μm. (B–E) EGFP:mRFP signal ratios at 1.5 h (B), 4 h (C), 8 h (D) and 12 h

(E) of ds*GFP* or ds*NICA* BPH treatment compared with no BPH control. EGFP was imaged at λ_{Ex}

= 500 nm and λ_{Em} = 540 nm. mRFP was imaged at λ_{Ex} = 570 nm and λ_{Em} = 620 nm. Values are

displayed as mean ± SEM ($n \geq 24$ circular areas of leaf sheath phloem with a diameter of 100 μm

with the feeding site at the center). (F–I) Expression of defense response genes, *OsWRKY45* (F),

OsWRKY13 (G), *OsNH1* (H) and *OsNH2* (I), in Nipponbare plants infested by ds*GFP* or ds*NICA*

BPH. Values are displayed as mean ± SEM of three biological replicates. Each biological

replicate represents pooled leaf sheaths from three individual rice plants fed by 20 5^{th} instar BPH

nymphs per plant. (J) Confocal microscopic images of Nipponbare plants expressing a ratiometric

cytoplasmic pH sensor (Cyto-pHusion) at 12 h after being transferred to Yoshida medium at pH of

2. Confocal images from plants grown in Yoshida medium with pH of 4 served as the control.

Scale bar = 15 μm. (K) EGFP:mRFP signal ratios. EGFP was imaged at λ_{Ex} = 500 nm and λ_{Em} =

540 nm. mRFP was imaged at λ_{Ex} = 570 nm and λ_{Em} = 620 nm. Values are displayed as mean ±

SEM ($n \geq 33$ calculation area per condition). (L) The 7^{th}-day survival rates of ds*GFP*- and

ds*NICA*-injected BPH insects feeding on Nipponbare growing in media with different pHs. Values

are displayed as mean ± SEM of 3 biological replicates (20 individual insects per each biological

replicate). Different letters indicate statistically significant differences analyzed by two-way

ANOVA (Tukey test, $P < 0.05$). Experiments were repeated three times with similar trends. See

also Figure S2 and Figure S3.

Figure 4. NICA dampens callose deposition and defense gene expression.

(A–F) Callose accumulation (bright blue fluorescence indicated by red arrows) on the sieve plates

of Nipponbare leaf sheaths (A and D) and *NICA*-OE leaf sheaths (B and E) plants at 72 h after

BPH feeding. P, phloem. The pictures were taken under a Zeiss Microscope. (A–B) Cross-

sections. Scale bar = 5 μm. (D–E) Longitudinal sections. Scale bar = 10 μm. (C and F) The bright

field views of cross and longitudinal phloem sections, respectively. (G) Total areas of callose

deposition in BPH-infested leaf sheaths of Nipponbare and *NICA*-OE. Each data point represents

the total areas of callose deposition found in 300 cross-sections of each experiment. Values are

displayed as mean ± SEM of four experiments. (H) Total number of callose deposits in BPH-

infested leaf sheaths of Nipponbare and *NICA*-OE. Each data point represents the total number of

callose deposits found in 300 cross-sections of each experiment. Values are displayed as mean ±

SEM of four experiments. (I–L) Relative expression levels of the callose synthase gene *OsGSL1*

(I), *OsGSL3* (J), *OsGSL5* (K) and *OsGns5* (L) in response to BPH feeding. Values are displayed

as mean ± SEM of 3 biological replicates. Each biological replicate represents pooled leaf sheath

form three individual rice plants fed by 20 5th instar BPH nymphs per plant. (M–P) Expression of

defense marker gene *OsWRKY45* (M), *OsWRKY13* (N), *OsNH1* (O) and *OsNH1* (P) in response

to BPH feeding. Values are displayed as mean ± SEM of 3 biological replicates. Each biological

replicate represents pooled leaf sheaths from three individual rice plants fed by 20 5th instar BPH

nymphs per plant. ns indicates no significant difference between treatments (Two-way ANOVA).

Different letters indicate statistically significant differences analyzed by two-way ANOVA (Tukey

test, $P < 0.05$). Experiments were repeated three times with similar trends.

STAR METHODS

Detailed methods are provided in the online version of this paper and include the following:

● KEY RESOURCES TABLE

● RESOURCE AVAILABILITY

○ Lead contact

○ Materials availability

○ Data and code availability

● EXPERIMENTAL MODEL AND SUBJECT DETAILS
○ Plant materials and insects

● METHOD DETAILS
○ Plant materials and insects
○ Transgenic rice
○ *In situ* mRNA hybridization analysis
○ Rice sheath sample preparation for LC-MS
○ RNA interference (RNAi)
○ BPH survival test on artificial diet
○ BPH survival test on rice
○ pH assay
○ RNA isolation and quantitative real-time PCR (qRT-PCR).
○ Callose deposition in rice sheath

● QUANTIFICATION AND STATISTICAL ANALYSIS
467
SUPPLEMENTAL INFORMATION
Supplemental Figure 1. Generation of *NICA-OE* and mutant *NICA-OE* plants, and survival rates
of BPH, related to Figure 2.
Supplemental Figure 2. Quantification of fluorescent signals in leaf sheaths of Nipponbare
plants expressing the cyto-pHusion ratiometric pH sensor in response to BPH feeding, related to
Figure 3.
Supplemental Figure 3. Defense response genes are modulated by ectopic pH manipulation in
rice, related to Figure 3.
Supplemental Figure 4. Phylogenic analysis of carbonic anhydrase proteins from representative
bacteria, insect, mammal, and plant species, related to Discussion section.
Supplemental Table 1. List of detected BPH-secreted salivary proteins, related to Figure 1.
479
480 **REFERENCES**
481

1. Jing, S.L., Zhao, Y., Du, B., Chen, R.Z., Zhu, L.L., and He, G.C. (2017). Genomics of
interaction between the brown planthopper and rice. Curr. Opin. Insect Sci. *19*, 82–87.
2. Sogawa, K. (1973). Feeding of the rice plant- and leafhoppers. Rev. Plant Prot. Res. *6*, 31–43.
3. Ling, K. C., Tiongco, E.R., and Aguiero, V.M. (1978). Rice ragged stunt, a new virus disease.
Plant Disease Report *62*, 701–705.

4. Zhou, S.X., Chen, M.T., Zhang, Y.B., Gao, Q., Noman, A., Wang, Q., Li, H., Chen, L., Zhou,
P.Y., Lu, J., *et al.* (2019). OsMKK3, a Stress-Responsive Protein Kinase, Positively Regulates
Rice Resistance to *Nilaparvata lugens* via Phytohormone Dynamics. Int. J. Mol. Sci. *20*, 3023–
3038.

5. Du, B., Chen, R.Z., Guo, J.P. and He, G.C. (2020). Current understanding of the genomic,
genetic, and molecular control of insect resistance in rice. Mol. Breeding *40*, 24.

6. Huang, H.J., Liu, W.C., Huang, X.H., Zhou, X., Zhuo, J.C., and Zhang, C.X. (2016). Screening
and Functional Analyses of *Nilaparvata lugens* Salivary Proteome. J. Proteome Res. *15*, 1883–
1896.

7. Rao, W., Zheng, X.H., Liu, B.F., Guo, Q., Guo, J.P., Wu, Y., Shangguan, X.X., Wang, H.Y., Wu,
D., Wang, Z.Z. *et al.* (2019). Secretome analysis and in planta expression of salivary proteins
identify candidate effectors from the brown planthopper, *Nilaparvata lugens*. Mol. Plant Microbe
Interact. *32*, 227–239.

8. Liu, X.Q., Zhou, H.Y., Zhao, J., Hua, H.X. and He, Y.P. (2016). Identification of the secreted
watery saliva proteins of the rice brown planthopper, *Nilaparvata lugens* (Stål) by transcriptome
and Shotgun LC-MS/MS approach. J. Insect Physiol. *89*, 60–69.

9. Wang, Y.Y., Wang, X.L., Yue, H.Y., Chen, R.Z., Zhu, L.L., He, R.F., and He, G.C. (2008).
Responses of two contrasting genotypes of rice to brown planthopper. Mol. Plant Microbe
Interact. *21*, 122–132.

10. Miles, P.W. (1999). Aphid saliva. Biol. Reviews *74*, 41–85.

11. Ji, R., Ye, W.F., Chen, H.D., Zeng, J.M., Li, H., Yu, H.X., Li, J.C., and Lou, Y.G. (2017). A
Salivary Endo-beta-1,4-Glucanase Acts as an Effector That Enables the Brown Planthopper to
Feed on Rice. Plant Physiol. *173*, 1920–1932.

12. Ye, W.F., Yu, H.X., Jian, Y.K., Zeng, J.M., Ji, R., Chen, H.D., and Lou, Y.G. (2017) A salivary
EF-hand calcium-binding protein of the brown planthopper *Nilaparvata lugens* functions as an
effector for defense responses in rice. Sci. Rep. *7*, 40498.

13. Gong, G., Yuan, L.Y., Li, Y.F., Xiao, H.X., Li, Y.F., Zhang, Y., Wu, W.J., and Zhang, Z.F.
(2022). Salivary protein 7 of the brown planthopper functions as an effector for mediating tricin
metabolism in rice plants. Sci. Rep. *12*, 3205.

14. Huang, H.J., Liu, C.W., Xu, H.J., Bao, Y.Y. and Zhang, C.X. (2017). Mucin-like protein, a
saliva component involved in brown planthopper virulence and host adaptation. J. Insect
Physiol. *98*, 223–230.

15. Shangguan, X.X., Zheng, J., Liu, B.F., Zhao, Y., Wang, H.Y., Wang, Z.Z., Guo, J.P., Rao,
W.W., Jing, S.L., Guan, W., *et al.* (2018). A Mucin-Like Protein of Planthopper Is Required for
Feeding and Induces Immunity Response in Plants. Plant Physiol. *176*, 552–565.

16. Meldrum N.U., and Roughton, F.J.W. (1933). Carbonic anhydrase. Its preparation and
properties. J. Physiol. *80*, 113–142.

17. Engineer, C.B., Ghassemain, M., Anderson, J.C., Peck, S.C., Hu, H.H., and Schroeder, J.I.
(2014). Carbonic anhydrases, EPF2 and a novel protease mediate CO2 control of stomatal
development. Nature *513*, 246–250.

18. Yu, X.J., McGourty, K., Liu, M., Unsworth, K.E. and Holden, D.W. (2010). pH sensing by
intracellular Salmonella induces effector translocation. Science *328*, 1040–1043.

19. Smith, K.S. and Ferry, J.G. (2000). Prokaryotic carbonic anhydrases. FEMS Microbiol. Rev.
24, 335–366.

20. Henry, R. P. (1996). Multiple roles of carbonic anhydrase in cellular transport and metabolism.
Annu. Rev. Physiol. *58*, 523–538.

21. Zhou, Y.L. Vroegop-Vos, I.A., Van Dijken, A.J.H., Van der Does, D., Zipfel, C., Pieterse,
C.M.J., Van Wees, S.C.M. (2020). Carbonic anhydrases CA1 and CA4 function in atmospheric
CO2-modulated disease resistance. Planta *251*, 75.

22. Xu, H.J., Xue, J., Lu, B., Zhang, X.C., Zhuo, J.C., He, S.F., Ma, X.F., Jiang, Y.Q., Fan, H.W.,
Xu, J.Y., *et al*. (2015). Two insulin receptors determine alternative wing morphs in planthoppers.
Nature *519*, 464–467.

23. Ghaffar, M.B., Pritchard, J. and Ford-Lloyd, B. (2011). Brown planthopper (*N. lugens* Stål)
feeding behaviour on rice germplasm as an indicator of resistance. PLoS One *6*, e22137.

24. Gjetting, K.S.K., Ytting, C.K., Schulz, A. and Fuglsang, A.T. (2012). Live imaging of intra- and
extracellular pH in plants using pHusion, a novel genetically encoded biosensor. J. Exp. Bot. *63*,
12.

25. Kesten, C., Gámez-Arjona, F.M., Menna, A., Scholl, S., Dora, S., Huerta, A.I., Huang, H.Y.,
Tintor, N., Kinoshita, T., Rep, M., *et al*. (2019). Pathogen-induced pH changes regulate the
growth-defense balance in plants. EMBO J. *38*, e101822.

26. Liu, L., Song, W., Huang, S., Jiang, K., Moriwaki, Y., Wang, Y., Men, Y., Zhang, D., Wen, X.,
Han, Z., *et al*. (2022). Extracellular pH sensing by plant cell-surface peptide-receptor
complexes. Cell *185*, 3341–3355.

27. Du, B., Zhang, W.L., Liu, B.F., Hu, J., Wei, Z., Shi, Z.Y., He, R.F., Zhu, L.L., Chen, R.Z., Han,
B., *et al*. (2009). Identification and characterization of *Bph14*, a gene conferring resistance to
brown planthopper in rice. Proc. Natl. Acad. Sci. USA *106*, 22163–22168.

28. Guo, J.P., Xu, C.X., Wu, D., Zhao, Y., Qiu, Y.F., Wang, X.X., Ouyang, Y.D., Cai, B.D., Liu, X.,
Jing, S.L. *et al*. (2018). *Bph6* encodes an exocyst-localized protein and confers broad
resistance to planthoppers in rice. Nat. Genet. *50*, 297–306.

29. Wang, Y., Cao, L., Zhang, Y., Cao, C., Liu, F., Huang, F., Qiu, Y., Li, R., and Lou, X. (2015).
Map-based cloning and characterization of *BPH29*, a B3 domain-containing recessive gene
conferring brown planthopper resistance in rice. J. Exp. Bot. *66*, 6035–6045.

30. Hu, L., Wu, Y., Wu, D., Rao, W., Guo, J., Ma, Y., Wang, Z., Shangguan, X., Wang, H., Xu, C.,
et al. (2017). The coiled-coil and nucleotide binding domains of BROWN PLANTHOPPER
RESISTANCE14 function in signaling and resistance against planthopper in rice. Plant Cell *29*,
3157–3185.

31. Fu, Z.Q. and Dong, X.N. (2013). Systemic acquired resistance: turning local infection into
global defense. Annu. Rev. Plant Biol. *64*, 839–863.

32. Innes, R. (2018). The Positives and Negatives of NPR: A Unifying Model for Salicylic Acid
Signaling in Plants. Cell *173*, 1314–1315.

33. He, J., Liu, Y.Q., Yuan, D.Y., Duan, M.J., Liu, Y.L., Shen, Z.J., Yang, C.Y., Qiu, Z.Y., Liu,
D.M., Wen, P.Z., *et al.* (2020). An R2R3 MYB transcription factor confers brown planthopper
resistance by regulating the phenylalanine ammonia-lyase pathway in rice. Proc. Natl. Acad.
Sci. USA *117*, 271–277.

34. Yoshida, S.F.D., Cock, J.H., and Gomez, K.A. (1976). Laboratory manual for physiological
studies of rice, 3rd edn. Manila, The Philippines: International Rice Research Institute, 61–64.

35. Hao, P.Y., Liu, C.X., Wang, Y.Y., Chen, R.Z., Tang, M., Du, B., Zhu, L.L., and He, G.C.
(2008). Herbivore-induced callose deposition on the sieve plates of rice: an important
mechanism for host resistance. Plant Physiol. *146*, 1810–1820.

36. Guo, J., Wang, H., Guan, W., Guo, Q., Wang, J., Yang, J., Peng, Y., Shan, J., Gao, M., Shi,
S., *et al.* (2023). A tripartite rheostat controls self-regulated host plant resistance to insects.
Nature *618*:799–807.

37. Huang, H.J., Wang, Y.Z., Li, L.L., Lu, H.B., Lu, J.B., Wang, X., Ye, Z.X., Zhang, Z.L., He, Y.J.,
Lu, G., *et al.* (2023). Planthopper salivary sheath protein LsSP1 contributes to manipulation of
rice plant defenses. Nat. Commun. *14*, 737.

38. Huang, H.J., Cui, J.R., Xia, X., Chen, J., Ye, Y.X., Zhang, C.X., and Hong, X.Y. (2019).
Salivary DNase II from *Laodelphax striatellus* acts as an effector that suppresses plant defense.
New Phytol. *224*, 860–874.

39. Elzinga, D.A., De Vos, M. and Jander, G. (2014). Suppression of plant defenses by a *Myzus*
persicae (green peach aphid) salivary effector protein. Mol. Plant Microbe Interact. *27*, 747–756.

40. Su, Q., Peng, Z., Tong, H., Xie, W., Wang, S., Wu, Q., Zhang, J., Li, C., and Zhang, Y. (2019).
A salivary ferritin in the whitefly suppresses plant defenses and facilitates host exploitation. J.
Exp. Bot. *70*, 3343–3355.

41. Fu, J., Shi, Y., Wang, L., Zhang, H., Li, J., Fang, J., Ji, R. (2020). Planthopper-Secreted
Salivary Disulfide Isomerase Activates Immune Responses in Plants. Front. Plant Sci. *11*,
622513.
42. Murphy, E. and De Smet, I. (2014). Understanding the RALF family: a tale of many species.
Trends Plant Sci. *19*, 664–671.
43. Abarca, A., Franck, C.M. and Zipfel, C. (2021). Family-wide evaluation of RAPID
ALKALINIZATION FACTOR peptides. Plant Physiol. *187*, 996–1010.
44. Haruta, M., Sabat, G., Stecker, K., Minkoff, B.B. and Sussman, M.R. (2014). A peptide
hormone and its receptor protein kinase regulate plant cell expansion. Science *343*, 408–411.
45. Koh, E.J. Zhou, L., Williams, D.S., Park, J., Ding, N., Duan, Y.P., Kang, B.H. (2012). Callose
deposition in the phloem plasmodesmata and inhibition of phloem transport in citrus leaves
infected with "*Candidatus Liberibacter asiaticus*". Protoplasma *249*, 687–697.
46. Ernst, A.M., Jekat, S.B., Zielonka, S., Muller, B., Neumann, U., Ruping, B., Twyman, R.M.,
Krzyzanek, V., Prufer, D., and Noll, G.A. (2012). Sieve element occlusion (SEO) genes encode
structural phloem proteins involved in wound sealing of the phloem. Proc. Natl. Acad. Sci. USA
109, E1980–1989.
47. Batailler, B., Lemaitre, T., Vilaine, F., Sanchez, C., Renard, D., Cayla, T., Beneteau, J.,
Dinant, S. (2012). Soluble and filamentous proteins in Arabidopsis sieve elements. Plant Cell
Environ. *35*, 1258–1273.
48. Jiang, Y.J., Zhang, C.X., Chen, R.Z. and He, S.Y. (2019). Challenging battles of plants with
phloem-feeding insects and prokaryotic pathogens. Proc. Natl. Acad. Sci. USA *116*: 23390–
23397.
49. Hattori, M., Komatsu, S., Noda, H. and Matsumoto, Y. (2015). Proteome Analysis of Watery
Saliva Secreted by Green Rice Leafhopper, Nephotettix cincticeps. PLoS One *10*, e0123671.
50. Guo, H., Zhang, Y.J., Li, C.W., Shi, Q.Y., Zhu-Salzman, K.Y., Ge, F., and Sun, Y.C. (2023).
Salivary carbonic anhydrase II in winged aphid morph facilitates plant infection by viruses. Proc.
Natl. Acad. Sci. USA *120*, e2222040120.
51. Yang, Y.L. and Sourjik, V. (2012). Opposite responses by different chemoreceptors set a
tunable preference point in Escherichia coli pH taxis. Mol. Microbiol. *86*, 1482–1489.
52. Hayashi, M., Fukuzawa, T., Sorimachi, H. and Maeda, T. (2005). Constitutive activation of the
pH-responsive Rim101 pathway in yeast mutants defective in late steps of the MVB/ESCRT
pathway. Mol. Cell. Biol. *25*, 9478–9490.
53. Shen, J., Zeng, Y.L., Zhuang, X.H., Sun, L., Yao, X.Q., Pimpl, P., Jiang, L.W. (2013).
Organelle pH in the Arabidopsis Endomembrane System. Mol. Plant *6*, 1419–1437.

54. Gaupels, F., Knauer, T. and van Bel, A.J. (2008). A combinatory approach for analysis of protein sets in barley sieve-tube samples using EDTA-facilitated exudation and aphid stylectomy. J. Plant Physiol. *165*, 95–103.

55. Du, B., Wei, Z., Wang, Z., Wang, X., Peng, X., Du, B., Chen, R., Zhu, L., and He, G. (2015). Phloem-exudate proteome analysis of response to insect brown plant-hopper in rice. J. Plant Physiol. *183*, 13–22.

A

B

```
  1  MELFFHISMI  VTIMVVSLTG  MGRTTRLFRS  RGGPMAPSEP

 41  EFPTELEGED  RFLVNPLNVD  EVLSFGPRAP  QGSSPGFPWD

 81  RSQLPQPQQS  PIDICRCKCK  SMTYQPLKLS  YAKTSRNDGI

121  VVEISNSGHG  VNLKVPKAEA  KPFLSGGMLS  NRYIYDNIHC

161  HWGRNDIMGS  EHFIDAESYP  MECHMVTFNE  KYGTINEALK

201  NKQDGVVVLT  WFYKLANEDN  KRLQNFLQYV  PSVRENGAKI

241  FRADQNALFF  PLEVLPINQE  YFAYPGSLTT  PPFTEAVTFI

281  ILPIAVGISK  DQLALLRTLM  DDKNPSKPVL  SNRREIQTLN

321  HRTVYCVKGR  VQAISREIPP  QQSNDVYNQI  HRYC
```

C

Phloem exudate collection

D

E

A

Relative transcript level (%)

- Control: a
- dsGFP: a
- dsNICA: b

B

Survival rate (%) — Artificial diet — ns

BPH infestation time (Day)

- dsGFP BPH
- dsNICA BPH

C

Survival rate (%)

BPH infestation time (Day)

- Nip-dsGFP
- Nip-dsNICA
- NICA-OE/dsGFP BPH
- NICA-OE/dsNICA BPH

P < 0.05

D

```
1   MELFFHISMI  VTIMVVSLTG  MGRTTRLFRS  RGGPMAPSEP  EFPTELEGED  RFLVNPLNVD
61  EVLSFGPRAP  QGSSPGFPWD  RSQLPQPQQS  PIDICRCKCK  SMTYQPLKLS  YAKTSRNDGI
121 VVEISNSGHG  VNLKVPKAEA  KPFLSGGMLS NRYIYDNIHC  HWGRNDIMGS  EHFIDAESYP
181 MECHMVTFNE  KYGTINEALK  NKQDGVVVLT  WFYKLANEDN  KRLQNFLQYV  PSVRENGAKI
241 FRADQNALFF  PLEVLPINQE  YFAYPGSLTT  PPFTEAVTFI  ILPIAVGISK  DQLALLRTLM
301 DDKNPSKPVL SNRREIQTLN HRTVYCVKGR  VQAISREIPP  QQSNDVYNQI  HRYC
```

E

Phloem exudate

Nip   NICA-YFP

— 60 kDa

α-GFP

F

N-terminal	Catalytic	C-terminal

1   87   158   active site   280   322   355

```
         158                                            185        268        280
NICA     I H C H W G R N D I M G S E H F I D A E S Y P M E C H M - - - - L T T P P F T E A V T F I
NICA-m1  I A C A W G R N D I M G S E H F I D A E S Y P M E C H M - - - - L T T P P F T E A V T F I
NICA-m2  I H C H W G R N D I M G S E H F I D A E S Y P M A C A M - - - - L T T P P F T E A V T F I
NICA-m3  I A C A W G R N D I M G S A H F I D A E S Y P M A C A M - - - - L A T P P F T E A V T A I
```

G

Survival rate (%)

- Nipponbare
- NICA-OE
- NICA-m1-OE
- NICA-m2-OE
- NICA-m3-OE

ns   ns

dsGFP   dsNICA
Day 1

H

Survival rate (%)

dsGFP (Day 15): a a a a a
dsNICA: b a a b b

dsGFP   dsNICA
Day 15