In vivo long-term voltage imaging by genetically encoded voltage
indicator reveals spatiotemporal dynamics of spinal cord neuronal
populations during development

Asuka Shiraishi[1], Ayane Hayashi[2], Narumi Fukuda[1], Mari Hishinuma[1], Hiroaki
Miyazawa[1], Sachiko Tsuda[1][§]

1 Division of Life Science, Graduate School of Science and Engineering, Saitama
University, 255 Shimo-Okubo, Sakura-ku, Saitama City, Saitama 338-8570,
Japan
2 Department of Regulatory Biology, Faculty of Science, Saitama University, 255
Shimo-Okubo, Sakura-ku, Saitama City, Saitama 338-8570, Japan

[§]Author for correspondence (stsuda@mail.saitama-u.ac.jp)

Abstract

A central question of brain development lies in how individual neurons emerge and organize communities to acquire various functions. Voltage imaging provides unique approaches to address this by enabling simultaneous, non-invasive, in vivo recording of voltage dynamics from a population of cells. Recently, genetically encoded voltage indicators (GEVIs) facilitate cell-type specific imaging of voltage dynamics. However, it has not been applied to brain development. Here, we applied ArcLight, a GEVI utilizing voltage-sensitive domain, to zebrafish and established experimental approaches for analyzing voltage and morphology during development, focusing on the spinal cord and cerebellum. We initially demonstrated that Arclight was widely distributed in the neural tissues. By voltage imaging, we successfully visualized the coordinated, spontaneous activity of spinal cord neurons in their early stage of development at a high spatiotemporal resolution, at subcellular and population levels. Hyperpolarization and subthreshold signals were also detected. Finally, long-term voltage imaging during development revealed the process of changes in voltage dynamics in neuron populations, which were accompanied by axonal outgrowth. Voltage imaging could greatly contribute to our understanding of the functional organization of the nervous system during development.

Introduction

The nervous system consists of a large number of neurons that perform various functions. During development, these neurons are generated and communicate with each other to form functional networks. For a deeper understanding of brain function and development, it is essential to understand how the brain is formed and organized. However, such attempts are often hindered by the vast number of neurons. Optical methods provide a powerful approach to solving these difficulties because light facilitates non-invasive access to many neurons, which is difficult with electrophysiological techniques.

Voltage imaging enables fast, direct, and simultaneous detection of membrane potentials in a population of neurons, providing an excellent system for circuit analysis [1-3]. Its high speed and directness allow the detection of action potentials, hyperpolarization, and subthreshold responses, which are difficult to examine using calcium imaging, partly because of the slow kinetics of the indicators and the dynamics of calcium [4, 5]. Voltage imaging began with activity recording using voltage-sensitive dyes (VSDs) [6]. Subsequently, various types of VSDs have been developed to improve the brightness, signal-to-noise ratio, and wider variations in color, expanding our understanding of network dynamics and development in various brain regions [7-9]. In recent years, the development of genetically encoded voltage indicators (GEVIs) has progressed rapidly, resulting in cell-type-specific voltage imaging, which is difficult to achieve with VSDs. This yielded GEVIs with faster kinetics and brighter fluorescence, overcoming prior weaknesses in the speed and signal-to-noise ratio [10-12]. There are several types of GEVIs, based on their structures. One is a mosaic protein that uses the voltage-sensitive domain of voltage-sensitive phosphatases [13-16]. Another class of GEVIs uses the voltage sensitivity of microbial rhodopsins [17-20]. These GEVIs have been applied to the study of brain function for various brain regions [21-25]. However, in most studies, GEVI voltage imaging was used to undermine the function of neural circuits, and very few attempts have been made to understand brain development.

Here, to examine the dynamics of neuronal populations and their development,

we applied ArcLight to zebrafish, and established a series of recording
approaches to expand the power of voltage imaging in developmental biology.
Zebrafish is an optimal model system for developmental biology and
neuroscience. The transparent and relatively small brain of zebrafish, together
with various transgenic lines, provides an excellent system for analyzing the
functional organization of the brain in various behavioral modalities by the whole-
brain recordings [26, 27]. Several GEVIs have been successfully used for voltage
imaging of the zebrafish brain, mainly to analyze the circuit functions [19, 20, 23,
24, 28]. In this study, we focus on the spinal cord and cerebellum, which are
suitable for studying neural circuit development because of their rapid
development, ease of observation, and accumulated information on circuit
organization [29, 30].

The spinal cord contains neural circuits that play fundamental roles in
locomotion. These circuits drive coordinated movements of the left and right sides
of the body by a central pattern generator, which is a fundamental element of
locomotion [31]. At an early stage of development, this coordinated activity
emerges as the spontaneous firing of spinal cord neurons, which is considered
important for circuit formation [32]. Although electrophysiological and optical
studies have elucidated the firing dynamics of these neurons, the mechanisms
by which this coordinated activity emerges remain to be understood [33-35]. This
transient activity, known as fetal movement in mammals, has been observed as
coiling movements during early development in zebrafish. In the spinal cord, this
process leads to periodic bursts of alternating populations of neurons on the left
and right sides [36, 37]. Previous studies have shown that such activity patterns
emerge from uncorrelated activities within a few hours of early development [34,
37-39]. However, the mechanisms by which individual neurons emerge and
constitute network function remain unclear. This is partly because previous
studies have utilized either electrophysiology or calcium imaging, creating a large
gap between precise information about the synaptic profiles of individual neurons
and the population information obtained by calcium imaging. To solve this issue,
we need to obtain voltage information from a population of neurons during the

course of circuit formation.

Cumulative information has been obtained about cerebellar circuits, which play

pivotal roles in motor coordination, learning, and cognition, and are evolutionally

highly conserved [29, 40]. Inhibition and high-frequency activity are prominent in

the cerebellum and contribute to various adaptive properties of the cerebellum,

including long-term depression, complex and simple spiking of Purkinje cells,

modularity of the cerebellar circuitry, and activity-dependent pruning during

development [41-45]. While VSD imaging has long been used to uncover

cerebellar circuit dynamics, GEVI in the cerebellum, especially in its

developmental process, is still limited [23, 42, 46].

Materials and Methods
115

Fish lines
116

Wild-type zebrafish (Danio rerio) with an RW genetic background or nacre line

were used. Tg(elavl3:GAL4-VP16), Tg(hspGFFDMC28C;UAS:RFP),

Tg(gSA2AzGFF152B;UAS:RFP), and Tg(UAS:lynRFP) fish lines have been

described previously and obtained from the National BioResource Project [23, 47,

48]. The Tg lines were crossed with the nacre line [49]. All procedures were

performed in accordance with a protocol approved by the Saitama University

Committee on Animal Research.

DNA Construction and establishment of the ArcLight transgenic line
125

To generate the pTol2-UAS-ArcLight plasmid, ArcLight-A242 sequences were

excited from the pCS2+/ArcLight-A242 plasmid (Addgene) and inserted into

pTol2-UAS-MCSF-polyA, in which a UAS and polyA sequence was integrated into

the pTol2-MCS plasmid [50]. pTol2-Aldoca-ArcLight-Q239-polyA construct was

generated using a Purkinje cell-specific promoter, aldorace C, and the ArcLight-

Q239 sequence. The ArcLight-Q239 and aldorace sequences were extracted

from pCS2+/ArcLight-Q239 and aldoca-5k-pGEM-T_easy, respectively, and

inserted into pTol2-UAS-MCSF-polyA pDNA. To establish pTol2-UAS-ArcLight

line, Tol2 plasmid DNA and transposase messenger RNA (mRNA) were co-

injected into one-cell-stage embryos. Genotyping was performed using the

primers ArcLight-A242 (5′-TCGACGGTATCGATAAGCTTGC-3′,5′-

CACCTCCCCCTGAACCTGAAACATA-3′).

Transient expression of ArcLight in the cerebellum and spinal cord

For the spinal cord, 25-ng/μl Tol2-UAS-ArcLight-A242-polyA plasmid DNA was

co-injected with 25-ng/μl *Tol2 transposase* mRNA into one-cell-stage embryos of

Tg(elavl3:GAL4-VP16) embryos, which express Gal4 in neurons [51] After the

injection, the embryos were incubated at 23°C and then transferred to 28°C by

18 hour post fertilization (hpf). ArcLight-positive embryos were screened at the

19-20 somite stage, followed by voltage imaging of the spinal cord at 20 to 24 hpf.

For the cerebellum, 5-40 ng/μl Tol2-Aldoca-ArcLight-Q239-polyA plasmid DNA

was co-injected with 25 ng/μl *Tol2 transposase* mRNA into one-cell-stage nacre

embryos. ArcLight-positive embryos were screened at 4 to 5 days post

fertilization (dpf), followed by voltage imaging at 5 to 6 dpf.

Confocal imaging

Zebrafish embryos were paralyzed in 0.02% tricaine and mounted on 2% low

melting-point agarose (Sigma-Aldrich). Optical sectioning was performed using

an Olympus FV1000 or a Nikon A1R (detector: A1-DU4) confocal microscope.

ImageJ, NIS-Elements (Nikon), and Imaris (Bitplane) software were used to

perform the image analysis.

Voltage imaging of the spinal cord

Embryos from 16 to 23 hpf were paralyzed with tubocurarine (0.5 mM, Sigma-

Aldrich), with their tail tips cut for better penetration of tubocurarine [37, 52]. Then,

embryos were mounted on 1.8% low melting-point agarose containing 0.1-0.2

mM tubocurarine. After embedding the embryos onto 1% agarose, an

extracellular solution (134 mM NaCl, 2.9 mM KCl, 2.1 mM $CaCl_2$, 1.2 mM $MgCl_2$,

10 mM HEPES, and 10 mM glucose, adjusted to pH 7.8 with NaOH) containing

150 mM tubocurarine was added. The agarose around the tail was removed using

a surgical blade to allow efficient transfer of the drug solution. For high-speed

voltage imaging, a fluorescence microscope (FN-1, Nikon) equipped with a

CMOS camera (ORCA-Flash4.0, Hamamatsu Photonics) was used with a

40×/0.8 NA water-immersion lens. Fluorescence images were acquired using

NIS-Elements (Nikon) software at approximately 16-200 Hz. High-speed confocal

scanning was performed using a confocal microscope (Nikon A1R) with a 25×/1.0

NA water-immersion lens. For drug treatment, tetrodotoxin (1 µM, TOCRIS) was

added to the extracellular solution using a peristatic pump (World Precision

Instruments). A thermo plate was used to keep the recording chamber at 28°C

(Tokai hit).

Voltage imaging of the cerebellum

Embryos from 5 to 9 dpf were paralyzed with 0.02% tricaine and transferred to

an extracellular solution. Next, embryos were mounted on 2% low melting-point

agarose. For imaging, a fluorescence microscope (FN-1, Nikon) equipped with a

CMOS camera (ORCA-Flash4.0, Hamamatsu Photonics) was used with a

40×/0.8 NA water-immersion lens. Fluorescent images were acquired using NIS-

Elements or HCImage (Hamamatsu Photonics) at approximately 50–200 Hz.

Image analysis

Images were analyzed using the NIS-Elements (Nikon), Fiji, and OriginPro

(Lightstone) programs as described previously [52]. To extract changes in

fluorescence intensity, regions of interest (ROIs) were manually set using NIS-

element software (Nikon). Normalized changes in fluorescence intensity in the

target regions (ROIs) were quantitatively analyzed by calculating dF/F in the ROIs

(dF/F0=[F(t)–F0]/F0, where F(t) and F0 are the fluorescence intensities at a

certain time point and initial time points, respectively). If the larvae/embryos

moved during the voltage imaging, the images were aligned using a Fiji program

(TurboReg plugin). Denoising and bleach corrections were also applied to some

data using OriginPro (Lightstone). Cross-correlation and hierarchical clustering

analyses were performed using OriginPro (Lightstone).

Results

Establishment of the ArcLight zebrafish line

To examine neuronal activity in the developing brain using voltage imaging, GEVI ArcLight was used to establish transgenic zebrafish lines via the GAL4-UAS and Tol2 transposon systems [27]. We generated a pTol2-UAS-ArcLight-A242-polyA construct and established a transgenic line (Fig. 1a, b). To test whether ArcLight was localized to the cellular membranes in Tg(UAS:ArcLight) fish, we observed Tg(elavl3:GAL4-VP16;UAS:ArcLight;UAS:lyn-RFP) embryos, in which gal4 was expressed in neurons, and the cellular membranes were labeled with Lyn-RFP. At 1 dpf, ArcLight was widely observed in the neural tube and co-localized with Lyn-RFP, indicating that ArcLight was properly distributed on the cellular membrane in the neural tube (Fig. 1b, c). The distribution of ArcLight was also examined in other brain regions, such as the cerebellum, using transgenic lines expressing gal4 specifically in cerebellar neurons. Confocal imaging showed that ArcLight was clearly expressed in granule cells and inferior olive neurons, which are key components of cerebellar circuits [40] (Supplementary Figs. S1 and S2). These results indicate that ArcLight is expressed in various neuron types in developing zebrafish.

ArcLight could detect neural activity at the population and single-cell levels

Next, we performed high-speed imaging of the spinal cord of the Tg(elavl3:GAL4-VP16;UAS:ArcLight) embryos. Spinal cord neurons show spontaneous firing at the early stage of development, and this coordinated activity of the neuronal populations begins at 17 hpf [34, 38, 39]. As shown in Figure 1, oscillatory changes in ArcLight fluorescence intensity were detected in the ventral region of somites 3–8 at 20 hpf at a frequency of 0.14±0.10 Hz (6 fish, Fig. 1d–g). Phases of fluorescent changes differed between the left and right sides of the spinal cord and occurred alternately, which is consistent with previous reports showing

periodic depolarization (PD) of spinal cord neurons (Warp et al., 2012). This
activity mostly disappeared after treatment with the sodium channel blocker
tetrodotoxin (TTX) (Fig. 1e-g; left side: control 0.14±0.10 Hz, TTX 0.01±0.02 Hz,
p<0.05, right side: control 0.14±0.10 Hz, TTX 0.01±0.02 Hz, p<0.05, Wilcoxon
signed-rank test, 6 fish). These data indicated that ArcLight can detect
spontaneous activity in developing zebrafish spinal cord neurons.
Next, we examined whether ArcLight enables the detection of neural activity at
the single-cell level. Figure 2 shows an example of the voltage imaging of the
spinal cord. Oscillatory activity was clearly detected in each neuron, and
differences in the size and pattern of activity between neighboring neurons were
visualized using ArcLight (Fig. 2a and b, 35 fish). In addition to large
depolarizations that closely resembled PD, slight membrane potential fluctuations
were also observed between the PDs (Fig. 2c). Their timing and patterns suggest
that they represent a subthreshold activity called a synaptic burst, which has been
reported in electrophysiological recordings (Saint-Amant & Drapeau, 2000).
Cross-correlation analysis revealed a strong positive correlation between the
ipsilateral neurons and a negative correlation between the contralateral neurons
(Fig. 2d). This result indicates that membrane potential changes are highly
synchronized between neurons on the ipsilateral side of the spinal cord at the
single-cell level, and that they alternate between the left and right sides.
Furthermore, the hierarchical clustering analysis suggested that populations of
neurons on the same side exhibited higher synchrony (Fig. 2e).
249

High-spatiotemporal voltage imaging at the subcellular in the spinal cord
Measuring membrane potential dynamics at the level of neuronal compartments
such as axons and dendrites is important for understanding information
processing in the brain [53]. To record the cell bodies and axons of the spinal cord
neurons, Tg(elavl3:GAL4-VP16;UAS:ArcLight) embryos between 20 and 23 hpf
were observed from their side (Fig. 3a). We focused on primary motor neurons
(PMNs), as they are easy to observe because of their distinguished morphology
(Fig. 3b). As shown in Figure 3, we successfully performed wide-field voltage

imaging and detected the membrane potential in the soma and proximo/distal
regions of the axons (13 fish). The soma and axons showed similar patterns of
depolarization, but the size of the membrane potential fluctuations tended to be
smaller in the axons than in the cell soma (Fig. 3b). Furthermore, recording at a
higher spatial resolution enabled illumination of filopodia dynamics and voltage
changes (Supplementary Fig. S3 and S4). Interestingly, we noticed that there
were areas within the axon where the amplitude of the membrane potential
dynamics was larger than that of the others (Supplementary Fig. S4).
266

ArcLight mosaic labeling enables simultaneous single-cell observation of
morphology and neuronal activity
Spinal cord circuits consist of many types of neurons whose position, morphology,
and physiological properties vary (Saint-Amant & Drapeau, 2001). To uncover the
circuit dynamics and development of neural circuits, it is essential to observe the
morphology and voltage dynamics of each component. For this purpose, the
aforementioned Tg(elavl3:GAL4-VP16;UAS:ArcLight) line has a weakness in that
a large number of neighboring neurons express ArcLight, making it challenging
to observe the morphology of individual cells. Therefore, we obtained both
recordings by expressing ArcLight in a mosaic manner.
By injecting UAS-ArcLight pDNA into Tg(elavl3:GAL4-VP16) fish, we obtained
embryos in which spinal cord neurons were sparsely labeled with ArcLight,
enabling clear detection of the dynamics of membrane potentials (Fig. 4a–c, 20
hpf, PMNs). We also succeeded in recording interneurons that were otherwise
difficult to identify (Fig. 4d and e). In addition to PD and synaptic burst-like activity,
some neurons exhibited hyperpolarization (Fig. 4e). In one hour, these neurons
changed their morphology; cell 2 extended a long axon; however, it did not show
depolarization (Supplementary Fig. S5).
We also attempted to perform voltage imaging of cerebellar Purkinje cells,
which play pivotal roles in the cerebellum; however, GEVI imaging of such cells
has not yet been reported. By injecting ArcLight DNA, which drives the expression
of ArcLight specifically in Purkinje cells, we obtained embryos with sparse

labeling of Purkinje cells (Supplementary Fig. S7a and b). Voltage imaging at 5
dpf revealed a change in fluorescence intensity, indicative of depolarization in
some Purkinje cells (Supplementary Fig. S7b–d). However, significant bleaching
makes long-term imaging difficult, and the membrane potential cannot be
recorded from multiple Purkinje cells. These results indicate that voltage imaging
using ArcLight enables measurement of the voltage dynamics of neurons at
various levels (population, single cell, and subcellular), providing a powerful
approach for analyzing neural circuits.
297
Developmental changes in the membrane potentials of spinal neurons
Next, we examined the developmental changes in voltage dynamics in neuron
populations. The periodic firing observed in the spinal cord of zebrafish embryos
corresponds to coiling behavior, which is homologous to fetal movements in
mammals and is suggested to be essential for circuit formation [32]. During
embryogenesis, this characteristic activity appears at approximately 17 hpf and
rapidly increases in frequency and amplitude until 20 hpf, followed by a gradual
decrease. However, the mechanism by which the cooperative activity of this
population of spinal neurons emerges and changes during development remains
unclear. To investigate voltage dynamics in populations of spinal cord neurons,
we performed voltage imaging of zebrafish embryos during this developmental
period.
As shown in Figure 5, voltage imaging was successfully performed for the first
time from the developing spinal cord continuously from 18 to 20 hpf (Fig. 5). At
18 hpf, a PMN located in the eight somite region had an axonal structure but
seldom exhibited burst activity. Instead, it often showed fluctuations in the voltage
signal (Fig. 5c, cell 1). This neuron gradually extended its axons, increasing the
number of burst firings at 18.5 hpf. By 19.5 hpf, the firing frequency quickly
reached 0.15 Hz. In the same embryo, we also obtained a recording from the
PMN located in 10 somite region (Fig. 5c, cell 2). At 18 hpf, cell 2 exhibited a short
protrusion near its soma, suggesting that it was about to extend its axons.

Interestingly, cell 2 appeared to have almost no activity at 18 hpf but started to
show depolarization at 18.5 hpf. However, the pattern of membrane potential
fluctuation was quite different from that of mature PMNs, with many cells showing
a slow increase in depolarization. Additionally, small fluctuations in membrane
potential were continuously detected, suggesting that they were subthreshold
dynamics of membrane potentials in PMNs (Fig. 5c). By 19.5 hpf, cell2 extended
the axon, while the frequency of bursts rapidly increased to reach 0.15 Hz. This
tendency was also observed in the soma of cell 2 (Supplementary Fig. S6). A
similar pattern of development was observed in other embryos (5 fish, 20 cells).
These findings suggest that the coordinated activity in the spinal cord during the
early stage develops through multiple steps: (1) no activity, (2) small membrane
potential fluctuations and slow depolarization, and (3) oscillatory activity that
coincides with axonal growth.

Next, we examined how the spontaneous activity in the spinal cord changes
over time. Voltage imaging of a population of spinal cord neurons was performed
from 20 to 23 hpf at 30-minute intervals (Fig. 6; Supplementary Fig. S8). The
results showed that the regular and rapid population activity observed at 20 hpf
decreased by 23 hpf, and there was also a change in the pattern, with the bursts
becoming longer in duration (Fig. 6b). Furthermore, at 20 hpf, synchronized
activity within ipsilateral neurons appeared regularly between the left and right
sides; however, at 23 hpf, this regularity decreased, and several consecutive
bursts of activity were observed on the same side of the spinal cord (Fig. 6b).
Cross-correlation analysis revealed that at 20 hpf, both of ipsilateral and
contralateral pairs of neurons showed a high degree of synchrony and periodicity
in their activity (Fig. 6c). Again, they exhibited out-of-phase activity. In contrast,
at 23 hpf, the periodicity in the ipsilateral and contralateral pairs, as well as the
coordination between the left and right sides, seemed to be greatly reduced. In
contrast, the contralateral pairs showed a reduction in synchrony, while the
ipsilateral pairs maintained synchrony (Fig. 6c).

To confirm the stage differences in this activity pattern, the ratio of alternating
activity between neurons located on the left and right sides of the spinal cord was

compared at 20–21 hpf and 23–24 hpf. At 20–21 hpf, almost all neurons showed

alternating left-right activity, while at 23–24 hpf, this was significantly reduced to

nearly 80% (Fig. 6d; 20–21 hpf: 98.9±1.10%; 23–24 hpf: 98.9±1.10%, 5 fish; 23–

24 hpf: 79.8%±5.71%, 4 fish; $p<0.05$, Mann–Whitney U test). In contrast, all fish

observed showed synchronous activity in the ipsilateral neurons at both

developmental stages. These results suggest that in developing zebrafish spinal

neurons, as the periodic activity acquired by 20 hpf gradually changes and

disappears, left-right coordination is lost, but the correlation between ipsilateral

neurons remains.

Discussion

GEVI voltage imaging is useful as it enables the recording of membrane

potentials from many specific neuronal cell types noninvasively without using

electrodes. However, it has not been applied to studying neurogenesis. In this

study, we established experimental methods to enhance voltage imaging in the

zebrafish cerebellum and spinal cord. These are important not only for analyzing

neural circuits, but also for revealing developmental mechanisms.

Previously, electrophysiological recordings have been used to characterize

membrane dynamics in the spinal cord. However, neurons were observed for a

short period due to the invasive nature of the recordings. In addition, recordings

were limited to single cells [54, 55]. Moreover, developmental analysis was

performed using calcium imaging; however, in principle, it is impossible to record

membrane potential dynamics using this method [37-39]. To the best of our

knowledge, this is the first report of visualizing membrane potential dynamics in

genetically defined neuronal populations during spinal cord development.

Similarly, we believe that this is the first example of long-term GEVI-based

imaging of neural circuit formation during development.

Voltage imaging of spinal cord neural circuits

In this study, we successfully established a transgenic zebrafish line expressing

ArcLight. Initially, we focused on the spinal cord: one of the earliest circuits to
emerge during development, consisting of a relatively limited number of neurons
[34]. Voltage imaging at 20 hpf showed that the spinal neuronal population
exhibited regular, alternating left-right synchronous activity. The obtained patterns
were clear, with similar patterns of voltage changes to those recorded by
electrophysiology, but with higher resolution than those previously recorded using
ASAP1 or calcium imaging (Miyazawa et al., 2018; Muto et al., 2011; Wan et al.,
2019; Warp et al., 2012). This is probably because ArcLight has faster kinetics
than the calcium sensor, and a higher signal-to-noise ratio than ASAP1[12, 56].
Previously, electrophysiological studies showed that most of the spinal cord
neurons at the stage exhibit periodic bursts of activity called Periodic
depolarization (PD) that drives tail-coiling behaviors (equivalent to human fetal
movements). Similar patterns of burst firing, TTX dependence, suggesting that
Na channels are important, and similar firing frequency all support the notion that
the observed depolarization represents PD.
In addition to these suprathreshold activities, subthreshold activities, such as
SB and hyperpolarization, were also detected by ArcLight. SB, which cannot be
recorded using Ca imaging because it does not involve firing and is a small signal
change, was captured by optical recording for the first time in this study. In the
early developmental spinal cord, the contribution of chemical synapses is limited;
many neurons have been shown to be connected by gap junctions [54]. ArcLight
imaging might be able to visualize spatial information on electrical as well as
chemical coupling, which is useful for circuit analysis. While further analysis is
needed, including simultaneous electrophysiological recordings, this is the first
non-invasive imaging of voltage dynamics in a spinal cord neurons population at
their early stage of development.
408
High-resolution voltage imaging
Compartment-level analysis is important because neuronal function is regulated
at the compartmental level, such as for dendritic computation and axonal
compartments. ArcLight allowed recording separately from the cell body and axon

compartments of PMNs. Whilst most axons exhibited more activity proximally
than distally, for some neurons, we also detected larger activity in the distal area.
This may be due in part to the non-uniform distribution of sodium channels.

In addition, filopodia and lamellipodia were labeled by ArcLight, with filopodia
also showing voltage dynamics. Further research is needed to understand the
role of voltage changes in filopodia; for instance, in synapse formation and
plasticity. Further extension of this technique to ratiometry voltage analysis is
desirable for accurate recording [57]. The ability to simultaneously and
noninvasively observe cell morphology and activity allows for a deeper
understanding of network dynamics and their formation.

Emergence and changes in spinal cord networks during development

Long-term voltage imaging in the spinal cord showed that regular cooperative
activity arises in parallel with axon outgrowth. For instance, PMNs were found to
mature through multiple steps: (1) no firing, (2) small depolarization (initiation of
axon outgrowth), (3) irregular cooperative activity, and (4) cooperative activity.
This process proceeds in the same order in neurons distributed along the roster-
caudal axis. Although it is known that spinal cord neurons without axonal
outgrowth do not fire, this is the first time that stage-dependent changes have
been demonstrated for the same population [54].

Notably, PMNs exhibited small fluctuations in membrane potentials before
starting regular depolarization. Ion channels and gap junctions are key in driving
membrane potential in early-stage spinal cord neurons [34], where the
contribution of chemical synapses is limited, and many neurons are connected
by gap junctions. Further research is needed to determine whether the observed
small depolarization reflects, to some extent, spikelets. This could suggest that
small fluctuation may be due to neuronal electrical coupling, as electrical
synapses are a powerful trigger of synchronous activity [41, 58, 59]. Furthermore,
evidence is still sparse on important questions; whether pattern generators exist,
where they are located, and how they facilitate network activity [39]. Our
approach with in vivo voltage imaging could provide further information.

Voltage imaging in the cerebellum

In the cerebellum, cumulative information on circuit dynamics was collected by voltage-sensitive dye imaging [60-62]. To further understand circuit dynamics, optical recording of membrane potentials in a cell-type-specific manner is desirable. However, no study has clearly described cell-type-specific voltage imaging using GEVIs, despite several systems being available for neuron-type-specific expression [48]. In this study, we obtained strong ArcLight expression in cerebellar granule cells and inferior olive neurons in our transgenic fish. As for Purkinje cells, a central neuron in the cerebellum, we obtained ArcLight positive cells using transient expression with a new construct, since strong expression was not observed in Tg fish. Using voltage imaging, spontaneous depolarization was detected in Purkinje cells. We assumed that this depolarization represented a complex spike (CS), one of the key firing patterns of Purkinje cells, since the frequency was similar to that of CSs detected by cell-attached electrophysiological recordings at the same developmental stage in zebrafish [63]. However, we could not detect simple spikes (SSs) in Purkinje cells; this is partly because SSs exhibit smaller amplitude and higher frequencies, making them more difficult to target with voltage imaging [64, 65]. To improve the signal-to-noise ratio of the recordings, stronger drivers for increasing GEVI expression, more sensitive sensors, and an improved illumination/recording system are desirable [12, 19, 66]. Although further validation with electrophysiology is required, this work will provide an important step forward in understanding information processing and development in the cerebellum.

Regarding GEVI expression, we noticed considerable differences in intensity between neuron types, similar to other GEVIs such as ASAP1 [23]. To expand GEVI voltage imaging for the analysis of various neuron types, it is important to examine this difference in expression efficiency and to solve this issue.

In conclusion, this is the first report of simultaneous and non-invasive recordings of membrane potentials from neuronal populations during

development using a GEVI. This approach allowed us to cultivate dynamic
changes in their spatiotemporal patterns of activity. Our findings demonstrate that
this experimental system is an effective tool for elucidating the mechanisms
underlying neural circuit formation during development. Simple neuronal circuits
connected by gap junctions are one of the key regulators in animal behavior, as
shown in crab circuits [67]. Based on the imaging system we established for
zebrafish, we would be able to further understand how complex activity and
behaviors emerge from small populations of neurons by combining cell-type-
specific recordings with manipulations of neuronal activity through optical,
electrophysiological, and genetic approaches. Furthermore, in recent years,
bioelectric signaling has gradually been found to be important for various
biological phenomena, including morphogenesis; GEVI will be optimal to further
investigate this area [68, 69].
488

Acknowledgement

We thank Drs M. Hibi, K. Kawakami, S. Higashijima, H. Okamoto, H. Kakinuma, and R. Omo for providing fish lines and a constructs, and J. Nakai, M. Ohkura, K. Ando, K. Yamasu, L. Cohen, K. Yamasu, and Y. Bando for technical supports and helpful discussion. This work was supported in part by JSPS KAKENHI (Grants-in-Aid for Scientific Research), JST Tenure Track program at Saitama University (SUTT), Strategic Research Program for Brain Science from Ministry of Education, Culture, Sports, Science, and Technology of Japan (ASTEP), Kao Foundation grant, Asahi Glass Foundation, and Kato-bio Foundation.

Author Contributions

S.A. and S.T. conceived and designed the experiments. S.A, A.H., N.F., M.H., K.M., and S.T. performed the experiments. S.A., A.H., N.F., H.M, and S.T wrote the manuscript. All authors reviewed the manuscript.

Additional Information

Competing interests

The authors declare no competing interests.

Data Availability

All data generated or analyzed in this study are included in the manuscript and supporting files.

Reference

1. Cohen, L.B. and B.M. Salzberg, *Optical measurement of membrane potential.* Rev Physiol Biochem Pharmacol, 1978. **83**: p. 35-88.

2. Loew, L.M., et al., *Evidence for a charge-shift electrochromic mechanism in a probe of membrane potential.* Nature, 1979. **281**(5731): p. 497-9.

3. Djurisic, M., et al., *Optical monitoring of neural activity using voltage-sensitive dyes.* Methods Enzymol, 2003. **361**: p. 423-51.

4. Lin, M.Z. and M.J. Schnitzer, *Genetically encoded indicators of neuronal activity.* Nat Neurosci, 2016. **19**(9): p. 1142-53.

5. Homma, R., et al., *Wide-field and two-photon imaging of brain activity with voltage- and calcium-sensitive dyes.* Philos Trans R Soc Lond B Biol Sci, 2009. **364**(1529): p. 2453-67.

6. Ross, W.N., et al., *Changes in absorption, fluorescence, dichroism, and Birefringence in stained giant axons: : optical measurement of membrane potential.* J Membr Biol, 1977. **33**(1-2): p. 141-83.

7. Yan, P., et al., *Palette of fluorinated voltage-sensitive hemicyanine dyes.* Proc Natl Acad Sci U S A, 2012. **109**(50): p. 20443-8.

8. Wuskell, J.P., et al., *Synthesis, spectra, delivery and potentiometric responses of new styryl dyes with extended spectral ranges.* J Neurosci Methods, 2006. **151**(2): p. 200-15.

9. Momose-Sato, Y., T. Nakamori, and K. Sato, *Spontaneous depolarization wave in the mouse embryo: origin and large-scale propagation over the CNS identified with voltage-sensitive dye imaging.* Eur J Neurosci, 2012. **35**(8): p. 1230-41.

10. Storace, D., et al., *Toward Better Genetically Encoded Sensors of Membrane Potential.* Trends Neurosci, 2016. **39**(5): p. 277-289.

11. Kannan, M., G. Vasan, and V.A. Pieribone, *Optimizing Strategies for Developing Genetically Encoded Voltage Indicators.* Front Cell Neurosci, 2019. **13**: p. 53.

12. Bando, Y., et al., *Comparative Evaluation of Genetically Encoded Voltage Indicators.* Cell Rep, 2019. **26**(3): p. 802-813 e4.

13. Akemann, W., et al., *Imaging neural circuit dynamics with a voltage-sensitive fluorescent protein.* J Neurophysiol, 2012. **108**(8): p. 2323-37.

14. Jin, L., et al., *Single action potentials and subthreshold electrical events imaged in neurons with a fluorescent protein voltage probe.* Neuron, 2012. **75**(5): p. 779-85.

15. St-Pierre, F., et al., *High-fidelity optical reporting of neuronal electrical activity with*

an ultrafast fluorescent voltage sensor. Nat Neurosci, 2014. **17**(6): p. 884-9.

16.   Tsutsui, H., et al., *Improving membrane voltage measurements using FRET with*
new fluorescent proteins. Nat Methods, 2008. **5**(8): p. 683-5.

17.   Hochbaum, D.R., et al., *All-optical electrophysiology in mammalian neurons*
using engineered microbial rhodopsins. Nat Methods, 2014. **11**(8): p. 825-33.

18.   Gong, Y., et al., *High-speed recording of neural spikes in awake mice and flies*
with a fluorescent voltage sensor. Science, 2015. **350**(6266): p. 1361-6.

19.   Abdelfattah, A.S., et al., *Bright and photostable chemigenetic indicators for*
extended in vivo voltage imaging. Science, 2019. **365**(6454): p. 699-704.

20.   Piatkevich, K.D., et al., *A robotic multidimensional directed evolution approach*
applied to fluorescent voltage reporters. Nat Chem Biol, 2018. **14**(4): p. 352-360.

21.   Nakajima, R. and B.J. Baker, *Mapping of excitatory and inhibitory postsynaptic*
potentials of neuronal populations in hippocampal slices using the GEVI, ArcLight.
J Phys D Appl Phys, 2018. **51**(50).

22.   Storace, D.A. and L.B. Cohen, *Measuring the olfactory bulb input-output*
transformation reveals a contribution to the perception of odorant concentration
invariance. Nat Commun, 2017. **8**(1): p. 81.

23.   Miyazawa, H., et al., *Optical interrogation of neuronal circuitry in zebrafish using*
genetically encoded voltage indicators. Sci Rep, 2018. **8**(1): p. 6048.

24.   Bohm, U.L., et al., *Voltage imaging identifies spinal circuits that modulate*
locomotor adaptation in zebrafish. Neuron, 2022. **110**(7): p. 1211-1222 e4.

25.   Yang, H.H., et al., *Subcellular Imaging of Voltage and Calcium Signals Reveals*
Neural Processing In Vivo. Cell, 2016. **166**(1): p. 245-57.

26.   Ahrens, M.B., et al., *Brain-wide neuronal dynamics during motor adaptation in*
zebrafish. Nature, 2012. **485**(7399): p. 471-7.

27.   Asakawa, K., et al., *Genetic dissection of neural circuits by Tol2 transposon-*
mediated Gal4 gene and enhancer trapping in zebrafish. Proc Natl Acad Sci U S
A, 2008. **105**(4): p. 1255-60.

28.   Yoshimatsu, T., et al., *Ancestral circuits for vertebrate color vision emerge at the*
first retinal synapse. Sci Adv, 2021. **7**(42): p. eabj6815.

29.   Hashimoto, M. and M. Hibi, *Development and evolution of cerebellar neural*
circuits. Dev Growth Differ, 2012. **54**(3): p. 373-89.

30.   McLean, D.L., et al., *A topographic map of recruitment in spinal cord.* Nature,
2007. **446**(7131): p. 71-5.

31.   Grillner, S., *Neurobiological bases of rhythmic motor acts in vertebrates.* Science,
1985. **228**(4696): p. 143-9.

32.   Menelaou, E., et al., *Embryonic motor activity and implications for regulating*
587   *motoneuron axonal pathfinding in zebrafish.* Eur J Neurosci, 2008. **28**(6): p. 1080-
588   96.

589   33.   Fidelin, K. and C. Wyart, *Inhibition and motor control in the developing zebrafish*
590   *spinal cord.* Curr Opin Neurobiol, 2014. **26C**: p. 103-109.

591   34.   Saint-Amant, L., *Development of motor rhythms in zebrafish embryos.* Prog Brain
592   Res, 2010. **187**: p. 47-61.

593   35.   O'Donovan, M.J., *The origin of spontaneous activity in developing networks of*
594   *the vertebrate nervous system.* Curr Opin Neurobiol, 1999. **9**(1): p. 94-104.

595   36.   Saint-Amant, L. and P. Drapeau, *Motoneuron activity patterns related to the*
596   *earliest behavior of the zebrafish embryo.* J Neurosci, 2000. **20**(11): p. 3964-72.

597   37.   Muto, A., et al., *Genetic visualization with an improved GCaMP calcium indicator*
598   *reveals spatiotemporal activation of the spinal motor neurons in zebrafish.* Proc
599   Natl Acad Sci U S A, 2011. **108**(13): p. 5425-30.

600   38.   Warp, E., et al., *Emergence of patterned activity in the developing zebrafish*
601   *spinal cord.* Curr Biol, 2012. **22**(2): p. 93-102.

602   39.   Wan, Y., et al., *Single-Cell Reconstruction of Emerging Population Activity in an*
603   *Entire Developing Circuit.* Cell, 2019. **179**(2): p. 355-372 e23.

604   40.   Ito, M., *Cerebellar circuitry as a neuronal machine.* Prog Neurobiol, 2006. **78**(3-
605   5): p. 272-303.

606   41.   Mann-Metzer, P. and Y. Yarom, *Electrotonic coupling interacts with intrinsic*
607   *properties to generate synchronized activity in cerebellar networks of inhibitory*
608   *interneurons.* J Neurosci, 1999. **19**(9): p. 3298-306.

609   42.   Cohen, D. and Y. Yarom, *Patches of synchronized activity in the cerebellar cortex*
610   *evoked by mossy-fiber stimulation: questioning the role of parallel fibers.* Proc
611   Natl Acad Sci U S A, 1998. **95**(25): p. 15032-6.

612   43.   Whittington, M.A., R.D. Traub, and J.G. Jefferys, *Synchronized oscillations in*
613   *interneuron networks driven by metabotropic glutamate receptor activation.*
614   Nature, 1995. **373**(6515): p. 612-5.

615   44.   Ito, M., *Cerebellar long-term depression.* Trends Neurosci, 1996. **19**(1): p. 11-12.

616   45.   Apps, R. and R. Hawkes, *Cerebellar cortical organization: a one-map hypothesis.*
617   Nat Rev Neurosci, 2009. **10**(9): p. 670-81.

618   46.   Tsuda, S., et al., *Probing the function of neuronal populations: combining*
619   *micromirror-based optogenetic photostimulation with voltage-sensitive dye*
620   *imaging.* Neurosci Res, 2013. **75**(1): p. 76-81.

47.   Kimura, Y., C. Satou, and S. Higashijima, *V2a and V2b neurons are generated*

by the final divisions of pair-producing progenitors in the zebrafish spinal cord.
Development, 2008. **135**(18): p. 3001-5.

48. Takeuchi, M., et al., *Establishment of Gal4 transgenic zebrafish lines for analysis*
of development of cerebellar neural circuitry. Dev Biol, 2015. **397**(1): p. 1-17.

49. Lister, J.A., et al., *nacre encodes a zebrafish microphthalmia-related protein that*
regulates neural-crest-derived pigment cell fate. Development, 1999. **126**(17): p.
3757-67.

50. Nakayama, Y., et al., *Comprehensive analysis of target genes in zebrafish*
embryos reveals gbx2 involvement in neurogenesis. Dev Biol, 2017. **430**(1): p.
237-248.

51. Kimura, Y., et al., *Hindbrain V2a neurons in the excitation of spinal locomotor*
circuits during zebrafish swimming. Curr Biol, 2013. **23**(10): p. 843-9.

52. Hiyoshi, K., et al., *In vivo wide-field voltage imaging in zebrafish with voltage-*
sensitive dye and genetically encoded voltage indicator. Dev Growth Differ, 2021.
63(8): p. 417-428.

53. London, M. and M. Hausser, *Dendritic computation.* Annu Rev Neurosci, 2005.
28: p. 503-32.

54. Saint-Amant, L. and P. Drapeau, *Synchronization of an embryonic network of*
identified spinal interneurons solely by electrical coupling. Neuron, 2001. **31**(6):
p. 1035-46.

55. Saint-Amant, L. and P. Drapeau, *Time course of the development of motor*
behaviors in the zebrafish embryo. J Neurobiol, 1998. **37**(4): p. 622-32.

56. Storace, D.A., et al., *Monitoring brain activity with protein voltage and calcium*
sensors. Sci Rep, 2015. **5**: p. 10212.

57. Kim, B.B., et al., *A red fluorescent protein with improved monomericity enables*
ratiometric voltage imaging with ASAP3. Sci Rep, 2022. **12**(1): p. 3678.

58. Epsztein, J., et al., *Impact of spikelets on hippocampal CA1 pyramidal cell activity*
during spatial exploration. Science, 2010. **327**(5964): p. 474-7.

59. de Solages, C., et al., *High-frequency organization and synchrony of activity in*
the purkinje cell layer of the cerebellum. Neuron, 2008. **58**(5): p. 775-88.

60. Roome, C.J. and B. Kuhn, *Simultaneous dendritic voltage and calcium imaging*
and somatic recording from Purkinje neurons in awake mice. Nat Commun, 2018.
9(1): p. 3388.

61. Rokni, D., R. Llinas, and Y. Yarom, *Stars and stripes in the cerebellar cortex: a*
voltage sensitive dye study. Front Syst Neurosci, 2007. **1**: p. 1.

62. Okumura, K., et al., *Optical measurement of neuronal activity in the developing*

cerebellum of zebrafish using voltage-sensitive dye imaging. Neuroreport, 2018.

29(16): p. 1349-1354.

63.   Hsieh, J.Y., et al., *Rapid development of Purkinje cell excitability, functional*

cerebellar circuit, and afferent sensory input to cerebellum in zebrafish. Front

Neural Circuits, 2014. **8**: p. 147.

64.   Knogler, L.D., A.M. Kist, and R. Portugues, *Motor context dominates output from*

purkinje cell functional regions during reflexive visuomotor behaviours. Elife,

2019. **8**.

65.   Harmon, T.C., et al., *Distinct responses of Purkinje neurons and roles of simple*

spikes during associative motor learning in larval zebrafish. Elife, 2017. **6**.

66.   Namikawa, K., et al., *Modeling Neurodegenerative Spinocerebellar Ataxia Type*

13 in Zebrafish Using a Purkinje Neuron Specific Tunable Coexpression System.

J Neurosci, 2019. **39**(20): p. 3948-3969.

67.   Gutierrez, G.J., T. O'Leary, and E. Marder, *Multiple mechanisms switch an*

electrically coupled, synaptically inhibited neuron between competing rhythmic

oscillators. Neuron, 2013. **77**(5): p. 845-58.

68.   Levin, M., *Bioelectric signaling: Reprogrammable circuits underlying*

embryogenesis, regeneration, and cancer. Cell, 2021. **184**(8): p. 1971-1989.

69.   Harris, M.P., *Bioelectric signaling as a unique regulator of development and*

regeneration. Development, 2021. **148**(10).

 Figure Legends

Figure 1

Transgenic zebrafish showing membrane-localized ArcLight and spontaneous

activity in the neural tube. (a) Schematic diagram of plasmid construct for ArcLight.

(b) Lateral views of ArcLight-expressing embryos at 1 dpf. ArcLight is distributed

widely in the neural tube of Tg(elavl3:GAL4-VP16;UAS:ArcLight) fish. (c) Dorsal

view of the neural tube of Tg(elavl3:GAL4-VP16;UAS:ArcLight;UAS:lyn-RFP)

embryos at 1 dpf. ArcLight (green) is co-localized with Lyn-RFP (red: cell

membranes) in the neural tube (arrowheads). Higher magnification images are

shown in the lower panel. (d-f) Spontaneous activity of spinal cord neurons is

detected by ArcLight imaging. (d) Dorsal view of the ventral spinal cord of Tg

(elavl3:GAL4-VP16;UAS:ArcLight-A242) fish at 20 hpf. The rostral side is to the

left, and the area between 3–8 somites is shown. A schematic diagram of the

imaging system is shown on the right. Changes in fluorescence of ArcLight

($-\Delta F/F_0$) are indicated by the pseudocolor scale shown at right. White

arrowheads indicate the activated neurons. Regions of interest (ROIs) located

between 5–7 somites are indicated by red (right side) and blue (left side)

rectangular in (d). Fluorescence changes of ArcLight in the ROIs are shown in

the right. The time point of the image is indicated by a black arrowhead. (f-g) The

spontaneous activity in the spinal cord mostly disappeared by tetrodotoxin (TTX)

treatment (6 fish, *: $p < 0.05$). The imaged plane and ROIs are in the same positions

as in (d).

Figure 2

Monitoring of spontaneous activity of individual cells in the spinal cord by ArcLight

imaging. (a) A fluorescence image of the ventral spinal cord of Tg(elavl3:GAL4-

VP16;UAS:ArcLight) fish at 21 hpf. ROIs are located at the 18 cells (red: right

side, blue: left side). Higher magnification views are shown on the right. (b)

Changes in ArcLight signals fluorescence of ArcLight at the two time points

indicated in (c). (c) Activity patterns of the 18 cells are shown. A higher

magnification view of the ArcLight signal of cell 4 is shown at the bottom. (b). (d,

e) Hierarchal clustering analysis of the voltage dynamics in the 18 cells shows

the synchronous activity in the ipsilateral neuron pairs (d), and also several

groups with higher synchrony within them.

Figure 3

Voltage imaging of spinal cord neurons at a high spatiotemporal resolution at the

cellular compartment level by ArcLight. (a) Lateral view of the spinal cord of

Tg(elavl3:GAL4-VP16;UAS:ArcLight) fish at 20 hpf. (b) Changes in ArcLight

signal are observed at soma and axons in three different ROIs shown in (a).

Higher resolution observations are shown in Supplementary Fig. S3 and S4.

Figure 4

Single cell labeling of neurons by ArcLight enables simultaneous recording of

voltage and morphology. (a) Schematic diagram of the co-injection experiment.

(b) A stacked confocal image of the spinal cord region of the injected embryo at

20 hpf. The left side view is shown. Arrowheads and arrows indicate the soma

and axons of ArcLight positive cells, respectively. (c) Voltage dynamics of a

primary motor neuron are indicated by a yellow arrowhead in (b). ROIs at soma

and axons are shown in red and blue, respectively. (d, e) Voltage imaging of the

spinal cord interneurons. A higher magnification view is shown in e. ArcLight

signals from three cells are shown on the right. In addition to periodic

depolarization, hyperpolarization (white arrowheads) and subthreshold-like

signals (black arrowheads) are observed.

Figure 5

Long-term voltage imaging of the spinal cord shows the emergence of population

activity. (a) Left views of the spinal cord of ArcLight positive fish from 18 hpf to

19.5 hpf. (b) Enlarged views of (a). (c) Voltage dynamics obtained from the axons

of the two cells shown in (a). Arrowheads indicate small depolarization.

Figure 6

Changes in firing properties in spinal cord neuron populations. (a) Dorsal view of

ArcLight positive spinal cord at 20 hpf. (b) Voltage dynamics of each neuron are

shown at 20 and 23 hpf. The position of ROIs is indicated in (a). (c) Cross-

correlation analysis of ipsilateral and contralateral pairs of spinal cord neurons at

20 and 23 hpf. (d) Probability of altering activity at 20-21 hpf and 23-24 hpf.

Supplementary Figure S1

ArcLight (green) is distributed in cerebellar granule cells (red). Dorsal view of the

cerebellum of Tg(hspGFFDMC152C;UAS:RFP;UAS:ArcLight) fish observed at 6

dpf. Higher magnification views are shown in the lower panel.

Supplementary Figure S2

ArcLight is expressed in inferior olivary neurons. (a) Dorsal view of the inferior

olive (IO) of Tg(hspGFFDMC28C;UAS:RFP;UAS:ArcLight) fish at 8 dpf.

Arrowheads in the cerebellum (CE) indicate ArcLight signals (green) in the

climbing fibers. (b) (left) Schematic diagram of the co-injection experiment. (right)

Transient expression of ArcLight in a single IO neuron. Higher magnification

views are shown in the right panel.

Supplementary Figure S3

Confocal observation of ArcLight expression and voltage dynamics in the spinal

cord. (a) Lateral views of Tg(elavl3:GAL4-VP16;UAS:ArcLight) embryo at 1 dpf.

Higher magnification images shown in the right indicate filopodia and lamerioidia

of primary motor neurons (arrowheads). (b) Voltage imaging of cell1 in (a). ROIs

were set at the axonal region. ArcLight detected depolarization at axons (black

arrowheads). The gray area is the period when the embryos moved.

Supplementary Figure S4

Voltage imaging of a primary motor neuron at higher spatial resolution. (a) Left views of the spinal cord at two different time points. Arrowheads indicate filopodia (white, yellow). A yellow arrowhead indicates a filopodium that shows a prominent extension. (b, c) An example of voltage dynamics detected at various positions in the axon. ROI4-6 were set on a filopodium. ROI 7 was set in a region where no clear structure was observed.

Supplementary Figure S5

Single-cell labeling by ArcLight shows the morphological changes of the spinal cord neurons. Images taken at 21 hpf and 22 hpf are shown. An arrowhead indicates a growth cone of the extending axon.

Supplementary Figure S6

ArcLight imaging in Purkinje cells. (a) (top) Schematic diagram ofpTo2-aldoca-ArcLightQ239-polyA construct. (bottom) ArcLight is expressed in Purkinje cells in the cerebellum (CE). Arrowheads and an arrow indicate Purkinje cell soma and dendrite, respectively. A higher magnification view is shown on the right. (b-d) ArcLight imaging exhibits depolarization in Purkinje cells. (b) ArcLight signal in the recorded Purkinje cells. (c) Depolarization signals are observed by ArcLight imaging (arrowheads). (d) Enlarged views of the traces are shown in (c).

Supplementary Figure S7

ArcLight imaging in Purkinje cells. (a) (top) Schematic diagram ofpTo2-aldoca-ArcLightQ239-polyA construct. (bottom) ArcLight is expressed in Purkinje cells in the cerebellum (CE). Arrowheads and an arrow indicate Purkinje cell soma and dendrite, respectively. A higher magnification view is shown on the right. (b-d)

804 ArcLight imaging exhibits depolarization in Purkinje cells. (b) ArcLight signal in
805 the recorded Purkinje cells. (c) Depolarization signals are observed by ArcLight
806 imaging (arrowheads). (d) Enlarged views of the traces are shown in (c).
807
808
809 **Supplementary Figure S7**
810 Long-term voltage imaging shows the changes in the firing patterns of the spinal
811 cord neurons population. (a) ROIs position of cells 1-6. (b) Voltage dynamics of
812 6 cells are shown from 20 to 23 hpf. Cells on the right and left sides of the spinal
813 cord are shown in red and blue, respectively.
814
815
816 **Supplementary Movie1**
817 Spontaneous activity of individual cells in the spinal cord is observed by ArcLight.
818 Dorsal view of the ventral spinal cord of Tg(elavl3:GAL4-VP16;UAS:ArcLight) fish
819 at 21 hpf.
820
821
822 **Supplementary Movie2**
823 Monitoring of spontaneous activity of spinal cord neurons, including primary
824 motor neurons by ArcLight. Left view of the spinal cord of Tg(elavl3:GAL4-
825 VP16;UAS:ArcLight) fish at 20 hpf.
826
827
828 **Supplementary Movie3**
829 Single-cell labeling and voltage imaging of spinal cord neurons using ArcLight.
830 Left view of the spinal cord of Tg(elavl3:GAL4-VP16) embryo at 21 hpf. A high-
831 resolution image is shown in Supplementary Figure S5.
832
833
834 **Supplementary Movie4**

835 Voltage dynamics of an axonal region of a primary motor neuron at a high spatial
836 resolution. Left view of the spinal cord Tg(elavl3:GAL4-VP16;UAS:ArcLight) fish
837 at 18.5 hpf.
838

Figure 1

a pTol2-UAS-ArcLight-A242-polyA

Tol2 UAS ArcLight-A242 pA Tol2

b

300 μm

c ArcLight   Lyn-RFP   Merge

30 μm

10 μm

d ArcLight

50 μm

lens

e -ΔF/F₀   control

f TTX

right

left

right

left

-ΔF/F₀
0.5 %

5 s

g

Figure 2

Figure 3

Figure 4

a

transposase mRNA
UAS-ArcLight pDNA

Tg(elavl3:GAL4)

b

50 µm

c

ROI1

10 µm

ROI 1

-ΔF/F$_0$
2%

5 s

d

ArcLight

50 µm

-ΔF/F$_0$

11.0

-ΔF/F$_0$ (%)

-0.5

e

cell 1

10 µm

cell 1

-ΔF/F$_0$
2%

5 s

Figure 5

a

ArcLight

18 hpf   Cell 1   2

18.5 hpf

19 hpf

19.5 hpf   50 µm

b

Cell 1   18 hpf   18.5 hpf   19 hpf   19.5 hpf

Cell 2   18 hpf   18.5 hpf   19 hpf   19.5 hpf

c

Cell 1   18 hpf   18.5 hpf   19 hpf   19.5 hpf

Cell 2   18 hpf   18.5 hpf   19 hpf   19.5 hpf

$-\Delta F/F_0$ 2%

5 s

Figure 6

a

cell 1   2   3   ArcLight

4   5   6   50 μm

b

20 hpf   23 hpf

Cell number

right   1, 2, 3

left   4, 5, 6

$-\Delta F/F_0$
3 %

5 s

c

20 hpf   — ipsilateral   — contralateral

23 hpf

Cross-correlation coefficient

Time (s)

cross-correlation coefficient

Time (s)

d

Probability of alternating activity (%)

*

20-21   23-24

Developmental stage (hpf)

Supplementary Figure S1

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4

Supplementary Figure S5

Supplementary Figure S6

Supplementary Figure S7

a

20 hpf
ArcLight

cell 1   2   3

4   5   6   50 µm

b

20 hpf

cell number

right
1
2
3

left
4
5
6

21 hpf

cell number

right
1
2
3

left
4
5
6

22 hpf

cell number

right
1
2
3

left
4
5
6

23 hpf

cell number

right
1
2
3

left
4
5
6

-ΔF/F$_0$
3 %

5 s

Supplementary Figure S8