1 # Designing patient-oriented combination therapies for acute myeloid
2 # leukemia based on efficacy/toxicity integration and bipartite network
3 # modeling

4 **Mehdi Mirzaie**[a,†]**, Elham Gholizadeh**[a,†]**, Juho J. Miettinen**[b]**, Filipp Ianevski**[b]**, Tanja Ruokoranta**[b]**, Jani Saarela**[b]**,
5 **MikkoManninen**[c]**, Susanna Miettinen**[d]**, Caroline A. Heckman**[e,*]**, and Mohieddin Jafari**[a,*]

6
[a]Department of Biochemistry and Developmental Biology, University of Helsinki, Helsinki, Finland; [b]Institute for Molecular Medicine Finland, Helsinki Institute of
Life Science,University of Helsinki, Helsinki, Finland; [c]Orton Orthopaedic Hospital, Helsinki, Finland; [d]Adult Stem Cell Group, Faculty of Medicine and Health
Technology, Tampere University, Tampere, Finland; [e]Institute for Molecular Medicine Finland - FIMM, HiLIFE - Helsinki Institute of Life Science, iCAN Digital
Precision Cancer Medicine Flagship, University of Helsinki, Helsinki, Finland

Key Points

- Ruxolitinib-ulixertinib and sapanisertib-LY3009120 have the best synergistic effects
on AML, with the least toxicity.
- This study's combinations destroy blasts without harming other healthy cells, unlike
standard chemotherapy, which is less specific.

Abstract

Acute myeloid leukemia (AML), a heterogeneous and aggressive blood cancer, does not
respond well to single-drug therapy. A combination of drugs is required to effectively treat
this disease. Computational models are critical for combination therapy discovery due to the
tens of thousands of two-drug combinations, even with approved drugs. While predicting
synergistic drugs is the focus of current methods, few consider drug efficacy and potential
toxicity, which are crucial for treatment success. To find effective new drug candidates, we
constructed a bipartite network using patient-derived tumor samples and drugs. The network
is based on drug-response screening and summarizes all treatment response heterogeneity as
drug response weights. This bipartite network is then projected onto the drug part, resulting in
the drug similarity network. Distinct drug clusters were identified using community detection
methods, each targeting different biological processes and pathways as revealed by
enrichment and pathway analysis of the drugs' protein targets. Four drugs with the highest
efficacy and lowest toxicity from each cluster were selected and tested for drug sensitivity
using cell viability assays on various samples. Results show that the combinations of
ruxolitinib-ulixertinib and sapanisertib-LY3009120 are the most effective with the least
toxicity and best synergistic effects on blasts. These findings lay the foundation for
personalized and successful AML therapies, ultimately leading to the development of drug
combinations that can be used alongside standard first-line AML treatment.

Keywords: Drug combination design; Acute myeloid leukemia (AML); Nominal data mining;
Bipartite network modeling; High-throughput drug screening; drug efficacy; drug toxicity

Introduction

Acute myeloid leukemia (AML) is an inter- and intra-tumor heterogeneous disease [1,2]. It is identified when the bone marrow (BM) contains at least 20% of blast cells of the myeloid lineage [3]. Traditional chemotherapeutics have limited efficacy in patients over the age of 65, with a survival rate of less than 25% at one year follow-up and less than 9% after 5 years [4]. Despite recent advances in genome sequencing, which enables researchers to identify a large number of mutations, we are still hampered by the absence of drugs which specifically target cancer-associated mutations [5]. On the other hand, the majority of AML patients do not have actionable mutations, and the link between cancer genotype, phenotype, and therapeutic function is poorly understood [6]. Even if we overcome the above difficulties and identify the exact mutations in genotype, monotherapy drug resistance will remain a major clinical complication [7]. Targeted anti-cancer compounds used in combination therapy have the potential to overcome resistance, improve patient response to current treatments, reduce dose-limiting single agent toxicity, and broaden the spectrum of available therapies [8].

Drug combination therapy offers the chance to suppress a number of pathways synergistically, including patient-specific cancer rescue pathways and phenotypic redundancy across heterogeneous cancer sub-clones [9]. The phenotypic effects of thousands of drug combinations can be evaluated in patient-derived cells and other pre-clinical model systems using high-throughput screening. However, because there are so many possible drug and dose combinations, large-scale multi-dose combinatorial screening is not recommended, due to the limited number of cells available from patient samples. Using the presented method in this study, researchers would be able to categorize the most important AML drugs into different clusters, each of which target proteins associated with various signaling pathways.

In our earlier research, we designed a systems pharmacology approach based on network

modeling to identify prospective drug combinations in AML [10]. To gain a deeper

understanding of the factors that govern drug response in AML patients, we utilized a unique

and extensive dataset obtained through drug response screening of samples from both AML

patients and healthy donors in Finland as part of our study. Our model accounts for the

efficacy and toxicity of drug response, which are simultaneously evaluated on patient and

healthy samples, respectively [11]. A weighted bipartite network composed of two parts,

chemical components, and patient samples was built to develop a drug combination strategy

using the screening outcomes of single drug responses on AML patient samples. This enables

researchers to directly access the phenotype of the patients' cancer cells through *ex vivo* drug

response data, and by using network modeling and clustering analysis, demonstrate the drugs

functionalities. Next, top drug combinations can be predicted based on phenotypic

responses of samples in each cluster. In addition, we used two different computational

resources, i.e., molecular biology annotations, and the chemical structure of drugs, to perform

intra-cluster homogeneity analysis. The top combinations for further study in current pre-

clinical drug combination screening are frequently chosen based on synergy alone, without

taking into account the combination efficacy and toxicity effects, despite the fact that these

are critical determinants of a therapy's clinical success [12].

Considering the importance of toxicity, in this study, we investigated the drug response

of AML patient and healthy donor samples to calculate both efficacy and toxicity,

respectively. The efficacy and toxicity of the drugs was assessed *ex vivo* on AML patient

samples using cell viability and flow cytometry assays. Since clinical symptoms in patients

are caused by blast cell accumulation in bone marrow [13], based on the assessment of CD

marker expressions, we suggested the most blast-specific combinations as promising

combinations for AML treatment. The novel combinations indicated in this study are effective on

blasts but have a reduced level of toxicity on other cell populations, whereas with standard

chemotherapy the majority of the blasts are killed together with other cell types.

Results

The entire workflow of this study is depicted in Figure 1 and detailed material and method

is presented in the supplementary information. The drug responses of 525 chemical

compound tested on 199 AML patients' bone marrow samples were obtained from the

FIMM AML data set[11]. The bipartite network was constructed using this data set, as

explained in the materials and methods section in the supplement. A bipartite network can

be projected into two different types of unipartite networks containing nodes of only one

type. The projection of the bipartite network, onto the "drug" node set is considered here,

called the drug similarity network. The Louvain community detection approach was used to

find drugs that behaved similarly in terms of drug response[14]. The results gave us two

communities (clusters) of drugs denoted by C_1 and C_2 with network sizes of 155 and 141,

respectively (Table S1).

Comparing AML drug clusters: Evaluating protein target pathways and chemical
structure similarity

We used two independent computational methods to determine how distinct the two

clusters are: the first identifies the significant difference between biological pathways of

drug protein targets in each cluster, and the second evaluates the chemical structure

similarity of drugs in each cluster. We constructed a drug-target network using the drug

target commons (DTC) database[15], which is also a bipartite network in which each link

connects drugs to their protein targets. Let T_1 and T_2 represent the set of protein targets of

drugs in the cluster C_1 and C_2, respectively, and T represents the union of T_1 and T_2. In this

study $|T_1| = 921, |T_2| = 842, and |T| = 1055.$

Proteins with a score S (explained in the supplementary information 1.3) greater than log

(2) are considered to be preferentially targeted by drugs in cluster 1, denoted by PPT1.

Similarly, PPT2 proteins have a score of less than $log(0.50)$. We performed GSEA (gene

set enrichment analysis) on PPT1 and PPT2 proteins based on their associated scoring

functions. As expected, the biological processes and signaling pathways affected by drugs

in Clusters 1 and 2 are distinct. This difference enables us to inhibit two different signaling

pathways using one combination. Drugs in cluster 1 (PPT1), such as LY3009120 (a Pan-

RAF inhibitor), predominantly target proteins associated with the RAF-MEK-ERK

signaling pathway. This pathway plays a crucial role in cell proliferation and growth,

indirectly influencing processes like cell-substrate adhesion and ion trans-membrane

transport, which are enriched in our analysis [16]. In contrast, JAK1/2 inhibitors like

ruxolitinib target JAK proteins, involved in cytokine signaling and immune responses,

impacting pathways related to neuroactive ligand-receptor interactions and the regulation

of actin cytoskeleton [17]. Drugs like birabresib, which target proteins in the bromodomain

and extra-terminal (BET) family, have a role in gene regulation through chromatin binding,

affecting gene expression and pathways related to chemical reactions and collagen

metabolism [18]. Plicamycin, which binds to guanine-cytosine-rich regions of DNA, may

influence gene expression and regulation, impacting pathways related to collagen

metabolism and other DNA-dependent processes (Figure 2 A) [19]. On the other hand,

proteins targeted by drugs in cluster 2 (PPT2) (silmitasertib, ulixertinib, sapanisertib, and

teniposide) are in the p53 signaling pathway, cell cycle, apoptosis, and pancreatic,

colorectal and chronic myeloid leukemia cancers and related to tumorigenesis and

progression pathways, including human immunodeficiency virus 1 infection [20-23].

We also performed ShinyGO [24] Gene Ontology and KEGG pathway enrichment analysis

on proteins that are merely targeted by drugs in one cluster. For this purpose, two protein

sets G1 and G2 were selected such that G1 includes proteins targeting by at least three

drugs in cluster 1 and at most two drugs in cluster 2, and similarly, G2, consist of

proteins that are mostly targeted by drugs in cluster 2 (at least three drugs in cluster 2

and at most two drugs in cluster 1). REVIGO was also used to summarize the enriched

GO terms, and the results are shown in Figure 2 and Tables S2 and S3. The cAMP

signaling pathway, lipids and atherosclerosis, steroid hormone biosynthesis, and rhythmic

processes and circadian rhythm are biological processes related to G1 proteins, which are

mostly targeted by LY3009120, birabresib, plicamycin, and ruxolitinib. Cell cycle, cellular

senescence, T-cell leukemia virus 1 infection and cell division, mitotic cell cycle, and

protein phosphorylation processes are related to G2 proteins, mostly targeted by

silmitasertib, ulixertinib, sapanisertib, and teniposide. Therefore, we demonstrate that the

protein targets of drugs in each cluster are on different pathways and biological processes.

To do homogeneity analysis of chemical structure of drugs, the dice similarity test was

used to show how structurally similar the drugs are in each cluster. This measurement

compares the number of chemical features shared by a pair of compounds to the average

size of the total number of features present. Pairwise similarities were calculated for

chemical compounds chosen from two drug clusters for inter-cluster comparison. Drugs

from different clusters are less similar than drugs from the same cluster, as shown in the

Figure 3A. According to the box plot, the inter-cluster similarities are less than the intra-

cluster similarities in both clusters. The results of the t-test imply that the mean of inter-

cluster similarities is less than the mean of intra-cluster similarities in clusters 1 and 2 (p-

value < 2.2e-16 for both t-test).

Combination selection: Balancing toxicity and efficacy across clusters

As a result, we demonstrated that clusters are well-separated and that the protein targets of

drugs in each cluster are involved in distinct pathways. In this novel combination strategy,

we aim to select two drugs from distinct clusters while taking both toxicity and efficacy

into account. The optimal combinations are those that have lower toxicity than the average

toxicity and higher efficacy values than the average efficacy value for all drugs. For each

drug, the average drug response of healthy and AML patient samples in the data set are

considered as toxicity and efficacy, respectively. We assume that the ideal drugs have no

inhibitory effect on healthy samples but significantly influence blast cells in AML patient

samples. We chose the top 5% of drugs whose toxicity is less than the average of all drug

toxicity and efficacy is greater than the average of all drug efficacy. Figure 3B depicts the

link between toxicity and efficacy values of 296 drugs on 81 samples. The top selected

small molecule in each cluster is summarized in Table 1 and Table S4. Four chemical

compounds from cluster 1 including birabresib, LY3009120, plicamycin, and ruxolitinib as

well as four drugs from cluster 2 including sapanisertib, silmitasertib, teniposide, and

ulixertinib were chosen for drug combination testing. According to our experimental

design, the combination of drugs within a single cluster is known as negative group or

intra-cluster, and the combination of drugs between clusters is considered as positive

group or inter-cluster.

Enhanced efficacy and reduced toxicity in inter-cluster drug combinations on AML
177
patient samples unveiled by cell viability drug screening

In the testing of all 16 inter-cluster and 12 intra-cluster combinations at five different

concentrations, the cell viability of sixteen AML patient samples and five healthy donor

samples were monitored. Patient samples with blast percentage more than 49% were

chosen for testing with the Cell Titer-Glo (CTG) assay (Table S5). The average inhibition

across dosages on 16 patient samples is regarded as efficacy, whereas the average

inhibition across dosages on healthy samples is regarded as toxicity. The drug

combinations with rectangular labels have higher efficacy and lower toxicity than the

median. The proportion test (p-value $=$ 0.006) revealed that the percentages of inter-

cluster drug combinations with high efficacy (efficacy higher than the third quantile of

efficacy values) and low toxicity (toxicities lower than the first quantile of toxicities) are

significantly more than random choices.

The synergy and combination ratio (CR) of drug combinations on AML and healthy

samples was then calculated using synergy scoring functions has [25], bliss [26], loewe [27], and

ZIP [28] (Figure 4 and S3). The same analysis was done on synergy scoring values, and

it was discovered that inter-cluster drug combinations differ significantly from random

choices (P-values shown in the Figure 4 A-F). The drug combinations shown with

rectangular labels have the highest synergy on AML patient samples, and the lowest

synergy on healthy samples. Table 2 summarizes all six plots and the significant drug

combinations according to different measures are highlighted by green (inter-cluster),

yellow and purple (intra-clusters). Following CTG analysis, consensus across synergy

scoring functions led to the selection of the five best drug combinations out of 28 to quantify

blast-specific drug responses with flow cytometry.

Cell subtype viability analysis highlights low toxicity of selected combinations

Using the CTG assay, we measure the general bone-marrow mononuclear cell (BM-MNC)

sensitivity, whereas with flow cytometry analysis we measure the number of live cells

among different cell populations. Following 72-hour treatment with the 5 selected

combinations on 3 different samples, viability of different cell subtypes of interest was

measured by flow cytometry. For each sample, there is a specific plate layout which can be

found in Figure S1. We used six cell surface markers (CD14, CD15, CD45, CD38, CD117,

CD34) (Table S6) to identify the major leukocyte populations present in the AML BM-

MNCs: monoblasts, myelocytes, leukemic blasts, leukemic stem cells, and myeloid

progenitor cells (Figure S2).

In the studied samples, the average of blasts out of CD45 positive leukocytes, was 70% in

DMSO, while on average 36% $\pm$ 16% of the blasts were killed by the combinations (Table

S7). Based on the results, the percentage of dead cells for all 5 combinations in

lymphocytes is considerably lower than 25% (Figure 5). More importantly selected

combinations have lower synergistic effect on lymphocytes compared to the blast

population, demonstrating the lower toxicity of combinations (Figure 7 and S3). The

combination of JAK1/2 inhibitor (ruxolitinib) with either ERK or CSNK2A1 inhibitor

has the highest efficacy and lowest toxicity which represents the important role of this

target in AML. Numerous studies show the significance of the JAK/STAT signaling

system in determining how hematopoietic cells react to various cytokines and growth

factors [29,30]. Recently there has been increased interest in different drug combinations with

ruxolitinib [31-34] and as our results show the combinations of this drug, by having the

lowest toxicity, seems to be promising for AML treatment.

Blast-specific drug responses in AML: Efficacy profiles of selected combinations

We were able to assess blast-specific drug combination responses and compare them to the

other combinations within different samples. Among the five tested combinations, two

combinations with ruxolitinib which targets JAK1/2 were among the most efficient

combinations. The combination of ruxolitinib with ulixertinib, an ERK inhibitor, exhibits

the strongest efficacy against blasts, according to the results. After treatment, the

combination induced 47% ± 13% cell death in blasts (Figure 5 and Table S7) with a more

synergistic effect on the blast population compared to the lymphocyte population

(Figure 7A). We depicted the gating of 1000 nM concentration of each drug on sample

AML_3 to better understand the impact of combination therapy vs. DMSO control and

single drug treated samples in Figure 6. The number of blast cells in the ruxolitinib and

ulixertinib treated well was reduced to 37%, showing the largest reduction compared to all

other treatments, as shown in Figure 6A. The second combination of ruxolitinib and

silmitasertib, a CSNK2A1 inhibitor, showed high efficacy on blasts. On average, this

combination induced death to almost half ± 14% of the blast population but had less effect

on lymphocytes (Figure 5 and Table S7). Additionally, this combination had a substantially

higher inhibition rate compared to each single drug and acted synergistically towards the

blast population (Figure 7A).

Given the importance of pan-RAF inhibition, we next examined LY3009120 in combination

with three other drugs. The samples used for the combination of LY3009120 and

sapanisertib (mTOR1/2 inhibitor), consist of 56% blast and the response for them is 40% ±

12% inhibition. To confirm that this combination is efficient, we analyzed the effect of

LY3009120 and sapanisertib combination with single treated and DMSO treated cells in

AML_3. In the combination treated sample, the blast cells were significantly reduced to

13% while in the individual drugs LY3009120 and sapanisertib reduced the blasts to 38%

and 75%, respectively (Figure   6B). These results indicate that this combination has

substantially higher inhibition rate compared to each single drug and a greater synergistic

effect on blasts than on lymphocytes (Figure 7A). Ulixertinib (ERK inhibitor) is the

second drug that is being used in combination with pan-RAF inhibitor. Patient samples

treated with this combination, on average, contain 60% blast cells and after treatment they

are reduced to 28% ± 14%. Finally, we tested the combination of LY3009120 with

silmitasertib, a CSNK2A1 inhibitor on three different samples. The average blast

population for these three samples is 62% and the response was 21% ± 5%. Overall, as

shown in Figure 5 and 7A, all combinations have very little impact on the lymphocyte

populations, demonstrating low toxicity, and significantly more impact on less differentiated

malignant cells, demonstrating the efficacy of the combinations.

Increased sensitivity of AML samples to combination therapies over single drugs,
regardless of genetic mutations and prognosis categories

There is a significant correlation between CTG assay and blast specific results, indicating that

reduction in cell number measured by CTG, is related to the malignant cell populations

(Figure 7B and 7C). The cell viability readout for a single drug is converted to a drug

sensitivity score (DSS) which is a drug sensitivity metric based on area under the dose-

response curve. A greater DSS indicates higher sensitivity[35]. Strikingly, by combining

selected inter-cluster drugs, the blasts were targeted, and combinations showed a synergistic

effect on this population (Figure 7A). Then in order to monitor the drug responses based on

genetic changes, we examined the existing mutations in selected samples that are the most

significant mutations among AML patients (Figure 7A) [36,37]. To evaluate the impact of the

combinations on samples bearing genetic alterations, some mutations that are frequently

found in AML patients were considered (Figure 7A). *FLT3* mutation is a well-known mutant

gene among AML patients and is represented in two samples. The other prevalent mutations

were *NPM1, GATA2, DNMT3A, TET2, KMT2A, NRAS, SMC3* and *SRSF2*. The combinations

induced a synergistic effect on the blast population, regardless of the genetic alterations. The

European Leukemia Network (ELN) classifies patients into three prognosis categories:

"favorable", "intermediate", or "adverse"[38]. AML patients are also classified based on

morphological features which is named French-American-British Classification (FAB) [39].

Regardless of sample type, we observed a synergistic effect following treatment. Importantly,

after therapy, we noticed a synergistic effect in all samples, indicating that these combinations

are effective at combating the heterogeneity of the AML disease. It has been demonstrated

that drugs should target the less differentiated leukemic blasts to achieve the best response in

patients [6]. Given these two facts—the presence of the most relevant mutations and the

prevalence of blast cells in the samples— the combinations seem to be promising for

treatment.

Efficacy and toxicity of the novel combinations compared to first-line treatment in AML

In the following analysis, we compared the proposed combinations in this study (ruxolitinib-

ulixertinib and LY3009120-sapanisertib) with two FDA-approved combinations (venetoclax-

azacitidine and venetoclax-cytarabine), as well as the investigational combination of

venetoclax-ruxolitinib. As illustrated in Figure 8, venetoclax-ruxolitinib demonstrates the

highest efficacy on both blast cells and lymphocytes compared to the other combinations.

This dual efficacy profile is a noteworthy advantage; however, it comes at the cost of

heightened toxicity, as indicated by our results.

Conversely, the novel combinations, ruxolitinib-ulixertinib and LY3009120-sapanisertib,

show comparable efficacy in targeting blast populations as the established combinations.

Notably, there is no significant difference in terms of efficacy (p-values are shown in Figure

8). However, these two combinations have a significant advantage in demonstrating lower

toxicity compared to first-line combinations, particularly for lymphocytes. The effect of

ruxolitinib-ulixertinib and LY3009120-sapanisertib on blast lymphocyte population is

significantly lower than all other combinations except for venetoclax-cytarabine (p-value =

0.25) which is not significant but still is lower. This reduction in toxicity suggests these

combinations can offer effective treatment while minimizing side effects associated with

prevalence therapies.

Discussion

In this study we employed a nominal data mining approach to construct a weighted bipartite

network for the selection of the most effective [10], as well as the least toxic drugs. The selected

combinations inhibit important signaling pathways and included pan-RAF inhibitors, JAK1/2

inhibitors, BET family inhibitors, topoisomerase II inhibitor, CSNK2A1 inhibitors, ERK

inhibitors, mTOR1/2 inhibitors, and DNA binding drugs.

The MAPK (RAS/RAF/MEK/ERK) signaling pathway is hyperactivated in AML patients,

leading to leukemogenesis, leukemia progression, and chemo resistance [40-43]. Since,

targeting RAS and ERK remains challenging, pan-RAF inhibitors are an interesting new

pharmacological class in this context [44,45]. Recent studies demonstrated that the pan-RAF

inhibitor LY3009120 induces growth inhibition and apoptosis in RAS-mutated AML cell

lines [46,47]. Another component of the MAPK pathway is ERK, and the ERK inhibitor

ulixertinib is a promising early efficacy agent for treating patients with tumors harboring

alterations in the MAPK pathway [48]. The ERK pathway is activated in approximately 30% of

all human cancers via *RAS, BRAF,* or *MAP2K1 (MEK1)* activating mutations and thus

targeted therapeutics are vital to treating ERK pathway–activated human cancers [49]. Bhagwat

et al. demonstrated the efficacy of LY3214996 as an ERK inhibitor to delay or reverse

resistance to BRAF and MEK inhibitor therapy. They also demonstrated a synergistic effect

of the combination of LY3214996 with a pan-RAF inhibitor, LY3001920, in a KRAS-mutant

colorectal cancer xenograft model [50]. Moreover, here we identified ulixertinib and

LY3009120 as an inter-cluster combination with high efficacy and low toxicity. Upon further

analysis, we found two combinations of ulixertinib and three combinations of LY3009120

among the five most efficient and synergistic combinations.

The other signaling pathway is JAK signaling, which plays critical roles in several important

intracellular signaling pathways [51,52]. In a number of leukemias, aberrations of the JAK/STAT

pathway and constitutive STAT activation have been well described [53]. Ruxolitinib has been

approved as a non-selective JAK1/2 inhibitor for patients with intermediate to high risk

primary or secondary myelofibrosis [54]. Ruxolitinib, by inhibiting JAK 1/2, reduce activation

of JAK-signal transducers and lower down the transcription (STAT) signaling [55]. Based on

our computational analysis, CTG results, and flow cytometry analysis, we revealed that

combinations of ruxolitinib with ulixertinib and silmitasertib are the most blast-specific

compounds, while having minor effect on lymphocytes in AML *ex vivo* screening.

Next, we examined the effect of Bromodomain and Extra-Terminal motif (BET) protein

inhibitors on cancer cells as BET proteins are often altered in cancer. While bromodomain

inhibition is effective, not all patients benefit from it and become resistant over time [56].

Researchers recently reported the efficacy of combined JAK/BET inhibition on cytokine

production, BM fibrosis, and tumor burden *in vivo* [57], leading to complete reversal of reticulin

fibrosis in mice. In our model, this drug was classified among effective drugs in cluster 1

and based on the CTG assay, in combination with sapanisertib, ulixertinib, and teniposide

can be potential compound for the treatment of AML patients.

The other drug that was selected is teniposide which is a chemotherapy medicine used to treat

solid tumors and leukemia, typically in combination with other antineoplastic agents. This

agent causes DNA damage in proliferating cells by inhibiting topoisomerase II. Recently, it

has been proven that teniposide not only inhibits topoisomerase II but also inhibits MYB,

transcription factor, activity and expression [58]. Topoisomerase II inhibitors' ever-evolving

role in the treatment of AML has led to an increase in their utilization in combination

therapies [59]. Here in this study, we identified the combination of teniposide and birabresib

among the most effective compounds with low toxicity.

The next drug is silmitasertib that inhibits CSNK2A1. Studies showed increased

expression of this kinase in AML cells, which contributes to several biological processes

related to cancer progression such as apoptosis, cell growth, proliferation and transcription [60].

Silmitasertib is currently being evaluated in clinical trials for the treatment of multiple

malignancies [61]. Hereby, considering the importance of pan-RAF and CSNK2A1

inhibitors, targeting these two pathways simultaneously might be a promising treatment.

Additionally, as we suggested, the combination of silmitasertib and ruxolitinib could be

considered as potential combinations for treating AML patients.

We then, studied the effect of plicamycin, which inhibits RNA synthesis by binding to GC-

rich regions of DNA. Plicamycin has potent anti-tumor effects, both as mono-therapy and in

combination with anti-tumor drugs [62]. A highly selective adenosine triphosphate-

competitive inhibitor of mTOR kinase, named sapanisertib has been also included in this

study. Sapanisertib has demonstrated antitumor activity in AML by inhibiting both

mTORC1 and mTORC2 [63]. Sapanisertib in combination with LY3009120, birabresib or

ulixertinib has synergistic effect in AML samples (Figure 4).

Lastly, we thoroughly compared the two FDA-approved combinations (venetoclax-

azacitidine and venetoclax-cytarabine) and one investigational combination (venetoclax-

ruxolitinib) with the novel combinations suggested in this study, namely ruxolitinib-

ulixertinib and LY3009120-sapanisertib. Notably, the outcomes validated the comparability

of the suggested combinations' efficacy with first-line combinations in this investigation.

There is no significant difference in efficacy on blast cells, between proposed combinations

and first-line AML combinations. Significantly, the toxicity of selected combinations is lower

than others, except for venetoclax-cytarabine, which indicates that proposed combinations

might offer both effective treatment and a reduced side effect compared to standard AML

combinations.

In summary, we proposed effective drug combinations for AML patients with the highest

efficacy and lowest toxicity based on nominal data mining method and *ex vivo* drug

sensitivity assay. The combinations were tested on three different samples, and the results

indicate that the combinations of ruxolitinib-ulixertinib and sapanisertib-LY3009120 have

the highest synergistic effect, the highest efficacy on blasts, and the lowest toxicity.

Although the approach of combining targeted agents suffers from cumulative toxicity effects

[64], we have demonstrated that our approach overcomes this limitation in designing a drug

combination in AML. Considering the importance of toxicity, in all steps we considered

toxicity as an important factor for the selection of combinations with the lowest effect on

healthy cells. Standard chemotherapy kills most of the blasts as well as other cell types and

has a high value of toxicity [6], while recommended combinations in this study are effective on

blasts, but have lower toxicity on other cell populations.

Acknowledgments

We thank the patients and donors who participated in the study. Drug screening was carried out at the
FIMM High Throughput Biomedicine Unit (HTB), which is hosted by the University of Helsinki and supported
by HiLIFE and Biocenter Finland. Alun Parsons is gratefully acknowledged for his technical support. We also
appreciate comments from Marc Baumann, Julio Saez-Rodriguez and Farnaz Barneh. This study was
financially supported by the Research Council in Finland [grant 332454 to M.J.; grants 320185, 334781, 357686
to C.A.H.], and Jane & Aatos Erkko Foundation [grant 220031 to M.J.], and the Sigrid Jusélius Foundation
[grant to C.A.H.].

Authorship Contributions

M.J. conceived of the study and designed research questions. M.J. and C.A.H. oversaw the scientific
direction, supervised the project, and assisted with the manuscript. Also C.A.H. contributed resources
including patient samples. M.M. led the computational and statistical analysis, interpreted the data, and
assisted with study design, while E.G., T.R., and J.S. designed and developed the experimental methods for
cell viability analysis. J.M. and F.I. contributed to the interpretation of the findings for drug sensitivity
analysis. M.M and S.M contributed bone marrow samples from healthy donors.
408
409   † M.M. contributed equally to this work with E.G.
410

Conflict of Interest

412   C.A.H. has received research funding from BMS/Celgene, Kronos Bio, Novartis, Oncopeptides, Orion
413   Pharma, WNTResearch, Glykos Finland, and IMI2 projects HARMONY and HARMONYPLUS, plus
414   personal fees from Amgen and Autolus unrelated to this work. The remaining authors declare no competing
415   interests.
416
417

Data availability statement

The data that support the findings of this study and all code for data analysis are openly available at https://github.com/jafarilab/DrugComb_AML.

References

Bowman, R. L., Busque, L. & Levine, R. L. Clonal hematopoiesis and evolution to hematopoietic malignancies. *Cell stem cell* **22**, 157--170 (2018).

Grzywa, T. M., Paskal, W. & Wodarski, P. K. Intratumor and intertumor heterogeneity in melanoma. *Translational oncology* **10**, 956--975 (2017).

Bennett, J. M. *et al.* Proposals for the classification of the acute leukaemias French-American-British (FAB) co-operative group. *British journal of haematology* **33**, 451--458 (1976).

Peipert, J. D. *et al.* Patient-reported outcomes predict overall survival in older patients with acute myeloid leukemia. *Journal of Geriatric Oncology* **13**, 935--939 (2022).

Daver, N. *et al.* New directions for emerging therapies in acute myeloid leukemia: The next chapter. *Blood cancer journal* **10**, 1--12 (2020).

Kuusanmki, H. & Lepp. Phenotype-based drug screening reveals association between venetoclax response and differentiation stage in acute myeloid leukemia. *Haematologica* **105**, 708 (2020).

Intlekofer, A. M. *et al.* Acquired resistance to IDH inhibition through trans or cis dimer-interface mutations. *Nature* **559**, 125--129 (2018).

Lopez, J. S. & Banerji, U. Combine and conquer: challenges for targeted therapy combinations in early phase trials. *Nature reviews Clinical oncology* **14**, 57--66 (2017).

Ianevski, A. *et al.* Patient-tailored design for selective co-inhibition of leukemic cell subpopulations. *Science advances* **7**, eabe4038 (2021).

Jafari, M. *et al.* Bipartite network models to design combination therapies in acute myeloid leukaemia. *Nature communications* **13**, 1--12 (2022).

Malani, D. *et al.* Implementing a Functional Precision Medicine Tumor Board for Acute Myeloid LeukemiaAML Functional Molecular Precision Medicine. *Cancer Discovery* **12**, 388--401 (2022).

Ianevski, A., Timonen, S., Kononov, A., Aittokallio, T. & Giri, A. K. SynToxProfiler: An interactive analysis of drug combination synergy, toxicity and efficacy. *PLoS computational biology* **16**, e1007604 (2020).

Sarma, A. *et al.* Expression of aberrant CD markers in acute leukemia: A study of 100 cases with immunophenotyping by multiparameter flowcytometry. *Cancer Biomarkers* **15**, 501--505 (2015).

Blondel, V. D., Guillaume, J.-L., Lambiotte, R. & Lefebvre, E. Fast unfolding of communities in large networks. *Journal of statistical mechanics: theory and experiment* **2008**, P10008 (2008).

Tang, J. *et al.* Drug target commons: a community effort to build a consensus knowledge base for drug-target interactions. *Cell chemical biology* **25**, 224--229 (2018).

Park, J. *et al.* PanRAF inhibitor LY3009120 is highly synergistic with lowdose cytarabine, but not azacitidine, in acute myeloid leukemia with RAS mutations. *Oncology letters* **22**, 1-11 (2021).

Klein, K., Stoiber, D., Sexl, V. & Witalisz-Siepracka, A. Untwining anti-tumor and immunosuppressive effects of JAK inhibitors—a strategy for hematological malignancies? *Cancers* **13**, 2611 (2021).

Gilan, O. *et al.* Selective targeting of BD1 and BD2 of the BET proteins in cancer and immunoinflammation. *Science* **368**, 387-394 (2020).

Albertini, V. *et al.* Novel GC-rich DNA-binding compound produced by a genetically engineered mutant of the mithramycin producer Streptomyces argillaceus exhibits improved transcriptional repressor activity: implications for cancer therapy. *Nucleic acids research* **34**, 1721-1734 (2006).

Blanco-Aparicio, C. & Carnero, A. Pim kinases in cancer: diagnostic, prognostic and treatment opportunities. *Biochemical pharmacology* **85**, 629-643 (2013).

21   Hayes, T. K. & Der, C. J. Targeting the Raf-MEK-ERK mitogen-activated protein
kinase cascade for the treatment of RAS mutant cancers. *Ras Superfamily Small G*
Proteins: Biology and Mechanisms 1: General Features, Signaling, 135-156 (2014).
22   Shimizu, T. *et al*. A phase 1 study of Sapanisertib (TAK-228) in East Asian patients
with advanced nonhematological malignancies. *Targeted oncology*, 1-10 (2022).
23   Economides, M. P., McCue, D., Borthakur, G. & Pemmaraju, N. Topoisomerase II
inhibitors in AML: past, present, and future. *Expert opinion on pharmacotherapy*
20, 1637-1644 (2019).
24   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for
animals and plants. *Bioinformatics* **36**, 2628--2629 (2020).
25   Berenbaum, M. C. What is synergy? *Pharmacological reviews* **41**, 93--141 (1989).
26   Bliss, C. I. The toxicity of poisons applied jointly 1. *Annals of applied biology* **26**,
585--615 (1939).
27   Loewe, S. The problem of synergism and antagonism of combined drugs.
Arzneimittelforschung **3**, 285--290 (1953).
28   Yadav, B., Wennerberg, K., Aittokallio, T. & Tang, J. Searching for drug synergy in
complex dose--response landscapes using an interaction potency model.
Computational and structural biotechnology journal **13**, 504--513 (2015).
29   Ferrajoli, A., Faderl, S., Ravandi, F. & Estrov, Z. The JAK-STAT pathway: a
therapeutic target in hematological malignancies. *Current cancer drug targets* **6**,
671--679 (2006).
30   Eghtedar, A. *et al*. Phase 2 study of the JAK kinase inhibitor ruxolitinib in patients
with refractory leukemias, including postmyeloproliferative neoplasm acute myeloid
leukemia. *Blood, The Journal of the American Society of Hematology* **119**, 4614--
4618 (2012).
31   Borate, U. *et al*. Novel Combination Therapy of Venetoclax and Ruxolitinib in the
Treatment of Patients with Relapsed/Refractory Acute Myeloid Leukemia. *Blood*
138, 2333 (2021).
32   Verbeke, D. *et al*. Ruxolitinib synergizes with dexamethasone for the treatment of T-
cell acute lymphoblastic leukemia. *Hemasphere* **3** (2019).
33   Dal Bello, R. *et al*. A multiparametric niche-like drug screening platform in acute
myeloid leukemia. *Blood Cancer Journal* **12**, 1--12 (2022).
34   Karjalainen, R. *et al*. JAK1/2 and BCL2 inhibitors synergize to counteract bone
marrow stromal cell--induced protection of AML. *Blood, The Journal of the*
American Society of Hematology **130**, 789--802 (2017).
35   Yadav, B. *et al*.   (2014).
36   Papaemmanuil, E. *et al*. Genomic classification and prognosis in acute myeloid
leukemia. *New England Journal of Medicine* **374**, 2209--2221 (2016).
37   Tyner, J. W. *et al*. Functional genomic landscape of acute myeloid leukaemia. *Nature*
562, 526--531 (2018).
38   Rausch, C. *et al*. Validation of the 2022 European Leukemianet genetic risk
stratification of acute myeloid leukemia. *Blood* **140**, 3408--3409 (2022).
39   Hasserjian, R. P. Controversies in the recent (2016) World Health Organization
classification of acute myeloid leukemia. *Best Practice & Research Clinical*
Haematology **34**, 101249 (2021).
40   Ricciardi, M. R. *et al*. Quantitative single cell determination of ERK phosphorylation
and regulation in relapsed and refractory primary acute myeloid leukemia
Tandem-duplicated Flt3 constitutively activates STAT5 and MAP kinase and introduces
autonomous cell growth in IL-3-dependent cell lines. *Leukemia* **19**, 1543--1549
(2005).
41   Steelman, L. S. *et al*. Roles of the Ras/Raf/MEK/ERK pathway in leukemia therapy.
Leukemia **25**, 1080--1094 (2011).
42   Chang, F. *et al*. Signal transduction mediated by the Ras/Raf/MEK/ERK pathway
from cytokine receptors to transcription factors: potential targeting for therapeutic
intervention. *Leukemia (08876924)* **17** (2003).

43   Kornblau, S. M. *et al.* Simultaneous activation of multiple signal transduction
pathways confers poor prognosis in acute myelogenous leukemia. *Blood* **108**, 2358--
2365 (2006).
44   Cox, A. D., Fesik, S. W., Kimmelman, A. C., Luo, J. & Der, C. J. Drugging the
undruggable RAS: Mission possible? *Nature reviews Drug discovery* **13**, 828--851
(2014).
45   Tambe, M. *et al.* Pan-RAF inhibition induces apoptosis in acute myeloid leukemia
cells and synergizes with BCL2 inhibition. *Leukemia* **34**, 3186--3196 (2020).
46   Khoury, J. D. *et al.* Pan-RAF inhibition shows anti-leukemic activity in RAS-mutant
acute myeloid leukemia cells and potentiates the effect of sorafenib in cells with FLT3
mutation. *Cancers* **12**, 3511 (2020).
47   Park, J. *et al.* Pan-RAF inhibitor LY3009120 is highly synergistic with low-dose
cytarabine, but not azacitidine, in acute myeloid leukemia with RAS mutations.
Oncology letters **22**, 1--11 (2021).
48   idecabtagene Vicleucel, C. A. R. T. C. T. FDA updates highlighting the Latest Cancer
Treatments. *Oncology Times* (2020).
49   Roberts, P. J. & Der, C. J. Targeting the Raf-MEK-ERK mitogen-activated protein
kinase cascade for the treatment of cancer. *Oncogene* **26**, 3291--3310 (2007).
50   Bhagwat, S. V. *et al.* ERK inhibitor LY3214996 targets ERK pathway--driven cancers:
a therapeutic approach toward precision medicine. *Molecular cancer therapeutics*
19, 325--336 (2020).
51   Ihle, J. N. & Kerr, I. M. Jaks and Stats in signaling by the cytokine receptor
superfamily. *Trends in Genetics* **11**, 69--74 (1995).
52   Rawlings, J. S., Rosler, K. M. & Harrison, D. A. The JAK/STAT signaling pathway.
Journal of cell science **117**, 1281--1283 (2004).
53   Lin, T. S., Mahajan, S. & Frank, D. A. STAT signaling in the pathogenesis and
treatment of leukemias. *Oncogene* **19**, 2496--2504 (2000).
54   Furqan, M., Mukhi, N., Lee, B. & Liu, D. Dysregulation of JAK-STAT pathway in
hematological malignancies and JAK inhibitors for clinical application. *Biomarker*
research **1**, 1--10 (2013).
55   McKeage, K. Ruxolitinib: a review in polycythaemia vera. *Drugs* **75**, 1773--1781
(2015).
56   Bechter, O. & Schffski, P. Make your best BET: The emerging role of BET inhibitor
treatment in malignant tumors. *Pharmacology & Therapeutics* **208**, 107479 (2020).
57   Kleppe, M. *et al.* Dual targeting of oncogenic activation and inflammatory signaling
increases therapeutic efficacy in myeloproliferative neoplasms. *Cancer cell* **33**, 29--
43 (2018).
58   Yusenko, M., Jakobs, A. & Klempnauer, K.-H. A novel cell-based screening assay for
small-molecule MYB inhibitors identifies podophyllotoxins teniposide and etoposide
as inhibitors of MYB activity. *Scientific reports* **8**, 1--11 (2018).
59   Economides, M. P., McCue, D., Borthakur, G. & Pemmaraju, N. Topoisomerase II
inhibitors in AML: past, present, and future. *Expert opinion on pharmacotherapy*
20, 1637--1644 (2019).
60   Kang, K.-W. *et al.* A Proteomic Approach to Understand the Clinical Significance of
Acute Myeloid Leukemia--Derived Extracellular Vesicles Reflecting Essential
Characteristics of Leukemia. *Molecular & Cellular Proteomics* **20** (2021).
61   Roffey, S. E. & Litchfield, D. W. CK2 Regulation: Perspectives in 2021. *Biomedicines*
9, 1361 (2021).
62   Chen, C. & Lan, M. S. A promoter-driven assay for INSM1-associated signaling
pathway in neuroblastoma. *Cellular Signalling* **76**, 109785 (2020).
63   Shimizu, T. *et al.* A Phase 1 Study of Sapanisertib (TAK-228) in East Asian Patients
with Advanced Nonhematological Malignancies. *Targeted oncology* **17**, 15--24
(2022).
64   Tannock, I. F. & Hickman, J. A. Molecular screening to select therapy for advanced
cancer? *Annals of Oncology* **30**, 661-663 (2019).

Figure legends

Fig. 1. Schematic outline of the study. Data pre-processing began after data collection, which was
followed by full matrix extraction, weighted bipartite network reconstruction, and computational validation.
After the selection of the best combinations, bone marrow and peripheral blood samples from both healthy
individuals (n=5) and AML patients (n= 16) were subjected to drug sensitivity assessment. For ATP-based
viability assay the study design contains 8 drugs and 28 combinations in 384-well plates, each drug with 5
different concentrations and two replicate. The single cell sensitivity assay using the iQue® Screener PLUS
flow cytometer was performed in 384-well plates to monitor drug effects on cell sub-types. The study
design contains 5 drugs and 3 combinations, all with two replicates and five concentrations. For
sapanisertib, the drug concentrations are 0.1, 1, 10, 100, and 1000 nM, and for all other drugs are 1, 10,
100, 1000, and 10000 nM.

Fig. 2. Sankey plot of enriched (A) KEGG signalling pathways and (B) GO biological processes
related to target protein clusters PPT1 and PPT2. Each rectangle on the right side represents a pathway
or biological process, and the size of each rectangle illustrates the degree of connectivity of each pathway.
Each biological process or pathway is represented by a unique color. GO and KEGG pathway enrichment
analysis on proteins that are merely targets by drugs in one cluster. G1 (G2) includes proteins that are
targeted by at least three drugs in cluster1 (cluster2) (155 and 141 drugs). (C) Biological processes (BPs) of
G1, (D) Biological processes (BPs) of G2, (E) KEGG pathway related to G1 proteins and (F) KEGG
pathway related to G2 proteins. The size of the node corresponds to number of genes, the x-axis is Fold
Enrichment and the color of bars indicates the negative logarithm of Fold Enrichment.

Fig. 3. Comparative analysis of dice similarity and drug efficacy-toxicity profiles in AML therapy.
(A)Box plot of dice similarity coefficient indices comparing intra-cluster 1 and 2 to inter-cluster compound
pairs. P-value is generated using Wilcoxon signed-rank test, shown in red color. (B) The toxicity and
efficacy of 296 drugs. Inset plot shows the relationship between toxicity and efficacy. Top five percent of
drugs whose toxicity is less than the average of all drug toxicity and whose efficacy is greater than the average
of all drug efficacy are in blue, and their name is shown in rectangle labels.

Fig. 4. Drug combinations' synergy scores on 16 AML samples and 5 healthy samples. X-axis
depicts the synergy on AML samples and the Y-axis represents the synergy on healthy samples. The
median inhibition on AML and healthy samples, respectively, is shown by the dashed lines in red and
blue. There are three groupings: clusters 1, 2, and inter-cluster, and the color of each dot indicates each of
these groups.

Fig. 5. The cell viability assay (CTG) and response of different cell populations to 5 selected
combinations using flow cytometry assay. Response signifies the percentage of dead cells following a 72-
hour treatment. For each combination three different samples, distinguished by the color of points, were
treated with three doses (10, 100, 1000), which are illustrated by different shapes of points. The colors in
each cell group facet corresponds to a specific drug combination.

Fig. 6. Flow cytometry scatter plots showing the effects of drug combinations on cell populations,
along with comparisons to DMSO and single drug treated samples. (A) This figure illustrates the effects
of ruxolitinib and ulixertinib combination and (B) LY3009120 and sapanisertib combination on blasts,
monocytic cells (CD14+) and lymphocytes after 72h drug treatment. Numbers are presenting the percentage
of cell counts in each cell population. The plot represents concentration of 1000 nM on sample AML_3.

Fig. 7.   Characteristics of drug responses and correlation analysis in AML treatment:
Comprehensive flow cytometry and CTG assessment. (A)Heat map showing characteristics of single
agent and combinations response measured by flow cytometry readout. Blast-specific response of single
drugs is highlighted according to drug sensitivity score (DSS) values with dark blue corresponding to high
DSS value and white to low DSS value. Blast-specific and lymphocyte-specific response combinations in
1000 nM are highlighted according to percentage of apoptotic/dead cells, with dark blue in blast and red for
lymphocyte corresponding to high percentage and white to a low percentage of apoptotic/dead cells. The
synergistic effect of the drug combination was assessed based on the HSA synergistic score in 1000 nM
on blast cells shown in blue and lymphocytes shown in red. (B) The correlation between responses
measured by CTG and flow cytometry on five single drugs ruxolitinib, silmitasertib, ulixertinib,
LY3009120, and sapanisertib, and (C) five drug combinations sapanisertib-LY3009120, ulixertinib-
ruxolitinib, silmitasertib-ruxolitinib, silmitasertib-LY3009120, and ulixertinib-LY300912.

Fig. 8. Flow cytometry assay of selected combinations compared to first-line AML combinations.
Response signifies the percentage of dead cells following 72-hour treatment. Each combination has been
tested on different samples at 50 nM concentration for venetoclax and 1000 nM for the other drugs. Red
asterisks define the average response for each combination and colored dots represent different samples.
Each panel also represents six p-values resulting from the Wilcoxon signed-rank tests to compare
statistically two proposed combinations with three other combinations (including two first-line treatments
and one investigational combination) for AML.

Table legends

Table 1. The selected chemical compounds from two clusters of drugs in the drug similarity
network.

Table 1. Selected drug combinations sorted by synergy scoring functions. Highlighted in
green (inter-cluster), yellow (intra-cluster 1), and purple (intra-cluster 2). Inh stands for the
average of inhibition of drug combinations on dosages, ZIP, hsa, bliss, and loewe are synergy
scorings and CR is the combination ratio.

Figure 1

- Cell viability analysis among different subpopulations
- Ranking drug combinations based on efficacy and toxicity analysis

Flow cytometry single cell-viability assay

- Incidence matrix generation
- Full submatrix extraction
- (sample: 81 Compound: 296)
- Min-max normalization

Preprocessing of drug response data set

- Synergy analysis of drug combinations
- Using consensus synergy analysis to narrow down the list of drug combinations ($n_1 = 2$, $n_2 = 3$)

Drug combination sensitivity analysis

FIMM AML

Nominal data mining

- Bipartite weighted network construction
- Similarity network projection
- Community detection

- Dice similarity analysis of drugs SMILE structure
- KEGG and GO enrichment analysis of protein targets

Intra-cluster homogenity analysis

- AML bone marrow samples (n = 16),
- Healthy samples (n = 5)
- Drug combinations (n = 28, 16 inter-cluster, 12 intra-cluster)

ATP-based bulk viability assay (CTG)

Inter-cluster combinations of drugs

- Shortlisting single drugs in each cluster ($n_1 = n_2 = 4$) with high efficacy and low toxicity for combinatorial design

Figure 2

A) KEGG enriched pathways

B) Biological Process

C) Biological Process: Protein targets G1

D) Biological Process: Protein targets G2

E) KEGG: Protein targets G1

F) KEGG: Protein targets G2

Figure 6

Figure 7

A)

Single drugs	AML sample	1	2	3	4	5
	Ruxolitinib			19.40	23.40	48.00
	Silmitasertib	7.40	1.70	2.30	3.90	5.00
	Ulixertinib	12.90	0.20	5.80	5.50	10.40
	LY3009120	17.10	8.50	13.10	15.40	38.70
	Sapanisertib	44.60	0.00	9.50		

DSS (FC Blast) — 0, 20, 40, 60; N/A

Drug combinations		1	2	3	4	5
	Sapa_LY30	92.10	49.50	85.00		
	Ulix-Ruxo			57.10	80.30	95.50
	Silm-Ruxo			45.50	63.80	93.50
	Silm-LY30	85.70	37.10		60.00	
	Ulix-LY30	89.60	33.10			94.70

%Apoptotic(FC Blast) — 0, 40, 80, 100; N/A

Drug combinations		1	2	3	4	5
	Sapa_LY30	12.89	25.51	16.55		
	Ulix-Ruxo			6.67	2.60	70.84
	Silm-Ruxo			14.76	1.55	59.68
	Silm-LY30	19.27	11.02		-14.02	
	Ulix-LY30	11.17	8.44			60.92

%Apoptotic(FC Lymphocytes) — 0, 40, 80, 100; N/A

Blast		1	2	3	4	5
	Sapa_LY30	9.35	15.60	19.06		
	Ulix-Ruxo			18.14	14.67	7.34
	Silm-Ruxo			5.88	0.00	4.92
	Silm-LY30	1.14	4.58		0.93	
	Ulix-LY30	4.47	0.62			6.27

HSA(Synergy) — 0, 5, 10, 20; N/A

Lymphocyte		1	2	3	4	5
	Sapa_LY30	6.38	12.89	6.03		
	Ulix-Ruxo			6.35	2.59	10.70
	Silm-Ruxo			14.75	1.54	-2.57
	Silm-LY30	12.74	12.75		0.00	
	Ulix-LY30	4.64	4.64			11.61

HSA(Synergy) — 0, 5, 10, 20; N/A

Molecular Profiling		1	2	3	4	5
	Disease Stage	Rf	R	D	D	R
	GATA2	N/A	Non-mutated	Non-mutated	Non-mutated	Mutated
	DNMT3A	N/A	Non-mutated	Non-mutated	Non-mutated	Mutated
	NPM1	N/A	Non-mutated	Non-mutated	Mutated	Mutated
	FLT3	N/A	Non-mutated	Non-mutated	Mutated	Mutated
	TET2	N/A	Mutated	Non-mutated	Non-mutated	Non-mutated
	NRAS	N/A	Mutated	Non-mutated	Non-mutated	Non-mutated
	SMC3	N/A	Mutated	Non-mutated	Non-mutated	Non-mutated
	SRSF2	N/A	Mutated	Non-mutated	Non-mutated	Non-mutated
	KMT2A	N/A	Non-mutated	Mutated	Non-mutated	Non-mutated
	CROCC	N/A	Non-mutated	Mutated	Non-mutated	Non-mutated
	FAB subtype	M5	N/A	M5	M2	M5
	Blast Ratio	0.16	0.76	0.76	0.94	0.89

Non-mutated; Mutated; N/A

B) Single Drugs — cor = 0.97 (FC–Blast vs CTG)

C) Drug Combinations — cor = 0.90 (FC–Blast vs CTG)

Table 1. The selected chemical compounds from two clusters of drugs in the drug similarity network.

Cluster 1	Cluster 2
LY3009120 (LY30)	Teniposide (Teni)
Ruxolitinib (Ruxo)	Silmitasertib (Silm)
Birabresib (Bira)	Ulixertinib (Ulix)
Plicamycin (Plic)	Sapanisertib (Sapa)

Table 1. Selected drug combinations sorted by synergy scoring functions. Highlighted in green (inter-cluster), yellow (intra-cluster 1), and purple (intra-cluster 2). Inh stands for the average of inhibition of drug combinations on dosages, ZIP, hsa, bliss, and loewe are synergy scorings and CR is the combination ratio.

Drug Combinations	Inh	ZIP	hsa	bliss	loewe	CR
LY3009120 , Silmitasertib		green	green	green	green	green
Ulixertinib , Ruxolitinib	green	green			green	
Silmitasertib , Ruxolitinib	green		green			
Sapanisertib , LY3009120		green		green		
Ulixertinib , LY3009120				green		green
Ulixertinib , Plicamycin		green				
Birabresib , Ulixertinib		green				
Birabresib , Teniposide	green					
LY3009120 , Ruxolitinib				yellow	yellow	
Sapanisertib , Ulixertinib						purple
Ulixertinib , Silmitasertib		purple				