1

**Inhibitory G proteins play multiple roles to polarize sensory hair cell**

**morphogenesis**

Amandine Jarysta[1], Abigail L.D. Tadenev[1], Matthew Day[1], Barry Krawchuk[1], Benjamin E.

Low[1], Michael V. Wiles[1], and Basile Tarchini[1,2] [#]

[1]The Jackson Laboratory, Bar Harbor, ME 04609, USA. [2]Tufts University School of

Medicine, Boston, MA 02111, USA.

[#]Corresponding author: basile.tarchini@jax.org

**ABSTRACT**

Inhibitory G alpha (GNAI or Gαi) proteins are critical for the polarized morphogenesis of sensory hair cells and for hearing. The extent and nature of their actual contributions remains unclear, however, as previous studies did not investigate all GNAI proteins and included non-physiological approaches. Pertussis toxin can downregulate functionally redundant GNAI1, GNAI2, GNAI3 and GNAO proteins, but may also induce unrelated defects. Here we directly and systematically determine the role(s) of each individual GNAI protein in mouse auditory hair cells. GNAI2 and GNAI3 are similarly polarized at the hair cell apex with their binding partner GPSM2, whereas GNAI1 and GNAO are not detected. In *Gnai3* mutants, GNAI2 progressively fails to fully occupy the subcellular compartments where GNAI3 is missing. In contrast, GNAI3 can fully compensate for the loss of GNAI2 and is essential for hair bundle morphogenesis and auditory function. Simultaneous inactivation of *Gnai2* and *Gnai3* recapitulates for the first time two distinct types of defects only observed so far with pertussis toxin: 1) a delay or failure of the basal body to migrate off-center in prospective hair cells, and 2) a reversal in the orientation of some hair cell types. We conclude that GNAI proteins are critical for hair cells to break planar symmetry and to orient properly before GNAI2/3 regulate hair bundle morphogenesis with GPSM2.

**INTRODUCTION**

Developing sensory hair cells (HC) in the inner ear undergo a complex polarization process to detect and interpret mechanical stimuli, including sound. Each mature HC is able to detect stimuli in a directional manner by developing an asymmetric brush of actin-based membrane protrusions, or stereocilia: the hair bundle. Neighboring HC also adopt concerted orientations to align their hair bundles, a property known as planar cell polarity (PCP). One subclass of heterotrimeric guanine nucleotide-binding (G) protein was intimately associated with different levels of mouse HC polarization: inhibitory G alpha subunits (GNAI1, GNAI2, GNAI3 and GNAO; collectively GNAI or G$\alpha$i) (Figure 1A). However, several roles proposed for GNAI proteins have not been validated physiologically, and the individual contribution of each GNAI remains uncertain.

HC polarization along the epithelial plane starts with the off-center migration of the basal body and its associated primary cilium, the kinocilium (Denman-Johnson & Forge, 1999; Mbiene & Sans, 1986; Tilney et al, 1992). At that stage, Regulator of G protein Signaling 12 (RGS12) is required for GNAI and the G protein signaling modulator 2 (GPSM2) scaffold to form a polarized complex at the apical membrane on the side of the off-center basal body (Akturk et al, 2022; Ezan et al, 2013; Tarchini et al, 2013). GPSM2-GNAI is best known as a highly conserved protein complex orienting the mitotic spindle during progenitor divisions (Du et al, 2001; Schaefer et al, 2001; Schaefer et al, 2000; Woodard et al, 2010). In post-mitotic HC, GPSM2-GNAI first excludes microvilli and microvilli-derived stereocilia from the portion of the HC surface where it resides, the expanding bare zone (Figure 1B). Bare zone expansion then pushes back the basal body/kinocilium from a more eccentric position near the lateral junction to a less eccentric position at the vertex of the forming hair bundle. Later, GPSM2-GNAI becomes enriched at the distal tip of row 1 stereocilia abutting the bare zone (Tarchini et al, 2016). In these stereocilia, GPSM2-GNAI is a module of the elongation complex that also comprises MYO15A, WHRN and EPS8 (Mauriac et al, 2017; Tadenev et al, 2019). GPSM2-GNAI is required at row 1 tips for boosting enrichment of other elongation complex partners, and presumably actin incorporation, compared to further stereocilia rows. GPSM2-GNAI thus

confers row 1 its tallest identity and the hair bundle its asymmetric graded-height
morphology.

Pertussis toxin (ptx) has been extensively used as a tool to ADP-ribosylate the GNAI
subunit and dissociate heterotrimeric Gαiβγ protein complexes from G protein-coupled
receptors (GPCR) to inactivate downstream signaling (Locht et al, 2011). *In vivo*
expression of ptx catalytic subunit (ptx-S1 or ptxA) prevents normal enrichment and
polarization of GPSM2-GNAI in developing HC (Tadenev et al, 2019; Tarchini et al, 2013),
suggesting that ADP-ribosylation directly or indirectly inhibits GPSM2-GNAI function as
well. Ptx provokes immature-looking hair bundles with severely stunted stereocilia,
mimicking defects in *Gpsm2* mutants and *Gnai2; Gnai3* double mutants (Beer-Hammer
et al, 2018; Mauriac et al, 2017; Tadenev et al, 2019; Tarchini et al, 2016). In contrast,
stereocilia height is more variably reduced in *Gnai3* single mutants, with defects more
severe at the cochlear base (Beer-Hammer et al, 2018; Mauriac et al, 2017). This can
explain why hearing loss is more severe at high frequencies in *Gnai3* mutants, but
profound at all frequencies in a *ptxA* model and in mutants lacking GPSM2 or both GNAI2
and GNAI3 (Beer-Hammer et al, 2018; Mauriac et al, 2017; Tarchini et al, 2016).
Surprisingly, before affecting hair bundle differentiation at postnatal stages, ptx also
causes two distinct defects in HC polarization at embryonic stages.

First, one study reported that a high dose of ptx in cultured explants of the developing
cochlea results in a low proportion of symmetrical HC with a central kinocilium surrounded
by a rounded hair bundle (Figure 1C) (Ezan et al, 2013). Because GPSM2-GNAI recruits
partners to pull on astral microtubules during mitotic spindle orientation, the authors
proposed that GPSM2-GNAI functions similarly and triggers the basal body off-center
migration when post-mitotic HC break planar symmetry. However, this hypothesis has not
been validated *in vivo* to date. Studies where *Gpsm2* (Bhonker et al, 2016; Ezan et al,
2013; Mauriac et al, 2017; Tarchini et al, 2013), *Gnai3* (Beer-Hammer et al, 2018; Ezan
et al, 2013; Mauriac et al, 2017) or *Gnai2; Gnai3* (Beer-Hammer et al, 2018) were
inactivated did not report symmetrical HC. In addition, mouse strains expressing ptxA *in*

*vivo* also did not produce symmetrical cochlear HC (Tarchini et al, 2013; Tarchini et al,
2016).

Second, ptx experiments induced striking HC misorientation. In the cochlea,
misorientation manifested as a 180° inversion of outer HC in the first and second row
(OHC1-2) whereas inner HC (IHC) and OHC3 were much less affected (Figure 1C) (Ezan
et al, 2013; Kindt et al, 2021; Tarchini et al, 2013). In the vestibular system, ptxA
expression abrogated the line of polarity reversal and thus the mirror-image HC
organization characteristic of macular organs, the utricle and saccule (Jiang et al, 2017;
Kindt et al, 2021). Normal orientation reversal was also lost upon inactivating two
endogenous mouse proteins: the transcription factor EMX2 in the maculae (Jiang et al,
2017) and the orphan GPCR GPR156 in cochlear OHC1-2 and in the maculae (Kindt et
al, 2021). Together, these recent studies uncovered an EMX2>GPR156>GNAI signaling
cascade that secures a normal pattern of HC orientation by reversing *Emx2*-positive HC.
GNAI signals downstream of GPR156 to reverse the orientation of the basal body
migration in *Emx2*-positive compared to *Emx2*-negative HC (Tona & Wu, 2020) (reviewed
in (Tarchini, 2021)). While ptx impact on orientation thus appears to be physiologically
relevant, HC misorientation was surprisingly not reported in single *Gnai3* or double *Gnai2;*
*Gnai3* mutants (Beer-Hammer et al, 2018; Mauriac et al, 2017).

In summary, the importance of the GPSM2-GNAI complex for hair bundle development
is well-established, but multiple discrepancies cast a doubt on whether GNAI proteins
also assume earlier polarization roles. Specific questions include whether GNAI proteins
participate in the mechanism that pushes the basal body away from the cell center, and
in the distinct EMX2>GPR156 mechanism that makes a binary decision on the direction
of this push (Figure 1B). Furthermore, it remains unclear whether GNAI2 and GNAI3
adopt similar or distinct distributions in HC, and whether GNAI1 or GNAO also participate
in these processes.

In this study, we embarked on a systematic analysis of single and combined *Gnai1*,
*Gnai2*, *Gnai3* and *Gnao1* mouse mutants to solve the actual role(s) of GNAI/O proteins

during HC differentiation. Our results confirm that GNAI3 is the only GNAI/O protein

required for normal hair bundle morphogenesis and normal auditory brainstem response

(ABR) thresholds. In absence of GNAI3, GNAI2 can fully compensate at embryonic

stages but is not enriched with GPSM2 long enough to ensure normal hair bundle

morphogenesis at postnatal stages. We directly demonstrate that GNAI proteins have two

early polarization roles independent of GPSM2 during embryogenesis. In sum, GNAI

function is instrumental for HC to a) break planar symmetry, b) adopt a proper binary

orientation along the PCP axis downstream of EMX2 and GPR156, and c) elongate and

organize stereocilia into a functional hair bundle with GPSM2.

**RESULTS**

**A near-comprehensive collection of *Gnai/o* mouse mutants**

In order to interrogate the individual and combined roles of all inhibitory G proteins during HC differentiation, we obtained or generated mouse strains to build a collection of single and double *Gnai/o* mutants. Single *Gnai1* and *Gnai3* mutants were derived from the *Gnai1^{tm1Lbi}; Gnai3^{tm1Lbi}* double mutant strain (hereafter *Gnai1^{neo}; Gnai3^{neo}*) (Jiang et al, 2002) by segregating individual mutations upon breeding (see Methods and Supp. Table 1 for details on all strains). We generated a new constitutive *Gnai2* mutant strain (*Gnai2^{del}*) carrying a deletion of exons 2 to 4 (Figure 2 Supp. 1A; see Methods). Finally, we obtained and derived two *Gnao1* mutant strains: a constitutive inactivation allele where a neomycin cassette disrupts exon 6 (*Gnao1^{neo}*) (Jiang et al, 1998) and the conditional inactivation allele *Gnao1^{tm1c(EUCOMM)Hmgu}* (hereafter *Gnao1^{flox})*. As simultaneous constitutive loss of GNAI1 and GNAI2 was reported as viable (Plummer et al, 2012), we established a *Gnai1^{neo}; Gnai2^{del}* double mutant strain in addition to *Gnai1^{neo}; Gnai3^{neo}* (Jiang et al, 2002). In contrast, double inactivation of *Gnai2; Gnai3* is lethal around Embryonic (E) day 10.5, before HC are born (Gohla et al, 2007). Consequently, we generated a new *Gnai3^{flox}* strain by flanking exons 2 and 3 with *loxP* sites (Figure 2 Supp. 1B; see Methods). We then generated conditional *FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}* double mutants where *Gnai3* inactivation occurs as early as E8.5 in the otic vesicle (Hebert & McConnell, 2000). Investigating all *Gnai/o* strains in the same genetic background was unrealistic for feasibility and lethality reasons. We reasoned that apical HC development is probably highly constrained and less likely to be influenced by genetic heterogeneity compared to susceptibility to disease, for example.

As two of the three defects directly attributed to GNAI/O dysfunction were only observed using pertussis toxin (Figure 1C), the *Gnai/o* strains above needed to be compared to a strain expressing ptxA in HC. We used our *R26^{LSL-myc:ptxA}* strain (hereafter *LSL-myc:ptxA*) expressing N-terminal myc-tagged ptxA upon Cre recombination (Figure 2 Supp. 1C) (Tarchini et al, 2016). Because a related *R26^{LSL-ptxAa:myc}* strain carrying a C-terminal myc tag (Regard et al, 2007) caused milder HC misorientation defects than *LSL-myc:ptxA* in

the vestibular system (Jiang et al, 2017; Kindt et al, 2021), we wondered whether the myc

tag could weaken toxin activity even perhaps when located N-terminal. Consequently, we

generated a new strain, *R26^{DIO-ptxA}* (hereafter *DIO-ptxA*), where untagged ptxA is flanked

by double-inverted *lox* sites and flipped from the non-coding to the coding strand upon

Cre recombination (Figure 2 Supp. 1D; see Methods) (Schnutgen et al, 2003). In *DIO-*

*ptxA, ptxA* expression is driven by a strong artificial CAG promoter, and not by the

endogenous *R26* promoter as in previous strains (Regard et al, 2007; Tarchini et al,

2016). We bred *LSL-myc:ptxA* and *DIO-ptxA* either with *FoxG1*-Cre active in otic

progenitors (Hebert & McConnell, 2000), or alternatively with *Atoh1-Cre* (Matei et al,

2005) to limit GNAI/O inhibition to post-mitotic HC. Supplementary Table 1 summarizes

the strains used in this study, their origin, genetic background and viability.

**Only GNAI3 is required for hair bundle morphogenesis yet GNAI2 participates**

To investigate the role of individual GNAI/O proteins in HC development, we imaged hair

bundles in 3-week-old mutant and control littermate mice at the mid-cochlear position

using scanning electron microscopy (SEM). Single *Gnai1* or *Gnai2* mutants as well as

double *Gnai1; Gnai2* mutants did not show overt defects in OHC or IHC (Figure 2A). In

single *Gnai3* mutants by contrast, some OHC hair bundles appeared truncated, and IHC

displayed variably shortened row 1 stereocilia as well as supernumerary stereocilia rows

(Figure 2A-B), as previously described (Mauriac et al, 2017). The same defects were

observed in double *Gnai1; Gnai3* mutants and, as reported previously, in the *LSL-*

*myc:ptxA* model (Figure 2A-B) (Tadenev et al, 2019). A constitutive (*Gnao1^{neo}*) or

conditional (*Atoh1-Cre; Gnao1^{flox}*) inactivation of *Gnao1* did not produce overt apical HC

defects (Figure 2 Supp. 1E-F). Conditional inactivation of *Gnao1* also did not enhance

defects in the *Gnai1; Gnai3* mutant background (Figure 2 Supp. 1F).

In addition, *myc:ptxA* expression inverted the orientation of OHC1-2s (Figure 2A), as

expected since ptxA inactivates the EMX2>GPR156>GNAI signaling cascade that

defines HC orientation along the PCP axis (Kindt et al, 2021; Tarchini et al, 2013; Tarchini

et al, 2016). However, a defect in HC orientation was not observed in single *Gnai1*, *Gnai2*,

*Gnai3* mutants, or in double *Gnai1*; *Gnai2* and *Gnai1*; *Gnai3* mutants (Figure 2A). Despite

extensive breeding, we could only obtain one adult animal carrying a double *Gnai2; Gnai3*
inactivation due to postnatal lethality (*FoxG1-Cre; Gnai2^{del/del}; Gnai3^{flox/flox}*; see Supp.
Table 1). Remarkably, this unique specimen not only recapitulated stereocilia stunting
and extra stereocilia rows observed in the *Gnai3^{neo}, Gnai1^{neo}; Gnai3^{neo}* and *LSL-myc:ptxA*
models (Figure 2A-B), but apparently also OHC1-2 misorientation only observed to date
in the *LSL-myc:ptxA* and *Gpr156* mutants (Figure 2A) (Kindt et al, 2021). This result
suggests that endogenous GNAI proteins are involved in HC orientation, and that GNAI2
can function in the EMX2>GPR156>GNAI signaling cascade to secure proper OHC1-2
orientation in *Gnai1; Gnai3* double mutants. This point is verified and expanded below
when neonate HC orientation is addressed.

To acquire a quantitative view of *Gnai* mutant defects and help comparisons, we first
focused on IHC and measured row 1 stereocilia height and width, as well as the number
of stereocilia in row 1 and the number of rows in the bundle in all strains (Figure 2C-F).
This analysis uncovered a *Gnai3^{neo} < Gnai1^{neo}; Gnai3^{neo} < LSL-myc:ptxA < FoxG1-Cre;*
*Gnai2^{del}; Gnai3^{flox}* allelic series along which a) defects increased in severity, as
manifested by increased variability and decreased averages (row 1 height; Figure 2C),
and b) new defects appeared (excess row 1 stereocilia only observed in *LSL-myc:ptxA*
*and Gnai2^{del}; Gnai3^{flox}*; Figure 2E). The phenotypic series moved towards increasingly
immature-looking hair bundles, and increasingly mimicked severe defects in *Gpsm2*
mutants (Mauriac et al, 2017; Tadenev et al, 2019; Tarchini et al, 2016).

To quantify and compare hair bundle truncations in OHC, we measured the length of the
half-bundle 'wings' on each side of the central vertex. We limited this analysis to mutant
strains where hair bundles retained a recognizable vertex, thus excluding *LSL-myc:ptxA*
and *FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}*. We then plotted paired left and right values for each
OHC and tested length variance for each mutant (Figure 2G; Figure 2 Supp. 1G). In
*Gnai1*, *Gnai2* and *Gnai1; Gnai2* mutants, the two wings of each hair bundle had similar
lengths, and variance was similar to control littermates. In contrast, variable truncation of
one wing in *Gnai3* and *Gnai1; Gnai3* mutants resulted in significantly higher length
variance compared to controls (Figure 2G; Figure 2 Supp. 1G). In both *Gnai3* and *Gnai1;*

*Gnai3* mutants, 49% of hair bundles had wings of more different lengths than the worst

OHC outlier in littermate controls (see Source Data file).

In conclusion, all GNAI/O proteins are not equally involved in hair bundle morphogenesis,

with GNAI3 playing a particularly prominent role. GNAI2 makes a clear contribution since

*Gnai3* mutant stereocilia defects dramatically increase in severity when GNAI2 is also

absent in *Gnai2; Gnai3* double mutants. These results confirm previous conclusions

(Beer-Hammer et al, 2018). In addition, we largely rule out that GNAO is involved in apical

HC differentiation. More severe defects in *Gnai1; Gnai3* double mutants compared to

*Gnai3* single mutants may indicate that GNAI1 plays a subtle role. However, we cannot

rule out differences in genetic background as the underlying cause since *Gnai3^{neo}* is in

mixed (29S1/SvImJ; C57BL/6J) and *Gnai1^{neo}; Gnai3^{neo}* is in pure 129S1/SvImJ

background (Supp. Table 1). GNAI1 thus has at best a minimal role in hair bundle

development.

**Only GNAI3 is required for normal auditory brainstem thresholds**

To pair morphological defects with auditory function, we next tested Auditory Brainstem

Response (ABR) in mutants and control littermates at 3 to 4 weeks of age. Anesthetized

animals were presented with pure tone stimuli of decreasing sound pressure intensity (dB

SPL) at 8, 16, 32 and 40 kHz and ABRs were recorded with a subcutaneous probe (see

Methods). Mirroring their overtly normal apical HC morphology, *Gnai1*, *Gnai2* and *Gnai1;*

*Gnai2* mutants showed thresholds comparable to littermate controls at all frequencies

(Figure 3A-C). Similarly, constitutive (*Gnao1^{neo}*) or conditional (*Atoh1-Cre; Gnao1^{flox}*) loss

of GNAO did not alter ABR thresholds (Figure 3 Supp. 1A-B). In contrast, *Gnai3* mutants

were profoundly deaf at 32 and 40 kHz and displayed significantly elevated thresholds at

8 and 16 kHz compared to littermate controls (Figure 3D). *Gnai1; Gnai3* double mutants

shared a similar ABR profile as *Gnai3* single mutants, with apparently higher thresholds

at 8 and 16 kHz although these two strains were not compared as littermates (Figure 3D).

Littermate controls for *Gnai1; Gnai3* double mutants were homozygote for *Gnai1^{neo}*

(*Gnai1^{neo/neo}; Gnai3^{neo/+}*) and appeared to have higher thresholds than littermate controls

for *Gnai3* mutants (*Gnai3^{neo/+}*) at 32 and 40 but not 8 and 16 kHz, whereas loss of GNAI1

did not impact auditory thresholds on its own (Figure 3A). These results match possibly

more severe hair bundle defects when GNAI1 is inactivated along with GNAI3 (Figure

2C-F), and here again, could reflect either a difference in genetic background or a limited

role for GNAI1.

In summary, we confirm that GNAI3, but not GNAI2, is essential for proper auditory

thresholds, as previously proposed (Beer-Hammer et al, 2018). We also clarify that

GNAO does not participate. Beer-Hammer and colleagues previously showed that

inactivating GNAI2 worsens hearing loss in the *Gnai3* null background, as tested in their

*FoxG1-Cre; Gnai2^{flox/flox}; Gnai3^{flox/flox}* adults (Beer-Hammer et al, 2018). Using a

comparable yet distinct model (*FoxG1-Cre; Gnai2^{del/del}; Gnai3^{flox/flox}*), we were largely

unable to obtain young adults and thus could not perform ABR. This suggests that our

model is more severely affected, and would probably lack ABRs, similar to *FoxG1-Cre;*

*Gnai2^{flox/flox}; Gnai3^{flox/flox}* (Beer-Hammer et al, 2018) and *LSL-myc:ptxA* (Tarchini et al,

2016).

**Distribution of individual GNAI proteins in neonate hair cells**

Next, we attempted to define how individual GNAI/O proteins localize in neonate HC.

GNAI/O proteins are close paralogs, with mouse and human GNAI1 and GNAI3 most

similar in the GNAI group, and GNAO more divergent compared to all GNAI (Figure 1A).

Validating and using specific antibodies would thus be challenging. Instead, we used one

commercial antibody raised against GNAI3 (scbt"GNAI3") and one antibody raised

against GNAI2 (pt"GNAI2") and systematically immunolabeled our mouse model

collection to tease apart protein-specific behavior.

Both antibodies produced the familiar GNAI pattern of enrichment at the bare zone and

at row 1 stereocilia tips (Figure 4; arrows and arrowheads, respectively) (Tarchini et al,

2013; Tarchini et al, 2016). Both antibodies revealed generally normal GNAI enrichment

in *Gnai1* (Figure 4A), *Gnai2* (Figure 4B), and *Gnai1; Gnai2* mutants (Figure 4C). This

indicates that pt"GNAI2" is not specific for GNAI2 and can also detect GNAI3. In *Gnai3*

and *Gnai1; Gnai3* mutants, both antibodies still detected GNAI protein at the bare zone

and at stereocilia tips although some HC displayed incomplete enrichment (Figure 4D-

E; detailed analysis below, Figure 5). This indicates that scbt"GNAI3" is not specific for

GNAI3 and can also detect GNAI2. Finally, neither antibody detected consistent signal

over background in *Gnai2; Gnai3* double mutants where GNAI1 is the only GNAI protein

remaining (Figure 4F; *FoxG1-Cre; Gnai2^{del/del}; Gnai3^{flox/flox}*). To test experimentally

whether these antibodies can detect GNAI1, we first electroporated *Gnai1*, *Gnai2* or

*Gnai3* constructs in E13.5 inner ears and cultured the cochlea for 6 days before

immunolabeling. Overexpressed *Gnai1* was efficiently detected by pt"GNAI2" but not by

scbt"GNAI3" whereas, as expected, overexpressed *Gnai3* was detected by scbt"GNAI3"

(Figure 4 Supp. 1A). We next used pt"GNAI2" to immunolabel the gallbladder epithelium

where *Gnai1* expression was specifically reported (mousephenotype.org; *LacZ* reporter

in *Gnai1^{tm1a(EUCOMM)Wtsi}* strain). Signals were visibly reduced in *Gnai1* mutants compared

to littermate controls (Figure 4 Supp. 1B).

Together, these results indicate that a) GNAI2 and GNAI3 share the same polarized

distribution pattern at the bare zone and at stereocilia tips, and b) GNAI1 is absent or

hidden by background signals at the HC apical surface since it can be detected by the

pt"GNAI2" antibody in other contexts. Loss of GNAO in the *Gnao1^{neo}* or *Atoh1-Cre;*

*Gnao1^{flox}* models did not alter signals obtained with scbt"GNAI3" (Figure 4 Supp. 1C-D).

Moreover, a GNAO antibody produced unpolarized apical signals that proved unspecific

since they were unchanged in *Atoh1-Cre; Gnao1^{flox/flox}* mutants (Figure 4 Supp. 1E). Thus,

GNAO likely does not contribute to HC polarization, as also suggested by normal hair

bundles and normal ABR thresholds in *Gnao1* mutants.

**GNAI2 only partially spans the bare zone and stereocilia tips and partially rescues**

**hair bundle development in postnatal hair cells lacking GNAI3**

We next took a closer look at incomplete GNAI enrichment in *Gnai3* and *Gnai1; Gnai3*

mutant HC (Figure 4D-E). In both models, we observed an identical outcome at the P0

cochlear base where remaining GNAI was unable to fully and consistently occupy the HC

sub-domains when GNAI3 was missing (Figure 5A-B; bare zone, arrows; stereocilia tips,

arrowheads). In *Gnai1;* Gnai3 double mutants, the GNAI protein detected is by default

GNAI2, and this is likely also true in single *Gnai3* mutants since GNAI1 is not observed in HC (Figure 4F). Because in all cases GPSM2 co-localized with GNAI2 (Figure 5A-B), these results demonstrate that both GNAI2 and GNAI3 can form a complex with GPSM2 in HC. Intriguingly, the absence of GPSM2-GNAI2 on one side of the bare zone at P0 appeared to coincide with its absence at stereocilia tips on the corresponding side (Figure 5A-B). We thus quantified GNAI2 intensity at the bare zone and at stereocilia tips in half-OHC in *Gnai1; Gnai3* mutants. While as expected control OHC showed little variation in GNAI enrichment in either compartment (Figure 5 Supp. 1A), mutant OHC showed highly variable signals that were significantly correlated between the bare zone and tips in the same half-OHC (Figure 5C). Analyzing HC at different stages and tonotopic positions clarified that GNAI2 is progressively unable to compensate for missing GNAI3. At E18.5, the GPSM2-GNAI2 complex could still occupy the totality of the bare zone in *Gnai1; Gnai3* mutants (Figure 5 Supp. 1B), suggesting that GNAI2 fully compensates for the loss of GNAI3 in embryonic HC. At later stages of HC differentiation, partial compensation by GNAI2 observed at P0 (Figure 5A-C) evolved into a lack of compensation by P6 at the cochlear base, where GNAI2 was no longer detected consistently at the tips of stunted stereocilia in mutant IHC (Figure 5D). In contrast, less mature P6 IHC at the cochlear mid position retained partial GNAI2 enrichment correlated between the bare zone and tips (Figure 5E), as seen at P0 (Figure 5A-B).

The progressive inability of GNAI2 to cover for GNAI3 in individual HC helps explain the unique profile of apical defects in *Gnai3* and *Gnai1; Gnai3* mutants. Loss of GPSM2-GNAI2 in one wing of the OHC hair bundle led to stereocilia degeneration by P8 (Figure 5F). One-sided loss of global GNAI function at stereocilia tips is thus likely the origin of truncated hair bundle wings observed in adults *Gnai3* and *Gnai1; Gnai3* mutant OHC (Figure 2G). We divided the P8 OHC apical surface in two halves based on the position of the basal body at the hair bundle vertex, and measured the length of each hair bundle wing as well as the total apical surface area in the same HC half in *Gnai1; Gnai3* mutants (Figure 5F). We found a significant correlation between these two values (Figure 5G; control graph in Figure 5 Supp. 1C), providing further evidence that loss of stereocilia prompts a corresponding loss of flat HC surface area on the same OHC side (Etournay

et al, 2010). Finally, we asked whether in time GNAI2 is lost in all stereocilia and along

the entire cochlea in *Gnai1; Gnai3* mutants. This proved not to be the case, as GNAI2

could still be detected at stereocilia tips in P28 OHC and IHC, although in low and variable

amounts compared to GNAI tip signals in littermate controls (Figure 5 Supp. 1D-E).

In conclusion, GNAI2 could provide a low dose of GNAI protein at stereocilia tips when

GNAI3 is missing and preserve elongation and height to some extent. Variable GNAI2

amounts thus likely explain why IHC stereocilia have variably reduced heights in absence

of GNAI3 (Figure 2B-C; Figure 5D-E), unlike in *Gpsm2* or *LSL-myc:ptxA* mutants where

they are more uniformly stunted (Beer-Hammer et al, 2018; Mauriac et al, 2017; Tadenev

et al, 2019; Tarchini et al, 2016). Loss of GNAI2 signals that coincides at the bare zone

and at stereocilia tips on the same HC side adds to previous evidence suggesting that

bare zone enrichment is essential for GPSM2-GNAI trafficking to adjacent row 1

stereocilia (Akturk et al, 2022; Jarysta & Tarchini, 2021; Tarchini et al, 2016).

**Combined loss of GNAI2 and GNAI3 delays and de-polarizes bare zone expansion**

**with drastic consequences on stereocilia distribution**

Since GNAI2 and GNAI3 show functional redundancy and are the most important GNAI/O

proteins for hair bundle differentiation, we next focused on characterizing early HC

development in *Gnai2; Gnai3* double mutants (*Foxg1-Cre; Gnai2$^{del/del}$; Gnai3$^{flox/flox}$*) and

comparing defects to those observed previously with ptx. Unlike at adult stages, *Gnai2;*

*Gnai3* double mutants were obtained in close to Mendelian proportions at P0 (see Supp.

Table 1). For this goal, we used the new *DIO-ptxA* allele in case the myc tag hindered

ptxA activity in the *LSL-myc:ptxA* allele. We first validated the new *Gnai3$^{flox}$ and DIO-ptxA*

alleles (Figure 2 Supp. 1B, D). As expected, GNAI signals at the bare zone and stereocilia

tips were normal in *Gnai3$^{flox/flox}$* homozygotes but showed a distinctive incomplete pattern

along with stunted stereocilia upon Cre recombination (Figure 6 Supp. 1A), as in

constitutive mutants (Figure 5A). In fact, the new *DIO-ptxA* model produced identical

apical HC defects to the earlier *LSL-myc:ptxA* strain in single HC (Kindt et al, 2021;

Tarchini et al, 2016) (Figure 6 Supp. 1B). The fraction of inverted OHC1 in the *DIO-ptxA*

model was lower than in the *LSL-myc:ptxA* model when using the postmitotic HC driver

*Atoh1-Cre* (Figure 6 Supp. 1C) but identically encompassed 100% of OHC1 with the early *FoxG1-Cre* driver (Figure 6 Supp. 1D). Similar apical HC defects between strains indicate that the N-terminal myc tag and mild *R26* promoter do not limit ptxA activity in *LSL-myc:ptxA*. For comparison purposes, we used the *FoxG1-Cre* driver to inactivate GNAI2/GNAI3 and to express ptxA, the same driver used by Beer-Hammer and colleagues (Beer-Hammer et al, 2018).

Inactivating GNAI2 and GNAI3 abolished the bare zone at E17.5 based on abnormally uniform F-actin signals at the HC surface compared to controls where low F-actin matched GNAI signals (Figure 6A-B; arrows point to the bare zone). In contrast, E17.5 *DIO-ptxA* HC had developed a distinct bare zone (Figure 6C, arrows). Measuring its surface area by HC type confirmed that the bare zone was virtually absent in *Gnai2; Gnai3* double mutants but only trended as reduced in ptxA-expressing OHC (Figure 6D). By P0, most *Gnai2; Gnai3* mutant HC at the cochlear base had developed a region lacking microvilli or stereocilia, but its position at the apical surface was highly irregular, complementing extremely dysmorphic hair bundles (Figure 6E-F; arrows point to bare regions). In contrast, P0 *DIO-ptxA* HC displayed largely coherent hair bundles and a polarized bare zone (Figure 6G, arrows). Quantifications revealed a significantly reduced bare area in P0 *Gnai2; Gnai3* mutants compared to littermate controls (Figure 6H), showing that bare zone emergence and expansion is greatly delayed and deregulated in this model (compare P0 in Figure 6H and E17.5 in Figure 6D). In P0 *DIO-ptxA* HC, the bare zone surface area was largely comparable to controls, suggesting that a slight delay in expansion is progressively corrected in time in this model (compare P0 in Figure 6H and E17.5 in Figure 6D). These results best illustrate to date the importance of GPSM2-GNAI for bare zone emergence, expansion and polarized positioning. While both mutant models consistently delay bare zone expansion, differences in timing and severity suggest that ptxA is not as effective as the *Gnai2; Gnai3* double mutant to inhibit GPSM2-GNAI function. This is further underscored by severe stereocilia distribution defects observed in *Gnai2; Gnai3* but not *DIO-ptxA* mutants.

As reported previously (Kindt et al, 2021; Tarchini et al, 2013; Tarchini et al, 2016), the

orientation of OHC1-2 expressing ptxA was inverted compared to controls (Figure 6C, G).

Our single *Gnai2; Gnai3* adult mutant sample also appeared to have inverted OHC1

based on hair bundle morphology (Figure 2A). However, HC orientation proved

challenging to assess in neonate *Gnai2; Gnai3* mutants due to highly dysmorphic hair

bundles (Figure 6A-B; E-F). We thus used acetylated tubulin (AcTub) and pericentrin

(PCNT) as markers for the kinocilium and the basal body, respectively. This showed that

in *Gnai2; Gnai3,* but not in *DIO-ptxA* mutants, the basal body and kinocilium were often

in an approximately central position surrounded partially or entirely by stereocilia (Figure

6E-F). Rounded perinatal hair bundles are a hallmark of HC where the early off-center

migration of the basal body, hence symmetry breaking, is defective.

To distinguish symmetry-breaking from HC orientation, we next used the position of the

basal body at the base of the kinocilium to derive both HC eccentricity and HC orientation.

This strategy helped compare the *Gnai2; Gnai3* and *DIO-ptxA* mouse models, and

identified two early functions for GNAI before its association with GPSM2 for stereocilia

elongation.

**GNAI proteins drive the off-center migration of the basal body**

We used the position of the basal body to measure HC eccentricity as a metric for

cytoskeleton asymmetry. Eccentricity was calculated as a ratio reporting how far away

from the cell center the basal body was positioned (see schematic in Figure 7). A perfectly

symmetrical HC with a central basal body would thus have an eccentricity close to 0

whereas a ratio close to 1 would indicate that the off-center basal body is juxtaposed to

the apical junction. Because HC maturation progresses in time and along the tonotopic

axis (cochlear apex to base gradient of increasing maturity), we measured eccentricity at

E17.5 (base and mid cochlear position) and at P0 (base, mid and apex) to understand

how eccentricity progressed at the HC population level.

Control HC at E17.5 had already broken symmetry at the mid position, with eccentricity

averaging ~0.5 and increasing at the more mature base position (Figure 7A). At E17.5

mid and base positions, *Gnai2; Gnai3* mutant HC showed a significantly reduced
eccentricity, suggestive of a delay in the off-center migration of the basal body. We
arbitrarily defined 0.25 as a threshold below which HC were considered near-symmetrical.
Symmetrical HC were not observed in controls but represented up to ~29% of IHC and
~34% of OHC in *Gnai2; Gnai3* mutants at E17.5 (detailed in Figure 7A by stage, position
and HC type; red highlights). By P0, the proportion of symmetrical cells in mutants had
decreased at all positions for OHC (3.3-15.2%), but remained high or increased for IHC
(21.5-42.4%; Figure 7A).

The situation was radically different in the *DIO-ptxA* model. First, no symmetrical HC was
observed at any stage or position in mutants (Figure 7B). In the *DIO-ptxA* model, the
distribution of eccentricity values in IHC was generally similar to controls. By contrast,
OHC eccentricity trended as higher in *DIO-ptxA* compared to littermate controls, with the
difference becoming less significant with OHC differentiation (Figure 7B). Defective bare
zone expansion (Figure 6D, H) can help explain the distinct eccentricity defects in the
*DIO-ptxA* (transiently higher eccentricity) and *Gnai2; Gnai3* (lower eccentricity) mouse
models. The position of the basal body is the sum of two opposite movements (Figure
1B): a) the early off-center migration that brings the basal body in the vicinity of the lateral
junction, and b) the subsequent relocalization towards the cell center upon bare zone
expansion that brings the basal body in contact with the forming hair bundle (Tarchini et
al, 2013). In *DIO-ptxA*, transiently increased eccentricity in OHC is likely the outcome of
a normal off-center basal body migration combined with delayed bare zone expansion
(Figure 6D, H). In *Gnai2; Gnai3* mutants by contrast, decreased eccentricity in OHC
(Figure 7A) stems from defective off-center migration of the basal body combined with an
initially absent (E17.5), and then reduced (P0), bare region (Figure 6D, H). When an
unpolarized bare region eventually emerges in *Gnai2; Gnai3* mutants (Figure 6H), it
impacts basal body position without directionality, leading to highly variable eccentricity
values compared to controls (Figure 7A), and thus variably dysmorphic hair bundles. The
apparent increase in symmetrical IHC over time in *Gnai2; Gnai3* mutants (Figure 7A) may
result from the delayed expansion of the bare region, which will relocalize the basal body
centrally in a proportion of IHC.

In conclusion, symmetry-breaking (Figure 7), bare zone emergence and expansion and stereocilia distribution (Figure 6) are all severely impaired in *Gnai2; Gnai3* mutants. In ptxA mutants in contrast, symmetry-breaking occurs normally, bare zone expansion is only delayed and stereocilia distribution is less affected. It follows that ptxA does not achieve a loss of GNAI2/GNAI3 function as extensive as in *FoxG1-Cre; Gnai2; Gnai3* mutants. This is probably because ptxA only downregulates and does not inactivate GNAI/O proteins, and because ptxA substrates also include GNAI1 and GNAO that could be active in other contexts than polarization in HCs.

## GNAI proteins orient hair cells laterally

GNAI proteins were proposed to signal downstream of the GPR156 receptor to regulate HC orientation in auditory and vestibular organs (Jiang et al, 2017; Kindt et al, 2021). In HC expressing the transcription factor EMX2, GPR156-GNAI reverses the orientation of the off-center basal body migration, and thus HC orientation, compared to *Emx2*-negative HC (Tona & Wu, 2020). To date, however, the HC misorientation profile observed in *Emx2* or *Gpr156* mutants was only recapitulated using ptxA (Figure 1C) (Kindt et al, 2021). If ptxA-based orientation defects are physiologically relevant to GNAI function, they might be observed in *Gnai2; Gnai3* mutants as well. As mentioned above, a caveat to test this assumption is that highly dysmorphic hair bundles in *Gnai2; Gnai3* mutants obscure HC orientation.

To circumvent this limitation, we first excluded near-symmetrical HC in the *Gnai2; Gnai3* mutant datasets since an asymmetric cytoskeleton is pre-requisite for HC to exhibit a defined orientation (excluded: eccentricity<0.4 at E17.5 and <0.25 at P0). Next, we used the position of the off-center basal body to infer the orientation of the remaining asymmetric HC, ignoring the shape of the hair bundle. We measured the angle formed by a vector running from the HC center to the basal body relative to the cochlear longitudinal axis ($\alpha$, see schematic in Figure 8). Angles were plotted in circular histograms where 0° points to the cochlear base and 90° to the lateral edge. At E17.5 at the cochlear mid position, *DIO-ptxA* showed the graded pattern of OHC misorientation observed as

early as these cells break planar symmetry (Kindt et al, 2021), with inverted OHC1 and

OHC2 and imprecisely oriented IHC and OHC3 (Figure 8A). *DIO-ptxA* OHC1-2

maintained this misorientation profile at the E17.5 base and at P0 (Figure 8B-D) as well

as in adults (Figure 2A).

At the E17.5 cochlear mid, many but not all OHC1 were strikingly inverted as well in

*Gnai2; Gnai3* mutants (Figure 8A). The majority of more mature OHC1 at the E17.5

cochlear base were inverted (Figure 8B), as were P0 OHC1 at all positions (Figure 8C-

D; Figure 8 Supp. 1A). Loss of GNAI2 and GNAI3 can thus recapitulate inverted OHC1-

2 observed in the *DIO-ptxA* model with what appears to be a delay. In contrast however,

OHC2 were generally oriented laterally (90°) at E17.5 and at the P0 apex and mid in

*Gnai2; Gnai3* mutants (Figure 8A-C; Figure 8 Supp. 1A). OHC2 only adopted inverted

orientation characteristic of the *DIO-ptxA* model at the P0 base where HC are more

mature (Figure 8D). As OHC2 mature later than OHC1 (Anniko, 1983), this again

suggests a delay where GPR156-GNAI signaling can initially reverse OHC1-2 normally,

but where a lateral orientation cannot be maintained and OHC1-2 ultimately become

inverted as in the *DIO-ptxA* model.

*FoxG1-Cre* is expressed at the otocyst stage in cochlear progenitors (Hebert &

McConnell, 2000) and globally downregulates GNAI/O proteins early on in the *DIO-ptxA*

model. However, *FoxG1-Cre*-mediated deletion at the *Gnai3* locus in *Foxg1-Cre*;

*Gnai2$^{del/del}$*; *Gnai$^{flox/flox}$* mutants might preserve some functional GNAI3 proteins for a

longer time and explain the apparent delay in OHC1-2 misorientation. To test this idea,

we delayed GNAI downregulation by ptxA and asked whether this would result in a milder,

delayed OHC1-2 misorientation phenotype as seen in *Gnai2; Gnai3* mutants. For that

goal, we used the *Atoh1-Cre* driver which is only active in postmitotic HC (Matei et al,

2005). Strikingly, while P0 OHC1 were inverted in *Atoh1-Cre; DIO-ptxA* as in *FoxG1-Cre;*

*DIO-ptxA*, OHC2 generally pointed laterally (90°) in *Atoh1-Cre; DIO-ptxA* as in *Gnai2;*

*Gnai3* mutants at the E17.5 base and the P0 apex (Figure 8 Supp. 1B). However, a large

proportion of OHC2 were inverted at the P0 base, supporting the hypothesis of a delayed

inversion following delayed GNAI loss-of-function.

Together, these results first show that the *ptxA* misorientation pattern is present in *Gnai2;*
*Gnai3* double mutants. Differences in severity between models can be explained by
different timing of GNAI inactivation. Of note, the only surviving *Gnai2; Gnai3* double
mutant we obtained suggests that OHC1-2 inversion is maintained in adults (Figure 2A),
as in *ptxA* and *Gpr156* mutants (Figure 2A) (Kindt et al, 2021). Endogenous GNAI proteins
are thus integral for OHC1-2 to reverse and adopt a proper lateral orientation during
development, and at least transiently to maintain this lateral orientation. Second, these
results also suggest that the GNAI activities required for symmetry-breaking, lateral HC
orientation and for hair bundle morphogenesis are qualitatively different (Figure 9). *Gnai2;*
*Gnai3* mutants are more severely affected considering symmetry-breaking and hair
bundle morphogenesis, whereas *DIO-ptxA* mutants appear more severely affected
considering OHC1-2 orientation. Of note however, IHC and OHC3 were severely
misoriented in *Gnai2; Gnai3* mutants at all stages and positions analyzed, but much less
affected in *DIO-ptxA* mutants (Figure 8; Figure 8 Supp. 1A). We discuss below how
different GNAI protein identity, dose, timing and upstream regulators may underlie
different roles (Figure 9) and explain differences in phenotype across mutant models.

**DISCUSSION**

By examining a large collection of single and combined mutations in *Gnai/o* genes (*Gnai1*, *Gnai2*, *Gnai3*, *Gnao1*), we assign here three distinct roles for GNAI proteins during the apical polarization of a developing HC (Figure 9): a) to drive the centrifugal migration of the basal body when a prospective HC breaks planar symmetry and b) to orient this migration along the PCP axis in a binary manner, and c) to position and elongate stereocilia during hair bundle morphogenesis. Key results in the study include demonstrating that endogenous GNAI proteins indeed serve early functions (a) and (b), since to date only hair bundle defects (c) were reported in knock-out mouse models where *Gnai* genes were targeted (Beer-Hammer et al, 2018; Mauriac et al, 2017). Previous studies suggested that GNAI proteins partner with different regulators to fulfill different roles, organizing and elongating stereocilia by binding the scaffold GPSM2 (c), and reversing HC orientation in *Emx2*-expressing HC downstream of the receptor GPR156 (b).

**Different GNAI/O identity, dosage and timing underlie different roles**

Interestingly, besides involving different regulators, each role also appears to involve different dosage and specific identity of the underlying GNAI protein(s) at different times during HC development (Figure 9). For postnatal hair bundle morphogenesis (c), GNAI proteins are sequestered in the GDP-bound state by GPSM2, forming a complex highly enriched at the bare zone and at row 1 stereocilia tips that can be reliably immunodetected (Akturk et al, 2022; Kindt et al, 2021). In these two sub-cellular compartments, the identity of the GNAI proteins at work matters, with GNAI3 playing a required and prominent role while GNAI2 is important yet not required. Although one of our GNAI antibodies can detect GNAI1, we did not observe GNAI1 signals in HC (Figure 4), and *Gnai1* mutants did not exhibit significant hair bundle defects or ABR threshold shifts (Figure 2; Figure 3A). On the other hand, we observed a trend towards more severe stereocilia defects and more elevated ABR thresholds in *Gnai1; Gnai3* compared to single *Gnai3* mutants (Figure 2C-E; Figure 3D). Together, these results suggest that GNAI1 plays a minor, or no role in hair bundle development, with differences in the genetic

background between strains possibly explaining the latter results. Similarly, we did not observe HC or auditory defects in absence of GNAO.

In stark contrast with its inability to rescue hair bundle morphogenesis (c), GNAI2 can fully rescue the off-center migration of the basal body (a) and its direction which defines HC orientation (b) when both GNAI1 and GNAI3 are constitutively absent. These two early roles are qualitatively different because they do not strictly reflect a difference in timing (a and b occur at the same time) or GNAI/O dose. Symmetry-breaking (a) is affected in *Gnai2; Gnai3* double mutants but is intact in the *ptxA* model despite early (*FoxG1-Cre*) and high (caggs promoter) expression of untagged pertussis toxin. On the contrary, orientation of OHC1-2 (but not IHC and OHC3) is most severely affected in the *DIO-ptxA* model, with *Gnai2; Gnai3* mutants showing delayed defects. PtxA downregulates but does not abolish GNAI/O proteins, and we thus conclude that ptxA cannot achieve a loss of GNAI2 and GNAI3 function as extensive as in double targeted mutants. Among GNAI/O proteins, GNAI2 and GNAI3 may be more specifically required for HC to break symmetry, but other GNAI/O proteins and/or other factors may participate as well since a majority of HC eventually break symmetry even in *Gnai2; Gnai3* mutants. On the other hand, GPR156 is known to signal via heterotrimeric GNAI proteins (Watkins & Orlandi, 2021), and we speculate that any GNAI/O protein may equally relay signals to regulate HC orientation. PtxA would thus be a particularly effective way to disrupt the orientation role (b), with GNAI1 and/or GNAO partially rescuing orientation in *Gnai2; Gnai3* mutants. Our results also suggest that GNAI/O proteins are not only required for OHC1-2 to adopt a normal lateral instead of medial orientation upon symmetry-breaking (normal GPR156-driven reversal; Figure 9), but also to maintain this lateral orientation, at least transiently. Delayed GNAI/O inactivation indeed appears to result in initially normal OHC1-2 lateral orientation followed by a switch to medial orientation in a proportion of cells (abnormal inversion; Figure 8; Figure 8 Supp. 1B).

Regarding dosage, we propose that embryonic activities (a, b) run on low GNAI/O amounts compared to hair bundle functions (c). Of note, GNAI/O proteins are not immunodetected along with GPR156 in HC (b) but are easily detected with GPSM2 at the

bare zone and stereocilia tips (Akturk et al, 2022; Kindt et al, 2021). This may be because

GNAI/O proteins in heterotrimeric G protein complexes cycle dynamically between a GDP

and GTP-bound state at low dose and escape immunodetection. In this context, it should

be noted that a GNAI/O regulator for symmetry-breaking and its mode of action remain

unknown (Figure 9, role a).

A time- and dosage- dependence for early GNAI roles can explain why our conditional

*Gnai2; Gnai3* double mutant model is more severely affected than a comparable

conditional *Gnai2; Gnai3* model where only hair bundle defects (c) were reported (Beer-

Hammer et al, 2018). While Beer-Hammer and colleagues used *FoxG1-Cre* as a driver

as we did, combined GNAI2 and GNAI3 inactivation necessitated Cre recombination at

four floxed loci (*Gnai2$^{flox/flox}$; Gnai3$^{flox/flox}$*). In contrast, our model necessitates

recombination at two loci only (*Gnai2$^{del/del}$; Gnai3$^{flox/flox}$*). We thus conclude that our model

achieves an earlier and further loss of GNAI2/GNAI3 activity. This hypothesis is supported

by extensive postnatal lethality in our model, whereas Beer-Hammer and colleagues

could analyze *Gnai2; Gnai3* double mutants as adults. Our work thus usefully reconciles

previous reports about GNAI functions in HC, notably showing that inactivating

endogenous GNAI proteins does produce early defects (a,b) so far only observed with

ptx. We thus validate all HC defects observed with ptx as physiologically relevant and

specific to GNAI/O function.

**Hair cell break of symmetry**

Although cochlear HC expressing ptxA undergo normal symmetry-breaking in vivo (this

work and (Kindt et al, 2021; Tadenev et al, 2019; Tarchini et al, 2013; Tarchini et al,

2016)), a fraction of utricular and saccular HC expressing myc:ptxA showed an

abnormally central basal body (Kindt et al, 2021). This suggests that a higher dose of

GNAI/O is required for the off-center migration of the basal body in vestibular compared

to cochlear HC, making vestibular HC more susceptible to ptxA. "Ciliary" proteins are

required for ciliogenesis, including kinocilium formation and maintenance, and also play

non-ciliary functions (May-Simera et al, 2015; Sipe & Lu, 2011). Inactivation of

intraflagellar transport protein IFT88 was reported to result in a central basal body and

circular hair bundles in ~10% of OHC (Jones et al, 2008), but the underlying mechanism

was not elucidated. A low proportion of symmetrical HC was also reported in absence of

the CD2 isoform of the protocadherin PCDH15 that forms inter-stereocilia and kinocilium-

stereocilia fibrous links during embryogenesis (Webb et al, 2011). It remains unclear

whether GNAI/O activity participates in *Ift88* or *Pcdh15* mutant defects, or whether GNAI

could be active at the basal body or the kinocilium.

**Hair cell orientation**

The planar cell polarity (PCP) axis is defined by opposite asymmetric enrichment of the

core PCP transmembrane proteins VANGL2 and FZD3/6 and their specific cytosolic

partners at the apical junction of HC and support cells (Deans, 2013; Montcouquiol &

Kelley, 2019; Tarchini & Lu, 2019). Previous work showed that regional *Emx2* expression

reverses how the HC basal body migrates relative to uniform and early-set core PCP

landmarks, notably establishing the line of polarity reversal in the utricle and saccule.

EMX2 activates the GPR156 receptor by triggering its polarized enrichment at the apical

HC junction, where downstream GNAI/O signaling appears to reverse how core PCP

cues are interpreted and to repel the basal body (Figure 9, role b) (Kindt et al, 2021).

EMX2 achieves this goal by preventing expression of the kinase STK32A that suppresses

apical enrichment and polarization of the GPR156 protein (Jia et al, 2023). It is important

to note that HC lacking GPR156 have a normal apical cytoskeleton, including a normally

formed hair bundle, and only show orientation defects (Kindt et al, 2021). This indicates

that the centrifugal migration of the basal body and the direction of this migration that

defines early HC orientation are two distinct molecular mechanisms although both involve

GNAI/O proteins (Figure 9; a *versus* b). Similarly, HC lacking GPSM2 are not inverted in

orientation as in *Emx2*, *Gpr156*, *Gnai2; Gnai3* or *ptxA* mutants (Bhonker et al, 2016; Ezan

et al, 2013; Tarchini et al, 2013). This indicates that bare zone enrichment of the GPSM2-

GNAI complex is a mechanism to shape and polarize the growth of the hair bundle, but

not to migrate off-center or orient the basal body. GPSM2-GNAI is polarized on the side

of the off-center basal body but occupies the HC apical membrane and not the apical

junction where core PCP proteins regulate HC orientation (Siletti et al, 2017; Tarchini et

al, 2013).

While both GPR156 (Greene et al, 2023; Ramzan et al, 2023) and GPSM2 (Doherty et

al, 2012; Walsh et al, 2010) have been identified as human deafness genes, there is

currently no evidence implicating a GNAI/O protein. This is most likely because all

GNAI/O proteins, including GNAI3 which is more specifically required in HC for hair

bundle morphogenesis (Figure 9), play ubiquitous and critical signaling roles in the

context of heterotrimeric protein signaling across many cell types.

**METHODS**

**Mouse strains and husbandry**

All mouse strains used in this work and strain-related information is summarized in Supplementary Table 1. All primers used for genotyping are indicated in Supplementary Table 2 by strain.

Strains from The Jackson Laboratory repository: *Gnai1^neo; Gnai3^neo* (*Gnai1^tm1Lbi; Gna3^tm1Lbi;* MGI: 5560183) (Jiang et al, 2002) carries a neo cassette in *Gnai1* exon 3 and a neo cassette replacing part of intron 5 and exon 6 in *Gnai3* in the 129S1/SvlmJ background. The single *Gnai1^neo* and single *Gnai3^neo* alleles were segregated from *Gnai1^neo; Gnai3^neo* by breeding with C57BL/6J mice and are consequently on a mixed 129S1/SvlmJ: C57BL/6J background. *Gnao1^neo* (*Gnao1^tm1Lbi*; MGI: 2152685) carries a neo cassette in *Gnao1* exon 6 in the 129S1/SvlmJ background (Jiang et al, 2002). The two *Cre* strains used in this work are *Atoh1-Cre* (*Tg(Atoh1-cre)1Bfri;* MGI: 3775845) expressing Cre in HC from E14.5 (Matei et al, 2005) and *FoxG1-Cre* (*Foxg1^tm1(cre)Skm*; MGI: 1932522) expressing Cre in the prospective otic vesicle from E8.5 (Hebert & McConnell, 2000).

Consortium strain: *Gnao1^flox* (*Gnao1^tm1c(EUCOMM)Hmgu*) was derived from the EUCOMM strain *Gnao1^tm1a(EUCOMM)Hmgu* (MGI: 4456727; "KO first, conditional ready"). The *Gnao1^flox* conditional allele was produced via FLP-mediated recombination (FLP strain MGI: 4830363) to remove the FRT-flanked *LacZ-neo* cassette, leaving exon 3 floxed. *Gnao1^flox* is on C57BL/6N background.

Constitutive *Gnai2^del* inactivation (see Figure 2 Supp. 1A). *Gnai2^del* (*Gnai2^em1Btar;* MGI: 6466534) was generated using delivery of CRISPR-Cas9 reagents in mouse zygotes via electroporation. The following guide RNAs (gRNAs) were used to delete exon 2, exon 3 and part of exon 4 in the C57BL/6J background. Upstream gRNA: CTGCCCTCTGTTCCAGGTGC, downstream gRNA: ATGCTTCCTGAAGACCTGTC. The resulting 681 bp deletion encompassed TGCTGGAGAGTCAGGGAAGA...CCTGAAGACCTGTCCGGTGT. The electroporation

mixture consisted of the gRNAs with AltR- *Streptococcus pyogenes* Cas9 (SpCas9) V3

(Integrated DNA Technologies #1081059) in embryo-tested TE buffer (pH 7.5).

Electroporation of zygotes was performed as described in (Qin et al, 2015). In order to

segregate away potential non-specific mutations, founders were bred for two generations

with C57BL/6J animals to generate a N2 heterozygote stock. Because *Gnai2^{del}*

homozygotes showed low viability, this strain was used in a mixed C57BL/6J:FVB/J

background that improved the proportion of homozygotes (see Supp. Table 1).

<u>Conditional *Gnai3^{flox}* inactivation</u> (see Figure 2 Supp. 1B). A plasmid-based donor vector

was cloned to flank *Gnai3* exon 2 and 3 with *loxP* sites using the same CRISPR/Cas9

system as described above for *Gnai2^{del.}* A fragment carrying a *loxP*, a genomic region

including exon 2-3, the restriction site ClaI, a second *loxP* and the restriction site XhoI

was synthetized (Genscript) and cloned between 5' and 3' homology arms amplified by

PCR using Gibson assembly. The following gRNAs were used and define the extremities

of the floxed region: upstream, AGCTCACCAAAATTCCCATT,

TAGGGGATATAGATCCAAAT, downstream, TTCCAGGACTCTGCATGCGT,

TACCGACGCATGCAGAGTCC. The gene editing reagents (gRNAs, donor vector,

SpCas9) were microinjected in zygotes as described in (Qin et al, 2015). To confirm

insertion at the *Gnai3* locus, we performed long-range PCRs with a genomic primer

located outside the homology arm on each side: *5' long-range:* F(external)_

TACTGAGATGAGAGACTGAGGG and R (internal)_TGGCTGACATCCTTTGATGGAC.

The 3070 bp product was digested with XhoI, producing 2,551 + 519 bp fragments upon

donor insertion (flox allele). 3' long-range: F

(internal)_TGAAAGGTAAAGGCAACGTGAG and R

(external)_TGTGAGACAGGGTCTCTCTTTG. The 2,966 bp product was digested with

XhoI, producing 2,000 + 966 bp fragments upon donor insertion (flox allele). In order to

segregate away potential non-specific mutations, founders were bred for two generations

with C57BL/6J animals to generate a N2 heterozygote stock.

<u>Pertussis toxin catalytic subunit (ptxA)-expressing strains </u>(see Figure 2 Supp. 1C-D). The

*LSL-myc:ptxA* strain at the *ROSA26* (*R26*) locus was described previously

(*Gt(ROSA)26Sor*[em1(ptxA)Btar]; MGI: 6163665) (Tarchini et al, 2016). The new *CAG-DIO-ptxA* strain line (see Figure 2 Supp. 1D) was generated at the *R26* locus using the Bxb1 attP(GT) integrase technology and *C57BL/6J-Gt(ROSA)26Sor*[<em2Mvw>/Mvw] (MGI: 6757188) as the host strain (Low et al, 2022). The donor vector consisted in a CAG promoter followed by the flipped ptxA coding sequence flanked by double inverted lox sites (loxP and lox2272; Figure 2 Supp. 1D) and a bGHpA sequence. In order to exclude the prokaryotic vector backbone, the donor vector was prepared as a minicircle using the MC-Easy Minicircle Production kit (SystemBio, MN920A-1). Briefly, the donor plasmid insert was cloned into a Minicircle Cloning Vector using PhiC31 integrase before transformation into the ZYCY10P3S2T *E. coli* minicircle producer strain, which after induction with arabinose, results in the generation of the donor minicircle. To eliminate parental plasmid contamination, a restriction digest was performed, followed by MC-safe DNase treatment. The minicircle DNA was then purified by phenol-chloroform extraction and reconstituted in microinjection buffer (10 mM Tris; 0.1 mM EDTA pH7.5). The *CAG-DIO-ptxA* strain was generated by pronuclear microinjection of the Bxb1 Integration reagents directly into zygotes of the host strain. These reagents included the Bxb1 mRNA (Trilink) at 100ng/µl, RNasin (Promega) at 0.2U/µl and the donor minicircle DNA at 10ng/µl combined in microinjection buffer. To confirm successful integration of the 3,212bp transgene, as well as identify random transgenics, the initial screening was performed using a four-PCR strategy. The In/Out Left and Right (IOL, IOR) PCRs, which bridge the recombined attachment sites, each contain one primer specific to the *R26* locus and a second primer designed against the inserted sequence. The Transgene (TG) PCR amplifies a non-genomic sequence in the insert, and the Off-Target Integration (OTI) PCR was designed to detect non-recombined minicircle integration, presumably outside of the *R26* locus.

IOL PCR (F1/R1), F1: GTCGCTCTGAGTTGTTATCAGT, R1: GCCAAGTAGGAAAGTCCCATAA (720 bp, WT=no band). IOR PCR (F2/R2), F2: GGTGATGCCGTTGTGATAGA, R2: TGTGGGAAGTCTTGTCCCTCCAAT (1,013 bp, WT=no band). TG PCR (F2/R3) F2: 5'-GGTGATGCCGTTGTGATAGA, R3: CCACTTCATCGGCTACATCTAC (214 bp, WT=no band). OTI PCR (F3/R1) F3: GGGAGGATTGGGAAGACAATAG, R1: GCCAAGTAGGAAAGTCCCATAA (578 bp,

WT=no band). 33 mice were born following 6 transfers totaling 117 injected embryos (28% survival), and 1/33 (3%) was identified as a founder and crossed to C57BL6/J to generate N1 offspring. The In/Out PCR confirmed that a copy of the transgene was inserted at the *R26* locus, but the OTI strategy also gave a product, even after breeding for multiple generations. As breeding would have segregated a separate, random integration of the transgene, we performed Nanopore-based Cas9-targeted sequencing at the *R26* locus (Gilpatrick et al, 2020; Low et al, 2022). This revealed that, as seen in some cases previously (Low et al, 2022), tandem insertion had occurred in this founder Specifically, 3 consecutive copies of the *CAG-DIO-ptxA* transgene were inserted at the *R26* locus in this strain. This did not impact specific Cre-based expression of ptxA because phenotypes observed in this new strain were comparable to the previous *LSL-myc:ptxA* strain (Figure 6 Supp.1 B-D).

Experimental animals in the study ranged in age between E17.5 and P29 as indicated in each figure. Male and females were systematically included but sex was not tracked except for Auditory Brainstem Recordings because there is no evidence that sex influences HC orientation or hair bundle morphogenesis. Animals were maintained under standard housing conditions (14h light / 10h dark cycle, ambient temperature and normal humidity). All animal work was reviewed for compliance and approved by the Animal Care and Use Committee of The Jackson Laboratory (Animal Use Summary AUS #14012).

**Scanning Electron Microscopy (SEM)**

Temporal bones were isolated, punctured at the cochlear apex, and fixed by immersion for at least one overnight at 4°C in 2.5% glutaraldehyde (Electron Microscopy Sciences; 16200) and 4% paraformaldehyde (Electron Microscopy Sciences; 15710) in 1mM $MgCl_2$, 0.1M Sodium Cacodylate, 20mM $CaCl_2$. Samples were rinsed and decalcified overnight in 4% EDTA. The auditory epithelium was then dissected into 3 pieces (cochlear base, mid, and apex) before progressive dehydration in an ethanol series (30-50-70-80-90-100%, at least 20 minutes per step) and chemical drying with hexamethyldisilazane (HMDS; Electron Microscopy Sciences 50-243-18). Dry samples were mounted on

aluminum stubs using double-sided carbon tape and sputter-coated with gold-palladium

before imaging on a Hitachi 3000N VP electronic microscope at 20kV.

**Immunofluorescence and antibodies**

For embryonic and postnatal stages, temporal bones were immediately dissected to

expose the sensory epithelium and fixed in paraformaldehyde (PFA 4%; Electron

Microscopy Sciences; 15710) for 1h at 4°C. After fixation, the tectorial membrane was

removed, and samples were permeabilized and blocked in PBS with 0.5% Triton-X100

and bovine serum albumin (1%) for at least at 1 hour at room temperature. For adult

stages, temporal bones were isolated and punctured at the cochlear apex to facilitate

access of the fixative. Samples were then immersion-fixed in PFA 4% for 1 hour at 4°C,

rinsed in PBS, and incubated overnight in 4% EDTA for decalcification. Cochleae were

next dissected in 3 pieces (cochlear base, mid, and apex), before permeabilization and

blocking as described above. Primary and secondary antibodies were incubated

overnight at 4°C in PBS with 0.025% sodium azide. Fluorescent dye-conjugated

phalloidin was added to secondary antibodies. Samples were washed 3 times in PBS +

0.05% Triton-X100 after each antibody incubation and post-fixed in PFA 4% for at least

1 hour at room temperature. Samples were then mounted flat on microscopy slides

(Denville M102) using Mowiol as mounting medium (Calbiochem/MilliporeSigma

4759041), either directly under a 18x18mm #1.5 coverglass (VWR 48366-045) (postnatal

cochleae) or using one layer of office tape as a spacer (adult cochleae). Mowiol (10%

w/v) was prepared in 25%(w/v) glycerol and 0.1M Tris-Cl pH8.5. Primary antibodies used

were:

Rabbit anti-GNAI2, pt"GNAI2" (Proteintech, 11136-1-AP); the antigen is the full human

GNAI2 protein

Rabbit anti-GNAI3, scbt"GNAI3" (Santa Cruz Biotechnology, sc-262); the antigen is

undisclosed but corresponds to a C-terminal region of the rat GNAI3 protein

Rabbit anti-GNAO (Proteintech, 12635-1-AP)

Mouse anti-acetylated alpha tubulin (Santa Cruz Biotechnology scbt-23950)

Rabbit anti-GPSM2 (Sigma, A41537)

Goat anti-GPSM2 (Thermofisher, PA5-18646)

Rabbit anti-Pericentrin/PCNT (Biolegend/Covance, PRB-432C)

Rat anti-ZO1 (Developmental Studies Hybridoma Bank, R26.4C)

Secondary antibodies from ThermoFisher Scientific were raised in donkey and
conjugated to Alexa Fluor (AF) 488, 555, or 647 (donkey anti-rabbit AF555 (A-31572),
AF647 (A-31573), donkey anti-mouse AF647 (A-31571), donkey anti-rat AF488 (A-
21208), donkey anti-goat AF555 (A-21432), AF647 (A-21447)). Fluorescent conjugated
phalloidins used to reveal F-actin were from ThermoFisher Scientific (AF488 (A12379);
AF555 (A34005)) and Sigma-Aldrich (FITC (P5282)).

**Inner ear electroporation and cochlear culture**

Inner ears from wild-type animals (mixed C57BL/6J-FVB/NJ background) were harvested
at E13.5 in HBSS/5mM Hepes. Caggs plasmid vectors (CMV early enhancer, beta-actin
promoter) driving either mouse *Gnai1*, *Gnai2* or *Gnai3* cDNA expression were mixed with
Fast Green FCF (0.05%; Sigma, F7252) and injected at 2.5mg/ml in the cochlear duct
using a Wiretroll capillary and plunger (Drummond, 53507-426). Injected inner ears were
next electroporated (BTX ECM830; 27V, 27 ms, 6 square pulses at 950 ms intervals), the
condensed mesenchyme was dissected away and the soft cochlear labyrinth was
embedded in 50% Matrigel (Corning, CB40234). Cochlear explants were cultured for 6
days in DMEM with 10% fetal bovine serum and 10 µg/ml ciprofloxacin (Sigma, 17850).
Explants were then fixed in 4% PFA for 15 min at room temperature before being
processed for immunolabeling.

**Sample cohorts, image acquisition and analysis**

All quantifications include at least three animals per genotype. All graphs or their legends
indicate the animal cohort size (N) as well as the number of HC or stereocilia (n) analyzed.
Further details can be found in the Supplementary Source Data file where tabs are
referencing figure panels in order. When an experimental outcome was not quantified, at
least 3 mutant and 3 control littermates encompassing two or more litters were analyzed,
and figure panels illustrate a representative outcome observed in all samples of the same

genotype. A single exception is adult *Foxg1-Cre; Gnai2$^{del/del}$; Gnai3$^{flox/flox}$* data in Fig. 2A-

F where a single mutant animal was analyzed since we failed to obtain others at 3 weeks

of age in spite of extensive breeding (see Supp. Table 1).

Confocal images were captured with a LSM800 line scanning confocal microscope, a 63x

NA1.4 oil objective, the Airyscan detector in confocal mode (except in Figure 5 Supp.1 D,

E where the Airyscan detector was used in Airyscan mode) and the Zen 2.3 or Zen 2.6

software (Carl Zeiss AG). Raw Airyscan images in Figure 5 Supp.1 D, E were processed

in Zen 2.6 selecting for automatic strength. Unless stated otherwise in the legends,

images show a single optical z plane. To quantify HC eccentricity (Figure 7), images were

captured with a Leica DM5500B widefield microscope, a 63x oil objective, a Hamamatsu

ORCA-Flash4.0 sCMOS camera and the Leica Application Suite (LasX) software (Leica

Microsystems). All confocal images in the same experiment were acquired using the

same laser intensity and gain, and were then processed in Adobe Photoshop (CC2020)

where the same image treatment was applied across conditions.

To measure stereocilia length and width in IHC (Figure 2C-D), SEM samples were imaged

laterally at 10,000x magnification and with an appropriate tilt (from 0 to 30°) to bring

stereocilia parallel to the imaging plane and minimize parallax. To quantify the number of

rows in adult IHC and the number of stereocilia in their first row (Figure 2E, F), SEM

samples were imaged medially at 5,000x magnification. To measure the length of OHC

hair bundle wings (Figure 2G, Figure 2 Supp. 1E), OHC were imaged top down at 5,000x.

Stereocilia width was measure at half-length and all measurements were done with the

straight-line tool in Fiji.

To quantify GNAI signal intensity in *Gnai1$^{neo}$; Gnai3$^{neo}$* mutants (Figure 5C, Figure 5

Supp. 1A), Z-stack series were acquired at the mid cochlear position. A single Z slice was

chosen at the bare zone level, another one at the stereocilia tip level, each based on

strongest signal in that compartment. The vertex of the V-shaped hair bundle was used

to divide the HC apical surface in two equal halves. In each half (left and right), regions

of interest (ROIs) were drawn using the polygon selection tool in Fiji to encompass all

signals in that apical compartment, and mean gray values were measured. For each
image, background signal was measured and averaged, and subtracted from all
measurements in the same image.

To quantify surface area of the bare zone or total surface area in half-OHC as well as the
length of half hair bundles (Figure 5G, Figure 5 Supp. 1C; Figure 6D, H), Z-stack series
were acquired at the cochlear base and a single Z slice was selected at apical junction
level using ZO1 (Figure 5G, Figure 5 Supp. 1C) or F-actin (Figure 6D, H) as reference.
To measure total apical surface area and bundle length in half-OHC, the pericentrin-
stained basal body was used to divide the apical surface in two halves (Figure 5G, Figure
5 Supp. 1C). The ZO1-positive cell outline along with the dividing line at the basal body
level were used as reference to measure surface area with the polygon selection tool in
Fiji. To quantify the bare zone surface area (Figure 6D, H), a ROI was drawn around the
total apical surface lacking phalloidin (F-actin) signals. The length of the F-actin stained
hair bundle was measured using the straight-line tool in Fiji.

To determine HC eccentricity (Figure 7), the geometrical center of the HC apical surface
was determined as the intersection of two orthogonal lines representing the maximal
diameter of the cell along the cochlear longitudinal and radial axes. A vector (BB) was
drawn from the center to the pericentrin-labeled basal body, and eccentricity was
calculated as the ratio of BB length over the cell radius (r) along the same trajectory using
F-actin-labeled apical junction as landmark (see Figure 7). To determine cell orientation
(Figure 8 and Figure 8 Supp. 1-2), the angle ($\alpha$) separating the longitudinal axis of the
organ of Corti from the BB vector was measured with the angle tool in Fiji (see Figure 8).
Both right and left cochleae were used and angles were measured so that 0° pointed
toward the cochlear base and 90° towards the cochlear periphery (lateral). Eccentricity
and angles were measured at the cochlear positions indicated (base ~20%, mid ~50%,
apex ~80% of the cochlear length starting from the base).

**Auditory Brainstem Response (ABR) tests**

All tests were performed in a sound-attenuating chamber, and body temperature of the anesthetized animals was maintained at 37°C using a heating pad (FHC Inc.). Animals from all strains except *Atoh1-Cre; Gnao1^flox* (Figure 3 Supp. 1B, see below) were anesthetized with a mix of ketamine and xylazine (1 mg and 0.8 mg per 10g of body weight, respectively) and tested using the RZ6 Multi-I/O Processor System coupled to the RA4PA 4-channel Medusa Amplifier (Tucker-Davis Technologies). ABRs were recorded after binaural stimulation in an open field by tone bursts at 8, 16, 32, and in some cases 40 kHz generated at 21 stimuli/second, and a waveform for each frequency/dB level was produced by averaging the responses from 512 stimuli. Subdermal needles were used as electrodes, the active electrode inserted at the cranial vertex, the reference electrode under the left ear and the ground electrode at the right thigh.

*Atoh1-Cre; Gnao1^flox* animals (Figure 3 Supp. 1B) were anesthetized with tribromoethanol (2.5mg per 10g of body weight) and tested with the Smart EP evoked potential system from Intelligent Hearing Systems (IHS). ABR were recorded after single ear stimulation (right ear), using ear tubes speakers (ER3C insert earphones, IHS) delivering tone bursts at 8, 16, and 32 kHz generated at 40 stimuli/second. Electrodes were positioned as previously described except for the ground electrode that was placed at the base of the tail. ABR thresholds were obtained for each frequency by reducing the sound pressure level (SPL) by 5 decibels (dB) between 90 and 20 dB to identify the lowest level at which an ABR waveform could be recognized. We compared waveforms by simultaneously displaying 3 or more dB levels on screen at the same time.

**Statistical Analyses**

All data were plotted in Prism 9 (GraphPad), except for circular diagrams of HC orientation (Figure 8; Figure 8 Supp. 1-2) that were generated with R (4.2.2) and Rstudio (2022.12.0+353).

All data except for angles in circular diagrams (Figure 8 and Figure 8 Supp. 1-2) were plotted individually. Distribution was framed with 25-75% whisker boxes where exterior lines show the minimum and maximum, the middle line represents the median, and + represents the mean. Potential differences in data distribution between genotypes were

tested for significance using nested (hierarchical) t-test except for ABR thresholds (Figure 3 and Figure 3 Supp. 1) where a 2-way ANOVA with Sidak's multiple comparison post-hoc test was used. Nested t-tests help avoid pseudoreplication by taking into consideration the data structure, here specifically variance in each animal (Eisner, 2021; Krey et al, 2023). OHC half-bundle lengths (Figure 2G, Figure 2 Supp. 1E) were plotted as paired left and right values in the same HC and a potential difference in data variance between genotypes was tested using a F-test. GNAI signal intensity at the OHC bare zone and stereocilia tips (Figure 5C, Figure 5 Supp. 1A) as well as OHC surface area and half-bundle length (Figure 5G, Figure 5 Supp. 1C) were plotted and a simple linear regression curve was calculated and drawn for each pair of datasets. A correlation between variables was addressed with Pearson's correlation test. Exact p-values were indicated on each graph when non-significant (p>0.05) and otherwise summarized as follows: p<0.0001 ****, p<0.001 ***, p<0.01**, p<0.05*. Details of statistical results for all figures can be found in the Supplementary Data file.

Angle frequency distribution (Figure 8 and Figure 8 Supp. 1-2) was plotted in circular diagrams using the R package dyplr and the coord_polar function of the ggplot2 package to organize data and produce the graphs, respectively. The angle formed by the red line indicates the circular mean and the length of the arc at the end of the red line indicates the mean circular deviation. Both values were obtained using the colstats function in the R circular package. All scripts used in this work will be posted on github.

**ACKNOWLEDGMENTS**

We are grateful to Elli Hartig for reading and commenting on the manuscript. We thank Simon Lesbirel for nanopore-based Cas9-targeted sequencing of the *DIO-ptxA* strain. We are grateful to The Jackson Laboratory Genome Engineering Technology and Reproductive Science services for their help with generating the *Gnai2^{del}* and *Gnai3^{flox}* strains, and cryorecovering the *Gnai1^{neo}; Gnai3^{neo}* and *Gnao1^{neo}* strains, respectively. A.J. was supported by a postdoctoral fellowship from Fondation Pour l'Audition (2018-2020;

1008   FPA RD-2018-3). This work was supported by the National Institute on Deafness and

1009   Other Communication Disorders grants R01DC015242 and R01DC018304 (to B.T.).

1010

**AUTHOR CONTRIBUTION**

B.T. conceived the study. B.T., A.J and A.L.T designed, performed and analyzed

experiments. M.D. provided technical help with mouse colony maintenance, sample

processing and genotyping. B. K. wrote the R script to generate the circular histograms.

B.E.L and M.V.W. helped generate the *DIO-ptxA* strain using BxB1 recombination

technology. B.T supervised the work. B.T. and A.J. wrote the manuscript. B.T. secured

funding.

1018

1019   **COMPETETING INTERESTS**

1020   The authors declare no competing interest.

1021

1022   **DATA AVAILABILITY**

1023   The data in all graphical representations are included in a single Source Data file. Original

1024   material used for quantifications and to build figures will be deposited in a publicly

1025   accessible repository.

1026

1027

1028

1029

**REFERENCES**

Akturk, A., Day, M. & Tarchini, B. (2022) RGS12 polarizes the GPSM2-GNAI complex to organize and elongate stereocilia in sensory hair cells. *Sci Adv*, 8(42), eabq2826.

Anniko, M. (1983) Postnatal maturation of cochlear sensory hairs in the mouse. *Anat Embryol (Berl)*, 166(3), 355-68.

Beer-Hammer, S., Lee, S. C., Mauriac, S. A., Leiss, V., Groh, I. A. M., Novakovic, A., Piekorz, R. P., Bucher, K., Chen, C., Ni, K., Singer, W., Harasztosi, C., Schimmang, T., Zimmermann, U., Pfeffer, K., Birnbaumer, L., Forge, A., Montcouquiol, M., Knipper, M., Nurnberg, B. & Ruttiger, L. (2018) Galphai Proteins are Indispensable for Hearing. *Cell Physiol Biochem*, 47(4), 1509-1532.

Bhonker, Y., Abu-Rayyan, A., Ushakov, K., Amir-Zilberstein, L., Shivatzki, S., Yizhar-Barnea, O., Elkan-Miller, T., Tayeb-Fligelman, E., Kim, S. M., Landau, M., Kanaan, M., Chen, P., Matsuzaki, F., Sprinzak, D. & Avraham, K. B. (2016) The GPSM2/LGN GoLoco motifs are essential for hearing. *Mamm Genome*, 27(1-2), 29-46.

Deans, M. R. (2013) A balance of form and function: planar polarity and development of the vestibular maculae. *Semin Cell Dev Biol*, 24(5), 490-8.

Denman-Johnson, K. & Forge, A. (1999) Establishment of hair bundle polarity and orientation in the developing vestibular system of the mouse. *J Neurocytol*, 28(10-11), 821-35.

Doherty, D., Chudley, A. E., Coghlan, G., Ishak, G. E., Innes, A. M., Lemire, E. G., Rogers, R. C., Mhanni, A. A., Phelps, I. G., Jones, S. J., Zhan, S. H., Fejes, A. P., Shahin, H., Kanaan, M., Akay, H., Tekin, M., Consortium, F. C., Triggs-Raine, B. & Zelinski, T. (2012) GPSM2 mutations cause the brain malformations and hearing loss in Chudley-McCullough syndrome. *Am J Hum Genet*, 90(6), 1088-93.

Du, Q., Stukenberg, P. T. & Macara, I. G. (2001) A mammalian Partner of inscuteable binds NuMA and regulates mitotic spindle organization. *Nat Cell Biol*, 3(12), 1069-75.

Eisner, D. A. (2021) Pseudoreplication in physiology: More means less. *J Gen Physiol*, 153(2).

Etournay, R., Lepelletier, L., Boutet de Monvel, J., Michel, V., Cayet, N., Leibovici, M., Weil, D., Foucher, I., Hardelin, J. P. & Petit, C. (2010) Cochlear outer hair cells undergo an apical circumference remodeling constrained by the hair bundle shape. *Development*, 137(8), 1373-83.

Ezan, J., Lasvaux, L., Gezer, A., Novakovic, A., May-Simera, H., Belotti, E., Lhoumeau,
A. C., Birnbaumer, L., Beer-Hammer, S., Borg, J. P., Le Bivic, A., Nurnberg, B., Sans, N.
& Montcouquiol, M. (2013) Primary cilium migration depends on G-protein signalling
control of subapical cytoskeleton. *Nat Cell Biol*, 15(9), 1107-15.

Gilpatrick, T., Lee, I., Graham, J. E., Raimondeau, E., Bowen, R., Heron, A., Downs, B.,
Sukumar, S., Sedlazeck, F. J. & Timp, W. (2020) Targeted nanopore sequencing with
Cas9-guided adapter ligation. *Nat Biotechnol*, 38(4), 433-438.

Gohla, A., Klement, K., Piekorz, R. P., Pexa, K., vom Dahl, S., Spicher, K., Dreval, V.,
Häussinger, D., Birnbaumer, L. & Nürnberg, B. (2007) An obligatory requirement for the
heterotrimeric G protein Gi3 in the antiautophagic action of insulin in the liver. *Proc Natl*
*Acad Sci U S A*, 104(8), 3003-8.

Greene, D., Genomics England Research, C., Pirri, D., Frudd, K., Sackey, E., Al-Owain,
M., Giese, A. P. J., Ramzan, K., Riaz, S., Yamanaka, I., Boeckx, N., Thys, C., Gelb, B.
D., Brennan, P., Hartill, V., Harvengt, J., Kosho, T., Mansour, S., Masuno, M., Ohata, T.,
Stewart, H., Taibah, K., Turner, C. L. S., Imtiaz, F., Riazuddin, S., Morisaki, T.,
Ostergaard, P., Loeys, B. L., Morisaki, H., Ahmed, Z. M., Birdsey, G. M., Freson, K.,
Mumford, A. & Turro, E. (2023) Genetic association analysis of 77,539 genomes reveals
rare disease etiologies. *Nat Med*.

Hebert, J. M. & McConnell, S. K. (2000) Targeting of cre to the Foxg1 (BF-1) locus
mediates loxP recombination in the telencephalon and other developing head structures.
*Dev Biol*, 222(2), 296-306.

Jarysta, A. & Tarchini, B. (2021) Multiple PDZ domain protein maintains patterning of the
apical cytoskeleton in sensory hair cells. *Development*, 148(14).

Jia, S., Ratzan, E. M., Goodrich, E. J., Abrar, R., Heiland, L., Tarchini, B. & Deans, M. R.
(2023) The dark kinase STK32A regulates hair cell planar polarity opposite of EMX2 in
the developing mouse inner ear. *Elife*, 12.

Jiang, M., Gold, M. S., Boulay, G., Spicher, K., Peyton, M., Brabet, P., Srinivasan, Y.,
Rudolph, U., Ellison, G. & Birnbaumer, L. (1998) Multiple neurological abnormalities in
mice deficient in the G protein Go. *Proc Natl Acad Sci U S A*, 95(6), 3269-74.

Jiang, M., Spicher, K., Boulay, G., Martin-Requero, A., Dye, C. A., Rudolph, U. &
Birnbaumer, L. (2002) Mouse gene knockout and knockin strategies in application to
alpha subunits of Gi/Go family of G proteins. *Methods Enzymol*, 344, 277-98.

Jiang, T., Kindt, K. & Wu, D. K. (2017) Transcription factor Emx2 controls stereociliary
bundle orientation of sensory hair cells. *Elife*, 6.

Jones, C., Roper, V. C., Foucher, I., Qian, D., Banizs, B., Petit, C., Yoder, B. K. & Chen,
P. (2008) Ciliary proteins link basal body polarization to planar cell polarity regulation. *Nat*
*Genet*, 40(1), 69-77.

Kindt, K. S., Akturk, A., Jarysta, A., Day, M., Beirl, A., Flonard, M. & Tarchini, B. (2021)
EMX2-GPR156-Galphai reverses hair cell orientation in mechanosensory epithelia. *Nat*
*Commun*, 12(1), 2861.

Krey, J. F., Chatterjee, P., Halford, J., Cunningham, C. L., Perrin, B. J. & Barr-Gillespie,
P. G. (2023) Control of stereocilia length during development of hair bundles. *PLoS Biol*,
21(4), e3001964.

Locht, C., Coutte, L. & Mielcarek, N. (2011) The ins and outs of pertussis toxin. *FEBS J*,
278(23), 4668-82.

Low, B. E., Hosur, V., Lesbirel, S. & Wiles, M. V. (2022) Efficient targeted transgenesis
of large donor DNA into multiple mouse genetic backgrounds using bacteriophage Bxb1
integrase. *Sci Rep*, 12(1), 5424.

Matei, V., Pauley, S., Kaing, S., Rowitch, D., Beisel, K. W., Morris, K., Feng, F., Jones,
K., Lee, J. & Fritzsch, B. (2005) Smaller inner ear sensory epithelia in Neurog 1 null mice
are related to earlier hair cell cycle exit. *Dev Dyn*, 234(3), 633-50.

Mauriac, S. A., Hien, Y. E., Bird, J. E., Carvalho, S. D., Peyroutou, R., Lee, S. C., Moreau,
M. M., Blanc, J. M., Geyser, A., Medina, C., Thoumine, O., Beer-Hammer, S., Friedman,
T. B., Ruttiger, L., Forge, A., Nurnberg, B., Sans, N. & Montcouquiol, M. (2017) Defective
Gpsm2/Galphai3 signalling disrupts stereocilia development and growth cone actin
dynamics in Chudley-McCullough syndrome. *Nat Commun*, 8, 14907.

May-Simera, H. L., Petralia, R. S., Montcouquiol, M., Wang, Y. X., Szarama, K. B., Liu,
Y., Lin, W., Deans, M. R., Pazour, G. J. & Kelley, M. W. (2015) Ciliary proteins Bbs8 and
Ift20 promote planar cell polarity in the cochlea. *Development*, 142(3), 555-66.

Mbiene, J. P. & Sans, A. (1986) Differentiation and maturation of the sensory hair bundles
in the fetal and postnatal vestibular receptors of the mouse: a scanning electron
microscopy study. *J Comp Neurol*, 254(2), 271-8.

Montcouquiol, M. & Kelley, M. W. (2019) Development and Patterning of the Cochlea:
From Convergent Extension to Planar Polarity. *Cold Spring Harb Perspect Med*.

Plummer, N. W., Spicher, K., Malphurs, J., Akiyama, H., Abramowitz, J., Nürnberg, B. &
Birnbaumer, L. (2012) Development of the mammalian axial skeleton requires signaling

through the Gα(i) subfamily of heterotrimeric G proteins. *Proc Natl Acad Sci U S A*,
109(52), 21366-71.

Qin, W., Dion, S. L., Kutny, P. M., Zhang, Y., Cheng, A. W., Jillette, N. L., Malhotra, A.,
Geurts, A. M., Chen, Y. G. & Wang, H. (2015) Efficient CRISPR/Cas9-Mediated Genome
Editing in Mice by Zygote Electroporation of Nuclease. *Genetics*, 200(2), 423-30.

Ramzan, M., Bozan, N., Seyhan, S., Zafeer, M. F., Ayral, A., Duman, D., Bademci, G. &
Tekin, M. (2023) Novel GPR156 variants confirm its role in moderate sensorineural
hearing loss. *Sci Rep*, 13(1), 17010.

Regard, J. B., Kataoka, H., Cano, D. A., Camerer, E., Yin, L., Zheng, Y. W., Scanlan, T.
S., Hebrok, M. & Coughlin, S. R. (2007) Probing cell type-specific functions of Gi in vivo
identifies GPCR regulators of insulin secretion. *J Clin Invest*, 117(12), 4034-43.

Schaefer, M., Petronczki, M., Dorner, D., Forte, M. & Knoblich, J. A. (2001) Heterotrimeric
G proteins direct two modes of asymmetric cell division in the Drosophila nervous system.
*Cell*, 107(2), 183-94.

Schaefer, M., Shevchenko, A., Shevchenko, A. & Knoblich, J. A. (2000) A protein complex
containing Inscuteable and the Galpha-binding protein Pins orients asymmetric cell
divisions in Drosophila. *Curr Biol*, 10(7), 353-62.

Schnutgen, F., Doerflinger, N., Calleja, C., Wendling, O., Chambon, P. & Ghyselinck, N.
B. (2003) A directional strategy for monitoring Cre-mediated recombination at the cellular
level in the mouse. *Nat Biotechnol*, 21(5), 562-5.

Siletti, K., Tarchini, B. & Hudspeth, A. J. (2017) Daple coordinates organ-wide and cell-
intrinsic polarity to pattern inner-ear hair bundles. *Proc Natl Acad Sci U S A*, 114(52),
E11170-E11179.

Sipe, C. W. & Lu, X. (2011) Kif3a regulates planar polarization of auditory hair cells
through both ciliary and non-ciliary mechanisms. *Development*, 138(16), 3441-9.

Tadenev, A. L. D., Akturk, A., Devanney, N., Mathur, P. D., Clark, A. M., Yang, J. &
Tarchini, B. (2019) GPSM2-GNAI Specifies the Tallest Stereocilia and Defines Hair
Bundle Row Identity. *Curr Biol*, 29(6), 921-934 e4.

Tarchini, B. (2021) A Reversal in Hair Cell Orientation Organizes Both the Auditory and
Vestibular Organs. *Front Neurosci*, 15, 695914.

Tarchini, B., Jolicoeur, C. & Cayouette, M. (2013) A molecular blueprint at the apical
surface establishes planar asymmetry in cochlear hair cells. *Dev Cell*, 27(1), 88-102.

Tarchini, B. & Lu, X. (2019) New insights into regulation and function of planar polarity in
the inner ear. *Neurosci Lett*, 709, 134373.

Tarchini, B., Tadenev, A. L., Devanney, N. & Cayouette, M. (2016) A link between planar
polarity and staircase-like bundle architecture in hair cells. *Development*, 143(21), 3926-
3932.

Tilney, L. G., Tilney, M. S. & DeRosier, D. J. (1992) Actin filaments, stereocilia, and hair
cells: how cells count and measure. *Annu Rev Cell Biol*, 8, 257-74.

Tona, Y. & Wu, D. K. (2020) Live imaging of hair bundle polarity acquisition demonstrates
a critical timeline for transcription factor Emx2. *Elife*, 9.

Walsh, T., Shahin, H., Elkan-Miller, T., Lee, M. K., Thornton, A. M., Roeb, W., Abu
Rayyan, A., Loulus, S., Avraham, K. B., King, M. C. & Kanaan, M. (2010) Whole exome
sequencing and homozygosity mapping identify mutation in the cell polarity protein
GPSM2 as the cause of nonsyndromic hearing loss DFNB82. *Am J Hum Genet*, 87(1),
90-4.

Watkins, L. R. & Orlandi, C. (2021) In vitro profiling of orphan G protein coupled receptor
(GPCR) constitutive activity. *Br J Pharmacol*, 178(15), 2963-2975.

Webb, S. W., Grillet, N., Andrade, L. R., Xiong, W., Swarthout, L., Della Santina, C. C.,
Kachar, B. & Muller, U. (2011) Regulation of PCDH15 function in mechanosensory hair
cells by alternative splicing of the cytoplasmic domain. *Development*, 138(8), 1607-17.

Woodard, G. E., Huang, N. N., Cho, H., Miki, T., Tall, G. G. & Kehrl, J. H. (2010) Ric-8A
and Gi alpha recruit LGN, NuMA, and dynein to the cell cortex to help orient the mitotic
spindle. *Mol Cell Biol*, 30(14), 3519-30.

1183

1184

**Figure 1. Summary of GNAI-related functions proposed previously in hair cells. A)** Phylogenetic tree of GNAI/O proteins with percent amino acid identity (mouse). **B)** Apical HC differentiation from symmetry breaking to hair bundle development. The distribution of the GPSM2-GNAI complex at the bare zone and stereocilia tips is indicated in orange. Arrows indicate off-center (left) and then inward (middle) movements of the basal body. **C)** Defects observed with pertussis toxin (ptx) or when inactivating GNAI proteins. Defective off-center migration of the basal body and inverted OHC1-2 were only observed with ptx, respectively in cochlear explants (in vitro) and by expressing the ptx catalytic subunit (ptxA) in vivo. Mouse knock-out (KO)s of *Gnai* genes were to date only reported to affect hair bundle morphogenesis. Known GNAI regulators that produce similar defects when inactivated are indicated on top for each type of defect. DKO, double KO.

**Figure 2. Individual GNAI proteins make different contributions to hair bundle development.**
**A-B)** SEM images of representative OHC (A) and IHC (B) in 3-week old animals at the cochlear mid. *Gnai1^{neo}*, *Gnai2^{del}*, and *Gnai1^{neo}; Gnai2^{del}* mutants show apparently normal hair bundles in both HC types. In contrast, *Gnai3^{neo}* and *Gnai1^{neo}; Gnai3^{neo}* mutants show defects in both HC types, including truncated hair bundles in OHC (arrow), as well as supernumerary rows of stunted (hollow arrowheads) or variable height stereocilia (full arrowheads) in IHC. In addition, in *FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}* and *Atoh1-Cre; LSL-myc:ptxA* mutants, OHC1-2s are severely misoriented. **C-F)** Quantification of various hair bundle features in 3-week old IHC at the cochlear mid. Each mutant strain is compared to littermate controls (in black; exact genotypes detailed in Source Data file). At least 3 animals, 17 IHC and 108 stereocilia are represented per condition, except for *FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}* where we could only obtain a single adult animal due to postnatal lethality. Nested (hierarchical) t-test sorted by animal; $p<0.0001$****, $p<0.001$***, $p<0.01$**, $p<0.05$*; non-significant p-values are indicated. Detailed cohort sizes and statistics can be found in Source Data file. **G)** SEM images of representative OHC showing a truncated hair bundle (arrow). Lengths of the left and right wing of the hair bundle were measured and plotted as paired values for the same OHC. A littermate control graph is only shown for *Gnai3* mutants (*Gnai3^{neo/+}* controls). Littermate control graphs for the other mutants can be found in Fig. 2 Supp. 1G. p-values are for a F-test of variance of pooled left and right wing lengths compared to littermate controls. At least 3 animals and 88 OHC are represented per genotype. Only *Gnai3* and *Gnai1; Gnai3* mutants show truncated hair bundles and a significant p value ($p<0.05$). Scale bars are 10μm (A) and 2μm (B, G). OHC, outer hair cell; IHC, inner hair cell.

**Figure 2 Supplement 1. A-D)** New mouse strains generated in this study are *Gnai2del* (A), *Gnai3flox* (B) and *R26DIO-ptxA* (D; see Methods for details). *R26LSL-myc:ptxA* was published previously and is used in this study as well (C). **E-F)** Loss of GNAO has no obvious impact on apical HC morphology or HC orientation. Representative confocal images of phalloidin-stained *Gnao1* constitutive mutants at P21 (E; *Gnao1neo*) and scanning electron microscopy images of conditional *Gnao1* inactivation in the *Gnai1; Gnai3* double mutant background (F; *Gnao1flox*). In F, top panels compare control (left) and conditional

*Gnao1* inactivation (right) when GNAI1 and GNAI3 function is preserved. Bottom panels compare control (left) and conditional *Gnao1* inactivation (right) when GNAI1 and GNAI3 are inactivated. Note how HC-specific inactivation of GNAO using the *Atoh1-Cre* driver does not obviously enhance defects observed in *Gnai1; Gnai3* double mutants (F, bottom). **G)** Graphs showing the length of the left and right wing of 3-week-old OHC hair bundles paired by cell. Graphs for control littermates that were omitted in Fig. 2G are added here (left) next to repeated mutant graphs (right). p-values are for a F-test of variance of pooled left and right wing lengths compared to littermate controls. At least 3 animals and 88 OHC are represented per genotype. Only *Gnai3$^{neo}$* (Fig. 2G) and *Gnai1$^{neo}$*; *Gnai3$^{neo}$* mutants show truncated hair bundles and a significant p value ($p < 0.05$).

**Figure 3. Loss of GNAI3 leads to hearing loss most severe at high frequencies. A-D)** ABR thresholds at 8, 16, 32, and 40 kHz for *Gnai1^neo^* (A), *Gnai2^del^* (B), *Gnai1^neo^; Gnai2^del^* (C), and *Gnai3^neo^* and *Gnai1^neo^; Gnai3^neo^* (D) mutants tested between P21 and P29. Boxplots are framed with 25-75% whisker boxes where exterior lines show the minimum and maximum values, the middle line represents the median, and + represent the mean. A plotted value of 100 dB indicates animals that did not respond to 90 dB stimuli. In (C), controls are a pool of *Gnai1^+/+^; Gnai2^del/+^*, *Gnai1^neo/+^; Gnai2^+/+^* and *Gnai1^neo/+^; Gnai2^del/+^* animals. N indicates the number of animals of both sexes tested. Two-way ANOVA with Sidak's multiple comparison. p<0.0001****, p<0.001***; non-significant p-values are indicated. p-values in orange were obtained comparing non-littermate animals and suggest possibly raised thresholds when GNAI1 is inactivated in addition to GNAI3, or due to a difference in genetic background (see text). kHz, kiloHertz, dB SPL, decibel sound pressure level.

**Figure 3 Supplement 1. A-B)** ABR thresholds at 8, 16, 32 kHz for constitutive *Gnao1* (A) and conditional *Atoh1-Cre; Gnao1^flox/flox^* (B) mutants tested between P20 and P25. Boxplots are framed with 25-75% whisker boxes where exterior lines show the minimum and maximum values, the middle line represents the median, and + represent the mean. N indicates the number of animals tested. In (B), because the original animal cohort was established to test whether the loss of GNAO could enhance defects in *Gnai1; Gnai3* double mutants (see Fig. 2 Supp. 1F), controls include the following genotypes: *Atoh1-Cre; Gnao1^flox/+^, Gnao1^flox/flox^, Gnai1^neo/+^; Gnai3^neo/+^, Gnai1^neo/+^, Atoh1-Cre; Gnao1^flox/+^; Gnai1^neo/neo^; Gnai3^neo/+^* and *Gnai1^neo/neo^; Gnai3^neo/+^* where other alleles are wild-type. Mutants include the following: *Atoh1-Cre; Gnao1^flox/flox^; Gnai3^neo/+^* and *Atoh1-Cre; Gnao1^flox/flox^; Gnai1^neo/neo^; Gnai3^neo/+^* where other alleles are wild-type. Cohort details can be found in Source Data file. Two-way ANOVA with Sidak's multiple comparison. non-significant p-values are indicated. kHz, kiloHertz, dB SPL, decibel sound pressure level.

**Figure 4. Systematic immunolabeling of GNAI proteins in *Gnai* mutant strains. A-F)** Two different antibodies (scbt"GNAI3" and pt"GNAI2") were used to label the auditory epithelium at P0-P3. GNAI proteins were detected at the bare zone (arrows) and at stereocilia tips (arrowheads). Neither antibody is specific for its protein target, as scbt"GNAI3" is able to detect GNAI2 (E) and pt"GNAI2" is able to detect GNAI3 (C). Note how no apical GNAI signal is visible with either antibody in *Gnai2; Gnai3* double mutants (F), showing that GNAI1 is not enriched apically in HC (see Fig. 4 Supp. 1A-B for evidence that pt"GNAI2" detects GNAI1). When the identity of the GNAI protein detected is unambiguous based on genotype, it is made explicit in orange. Scale bars are 10μm.

**Figure 4 Supplement 1. GNAI1 detection by Proteintech GNAI2 antibody, GNAI protein distribution in *Gnao1* mutants and lack of evidence for GNAO enrichment at the hair cell apex.** **A)** Cochlear explants electroporated at E13.5 with the Gnai constructs indicated and cultured for 6 days before immunolabeling with Proteintech (pt"GNAI2"; left) or Santa Cruz Biotechnology (scbt"GNAI3"; right) GNAI antibodies (maximum projections). pt"GNAI2" efficiently detects ectopically expressed GNAI1 (and GNAI2, as expected), but scbt"GNAI3" only weakly detects GNAI1 (and efficiently detects GNAI3, as expected). The greater epithelial ridge (GER) is imaged with IHC on top. Caggs: CMV early enhancer/beta-actin promoter expression vector. Results are representative of 3 independent explants. **B)** pt"GNAI2" immunolabeling of the adult gallbladder epithelium (maximum projection) where specific *Gnai1* expression was reported in the *Gnai1*<sup>tm1a(EUCOMM)Wtsi</sup> mouse strain (International Mouse Phenotyping Consortium; mousephenotype.org). Apical signals are reduced in *Gnai1* mutants, confirming that the

pt"GNAI2" antibody can detect endogenous GNAI1. **C-D)** GNAI (scbt"GNAI3" antibody; see Fig. 4) immunolabeling in P5 *Gnao1* constitutive (C) and *Atoh1-Cre; Gnao1$^{flox}$* conditional (D) mutants at the cochlear base. Enrichment of GNAI (C-D) and GPSM2 (D) at the bare zone and stereocilia tips is unchanged in absence of GNAO. **E)** GNAO immunolabeling at the P3 cochlear mid produces signals that are unspecific since they are unchanged in conditional *Gnao1* mutants. Scale bars are 10μm (A, C-E) and 25μm (B).

**Figure 5. GNAI2 only partially rescues loss of GNAI3 in individual postnatal hair cells. A-B)** GNAI (pt"GNAI2" antibody; see Fig. 4) and GPSM2 co-immunolabeling in P0 *Gnai3ᵐᵉᵒ* (A) and *Gnai1ᵐᵉᵒ; Gnai3ᵐᵉᵒ* (B) animals at the cochlear base. Boxed regions are magnified on the right. In both mutants, incomplete GNAI patterns are observed at the bare zone (arrow) and stereocilia tips (arrowheads). Remaining GNAI signals must reflect GNAI2 in (B). **C)** Correlation plot of GNAI signal intensity at the bare zone and tips in half-OHC at the P2 cochlear mid. Presence or absence of GNAI is remarkably correlated spatially between bare zone and tips in the same half-OHC. N=3 animals, n=37 OHC,

Pearson correlation with best fit (red line; plot for control littermates can be found in Fig. 5 Supp. 1A). **D-E)** GNAI (pt"GNAI2" antibody) immunolabeling in P6 *Gnai1^{neo}; Gnai3^{neo}* animals. Boxed IHC regions are magnified below. Loss of GNAI2 progresses with HC differentiation, with largely absent IHC signals at the P6 cochlear base (D) but partial rescue on one side of the cell at the P6 mid (E), as observed at the cochlear base at P0 (A-B). **F)** ZO1 (apical junctions) and pericentrin (PCNT; basal body) immunolabeling in P8 OHC (maximum projection). The position of the basal body is used to determine the vertex (middle) of the original hair bundle and to draw a radial line separating each OHC into two halves. **G)** The length of each hair bundle wing (y axis) is graphed in relation to the corresponding apical surface area (x axis) in the same half-OHC. Truncated OHC wings correlate with reduced apical membrane area on the same side. N=3 animals, n=58 OHC, Pearson correlation with best fit (red line; plot for control littermates can be found in Fig. 5 Supp. 1C). A.U., arbitrary unit. Scale bars are 10μm.

**Figure 5 Supplement 1. GNAI2 fully rescues loss of GNAI3 at embryonic stages and is still detected in low amounts at stereocilia tips in adult hair cells lacking GNAI3. A)** Control correlation plot for *Gnai1; Gnai3* double mutants in Fig. 5C. GNAI signal intensity at the bare zone and stereocilia tips in control half-HC at the P2 cochlear base. N=3 animals, n=40 OHC, Pearson correlation with best fit (red line). The graph shows relatively uniform GNAI enrichment in each sub-cellular compartment, unlike in *Gnai1; Gnai3* double mutants (Fig. 5C). **B)** GNAI (pt"GNAI2" antibody; see Fig. 4) and GPSM2

co-immunolabeling at the E18.5 5 cochlear base. As sole remaining GNAI, GNAI2 can still encompass the entire bare zone in *Gnai1; Gnai3* double mutants and no obvious hair bundle defects are observed, unlike at P0 (Fig. 5A-B). The boxed region is magnified to the right. **C)** Control correlation plot for *Gnai1; Gnai3* double mutants in Fig. 5G. The position of the basal body was used to determine the vertex (middle) of the hair bundle and to draw a radial line separating each OHC into two halves. The length of each hair bundle wing (y axis) is graphed in relation to the corresponding apical surface area (x axis) in the same half-OHC. The plot shows relatively uniform wing lengths and half apical areas in littermate controls, unlike in *Gnai1; Gnai3* double mutants (Fig. 5G). N=3 animals, n=52 OHC, Pearson correlation with best fit (red line). **D-E)** GNAI (pt"GNAI2" antibody; see Fig. 4) immunolabeling in P28 adults (maximum projections). As sole remaining GNAI, GNAI2 can still be detected at low levels at stereocilia tips in *Gnai1; Gnai3* double mutants at the cochlear base, mid and apex (arrowheads). E shows higher magnification views of a single OHC1 (left) and IHC (right) at the cochlear mid. A.U., arbitrary unit. Scale bars are 10µm (B, D) and 5 µm (E).

E17.5

**A** control · *FoxG1-Cre; Gnai2^{del/del}; Gnai3^{flox/flox}*

**B** OHC1 · IHC

**C** control · *FoxG1-Cre; DIO-ptxA* · OHC1 · IHC

**D** E17.5 bare zone surface area

*FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}* · *FoxG1-Cre; DIO-ptxA*

controls · mutants

P0

**E** control · *FoxG1-Cre; Gnai2^{del/del}; Gnai3^{flox/flox}*

**F** OHC1 · IHC

**G** *DIO-ptxA* · *FoxG1-Cre; DIO-ptxA* · OHC1 · IHC

**H** P0 bare zone surface area

*FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}* · *FoxG1-Cre; DIO-ptxA*

controls · mutants

**Figure 6. Delayed bare zone expansion and severely dysmorphic hair bundles in absence of GNAI2 and GNAI3. A-B)** GNAI (scbt"GNAI3" antibody, see Fig. 4) and acetylated tubulin (AcTub; kinocilium) co-immunolabeling at the E17.5 cochlear base. Note how F-actin labeling (phalloidin) reveals a polarized bare zone marked by GNAI (arrows) in control but not in *Gnai2; Gnai3* double mutants. **C)** GPSM2 and AcTub co-immunolabeling at the E17.5 cochlear base. In contrast to *Gnai2; Gnai3* double mutants (A-B), *FoxG1-Cre; DIO-ptxA* mutants have a polarized bare zone (arrows) despite OHC1-2 adopting a reversed orientation (V brackets indicate OHC1 orientation). GPSM2 marks the bare zone in controls and is reduced in mutants. **D)** Graphs of bare zone surface area in E17.5 HC at the cochlear base. *FoxG1-Cre; Gnai2$^{del}$; Gnai3$^{flox}$*: controls (*Gnai2$^{del/+}$; Gnai3$^{flox/+}$* and *Gnai2$^{del/+}$; Gnai3$^{flox/flox}$*) N=3 animals, n=19 IHC, 23 OHC1, 19 OHC2, 21 OHC3; mutants N=3, n=23 IHC, 24 OHC1, 23 OHC2, 24 OHC3. *FoxG1-Cre; DIO-ptxA*: controls (Cre-negative *DIO-ptxA*) N=3, n=18 IHC, 18 OHC1, 21 OHC2, 18 OHC3; mutants N=3, n= 21 IHC, 20 OHC1, 21 OHC2, 9 OHC3. **E-G)** Pericentrin (PCNT) and AcTub co-immunolabeling at P0 at the cochlear base. Unlike at E17.5 (A-B, D), most P0 *Gnai2; Gnai3* double mutant HC have a bare region (E-F, arrows). This bare region is unpolarized and its abnormal shape reflects aberrant stereocilia distribution. In sharp contrast, ptxA mutants have normally shaped hair bundles and bare zones despite OHC1-2 adopting a reversed orientation (G). **H)** Graphs of bare zone surface area in P0 HC at the cochlear base. *FoxG1-Cre; Gnai2$^{del}$; Gnai3$^{flox}$*: controls (*Gnai2$^{del/+}$; Gnai3$^{flox/+}$*, *Gnai2$^{del/+}$; Gnai3$^{flox/flox}$* and *FoxG1-Cre; Gnai2$^{del/+}$; Gnai3$^{flox/+}$*) N=3, n=19 IHC, 23 OHC1, 21 OHC2, 21 OHC3; mutant N=3, n=21 IHC, 22 OHC1, 24 OHC2, 24 OHC3. *FoxG1-Cre; DIO-ptxA:* controls (Cre-negative *DIO-ptxA*) N=3, n=15 IHC, 23 OHC1, 18 OHC2, 15 OHC3; mutants N=3, n= 16 IHC, 24 OHC1, 18 OHC2, 19 OHC3. D, H: Nested (hierarchical) t-test sorted by animal; p<0.01 **, p<0.05 *; non-significant p-values are indicated. All *ptxA* samples are heterozygotes (*R26$^{DIO-ptxA/+}$*). Scale bars are 10μm (A, C (OHC), E, G (OHC)) and 5μm (B, C (IHC), F, G (IHC)).

**Figure 6 Supplement 1. Validation of the *Gnai3^flox* and *DIO-ptxA* mouse strains. A)** GNAI (pt"GNAI2" antibody; see Fig. 4) immunolabeling at the P0 cochlear base. Note how in absence of Cre recombinase, *Gnai3^flox/flox* HC have normal hair bundles and normal GNAI enrichment at the bare zone and stereocilia tips. In contrast, *Atoh1-Cre* produces incomplete GNAI enrichment (arrows; compare with Fig. 4D) and shortened stereocilia typical of constitutive *Gnai3* mutants. Boxed IHC regions are magnified below. **B-D)** Comparison of the two *ptxA* strains used in this study (see also Fig. 2 Supp. 1C-D). PtxA expression with the *Atoh1-Cre* driver produces comparable P4 apical HC defects in the *LSL-myc:ptxA* and *DIO-ptxA* strains (B). "Escaper" OHC1 that are not inverted in orientation (arrows) likely do not express Cre, or did not undergo recombination. **C)** When *Atoh1-Cre* induces ptxA expression in post-mitotic HC, the number of escaper OHC1 is higher in the *DIO-ptxA* strain. This is likely because *lox*-based genomic deletions (LSL) are favored over genomic inversions (DIO; see Fig. 2 Supp. 1C-D). **D)** In contrast to *Atoh1-Cre*, an earlier Cre driver (*FoxG1-Cre)* shows virtually no escapers in either ptxA strain. N and n indicate the number of animals and OHC1 analyzed, respectively. All *ptxA* samples in the study are heterozygotes (*R26^LSL-myc:ptxA/+* or *R26^DIO-ptxA/+*), and controls are Cre-negative heterozygotes. Scale bars are 10μm (A, B), 20μm (D).

**Figure 7. Loss of GNAI2 and GNAI3 provokes hair cell eccentricity defects absent in *ptxA* mutants. A-B)** Graphs of HC eccentricity representing the position of the basal body as a ratio of the radius (BB/r, top diagram). Data cover E17.5 mid and base and P0 apex, mid and base cochlear positions for each HC type. HC were considered near-symmetrical when their eccentricity ratio was lower than 0.25 (red zone). Only *FoxG1-Cre; Gnai2^{del}; Gnai3^{flox}* mutants harbor symmetrical HC. Their

proportion is indicated in the bar graphs on the right (A). Overall, the proportion of symmetrical cells tends to decrease in maturing OHC but remains high or increases in IHC. At least 3 animals and 39 cells per HC type are represented for each stage, cochlear position and genotype (detailed in Source Data file). Controls for *FoxG1-Cre; Gnai2$^{del/del}$; Gnai3$^{flox/flox}$* are *Gnai2$^{del/+}$; Gnai3$^{flox/+}$*, *Gnai2$^{del/+}$; Gnai3$^{flox/flox}$*, *FoxG1-Cre; Gnai2$^{del/+}$; Gnai3$^{flox/+}$* and *FoxG1-Cre; Gnai2$^{del/+}$; Gnai3$^{flox/flox}$*. Controls for *FoxG1-Cre; DIO-ptxA* are Cre-negative *DIO-ptxA* heterozygotes. Nested (hierarchical) t-test sorted by animal; p<0.0001 ****, p<0.01 **, p<0.05 *; non-significant p-values are indicated. Detailed cohort sizes and statistics can be found in Source Data file.

**Figure 8. Loss of GNAI2 and GNAI3 recapitulates hair cell orientation defects observed in *ptxA* mutants. A-D)** Circular histograms showing HC orientation (α) based on the position of the basal body (purple dot in top diagram) at the stage and cochlear position indicated. 0° is towards the cochlear base and 90° is lateral. HC represented have an eccentricity greater than 0.4 in A-B (E17.5), and 0.25 in C-D (P0) (see Fig. 7). Histograms show frequency distribution (10° bins) and red radial lines and arcs respectively indicate circular mean and circular mean deviation. In control cochleae, HC are tightly oriented laterally (90°) except for OHC3 that show a slight bias towards the cochlear apex (180°). As reported previously, ptxA expression inverts OHC1 and OHC2, and results in imprecise lateral orientation of OHC3. This phenotype is recapitulated in *Gnai2; Gnai3* double mutants with a

delay (compare least mature E17.5 mid (A) and most mature P0 base (D)). IHC also show severe misorientation in *Gnai2; Gnai3* double mutants, unlike in *ptxA* mutants. n, HC number in N=3-4 animals. Controls for *FoxG1-Cre; Gnai2^{del/del}; Gnai3^{flox/flox}* are *Gnai2^{del/+}; Gnai3^{flox/+}*, *Gnai2^{del/+}; Gnai3^{flox/flox}*, *FoxG1-Cre; Gnai2^{del/+}; Gnai3^{flox/+}* and *FoxG1-Cre; Gnai2^{del/+}; Gnai3^{flox/flox}*. Data at the P0 cochlear mid position can be found in Fig. 8 Supp. 1A. Histograms for littermate controls of *FoxG1-Cre; DIO-ptxA* mutants can be found in Fig. 8 Supp. 2. Detailed cohorts and results can be found in Source Data file.

hair cell orientation

**Figure 8 Supplement 1. Loss of GNAI2 and GNAI3 recapitulates hair cell orientation defects observed in *ptxA* mutants. A)** P0 mid-cochlea histograms that complement P0 apex and P0 base histograms in Fig. 8C-D. **B)** Comparison of OHC1 and OHC2 orientation in *FoxG1-Cre; Gnai2del/del;*

*Gnai3^(flox/flox)* (left diagrams; identical to Fig. 8A, C-D), *Atoh1-Cre; DIO-ptxA* (center diagrams) and *FoxG1-Cre; DIO-ptxA* mutants (right diagrams; identical to Fig. 8A, C-D) at the E17.5 base, P0 apex and P0 base. Post-mitotic Cre expression in *Atoh1-Cre* HC results in delayed *ptxA* expression and delayed inversion of OHC2 compared to earlier *FoxG1-Cre* expression in progenitor cells. This outcome is reminiscent of *Gnai2; Gnai3* mutants. At the E17.5 base and P0 apex (less mature), most OHC2s are inverted in the *FoxG1-Cre; DIO-ptxA* model only. At the P0 cochlear base (more mature), OHC2s are largely inverted in *Gnai2; Gnai3* mutants and in the *Atoh1-Cre; DIO-ptxA* model as well. Circular histograms show HC orientation ($\alpha$) based on the position of the basal body (purple dot in top diagram) at the stage and cochlear position indicated. 0° is towards the cochlear base and 90° is lateral/abneural. HC represented have an eccentricity greater than 0.4 (E17.5) or 0.25 (P0) . Histograms show frequency distribution (10° bins) and red radial lines and arcs respectively indicate the circular mean and the circular mean deviation. n, HC number in N=3-4 animals. Littermate control histograms for *Atoh1-Cre; DIO-ptxA* mutants (B) can be found in Fig. 8 Supp. 2F.

hair cell orientation

**A** E17.5 mid

**B** E17.5 base

**C** P0 apex

**D** P0 mid

**E** P0 base

**F**

**Figure 8 Supplement 2. Littermate control histograms for *ptxA* mutants. A-E)** Histograms showing littermate controls for *FoxG1-Cre; DIO-ptxA* mutants in Fig. 8 (A-C, E) and in Fig. 8 Supp. 1A (D; P0 mid), and littermate controls for *Atoh1-Cre; DIO-ptxA* mutants in Fig. 8 Supp. 1B (F). Note that mutant histograms are identical to Fig. 8 or Fig. 8 Supp. 1. Circular histograms show HC orientation ($\alpha$) based on the position of the basal body (purple dot in top diagram) at the stage and cochlear position indicated. 0° is towards the cochlear base and 90° is lateral/abneural. HC represented have an eccentricity above 0.4 at E17.5 and above 0.25 at P0. Histograms show frequency distribution (10° bins) and red radial lines and arcs respectively indicate the circular mean and the circular mean deviation. n, HC number in N=3-4 animals. Controls for *FoxG1-Cre; DIO-ptxA* and *Atoh1-Cre; DIO-ptxA* are Cre-negative *DIO-ptxA* heterozygotes*.* Detailed cohorts and results can be found in Source Data file.

**Figure 9. Summary and model: three roles validated *in vivo* for GNAI proteins during hair cell polarized morphogenesis.** GNAI proteins are required for the off-center migration of the basal body, and thus HC symmetry breaking (a). This activity is distinct from defining binary HC orientation downstream of GPR156 (b), as *Gpr156* mutant HC have off-center basal bodies and normal hair bundles. Finally, GNAI partner with GPSM2 to shape the hair bundle and elongate stereocilia (c). The specific identity and the dosage of GNAI proteins required differ for each role. GNAI3 is the principal architect of hair bundle development (c), with GNAI2 plays an important but progressively waning role. High amounts of GNAI3/GNAI2 are required with GPSM2. A lower dose of GNAI/O proteins is required for embryonic role, and we speculate that GNAI2 and GNAI3 play a prominent role for symmetry-breaking (a) whereas any GNAI/O protein may effectively signal downstream of GPR156 for proper HC orientation (b). A GNAI/O regulator for symmetry breaking remains to be identified.

**Supplementary Table 1. Mouse strain details.**

| Gene or model | Official name | Name in the study | MGI # | Reference | Allele description | Genetic background | Genetic background in the study | % surviving mutants (expected 25%) | % surviving mutants at weaning age |
|---|---|---|---|---|---|---|---|---|---|
| Gnai1 | Gnai1^tm1Lbi | Gnai1^neo | | Jiang et al, 2002 PMID: 36092739 | neo cassette inserted in exon 3 | | 129S1/SvImJ: C57BL/6J mix | *from het x het (expected 25%)* 16.4% (12 of 73 animals) | *from hom x het (expected 50%)* 45.8% (49 of 107 animals) |
| Gnai2 | Gnai2^em1Btar | Gnai2^del | | this study | CRISPR/Cas9 deletion of exons 2-3-4 | C57BL/6J | | 9.9% (15 of 152 animals) | 9.6% (5 of 52 animals) |
| | | | | | | C57BL/6J: FVB/NJ mix | | 19.7% (12 of 61 animals) | 46.7% (50 of 107 animals) |
| Gnai3 | Gnai3^tm1Lbi | Gnai3^neo | | Jiang et al, 2002 PMID: 36092739 | part of intron 5 and exon 6 replaced with a neo cassette | | 129S1/SvImJ: C57BL/6J mix | 30% (12 of 40 animals) | 42% (63 of 150 animals) |
| | | Gnai3^flox | | this study | floxed exons 2-3 | C57BL/6J | | | |
| Gnao1 | Gnao1^tm1Lbi | Gnao1^neo | 2152685 | Jiang et al, 1998 PMID: 9501252 | neo cassette inserted in exon 6 | 129S1/SvImJ | | *from het x het (expected 25%)* 6.7% (3 animals out of 45) | |
| | Gnao1^tm1c(EUCOMM)Hmgu | Gnao1^flox | from 4456727 | KOMP | floxed exon 3 - *derived from Gnao1tm1a(EUCOMM)Hmgu via FLP-based removal of LacZneo cassette* | C57BL/6N | | *from Atoh1-Cre; het x het (expected 12.5% cKO)* 40% (10 Atoh1-Cre; flox hom of 25 animals) | *from Atoh1-Cre; hom x het (expected 25% cKO)* 30.7% (4 Atoh1-Cre; flox hom of 13 animals) |
| Gnai1; Gnai2 | Gnai1^tm1Lbi ; Gnai2^em1Btar | Gnai1^neo ; Gnai2^del | | Jiang et al, 2002 PMID: 36092739; this study | | | 129S1/SvImJ: C57BL/6J:FVB/NJ mix | *from het; het x het; het (expected 6.25%)* 5.8% (7 of 121 animals) | *from hom; het x hom; het (expected 25% dKO)* 6.9% (2 of 29 animals) |
| Gnai1; Gnai3 | Gnai1^tm1Lbi ; Gnai3^tm1Lbi | Gnai1^neo ; Gnai3^neo | 5560183 | Jiang et al, 2002 PMID: 36092739 | | | 129S1/SvImJ | *from hom; het x hom; het (expected 25% dKO)* 22.2% (20 of 90 animals) | *from hom; hom x hom; het (expected 50% dKO)* 43.2% (60 of 137 animals) |
| FoxG1-Cre; Gnai2; Gnai3 | | Gnai2^del ; Gnai3^flox | | this study | | | 129S1/SvImJ: C57BL/6J:FVB/NJ mix | *second generation crosses* 0% (0 out of 149 animals) | *first generation crosses* 0.3% (1 out of 333 animals) |
| ptxA | Gt(ROSA)26Sor^tm1(ptxA)Cgh | LSL-ptxA:myc | 3776029 | Regard et al, 2007 PMID: 17992256 | loxP-stop-loxP cassette before ptxA:myc | C57BL/6:129SvJ | | | |
| | Gt(ROSA)26Sor^em1(ptxA)Btar | LSL-myc:ptxA | 6163665 | Tarchini et al, 2016 PMID: 27660326 | loxP-stop-loxP cassette before myc:ptxA | C57BL/6J | | | |
| | | DIO-ptxA | | this study | double inverted lox sites flanking ptxA | C57BL/6J | | | |
| Atoh1-Cre | Tg(Atoh1-cre)1Bfri | Atoh1-Cre | 3775845 | Matei et al, 2005 PMID: 16145671 | randomly inserted transgene with Atoh1 enhancer | (C57BL/6 x CBA)F1 | | | |
| FoxG1-Cre | Foxg1^tm1(cre)Skm | FoxG1-Cre | 1932522 | Hebert et al, 2000 PMID: 10837119 | Cre knock-in in the FoxG1 gene | 129P2/OlaHsd-Hprtb-m3 | | | |

| Strain | | Primer sequence | Bands size (bp) | |
|---|---|---|---|---|
| *Gnao1 flox* | forward | 5' - TGGGTTTGATCCACAAGAGCGG  - 3' | | |
| | forward | 5' - AGACGTGAAGCAGTACAAGC - 3' | | |
| | reverse | 5' - TCTCCTCAGTAGGGAAGATGGC - 3' | | |
| | | | wt: 938; flox: 1119 | wt: 387; flox: 421 |
| *Gnao1 neo* | forward | 5' - TGGTGACCCATTCTGACTTC  - 3' | | |
| | forward | 5' - TCGCCTTCTATCGCCTTCTTGACG  - 3' | | |
| | reverse | 5' - TTGGCAGAGAAAGCCCAAAG  - 3' | | |
| | | | wt: 836 | neo: 461 |
| *Gnai1 neo* | forward | 5' - TCTAGGTGCTGGGGAATCTG  - 3' | | |
| | reverse | 5' - GAGACGTGCTACTTCCATTTGTC - 3' | | |
| | reverse | 5' - TGAGCTTGTAGGCACAGAGG - 3' | | |
| | | | wt: 1004 | neo: 580 |
| *Gnai2 del* | forward | 5' -GCACCATCGTCAAGCAGATG  - 3' | | |
| | forward | 5' - ATGGGTGGTTGTGAGCCATC - 3' | | |
| | reverse | 5' - GAGCTGGTATTCTCGTGAGC - 3' | | |
| | | | wt: 728 | wt: 1195; neo: 515 |
| *Gnai3 neo* | forward | 5' - CTTCTATCGCCTTCTTGACG - 3' | | |
| | forward | 5' -ACTCTGCTGACTTTGGCACC - 3' | | |
| | reverse | 5' - CTTTCATGCATTCGGTTCTG - 3' | | |
| | | | wt: 762 | neo: 450 |
| *Gnai3 flox* | forward | 5' - ATGAGGACGGCTATTCAGAG  - 3' | | |
| | reverse | 5' - CATCCTACTTCACAAACTGC - 3' | | XhoI digestion |
| | | | wt: 700; flox: 747 | wt: 700; flox: 342 + 404 |
| *LSL-myc:ptxA* | forward | 5' - AAAGTCGCTCTGAGTTGTTAT  - 3' | | |
| | reverse | 5' - GCGAAGAGTTTGTCCTCAACC - 3' | | |
| | reverse | 5' - GGAGCGGGAGAAATGGATATG - 3' | | |
| | | | wt: 603 | ptxA: 250 |
| *DIO-ptxA* | forward | 5' - GCACTTGCTCTCCCAAAGTC  - 3' | | |
| | reverse | 5' - GGAGCGGGAGAAATGGATATG - 3' | | |
| | reverse | 5' - CGTCATTGACGTCAATAGGG - 3' | | |
| | | | wt: 617 | ptxA: 677 |
| *Atoh1-Cre* | forward | 5' - AGAGCGGCTGACAATAGAGG  - 3' | | |
| | reverse | 5' - TTTCCATGAGTGAACGAACC - 3' | | |
| | | | wt: no band | cre: 850 |
| *FoxG1-Cre* | forward | 5' - GCGACAAGAAGAACGGCAAG - 3' | | |
| | forward | 5' - GCAAGAACCTGATGGACATG - 3' | | |
| | reverse | 5' - AAGCACTTGTTGAGGGACAG - 3' | | |
| | reverse | 5' - TTTCCATGAGTGAACGAACC - 3' | | |
| | | | wt: 202 | cre: 381 |

**Supplementary Table 2. Genotyping strategies.**