Title: Ongoing convergent evolution of a selfing syndrome threatens plant-pollinator interactions

Authors: Samson Acoca-Pidolle*, Perrine Gauthier, Louis Devresse, Virginie Pons, Pierre-Olivier Cheptou

Affiliation: Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), University of Montpellier, CNRS, EPHE, IRD, 34293 Montpellier, France

*Corresponding author, email address: samson.acoca-pidolle@cefe.cnrs.fr

ORCID: SAP https://orcid.org/0009-0002-8456-5588; PG https://orcid.org/0000-0001-7926-4647; LD https://orcid.org/0009-0006-0613-4343; POC https://orcid.org/0000-0002-5739-5176

Manuscript information:

- 800 words for Introduction, 2387 words for Materials and Methods, 1418 words for Results, 616 words for Discussion
- 3 figures in colour and 2 tables
- Supporting information includes three figures and two tables.

Summary

• Plant-pollinator interactions have evolved early in the angiosperm radiation.
Ongoing environmental changes are however leading to pollinator declines that
may cause pollen limitation to plants and change the evolutionary pressures shaping
plant mating systems.

• We used the resurrection ecology methodology to contrast ancestries and
contemporary descendants in four natural populations of the field pansy (*Viola*
arvensis) in the Paris region (France), a depauperate pollinator environment. We
combine population genetics analysis and phenotypic measurements and behavorial
tests on a common garden experiment.

• Population genetic analyses reveal 27% increase in realized selfing rates in the
field during this period. We documented trait evolution towards smaller and less
conspicuous corollas, reduced nectar production and reduced attractiveness to
bumble bees, with these trait shifts convergent across the four studied populations.

• We demonstrate the rapid evolution of a selfing syndrome in the four studied plant
populations, associated with a weakening of the interactions with pollinators over
the last three decades. This study demonstrates that plant mating systems can evolve
rapidly in natural populations in the face of ongoing environmental changes. The
rapid evolution towards a selfing syndrome may in turn further accelerate pollinator
declines, in an eco-evolutionary feedback loop with broader implications to natural
ecosystems.

Key words: Rapid evolution, mating system, selfing syndrome, plant-pollinator interactions, resurrection ecology, *Viola arvensis*

Introduction

The astonishing diversification of angiosperms, which started 100 billion years ago
(Benton et al. 2022), is commonly thought to have been stimulated and hastened by the
diversity of plant-pollinator interactions (Whittall and Hodges 2007). More than 80% of
extant angiosperms rely on animals for pollination (Ollerton et al. 2011), thus plants and

pollinators have a close relationship. Yet, previous studies have shown a decline of
pollinators (including bees, Anthopila) is ongoing across the world (Goulson et al. 2015,
Hallmann et al. 2017, Grab et al. 2019). A lack of pollinators can in turn directly affect
plant reproduction, and indeed parallel declines in insect-pollinated plants and insects have
been documented, over the last 50 years (Biesmeijer et al. 2006). At evolutionary time
scales, pollen limitation has been shown to represent a major selective pressure in the
evolution of self-fertilization (selfing) in theoretical evolutionary models (Lloyd 1992,
Thomann et al. 2013). In recent years, such a shift towards selfing has been experimentally
reproduced in the absence of pollinators (Bodbyl Roels and Kelly 2011, Gervasi and
Schiestl 2017, Ramos and Schiestl 2019) and illustrated in natural populations
impoverished in pollinators (Brys and Jacquemyn 2012). This evolution of selfing is
commonly associated with changes in reproductive traits such as a decrease in the distance
between male and female organs (herkogamy) (Bodbyl Roels and Kelly 2011, Brys and
Jacquemyn 2012, Ramos and Schiestl 2019) but also a decrease in floral traits mediating
plant-pollinator interactions, such as a decrease in corolla and petal size, and decreased
scent and nectar production (Brys and Jacquemyn 2012, Gervasi and Schiestl 2017, Ramos
and Schiestl 2019). A shift in mating system and traits leading to higher selfing, less
conspicuous and attractive flowers has been well described at the phylogenetic level
(Stebbins 1957, Sicard and Lenhard 2011). This set of trait changes characterizes the
evolution of a selfing syndrome (Ornduff 1969, Sicard and Lenhard 2011).

Faced with pollinator declines, natural populations are expected to evolve towards
reduced plant-pollinator interactions and increased autonomous selfing (Thomann et al.
2013). However, demonstrating trait evolution requires distinguishing between the genetic
and the environmental components of phenotypes, which is not straightforward in
longitudinal studies of natural populations. The "resurrection ecology" approach, which
involves growing dormant seeds collected from field populations multiple generations ago
alongside their naturally occurring descendants (Franks et al. 2007, Weider et al. 2018), is
a powerful methodology to detect such evolution (Weider et al. 2018). Indeed, this
approach enables the contrast between traits of ancestral versus recent genotypes in a
common environment, thus revealing the genetic components of temporal trait variation.
Here, we took advantage of an ancestral seed collection of the field pansy (*Viola arvensis*)

collated about two decades ago in the Paris region (France) to test whether this species'
mating system has evolved in response to recent pollinator declines.

The study species is an annual weed of intensive crops (mostly wheat and oilseed
rape) that only produces chasmogamous, zygomorphic, showy flowers and mixed selfing
rates (Scoppola and Lattanzi 2012, Cheptou et al. 2022). The length of the nectar spur and
observations *in natura* are indicative of pollination by long-tongued bees. For example,
Bombus terrestris is one of the wild pollinators observed. The study region has degraded
pollination services compared with the rest of France (Martin et al. 2019). The status of
pollinators or pollination services in such intensive agricultural areas has been reported as
declining in several studies over our timescale (Goulson et al. 2015, Grab et al. 2019, Potts
et al. 2010, Burkle et al. 2013, Raven and Wagner 2021, Janousek et al. 2023). Parts of
Belgium (some 100 to 200 km north of our study area) have experienced similar
agricultural dynamics as in our study region, and studies there indicate that 32.8% of bee
species are threatened or extinct based on comparisons with their presence before and after
1970 (Drossart et al. 2019). These studies also highlighted a more severe decline or higher
vulnerability of long-tongued bees (Biesmeijer et al. 2006, Goulson et al. 2015, Drossart
et al. 2019, Janousek et al. 2023).

In this study, we contrast ancestral (collated in the 1990s to early 2000s) and
contemporary descendants (collated in 2021) from four localities across a range of 100
kilometres in the study region, combining genetic and phenotypic approaches. After a
refresher generation (F1), phenotypic traits were measured in the second generation (F2)
in 2022 in an air-conditioned greenhouse. We quantified the evolution of traits mediating
plant-pollinator interactions as well as other morphological traits not directly involved in
plant-pollinator interactions. We also performed a plant pollinator preference experiment
by monitoring bumblebee visitation in mixed artificial populations containing genotypes
of the two different source generations. Together, these approaches allowed us to document

on the evolution of plant-pollinator interactions and evolution of plant mating system induced in recent environmental changes.

Material and methods

Study system and sampling

Viola arvensis is an annual species of wild pansy (Violaceae). It is commonly found in fields as a weed or in meadows. This species is described as self-compatible with a mixed mating system (Scoppola and Lattanzi 2012, Cheptou et al. 2022). Plants were sampled twice in agricultural fields in four locations of the Parisian Basin (see Table **S1**). The first sampling was done and conserved by the Conservatoire Botanique National de Bailleul for Crouy (Cr) and Lhuys (Lh) in the 1990's and by the Conservatoire Botanique National du Bassin Parisien for Commeny (Co) and Guernes (Gu) in the 2000's. Seeds were collected by sampling fruits on a minimum of 100 mothers picked randomly in each population. The second sampling was performed in exactly the same locations, in an area that overlapped the ancestral sampling area as much as possible, in February 2021 by randomly sampling 40 or more seedlings. The minimal sampling areas was of 2000 m^2 and correspond to cases were all the population is circumscribed on this area. This sampling scheme produced four couples, one for each location, of ancestral (A) and descendant (21) populations.

Experimental design

To avoid maternal and potential storage effects (Franks et al. 2018), a refresher generation (F1) was grown in common garden for plants issued from the different populations. As we have seeds and seedlings, common conditions started at seedling stage. For ancestral populations, in mid-November 2020, seed lots were placed in germination chambers in petri dishes with vermiculite under light at 15°C for 12h and in the dark at 6°C for 12h. Germination rates was of 50% to 65%. 32 seedlings of ancestral and descendant populations were transplanted in late February 2021 in open ground in the Baillarguet site of the experimental platform "Terrains d'Expériences" of the Labex Cemeb, 10 km north of Montpellier, France (43°40′53.92″N, 3°52′28.74″E). A single event of mortality was

observed in Cr21 population. Populations were separated from each other by insect-proof mesh greenhouses (Fig. **S1**). During full blooming, in mid-April, we introduced commercial bumblebee hives, Natupol from Koppert France® (Cavaillon, France), into the greenhouses to produce open-pollinated fruits that we collected.

This refresher generation was used to generate seeds produced in the same environment for the experimental generation (F2). This F1 generation allows us to generate a family design in the F2 generation. We randomly selected 20 mothers per population in the F1 generation to construct 20 families in the F2 generation. Each family is composed of five siblings (Fig. **S1**). In December 2021, 30 seeds per family were placed in germination chambers in petri dishes with vermiculite under light at 15°C for 12h and in the dark at 6°C for 12h. Germination rates per population range between 75% and 90%. Plants were placed in individual pots ($11 \times 11 \times 11$ cm^3) filled with standardized soil in late February and randomized in a greenhouse of the experimental platform "Terrains d'Expériences" of the Labex Cemeb, Montpellier, France. Watering was manual and done when needed. Three families did not germinate enough to obtain 5 plants and thus were composed of one, two or four siblings. In total, 792 plants were followed during this experiment, no early mortality was observed. When all the morphological measurement were performed, plants were moved outside to complete their life cycle.

<u>Population genetics</u>

32 F1-plants per populationwere genotyped using eleven microsatellite markers, 6 design for *Viola arvensis* (Cheptou et al. 2022) and 5 design on *Viola tricolor* (Latron et al. 2018), a closely related species (Scoppola and Lattanzi 2012). DNA was extracted from fresh leaves using Dneasy Plant Mini Kit from QIAGEN®. Analyses of metrics such as allelic richness, expected heterozygosity (H_e), using the R package genepop (Rousset 2023), were performed to look at a change in genetic diversity over time. A drastic decrease in allelic richness (or H_e) is indicative of genetic drift, strong population bottlenecks or a sampling bias. Heterozygosity deficiency (F_{IS}) analyses were performed, using the R package HierFstats (Goudet et al. 2022), on fourteen microsatellite markers. F_{IS} was used to estimate the selfing rate using the following formula: $s_{Fis} = \frac{2F_{IS}}{1 + F_{IS}}$. Confidence intervals

were obtained by bootstraps over loci of population (10,000). To test for a difference in

F_{IS}, we pooled individuals of the same location, independently of their age and performed

bootstraps to reconstruct two populations of 32 individuals. We calculated the F_{IS} of these

two reconstruct populations and reproduce the procedure 10 000 times per locality on each

locality. Then, we recorded the number of cases where the absolute value of the difference

in reconstruct F_{IS} is higher than the true difference in F_{IS} measured. This number divided

by 10 000 gives us a P-value for H_0: "True difference $= 0$". We performed a Fisher's

combined probability test on the P-values obtained for each locality (Fisher 1970) to test a

global increase in selfing rate between A and 2021 populations.

<u>Measurements of plant traits</u>

Plants traits were measured on the test generation (F2). All the morphometric

measurements (Fig. **S2**) were performed by a single experimenter during all the

experimentation, to avoid any experimenter effect, using a digital caliper to 0.01 mm.

Corolla length, labellum width, spur length, sepal length, number of guides and

anthesis duration were measured on the five first opened flowers of the main shoot of each

plant. We chose the first five opened flowers of the main shoot because more exhausted

measurements in the F1 showed that they capture the floral variance of an individual.

Measurements were made day+2 after anthesis, or day+3 when falling on weekends. The

opening of flowers was followed every working day to assess the anthesis duration. Floral

area approximation is the product of corolla length and labellum width. Sepal/corolla

lengths ratio is the ratio of sepal to corolla lengths. Around 4,000 flowers were followed.

Flowering date is the date of anthesis of the first flower of a plant. The number of flowers

opened per plant was counted each working day, from the opening of the first flower, for

one month and half, and two more times with one week and then two weeks delay between

counts. This number was integrated using trapezoidal rule and divided by the duration

between the flowering date of the plant and the last day of measurement, giving the average

floral display. Nectar volume was collected and cumulated from three flowers per plant

using 0.5 μL microcapillaries. It was measured randomly on individuals of the two

populations of the same location simultaneously to avoid differences in nectar production

caused by the time of measurement. Thus, for this measure, Locality effect (see "Statistical analysis") is confounded with the day of measurement. It was measured on half of the plants from each population, representing all the families, thus 400 plants were sampled.

Vegetative traits such as rosette diameter, leaf length and leaf width were measured at the flowering date of the plant. Leaf length and leaf width were measured on the biggest leaf at this time, and the product of these two measures gives leaf area approximation. Biomass was measured on dried naturally dead plants at the end of their life cycle.

Reproductive traits were obtained while counting and weighing all the seeds contained by fruits to the nearest 0.1 mg using a precision balance. When floral and leaf traits had been scored, plants were placed by groups of twenty in insect-proof mesh greenhouses which contained commercial *Bombus terrestris* bumblebee hives, Natupol from Koppert France® (see Bumblebee preferences experiment below). After 2 hours minimum, plants were moved outside in open pollination context. Youngest receptive flower and oldest non-receptive ones on the same shoot were marked before the exposition to pollinator to differentiate fruits issued from flower in zero-pollinator context (self-pollination) versus fruits produced in open pollination. Thus, open pollination fruits are issued from flowers produced later in the shoot. For each plant, just before the dehiscence of the fruit produced by self-pollination, we sampled the two fruits at the same time and recorded the difference in the flower position number separating collected fruits (minimum 1, by definition, to maximum 5, with a mean of 1.6). The sampling of the two fruits was performed at the same time because of the practical difficulty to follow 1,600 fruits independently. Thus, open pollination fruits are not fully developed and comparing weights between open pollination and self-pollination fruits has a slight bias. However, comparing each measure between populations is fully possible while differences in floral positions are distributed randomly between populations. The seed weight reported is the measure of the weight of seeds contained by each fruit divided by the number of seeds counted in the fruit. The ratio of the number of seeds produced by self-pollination to the number of seeds produced by open pollination is a proxy of pollen limitation due to the absence of pollinators, the less this limitation is the more the self-pollination ability is.

Bumblebee preferences experiment

Plants were exposed to commercial bumblebee hives, Natupol from Koppert France®, in
insect-proof mesh greenhouses once all the trait measures had been made. In each insect-
proof greenhouse, we placed, for the entire experiment, a bumblebee hive of around 30
workers with a queen, with nutrient reserves. We randomly disposed plants from the same
location in a group of 20 plants, composed of 10 plants from each ancestral and descendant
population randomly selected in one of the two insect-proof greenhouses. Pots were equally
spaced at 35 cm. The observer, stationed in the entrance of the insect-proof greenhouses,
selected a foraging worker bumblebee, and recorded each flower visit during 10 to 15
minutes. Greenhouse number (1, 2), position in the greenhouse (1-20) and plant identities
were also recorded. Experiments were performed between the 2nd and the 19th of May 2022.
We performed the experiment on 8 groups for each locality but just 6 and 7 observations
were kept for Lhuys and Crouy respectively, because no pollinators foraged enough time
in the discarded observations. On the 580 plants in our dataset, 3 are individuals exposed
two times for experimental purpose. During a 10-15 minutes test, 4.28 plants were not
visited in average and 124 in total for all the tests. In the analyses, we sorted out plants
visited and unvisited (see below). We calculated the proportion of visits per plant as the
ratio between the number of visits on a given plant divided by the total number of visits
during a flight.

Statistical analysis

Changes in traits were evaluated using linear discriminant analysis (LDA) and univariate
linear mixed models (LMMs). We performed the LDA over nine traits supposed to be
linked with reproductive system; corolla length, labellum width, floral area, number of
guides, anthesis duration, flowering date, volume of nectar, average floral display and ratio
of seeds produced in self-pollination to open pollination. For traits with repeated measures,
we used the means of the trait per individual. We normalized values by locality in order to
focus on Age effect. We predefined only the population (8 groups) in the LDA and not the
age. LDA gives linear combinations of traits (discriminant functions) that attempt to
capture differences among groups. In our case, it produced seven discriminant functions.
The two first functions represent 55% and 22% of the variation.

Univariate analyses were conducted using LMMs with each trait as response

variable, Age (ancestral or descendant), Locality (4 localities), and the interaction between

Age and Locality as fixed factors, and Family as random factor. In order to account for

repeated measures on the same individual for floral measurement, we add Individual as

random factor nested in Family. For seeds traits, we accounted for the difference in flower

position between the two fruits collected adding this difference as random factor nested in

family. When significant Age effect, analysis was followed by T-test comparison, on the

means of the trait per individual for repeated measures, of ancestral and descendant

populations of each locality. *p*-values were corrected using Bonferroni correction. Rosette

diameter, leaf length, width and area, nectar volume and ratio of seeds without/with

pollinators were log-transformed to approach normal distribution of residuals. In these

analyses, Age significant effect represents a signal of global evolution of traits between

ancestral and descendant populations, Locality significant effect, represents a difference in

traits between populations. A significant effect of the interaction between Age and Locality

represents a difference in the magnitude or in the sign of the evolution between localities.

In the case of a significant Age effect and a significant interaction effect, if evolution is

consistent (same sign of evolution) among localities and only the magnitude change, we

can hypothesize a directional selective force that selects a trait in a direction convergently

in every locality. In other cases, a significant interaction between Age and Locality

indicates local differences in selective pressures on the traits or alternatively genetic drift.

For bumblebee preference analyses, we first performed a GLMM to test the effect

of Age on the number of unvisited plants (proportions: CoA: 0.24, Co21: 0.35;

CrA: 0.26, Cr21: 0.26; GuA: 0.15, Gu21: 0.18; LhA: 0.13, Lh21: 0.12). We modelled it

with Age and Locality as fixed effect, Family, Greenhouse, Position nested in Greenhouse

and Date as random effects, with binomial distribution and logit link, as response variable

was a Boolean variable with TRUE for visited plants and FALSE otherwise. There was no

significant effect of Age in our dataset (N = 580 plants, estimates: AgeA-Age21 = 0.223,

$P = 0.296$). We performed further analysis with and without unvisited plants (N = 124) and

found similar pattern. To avoid non-normal distribution, we excluded unvisited plants

(N = 124) in the rest of the analysis. We performed a LMM with log-proportion of visits

per plant as response variable (N = 456), and the same structure as for other traits with Age,

Locality, and the interaction between Age and Locality as fixed effect, Family as random effect and we added Greenhouse, Position nested in Greenhouse and Date as random effects. We then performed T-test comparisons between Age for given Locality and corrected *p*-values using Bonferroni correction.

All our statistical analyses were performed using the R software (R Core Team, Version 4.2.1). LDA was performed using the "lda" function implemented in the package "MASS" (Venables and Ripley 2002). LMMs and GLMM were performed using the "lmer" and "glmer" functions implemented in the package "lme4" (Bates et al. 2015).

Results

The localities exhibited no significant differences in allelic richness nor genetic diversity between ancestral and descendant populations based on fifteen microsatellites markers (Table **1**). In the case of a resampling bias of the populations or a strong demographic bottleneck or genetic drift between the two samples, we predicted a decrease in allelic richness and genetic diversity. Our results do not support these *scenarii*. In contrast, we did observe a consistent and significant increase in selfing rates in all four localities between ancestral and descendant populations (Table **1**), with an average and significant increase of 27% (Fisher test, $\chi^2_8 = 40.36$, $P < 0.001$).

Descriptive multivariate analyses performed on eight floral traits associated with the mating system (corolla length, labellum width, floral area, number of guides, anthesis duration, flowering date, volume of nectar and average floral display) and the ability of plant to set seeds in the absence of pollinators (*i.e.* autonomous self-pollination), corrected for differences between localities, showed differentiation between ancestral and descendant populations. We choose these traits because they are usually associated with shifts in mating systems (Bodbyl Roels and Kelly 2011, Sicard and Lenhard 2011, Gervasi and Schiestl 2017, Ramos and Schiestl 2019). The descendent populations and ancestral populations form two distinct clusters in the trait space (Fig. **1**), revealing convergent directional changes of floral traits in all four localities.

Table 1. Allelic richness, genetic diversity and selfing rates in our populations (mean ± SD or [CI]). Average allelic richness and genetic diversity are based on fifteen

microsatellites markers and selfing rate on fourteen. These markers were genotyped on the thirty-two individuals per population of the F1 generation issued from natural populations. For selfing rate, confidence intervals (CI) were obtained by bootstraps on loci (10,000). Fisher's combined probability test gave a significant overall increase in F_{IS} ($\chi^2_8 = 40.36$, $P < 0.001$, see Material and methods).

The first two letters are the name of the locality (Co = Commeny; Cr = Crouy; Gu = Guernes; Lh = Lhuys). "A" ancestral population (collected in 2000 for Co, 1993 for Cr, 2001 for Gu and 1992 for Lh) and "21" descendant population (all collected in 2021).

Population	CoA Co21	CrA Cr21	GuA Gu21	LhA Lh21
Average allelic richness	2.55 ± 1.21 1.73 ± 0.79	3.09 ± 1.81 3.36 ± 1.86	2.64 ± 1.29 3.27 ± 1.56	2.82 ± 1.78 2.18 ± 1.08
Genetic diversity (H_e)	0.302 ± 0.275 0.099 ± 0.133	0.435 ± 0.263 0.437 ± 0.254	0.315 ± 0.211 0.327 ± 0.255	0.315 ± 0.291 0.245 ± 0.225
Selfing rate [95% CI] derived from heterozygosity deficiency, F_{IS}	0.77 [0.68-0.82] 0.89 [0.85-1]	0.66 [0.60-0.76] 0.70 [0.65-0.76]	0.44 [0.34-0.61] 0.81 [0.73-0.86]	0.33 [0.21-0.47] 0.88 [0.81-0.94]

Figure 1. Multivariate differences between traits of ancestral and descendant populations of *Viola arvensis*. Results of a multivariate analysis (linear discriminant analysis) performed on nine traits and 357 plants (almost equally distributed among groups; N_{min} = 36 individuals; 7 points not represented because out of scale). The first two functions represent 55% and 22% of the variation respectively.

The first two letters are the name of the locality (Co = Commeny; Cr = Crouy; Gu = Guernes; Lh = Lhuys). "A" (▲) ancestral population (collected in 2000 for Co, 1993 for Cr, 2001 for Gu and 1992 for Lh) and "21" (●) descendant population (all collected in 2021). Enlarged symbols are centroids of groups.

Linear models on the nineteen traits measured revealed consistent patterns (Table **2**). We detected a significant difference between ancestral and descendant populations for all the floral traits except sepal length (Table **2**, Table **S2**). Plants have evolved a 10% decrease in floral area (Fig. **2a**) through a combined decrease in the labellum width and flower length (Table **2**). Flowers also have fewer nectar-guides (Fig. **2b**), which are assumed to promote recognition of the path to floral rewards (a correlation between the

number of nectar-guides and the rate of visits inducing pollination has been reported for

the South African iris, *Lapeirousia oreogena* (Hansen et al. 2012)). In contrast, none of the

morphological traits presumed to be unlinked to pollinator attraction (vegetative traits and

sepal length) showed consistent changes (Fig. **2c**, Fig. **S3**, Table **2**). Based on the

resurrection approach, patterns of trait changes between ancestral and descendant

populations provide a signature of mating system and associated trait evolution. The lack

of changes in vegetative traits reduces the likelihood that these changes are due to

inbreeding depression (unlikely that it would affect only floral traits), especially because

late acting traits such as biomass exhibited no changes (Winn et al. 2011). In addition,

evolution in petal length but not in sepal (Fig. **S3**) or leaf length (Fig. **2c**), despite their

homologous origins (Breuninger and Lenhard 2010), provides strong evidence for a

specific selection against costly traits that mediate interactions with pollinators.

Table 2. Linear models: effects of time, locality, and their interaction on the changes

of floral, vegetative and reproductive traits. Values in parentheses are sample sizes.

All traits are shown and the degree of significance of each effect on each trait is reported

(LMM, *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$, • $P < 0.1$, n.s. $P > 0.1$, see Table **S2** for

full statistical values). (+) a positive effect (increase), (−) a negative effect (decrease), (~)

a difference of effect sign between localities (see Table **S2** for exact values).

Trait (sample size)	Age	Locality	Interaction of Locality and Age
Floral traits			
Corolla length (3955)	***(—)	**	***
Labellum width (3955)	***(—)	***	**
Floral area approximation (3955)	***(—)	***	**
Number of guides (3993)	***(—)	***	***
Average floral display (789)	**(—)	***	*
Anthesis duration (3991)	**(~)	***	***
Spur length (3954)	**(—)	***	**
Sepal length (3955)	n.s.	***	n.s.
Sepal/corolla lengths ratio (3955)	***(+)	***	**
Flowering date (792)	n.s.	n.s.	n.s.

Vegetative traits			
Rosette diameter (792)	n.s.	n.s.	n.s.
Leaf length (792)	n.s.	***	*
Leaf width (792)	n.s.	***	*
Leaf area approximation (792)	n.s.	***	*
Biomass (777)	n.s.	n.s.	n.s.
Reproductive traits			
Log-ratio of the number of seeds without/with pollinators (693)	•(+)	***	n.s.
Weight per seed with pollinators (656)	*(—)	***	n.s.
Weight per seed without pollinators (701)	**(—)	***	***
Plant-pollinator interaction traits			
Nectar volume (400)	***(—)	***	n.s.
Bumblebee visit preferences (456)	***(—)	n.s.	n.s.

Regarding reproductive traits, our results revealed a marginally significant

($\chi^2_1 = 3.201$, $P = 0.074$) increase over time in the ability to set seeds in the absence of

pollinators (Fig. **2d**) and a significant decrease in seed weight (Table **2**). Seed weight

represents maternal investment in individual progeny. This shift towards increased

reproductive insurance is associated with reduced seed investment, as observed in

previous studies (De Jong et al. 2005).

Our study also reveals that both locality and the interaction between locality and

age (ancestors or descendants) have a strong effect on trait variations. The locality effects

indicate spatial differentiation in measured traits, whereas the locality×age interaction

reveals a site dependent evolution. The latter interaction effects were different in

magnitude, but not in sign, between ancestors and descendants for all the traits, except

anthesis duration (Table **2**). Hence, evolution towards a selfing syndrome in the four

localities exhibits the same directional change *i.e.* evidence of convergent shifts in the four

independent populations (Fig. **1**) while differing in magnitude (indicative of slight

differences in selective pressures between localities).

Figure 2. Evolutionary changes in floral, reproductive and vegetative traits. We measured floral traits (**a** and **b**) in the first five developed flowers per individual; N ≈ 4,000). (**a**) Floral area (multiplication of labellum width × corolla length). (**b**) Number of attractive guides. (**c**) Rosette diameter, measured on each plant at the start of flowering. (**d**) Log-ratio of seeds produced in self-pollination compared to open pollination is a proxy of selfing ability, measured by collecting one fruit in self-pollination and one in open pollination per plant (N = 693).

The first two letters are the name of the locality (Co = Commeny; Cr = Crouy; Gu = Guernes; Lh = Lhuys). "A" (▲) ancestral population (collected in 2000 for Co, 1993 for Cr, 2001 for Gu and 1992 for Lh) and "21" (●) descendant population (all collected in 2021).

We found a consistent decrease of 20% on average in rewarding traits, as
represented by measurements of nectar volume (Fig. **3a**, Table **2**). Because floral nectar
production is costly (Pyke 1991) and only dedicated to rewarding pollinators, it is key to
maintenance of stable plant-pollinator interactions(Brandenburg et al. 2012). A decrease
in nectar production in our study species is thus evidence of a weakening of the interaction
between the plant and its pollinators.

Fig. 3. Change in plant-pollinator interactions. (**a**) Nectar production measured as the
sum of the volume in three flowers per plants on fifty plants per population (N = 400). (**b**)
Bumblebee preferences, measured as proportion of visits per plant to a mixed plantation of
10 plants of the ancestral and 10 of the descendant population of a single locality in a
greenhouse, exposed together to bumblebees. We recorded the number of visits to each
plant by a unique flying bumblebee for 10 to 15 minutes in 6 to 8 replicates per location
and divided it by the total number of visits during the flight (only plants visited once are
represented).

The first two letters are the name of the locality (Co = Commeny; Cr = Crouy; Gu
= Guernes; Lh = Lhuys). "A" (▲) ancestral population (collected in 2000 for Co, 1993 for
Cr, 2001 for Gu and 1992 for Lh) and "21" (●) descendant population (all collected in
2021).

We assessed how this evolution impacts plant-pollinator interactions from the point
of view of pollinators through a greenhouse preference experiment involving visits of the

long-tongued bumblebees *Bombus terrestris* to a mixture of ancestral and descendant plants. Bumblebees visited ancestral plants more frequently than they visited descendant plants, showing a preference towards the former that is consistent with their higher attractiveness and reward traits (Fig. **3b**, Table **2**).

Discussion

Overall, we found decreases in floral display, in floral area, in floral attractiveness, and in nectar production, as well as evidence for enhanced selfing and an increase in realized selfing rates that are all consistent with the ongoing evolution of a selfing syndrome (Ornduff 1969, Sicard and Lenhard 2011). Furthermore, we found such patterns to be convergent across the four populations investigated. This study thus provides a unique demonstration for an evolutionary change in plant mating systems over a short-time scale in natural populations. While angiosperm-pollinator interactions have evolved over the long-term, our study shows that current environmental changes can drive a rapid evolution towards a breakdown of such interactions.

Recurrent evolution towards selfing has been repeatedly documented in the wild, but to our knowledge, only over the long-term (Stebbins 1957, Barrett 2002). This evolutionary transition is classically considered to be irreversible (the "selfing as an evolutionary dead-end" hypothesis (Stebbins 1957, Igic and Busch 2013)) and higher extinction rates of such phyla have been reported (Goldberg et al. 2010). Evolution towards selfing could thus be driven by natural selection over the short-term but could impede long-term plant population survival (Cheptou 2019). The increase in selfing rate is indeed expected to have genomic impacts on individuals and populations through inbreeding (homozygosity), reduced effective population sizes and increased genetic drift. Consequently, selfers may have a reduced potential response to natural selection (Glémin and Ronfort 2013, Noël et al. 2017). The short-term evolution of selfing in *Viola arvensis* we document here is no-doubt facilitated by its initial mixed mating system (contrary to self-incompatible plants for example (Thomann et al. 2015)), nonetheless our results illustrate how mixed-mating species may evolve towards a higher reliance on self-pollination and reduced plant-pollinator interactions in the face of global change, in

particular pollinator declines. If this rapid transition towards a selfing syndrome reflects a broader trend among Angiosperms, it may reflect a concerning extinction debt (Goldberg et al. 2010).

Ongoing environmental changes – such as habitat destruction and fragmentation, agricultural land pollution and species' introductions – have been documented as causes of pollinator declines, both by affecting pollinator populations directly (Potts et al. 2010, Goulson et al. 2015), and indirectly through their impacts on plant community composition, affecting the quantity and diversity of nectar resources (Goulson et al. 2015, Baude et al. 2016). In addition, and as we illustrate here, floral nectar production can also evolve rapidly in plant species in response to a relaxed selective pressure associated with pollinator declines, prompting declines in nectar resources of a magnitude similar to those associated with changes in plant community composition (Baude et al. 2016). These decreases in nectar production may then reinforce pollinator declines if nectar levels fall below those necessary to sustain wild bee populations. Environmental changes may thus a double jeopardy to pollinator populations, as they become victims of both the changes themselves and of plant trait evolution (Weinbach et al. 2022). This in turn may result in an eco-evolutionary positive feedback loop that furthers pollinator declines, further reinforcing plant evolution towards a selfing syndrome. This can explain plant-pollinator network degradation as documented in a previous study (Burkle et al. 2013) and raises the concerning prospect of cascading effects in trophic networks in general, beyond plant-pollinator interactions.

In summary, our study highlights the potential of natural populations to respond quickly to environmental changes. However, such evolutionary responses may have impacts on ecological interactions, here plant-pollinator interactions, and potentially cascading trophic consequences in ecosystems. There is thus an urgent need to investigate whether these results are symptomatic of a broader pattern among angiosperms and their pollinators, and if so understanding whether there is a possibility to reverse this process and break this eco-evolutionary positive feedback loop.

Acknowledgments

We thank Pauline Durbin and Fabien Lopez for help in taking care of the plants during the experiment, Marie-Pierre Dubois for lab-work guidance, the Conservatoire Botanique National de Bailleul and the Conservatoire Botanique National du Bassin Parisien for the supply of seeds and information on sampling locations, Antoine Deverge Merdrignac and Anaïs Vignerol for help in data collection, Ana Rodrigues and John D. Thompson for improving this manuscript. Genotyping data was obtained using technical facilities of GenSeq of the "Institut des Sciences de l'Evolution de Montpellier" with the support of LabEx CeMEB an ANR "Investissements d'avenir" program (ANR-10-LABX-04-01).

Competing interests

Authors declare that they have no competing interests.

Author contributions

Conceptualization: POC, SAP; methodology: SAP, POC; lab work: LD, SAP; data collection: SAP, PG, VP, POC; analysis: SAP, LD; writing – original draft: SAP; writing – review & editing: POC, SAP

Data availability

All data are available on (*access will be available before publication*).

References

Barrett SCH. **2002**. The evolution of plant sexual diversity. *Nature Reviews Genetics* **3**: 274–284.

Bates D, Mächler M, Bolker B, Walker S. **2015**. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**.

Baude M, Kunin WE, Boatman ND, Conyers S, Davies N, Gillespie MAK, Morton

RD, Smart SM, Memmott J. **2016**. Historical nectar assessment reveals the fall

and rise of floral resources in Britain. *Nature* **530**: 85–88.

Benton MJ, Wilf P, Sauquet H. **2022**. The Angiosperm Terrestrial Revolution and the

origins of modern biodiversity. *New Phytologist* **233**: 2017–2035.

Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T,

Schaffers AP, Potts, SG, Kleukers R, Thomas CD, et al. 2006. Parallel

declines in pollinators and insect-pollinated plants in Britain and the Netherlands.

Science **313**: 351–354.

Bodbyl Roels SA, Kelly JK. **2011**. Rapid evolution caused by pollinator loss in

Mimulus guttatus: rapid evolution after pollinator loss. *Evolution* **65**: 2541–2552.

Brandenburg A, Kuhlemeier C, Bshary R. **2012**. Hawkmoth pollinators decrease seed

set of a low-nectar *Petunia axillaris* line through reduced probing time. *Current*

Biology **22**: 1635–1639.

Breuninger H, Lenhard M. **2010**. Control of tissue and organ growth in plants. *Current*

Topics in Developmental Biology (91 vol.). Elsevier. pp 185–220.

Brys R, Jacquemyn H. **2012**. Effects of human-mediated pollinator impoverishment on

floral traits and mating patterns in a short-lived herb: an experimental approach.

Functional Ecology **26**: 189–197.

Burkle LA, Marlin JC, Knight TM. **2013**. Plant-pollinator interactions over 120 years:

loss of species, co-occurrence, and function. *Science* **339**: 1611–1615.

Cheptou P, Imbert E, Thomann M. **2022**. Rapid evolution of selfing syndrome traits in

Viola arvensis revealed by resurrection ecology. *American Journal of Botany* **109**:

1838–1846.

Cheptou P-O. **2019**. Does the evolution of self-fertilization rescue populations or

increase the risk of extinction? *Annals of Botany* **123**: 337–345.

De Jong TJ, Van Dijk H, Klinkhamer PGL. **2005**. Hamilton's rule, imprinting and

parent-offspring conflict over seed mass in partially selfing plants: Parent-

offspring conflict over seed mass. *Journal of Evolutionary Biology* **18**: 676–682.

Drossart M, Rasmont P, Vanormelingen P, Dufrêne M, Folschweiller M, Pauly A,

Vereecken N, Vray S, Zambra E, D'Haeseleer J, et al. 2019. *Belgian Red List*

of Bees. Presse universitaire de l'Université de Mons, Mons, Belgique.

Fisher RA. **1970**. *Statistical Methods for Research Workers*. Darien, USA: Hafner

Publishing Company.

Franks SJ, Hamann E, Weis AE. **2018**. Using the resurrection approach to understand

contemporary evolution in changing environments. *Evolutionary Applications* **11**:

17–28.

Franks SJ, Sim S, Weis AE. **2007**. Rapid evolution of flowering time by an annual

plant in response to a climate fluctuation. *Proceedings of the National Academy of*

Sciences **104**: 1278–1282.

Gervasi DDL, Schiestl FP. **2017**. Real-time divergent evolution in plants driven by

pollinators. *Nature Communications* **8**: 14691.

Glémin S, Ronfort J. **2013**. Adaptation and maladaptation in selfing and outcrossing

species: new mutations versus standing variation. *Evolution* **67**: 225–240.

Goldberg EE, Kohn JR, Lande R, Robertson KA, Smith SA, Igić B. **2010**. Species

selection maintains self-incompatibility. *Science* **330**: 493–495.

Goudet J, Jombart T, Kamvar ZN, Archer E, Hardy O. **2022**. *hierfstat: Estimation*

and Tests of Hierarchical F-Statistics. R package 0.5-11. URL https://CRAN.R-
project.org/package=hierfstat.

Goulson D, Nicholls E, Botías C, Rotheray EL. **2015**. Bee declines driven by
combined stress from parasites, pesticides, and lack of flowers. *Science* **347**:
1255957.

**Grab H, Branstetter MG, Amon N, Urban-Mead KR, Park MG, Gibbs J, Blitzer EJ,
Poveda K, Loeb G, Danforth BN**. **2019**. Agriculturally dominated landscapes
reduce bee phylogenetic diversity and pollination services. *Science* **363**: 282–284.

**Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, Stenmans W,
Müller A, Sumser H, Hörren T, et al. 2017**. More than 75 percent decline over
27 years in total flying insect biomass in protected areas. *PLOS ONE* **12**:
e0185809.

Hansen DM, Van der Niet T, Johnson SD. **2012**. Floral signposts: testing the
significance of visual 'nectar guides' for pollinator behaviour and plant fitness.
Proceedings of the Royal Society B: Biological Sciences **279**: 634–639.

Igic B, Busch JW. **2013**. Is self-fertilization an evolutionary dead end? *New Phytologist*
198: 386–397.

**Janousek WM, Douglas MR, Cannings S, Clément MA, Delphia CM, Everett JG,
Hatfield RG, Keinath DA, Koch JBU, McCabe LM, et al. 2023**. Recent and
future declines of a historically widespread pollinator linked to climate, land
cover, and pesticides. *Proceedings of the National Academy of Sciences* **120**:
e2211223120.

Latron M, Arnaud J-F, Ferla H, Godé C, Duputié A. **2018**. Polymorphic nuclear

markers for coastal plant species with dynamic geographic distributions, the rock

samphire (*Crithmum maritimum*) and the vulnerable dune pansy (*Viola tricolor*

subsp. curtisii). *Molecular Biology Reports* **45**: 203–209.

Lloyd DG. **1992**. Self- and cross-fertilization in plants. II. The selection of self-

fertilization. *International Journal of Plant Sciences* **153**: 370–380.

Martin G, Fontaine C, Accatino F, Porcher E. **2019**. New indices for rapid assessment

of pollination services based on crop yield data: France as a case study.

Ecological Indicators **101**: 355–363.

Noël E, Jarne P, Glémin S, MacKenzie A, Segard A, Sarda V, David P. **2017**.

Experimental evidence for the negative effects of self-fertilization on the adaptive

potential of populations. *Current Biology* **27**: 237–242.

Ollerton J, Winfree R, Tarrant S. **2011**. How many flowering plants are pollinated by

animals? *Oikos* **120**: 321–326.

Ornduff R. **1969**. Reproductive biology in relation to systematics. *TAXON* **18**: 121–133.

Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. **2010**.

Global pollinator declines: trends, impacts and drivers. *Trends in Ecology &*

Evolution **25**: 345–353.

Pyke GH. **1991**. What does it cost a plant to produce floral nectar? *Nature* **350**: 58–59.

Ramos SE, Schiestl FP. **2019**. Rapid plant evolution driven by the interaction of

pollination and herbivory. *Science* **364**: 193–196.

Raven PH, Wagner DL. **2021**. Agricultural intensification and climate change are

rapidly decreasing insect biodiversity. *Proceedings of the National Academy of*

Sciences **118**: e2002548117.

Rousset F. **2023**. *genepop: Population Genetic Data Analysis Using Genepop*. R
package 1.2.2. URL https://CRAN.R-project.org/package=genepop.

Scoppola A, Lattanzi E. **2012**. *Viola* section *Melanium* (Violaceae) in Italy. New data
on morphology of *Viola tricolor*-Group. *Webbia* **67**: 47–64.

Sicard A, Lenhard M. **2011**. The selfing syndrome: a model for studying the genetic
and evolutionary basis of morphological adaptation in plants. *Annals of Botany*
107: 1433–1443.

Stebbins GL. **1957**. Self fertilization and population variability in the higher plants. *The*
American Naturalist **91**: 337–354.

Thomann M, Imbert E, Devaux C, Cheptou P-O. **2013**. Flowering plants under global
pollinator decline. *Trends in Plant Science* **18**: 353–359.

Thomann M, Imbert E, Engstrand RC, Cheptou P-O. **2015**. Contemporary evolution
of plant reproductive strategies under global change is revealed by stored seeds.
Journal of Evolutionary Biology **28**: 766–778.

Venables WN, Ripley BD. **2002**. *Modern applied statistics with S*. New York: Springer.

Weider LJ, Jeyasingh PD, Frisch D. **2018**. Evolutionary aspects of resurrection
ecology: Progress, scope, and applications-An overview. *Evolutionary*
Applications **11**: 3–10.

Weinbach A, Loeuille N, Rohr RP. **2022**. Eco-evolutionary dynamics further weakens
mutualistic interaction and coexistence under population decline. *Evolutionary*
Ecology **36**: 373–387.

Whittall JB, Hodges SA. **2007**. Pollinator shifts drive increasingly long nectar spurs in
columbine flowers. *Nature* **447**: 706–709.

Winn AA, Elle E, Kalisz S, Cheptou P-O, Eckert CG, Goodwillie C, Johnston MO,

Moeller DA, Ree RH, Sargent RD, et al. 2011. Analysis of inbreeding

depression in mixed-mating plants provides evidence for selective interference

and stable mixed mating. *Evolution* **65**: 3339–3359.

Supporting information

Fig. S1. Experimental design. Four localities were sampled two times, with a minimum

20 years between the two samplings. A first generation (F1), the refresher generation, was

performed in common garden. It was used to analyze population genetics and to produce a

family design of 20 families of 5 siblings per population for a second generation (F2), the

test generation, on which we measured traits.

Fig. S2. Trait measures. List of traits measured and delimitation of their measurements.

Traits represented in this figure are only traits which were directly measured on plants.

Fig. S3. Absence of consistent change in sepal length between 1990's-2000's-ancestors and 2021-descendants. We measure sepal length in 3,955 flowers over 792 individuals. The first two letters are the name of the locality (Co = Commeny; Cr = Crouy; Gu = Guernes; Lh = Lhuys). "A" (▲) ancestral population (collected in 2000 for Co, 1993 for Cr, 2001 for Gu and 1992 for Lh) and "21" (●) descendant population (all collected in 2021).

586 **Table S1. Position and year of the sampling of our populations.**

Location	Commeny (49°7'37.11"N, 1°53'46.30"E)	Crouy (49°24'24.31"N, 3°22'20.73"E)	Guernes (49° 1'10.75", 1°38'46.82"E)	Lhuys (49°16'35.40"N, 3°32'58.52"E)
Year of first sampling	2001	1993	2000	1992
Year of second sampling	2021	2021	2021	2021

Table S2. Trait differences (mean ± SD) among plants of our populations. Statistical analyses were performed using LMMs with Age, Locality and the interaction between the two as fixed effects and family and, when necessary, replication of measures on the same plant as random effects (see Material and methods). When the Age effect was significant (bold traits), analyses were followed by two-sides T-Tests with a Bonferroni correction performed between the ancestral and descendant populations by locality. Significant differences in traits between ancestors and descendants ($P < 0.05$) was highlighted by bold measures and sign of changes are given using arrows (↘ for a decrease in time and ↗ for an increase in time).

The first two letters are the name of the locality (Co = Commeny; Cr = Crouy; Gu = Guernes; Lh = Lhuys). "A" ancestral population (collected in 2000 for Co, 1993 for Cr, 2001 for Gu and 1992 for Lh) and "21" descendant population (all collected in 2021).

Trait	N	CoA / Co21	CrA / Cr21	GuA / Gu21	LhA / Lh21	Factor	Chisq	Chi df	P
Floral traits									
Corolla length (mm)	3955	**14.03 ± 1.09**	13.28 ± 1.45	14.06 ± 2.18	13.23 ± 1.47	**Age**	15.314	1	< 0.001
		12.09 ± 1.13	13.03 ± 1.88	13.66 ± 1.35	12.99 ± 1.46	Locality	13.384	3	0.004
		↘				Age × Locality	16.638	3	< 0.001
Labellum width (mm)	3955	**7.16 ± 0.94**	7.45 ± 0.99	**7.51 ± 1.16**	7.04 ± 0.76	**Age**	17.872	1	< 0.001
		6.03 ± 0.59	7.30 ± 0.92	**7.01 ± 0.88**	6.94 ± 0.67	Locality	30.047	3	< 0.001
		↘		↘		Age × Locality	14.725	3	0.002
Floral area approximation (mm²)	3955	**102.2 ± 19.9**	101.0 ± 22.5	**108.8 ± 33.1**	95.5 ± 21.2	**Age**	17.965	1	< 0.001
		74.3 ± 14.0	97.7 ± 23.5	**97.6 ± 20.5**	92.5 ± 18.2	Locality	17.979	3	< 0.001
		↘		↘		Age × Locality	14.989	3	0.002
Spur length (mm)	3954	**4.25 ± 0.22**	4.30 ± 0.24	4.30 ± 0.28	4.37 ± 0.19	**Age**	8.373	1	0.004
		4.00 ± 0.23	4.27 ± 0.27	4.23 ± 0.31	4.38 ± 0.23	Locality	42.226	3	< 0.001
		↘				Age × Locality	12.388	3	0.006
Sepal length (mm)	3955	13.92 ± 1.32	14.26 ± 1.51	12.64 ± 1.34	13.61 ± 1.19	Age	0.550	1	0.459
		13.55 ± 1.01	13.95 ± 1.51	12.71 ± 1.35	13.77 ± 1.32	Locality	63.509	3	< 0.001
						Age × Locality	2.7607	3	0.430
Sepal/corolla lengths ratio	3955	**0.998 ± 0.078**	1.088 ± 0.117	0.918 ± 0.120	**1.044 ± 0.102**	**Age**	14.137	1	< 0.001
		1.132 ± 0.099	1.093 ± 0.125	0.941 ± 0.107	**1.077 ± 0.083**	Locality	93.828	3	< 0.001
		↗			↗	Age × Locality	15.302	3	0.002

Trait	N					Effect		df	p
Number of guides	3993	**6.46 ± 0.98**	**7.38 ± 0.68**	6.94 ± 0.64	7.13 ± 0.48	**Age**	33.591	1	**< 0.001**
		4.89 ± 1.48	6.62 ± 0.94	6.83 ± 0.88	7.04 ± 0.42	**Locality**	112.889	3	**< 0.001**
		↘	↘			**Age × Locality**	30.675	3	**< 0.001**
Anthesis duration (days)	3991	**6.39 ± 0.85**	6.20 ± 1.06	**7.85 ± 1.56**	**7.00 ± 1.04**	**Age**	9.261	1	**0.002**
		5.07 ± 0.85	6.07 ± 0.91	7.27 ± 1.29	7.41 ± 0.81	**Locality**	137.555	3	**< 0.001**
		↘		↘	↗	**Age × Locality**	24.407	3	**< 0.001**
Flowering date (days after the first flowering)	792	20.31 ± 6.78	17.48 ± 8.78	19.58 ± 10.81	19.30 ± 7.57	Age	0.027	1	0.870
		18.54 ± 4.58	17.57 ± 7.82	20.23 ± 12.69	19.69 ± 9.13	Locality	3.783	3	0.286
						Age × Locality	1.007	3	0.800
Average floral display (open flowers per day)	789	**6.37 ± 2.39**	6.85 ± 2.59	7.89 ± 3.47	6.64 ± 2.31	**Age**	8.190	1	**0.004**
		4.64 ± 1.99	6.89 ± 2.87	6.95 ± 3.26	6.43 ± 2.17	**Locality**	32.182	3	**< 0.001**
		↘				**Age × Locality**	7.855	3	**0.049**

Vegetative traits

Trait	N					Effect		df	p
Rosette diameter (mm)	792	118.23 ± 42.75	109.80 ± 38.60	110.05 ± 40.29	101.12 ± 33.77	Age	0.194	1	0.660
		108.15 ± 28.11	108.08 ± 38.24	109.49 ± 42.85	109.11 ± 48.21	Locality	5.601	3	0.133
						Age × Locality	2.397	3	0.494
Leaf length (mm)	792	26.30 ± 6.38	24.10 ± 5.63	22.34 ± 5.18	21.57 ± 4.85	Age	1.226	1	0.268
		22.42 ± 3.95	24.28 ± 5.65	23.17 ± 6.07	21.94 ± 5.63	**Locality**	19.895	3	**< 0.001**
						Age × Locality	11.311	3	**0.010**
Leaf width (mm)	792	19.84 ± 3.77	18.67 ± 3.03	17.02 ± 2.76	17.58 ± 3.07	Age	2.499	1	0.114
		17.96 ± 2.56	18.94 ± 3.01	16.66 ± 2.91	17.66 ± 3.07	**Locality**	40.014	3	**< 0.001**
						Age × Locality	8.024	3	**0.046**
Leaf area approximation (mm^2)	792	541.3 ± 231.9	461.3 ± 173.6	391.3 ± 152.8	390.2 ± 147.1	Age	1.910	1	0.167
		410.5 ± 128.7	472.4 ± 177.6	398.5 ± 173.7	401.6 ± 173.0	**Locality**	27.414	3	**< 0.001**
						Age × Locality	10.559	3	**0.014**
Biomass (g)	777	7.243 ± 2.515	8.220 ± 2.712	8.027 ± 2.624	8.090 ± 2.901	Age	0.029	1	0.864
		8.034 ± 2.553	7.789 ± 2.747	7.606 ± 2.904	8.386 ± 3.066	Locality	3.490	3	0.322
						Age × Locality	4.850	3	0.183

Reproductive traits

Trait	N					Effect		df	p
Log-ratio of the number of seeds without/with pollinators	693	-0.039 ± 0.619	0.085 ± 0.659	-0.008 ± 0.775	-0.342 ± 0.546	Age	3.201	1	0.074
		-0.046 ± 0.711	0.169 ± 0.687	0.173 ± 0.757	-0.220 ± 0.601	**Locality**	38.344	3	**< 0.001**
						Age × Locality	1.671	3	0.643
Weight per seed with pollinators (mg)	656	0.446 ± 0.199	0.421 ± 0.153	0.301 ± 0.154	0.541 ± 0.146	**Age**	4.744	1	**0.029**
		0.385 ± 0.161	0.369 ± 0.146	0.300 ± 0.143	0.546 ± 0.142	**Locality**	212.885	3	**< 0.001**
						Age × Locality	5.999	3	0.112

	N					Factor	χ²	df	p
Weight per seed without pollinators (mg)	701	**0.689 ± 0.206** 0.565 ± 0.200 ↘	0.633 ± 0.169 0.585 ± 0.164	0.517 ± 0.187 0.501 ± 0.189	0.764 ± 0.137 0.788 ± 0.117	**Age** **Locality** **Age × Locality**	9.3644 226.2555 18.3624	1 3 3	**0.002** **< 0.001** **< 0.001**
Plant-pollinator interaction traits									
Nectar volume (μL)	400	0.512 ± 0.280 0.450 ± 0.251	**0.330 ± 0.218** **0.249 ± 0.215** ↘	**0.586 ± 0.470** **0.379 ± 0.260** ↘	0.248 ± 0.182 0.239 ± 0.213	**Age** **Locality** Age × Locality	12.605 75.120 4.636	1 3 3	**< 0.001** **< 0.001** 0.200
Bumblebee visit preferences (proportion of visit per plant)	456	**0. 083 ± 0. 008** **0. 056 ± 0. 007** ↘	0.073 ± 0.008 0.062 ± 0.010	0.068 ± 0.006 0.051 ± 0.005	0.061 ± 0.007 0.053 ± 0.006	**Age** Locality Age × Locality	14.722 2.027 1.950	1 3 3	**< 0.001** 0.567 0.583