Accounting for active populations provides a new perspective on the degree and mechanisms of

soil microbiome resilience

Samuel E. Barnett[1], Ashley Shade[2*]

[1]Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,

MI 48824, USA

[2] Université de Lyon, France; Research Group on "bacterial efflux and environmental

resistance," CNRS, INRAe, Ecole Nationale Véterinaire de Lyon and Université Lyon 1, UMR

5557 Écologie Microbienne, 16 rue Dubois 69622 Villeurbanne Cedex, FRANCE

Corresponding author: Ashley Shade

UMR5557 - Ecologie Microbienne Lyon

Université Claude Bernard Lyon 1

Bât. Mendel, 5eme étage

16 Rue Dubois

69622 Villeurbanne cedex, FRANCE

+33472431377

ashley.shade@cnrs.fr

Classification: Biological Sciences; Microbiology

Keywords: Microbial ecology, soil, resilience, fire

Abstract

Soil microbiomes are essential mediators of ecosystem services, and their resistance and

resilience to anthropogenic disturbances can have profound effects on soil function. However,

we know little about the determinants of soil microbiome resilience, especially in the context of

long-term (press) disturbances. Here, we leveraged the seven-decade-old underground coal seam

fire in Centralia, Pennsylvania, USA, to examine surface soil bacterial community resilience to

long-term heat disturbance. As the fire slowly advances along coal seams, formerly fire-affected

soils cool to ambient temperature enabling geographically localized comparisons of impacted

soils across a gradient of fire intensities with recovered and reference soils. To assess soil

bacterial microbiome response and resilience to the fire, we executed a seven-year field study,

annually sampling sites selected to capture the range of disturbance intensities. We quantified the

dynamics of the bacterial communities over time and across the fire-impact gradient and

additionally accounted for population activation by combining 16S rRNA with 16S rRNA gene

amplicon sequencing. Over the years, the fire-affected sites cooled, and their microbiome

structures recovered towards those of reference and recovered soils. However, the whole

community demonstrated greater resilience than the active subset, suggesting that resilience

could not solely be attributed to environmental selection and growth of active populations. We

therefore suggest that regional dispersal, persistence, and revival from the local seed bank

contribute to microbiome resilience. Importantly, this work indicates that DNA-based methods

that do not account for activity may overestimate microbiome resilience, with implications for

understanding microbiome functional recovery.

Significance statement

The climate crisis and anthropogenic pressure on the environment are fundamentally altering ecosystems and their functions. Many critical ecosystem functions, such as biogeochemical cycling, are facilitated by microbial communities. Understanding the responses and functional consequences of microbial communities to these long-term (*press*) disturbances requires consistent, interannual observations and a focus on the active microbial populations that contribute to the critical functions. This study leverages the seven-decade-old coal seam fire in Centralia, Pennsylvania, USA, to examine the resilience of soil bacterial microbiomes to an intense, long-term heat disturbance. It provides an interannual analysis of microbiome resilience along a fire-impact gradient and directly compares the dynamics of the active subset and the "whole" community (which also includes signals from dormant and deceased cells). This work provides evidence that immigration of inactive cells together with environmental selection can drive perceived microbiome recovery, and advances understanding of the mechanisms of environmental microbiome resilience.

Introduction

Soil microbial communities are vitally important to the biosphere's health by filling important roles in biogeochemical cycling (1) and plant health promotion (2, 3), among other ecosystem services. Recently, there has been a push to use microbial communities as indicators of soil and ecosystem health (4–7), particularly in contexts of changing environments and disturbance where the stability of microbiomes can provide insight into the short and long-term effects these events have on the ecosystem (4, 5). Community stability can be defined as the interplay between resistance (*i.e.,* the degree to which the community remains unchanged during and following a disturbance event) and resilience (*i.e.,* the rate at which the community returns to a pre-disturbance state) (8–10). Both microbial community resistance and resilience can vary substantially across soils and disturbances (10). The duration of a disturbance event, whether it is a pulse (*i.e.,* a discrete, short-term event) or press (*i.e.,* a long-term, continuous event), and whether the disturbance is compounded, may differentially affect the stability of the affected microbial community (9, 11). In light of the long-term environmental impacts of human activities such as industrial mining and climate change, microbial community stability under press disturbance is of particular concern in soil microbial and restoration ecology (9, 12, 13).

Soil microbial communities tend to be sensitive (low resistance) to intense press disturbances (9). However, the rate and degree of resilience of these communities can vary widely. The perceived variability in resilience may be due to complex, interacting, and poorly understood ecological and evolutionary mechanisms that determine resilience (9, 10), including the interplay of speciation, selection, and dispersal assembly processes (14, 15). Quantifications of microbiome resilience may also be complicated by the substantial pool of dormant microorganisms that persists in many environmental systems, but especially in soils (16).

Dormancy is a reversible state of extremely low metabolism (Lennon and Jones) that can allow

bacteria to survive under poor conditions or during disturbance (17–19). Some dormant bacterial

cells can remain viable for extended periods, and some can be selectively revived given

conducive growth conditions (17, 18). Because ecosystem response to disturbance reflects the

functions of active microbiome members (9, 20), it is important to understand and account for

the dynamics of active members as a more direct predictor of ecosystem resilience and functional

recovery.

The seven-decade-old coal mine fire in Centralia, Pennsylvania, USA, provides a unique

case study to examine the resilience of soil microbial communities to an intense, long-term

thermal disturbance (21). The temperatures of the surface soils above the coal seams that have

active fire fronts are continuously higher than ambient soil temperatures (22). Additionally, coal

combustion products percolate through the soil and open vents in the ground, altering soil

chemistry at the surface (22–24). The gradual advancement of the fire along the coal seams

allows for direct comparisons across a gradient of disturbance intensities that we defined by soil

temperature and across fire-affected, decades-old recovered, and unaffected reference soils, all

within a geographically localized area encompassing less than 0.35 km^2 (21). Furthermore, as

disturbance intensity changes over time at a sampling site, community changes can be tracked

and understood in the context of the dynamic warming and cooling that define this press

disturbance.

The overarching goal of this study was to understand the capacity of soil bacterial

communities to recover from an intense press disturbance and to identify underlying ecological

mechanisms that support resilience. Our previous work investigated microbiome responses over

the Centralia fire-impact gradient. It showed that the microbiomes of recovered soils converged

towards reference soils, thus demonstrating resilience, while the microbiomes of fire-affected

soils maintained distinct structures and were highly variable (25). We subsequently conducted a

follow-up laboratory study to control dispersal and to track activation of the Centralia soil

microbiome to heating and found that both reactivation from dormancy and immigration together

contributed to the observed resilience (26). These results motivated us to track activation

dynamics in Centralia to understand the contributions of active members to the community

disturbance response.

Here, we present a 7-year longitudinal study of the "whole" bacterial communities and

the subset of active members (27, 28) annually sampled from our original fire-impact gradient of

fire-affected, recovered, and reference sites in Centralia (Fig. 1A, Table S1). Whole bacterial

communities were assessed using 16S rRNA gene amplicons, which include contributions of

active, dormant, and recently deceased cells. The active subset of the whole community was

determined by filtering the dataset to include only the taxa that met a conservative, population-

level activity criterion based on their 16S rRNA to 16S rRNA gene ratios. Then, we applied

several specific methods to quantify microbiome recovery, particularly time-lag and trajectory

analysis (29, 30), where resilience is indicated by the slope of a time-lag regression and by the

strength of directionality in trajectory analysis (e.g., toward an undisturbed state). Thus, we

expected the strength of directionality and time-lag regression slope to correlate positively with

disturbance intensity. Directionality and time lag were also compared across fire categories and

between the whole and active partitions to determine their relative resilience. Building from

previous findings (19, 25), we hypothesized that soil bacterial communities are resilient to long-

term heat disturbance and that the observed resilience is largely attributable to directional

changes in the active populations, indicative of environmental selection and turnover as soils

cool.

Results

Soil temperature and pH vary over fire classification

As expected, fire-affected soils were generally hotter than recovered and reference soils.

Notably, the fire-affected soil temperatures decreased over the seven years, yet no consistent

changes over time were observed in either recovered or reference sites (Fig. 1B, Table S2; linear

mixed effects models). Soil pH did not change over time (Fig. S1A, Table S2; linear mixed

effects models), though it did overall differ across fire classifications (ANOVA: DF = 2, F =

14.54, p-value < 0.001). pH was lower in recovered sites than either reference or fire-affected

sites, and the pH range across fire-affected sites was large (Fig. S1B, posthoc Tukey's: p-values

< 0.05).

Bacterial diversity changes across fire classification and over time

We used 16S rRNA gene and 16S rRNA amplicon sequencing of co-extracted DNA and

reverse-transcribed RNA to assess community structure across our samples. DNA and RNA

sequences were processed identically with shared operational taxonomic units (OTUs) generated

de novo with 97% sequence identity. Non-bacterial OTUs were removed, and the reads were

rarefied to the lowest depth within the analysis: 161,171 for DNA-only analysis (*i.e.,* whole

community), 100,491 for DNA and RNA analysis (*i.e.,* active community subset).

To examine how bacterial communities varied over time within our sites, we examined

the within-site phylogenetic diversity (*i.e.,* Faith's phylogenetic diversity). We found increased

diversity over time in fire-affected soils for both whole communities and active subsets but no

increase in recovered or reference soils (Fig. S2; Table S3; linear mixed effects models). When

time and fire classification were ignored, diversity increased as soil temperature decreased for

whole communities and active subsets (Fig. 1C and D, Table S4; linear mixed effects models).

We next evaluated the variation in phylogenetic and compositional diversity between

sites (*i.e.,* abundance-weighted UniFrac distance). Variation across whole bacterial communities

was explained by fire classification, time, and their interaction (Fig. S3, Table 1; PERMANOVA

blocked by Site ID). Fire classification explained the most variation (18%), while time and the

interaction of fire classification and time explained about 6% and 4%, respectively. *Post hoc*

tests within each fire class showed that time only explained variation in community structure in

fire-affected sites (Fig S4, Table S5). *Post hoc* tests within each year showed fire classification

explaining structural variation in all years, but the variation explained (R^2) decreased between

2015 and 2018 (Fig. S5, Table S5). Structural variation between fire classifications could be

attributed to differences in beta-dispersion across fire classifications (ANOVA; df = 2, F =

34.722, p-value < 0.001). Dispersion across fire-affected soil was higher than recovered and

reference soils (Table S6).

For the active subset of the bacterial community, fire classification and time, but not their

interaction, explained variation in community structure (Fig. S6, Table 1; PERMANOVA

blocked by Site ID). As for the whole community, fire classification explained more variation in

community structure than time (12% and 6%, respectively). Similarly, beta dispersion varied

across fire classifications for the active community subset (ANOVA; df = 2, F = 17.536, p-value

< 0.001). Dispersion across fire-affected soil was again higher than recovered and reference

soils, though the effect was smaller than for the whole community (Table S6).

Constrained correspondence analysis (CCA) demonstrated that the measured edaphic

factors, particularly soil temperature, pH, and pH-associated properties (*e.g.,* Ca, Mg, and Fe

concentration), explained a large portion of the variation in community structure for both whole

communities (Fig. S7A; variation explained = 45.342%) and active subsets (Fig. S7B; variation

explained = 35.789%). Variation between fire-affected and both reference and recovered sites

was largely associated with soil temperature, as expected. In contrast, variation across and within

the reference and recovered sites were primarily associated with pH (Fig. S7).

Heat intensity drives community trajectories

For both the whole bacterial communities and active subsets, time-lag analysis showed

increasing community dissimilarity with an increasing difference by year for fire-affected sites

but no relationship for either recovered or reference soils (linear mixed effects models; Fig. 2A

and B, Table S7). Using maximum temperature as an indicator of overall disturbance intensity,

we found that the slopes of the time lag relationships are positively correlated to the maximum

soil temperature of the sites (Fig 2C and D; linear regression; whole community: $R^2 = 0.67$, p-

value < 0.001, active subset: $R^2 = 0.30$, p-value $= 0.011$). Notably, the relationship between the

time lag slope and maximum soil temperature was stronger for the whole community than the

active subset.

Similar to the time-lag analysis, for both the whole communities and active subsets, there

was higher directionality in the fire-affected sites than either recovered or reference sites, with no

significant difference between the latter classifications (Fig. 3A and B; Kruskal-Wallis rank sum

test; whole community: chi-squared = 9.04, df = 2, p-value = 0.01, active subset: chi-squared =

8.67, df = 2, p-value = 0.01; *post hoc* Dunn's tests). Community directionality was positively

correlated with maximum soil temperature, with a stronger relationship for the whole

communities relative to the active subsets (Figs. 3C and D; linear regression; whole community:

$R^2 = 0.55$, p-value < 0.001, active subset: $R^2 = 0.20$, p-value $= 0.036$).

To assess whether the bacterial communities were becoming more like the undisturbed

reference communities as soils cooled, we tested whether the distance in community structure

between disturbed soils (*i.e.,* fire-affected and reference soils) and recovered soils within years

was associated with disturbance intensity, defined here as the difference in soil temperature to

reference soil. We found that for both whole communities and active subsets, as the disturbance

intensity decreased, the communities were less dissimilar (Fig 4; linear mixed effects models).

However, the y-intercept was higher for the active subsets than whole communities, suggesting a

greater difference in community structure even when disturbance intensity is null.

Active members contribute little to structural shifts in the whole community

We calculated the percentage of Bray-Curtis dissimilarity between years within

individual sites attributable to the dynamics of active OTUs. To examine the influence of OTUs

that become inactive or newly active over time, we ran this analysis twice using OTUs active in

either the earlier or the later year of year pairing, respectively. Bray-Curtis dissimilarity was used

in this case, rather than abundance-weighted UniFrac, because it can be effectively partitioned to

a subset of OTUs as previously demonstrated (31). Surprisingly, for the majority, less than 50%

of the structural dissimilarity in the whole community over time was attributable to active OTUs.

For fire-affected sites, but not for recovered or reference sites, the dissimilarity attributed to

active taxa within a site over time decreased as the period between sampling years increased

(Fig. 5, Table S8; linear mixed effects models; p-values < 0.05). This trend was statistically

supported when selectively partitioning taxa active at the earlier and later timepoint in the year-

pairing.

Interannual OTU activity dynamics vary across fire-affected and unaffected soils

We next examined the breakdown of OTUs into distinct yearly activity patterns (*i.e.,* sets

of years in which they were identified to be active; Fig. 6 and Figs. S8 and S9). Only OTUs

active in at least five samples were included in this analysis, and activity within a year was

defined as activity in at least one site. Furthermore, reference and recovered sites were

categorically combined as minimal change over time in active community subsets was observed

for these microbiomes. For fire-affected sites (Fig. 6A), the richest activity patterns (*i.e.,* those

encompassing the most OTUs) tended to have consistent activity in later sampling years (*e.g.,*

2018-2021) but not for early sampling years (*e.g.,* 2015-2017). This trend suggests that many

observed OTUs may become active in later years for fire-affected sites, corresponding to cooling

soil temperatures and recovery. For reference and recovered soils (Fig. 6B), the richest activity

patterns had no clear annual progression or delineation. This lack of trend suggests more random

activity patterns in the soils that experienced ambient temperatures. Activity groups were

taxonomically diverse for both cases, with no obvious enrichment in taxa at the phylum level or

class level for *Proteobacteria*.

Community assembly processes for whole communities differ across fire classifications

We used the beta-nearest taxon index (βNTI) (32, 33) and the abundance-weighted Raup-

Crick index (RC_{bray}) (33) between sites within each year to determine the dominant processes

driving whole bacterial community assembly within each fire classification. Due to the

computational requirements for these calculations, DNA samples were first rarefied to 10,000

reads. While the majority of within-year, site-to-site comparisons indicate dominance of

homogeneous selection (βNTI < -2), phylogenetic turnover among fire-affected sites tended to

be higher than among recovered or reference sites (Fig. 7A; Kruskal-Wallis rank sum test: DF =

2, chi-squared = 68.399, p-value < 0.001; post hoc Dunn test) with many site pairs demonstrating

weak selection (-2 $\leq$ βNTI $\leq$ 2) or variable selection (βNTI > 2). This pattern was consistent

across seven years (Fig. 7D and S10). For these site pairs with weak selection, homogenizing

dispersal (RC_{bray} < -0.95), followed by drift alone (-0.95 $\leq$ RC_{bray} $\leq$ 0.95), tended to be the

dominant assembly process (Fig. 7D).

253   To ask if two of the most notable edaphic soil factors were associated with potential

selection pressures in this system, we examined linear relationships between βNTI and scaled

between-site differences in soil pH and temperature and their interaction between sites agnostic

of fire classification. Both edaphic factors showed statistically significant positive relationships

to βNTI (Fig. 7B and C, Table S9; linear regression: R^2 = 0.5031, p-value < 0.001). Since we

used scaled measures for differences in pH and temperature, we can directly compare their

coefficients to examine their relative influence on phylogenetic turnover. We found that the

coefficient for the difference in pH (2.63) is much larger than that of the difference in

temperature (0.42), suggesting a greater influence of pH on phylogenetic turnover than on

temperature. There was no statistically significant interaction between these two factors

suggesting unrelated influences of temperature and pH in selection (Table S9).

Discussion

We used long-term microbiome surveys across soils from Centralia, Pennsylvania, USA,

to examine the resilience of whole bacterial communities and active subsets impacted by heat

press disturbance from a subterranean coal seam fire. Over seven years, we observed general

cooling of the fire-affected sites (Fig. 1B), indicating a continuous reduction in disturbance

intensity. Cooling was not unexpected, as the fire is predicted to move deeper underground with

its advance, following the coal seam incline (22). While heat intensity decreased in most sites,

some other fire-induced environmental changes continued. For example, elevated local CO_2

concentrations and steam venting were sometimes observed even when soils were only a few

degrees Celsius above ambient temperature (Table S1). While fire disturbance was ongoing, the

observable change in intensity provided a unique opportunity to observe the transition and

recovery of the microbial communities during both the early stages of recovery (*i.e.,* cooling fire-

affected sites) and later stages (*i.e.,* recovered sites).

Temporal trends in bacterial diversity, both within-sample (alpha) and between-sample

(beta) revealed strong evidence for the resilience of the microbiomes in fire-affected sites and

both the whole communities and in the active subsets (Figs. 1C-D, S2-S7). In comparison,

communities in recovered and reference sites were relatively less dynamic and more stable over

the years. Applying the concepts from the stability basin analogy for community resilience (29),

both directionality and time-lag analyses (Figs. 2 and 3) suggested that the fire-affected soils

were shifting from their high-temperature, disturbed community state to a relatively stable,

recovered state. Comparatively, the recovered and reference soils maintained a relatively stable

state over time. Further, the recovered state for the fire-affected and recovered soils appears to be

like the reference soils that did not experience any heating from the fires (Fig. 4). These findings

expand and reinforce the previous results from the single sampling timepoint in Centralia, that

also indicated recovery of bacterial communities in Centralia soils (25).

Dispersal is an important microbiome assembly mechanism for soil (34–37). Across our

Centralia sites, when selection was weak ($-2 < \beta NTI < 2$), which occurred almost exclusively

across fire-affected sites, homogenizing dispersal ($RC_{bray} < -0.95$) was the dominant assembly

process (Fig. 6). Laboratory-based manipulations of reference soils from Centralia have

demonstrated that dispersal enhances the resilience of microbial communities to heat disturbance

as compared to controls without dispersal (26). Dispersal likely explains the increased within-site

diversity observed as fire-affected soils cool (*i.e.,* new taxa adding to the local species pool).

However, even when dispersal is high, immigrant taxa may not necessarily be active or become

established. Rather their DNA may remain extractable either in intact cells (dormant or dead) or

as extracellular DNA (eDNA) (38–40). There have been proposed models whereby dormancy

potentially aids in microbial dispersal capability (*e.g.,* microbial conveyor belt (41)), though

more evidence is needed to support these hypotheses (42). During the 45-week mesocosm

experiment that explicitly controlled dispersal (26), we observed that post-disturbance activation

of some of the dispersed taxa was possible and contributed to overall resilience. Building from

these collective results, we present an updated working hypothesis to integrate dispersal and

dormancy processes and suggest one potential mechanism of soil microbiome resilience. We

hypothesize that taxa lost to disturbance are progressively replaced by dispersed members from

the regional species pool (*i.e.,* undisturbed and recovered soils) as thermal selection weakens but

likely maintains dormancy while conditions remain suboptimal. This process effectively rebuilds

and accumulates the dormant seed bank but does not contribute immediately to functional

recovery. When conditions become favorable, members of the seed bank can activate and

progress along a recovery trajectory. Re-establishing the microbial seed bank after a disturbance

provides a management strategy that may be useful in bioremediation and active restoration of

soils (43, 44), but much more work is needed to test and understand these proposed mechanisms.

Patterns of OTU activity over time in fire-affected sites (Fig. 6) demonstrated that many

bacteria are likely to be activating (either after dispersal or from the local seed bank) in the fire-

affected soils over the last few years as the soils have cooled. Of the top 20 interannual activity

patterns, in terms of the number of OTUs displaying the same pattern of activity, most tend to

have detectable activity only in the later sampling years. In other words, many active taxa

become detectably active after one to three years of no detectable activity in fire-affected soil.

This is best exemplified by the second most numerous activity pattern that includes activity in

the years 2016-2021 but not 2015. In combined reference and recovered sites, on the other hand,

the top 20 patterns appear to have relatively random years where activity is not detected. Most of

these top 20 activity patterns include activity in the first few sampling years. Upon examination

of the phylum-level taxonomic breakdown of these activity patterns, we see that the most

numerous patterns are comprised of highly diverse phyla (Fig. 6). This taxonomic diversity

suggests that activation and perhaps recovery of the active community subset is indiscriminate

across the phylogenetic tree, or at least at high taxonomic levels.

Notably, the temporal changes within the whole community were greater than in the

active subset, indicating differences in resilience and recovery between the active and whole

communities. For example, time explains a greater portion of the structural variation for whole

communities than for active subsets, and no interaction was found between time and fire

classification for those active communities (Table 1). Similarly, the relationships between heat

intensity (*i.e.,* maximum temperature) and both time lag slopes and structural directionality were

weaker for active subsets than the whole communities. When considering variation in the whole

community structure within a site over time, active OTUs tended to explain a smaller percentage

of the community dissimilarity as the timescale increased, but only for fire-affected sites (Fig. 5).

This important finding indicates that over the long term (*i.e.,* multiple years), changes in the

structure of the active community subset accounts for less of the change in overall community

structure in recovering soils relative to more stable communities. Thus, putting all these results

together, we propose that while the overall bacterial community is largely resilient, the active

populations may recover slower. From this conclusion and the evidence of this study, we

hypothesize that dispersal enables bacterial taxa from the regional species pool to arrive and

possibly establish in the fire-affected soils as they cool, with a portion of immigrants remaining

dormant though persistent as disturbance intensity slowly wanes.

pH and associated edaphic properties are well-known drivers of bacterial community

diversity and soil assembly (35, 45–48). Apart from temperature, edaphic factors associated with

soil pH explained structural differences in the community (e.g., Ca, Mg, Fe, etc.; Fig. 2) (43).

Standardized phylogenetic turnover between sites (βNTI), a key indicator of selection (49, 50),

correlated to differences in soil pH (Fig 6), indicating pH-related selection pressures across

Centralia soils (35, 47). βNTI was also correlated to differences in soil temperature between

sites. However, this relationship was weaker than that of soil pH, and the interaction between

these edaphic factors was not statistically significant, suggesting that the selection pressure of

these factors differs (Table S9). From these trends in phylogenetic turnover, we propose that soil

pH and related factors are selection drivers maintaining bacterial communities across soils in

Centralia but that active coal fires alter the dominant assembly processes either by imposing

alternate selection pressures (*e.g.,* selection for thermotolerant traits) or by increasing

stochasticity (*e.g.,* dispersal and drift) among sites. Other press disturbances, such as land use

intensification, have been shown to interact with soil pH in driving variation in soil bacterial

community assembly (35). The acidic soils in the recovered and some fire-affected sites are

likely the result of the deposition of combustion and mining products such as ammonium, nitrate,

or sulfur (23–25). Soil acidification may be a legacy effect of mining (51, 52) and coal fires in

this area, potentially creating alternative stable states among the recovered communities rather

than a comprehensive, single recovery state (53).

Other mechanisms that are difficult to disentangle from our hypothesized model may also

be contributing to the patterns we observed. First, as observed across seasons from our HOBO

temperature monitors in five sites (Fig. S11), cyclical disturbance intensity patterns may produce

temporary conditions suitable for the growth or maintenance activity of otherwise fire-sensitive

taxa (11). Second, as soils are extremely diverse (54) non-detection of rare taxa is a distinct

possibility despite our deep sequencing, as demonstrated by the rarefaction curves (Fig. S12).

Third, extracellular DNA (eDNA) may contribute to the divergence in the whole community and

active subset dynamics between fire-affected and unheated soils because eDNA in soil degrades

more rapidly at higher temperatures (55). However, a large influence of eDNA on the observed

interannual patterns is not expected due to the short lifespan of nucleic acids in soils in general.

Broadly, our findings and hypothesized model for bacterial community recovery from

press disturbance have implications for understanding ecosystem recovery to large, long-term

stress. We interpret the observed relatively high resilience of the whole community with

somewhat lower resilience of the active subset to indicate that, while microbial community

functional potential may recover rapidly along with the demographic recovery of members lost

to the disturbance, the realized activity of these communities may take longer to recover because

it is harbored within the dormant pool. As only the active subset of the community performs

ecosystem functions and services, it is likely that biologically mediated soil processes, such as

biogeochemical cycling, may remain altered for some period after disturbance intensity wanes,

especially for functions with low redundancy among populations. If ecosystem function is used

to measure ecosystem recovery (12), then whole bacterial community surveys will provide a

biased view of the recovery progress. However, our findings provide a positive outlook on the

recovery potential of heavily anthropogenically disturbed soils. With the growing importance of

restoration ecology in the context of anthropogenetic press disturbances and climate change,

understanding the stability and potential for recovery of soil microbial communities will help to

predict changes to vital ecosystem functions and develop methods of microbiome restoration.

Materials and Methods

Study site, soil collection, and processing

Centralia is located in Columbia County, Pennsylvania, overlying an anthracite seam of

the Buck Mountain coal bed (22). This coal seam has been continuously burning since 1962,

with at least two fire fronts moving at about two m/yr. While the coal seam and fire lie about 46

m below the surface, heat from the fire and combustion gasses make their way to the surface,

significantly increasing surface soil temperatures in some areas and producing observable steam

vents (22). Based on previous experiments, we selected 23 sites along the two main fire fronts

(19, 25). Soils were collected from each site from 2015 to 2021 in the first or second week of

October. Four sites were removed from the experiment before analysis due to exhaustion of

available soil, a landslide, or exceedingly low DNA concentrations in the collected soil.

Where possible, 20 cm deep and 5.1 cm in diameter soil cores were collected using

ethanol-sterilized PVC soil corers. When coring was impossible, primarily in vents, the soil was

collected with an ethanol-sterilized trowel. Collected soil was homogenized and sieved to 4 mm.

Soil for nucleic acid extraction was immediately transferred to a sterile 50 ml falcon tube and

flash-frozen in liquid N_2. The time from soil collection to flash freezing was generally under 4

minutes. Flash-frozen soils were stored at -20˚C or on ice while transported to the lab and then

stored at -80˚C until nucleic acid extraction. The soils used for soil chemistry analyses were

transferred to a Whirl-Pak bag and stored on ice or at 4˚C while transported to the lab. Soil

moisture was measured gravimetrically with ~10 g of soil. The Soil and Plant Nutrition Lab at

Michigan State University (East Lansing, MI, USA; https://www.canr.msu.edu/spnl) processed

all other soil chemistry analyses according to standard protocols. Just before soil collections, soil

core temperature and air temperatures were measured using ethanol-sterilized electronic

thermometers, and CO_2 venting measurements were measured using an EGM-5 portable CO_2 gas

analyzer (PP Systems). Daily soil temperatures were monitored at five sites (Table S1) using

HOBO 64K pendant Temperature data loggers (UA-001-64; ONSET) buried ~10cm below the

surface near the coring site and recovered each year.

DNA/RNA extraction and processing

DNA and RNA were coextracted from 0.25 g of soil using the Griffiths et al. phenol-

chloroform protocol (56), with 0.1 mm zirconium bead containing BeadBug homogenizer tubes

(#Z763764; Benchmark Scientific) for bead beating. DNA and RNA aliquots of extracted

nucleic acid were stored at -80˚C for further processing. DNA and RNA processing was

performed within a month of extraction to prevent excessive RNA degradation. DNA

concentration was determined using Qubit dsDNA BR Assay Kit (#Q32853; Invitrogen). If

possible, DNA aliquots were diluted to ~2 ng/µl DNA, and RNA aliquots were diluted to ~200

ng/µl DNA before further processing. RNA concentration was determined using Qubit RNA HS

Assay Kit (#Q32852; Invitrogen). DNA was degraded using the TURBO DNA-free kit

(#AM1907; Invitrogen) following the rigorous DNase treatment protocol for the RNA aliquot.

cDNA was generated from RNA using the SuperScript III Reverse Transcriptase kit (#18080093;

Invitrogen) following the standard protocol. Both mock community (57) and nuclease-free water

controls were performed for each batch of nucleic acid extraction, and all controls were

processed along with samples.

Amplification and sequencing of DNA and cDNA were performed by the Research

Technology Support Facility at Michigan State University (East Lansing, MI, USA).

Amplification via PCR was accomplished using primers targeting the V4 16S rRNA gene region

with 515f = GTGCCAGCMGCCGCGGTAA and 806r = GGACTACHVGGGTWTCTAAT

(58). The PCR mixture consisted of 7.5 µl 2X Dream Taq master mix (#K1071; Thermo

Scientific), 6.5 µl 0.5 µM primer mix, and 1 µl template DNA or cDNA. PCR was performed

using the following protocol: 95˚C for 3 min, 30 cycles of 95˚C for 45 s, 50˚C for 60 s, 72˚C for

90 s, and finally 72˚C for 10 min. During amplification, dual indexing barcodes were added, and

samples were pooled into two libraries following cleanup. Sequencing was performed on an

Illumina MiSeq with the 2x250 paired-end V2 kit. Triplicate sequencing runs were performed for

both DNA libraries, while single runs were performed for both RNA libraries.

Sequence processing

Raw, demultiplexed sequencing reads are available through the NCBI short read archive

(BioProject PRJNA973689). Individually for each sequencing run, raw reads were first trimmed

using cutadapt (v1.18) (59) to remove remaining primer sequences. Paired-end reads were then

merged using PEAR (v0.9.6) (60) with a maximum length of 600 bp and a minimum overlap of

20 bp. Paired reads with a maximum expected error greater than one and/or containing unknown

bases were removed using usearch (v10.0.240_i86linux64) (61). Filtered sequences were aligned

to the SILVA SEED reference alignment (v132), and poorly aligned sequences were removed

along with those containing homopolymers with a length of 8 or greater using mothur (v1.40.4)

(62). At this point, quality-filtered sequences from all sequencing runs, including DNA and

cDNA libraries, were pooled into a master dataset. Final operational taxonomic units (OTU)

were then clustered *de novo* at 97% sequence identity using usearch (61) with chimera removal.

Representative OTU sequences in FASTA format are available through figshare (DOI:

10.6084/m9.figshare.23060354). OTU taxonomy was assigned based on the SILVA reference

database (v132) using sintax from vsearch (v2.15.2_linux_x86_64) (63) with a probability cutoff

of 0.8. Following the removal of non-bacterial OTUs (*i.e., Archaea*, mitochondria, chloroplast,

and unclassified domain), a phylogenetic tree was generated using FastTree (v2.1.10) (64) with

Sulfolobus (X90478) included as an outgroup aligned with ssu-align (v0.1.1) (65). The sample

metadata, OTU table, taxonomy table, and phylogenetic tree were combined into a phyloseq (66)

object available through figshare (DOI: 10.6084/m9.figshare.23060354). The sequence

processing code is available at https://github.com/ShadeLab/Centralia_RNA_DNA_multiyear.

We defined bacterial OTUs based on 16S rRNA gene sequence clusters at 97% sequence

identity, including DNA and RNA sequence sets. OTUs were used rather than ASVs to avoid the

potential artificial splitting of bacterial populations into multiple clusters (67, 68) and to account

for potential increased errors due to the additional PCR step in reverse transcribing RNA.

Statistical analyses.

All statistical analyses were conducted using R (version 4.1.1) (69). The analysis code is

available at https://github.com/ShadeLab/Centralia_RNA_DNA_multiyear. Change in soil

temperature and pH over time was examined within fire classifications with linear mixed effects

(LME) models with package nlme (70) using either soil core temperature or pH as the response

variable, time as the fixed effect, and site ID as the random effect. Testing broad differences in

soil pH across fire classifications was accomplished using ANOVA and post hoc Tukey's test.

OTU counts from triplicate libraries were added together for DNA samples, and samples

with less than 100,000 reads were removed. RNA samples with less than 10,000 reads were also

removed. Before whole community analysis (DNA samples only), entire sites with less than

three years of adequate DNA sequencing were removed, and the resulting OTU table was

rarefied to 161,171 reads to account for the wide range of sequencing depths. For active bacterial

community analysis, DNA samples without a corresponding well-sequenced RNA sample and

entire sites with less than three years of adequate DNA and RNA sequencing were removed, and

the resulting OTU table was rarefied to 100,491 reads. OTUs found in an RNA sample but not

the corresponding DNA sample were given a count of 1 in the DNA sample (27, 28). OTUs with

an RNA: DNA ratio $\geq$ 1 in each paired RNA and DNA sample were then defined as active (27,

28). Further analysis of the active community subset used the modified OTU tables from the

DNA samples but only included active OTUs (*i.e.,* zero count wherever DNA and cDNA count

was zero or RNA: DNA ratio was < 1).

DNA and cDNA sequence set rarefaction produced 50,242 OTUs across both datasets,

38,020 (75.7%) of which were considered active in at least one sample. For each sample, $50.5 \pm$

0.6% of these active OTUs could be considered phantom (*i.e.,* cDNA reads found, but no reads

in the corresponding DNA samples after rarefaction). However, this phantom percentage drops

to $35.6 \pm 0.7\%$ when considering unrarefied DNA reads and $2.1 \pm 0.1\%$ when considering the

entire DNA dataset regardless of the sample (Fig. S13), providing evidence that many phantom

taxa within the rarefied dataset are likely biological sequences and not entirely artifacts.

Alpha diversity was measured as Faith's phylogenetic diversity indices. Change in alpha

diversity over time was examined within fire classifications with LME models, using the

diversity indices as the response variable, time as the fixed effect, and site ID as the random

effect. Again, differences in alpha diversity across soil temperatures were examined regardless of

fire classification using LME models with the diversity indices as the response variable, soil core

temperature as the fixed effect, and site ID as the random effect.

Beta diversity was measured using abundance-weighted UniFrac distance calculated with

package phyloseq (66). Variation in community structure across fire classification, sampling

year, and their interaction was performed using PERMANOVA blocked by site ID. *Post hoc*

analysis of variation in community structure across fire classifications within each year or across

years within each fire class was also performed using PERMANOVA blocked by site ID and

adjusting p-values for multiple comparisons using the Benjamini-Hochberg procedure. Variation

in community structure explained by edaphic factors and temperature was examined using a

constrained correspondence analysis with package phyloseq (66). Stepwise model variable

selection was accomplished with function ordistep from the vegan package.

Time-lag analysis was performed separately for each fire classification using LME

models with samples paired within each site, abundance-weighted UniFrac distance across sites

as the response variable, the square root of timepoint difference as the fixed effect, and site ID as

the random effect. Slopes of simple time-lag linear regressions for each site were then regressed

to maximum soil temperature at each site, regardless of fire classification, using a linear model.

Community trajectory analyses were performed using package ecotraj (71, 72). Trajectory

directionality was first compared across fire classifications using a Kruskall-Wallis test and *post*

hoc pairwise Dunn's tests with p-values adjusted for multiple comparisons using the Benjamini-

Hochberg procedure. Trajectory directionality was also regressed to maximum soil temperature

at each site, regardless of fire classification, using a linear model.

We calculated the percentage of the Bray-Curtis dissimilarity across sampling time points

within each site attributed to active taxa using methods from Shade *et al.* 2014 (31). In this case,

whole community OTU tables were those of the DNA samples modified for RNA: DNA ratio

analyses, including phantoms and 100,491 read rarefaction. This analysis was run separately for

OTUs active in the first timepoint of each timepoint pair (*e.g.,* 2016 when comparing a site

between 2016 and 2020) and in the second timepoint of each pair (*e.g.,* 2020 when comparing a

site between 2016 and 2020). The percentage of the Bray-Curtis dissimilarity attributed to active

taxa was compared across fire classifications using a Kruskall-Wallis test and *post hoc* pairwise

Dunn's tests with p-values adjusted for multiple comparisons using the Benjamini-Hochberg

procedure. These attributed percentages were also examined across time differences between

samplings within each fire class using LME models with the percentage of Bray-Curtis

dissimilarity as the response variable, the difference in sampling year as the fixed effect, and site

ID as the random effect.

Upset plots were made using custom code based on the R package UpSetR (73). For

these plots, reference and recovered sites were combined as an overall "no-fire" group, and all

sites within a year were combined for the new fire-affected and no-fire groups. Thus, an OTU

was considered active within a specific year if it was active in at least one site within that year.

Only OTUs found to be active in at least five years were included in this analysis.

Community assembly processes contributing to the structure of the whole bacterial

communities were accomplished using calculations of βNTI, a standardized measure of between-

sample phylogenetic turnover indicative of deterministic assembly processes, and RC_{bray}, a

standardized measure of species turnover between samples that combined with βNTI is

indicative of stochastic community assembly processes (49, 50). Before calculations, the DNA

sample OTU table was rarefied to 10,000 reads to reduce computational load. βNTI and RC_{bray}

calculations were conducted for site pairs within a year, not across years. βNTI was calculated

using the big data calculation method from package iCAMP (74). For site pairs, βNTI < -2

indicates assembly explained by homogeneous selection, βNTI > 2 indicates assembly explained

by variable selection, and |βNTI| ≤ 2 indicates assembly explained by stochastic process or weak

selection. For those site pairs with |βNTI| ≤ 2, RC_{bray} was calculated using code modified from

Barnett *et al*. 2020 (35). For these site pairs, RC_{bray} < -0.95 indicates assembly explained by

homogenizing dispersal, RC_{bray} > 0.95 indicates assembly explained by limited dispersal, and

|RC_{bray}| < 0.95 indicates assembly explained by drift alone. βNTI was performed using the

phylogenetic tree including all OTUs regardless of sampling year, assuming all OTUs recovered

are present in some capacity within the regional species pool. In contrast, RC_{bray} used separate

metacommunity abundance profiles for each sampling year to account for potential yearly

changes in the structure of the regional species pool. βNTI values for site pairs within each fire

classification were compared across these classes using a Kruskall-Wallis test and *post hoc*

pairwise Dunn's tests with p-values adjusted for multiple comparisons using the Benjamini-

Hochberg procedure. βNTI values for site pairs were examined in relation to the differences in

soil temperature, the difference in soil pH, and their interaction using a linear model after scaling

and centering both factors to make them directly comparable.

Acknowledgments

We thank Tammy Tobin-Janzen for Centralia site consultation and field support and

Keara L Grady for superb coordination of fieldwork logistics and organization of multi-year soil

procurement and processing. We additionally thank SH Lee, JW Sorensen, TK Dunivin, JL

Chodkowski, N Stopnisek, and M Mechan Llontop for their enthusiastic participation in the

fieldwork. This work was supported by the U.S. National Science Foundation CAREER award

#1749544 to AS.

References

1.   E. L. Madsen, Microorganisms and their roles in fundamental biogeochemical cycles.

Curr. Opin. Biotechnol. **22**, 456–464 (2011).

2.   T. R. Turner, E. K. James, P. S. Poole, The plant microbiome. *Genome Biol.* **14**, 209

(2013).

3.   R. Mendes, P. Garbeva, J. M. Raaijmakers, The rhizosphere microbiome: significance of

plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS*

Microbiol. Rev. **37**, 634–663 (2013).

4.   A. H. C. van Bruggen, *et al.*, One Health - Cycling of diverse microbial communities as a

connecting force for soil, plant, animal, human and ecosystem health. *Sci. Total Environ.*

664, 927–937 (2019).

5.   A. Dubey, *et al.*, Soil microbiome: a key player for conservation of soil health under

changing climate. *Biodivers. Conserv.* **28**, 2405–2429 (2019).

6.   J. Lehmann, D. A. Bossio, I. Kögel-Knabner, M. C. Rillig, The concept and future

prospects of soil health. *Nat. Rev. Earth Environ.* **1**, 544–553 (2020).

7.   R. C. Wilhelm, H. M. van Es, D. H. Buckley, Predicting measures of soil health using the

microbiome and supervised machine learning. *Soil Biol. Biochem.* **164**, 108472 (2022).

8.   S. L. Pimm, The complexity and stability of ecosystems. *Nature* **307**, 321–326 (1984).

9.   A. Shade, *et al.*, Fundamentals of Microbial Community Resistance and Resilience .

Front. Microbiol.   **3** (2012).

10.   B. S. Griffiths, L. Philippot, Insights into the resistance and resilience of the soil microbial

community. *FEMS Microbiol. Rev.* **37**, 112–129 (2013).

11.   P. Laurent, G. B. S., L. Silke, Microbial Community Resilience across Ecosystems and

Multiple Disturbances. *Microbiol. Mol. Biol. Rev.* **85**, e00026-20 (2021).

12.   J. A. Harris, Measurements of the soil microbial community for estimating the success of

restoration. *Eur. J. Soil Sci.* **54**, 801–808 (2003).

13.   R. D. Bardgett, T. Caruso, Soil microbial community responses to climate extremes:

resistance, resilience and transitions to alternative states. *Philos. Trans. R. Soc. B Biol. Sci.*

375, 20190112 (2020).

14.   M. Vellend, Conceptual Synthesis in Community Ecology. *Q. Rev. Biol.* **85**, 183–206

(2010).

15.   D. R. Nemergut, *et al.*, Patterns and Processes of Microbial Community Assembly.

Microbiol. Mol. Biol. Rev. **77**, 342–356 (2013).

16.   Z. T. Aanderud, S. E. Jones, N. Fierer, J. T. Lennon, Resuscitation of the rare biosphere

contributes to pulses of ecosystem activity. *Front. Microbiol.* **6**, 24 (2015).

17.   S. E. Jones, J. T. Lennon, Dormancy contributes to the maintenance of microbial diversity.

Proc. Natl. Acad. Sci. **107**, 5881 LP – 5886 (2010).

18.   J. T. Lennon, S. E. Jones, Microbial seed banks: the ecological and evolutionary

implications of dormancy. *Nat. Rev. Microbiol.* **9**, 119–130 (2011).

19.   P. J. Kearns, A. Shade, Trait-based patterns of microbial dynamics in dormancy potential

and heterotrophic strategy: case studies of resource-based and post-press succession.

ISME J. **12**, 2575–2581 (2018).

20.   S. D. Allison, J. B. H. Martiny, Resistance, resilience, and redundancy in microbial

communities. *Proc. Natl. Acad. Sci.* **105**, 11512–11519 (2008).

21.   A. Shade, Understanding Microbiome Stability in a Changing World. *mSystems* **3**,

e00157-17 (2018).

22.   J. M. Elick, Mapping the coal fire at Centralia, Pa using thermal infrared imagery. *Int. J.*

Coal Geol. **87**, 197–203 (2011).

23.   T. Tobin-Janzen, *et al.*, Nitrogen changes and domain Bacteria ribotype diversity in soils

overlaying the Centralia, Pennsylvania underground coal mine fire. *Soil Sci.* **170** (2005).

24.   C. Janzen, T. Tobin-Janzen, "Microbial Communities in Fire-Affected Soils" in

Microbiology of Extreme Soils, P. Dion, C. S. Nautiyal, Eds. (Springer Berlin Heidelberg,

2008), pp. 299–316.

25.   S.-H. Lee, J. W. Sorensen, K. L. Grady, T. C. Tobin, A. Shade, Divergent extremes but

convergent recovery of bacterial and archaeal soil communities to an ongoing

subterranean coal mine fire. *ISME J.* **11**, 1447–1459 (2017).

26.   J. W. Sorensen, A. Shade, Dormancy dynamics and dispersal contribute to soil

microbiome resilience. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190255 (2020).

27.   E. Blagodatskaya, Y. Kuzyakov, Active microorganisms in soil: Critical review of

estimation criteria and approaches. *Soil Biol. Biochem.* **67**, 192–211 (2013).

28.   B. A. W., K. P. J., S. Ashley, 16S rRNA/rRNA Gene Ratios and Cell Activity Staining

Reveal Consistent Patterns of Microbial Activity in Plant-Associated Soil. *mSystems* **4**,

e00003-19 (2019).

29.   K. A. Lamothe, K. M. Somers, D. A. Jackson, Linking the ball-and-cup analogy and

ordination trajectories to describe ecosystem stability, resistance, and resilience.

Ecosphere **10**, e02629 (2019).

30.   B. E. Beisner, D. T. Haydon, K. Cuddington, Alternative stable states in ecology. *Front.*

Ecol. Environ. **1**, 376–382 (2003).

31.   A. Shade, *et al.*, Conditionally Rare Taxa Disproportionately Contribute to Temporal

Changes in Microbial Diversity. *MBio* **5**, e01371-14 (2014).

32.   J. C. Stegen, *et al.*, Quantifying community assembly processes and identifying features

that impose them. *Isme J.* **7**, 2069 (2013).

33.   F. Dini-Andreote, J. C. Stegen, J. D. van Elsas, J. F. Salles, Disentangling mechanisms

that mediate the balance between stochastic and deterministic processes in microbial

succession. *Proc. Natl. Acad. Sci.* **112**, E1326 LP-E1332 (2015).

34.   M. B. N. Albright, J. B. H. Martiny, Dispersal alters bacterial diversity and composition in

a natural community. *ISME J.* **12**, 296–299 (2018).

35.   S. E. Barnett, N. D. Youngblut, D. H. Buckley, Soil characteristics and land-use drive

bacterial community assembly patterns. *FEMS Microbiol. Ecol.* **96**, fiz194 (2020).

36.   S. E. Evans, L. P. Bell-Dereske, K. M. Dougherty, H. A. Kittredge, Dispersal alters soil

microbial community response to drought. *Environ. Microbiol.* **22**, 905–916 (2020).

37.   K. E. Walters, *et al.*, Routes and rates of bacterial dispersal impact surface soil

microbiome composition and functioning. *ISME J.* (2022) https:/doi.org/10.1038/s41396-

022-01269-w.

38.   E. M. Morrissey, *et al.*, Dynamics of extracellular DNA decomposition and bacterial

community composition in soil. *Soil Biol. Biochem.* **86**, 42–49 (2015).

39.   P. Carini, *et al.*, Relic DNA is abundant in soil and obscures estimates of soil microbial

diversity. *Nat. Microbiol.* **2**, 16242 (2017).

40.   J. T. Lennon, M. E. Muscarella, S. A. Placella, B. K. Lehmkuhl, How, when, and where

relic DNA affects microbial diversity. *MBio* **9**, e00637-18 (2018).

41.   M. Mestre, J. Höfer, The Microbial Conveyor Belt: Connecting the Globe through

Dispersion and Dormancy. *Trends Microbiol.* **29**, 482–492 (2021).

42.   M. J. Choudoir, K. M. DeAngelis, A framework for integrating microbial dispersal modes

666   into soil ecosystem ecology. *iScience* **25**, 103887 (2022).

667   43.   B. Moreira-Grez, *et al.*, Reconditioning Degraded Mine Site Soils With Exogenous Soil

668   Microbes: Plant Fitness and Soil Microbiome Outcomes   . *Front. Microbiol.*   **10** (2019).

669   44.   W. L. King, T. H. Bell, Can dispersal be leveraged to improve microbial inoculant

670   success? *Trends Biotechnol.* **40**, 12–21 (2022).

671   45.   N. Fierer, R. B. Jackson, The diversity and biogeography of soil bacterial communities.

672   *Proc. Natl. Acad. Sci.* **103**, 626–631 (2006).

673   46.   A. A. Malik, B. C. Thomson, A. S. Whiteley, M. Bailey, R. I. Griffiths, Bacterial

674   Physiological Adaptations to Contrasting Edaphic Conditions Identified Using Landscape

675   Scale Metagenomics. *MBio* **8**, e00799-17 (2017).

676   47.   B. M. Tripathi, *et al.*, Soil pH mediates the balance between stochastic and deterministic

677   assembly of bacteria. *ISME J.* **12**, 1072–1083 (2018).

678   48.   J. B. Yavitt, C. A. Roco, S. J. Debenport, S. E. Barnett, J. P. Shapleigh, Community

679   Organization and Metagenomics of Bacterial Assemblages Across Local Scale pH

680   Gradients in Northern Forest Soils. *Microb. Ecol.* (2020) https:/doi.org/10.1007/s00248-

681   020-01613-7.

682   49.   J. C. Stegen, X. Lin, A. E. Konopka, J. K. Fredrickson, Stochastic and deterministic

683   assembly processes in subsurface microbial communities. *ISME J.* **6**, 1653–1664 (2012).

684   50.   J. C. Stegen, X. Lin, J. K. Fredrickson, A. E. Konopka, Estimating and mapping

685   ecological processes influencing microbial community assembly. *Front. Microbiol.* **6**, 370

686   (2015).

51.   K. C. Rice, J. S. Herman, Acidification of Earth: An assessment across mechanisms and

scales. *Appl. Geochemistry* **27**, 1–14 (2012).

52. H. Wang, J. Zhang, L. Zhang, J. Wang, Z. Xu, Gas Emission and Soil Chemical Properties

Associated with Underground Coal Fires, Wuda Coalfield, Inner Mongolia, China. *Nat.*

Resour. Res. **29**, 3973–3985 (2020).

53. P. D. de Quadros, *et al.*, Coal mining practices reduce the microbial biomass, richness and

diversity of soil. *Appl. Soil Ecol.* **98**, 195–203 (2016).

54. M. R. Rondon, R. M. Goodman, J. Handelsman, The Earth's bounty: assessing and

accessing soil microbial diversity. *Trends Biotechnol.* **17**, 403–409 (1999).

55. S. H. Sirois, D. H. Buckley, Factors governing extracellular DNA degradation dynamics

in soil. *Environ. Microbiol. Rep.* (2018) https:/doi.org/10.1111/1758-2229.12725.

56. R. I. Griffiths, A. S. Whiteley, A. G. O'Donnell, M. J. Bailey, Rapid Method for

Coextraction of DNA and RNA from Natural Environments for Analysis of Ribosomal

DNA- and rRNA-Based Microbial Community Composition. *Appl. Environ. Microbiol.*

66, 5488–5491 (2000).

57. J. Colovas, A. F. Bintarti, M. E. Mechan Llontop, K. L. Grady, A. Shade, Do-it-Yourself

Mock Community Standard for Multi-Step Assessment of Microbiome Protocols. *Curr.*

Protoc. **2**, e533 (2022).

58. J. J. Kozich, S. L. Westcott, N. T. Baxter, S. K. Highlander, P. D. Schloss, Development

of a dual-index sequencing strategy and curation pipeline for analyzing amplicon

sequence data on the MiSeq Illumina sequencing platform. *Appl. Environ. Microbiol.* **79**,

5112–5120 (2013).

59. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10 (2011).

60.   J. Zhang, K. Kobert, T. Flouri, A. Stamatakis, PEAR: a fast and accurate Illumina Paired-

End reAd mergeR. *Bioinformatics* **30**, 614–620 (2014).

61.   R. C. Edgar, Search and clustering orders of magnitude faster than BLAST.

Bioinformatics **26**, 2460–2461 (2010).

62.   P. D. Schloss, *et al.*, Introducing mothur: open-source, platform-independent, community-

supported software for describing and comparing microbial communities. *Appl. Environ.*

Microbiol. **75**, 7537–7541 (2009).

63.   T. Rognes, T. Flouri, B. Nichols, C. Quince, F. Mahé, VSEARCH: a versatile open source

tool for metagenomics. *PeerJ* **4**, e2584 (2016).

64.   M. N. Price, P. S. Dehal, A. P. Arkin, FastTree: Computing large minimum evolution

trees with profiles instead of a distance matrix. *Mol. Biol. Evol.* **26**, 1641–1650 (2009).

65.   E. P. Nawrocki, "Structural RNA Homology Search and Alignment using Covariance

Models," Washington University in Saint Louis. (2009).

66.   P. J. McMurdie, S. Holmes, phyloseq: An R package for reproducible interactive analysis

and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

67.   S. I. Glassman, J. B. H. Martiny, Broadscale Ecological Patterns Are Robust to Use of

Exact Sequence Variants versus Operational Taxonomic Units. *mSphere* **3**, e00148-18

(2018).

68.   P. D. Schloss, Amplicon Sequence Variants Artificially Split Bacterial Genomes into

Separate Clusters. *mSphere* **6**, e00191-21 (2021).

69.   R Core Team, R: A language and environment for statistical computing (2018).

70.   J. Pinheiro, D. Bates, S. DebRoy, D. Sarkar, R Core Team, nlme: Linear and Nonlinear

Mixed Effects Models (2020).

71.   M. De Cáceres, *et al.*, Trajectory analysis in community ecology. *Ecol. Monogr.* **89**,

735   e01350 (2019).

736   72.   A. Sturbois, *et al.*, Extending community trajectory analysis: New metrics and

737   representation. *Ecol. Modell.* **440**, 109400 (2021).

738   73.   J. R. Conway, A. Lex, N. Gehlenborg, UpSetR: an R package for the visualization of

739   intersecting sets and their properties. *Bioinformatics* **33**, 2938–2940 (2017).

740   74.   D. Ning, *et al.*, A quantitative framework reveals ecological drivers of grassland microbial

741   community assembly in response to warming. *Nat. Commun.* **11**, 4717 (2020).

Community type	Factor	Degrees of freedom	Sum of Squares	R^2	F value	p-value
Whole community	Fire Classification	2	0.497	0.180	12.811	0.001
	Year	6	0.157	0.057	1.350	0.001
	Fire Classification: Year	12	0.110	0.040	0.474	0.002
	Residual	103	1.997	0.723		
	Total	123	2.762	1.000		
Active community	Fire Classification	2	0.343	0.124	6.793	0.001
	Year	6	0.158	0.057	1.042	0.002
	Fire Classification: Year	12	0.184	0.067	0.608	0.18
	Residual	82	2.068	0.751		
	Total	102	2.753	1.000		

744
745 **Table 1:** PERMANOVA results comparing whole bacterial community or active subset structure
746 across samples based on abundance weighted UniFrac distance blocked by site ID.
747

Figure 1: Across the fire-affected sites, soils tended to cool over time, and along with this cooling, bacterial alpha diversity increased. A) Map of collection sites in Centralia, Pennsylvania, USA. B) Tracking of soil temperature (˚C) over the seven sampling years for each site showed general soil cooling in fire-affected sites. C) Whole bacterial community phylogenetic diversity (Faith's PD) decreased as soil core temperature increased. D) Active bacterial community subset phylogenetic diversity decreased as soil core temperature increased. For all plots, colored points and lines indicate site and fire classification. For plots B-D, white and black lines indicate linear mixed-effects regressions using site ID as the random effect, with solid white lines indicating statistically significant slopes (p-value < 0.05) and dashed lines indicating non-statistical significance (p-value ≥ 0.05). Full statistics are found in tables S2-S4, and phylogenetic diversity trends over time are found in Fig. S2.

767
768
Figure 2: Time-lag analysis demonstrated that variation in both whole bacterial community and
active subset structure between samplings of the same site increased as the time difference
between sampling increased. However, this trend was only statically significant for fire-affected
sites. A) Whole community time-lag analysis. B) Active community subset time-lag analysis. C)
Slopes of whole community time-lag regressions made individually for sites increased as
maximum soil temperature (proxy of disturbance intensity) increased across sites. D) Slopes of
active community subset time-lag regressions made individually for sites increased as maximum
soil temperature increased across sites. Points indicate site ID and fire classification. White and
black lines indicate linear mixed-effects regressions using site ID as the random effect (A and B)
and linear regression (C and D) with solid white lines indicating statistically significant slopes
(p-value < 0.05) and dashed lines indicating non-statistical significance (p-value ≥ 0.05). Full
statistics for the linear mixed effects regressions are provided in Table S7.
781
782

783
784
Figure 3: Trajectory analysis demonstrated that directionality in both whole bacterial
community (A) and active subset (B) structural shifts over time were generally highest in fire-
affected soils compared to recovered and reference soils. Post-hoc pairwise Dunn's test p-values
< 0.05 are displayed above statistical significance indicator bars. Directionality in both whole
bacterial community (C) and active subset (D) structural shifts increased as maximum soil
temperature within sites increased. White lines indicate linear regressions, which had statistically
significant slopes (p-value < 0.05). For all plots, points indicate site ID and fire classification.
792
793

794
795
Figure 4: For both (A) whole bacterial communities and (B) active subsets, warmer soils were
more structurally distinct from the undisturbed reference soils. Structural dissimilarity was
calculated as the mean weighted UniFrac distance between fire-affected and recovered sites and
the three reference sites paired within the sampling year. White lines indicate linear mixed-
effects regressions using site ID as the random effect, which had statistically significant slopes
(p-value < 0.05). Points indicate site ID and fire classification (fire affected or recovered,) and
error bars represent the standard error for the distance between a tested sample and the three
paired reference samples.
804

806
807
Figure 5: For fire-affected sites, the whole bacterial community structural dissimilarity
attributable to the active subset decreases as the time between sampling time points increases. In
other words, less of the whole community dissimilarity is attributable to the active subset across
larger timescales for fire-affected sites but not for recovered or reference sites. This holds across
site pairs for OTUs active in (A) the early timepoint (*e.g.,* 2016 when comparing 2016 and 2020)
and in (B) the later timepoint (*e.g.,* 2020 when comparing 2016 and 2020). Structural
dissimilarity over time is measured by the Bray-Curtis dissimilarity between years within a site,
partitioning OTUs active in either the early or late time points of each pair. White and black lines

indicate linear mixed-effects regressions using site ID as the random effect, with solid white
lines indicating statistically significant slopes (p-value < 0.05) and dashed lines indicating non-
statistical significance (p-value ≥ 0.05).
819
820

821
822
Figure 6: UpSet plots of the top 20 (as determined by OTU membership) activity across years
groups for (A) fire-affected sites and (B) combined recovered and reference sites. All sites
sampled each year were combined such that OTUs were defined as active if they had an RNA:
DNA ≥ 1 in at least one site within that year. The bottom right plots show the activity patterns of
the top 20 groups, with dots indicating activity within that year and groups ordered by OTU
membership. The top vertical bar chart shows the number of OTUs within each activity group,
colored by the top 10 phyla (class for *Proteobacteria*). The bottom left horizontal bar chart

shows the number of active OTUs within each year, colored by their membership in the top 20 activity groups. White sections of the bars indicate OTUs from the other 107 groups (see Figs. S8 and S9 for all 127 activity groups). For the fire-affected sites (A), most of the top 20 activity groups do not have detectable activity in the first few sampled years, while most of the top 20 activity groups for reference and recovered sites (B) have detectable activity during those years.

Figure 7: whole bacterial community assembly dynamics across sites as measured with standardized phylogenetic and structural turnover. A) βNTI is generally higher between fire-affected sites than recovered or reference sites whose site pairs almost always have a βNTI < 2, indicating homogeneous selection. Post hoc pairwise Dunn's test p-values < 0.05 are displayed

above statistical significance indicator bars. B) Regardless of fire classification, βNTI tended to
increase as the difference in soil pH between sites increased, indicating pH-related selection
pressures in these soils. C) Regardless of fire classification, βNTI tended to increase as the
difference in soil temperature between sites increased, indicating some temperature-related
selection pressures in these soils. Linear regressions, including these two relationships and their
interaction, are provided in Table S9. D) Percentage of site pairs within each fire classification
and year for which community assembly can be attributed to the five assembly processes
examined with βNTI and RC_{bray}. The numbers of site pairs are indicated by white numbers at the
bottom of each bar. All site pairs were within each year, not across years.