Birth of immunity in early life: stepwise emergence of resistance and immunity to complex molecular parasites via progressively degenerating hyperparasites

Short title:

parasite immunity from hyperparasitism

Authors:

Magnus Pirovino[1], Christian Iseli[2], Joseph A. Curran[3], Bernard Conrad[4*]

Affiliations: [1]OPIRO Consulting Ltd., FL-9495 Triesen, Principality of Liechtenstein, [2]Bioinformatics Competence Center, EPFL and Unil, CH-1015 Lausanne, [3]Department of Microbiology and Molecular Medicine, and Institute of Genetics and Genomics of Geneva (iGE3), Geneva University, CH-1211 Genève 4, [4]Genesupport, Avenue de Sévelin 18, CH-1004 Lausanne

*Correspondence to: bernard.conrad@genesupport.ch

Autocatalytic networks likely played a central role in the shift from inanimate systems to protocells.
Using an established hyperparasite framework, we model the emergence of antiparasite immunity
using the Lotka Volterra equations. It builds on autocatalytic, increasingly complexed RNA networks
reflecting the historical exposure to ubiquitous molecular parasites. First, upon parasite encounter,
specific catalysts providing parasite resistance appear in primordial autocatalytic cycles, supplanted
by promiscuous, efficiently self-replicating, but parasite sensitive ribozymes. This is supported by
catalyst tradeoff analysis. Substrate promiscuity confers high catalytic activity, but entails parasite
exposure, higher substrate specificity implies relatively lower activity, yet it offers parasite resistance.
Second, sustained parasite influx generates hyperparasite cycles, reactivating the original, parasite-
specific catalyst-subunit that now once more provides parasite resistance, in a homeostatically
stabilized habitat, empirically consistent with phage satellites and defective RNA viruses induced
across a range of parasite loads. Third, under continuous exposure to progressively elaborate parasite
populations, hyperparasites degenerate and embody antiparasite immunity. Again, the observed
triggering of antiviral immunity by phage satellite encoded antiphage systems and defective RNA
virus genomes are in agreement with this model. As such, it offers an attractive and unifying theory
for the spontaneous birth of immunity in early life.

Introduction

The multistep transition process from inanimate to living systems has been dated to >3.9 Ga [1]. A
non-exhaustive list of thresholds that needed to be progressively surmounted during this process has
been drawn, comprising, among others, chirality symmetry breaking, spontaneous polymerization,
self-assembly of compartments, and particularly the catalytic closure threshold [2]. One central pillar
in the theory of how early life could have materialized pertains to the "RNA world". This stipulates
that the initial informational polymer was RNA, and that the same molecule was also the
fundamental catalyst [3,4]. The underlying concept of autocatalytic sets was laid out more than half a
century ago [5,6]. It was subsequently refined [7] and validated in a proof of principle experiment
showing that RNA fragments could build self-replicating ribozymes from individual RNA fragments via
cooperative catalytic cycles [8]. In parallel, an RNA polymerase ribozyme was generated by *in vitro*
evolution that exhibited a high level of processivity [9]. Collectively, both theoretical considerations
and experimental evidence support the view of a gradual evolution towards RNA self-catalysis. This
started with self-assembly, followed by the formation of autocatalytic networks using self-assembled
substrates. From this emerged different ribozymes (recombinases, ligases) and template based
ligation, leading to a template based, self-replicase catalyzed assembly of monomers [10].

Prebiotic heredity in chemical networks is theorized to comprise variable autocatalytic cycles
that form a core (genotype) and a non-catalytic periphery (phenotype), allowing limited heredity and
selectability (11). Complexed mixtures of cross-replicating RNA enzymes were shown to undergo self-
sustained Darwinian evolution with the appearance of both, highly active enzymes that maintained
the population, and poorly active enzyme passengers [12]. In conclusion, the coexistence of prebiotic
heredity with an emerging, template-based canonical Darwinian heredity appears plausible [10].

The ribozyme catalyzed RNA-polymerization process is inherently error prone [9], generating
variant sequence space mutants called a quasispecies [13]. Mutants vary in frequency, length,
sequence and structural-complexity, as well as in ribozyme activity, thus providing selectable traits in
the dimensional fitness landscape, epitomizing the mutation-selection balance [14]. This variant
sequence-space intrinsically implies the obligate presence and evolutionary persistence of the
molecular parasites associated with all life forms, as argued essentially for thermodynamic reasons
[15]. In precellular life, the most abundant class of genetic parasites were most likely capsidless.
These subsequently gave rise to capsidless selfish genetic elements and to capsid-encoding viruses
that diverged from cellular, ribosome-encoding organisms [16]. Unsurprisingly, nucleic acid
polymerase evolution also most likely bifurcated along the same paths based on distinct RNA
recognition motifs [17].

It has been noted within the realm of the virosphere - in all Baltimore classes [18] - that
genetic parasites tend to be parasitized, typically forming a tripartite relationship. This is exemplified
by bacteriophages (phages), notorious parasites of bacteria that are themselves parasitized by phage
satellites [19,20]. Other typical threesome associations are observed in RNA viruses and
Nucleocytoplasmic Large DNA viruses (NCLDVs): for RNA viruses, it comprises the host, the RNA virus
and defective viral genomes [21], for NCLDLVs it is the eukaryotic host, the NCLDV and its dedicated
virophage from the same group of Bamfordviruses [22].

We previously developed a hyperparasite framework based on the Lotka Volterra equations
[23], demonstrating that superior homeostasis arose from the tripartite configuration in a given
habitat, namely, the host, the parasite and the hyperparasite, i.e. the parasite of the parasite. The
aforementioned empirical evidence lends substantial credit to this. This reasoning has now led us to
extend the model, building on complexed autocatalytic RNA networks with prebiotic heredity
reflecting the historical parasite exposure, and showing, how parasite resistance and antiparasite
immunity can be selected for and gradually emerge from progressively degenerating hyperparasites
in an intrinsic fashion in early life, in hosts exposed to increasingly more complexed parasite
populations.

97
98

Methods

Definitions and Terms

In what follows we assume the following: the Malthusian fitness r of a replicating (biochemical)

system (or species) is defined by $\frac{d}{dt} R = \dot{R} = r\,R$, where R denotes the population size of the

replication sys-tem or species. Evolutionary dynamics between two or more interacting species or

systems, e.g. $R, S, P, \ldots$ are described by the following type of replicator differential equation e.g. for

R:

$\dot{R} =$ "(A) terms of dynamics using habitat resources" x "(B) term of habitat resource restriction" + "(C)

terms of dynamics not using habitat resources"

Terms A include both "terms of first order kinetics" and "terms of second order kinetics"

— "Terms of first order kinetics" are typically natural growth or decay rates, e.g. $\dot{R} = r\,R$. If $r >$

0, growth process described by rR eats into resources, whereas if $r < 0$, r is a decay rate, and

de-cay dynamics does not.

Terms of second order kinetics are derived from encounters between different species. For

instance,   the   term   "bRS"   in   "$\dot{R} = -aR + bRS$" has   the   following   meaning:

$bRS =$

$(probabiliy\ of\ an\ encounter\ between\ members\ of\ R\ and\ S)$ x $(impact\ of\ encounter\ on\ the\ fitness\ of\ R)$.

Here we assume, according to the Lotka Volterra calculus that the probability of an encounter

between members of R and S is proportional to the product RS.

The term B of resource restriction is used in a habitat of limited resources. For instance, in the

classic logistic equation $\dot{R} = r\,R\left(1 - \frac{R}{K}\right)$ the term $\left(1 - \frac{R}{K}\right)$ indicates that the habitat has limited

resources supporting a maximum population K for R.

Since every encounter involves resource consuming, we assume here, that all terms of second

order kinetics are subject to resource restrictions and therefore need to be multiplied with the

term B.

Example of a host-parasite interaction (R=host, P=parasite):

$$\dot{R} = (a\,R - bPR)\left(1 - \frac{R+P}{K}\right)$$

$$\dot{P} = -c\,P + dRP\left(1 - \frac{R+P}{K}\right)$$

125   where $a > 0, b > 0, c > 0, d > 0$. The first order dynamics growth process aR and all the second

126   order processes eat into the habitat's resources, whereas the first order decay process $-c\,P$ does

127   not. The habitat resources restriction is represented by the term $\left(1 - \frac{R+P}{K}\right)$, indicating that a

128   maximum population size of $K = R + P$ can be nourished by the habitat and that both, S and P,

129   compete for the same resources.

Definition of catalysts, ribozymes and enzymes

We use the term _catalyst_, sometimes also _perfect catalyst_, for any substance that accelerates the rate of a reaction without undergoing any change during the process. We call an _imperfect catalyst_ a substance that accelerates the rate of a reaction, but that may undergo a change during the process; antibodies in the transition from hyperparasitism to antiparasitism are examples of imperfect catalysts (see process (3'), Suppl. Inf.). C _atalysts_ displaying relatively higher substrate specificity generally have relatively lower catalytic activity, for instance the protoreplicase. A subgroup of catalysts is called here _enzymes_, if they are generated based on RNA templates. An enzyme may be either a _ribozyme_, if it is also used as an RNA template to generate other enzymes, or a _pure enzyme_ if it cannot be used as an RNA template for other polymerization processes. The term _ribozyme_ encompasses different enzymatic activities, such as recombinase, ligase and polymerase. Recombinases and ligases predominate in early protocycles, while polymerases appear only in advanced networks.

Model $1^{(1)}$: habitat with one host autocatalytic cycle exposed to parasite invasion

$$\frac{dR_2}{dt} = a\,R_1 R_1 \Pi - d_{R_2} R_2$$

$$\frac{dR_1}{dt} = (b\,R_2 R_1 - \delta R_2 P)\Pi - d_{R_1} R_1$$

$$\frac{dP}{dt} = c R_2 P \Pi + \alpha - d_P P,$$

where a, b and c are positive constants, d_{R_i}, d_P are the decay rates of populations R_i, $i = 1, 2$, and of parasite P absent its host R_2; δ is parasite's P rate of exploitation of the host replicase R_2; and the habitat restriction term is defined as $\Pi = \left(1 - \frac{R_1 + R_2 + P/q_P}{K}\right)$, where $q_P > 1$, since typically parasites P are more efficient using the resources of the habitat than their hosts R. α is the parasite invasion parameter, i.e., per unit of time the habitat is invaded by α population units of parasites P.

Model $1^{(0)}$ habitat with primordial autocatalytic cycle, exposed to template invasion P (to become parasite later in evolution) and protoenzyme E_1:

$$\frac{dP}{dt} = \alpha - d_P P$$

$$\frac{dE_1}{dt} = \beta - d_{E_1} E_1$$

$$\frac{dE_2}{dt} = a E_1 F_1 \Pi - d_{E_2} E_2$$

$$\frac{dF_1}{dt} = bE_2 P \Pi - d_{F_1} F_1,$$

where

E_2 is the protoreplicase produced by this primitive autocatalytic cycle,

F_1 is the prototemplate produced by this primitive autocatalytic cycle,

α is the inflow rate of RNA-template P into the habitat,

β is the inflow rate of protoenzyme/catalyst E_1 into the habitat,

a and b are positive constants,

d_{E_i}, d_{F_1}, d_P are the decay rates of enzyme populations $E_i, i = 1, 2$, of prototemplate d_{F_1}, and of

RNA-template P,

$\Pi = \left(1 - \frac{E_2 + F_1/q_{F_1}}{K}\right)$ is the habitat restriction term where $q_{F_1} > 1$, since we assume that proto-

templates F_1 are more efficient using the resources of the habitat than the more complex

protoreplicase E_2.

Model 1$^{(1+)}$: stabilized host-parasite-hyperparasite cycle

$$\frac{dR_2}{dt} = (a\,R_1 R_1 - \delta_2 R_1 F_1)\Pi - d_{R_2} R_2$$

$$\frac{dR_1}{dt} = (b\,R_2 R_1 - \delta_1 R_2 P)\Pi - d_{R_1} R_1$$

$$\frac{dP}{dt} = (cR_2 P - \delta_3 F_2 P)\Pi + \alpha - d_P P$$

$$\frac{dF_1}{dt} = (d\,F_2 P + \beta R_2 P)\Pi - d_{F_1} F_1$$

$$\frac{dF_2}{dt} = (e\,R_1 F_1)\Pi - d_{F_2} F_2,$$

where

a, b, c, d, e are positive (reproduction) constants,

R_1 is a ribozyme of the autocatalytic host cycle,

R_2 is the replicase, i.e. another ribozyme (or may be also pure enzyme since it is not used as

template) of the autocatalytic host cycle,

P is the parasite, exploiting replicase R_2 of the host cycle (exploitation parameter δ_1),

F_1 is the prototemplate exploiting the host R_1 (exploitation parameter δ_2) for protoreplicase F_2

production, triggered upon encounter of R_2 with P (trigger parameter β) and further produced by

protoreplicase F_2 upon encounter with P,

F_2 is the protoreplicase (pure enzyme or ribozyme) producing F_1 upon encounter with parasite temp-

lates P ($F_2 = E_2$ from the primitive autocatalytic cycle of Model $1^{(0)}$), thus, exploiting template of

parasite P (exploitation parameter δ_3),

d_{R_i}, d_{F_i}, d_P are the decay rates of populations $R_i, F_i, i = 1, 2$, and of P, absent new inflow or

reproduction in the habitat,

α is the inflow rate of RNA template P into the habitat,

β is trigger parameter of replicase R_2 producing F_1 by encounter with parasite P,

$\Pi = \left(1 - \frac{R_1 + R_2 + P/q_P + F_1/q_{F_1} + F_2/q_{F_2}}{K}\right)$ is the habitat restriction term where $q_P, q_{F_1}, q_{F_2} > 1$, since we

assume that parasitizing cycles of P and F_1, F_2 are more efficient using the resources of the habitat

than the more complex host cycle R_1, R_2.

Model $1^{(2)}$: integrated host-parasite-hyperparasite cycle without new parasite inflow

The simulation results of Model $1^{(2)}$ are realized as follows: Model $1^{(1+)}$ provides the starting basis,

with arbitrary initial populations to establish a stabilized situation, given a moderate parasite inflow

with $\alpha > 0$ (Fig. 3lefthand part). Subsequently, Model $1^{(1+)}$ is adapted to Model $1^{(2)}$. All the

parameters of Model $1^{(2)}$ are the same as those of Model $1^{(1+)}$, with the exception of the parasite

inflow, now $\alpha = 0$. As initial conditions for the populations of Model $1^{(2)}$, we assume the terminal

populations of the simulations of Model $1^{(2)}$, as calculated for Fig. 3 at the beginning of the righthand

part. The larger fluctuations of the population at the beginning of the righthand part of Fig. 3 stem

from the transition from Model $1^{(1+)}$ to Model $1^{(2)}$, where the parasite inflow abruptly decreases

from $\alpha > 0$ to $\alpha = 0$. The fact that parasites in Model $1^{(2)}$ are now reproduced without new inflow

shows that they can be viewed as fully integrated into the autocatalytic network, realized in $1^{(2)}$ (see

Suppl. Inf.).

Model $1^{(1-)}$: transition from hyperparasitism to antiparasitism

$$\frac{dR_2}{dt} = (a\,R_1 R_1 - \delta_2 R_1 F_1)\Pi - d_{R_2} R_2$$

$$\frac{dR_1}{dt} = (b\,R_2 R_1 - \delta_1 R_2 P)\Pi - d_{R_1} R_1$$

$$\frac{dP}{dt} = (cR_2 P - \delta_3 F_2 P)\Pi + \alpha - \theta_P F_2 P - d_P P$$

$$\frac{dF_1}{dt} = (d\,F_2 P + \beta R_2 P)\Pi - d_{F_1} F_1$$

$$\frac{dF_2}{dt} = (e\,R_1 F_1)\Pi - \theta_{F_2} F_2 P - d_{F_2} F_2,$$

Where all the parameters are the same as in Model $1^{(1+)}$, with the exception that now two additional

parameters are added, namely θ_P and θ_{F_2}. The term $\theta_P F_2 P$ denotes the number of population units

of parasites P made unviable per unit of time during the encounter of protoreplicase F_2 with P, now

also functioning as antibody. In addition, the term $\theta_{F_2} F_2 P$ denotes the number of population units of

protoreplicase/antibody F_2 made unviable per unit of time during the encounter of

protoreplicase/antibody F_2 with P. For instance, if $\theta_p = \theta_{F_2}$, then the same number of antibodies and

parasites are made unviable during their encounter. In the evolutionary transition from

hyperparasitism to antiparasitism F_2 can still act both ways, as imperfect protoreplicase and as

antibody directed against the parasite P. We call it *imperfect* protoreplicase or imperfect catalyst,

because it still produces/catalyzes prototemplate F_1 upon encounter with template P. But now F_2

may be damaged during this process, therefore cannot be called a *perfect* catalyst anymore. The

replication differential equations of Model $1^{(1-)}$ realizing a pure antiparasitism situation are the

same as in Model $1^{(1-)}$ of Fig. 4A, except that the reproduction parameters hyperparasite d and δ_3

are now set to zero: $\delta_3 = d = 0$.

Results

In a stepwise and gradual process outlined in the Supplementary Information (proofs, Suppl. Inf.) and

Methods sections, we lay out using the Lotka Volterra equations, how simple autocatalytic cycles are

initially built and how over time they become increasingly more elaborated as a result of molecular

parasite exposure. The coexistence of prebiotic heredity with an emerging canonical, template-based

Darwinian heredity [10-12] allows for the complexification of the quasispecies that in turn reflects its

historical parasite exposure. The previously established tripartite habitat framework featuring a host,

a parasite that is itself parasitized (hyperparasite; [23]), is instrumental for homeostasis and the

emergence of resistance, and finally immunity to the increasingly more exhaustive parasite

populations.

Autocatalytic cycles in transitional early RNA-settings

The seminal duality of the RNA world, namely the fact that the same RNA molecule provides both the

template and the self-replicating catalyst [3,4] constitutes the foundation of the initial autocatalytic

cycles (replication equations (1.1), (1.2), forming a two-step autocatalytic cycle; proof outlined in

Suppl. Inf.). Since the template and the enzyme/ribozyme remain in principle unchanged during this

process, both can be viewed as catalysts, as long as the molecules do not undergo deleterious loss of

function mutations.

Parasitism in autocatalytic cycles

Molecular parasites are an inherent feature of the replication process, they are inseparably

associated with all life forms [24], for a number of reasons, the most important being the accuracy-

rate tradeoff [25-26]. Average enzymes operate orders of magnitude (k_{cat}/K_M of $\sim 10^5$ s^{-1}M^{-1}) below

the diffusion rate (10^8-10^9 s^{-1}M^{-1}), which implies promiscuous substrate binding, high catalytic activity,

and as a secondary tradeoff a high mutation rate with a large variance in the quasispecies (speed

being more important than enzyme accuracy; [26,27]). In an initial autocatalytic cycle, self-replicating

ribozymes emerge (ribozyme R_2 acting on a replicating template R_1; proof outline in Suppl. Inf.;

Model $1^{(1)}$ in Methods). The catalyst tradeoff opposes specific, but catalytically relatively less active

molecules, and promiscuous, yet catalytically highly efficient molecules [26]. Therefore, the

catalytically very efficient and promiscuous R_2 will inevitably be parasitized by RNA templates P.

Figure 1 illustrates the population sizes of the habitat constituents under high (Fig. 1A), and low (Fig.

1B) parasite exposure. As expected, high parasite loads overwhelm the habitat with P templates, an

observation that begs a solution.

A stable, primordial autocatalytic cycle in the presence of parasites

In a primordial habitat with abundant inflow of molecular RNA species including parasites, a

quasispecies will eventually emerge with the central feature that it also comprises molecules with

superior parasite-specificity, which implies relatively lower catalytic activity, and confers parasite

resistance (protoreplicase E_2, prototemplate F_1; equations and proofs outlined in the Suppl. Inf.).

This composite feature progressively arises in successive rounds of autocatalytic cycles, initially

comprising the encounter of the catalyst E_1, provided by a neighboring habitat, with F_1 evolved

autocatalytically in the habitat, producing the primordial catalyst E_2, that is capable of amplifying F_1

but not P. It is the twofold consequence of the primordial parasite-specificity of E_2, and of the

complexity of P that exceeds the processivity of the protoreplicase. As a corollary and as displayed in

Fig. 2, selection of such molecules confers considerable resistance to parasite invasion, and restores

host template and catalyst abundance (Model $1^{(0)}$, Methods). There is both, strong theoretical and

experimental evidence to support these findings. Indeed, while it is generally assumed that catalysts

arising in early autocatalytic cycles have low activity and high promiscuity [25], an in depth physico-

chemical analysis of how enzymes meet demands of selectivity and accuracy points towards

accuracy-rate tradeoffs [26]. Selectivity (accuracy, specificity) implies the ability to discriminate

between two templates, provided both are present [26], and rate denotes speed [25,26]. Ground

state discrimination means that specificity is achieved mainly through substrate binding, which

imposes strong accuracy-rate tradeoffs [26]. Improvements in selectivity mediated by tighter cognate

substrate binding invariably lead to lower catalytic efficiency (parallel decreases in the constants for

the cognate substrates $K_M{}^{cog}$ and $k_{cat}{}^{cog}$; [26]). This is precisely what the model stipulates, the

protoreplicase has a primal parasite-selectivity, which comes at the cost of a relatively reduced

catalytic activity, yet it allows for adequate host molecule-, but not parasite-amplification. In

conclusion, if catalysts with relatively reduced function in the quasispecies confer parasite resistance,

yet restore host catalyst and template abundance, they will be positively selected [14].

Evolution of the primordial autocatalytic cycle into a full-fledged autocatalytic cycle

Natural selection in the mutant quasispecies space will ultimately produce, starting from E_2, a fully

efficient, self-replicating ribozyme (replicase R_2) that is bestowed with high catalytic activity [9],

capable of generating all enzymes and templates in the habitat (equations and proofs outlined in the

Suppl. Inf.). However, this has a cost. R_2 is promiscuous, catalytically highly efficient and therefore

exposed to parasite invasion, which was not the case for its primordial catalyst precursor with

relatively lower activity (protoreplicase E_2; Model $1^{(1+)}$ stabilized host-parasite-hyperparasite cycles,

Methods).

At this point, the protoreplicase E_2-subdomain of R_2 confers P selectivity, restoring parasite

resistance and rescuing the abundance of the partial catalyst and its templates, even with a moderate

parasite inflow ($\alpha > 0$, Fig. 3 left panel). This process has as its central feature a hyperparasite cycle

(equations and proofs outlined in the Suppl. Inf.). It is further noteworthy that F_2 (formerly E_2) is still

produced (Fig. 3), and is acting additively to R_2, which is important to generate antiparasitism later.

Under permanent and moderate, thus manageable parasite inflow ($P^{special}$, Suppl. Inf.), an

increasingly ramified network of autocatalytic cycles arrives in the habitat that persists even when

the parasite exposure ceases, constituting a historical record of past parasite exposures ($\alpha = 0$, Fig. 3

right panel).

There is considerable evidence in favor of the essential components of this model, in

particular the catalyst accuracy-rate tradeoff and the hyperparasite cycle. Firstly, a solid body of

theoretical and experimental evidence supports catalyst accuracy-rate tradeoffs [25,26]. Secondly,

metagenomic studies demonstrate the virtually universal presence of tripartite constituents across

molecular parasite habitat classes, most persuasively for bacteria-phages-phage satellites [19,20], for

RNA viruses partnering the host- virus-defective interfering RNAs [21], and for NCLDVs, with

eukaryotic host-NCLDV-virophage [22]. Thirdly, additional support comes from the observation that

hyperparasites are induced across the range of parasite concentrations, and especially at high

multiplicity infection for RNA viruses [28,29].

Emergence of antiparasitism in autocatalytic cycles

Under conditions, where the sheer number and complexity of incoming parasite populations would

overwhelm the carrying capacity of the habitat, a degenerative process is initiated that erodes

catalysis. Formally, the hyperparasite cycle degenerates to an antiparasite cycle (equations and proofs

outlined in the Suppl. Inf.; Model $1^{(1-)}$, Methods). In Figures 4 and 5, the transition from

hyperparasitism to antiparasitism is illustrated, in the sense that both coexist at equal activity, partial

catalyst and non-catalytic antiparasitic activities (Fig. 4A, see Methods for parameters). In Figure 4B,

antiparasitism is fully implemented; hyperparasitism has been fully replaced by the non-catalytic

antiparasitism (reproduction parameters hyperparasite d and δ_3 are set to zero: $\delta_3 = d = 0$), making

the difference between pure antiparasitism, and the transition from hyperparasitism to

antiparasitism minor. In Figure 4B, the population fluctuations are more volatile than in Figure 4A, but

pure antiparasitism does effectively stabilize the autocatalytic cycle, without a hyperparasite

component. Figure 5 shows the gradual transition from pure hyperparasitism (top panel) to

antiparastism (botton panel), in four successive steps (0, 0.25, 05. 0.75 and 1), progressively replacing

hyperparasitisim with antiparasitism. Therefore, we conclude that antiparasitism can replace

hyperparasitism in situations with a stronger and highly diverse parasite inflow that would overrun

the carrying capacity of the autocatalytic cycle in the habitat.

Discussion

We have outlined in a stepwise fashion, how increasingly elaborate autocatalytic cycles gradually

emerge in an early life RNA-habitat. Initially, catalysts with relatively lower catalytic activity are

selected, because they are parasite-specific and confer resistance (primordial autocatalytic cycles).

Soon, complexed mixtures evolve, featuring self-replicating catalysts of high catalytic activity that

become stabilized through the selection and integration of hyperparasite cycles, including relatively

less active catalyst-subunits that confer powerful parasite resistance (stabilized host-parasite-

hyperparasite cycles). One quintessential building block materializes via prebiotic heredity,

encompassing a core with variable autocatalytic cycles (genotype), and a non-catalytic periphery

(phenotype). Conjointly with the mutation-selection balance, natural selection can operate on the

genotype rendering possible complexed networks that reflect the history of parasite exposure. Under

conditions in which there is the influx of overwhelmingly broad and complexed parasite populations

into the habitat, the carrying capacity of the autocatalytic networks are exceeded, and a non-

catalytic, degenerative process of hyperparasite cycles ensues that leads to the appearance of anti-

parasitism, and the birth of immunity in early life.

One of the strong features of this model is the solid empirical evidence supporting it. First of
all, the catalyst tradeoff analyses [26] applied to autocatalytic RNA networks [25] predict the
emergence of increasingly active ribozymes, binding promiscuity and high catalytic activity being
balanced with specificity and relatively lower activity, which is precisely what the model shows.
Moreover, tripartite microbial populations composed of a host, a parasite and a hyperparasite are
virtually ubiquitous, as evidenced for bacteria-phages-phage satellites [19,20], host-RNA viruses-
defective interfering RNA viruses [21], and for eukaryotic host-NCLDVs-virophages [22,30]. We
previously developed a hyperparasite framework based on the Lotka Volterra equations that
corroborates the superior homeostatic viability of tripartite habitats [23]. It thus becomes credibly
likely, that early life networks of habitats with RNA populations were constructed following this
architectural logic. Second, prebiotic heredity has a theoretical founding [8] and experimental
support [9,10,31]. Perhaps concepts based on information theory [32] could more appropriately
capture the meaning attached to the genotypic core composed of autocatalytic cycles, and its
exchange with the world of selfish genetic elements, as we describe it here in our model. Third, it is
striking that the appearance of microbial immunity is associated with hyperparasites. Indeed, several
recent studies show that phages and their satellites encode a set of diverse antiphage systems
[19,33], and defective interfering RNAs activate innate immunity [21,28]. Phage satellite-encoded
immune systems protect bacteria from phage predation and were proposed to be an integral innate
immune component [34]. This notion could easily be extended to omnipresent, bona fide
hyperparasites, such as endogenous retroviruses [35,36]. In conclusion, our model provides an
overarching theoretical foundation for the notorious association of immunity with the parasite-
hyperparasite sphere that is rooted in early life.

Furthermore, it should be noted that extant RNA viral quasispecies most closely resemble the
primordial RNA populations described here. Large populations exhibiting fast replication recapitulate
the fundamental tradeoff, favoring speed over polymerase accuracy [27]. High mutation rates enable
the emergence and maintenance of substantial genetic variation, which leaves marks of the past
ecological history in viral genomes via selective sweeps [37]. Moreover, in all contemporary tripartite
models the host diverged to cellular life forms carrying polymerases phylogenetically distinct from
those prevalent in viruses and selfish genetic elements [17], thus severely limiting genetic exchange.
This represented a marked change from the primordial tripartite populations that freely swapped
catalysts.

This model has by design intrinsic limitations, notably the fact that the conclusions have not
yet been experimentally verified. Some components of the model may even not be amenable to
realistic experimentation, especially highly efficient self-replicating RNA catalysts [9,10]. Along the
same line, it is impossible to infer causation from statistical associations, particularly the pervasive

horizontal gene transfer associated with selfish genetic elements blurs the lines. These aspects will be
addressed in future studies.

References

1   Betts H. C., Puttick M. N., Clark J. W., Williams T. A., Donoghue P. C. J., et al. Integrated
genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nat.*
Ecol. Evol. **2**, 1556-1562 (2018).

2   Jeancolas C, Malaterre C, Nghe P. Thresholds in Origin of Life Scenarios. *iScience* **23**,
101756; 10.1016/j.isci.2020.101756 (2020).

3   Robertson M. P., Joyce G. F. The origins of the RNA world. *Cold Spring Harb. Perspect. Biol.*
4, a003608; 10.1101/cshperspect.a003608 (2012).

4   Joyce G. F, Szostak J. W. Protocells and RNA Self-Replication. *Cold Spring Harb. Perspect.*
Biol. **10**, a034801; 10.1101/cshperspect.a034801 (2018).

5   Kauffman, S. A. Cellular homeostasis, epigenesis and replication in randomly aggregated
macromolecular systems. *J. Cybernet.* **1**, 71-96 (1971).

6   Eigen, M. Self-organization of matter and the evolution of biological macromolecules.
Naturwissenschaften **58**, 465-523 (1971).

7   Kauffman S. A. Autocatalytic sets of proteins. *J. Theor. Biol.* **119**, 1-24 (1986).

8   Vaidya N., Manapat M. L., Chen I.A., Xulvi-Brunet R., Hayden E. J., et al. Spontaneous
network formation among cooperative RNA replicators. *Nature.* **491**, 72-77 (2012).

9   Tjhung K. F., Shokhirev M. N., Horning D. P., Joyce G. F. An RNA polymerase ribozyme that
synthesizes its own ancestor. *Proc. Natl. Acad. Sci. U S A.* **117**, 2906-2913 (2020).

10   Pavlinova P., Lambert C. N., Malaterre C., Nghe P. Abiogenesis through gradual evolution
of autocatalysis into template-based replication. *FEBS Lett.* **597**, 344-379 (2023).

11   Vasas V., Fernando C., Santos M., Kauffman S., Szathmáry E. Evolution before genes. *Biol.*
Direct. **7**, 10.1186/1745-6150-7-1 (2012).

12   Sczepanski J. T., Joyce G. F. Synthetic evolving systems that implement a user-specified
genetic code of arbitrary design. *Chem. Biol.* **19**, 1324-32 (2012).

13   Domingo E., García-Crespo C., Perales C. Historical Perspective on the Discovery of the
Quasispecies Concept. *Annu. Rev. Virol.* **8**, 51-72 (2021).

14   Bull J. J., Meyers L. A., Lachmann M. Quasispecies made simple. *PLoS Comput Biol.* **1**, e61;
10.1371/journal.pcbi.0010061 (2005).

15   Koonin E. V., Wolf Y. I., Katsnelson M. I. Inevitability of the emergence and persistence of
407   genetic parasites caused by evolutionary instability of parasite-free states. *Biol. Direct.* **12**,
408   31; 10.1186/s13062-017-0202-5 (2017).

409   16   Koonin E. V., Dolja V. V. Virus world as an evolutionary network of viruses and capsidless
410   selfish elements. *Microbiol. Mol. Biol. Rev.* **78**, 278-303 (2014).

411   17   Koonin E. V., Krupovic M., Ishino S., Ishino Y. The replication machinery of LUCA: common
412   origin of DNA replication and transcription. *BMC Biol.* **18**, 61; 10.1186/s12915-020-00800-
413   9 (2020).

414   18   International Committee on Taxonomy of Viruses Executive Committee. The new scope of
415   virus taxonomy: partitioning the virosphere into 15 hierarchical ranks. *Nat. Microbiol.* **5**,
416   668-674 (2020).

417   19   de Sousa J. A. M., Fillol-Salom A., Penadés J. R., Rocha E. P. C. Identification and
418   characterization of thousands of bacteriophage satellites across bacteria. *Nucleic Acids*
419   *Res.* **51**, 2759-2777 (2023).

420   20   Eppley J. M., Biller S. J., Luo E., Burger A., DeLong E. F. Marine viral particles reveal an
421   expansive repertoire of phage-parasitizing mobile elements. *Proc. Natl. Acad. Sci. U S A.*
422   **119**, e2212722119; 10.1073/pnas.2212722119 (2022).

423   21   Vignuzzi M., López C. B. Defective viral genomes are key drivers of the virus-host
424   interaction. *Nat. Microbiol.* **4**, 1075-1087 (2019).

425   22   Katzourakis A., Aswad A. The origins of giant viruses, virophages and their relatives in
426   host genomes. *BMC Biol.* **12**, 51; 10.1186/s12915-014-0051-y (2014).

427   23   Conrad B., Iseli C., Pirovino M. Energy-harnessing problem solving of primordial life:
428   Modeling the emergence of catalytic host-nested parasite life cycles. *PLoS One.* **18**,
429   e0281661; 10.1371/journal.pone.0281661 (2023).

430   24   Iranzo J., Puigbò P., Lobkovsky A. E., Wolf Y. I., Koonin E. V. Inevitability of Genetic
431   Parasites. *Genome Biol. Evol.* **8**, 2856-2869 (2016).

432   25   Janzen E., Blanco C., Peng H., Kenchel J., Chen I. A. Promiscuous Ribozymes and Their
433   Proposed Role in Prebiotic Evolution. *Chem. Rev.* **120**, 4879-4897 (2020).

434   26   Tawfik D. S. Accuracy-rate tradeoffs: how do enzymes meet demands of selectivity and
435   catalytic efficiency? *Curr. Opin. Chem. Biol.* **21**, 73-80 (2014).

436   27   Fitzsimmons W. J., Woods R. J., McCrone J. T., Woodman A., Arnold J. J., et al. A speed-
437   fidelity trade-off determines the mutation rate and virulence of an RNA virus. *PLoS Biol.*
438   **16**, e2006459; 10.1371/journal.pbio.2006459 (2018).

439   28   Genoyer E., López C. B. The Impact of Defective Viruses on Infection and Immunity. *Annu.*
440   *Rev. Virol.* **6**, 547-566 (2019).

29 Ibarra-Chávez R., Brady A., Chen J., Penadés J. R., Haag A. F. Phage-inducible
chromosomal islands promote genetic variability by blocking phage reproduction and
protecting transductants from phage lysis. *PLoS Genet.* **18**, e1010146;
10.1371/journal.pgen.1010146 (2022).
30 Meng L., Endo H., Blanc-Mathieu R., Chaffron S., Hernández-Velázquez R., et al.
Quantitative Assessment of Nucleocytoplasmic Large DNA Virus and Host Interactions
Predicted by Co-occurrence Analyses. *mSphere.* **6**, e01298-20; 10.1128/mSphere.01298-
20 (2021).
31 Yeates J. A. M., Nghe P., Lehman N. Topological and thermodynamic factors that influence
the evolution of small networks of catalytic RNA species. *RNA.* **23**, 1088-1096 (2017).
32 Koonin E. V. The meaning of biological information. *Philos. Trans. A Math. Phys. Eng. Sci.*
374, 20150065; 10.1098/rsta.2015.0065 (2016).
33 Rousset F., Depardieu F., Miele S., Dowding J., Laval A. L., et al. Phages and their satellites
encode hotspots of antiviral systems. *Cell Host Microbe.* **30**, 740-753 (2022).
34 Fillol-Salom A., Miguel-Romero L., Marina A., Chen J., Penadés J. R. Beyond the CRISPR-
Cas safeguard: PICI-encoded innate immune systems protect bacteria from bacteriophage
predation. *Curr. Opin. Microbiol.* **56**, 52-58 (2020).
35 Stauffer Y., Marguerat S., Meylan F., Ucla C., Sutkowski N., et al. Interferon-alpha-induced
endogenous superantigen. a model linking environment and autoimmunity. *Immunity.*
15, 591-601 (2001).
36 Malfavon-Borja R., Feschotte C. Fighting fire with fire: endogenous retrovirus envelopes
as restriction factors. *J. Virol.* 89, 4047-4050 (2015).
37 Zhao L., Seth-Pasricha M., Stemate D., Crespo-Bellido A., Gagnon J., et al. Existing Host
Range Mutations Constrain Further Emergence of RNA Viruses. *J. Virol.* 93, e01385-18;
10.1128/JVI.01385-18 (2019).

466
467
468 **Acknowledgements**
469 We thank Thomas Curran for discussions.
470
471
Author contributions

Conception and design of the work: MP and BC. Data acquisition MP; data analysis MP, CI;
interpretation of data MP, CI, JAC, BC. Creation of new software used in the work: MP, CI. Drafted the
work and revised it: BC, MP and JAC.

Data availability

The data, including scripts, is fully made available at the following sites (https://github.com/BICC-
UNIL-EPFL/parasite-immunity-from-hyperparasitism [github.com];
https://doi.org/10.5281/zenodo.7970788 [doi.org]).

Competing interests

The authors declare no competing interests.

515   **Figures and Figure Legends**

Fig. 1

516
517

518   **Figure 1. Habitat with one autocatalytic host cycle, variably parasite exposed**

519   Fig. 1A. The catalysts in an autocatalytic cycle (ribozyme R_2, template R_1) exposed to incoming

520   parasites P are promiscuous and hence also amplify P. Under high parasite exposure, parasites will

521   therefore outcompete host molecules.

522   Fig. 1B. Under weak parasite exposure, the host molecules R_1 and R_2 prevail.

Fig. 2

533
534

Figure 2. Habitat with primordial autocatalytic cycle

Parasite-specific catalysts with relatively lower catalytic activity in the habitat (E_2 protoreplicase) will confer parasite resistance and thus ensure abundance of habitat molecules in the presence of parasite inflow. Host molecules are expectedly less prevalent compared to Fig. 1B, at least prototemplate F_1.

Fig. 3

557
558

Figure 3. Stabilized host-parasite-hyperparasite cycle with variable parasite inflow

Fig. 3 left panel (Model 1(1+)). A habitat with highly efficient catalysts (self-replicase R_2, template R_1) becomes stabilized with moderate parasite inflow via its relatively less active catalyst-subunit E_2 that confers parasite resistance. F_2 (formerly E_2 in Fig. 2) is generated as well, it acts additively with $R2$ and will be important for antiparasitism.

Fig. 3 right panel (Model 1(2)). Under constant, moderate and thus manageable parasite inflow ($P^{special}$, Suppl. Inf.), the hyperparasite cycle mutates and becomes integrated into the habitat, even when parasite inflow ceases (prebiotic heredity and mutation-selection balance). Host molecule prevalence is comparable to a habitat without parasites (Fig. 1B).

Fig. 4A

580
581

Figure 4. Transition from hyperparasitism to antiparasitism under strong parasite inflow

Fig. 4A Under strong, permanent exposure to complex parasite populations the hyperparasite cycle partially loses its catalytic activity and becomes antiparasitic. The protoreplicase F_2 now continues to function as a partially active catalyst, but also exerts a non-catalytic antiparasitic function (antibody). The parameters in Fig. 4A are set to have these two activities equally represented.

Fig. 4B

603
604

Fig. 4B The parameters are now set to completely shift to antiparasitism (fully degenerated hyperparasite cycle, completely non-catalytic). Antiparasitism has fully replaced hyperparasitism, and the habitat is still stabilized, yet with stronger fluctuations.

Fig. 5

625
626

Figure 5. Gradual transition from hyperparasitism to antiparasitism in four steps

The stepwise transition from pure hyperparasitism (top panel) to pure antiparasitism (botton panel)

in four steps is shown (0, 0.25. 0.5, 0.75 and 1).