Biomathematical model of prebiotic autocatalytic RNA networks shows birth

of adaptive immunity from degenerating molecular parasite catalysts

Short title:

degenerating molecular parasites seed immunity

Authors:

Magnus Pirovino[1], Christian Iseli[2], Joseph A. Curran[3], Bernard Conrad[4*]

Affiliations: [1]OPIRO Consulting Ltd., FL-9495 Triesen, Principality of Liechtenstein, [2]Bioinfor­matics Competence Center, EPFL and Unil, CH-1015 Lausanne, [3]Department of Microbiology and Molecular Medicine, and Institute of Genetics and Genomics of Geneva (iGE3), Geneva University, CH-1211 Genève 4, [4]Genesupport, Avenue de Sévelin 18, CH-1004 Lausanne

*Correspondence to: bernard.conrad@genesupport.ch

Catalysis and specifically autocatalysis are the quintessential building blocks of life. However,
autocatalytic networks are necessary but not sufficient for life-like properties such as self-propagation
and adaptation to emerge. We previously showed using Lotka-Volterra equations that a tripartite,
early life cycle comprising a host-template, molecular parasites and hyperparasites (parasites of
parasites) conferred superior stability to the habitat. Here we build on this model, and on the seminal
notion that chemical networks and ecologically interacting biological species are akin, implying an
instructive equational analogy between catalysis and ecology, both accurately modelled by equations
from the Lotka-Volterra family. Template-based autocatalytic RNA networks seed the spontaneous
emergence of immunity against molecular parasites, with degenerating molecular parasite catalysts
being the founding seed of the homeostatic process. First, upon molecular parasite encounter,
specific ribocatalysts providing parasite resistance appear in primordial autocatalytic cycles,
supplanted by promiscuous, efficiently self-replicating, but parasite sensitive ribozymes. This is
supported by catalyst tradeoff analysis: substrate promiscuity confers high catalytic activity, but
entails parasite exposure, higher substrate specificity implies relatively lower activity, yet it offers
parasite resistance. Second, sustained parasite influx generates hyperparasite cycles, memorizing
parasite-specific catalyst-subunits that now once more provide parasite resistance, in a
homeostatically stabilized habitat. Third, under continuous exposure to progressively elaborate
parasite populations, hyperparasite catalysts degenerate and embody antiparasite immunity. The
observed triggering of microbial immunity by hyperparasitic microbial genomes are in agreement
with this model. As such, it offers an attractive and unifying theory for the spontaneous birth of
immunity in early life.
50
51

Author Summary

The quintessential components of life comprise a potent acceleration of naturally occurring, but
improbable chemical reactions (catalysis), and the arrangement of accelerated chemical reactions in
closed loops (autocatalytic sets). This is required, but is not sufficient for such networks to self-
propagate and adapt. We previously developed an early life model for a stabilized habitat of
networks, using a family of equations shown to accurately model both chemical reactions and the
interactions of biological species. This model became stabilized only if the molecular host species was
parasitized, an event considered unavoidable, and at the same time, the parasite itself was also
parasitized. We now built on that model, using the same family of equations, by letting individual
autocatalytic sets interact. The new results show that primitive autocatalytic networks emerge
spontaneously, but inefficiently and resist parasites. With time, they become more efficient, which

renders them parasite-sensitive. Upon continuous external parasite invasion, the primitive, parasite-

resistant catalysts (hyperparasites) that still persist are recruited, conferring stability to the habitat. If

the habitat is further overwhelmed with parasites, it selects the elegant solution of letting the

parasite-resistant catalysts degenerate, which now produces immunity to those parasites

("antibodies"). This represents the birth of adaptive immunity.

Introduction

The multistep transition process from inanimate to living systems has been dated to >3.9 Ga [1]. A

non-exhaustive list of thresholds that needed to be progressively surmounted during this process has

been drawn, comprising, among others, chirality symmetry breaking, spontaneous polymerization,

self-assembly of compartments, and particularly the catalytic closure threshold [2]. While

autocatalysis is the core ingredient of life, it is not sufficient. The ability of prebiotic networks to self-

propagate and adapt hinges upon multiple, sequential interactions of autocatalytic subsystems

among themselves and with the environment, creating higher order structures [3-7]. In one of those

models, pairs of autocatalytic cycles following chemical mass action kinetics [8] exhibit the superior

complexity-building behaviors usually attributed to biological species, for instance competitive,

predator-prey or mutualistic relationships [9]. Predator-prey type autocatalytic cycles display a

dynamics similar to the Lotka-Volterra (LV) model [9].

One central pillar in the theory of how early life could have materialized pertains to the "RNA

world". This stipulates that the initial informational polymer was RNA, and that the same molecule

was also the fundamental catalyst [10]. The underlying concept of autocatalytic sets was laid out

more than half a century ago [11,12]. It was subsequently refined [3] and validated in a proof of

principle experiment showing that RNA fragments could build self-replicating ribozymes from

individual RNA fragments via cooperative catalytic cycles [13]. In parallel, an RNA polymerase

ribozyme was generated by *in vitro* evolution that exhibited a high level of processivity [14].

Collectively, both theoretical considerations and experimental evidence support the view of a gradual

evolution towards RNA self-catalysis. This started with self-assembly, followed by the formation of

autocatalytic networks using self-assembled substrates. From this emerged different ribozymes

(recombinases, ligases) and template based ligation, leading to a template based, self-replicase

catalyzed assembly of monomers [15].

RNA self-catalysis is unthinkable without access to generic energy levels. It is commonly

argued that autocatalytic chemical networks preceded template-based autocatalysis [16]. Indeed,

proto-metabolic reactions without metals and enzymes have been demonstrated, suggesting the

existence of an ancestral analogue of the core reverse Krebs cycle [17]. By force of circumstances,
autocatalytic metabolic and template-based networks are bound to have joined forces to render the
emergence of protocells sufficiently likely [18].

The ribozyme catalyzed RNA-polymerization process is inherently error prone [10,14], generating variant sequence space mutants called a quasispecies [19,20]. This variant sequence-space intrinsically implies the obligate presence and evolutionary persistence of the molecular parasites associated with all life forms, as argued essentially for thermodynamic reasons [21,22]. We previously developed a prebiotic hyperparasite framework based on the LV-equations [23], demonstrating that superior homeostasis arose from the tripartite configuration within a given molecular habitat, namely, the molecular host, the parasite and the hyperparasite, i.e. the parasite of the parasite. The empirical evidence of highly prevalent tripartite communities in the virosphere lends substantial credit to this notion [24-27]. The LV-equations have become the default tool to quantitatively assess interactions in microbial communities [28]. However, it is often forgotten that Alfred Lotka initially modelled catalysis and autocatalysis [29]. Based on his equations modeling chemical reactions that transform a substrate [S] into a product [P] via two intermediates, X and Y, if the reactions producing X and Y are assumed to be autocatalytic, then the resulting ordinary differential equation is the classical LV-predator-prey system [30]. This stoichiometry corresponds to the Michaelis and Menten mechanism - the universal formalism for catalysis - that considers fully reversible reactions $S + E \rightleftharpoons B1 \rightleftharpoons B2 \rightleftharpoons P + E$; ($B1$ and $B2$ are the intermediates equivalent to X and Y) [31]. In fact, LV-equations commonly used in ecology, the Rosenzweig-MacArthur model with Holling type II response [32,33], are analogous to the Michaelis-Menten mechanism in enzyme kinetics [34]. Jointly, these considerations lay the ground for modelling catalysis, autocatalysis and interactions within primordial, template-based ecological RNA communities with one single biomathematical tool, namely the LV-model. This reasoning has led us to extend the initial tripartite model, building on multiple, sequentially interacting autocatalytic RNA networks memorizing the historical parasite exposure. We model, how molecular parasite resistance and antiparasite immunity can be selected for and gradually emerge from progressively degenerating hyperparasites in an intrinsic fashion in early life, in hosts exposed to increasingly more complexed parasite populations.

125
126

Results

In a stepwise and gradual process described in the Supplementary Information (proofs, Suppl. Inf.)
and Methods sections, we lay out using the LV-equations, how simple template-based RNA
autocatalytic cycles are initially built and how over time they become increasingly more elaborated as
a result of molecular parasite exposure. Metabolic autocatalytic networks are assumed to be present

and to provide sufficient energy [23]. The coexistence of prebiotic ecology/heredity [35,36] with an

emerging canonical, template-based Darwinian heredity allows for the complexification of the

quasispecies that in turn reflects its historical parasite exposure. The previously established tripartite

habitat framework featuring a template-based RNA host, a molecular parasite that is itself parasitized

(hyperparasite; [23]), is instrumental for homeostasis and the emergence of resistance, and finally

immunity to the increasingly more exhaustive molecular parasite populations.

Autocatalytic cycles in transitional early RNA-settings

The seminal duality of the RNA world, namely the fact that the same RNA molecule provides both the

template and the self-replicating catalyst [10] constitutes the foundation of the initial autocatalytic

cycles (replication equations (1.1), (1.2), forming a two-step autocatalytic cycle; proof outlined in

Suppl. Inf.). Since the template and the enzyme/ribozyme remain in principle unchanged during this

process, both can be viewed as catalysts, as long as the molecules do not undergo deleterious loss of

function mutations.

Molecular parasitism in autocatalytic cycles

Molecular parasites are an inherent feature of the replication process, they are inseparably

associated with all life forms [21,22], for a number of reasons, the most important being the

accuracy-rate tradeoff [37,38]. Average enzymes operate orders of magnitude (k_{cat}/K_M of $\sim10^5$ s^{-1}M^{-1})

below the diffusion rate (10^8-10^9 s^{-1}M^{-1}), which implies promiscuous substrate binding, high catalytic

activity, and as a secondary tradeoff a high mutation rate with a large variance in the quasispecies

(speed being more important than enzyme accuracy; [39]). In an initial autocatalytic cycle, self-

replicating ribozymes emerge (ribozyme R_2 acting on a replicating template R_1; proof outline in

Suppl. Inf.; Model 1$^{(1)}$ in Methods). The catalyst tradeoff opposes specific, but catalytically relatively

less active molecules, and promiscuous, yet catalytically highly efficient molecules [37,38]. Therefore,

the catalytically very efficient and promiscuous R_2 will inevitably be parasitized by RNA templates P.

Figure 1 illustrates the population sizes of the habitat constituents under high (Fig. 1A), and low (Fig.

1B) parasite exposure. As expected, high parasite loads overwhelm the habitat with P templates, an

observation that begs a solution.

A stable, primordial autocatalytic cycle in the presence of molecular parasites

In a primordial habitat with abundant inflow of molecular RNA species including parasites, a

quasispecies will eventually emerge with the central feature that it also comprises molecules with

superior parasite-specificity, which implies relatively lower catalytic activity, and confers parasite

resistance (protoreplicase E_2, prototemplate F_1; equations and proofs outlined in the Suppl. Inf.).

This composite feature progressively arises in successive rounds of autocatalytic cycles, initially

comprising the encounter of the catalyst E_1, provided by a neighboring habitat, with F_1 evolved

autocatalytically in the habitat, producing the primordial catalyst E_2, that is capable of amplifying F_1

but not P. It is the twofold consequence of the primordial parasite-specificity of E_2, and of the

complexity of P that exceeds the processivity of the protoreplicase. As a corollary and as displayed in

Fig. 2, selection of such molecules confers considerable resistance to parasite invasion, and restores

host template and catalyst abundance (Model $1^{(0)}$, Methods). There is both, strong theoretical and

experimental evidence to support these findings. Indeed, while it is generally assumed that catalysts

arising in early autocatalytic cycles have low activity and high promiscuity [37], an in depth physico-

chemical analysis of how enzymes meet demands of selectivity and accuracy points towards

accuracy-rate tradeoffs [38]. Selectivity (accuracy, specificity) implies the ability to discriminate

between two templates, provided both are present [37,38], and rate denotes speed [39]. Ground

state discrimination means that specificity is achieved mainly through substrate binding, which

imposes strong accuracy-rate tradeoffs [38]. Improvements in selectivity mediated by tighter cognate

substrate binding invariably lead to lower catalytic efficiency (parallel decreases in the constants for

the cognate substrates K_M^{cog} and k_{cat}^{cog}; [38]). This is precisely what the model stipulates, the

protoreplicase has a primal parasite-selectivity, which comes at the cost of a relatively reduced

catalytic activity, yet it allows for adequate host molecule-, but not parasite-amplification. In

conclusion, if catalysts with relatively reduced function in the quasispecies confer parasite resistance,

yet restore host catalyst and template abundance, they will be positively selected [35].

Evolution of the primordial autocatalytic cycle into a full-fledged autocatalytic cycle

Natural selection in the mutant quasispecies space will ultimately produce, starting from E_2, a fully

efficient, self-replicating ribozyme (replicase R_2) that is bestowed with high catalytic activity [14],

capable of generating all enzymes and templates in the habitat (equations and proofs outlined in the

Suppl. Inf.). However, this has a cost. R_2 is promiscuous, catalytically highly efficient and therefore

exposed to parasite invasion, which was not the case for its primordial catalyst precursor with

relatively lower activity (protoreplicase E_2; Model $1^{(1+)}$ stabilized host-parasite-hyperparasite cycles,

Methods).

At this point, the protoreplicase E_2-subdomain of R_2 confers P selectivity, restoring parasite

resistance and rescuing the abundance of the partial catalyst and its templates, even with a moderate

parasite inflow ($\alpha > 0$, Fig. 3 left panel). This process has as its central feature a hyperparasite cycle

(equations and proofs outlined in the Suppl. Inf.). It is further noteworthy that F_2 (formerly E_2) is still

produced (Fig. 3), and is acting additively to R_2, which is important to generate antiparasitism later.

Under permanent and moderate, thus manageable parasite inflow ($P^{special}$, Suppl. Inf.), an
increasingly ramified network of autocatalytic cycles arrives in the habitat that persists even when
the parasite exposure ceases, constituting a historical record of past parasite exposures ($\alpha = 0$, Fig. 3
right panel).

There is considerable evidence in favor of the essential components of this model, in
particular the catalyst accuracy-rate tradeoff and the hyperparasite cycle. Firstly, a solid body of
theoretical and experimental evidence supports catalyst accuracy-rate tradeoffs [37,38]. Secondly,
metagenomic studies demonstrate the virtually universal presence of tripartite constituents across
molecular parasite habitat classes, most persuasively for bacteria-phages-phage satellites [24,25], for
RNA viruses partnering the host- virus-defective interfering RNAs [26], and for NCLDVs, with
eukaryotic host-NCLDV-virophage [27]. Thirdly, additional support comes from the observation that
hyperparasites are induced across the range of parasite concentrations, and especially at high
multiplicity infection for RNA viruses [40,41].

Emergence of antiparasitism in autocatalytic cycles

Under conditions, where the sheer number and complexity of incoming molecular parasite
populations would overwhelm the carrying capacity of the habitat, a degenerative process is initiated
that erodes catalysis. Formally, the hyperparasite cycle degenerates to an antiparasite cycle
(equations and proofs outlined in the Suppl. Inf.; Model $1^{(1-)}$, Methods). In Figures 4 and 5, the
transition from hyperparasitism to antiparasitism is illustrated, in the sense that both coexist at equal
activity, partial catalyst and non-catalytic antiparasitic activities (Fig. 4A, see Methods for
parameters). In Figure 4B, antiparasitism is fully implemented; hyperparasitism has been fully
replaced by the non-catalytic antiparasitism (reproduction parameters hyperparasite d and δ_3 are set
to zero: $\delta_3 = d = 0$), making the difference between pure antiparasitism, and the transition from
hyperparasitism to antiparasitism minor. In Figure 4B, the population fluctuations are more volatile
than in Figure 4A, but pure antiparasitism does effectively stabilize the autocatalytic cycle, without a
hyperparasite component. Figure 5 shows the gradual transition from pure hyperparasitism (top
panel) to antiparastism (botton panel), in four successive steps (0, 0.25, 05. 0.75 and 1), progressively
replacing hyperparasitism with antiparasitism. Therefore, we conclude that antiparasitism can replace
hyperparasitism in situations with a stronger and highly diverse parasite inflow that would overrun
the carrying capacity of the autocatalytic cycle in the habitat.

Discussion

We have outlined in a stepwise fashion, how increasingly elaborate autocatalytic cycles gradually emerge in an early life RNA-habitat. Initially, catalysts with low catalytic activity are selected, because they are parasite-specific and confer resistance (primordial autocatalytic cycles). Soon, complexed mixtures evolve, featuring self-replicating catalysts of high catalytic activity that become stabilized through the selection and integration of hyperparasite cycles, including relatively less active catalyst-subunits that confer parasite resistance (stabilized host-parasite-hyperparasite cycles). Under conditions in which there is the influx of broad and complexed molecular parasite populations into the habitat, the carrying capacity of the autocatalytic networks are exceeded, and a non-catalytic, degenerative process of hyperparasite cycles ensues that leads to the appearance of antiparasitism, and the birth of immunity in early life.

One of the strong features of this model is the solid empirical evidence supporting it. First of all, the catalyst tradeoff analyses [38] applied to autocatalytic RNA networks predicts the emergence of increasingly active ribozymes, with binding promiscuity and high catalytic activity being balanced with specificity and relatively lower activity, which is precisely what the model shows. Prebiotic models indeed support initially fast replicating communities with high molecular diversity, evolving later into more faithful replicators with a narrower repertoire [42]. Moreover, tripartite microbial populations composed of a host, a parasite and a hyperparasite are virtually ubiquitous in the virosphere, as evidenced for bacteria-phages-phage satellites [24,25], host-RNA viruses-defective interfering RNA viruses [26], and eukaryotic host-NCLDVs-virophages [27]. We previously developed a hyperparasite framework based on the LV-equations that corroborates the superior homeostatic viability of tripartite habitats [23]. It thus becomes likely, that early life network habitats with RNA populations were constructed following this architectural logic.

Second, some sort of trans-hierarchical self-organization has to be stipulated at this stage of the early RNA world [43], ideas conceptualized under various designations, such as prebiotic heredity or ecology, and autopoiesis [35,36]. The early RNA world for instance implies lipid-vesicle mediated compartmentalization and supramolecular selection [35]. Applied to our findings, ribozymes conferring parasite resistance are selected early on, survive autopoietically within the quasispecies, and can thus be reselected at later stages, once more efficient catalysts expose the community to waves of threatening parasite invasions.

Third, it is striking that the appearance of microbial immunity is associated with hyperparasites. Indeed, several recent studies show that phages and their satellites encode a set of diverse antiphage systems [24,44], and defective interfering RNAs activate innate immunity [26,40]. Phage satellite-encoded immune systems protect bacteria from phage predation and were proposed to be an integral innate immune component [45]. This notion could easily be extended to omnipresent, bona fide hyperparasites, such as endogenous retroviruses [46,47]. In conclusion, our

model provides an overarching theoretical foundation for the notorious association of immunity with

the parasite-hyperparasite sphere that is rooted in early life.

Furthermore, it should be noted that extant RNA viral quasispecies most closely resemble the

primordial RNA populations described here. Large populations exhibiting fast replication recapitulate

the fundamental tradeoff, favoring speed over polymerase accuracy [39]. High mutation rates enable

the emergence and maintenance of substantial genetic variation, which leaves marks of the past

ecological history within viral genomes via selective sweeps [48]. Moreover, in all contemporary

tripartite models the host diverged to cellular life forms carrying polymerases phylogenetically

distinct from those prevalent in viruses and selfish genetic elements [49], thus severely limiting

genetic exchange. This represented a marked change from the primordial tripartite populations that

freely swapped catalysts. The fact that the integration of the virosphere into early RNA world

scenarios has been proposed as an important future agenda issue further valorizes our model [43].

Conceptually, our model capitalizes upon one recent and new school of thought in

autocatalytic networks, namely the fact that chemical networks behave as ecological species

interacting with one another [9]. Since the common underlying physical principle in catalysis and

ecological interactions is the mass action law [8], it is not surprising that the biomathematical tools

most accurately capturing the quantitative nature of these interactions are also equivalent, namely

equations from the LV-family. Probably one of the most significant merits of our model founded in

the equivalence of autocatalysis and ecological interactions is the fact that catalytic processes by their

very nature can degenerate and become non-catalytic, and that a degenerative process has

innovative power. We can summarize the sequence of events as, i) the catalytic process that is

inherently error-prone and generates a quasispecies comprising molecular parasites; ii) sequential

ecological interactions between host and parasite catalysts stabilize the habitat, forming tripartite

ecological communities of host-parasites-hyperparasites; iii) a subgroup of hyperparasite catalysts in

such habitats, when overwhelmed with invading parasite waves, can degenerate and create adaptive

immunity against those parasites. Thus, in essence, degenerating catalysts create a new, homeostatic

function that allows the habitat to adapt.

This model has by design intrinsic limitations, notably the fact that certain conclusions have

yet to be experimentally verified [13,14]. Along the same line, it is impossible to infer causation from

statistical associations, in particular the pervasive horizontal gene transfer associated with selfish

genetic elements blurs the lines. These aspects will be addressed in future studies.

Methods or Models

Definitions and Terms

In what follows we assume the following: the Malthusian fitness r of a replicating (biochemical)

system (or species) is defined by $\frac{d}{dt} R = \dot{R} = r\,R$, where R denotes the population size of the

replication sys-tem or species. Evolutionary dynamics between two or more interacting species or

systems, e.g. $R, S, P, \ldots$ are described by the following type of replicator differential equation e.g. for

R:

$\dot{R} =$ "(A) terms of dynamics using habitat resources" x "(B) term of habitat resource restriction" + "(C)

terms of dynamics not using habitat resources"

Terms A include both "terms of first order kinetics" and "terms of second order kinetics"

— "Terms of first order kinetics" are typically natural growth or decay rates, e.g. $\dot{R} = r\,R$. If $r >$

0, growth process described by rR eats into resources, whereas if $r < 0$, r is a decay rate, and

de-cay dynamics does not.

Terms of second order kinetics are derived from encounters between different species. For

instance, the term "bRS" in "$\dot{R} = -aR + bRS$" has the following meaning:

$bRS =$

$(probabiliy\ of\ an\ encounter\ between\ members\ of\ R\ and\ S)$ x $(impact\ of\ encounter\ on\ the\ fitness\ of\ R)$.

Here we assume, according to the Lotka Volterra calculus that the probability of an encounter

between members of R and S is proportional to the product RS.

The term B of resource restriction is used in a habitat of limited resources. For instance, in the

classic logistic equation $\dot{R} = r\,R\left(1 - \frac{R}{K}\right)$ the term $\left(1 - \frac{R}{K}\right) = B$ indicates that the habitat has

limited resources supporting a maximum population K for R.

Since every encounter involves resource consuming, we assume here, that all terms of second

order kinetics are subject to resource restrictions and therefore need to be multiplied with the

term B.

Example of a host-parasite interaction (R=host, P=parasite):

$$\dot{R} = (a\,R - bPR)\left(1 - \frac{R + P}{K}\right)$$

$$\dot{P} = -c\,P + dRP\left(1 - \frac{R + P}{K}\right)$$

where $a > 0, b > 0, c > 0, d > 0$. The first order dynamics growth process aR and all the second

order processes eat into the habitat's resources, whereas the first order decay process $-c\,P$ does

not. The habitat resources restriction is represented by the term $\left(1 - \frac{R+P}{K}\right)$, indicating that a

maximum population size of $K = R + P$ can be nourished by the habitat and that both, S and P,

compete for the same resources.

Definition of catalysts, ribozymes and enzymes

We use the term *catalyst*, sometimes also *perfect catalyst*, for any substance that accelerates the

rate of a reaction without undergoing any change during the process. We call an *imperfect*

catalyst a substance that accelerates the rate of a reaction, but that may undergo a change

during the process; antibodies in the transition from hyperparasitism to antiparasitism are

examples of imperfect catalysts (see process $(3')$, Suppl. Inf.). Catalysts displaying relatively

higher substrate specificity generally have relatively lower catalytic activity, for instance the

protoreplicase. A subgroup of catalysts is called here *enzymes*, if they are generated based on

RNA templates. An enzyme may be either a *ribozyme*, if it is also used as an RNA template to

generate other enzymes, or a *pure enzyme* if it cannot be used as an RNA template for other

polymerization processes. The term *ribozyme* encompasses different enzymatic activities, such as

recombinase, ligase and polymerase. Recombinases and ligases predominate in early

protocycles, while polymerases appear only in advanced networks.

Model 1$^{(1)}$: habitat with one host autocatalytic cycle exposed to parasite invasion

$$\frac{dR_2}{dt} = a\, R_1 R_1 \Pi - d_{R_2} R_2$$

$$\frac{dR_1}{dt} = (b\, R_2 R_1 - \delta R_2 P)\Pi - d_{R_1} R_1$$

$$\frac{dP}{dt} = c R_2 P \Pi + \alpha - d_P P,$$

where a, b and c are positive constants, d_{R_i}, d_P are the decay rates of populations $R_i, i = 1, 2$, and

of parasite P absent its host R_2; δ is parasite's P rate of exploitation of the host replicase R_2; and the

habitat restriction term is defined as $\Pi = \left(1 - \frac{R_1 + R_2 + P/q_P}{K}\right)$, where $q_P > 1$, since typically parasites

P are more efficient using the resources of the habitat than their hosts R. α is the parasite invasion

parameter, i.e., per unit of time the habitat is invaded by α population units of parasites P.

Model 1$^{(0)}$ habitat with primordial autocatalytic cycle, exposed to template invasion P (to become

parasite later in evolution) and protoenzyme E_1:

$$\frac{dP}{dt} = \alpha - d_P P$$

$$\frac{dE_1}{dt} = \beta - d_{E_1} E_1$$

$$\frac{dE_2}{dt} = a E_1 F_1 \Pi - d_{E_2} E_2$$

$$\frac{dF_1}{dt} = b E_2 P \Pi - d_{F_1} F_1,$$

where

E_2 is the protoreplicase produced by this primitive autocatalytic cycle,

F_1 is the prototemplate produced by this primitive autocatalytic cycle,

α is the inflow rate of RNA-template P into the habitat,

β is the inflow rate of protoenzyme/catalyst E_1 into the habitat,

a and b are positive constants,

d_{E_i}, d_{F_1}, d_P are the decay rates of enzyme populations E_i, $i = 1, 2$, of prototemplate d_{F_1}, and of

RNA-template P,

$\Pi = \left(1 - \dfrac{E_2 + {}^{F_1}/q_{F_1}}{K}\right)$ is the habitat restriction term where $q_{F_1} > 1$, since we assume that proto-

templates F_1 are more efficient using the resources of the habitat than the more complex

protoreplicase E_2.

Model $1^{(1+)}$: *stabilized host-parasite-hyperparasite cycle*

$$\frac{dR_2}{dt} = (a\,R_1 R_1 - \delta_2 R_1 F_1)\Pi - d_{R_2} R_2$$

$$\frac{dR_1}{dt} = (b\,R_2 R_1 - \delta_1 R_2 P)\Pi - d_{R_1} R_1$$

$$\frac{dP}{dt} = (c R_2 P - \delta_3 F_2 P)\Pi + \alpha - d_P P$$

$$\frac{dF_1}{dt} = (d\,F_2 P + \beta R_2 P)\Pi - d_{F_1} F_1$$

$$\frac{dF_2}{dt} = (e\,R_1 F_1)\Pi - d_{F_2} F_2,$$

where

a, b, c, d, e are positive (reproduction) constants,

R_1 is a ribozyme of the autocatalytic host cycle,

R_2 is the replicase, i.e. another ribozyme (or may be also pure enzyme since it is not used as

template) of the autocatalytic host cycle,

P is the parasite, exploiting replicase R_2 of the host cycle (exploitation parameter δ_1),

F_1 is the prototemplate exploiting the host R_1 (exploitation parameter δ_2) for protoreplicase F_2

production, triggered upon encounter of R_2 with P (trigger parameter β) and further produced by

protoreplicase F_2 upon encounter with P,

F_2 is the protoreplicase (pure enzyme or ribozyme) producing F_1 upon encounter with parasite temp-

lates P ($F_2 = E_2$ from the primitive autocatalytic cycle of Model $1^{(0)}$), thus, exploiting template of

parasite P (exploitation parameter δ_3),

d_{R_i}, d_{F_i}, d_P are the decay rates of populations $R_i, F_i, i = 1, 2$, and of P, absent new inflow or

reproduction in the habitat,

α is the inflow rate of RNA template P into the habitat,

β is trigger parameter of replicase R_2 producing F_1 by encounter with parasite P,

$\Pi = \left(1 - \frac{R_1 + R_2 + P/q_P + F_1/q_{F_1} + F_2/q_{F_2}}{K}\right)$ is the habitat restriction term where $q_P, q_{F_1}, q_{F_2} > 1$, since we

assume that parasitizing cycles of P and F_1, F_2 are more efficient using the resources of the habitat

than the more complex host cycle R_1, R_2.

Model $1^{(2)}$: integrated host-parasite-hyperparasite cycle without new parasite inflow

The simulation results of Model $1^{(2)}$ are realized as follows: Model $1^{(1+)}$ provides the starting basis,

with arbitrary initial populations to establish a stabilized situation, given a moderate parasite inflow

with $\alpha > 0$ (Fig. 3 lefthand part). Subsequently, Model $1^{(1+)}$ is adapted to Model $1^{(2)}$. All the

parameters of Model $1^{(2)}$ are the same as those of Model $1^{(1+)}$, with the exception of the parasite

inflow, now $\alpha = 0$. As initial conditions for the populations of Model $1^{(2)}$, we assume the terminal

populations of the simulations of Model $1^{(2)}$, as calculated for Fig. 3 at the beginning of the righthand

part. The larger fluctuations of the population at the beginning of the righthand part of Fig. 3 stem

from the transition from Model $1^{(1+)}$ to Model $1^{(2)}$, where the parasite inflow abruptly decreases

from $\alpha > 0$ to $\alpha = 0$. The fact that parasites in Model $1^{(2)}$ are now reproduced without new inflow

shows that they can be viewed as fully integrated into the autocatalytic network, realized in $1^{(2)}$ (see

Suppl. Inf.).

Model $1^{(1-)}$: *transition from hyperparasitism to antiparasitism*

$$\frac{dR_2}{dt} = (a\,R_1 R_1 - \delta_2 R_1 F_1)\Pi - d_{R_2} R_2$$

$$\frac{dR_1}{dt} = (b\,R_2 R_1 - \delta_1 R_2 P)\Pi - d_{R_1} R_1$$

$$\frac{dP}{dt} = (cR_2 P - \delta_3 F_2 P)\Pi + \alpha - \theta_P F_2 P - d_P P$$

$$\frac{dF_1}{dt} = (d\,F_2 P + \beta R_2 P)\Pi - d_{F_1} F_1$$

$$\frac{dF_2}{dt} = (e\,R_1 F_1)\Pi - \theta_{F_2} F_2 P - d_{F_2} F_2,$$

Where all the parameters are the same as in Model $1^{(1+)}$, with the exception that now two additional

parameters are added, namely θ_P and θ_{F_2}. The term $\theta_P F_2 P$ denotes the number of population units

of parasites P made unviable per unit of time during the encounter of protoreplicase F_2 with P, now

also functioning as antibody. In addition, the term $\theta_{F_2} F_2 P$ denotes the number of population units of

protoreplicase/antibody F_2 made unviable per unit of time during the encounter of

protoreplicase/antibody F_2 with P. For instance, if $\theta_p = \theta_{F_2}$, then the same number of antibodies and
parasites are made unviable during their encounter. In the evolutionary transition from
hyperparasitism to antiparasitism F_2 can still act both ways, as imperfect protoreplicase and as
antibody directed against the parasite P. We call it *imperfect* protoreplicase or imperfect catalyst,
because it still produces/catalyzes prototemplate F_1 upon encounter with template P. But now F_2
may be damaged during this process, therefore cannot be called a *perfect* catalyst anymore. The
replication differential equations of Model $1^{(1-)}$ realizing a pure antiparasitism situation are the
same as in Model $1^{(1-)}$ of Fig. 4A, except that the reproduction parameters hyperparasite d and δ_3
are now set to zero: $\delta_3 = d = 0$.

418
419

Acknowledgements

421 We thank Thomas Curran for discussions.

422
423

References

1. Betts HC, Puttick MN, Clark JW, Williams TA, Donoghue PCJ, Pisani D. Integrated genomic and
fossil evidence illuminates life's early evolution and eukaryote origin. Nat Ecol Evol 2018; 2:
1556-1562. doi: 10.1038/s41559-018-0644-x. Epub 2018 Aug 20. PMID: 30127539
2. Jeancolas C, Malaterre C, Nghe P. Thresholds in Origin of Life Scenarios. iScience 2020; 23:
101756. doi: 10.1016/j.isci.2020.101756. PMID: 33241201
3. Kauffman SA. Autocatalytic sets of proteins. J Theor Biol 1986; 119: 1-24. doi: 10.1016/s0022-
5193(86)80047-9. PMID: 3713221
4. Hordijk W, Hein J, Steel M. Autocatalytic Sets and the Origin of Life. Entropy. 2010; 12: 1733-
1742. https://doi.org/10.3390/e12071733
5. Shapiro R. Small molecule interactions were central to the origin of life. Q Rev Biol. 2006; 81:
105-25. doi: 10.1086/506024. PMID: 16776061
6. Ruiz-Mirazo K, Briones C, de la Escosura A. Prebiotic systems chemistry: new perspectives for the
origins of life. Chem Rev 2014: 114: 285-366. doi: 10.1021/cr2004844. Epub 2013 Oct 31. PMID:
24171674
7. Peng Z, Linderoth J, Baum DA. The hierarchical organization of autocatalytic reaction networks
and its relevance to the origin of life. PLoS Comput Biol 2022; 18: e1010498. doi:
10.1371/journal.pcbi.1010498. PMID: 36084149

8.   Voit EO, Martens HA, Omholt SW. 150 years of the mass action law. PLoS Comput Biol 2015;

11:e1004012. doi: 10.1371/journal.pcbi.1004012. PMID: 25569257

9.   Peng Z, Plum AM, Gagrani P, Baum DA. An ecological framework for the analysis of prebiotic

chemical reaction networks. J Theor Biol 2020; 507: 110451. doi: 10.1016/j.jtbi.2020.110451.

Epub 2020 Aug 12. PMID: 32800733

10. Joyce GF, Szostak JW. Protocells and RNA Self-Replication. Cold Spring Harb Perspect Biol 2018;

10: a034801. doi: 10.1101/cshperspect.a034801. PMID: 30181195

11. Kauffman SA. Cellular homeostasis, epigenesis and replication in randomly aggregated

macromolecular systems. J Cybernet 1971; 1: 71-96.

12. Eigen M. Self-organization of matter and the evolution of biological macromolecules.

Naturwissenschaften 1971; 58: 465-523.

13. Vaidya N, Manapat ML, Chen IA, Xulvi-Brunet R, Hayden EJ, Lehman N. Spontaneous network

formation among cooperative RNA replicators. Nature 2012; 491: 72-7. doi:

10.1038/nature11549. Epub 2012 Oct 17. PMID: 23075853

14. Tjhung KF, Shokhirev MN, Horning DP, Joyce GF. An RNA polymerase ribozyme that synthesizes

its own ancestor. Proc Natl Acad Sci U S A 2020; 117: 2906-2913. doi: 10.1073/pnas.1914282117.

Epub 2020 Jan 27. PMID: 31988127

15. Pavlinova P, Lambert CN, Malaterre C, Nghe P. Abiogenesis through gradual evolution of

autocatalysis into template-based replication. FEBS Lett 2023; 597: 344-379. doi: 10.1002/1873-

3468.14507. Epub 2022 Nov 17. PMID: 36203246

16. Xavier JC, Hordijk W, Kauffman S, Steel M, Martin WF. Autocatalytic chemical networks at the

origin of metabolism. Proc Biol Sci 2020; 287: 20192377. doi: 10.1098/rspb.2019.2377. Epub

2020 Mar 11. PMID: 32156207

17. Stubbs RT, Yadav M, Krishnamurthy R, Springsteen G. A plausible metal-free ancestral analogue

of the Krebs cycle composed entirely of α-ketoacids. Nat Chem 2020; 12: 1016-1022. doi:

10.1038/s41557-020-00560-7. Epub 2020 Oct 12. PMID: 33046840

18. Takeuchi N, Hogeweg P, Koonin EV. On the origin of DNA genomes: evolution of the division of

labor between template and catalyst in model replicator systems. PLoS Comput Biol 2011;

7:e1002024. doi: 10.1371/journal.pcbi.1002024. Epub 2011 Mar 24. PMID: 21455287

19. Domingo E, García-Crespo C, Perales C. Historical Perspective on the Discovery of the

Quasispecies Concept. Annu Rev Virol 2021; 8: 51-72. doi: 10.1146/annurev-virology-091919-

105900. PMID: 34586874

20. Bull JJ, Meyers LA, Lachmann M. Quasispecies made simple. PLoS Comput Biol 2005; 1: e61. doi:

10.1371/journal.pcbi.0010061. PMID: 16322763

21.   Koonin EV, Wolf YI, Katsnelson MI. Inevitability of the emergence and persistence of genetic

parasites caused by evolutionary instability of parasite-free states. Biol Direct 2017; 12: 31. doi:

10.1186/s13062-017-0202-5. PMID: 29202832

22.   Iranzo J, Puigbò P, Lobkovsky AE, Wolf YI, Koonin EV. Inevitability of Genetic Parasites. Genome

Biol Evol 2016; 8: 2856-2869. doi: 10.1093/gbe/evw193. PMID

23.   Conrad B, Iseli C, Pirovino M. Energy-harnessing problem solving of primordial life: Modeling the

emergence of catalytic host-nested parasite life cycles. PLoS One 2023; 18: e0281661. doi:

10.1371/journal.pone.0281661. PMID: 36972235

24.   de Sousa JAM, Fillol-Salom A, Penadés JR, Rocha EPC. Identification and characterization of

thousands of bacteriophage satellites across bacteria. Nucleic Acids Res 2023; 51: 2759-2777.

doi: 10.1093/nar/gkad123. PMID: 36869669

25.   Eppley JM, Biller SJ, Luo E, Burger A, DeLong EF. Marine viral particles reveal an expansive

repertoire of phage-parasitizing mobile elements. Proc Natl Acad Sci U S A 2022; 119:

e2212722119. doi: 10.1073/pnas.2212722119. Epub 2022 Oct 18. PMID: 36256808

26.   Vignuzzi M, López CB. Defective viral genomes are key drivers of the virus-host interaction. Nat

Microbiol 2019; 4: 1075-1087. doi: 10.1038/s41564-019-0465-y. Epub 2019 Jun 3. PMID:

31160826

27.   Katzourakis A, Aswad A. The origins of giant viruses, virophages and their relatives in host

genomes. BMC Biol 2014; 12: 51. doi: 10.1186/s12915-014-0051-y. PMID: 25184667

28.   Davis JD, Olivença DV, Brown SP and Voit EO. Methods of quantifying interactions among

populations using Lotka-Volterra models. Front Syst Biol 2022; 2: 1021897. doi:

10.3389/fsysb.2022.1021897

29.   Lotka AJ. Contribution to the Theory of Periodic Reactions. J Phys Chem 1910; 14: 271–274.

30.   Boros B, Hofbauer J, Müller S, Regensburger G. The Center Problem for the Lotka Reactions with

Generalized Mass-Action Kinetics. Qual Theory Dyn Syst 2018; 17: 403-410. doi:

10.1007/s12346-017-0243-2. Epub 2017 Jun 13. PMID: 30636938

31.   Michaelis L, Menten ML, Johnson KA, Goody RS. The original Michaelis constant: translation of

the 1913 Michaelis-Menten paper. Biochemistry 2011; 50: 8264-9. doi: 10.1021/bi201284u.

Epub 2011 Sep 9. PMID: 21888353

32.   Kooi BW, Poggiale JC. Modelling, singular perturbation and bifurcation analyses of bitrophic food

chains. Math Biosci 2018; 301: 93-110. doi: 10.1016/j.mbs.2018.04.006. Epub 2018 Apr 22.

PMID: 29684407

33.   Poggiale JC, Aldebert C, Girardot B, Kooi BW. Analysis of a predator-prey model with specific time

scales: a geometrical approach proving the occurrence of canard solutions. J Math Biol 2020; 80:

39-60. doi: 10.1007/s00285-019-01337-4. Epub 2019 Feb 20. PMID: 30788562

34. Eilertsen J, Roussel MR, Schnell S, Walcher S. On the quasi-steady-state approximation in an
open Michaelis-Menten reaction mechanism. AIMS Math 2021; 6: 6781-6814. doi:
10.3934/math.2021398. Epub 2021 Apr 21. PMID: 34142000

35. Wieczorek R. On prebiotic ecology, supramolecular selection and autopoiesis. Orig Life Evol
Biosph 2012; 42:445-50; discussion 451-2. doi: 10.1007/s11084-012-9306-1. PMID: 23132761

36. Igamberdiev AU, Kleczkowski LA. Toward understanding the emergence of life: A dual function of
the system of nucleotides in the metabolically closed autopoietic organization. Biosystems 2023;
224: 104837. doi: 10.1016/j.biosystems.2023.104837. Epub 2023 Jan 14. PMID: 36649884

37. Janzen E, Blanco C, Peng H, Kenchel J, Chen IA. Promiscuous Ribozymes and Their Proposed Role
in Prebiotic Evolution. Chem Rev 2020; 120: 4879-4897. doi: 10.1021/acs.chemrev.9b00620.
Epub 2020 Feb 3. PMID: 32011135

38. Tawfik DS. Accuracy-rate tradeoffs: how do enzymes meet demands of selectivity and catalytic
efficiency? Curr Opin Chem Biol 2014; 21:73-80. doi: 10.1016/j.cbpa.2014.05.008. Epub 2014 Jun
20. PMID: 24954689

39. Fitzsimmons WJ, Woods RJ, McCrone JT, Woodman A, Arnold JJ, Yennawar M, Evans R, Cameron
CE, Lauring AS. A speed-fidelity trade-off determines the mutation rate and virulence of an RNA
virus. PLoS Biol 2018; 16: e2006459. doi: 10.1371/journal.pbio.2006459. PMID: 29953453

40. Genoyer E, López CB. The Impact of Defective Viruses on Infection and Immunity. Annu Rev Virol.
2019; 6: 547-566. doi: 10.1146/annurev-virology-092818-015652. Epub 2019 May 13. PMID:
31082310).

41. Ibarra-Chávez R, Brady A, Chen J, Penadés JR, Haag AF. Phage-inducible chromosomal islands
promote genetic variability by blocking phage reproduction and protecting transductants from
phage lysis. PLoS Genet 2022; 18: e1010146. doi: 10.1371/journal.pgen.1010146. PMID:
35344558

42. Markovitch O, Lancet D. Multispecies population dynamics of prebiotic compositional
assemblies. J Theor Biol 2014; 357: 26-34. doi: 10.1016/j.jtbi.2014.05.005. Epub 2014 May 14.
PMID: 24831416

43. Fine JL, Pearlman RE. On the origin of life: an RNA-focused synthesis and narrative. RNA 2023;
29: 1085-1098. doi: 10.1261/rna.079598.123. Epub 2023 May 4. PMID: 37142437

44. Rousset F, Depardieu F, Miele S, Dowding J, Laval AL, Lieberman E, Garry D, Rocha EPC, Bernheim
A, Bikard D. Phages and their satellites encode hotspots of antiviral systems. Cell Host Microbe
2022; 30: 740-753.e5. doi: 10.1016/j.chom.2022.02.018. Epub 2022 Mar 21. PMID: 35316646

45. Fillol-Salom A, Miguel-Romero L, Marina A, Chen J, Penadés JR. Beyond the CRISPR-Cas
safeguard: PICI-encoded innate immune systems protect bacteria from bacteriophage predation.

Curr Opin Microbiol 2020; 56: 52-58. doi: 10.1016/j.mib.2020.06.002. Epub 2020 Jul 9. PMID:
32653777
46. Stauffer Y, Marguerat S, Meylan F, Ucla C, Sutkowski N, Huber B, Pelet T, Conrad B. Interferon-
alpha-induced endogenous superantigen. a model linking environment and autoimmunity.
Immunity 2001; 15: 591-601. doi: 10.1016/s1074-7613(01)00212-6. PMID: 11672541
47. Malfavon-Borja R, Feschotte C. Fighting fire with fire: endogenous retrovirus envelopes as
restriction factors. J Virol 2015; 89: 4047-50. doi: 10.1128/JVI.03653-14. Epub 2015 Feb 4. PMID:
25653437
48. Zhao L, Seth-Pasricha M, Stemate D, Crespo-Bellido A, Gagnon J, Draghi J, Duffy S. Existing Host
Range Mutations Constrain Further Emergence of RNA Viruses. J Virol 2019; 93: e01385-18. doi:
10.1128/JVI.01385-18. PMID: 30463962
49. Koonin EV, Dolja VV. Virus world as an evolutionary network of viruses and capsidless selfish
elements. Microbiol Mol Biol Rev 2014; 78: 278-303. doi: 10.1128/MMBR.00049-13. PMID:
24847023

559
560

561 **Supporting information captions**

562 *Self-replicating RNA molecules*

563 Let us consider a set of populations $\{R_1, R_2, \ldots, R_n\}$ of different RNA molecules or enzymes in a

564 primordial RNA setting. Enzymatic functions are exerted by RNA molecules called ribozymes.

565 Ribozymes and other enzymes are forming a catalytic cycle, if

$$R_1 \to^c R_2 \to^c R_3 \to^c \ldots \to^c R_n \to^c R_1,$$

where the symbol "$A \to^c B$" means: A catalyzes B.

In this setting, RNA A can display two different functions, the coding and the enzymatic function.

RNA-molecules without enzymatic activity have at least a coding potential. Let us call the coding

function "template" of RNA molecule. A pure enzyme not bearing a coding function lacks the ability

to serve as template. Let us further assume that in this instance successful enzymatic activities

require both enzyme and template. An enzymatic activity is thus exerted by a ribozyme or a pure

enzyme upon encounter with an RNA template. For example, consider the following equations:

$$\frac{dR_2}{dt} = a\, R_1 R_1 \Pi - d_{R_2} R_2 \quad (1.1)$$

$$\frac{dR_1}{dt} = b\, R_2 R_1 \Pi - d_{R_1} R_1 \quad (1.2),$$

573 where a and b are positive constants, d_{R_i} are the decay rates of populations R_i, $i = 1, 2$, and

$$\Pi = \left(1 - \frac{R_1 + R_2}{K}\right)$$

the habitat restriction constant. K is the maximum number of R_1 or R_2 molecules, the habitat can
feed.

Replication equations (1.1), (1.2) form a two-step autocatalytic cycle. In (1.1), R_1 serves both as en-
zyme and template, in (1.2) R_2 serves as enzyme and R_1 as template, i.e. upon encounter of two
different molecules R_1, one serving as template and the other as enzyme, a ribozyme or pure enzyme
R_2 is produced/catalyzed, and upon encounter of a molecule R_1 serving as template with a molecule
R_2 serving as enzyme the original ribozyme R_1 is produced, forming a complete replication cycle. We
symbolize this autocatalytic process by:

$$R_1 \Rightarrow (R_1 \leftrightarrow R_1) \to^c R_2 \Rightarrow (R_2 \leftrightarrow R_1) \to^c R_1 \quad (1),$$

582 where the symbol "$\Rightarrow$" means *"enables"* and the symbol "$A \leftrightarrow B$" means *"encounter of A with B"*
583 and, as already stated above the symbol "$\to^c$" means *"catalyzes"*.

584 Thus, autocatalytic cycles can take the form:

$$R_1 \Rightarrow (R_1 \leftrightarrow R_1) \to^c R_2 \Rightarrow (R_2 \leftrightarrow R_1) \to^c R_3 \Rightarrow (R_3 \leftrightarrow R_1) \to^c \ldots \to^c R_n \Rightarrow (R_n \leftrightarrow R_1) \to^c R_1.$$

585 In this example, only R_1 is used as a template during every enzymatic subactivity of the autocatalytic
586 reproduction cycle; every enzymatic activity produces a new ribozyme or a new pure enzyme
587 $R_i, i = 1, \ldots n$.

588 Rather than producing a further ribozyme or a pure enzyme, an enzymatic activity can also produce a
589 further template, e.g.

$$R_1 \Rightarrow (R_1 \leftrightarrow R_1) \to^c R_2 \Rightarrow (R_2 \leftrightarrow R_1) \to^c R_3 \Rightarrow (R_2 \leftrightarrow R_3) \to^c R_4 \Rightarrow (R_4 \leftrightarrow R_1) \to^c R_1$$

reads as follows: a ribozyme R_1 enables the encounter with another copy of itself, one serving as
enzyme, the other as template, and both producing a new enzyme R_2. Upon encounter with a
template R_1, R_2 catalyzes a new template R_3, and upon encounter with this new template a new
enzyme R_4, which in turn catalyzes the final reproduction of a new ribozyme molecule R_1 using
another molecule R_1 as template.

Parasitism in autocatalytic cycles

To outline the problem of parasitism, e.g. in the autocatalytic cycle (1), the ribozyme R_2, which may
also be a pure enzyme, is acting as a replicase for R_1. Such a replicase is promiscuous and highly
active, therefore, R_2 will inevitably be parasitized by parasitic RNA-templates P:

$$\frac{dP}{dt} = cR_2 P\Pi - d_P P \quad (2.1),$$

600 or, written in above introduced technology template format:

601 $$P \Rightarrow (R_2 \leftrightarrow P) \to^c P \quad (2).$$

602 Here again, c is a positive constant, $d_P > 0$ is the decay rate of parasite P absent its host R_2, and the
603 habitat restriction term Π extends to

$$\Pi = \left(1 - \frac{R_1 + R_2 + {}^P/q_P}{K}\right),$$

where $q_P > 1$, since typically parasites P are more efficient at using the resources of the habitat
compared to their hosts R. A ribozyme or pure enzyme molecule R_2 that is used as the replicase for P
cannot at the same time be used as the replicase for R_1, thus P parasitizes the whole autocatalytic
cycle (1).

A stable, primordial autocatalytic cycle (1^{origin}) ***in the presence of parasites***
Let us assume the primordial habitat has an abundant inflow of RNA templates P of relatively
advanced complexity from external sources, and also an abundant inflow of a primary protoenzyme
E_1. In this mixture, containing P and E_1, we assume that the following primordial autocatalytic cycle
has already evolved. The cycle produces a prototemplate F_1 and a protoreplicase E_2 obeying the
following autoca-talytic equations:

$$\frac{dE_2}{dt} = e\, E_1 F_1 \Pi - d_{E_2} E_2 \quad (1.1^{origin})$$

$$\frac{dF_1}{dt} = f\, E_2 P\, \Pi - d_{F_1} F_1 \quad \left(1.2^{origin}\right),$$

615   where e and f are positive constants and d_{E_2}, d_{F_1} are, again decay rates as the ones above. And let
616   the habitat restriction term Π be accordingly defined to fit this situation. Thus, protoenzyme E_1 and
617   temp-lates P are given through inflow into the habitat, and ribozyme E_2, which may also be a pure
618   enzyme bearing no template function, and prototemplate F_1 are autocatalytically reproduced.
619   This original primitive autocatalytic process reads in above format as follows:

$$F_1 \Rightarrow (E_1 \leftrightarrow F_1) \rightarrow^c E_2 \Rightarrow (E_2 \leftrightarrow P) \rightarrow^c F_1 \quad (1^{origin}),$$

The protoenzyme E_1 produces a protoreplicase E_2 encoded by prototemplate F_1. The fact that E_2 is
only a protoreplicase means that is not yet capable of efficiently replicating a complex RNA template
P. But provided P is above a given threshold complexity, the protoreplicase E_2 can at least produce
copies of the prototemplate F_1. As long as the habitat has a minimal threshold concentration of both,
acceptably complexed templates P and protoenzymes E_1, the archaic autocatalytic cycle (1^{origin})
can thrive and produce further copies of F_1 and E_2. Remarkably, by design the primitive autocatalytic
cycle is immune against parasite invasion by P trying to exploit the protoreplicase E_2 for its own
replication. On the contrary, the primordial autocatalytic cycle (1^{origin}) depends on template
invasion P – in other words parasite invasion – for its own reproduction.

Evolution of the primordial autocatalytic cycle (1^{origin}) ***into a full-fledged autocatalytic cycle*** (1)

We surmise that the host cycle (1) is capable of prebiotic heredity [10-12], that the primordial

autoca-talytic cycle (1^{origin}) has been conserved in the quasispecies, and therefore can again be

selected for upon parasite invasion. Therefore, the protoreplicase E_2 can be reselected from the

quasispecies. This triggered process can take the following form:

$$R_2 \Rightarrow (R_2 \leftrightarrow P) \to^c F_1 \Rightarrow (R_1 \leftrightarrow F_1) \to^c E_2 \quad (1\&2 \to 3)$$

The fully functional replicase R_2 is designed to replicate R_1 according to (1.2), but it can also be

exploited to replicate genetic parasites P according to (2). Upon encounter with a parasitic template

P, it not only replicates P but also produces via its catalyst subunit E_2 a prototemplate F_1, as did its

parent E_2 replicase. What is true for the replicase R_2 – that it is more efficient than its parent

catalyst, the protoreplicase E_2 – is also true for ribozyme R_1. It is catalytically more efficient than

protoenzyme E_1 it had replaced during evolution. R_1 can not only produce a full-fledged replicase R_2

given another copy of itself as a template, but via its protoenzyme E_1-subunit it reproduces the

protoreplicase E_2 upon encounter with prototemplate F_1:

$$F_1 \Rightarrow (R_1 \leftrightarrow F_1) \to^c E_2 \Rightarrow (E_2 \leftrightarrow P) \to^c F_1 \quad (3).$$

In order to name the processes:

(1) is the host cycle

(2) is the parasite cycle exploiting the host cycle

(3) is the hyperparasite cycle, parasitizing the host (exploiting ribozyme R_1 for protoreplicase E_2

production), and parasitizing the parasite P (exploiting its template for prototemplate F_1 production)

triggered throughout the process ($1\&2 \to 3$).

Prebiotic heredity of autocatalytic cycles to historical parasite exposure

In an environment with a constant – i.e. moderate, thus manageable – inflow of invasive parasites the

full-fledged autocatalytic host-parasite-hyperparasite cycle $(1), (2), (3)$ together with the trigger

process ($1\&2 \to 3$) becomes increasingly integrated. If the nature of the constant inflow is always of

the same special type of invasive parasites $P^{special}$, then the hyperparasite process (3) as well as the

trigger process ($1\&2 \to 3$) can mutate/adapt to processes that are fully adapted to $P^{special}$:

$$\left(3^{special}\right): \quad F_1^{special} \Rightarrow \left(R_1 \leftrightarrow F_1^{special}\right) \to^c E_2^{special} \Rightarrow \left(E_2^{special} \leftrightarrow P^{special}\right) \to^c F_1^{special}$$

$$\left(1\&2 \to 3^{special}\right): R_2 \Rightarrow \left(R_2 \leftrightarrow P^{special}\right) \to^c F_1^{special} \Rightarrow \left(R_1 \leftrightarrow F_1^{special}\right) \to^c E_2^{special}$$

It is reasonable to assume that the host-parasite-hyperparasite process $(1), (2), (3)$ together with

the trigger process ($1\&2 \to 3$) mutates under permanent special parasite inflow $P^{special}$ to an

adapted process $(1), (2), (3^{special})$, together with the adapted trigger process $\left(1\&2 \to 3^{special}\right)$.

This network of newly formed autocatalytic cycles reproduces all its elements. All adapted elements

$R_1, R_2, P^{special}, F_1^{special}, E_2^{special}$ reproduce, even when the special external parasites $P^{special}$ cease

to invade the habitat.

With the above we now can say: the primordial autocatalytic cycle $(1)^{(0)}$ has mutated to the

cycle $(1)^{(1)}$, which in turn mutates to the cycle $(1)^{(2)}$, after having faced a longer term invasion of

the same type of parasites $P^{special}$.

With this setup evolution exhibits two salient properties:

- The systems $(1)^{(0)}$, $(1)^{(1)}$, and $(1)^{(2)}$ can be viewed as increasing complexed networks of

autonomous autocatalytic cycles. They can be seen as the first steps in kickstarting evolution

towards complexed living systems.

- Prebiotic heredity [10-12] allows the cycle-network to preserve the complete record of its

historical environmental exposure to parasites:

○ $(1)^{(2)}$ preserves the record of $(1)^{(1)}$, especially the type of its infection $P^{special}$, and

○ $(1)^{(1)}$ preserves the record of its origin $(1)^{(0)}$ being immune against a broad range of

para-site infections P, and

○ $(1)^{(0)}$ preserves the record of immunity against a broad range of parasite infections P.

As evolution progresses, the autocatalytic cycles $(1)^{(2)}$ and $(1)^{(1)}$ can now be invaded by new

parasites, which can be integrated into more complexed systems, e.g. $(1)^{(3)}$ or $(1)^{(2')}$, which in turn

always keep track of the entire historical parasite exposure.

Emergence of antiparasitism in autocatalytic cycles

Let us assume the autocatalytic cycle $(1)^{(1)}$ is permanently invaded by many different types of

parasites P, to which it was not yet been exposed historically. The integration of every special

subtype of these parasites would not be viable and lead to a complexity overflow in the evolving

autocatalytic network. As first line of defense, the emergence of the autocatalytic host-parasite-

hyperparasite cycle $(1)^{(1+)} = \{(1), (2), (3), (1\&2 \rightarrow 3)\}$, as previously described would be able to

stabilize the situation, if infection P is not too overwhelming, and as time advances not too diverse.

For a more sustained, overwhelming and diverse parasite attack, another type of immunity would be

adapted, although it also originates from the already established host-parasite-hyperparasite

stabilization $(1)^{(1+)}$:

In the original primitive autocatalytic cycle (1^{origin}), $F_1 \Rightarrow (E_1 \leftrightarrow F_1) \rightarrow^c E_2 \Rightarrow (E_2 \leftrightarrow$

$P) \rightarrow^c F_1$, it is essential the both encounters $(E_1 \leftrightarrow F_1)$ and $(E_2 \leftrightarrow P)$ preserve their catalytic nature

during the whole process, since otherwise – e.g. if they are damaged, this will heavily reduce their

respective population sizes – the autocatalytic nature of (1^{origin}) could break down rapidly.

However, this is not the case for a full-fledged autocatalytic host-parasite-hyperparasite trio

$(1), (2), (3)$ in $(1)^{(1+)}$ that has already evolved. The hyperparasite $(3), F_1 \Rightarrow (R_1 \leftrightarrow F_1) \to^c E_2 \Rightarrow$

$(E_2 \leftrightarrow P) \to^c F_1$, represents the host's (1) evolutionary heredity from its primordial origin (1^{origin}).

If in this hyperparasite cycle (3), the encounter of e.g. the protoenzyme E_2 with the genetic parasite

P loses its catalytic nature, then the autocatalytic nature of the full-fledged cycle $(1), (2), (3)$ in

$(1)^{(1+)}$ would not break down. If P, or both P and E_2 are damaged during their encounter, this would

only lead to a strengthening of the host cycle (1), which is viable also in the absence of parasite and

hyperparasite cycles $(2), (3)$. As a result, the cycles $(2), (3)$ would not break down, given the fact

that the habitat still faces a sustained genetic parasite invasion P. Thus, the full-fledged cycle

$(1), (2), (3)$ would easily survive if the encounter of the protoreplicase E_2 with the genetic parasite P

would lose its catalytic nature. On the contrary, if P would be damaged, the stability, i.e. the fitness of

$(1)^{(1)}$ and with it $(1)^{(1+)}$, would increase further. Under such circumstances, we expect evolution to

establish a degenerative process of the $(E_2 \leftrightarrow P)$ encounter. We write for such a degenerated

encounter:

$$E_2 \leftrightarrow^d P,$$

707   meaning that during the encounter of E_2 with P, both elements, E_2 and P, or at least the

708   template P, are damaged and made unviable:

$$F_1 \Rightarrow (R_1 \leftrightarrow F_1) \to^c E_2 \Rightarrow \left(E_2 \leftrightarrow^d P\right) \to F_1 \quad (3')$$

Here the symbol "$\to$" means "produces". But this "production" is not catalytic anymore, since the

"producers" are damaged during the process. For an illustration of this situation where hyper-

parasitism coexists with antiparasitism (i.e. F_1 production upon encounter of E_2 with P exploiting

P as template) see Fig. 4A and Fig. 5.

The antiparasite process can now degenerate even further to $(3'')$, where the encounter

of prototemplate E_2, which we now call *antibodies,* with parasite P loses its ability to produce

the prototemplate F_1:

$$F_1 \Rightarrow (R_1 \leftrightarrow F_1) \to^c E_2 \Rightarrow \left(E_2 \leftrightarrow^d P\right) \quad (3'')$$

The ability of the antibody E_2 to produce a prototemplate F_1 upon encounter with the parasite

template P is not an evolutionary necessary one in this situation. Prototemplates F_1 are only

needed as long as parasites P exploit replicase R_2 for reproduction, but replicase R_2 already

triggers F_1-production upon encounter with parasites P through the trigger process $(1\&2 \to 3)$.

Therefore, F_1-production through the antiparasite process $(3'')$ becomes redundant.

Author contributions

Conception and design of the work: MP and BC. Data acquisition MP; data analysis MP, CI, BC;

interpretation of data MP, CI, JAC, BC. Creation of new software used in the work: MP, CI. Drafted the

work and revised it: BC, MP and JAC.

Data availability

The data, including scripts, is fully made available at the following sites (https://github.com/BICC-

UNIL-EPFL/parasite-immunity-from-hyperparasitism [github.com];

https://doi.org/10.5281/zenodo.7970788 [doi.org]).

Competing interests

The authors declare no competing interests.

760 **Figures and Figure Legends**

Fig. 1

761
762

763 **Figure 1. Habitat with one autocatalytic host cycle, variably parasite exposed**

764 Fig. 1A. The catalysts in an autocatalytic cycle (ribozyme R_2, template R_1) exposed to incoming

765 parasites P are promiscuous and hence also amplify P. Under high parasite exposure, parasites will

766 therefore outcompete host molecules.

767 Fig. 1B. Under weak parasite exposure, the host molecules R_1 and R_2 prevail.

Fig. 2

778
779

780 **Figure 2. Habitat with primordial autocatalytic cycle**

781 Parasite-specific catalysts with relatively lower catalytic activity in the habitat (E_2 protoreplicase) will

782 confer parasite resistance and thus ensure abundance of habitat molecules in the presence of

783 parasite inflow. Host molecules are expectedly less prevalent compared to Fig. 1B, at least

784 prototemplate F_1.

Fig. 3

802
803

Figure 3. Stabilized host-parasite-hyperparasite cycle with variable parasite inflow

Fig. 3 left panel (Model 1(1+)). A habitat with highly efficient catalysts (self-replicase R_2, template R_1) becomes stabilized with moderate parasite inflow via its relatively less active catalyst-subunit E_2 that confers parasite resistance. F_2 (formerly E_2 in Fig. 2) is generated as well, it acts additively with $R2$ and will be important for antiparasitism.

Fig. 3 right panel (Model 1(2)). Under constant, moderate and thus manageable parasite inflow ($P^{special}$, Suppl. Inf.), the hyperparasite cycle mutates and becomes integrated into the habitat, even when parasite inflow ceases (prebiotic heredity and mutation-selection balance). Host molecule prevalence is comparable to a habitat without parasites (Fig. 1B).

Fig. 4A

825
826

827 **Figure 4. Transition from hyperparasitism to antiparasitism under strong parasite inflow**

828 Fig. 4A Under strong, permanent exposure to complex parasite populations the hyperparasite cycle

829 partially loses its catalytic activity and becomes antiparasitic. The protoreplicase F_2 now continues to

830 function as a partially active catalyst, but also exerts a non-catalytic antiparasitic function (antibody).

831 The parameters in Fig. 4A are set to have these two activities equally represented.

Fig. 4B

848
849

850 Fig. 4B The parameters are now set to completely shift to antiparasitism (fully degenerated

851 hyperparasite cycle, completely non-catalytic). Antiparasitism has fully replaced hyperparasitism, and

852 the habitat is still stabilized, yet with stronger fluctuations.

Fig. 5

870
871

Figure 5. Gradual transition from hyperparasitism to antiparasitism in four steps

The stepwise transition from pure hyperparasitism (top panel) to pure antiparasitism (botton panel)

in four steps is shown (0, 0.25. 0.5, 0.75 and 1).