Cation permeability and pore dynamics in TRPV1 ion channels.

Miriam García-Ávila[1], Javier Tello-Marmolejo[1,2], Tamara Rosenbaum[3] and León D.
Islas[1*]

From:

[1]Department of Physiology, School of Medicine, UNAM, Mexico City, 04510, Mexico.

[2]Current affiliation: Department of Physics, University of Gothenburg, 405 30
Gothenburg, Sweden.

[3]Department of Cognitive Neuroscience, Instituto de Fisiología Celular, UNAM, Mexico
City, 04510, Mexico.

*Correspondence to:

León D. Islas

Department of Physiology

School of Medicine

Universidad Nacional Autónoma de México (UNAM)

Circuito Escolar S/N, Ciudad Universitaria

Ciudad de México, 04510, México

leon.islas@gmail.com

ORCIDs

Javier Tello-Marmolejo: https://orcid.org/0000-0002-3665-793

L.D. Islas: https://orcid.org/0000-0002-7461-5214

Miriam García-Ávila: https://orcid.org/0000-0002-5326-7302

Tamara Rosenbaum: https://orcid.org/0000-0002-4791-3195

Abstract

The Transient Receptor Vanilloid (TRPV) 1 is a non-selective ion channel which is activated by several chemical ligands and heat. We have previously shown that activation of TRPV1 by different ligands result in single-channel openings with different conductance, suggesting that the selectivity filter is highly dynamic. TRPV1 is weakly voltage-dependent, here we sought to explore whether the permeation of different monovalent ions could influence the voltage-dependence of this ion channel. By using single-channel recordings, we show that TRPV1 channels undergo rapid transitions to closed states that are directly connected to the open state, which may result from structural fluctuations of their selectivity filters. Moreover, we demonstrate that the rates of these transitions are strongly influenced by the permeant ion, suggesting that ion permeation regulates the voltage dependence of these channels.

Introduction

Transient receptor potential vanilloid (TRPV) are members of a superfamily of voltage-gated-like ion channels (Yu and Catterall, 2004). The first member of the TRPV family to be identified was TRPV1, which is a non-selective cationic channel involved in thermal sensation and which participates in the physiology of pain and inflammation (Rosenbaum et al., 2022). TRPV1 channels assemble as homotetramers, and each subunit contains large intracellular N-and C-terminal domains of complex architecture (Liao et al., 2013), and a transmembrane domain formed by six alpha helical membrane-spanning segments. Within this transmembrane domain, the transmembrane helices S1 to S4 form a helix bundle constituting a voltage sensor-like domain (VSLD), that does not seem to function as the voltage-sensor domains (VSD) present in canonical voltage-activated channels (Palovcak et al., 2015a), but which contains multiple binding sites for activating and regulatory molecules (Cao, 2020).

The pore, which contains a selectivity filter and an activation gate, is formed by the S5 segment, a pore loop and a helix motif and the S6 segment (Liao et al., 2013).

TRPV1 can be activated by varied stimuli such as capsaicin, external pH < 5.4, heat, various ligands and voltage (Caterina and Julius, 2001; Jordt et al., 2000; Matta and Ahern, 2007; Grandl et al., 2010; Rosenbaum et al., 2022). Interactions with these stimuli lead to widening of the S6 activation gate, culminating in channel opening and ion permeation (Salazar et al., 2009). Importantly, these activating factors are allosterically coupled to the channel gate and also between them (Jara-Oseguera and Swartz, 2015).

Several structures of TRPV1 and related channels (TRPV1-4) have been solved by X-ray crystallography and cryoEM methods (Liao et al., 2013; Zubcevic et al., 2018; Shimada et al., 2020; Deng et al., 2018). These structures show that in the presence of activators, the selectivity filter (SF) can adopt varying conformations, suggesting that in these channels the SF is a highly dynamic structure (Zubcevic et al., 2018). Functional experiments have shown that, while not an activation gate, the SF shows dynamic accessibility changes (Jara-Oseguera et al., 2019). It has also been suggested that voltage-induced conformational changes in this structure underlie the very small voltage dependence of activation (Yang et al., 2020).

The dynamic character of the selectivity filter of TRPV1 has not been studied functionally, although several lines of evidence point towards this characteristic. For example, different agonists of TRPV1 can open the channel to distinct single-channel conductance levels (Canul-Sánchez et al., 2018; Geron et al., 2018). We decided to characterize the effect of the permeant ions on gating of TRPV1, with emphasis on the voltage-dependence of activation. We observed very rapid transitions to closed states that are directly connected to the open state and that might represent structural fluctuations of the SF. The rates of these transitions are strongly influenced by the permeant ion, in a manner that is consistent with ion transport being involved in the voltage dependence of these channels.

Methods

Cell culture and transfection.

We used the HEK293 cell line from American Type Culture Collection (ATCC; RRID: CVCL_0045) for heterologous expression of TRPV1. Cells were grown in Dulbecco´s Modified Eagle Medium (DMEM, Invitrogen) with the addition of 10% fetal bovine serum (Invitrogen) and 1% of penicillin-streptomycin (Invitrogen) (referred to as supplemented DMEM). Cultures were incubated at 37°C, in a 5 % CO_2 atmosphere. Every 3 or 4 days the cells were washed with PBS, followed by treatment with 1 ml of 0.05% trypsin-EDTA (Invitrogen) for 2 min, then 1 ml of supplemented DMEM was added. Subsequently, the cells were mechanically dislodged and 80 µl of the cell suspension were reseeded in 30 mm culture dishes containing 5 x 5 mm glass coverslips at the bottom and 2 ml of supplemented DMEM. After 1-2 days in culture, cells were cotransfected with pcDNA3.1 plasmid with WT rat TRPV1 (rTRPV1) and pEGFP-N1 plasmid containing the eGFP fluorescent protein using the jetPEI transfection reagent (Polyplus), according to manufacturer's instructions. Single-channel recordings were done one day after transfection, while macroscopic recordings were performed two or more days after transfection.

Electrophysiological recordings.

The experiments were carried out in the inside-out configuration of the patch-clamp recording technique (Hamill et al., 1981). Recordings were performed using a pipette (extracellular) solution consisting of 130 mM NaCl, 10 mM HEPES and 5 mM EGTA, pH 7.4, and bath (intracellular) solutions in the pipette with either 130 mM NaCl, KCl, RbCl, LiCl, NH_4Cl or CsCl (depending on the experiment, as indicated in the figure legends), 10 mM HEPES, and 5 mM EGTA, pH 7.4.

Currents were recorded with an EPC-10 patch-clamp amplifier (RRID:SCR_018399, HEKA Elektronik) controlled by PatchMaster software (RRID:SCR_000034, HEKA Elektronik). Borosilicate capillaries (Sutter instruments) were used to fabricate patch-

pipettes, which had a resistance after fire polishing of 4-6 MΩ for macroscopic
currents and 8-10 MΩ for single-channel recordings.

Macroscopic currents were filtered at 1 kHz (low-pass Bessel), sampled at 40 kHz and
single-channel currents filtered at 5 kHz, sampled at 50 kHz.

Analysis of macroscopic current recordings.

The macroscopic conductance in each permeant ion condition was determined from
the current I, applying Ohm's law according to:

$$G = \frac{I}{V - V_{rev}}$$

where V is the test voltage and V_{rev} is the reversal potential. The conductance was
normalized to the maximum G_{max} and plotted as a function of voltage and was fit to a
Boltzmann function:

$$G/Gmax = \frac{1}{1+e^{-z(V-V_{1/2})/(K_B T)}} \qquad \text{Eq. 1}$$

where z is the apparent valence of gating in e_o, $V_{1/2}$ is the potential where G/G_{max}=0.5,
K_B is Boltzmann's constant and T the temperature in Kelvin.

Analysis of single-channel recordings.

Single-channel recordings were analyzed to determine the open probability, P_o.
Openings were detected employing the 50% threshold crossing method (Colquhoun
and Sigworth, 1995a). Single-channel traces were interpolated with a cubic spline
function before event detection. Open dwell time (d_o) and amplitude (a_o) of open
events and dwell times of closed events (d_c) were compiled and used to determine P_o
as:

$$P_o = \frac{d_o}{d_o+d_c} \qquad \text{Eq. 2}$$

The dwell times were also used to construct amplitude vs. open dwell time
relationships.

Analysis of fast single-channel current fluctuations.

Fast current fluctuations are apparent in single-channel records both as increases in
the open-channel noise and as incomplete closing transitions. Since the half-amplitude
crossing method is not appropriate to study these extremely fast events, we chose to
analyze these fast kinetics by studying the all-points amplitude histograms compiled
from bursts of channel activity, employing an extension of the Beta-distribution
analysis developed by (Schroeder, 2015), which is applicable to systems with more
than two states. This analysis relies on fitting of the all-points amplitude histograms to
theoretical distributions derived from time series of simulated channel activity from
specific kinetic models. These fits yield forward and backward rate constants that
were analyzed over varying conditions of permanent ion and voltages.

Model fitting.

A simulated time series of channel openings and closings from specific Markov models
of gating was first generated employing a Monte Carlo method. For exponentially
distributed dwell times, any particular, random duration of value d_i, in any closed or
open state, i, is calculated from rate constants as in:

$$d_i = -\frac{1}{\sum_{i=1}^{j \neq i} k_{ij}} ln R_n \qquad \text{Eq. 3}$$

Here, k_{ij} are the rate constants leaving state i into state j and R_n is a uniformly
distributed random number from 0 to 1. Transition probabilities between sates i and j,
p_{ij} were calculated according to:

$$p_{ij} = k_{ij} / \sum_{i=1}^{j \neq i} k_{ij}$$

Once a time series is simulated, it is filtered by convolution with the impulse response
of a Gaussian filter given by (Colquhoun and Sigworth, 1995b):

$$H(t) = \frac{1}{2\pi^{1/2}\sigma_g} exp\left(\frac{-t^2}{2\sigma_g^2}\right) \qquad \text{Eq. 4}$$

σ_g is related to the filter corner frequency f_c by:

$$\sigma_g = \frac{\sqrt{(ln2)}}{2\pi f_c}.$$

Simulated and filtered data are converted to an all-points amplitude histogram which

is then fitted to the experimental histogram using a least-squares procedure and the

fit parameters used to simulate a new time series until the fit error is minimized.

Experimental amplitude histograms were fitted to simulated histograms resulting

from several models considering one open estate and two or more closed states with

appropriate voltage dependent rate constants.

Fit to rate constants.

The voltage-dependence of each transition was calculated by plotting the rate

constants as a function of voltage and by fitting the data to the following function:

$k = k(0) \cdot e^{\left(\frac{z\,V}{K_B T}\right)}$ Eq. 5

Here, $k(0)$ is the rate constant at 0 mV, z is the apparent charge associated with the

transition, K_B is the Boltzmann constant, T is the temperature (298°K) and V is the

voltage applied.

Fast transitions showed saturation at high voltages and were thus fitted using a

subtraction of two exponential functions:

$k = k(0)_1 \cdot e^{\left(\frac{z_1 V}{K_B T}\right)} - k(0)_2 \cdot e^{\left(\frac{z_2 V}{K_B T}\right)}$ Eq. 6

Data were analyzed with Patchmaster (HEKA) and custom-written programs in Igor

Pro 8 (RRID: SCR_000325, Wave-Metrics) and Python 3 using Jupyter Notebooks

(RRID:SCR_008394).

Statistical tests.

Statistical comparisons were made by using one-way ANOVA with significance

determined from Tukey´s post hoc test. *p < 0.05 indicates statistical significance.

Results

In these experiments, we will focus on outward currents carried by varying permeant

ions, in the nominal absence of divalent ions and presence of a constant extracellular

concentration of NaCl. These conditions were chosen since it has been shown that Ca^{2+}

can permeate, block and produce Ca^{2+}-calmodulin mediated desensitization of TRPV1

(Rosenbaum et al., 2004; Numazaki et al., 2003) and that extracellular Na^+ ions are

required for normal gating in TRPV1 (Jara-Oseguera et al., 2016).

Permeation and voltage dependence of TRPV1.

Macroscopic currents through TRPV1 were activated by a saturating concentration of

capsaicin of 10 μM and recorded in asymmetric ion concentrations, with Na^+ always

present at 130 mM in the extracellular face of the channel. Outward currents were

carried by the test ion and were always preceded by recording of current in

symmetric Na^+ in the same patch, allowing the results to be compared over multiple

patches, taking as reference the condition in which the channel permeates Na^+ in both

directions, for example, red traces for only Na^+ vs. blue traces for Na^+/Li^+(Figure 1A).

As can be seen from I-V curves normalized to the current in Na^+ at 130 mV, TRPV1

permeates monovalent ions (Figure 1B) (Caterina et al., 1997a). Outward macroscopic

currents are larger when the permeant ion has a larger ionic radius like Cs^+ (1.6 $\dot{A}$).

However, the reversal potential was not statistically different from zero for each ion

tested (Figure 1B), showing that the channel does not select between monovalent

ions. Conductance (G) vs. voltage (V) relationships, again, normalized to G_{Na}, show that

the magnitude of the maximal conductance at positive voltages (G_{max}) depends on the

permeant ion. These G-V curves were fitted to Eq. 1, to estimate apparent charge, z

and voltage of half activation $V_{1/2}$. Besides the effect on G_{max}, the values of z and $V_{1/2}$,

are different for each of the permeant ions studied; specifically, there is a significant

difference in the ΔΔG with Li^+ (Compared to Na^+, Figure 1, D and E), which is brought

about by a significant increase in z (increased steepness of the G-V curve) and a

negative shift of the $V_{1/2}$.

To understand why the macroscopic conductance is affected by the permeant ion, we

performed single-channel recordings in identical ionic conditions. Our single-channel

recordings in the inside-out configuration show a clear increase in the microscopic

conductance when larger ions permeate. Based on the size of the single-channel

conductance, relative to the single-channel conductance of Na^+ current in the same

patch, we can estimate a permeability sequence as: $Li^+ < Na^+ < K^+ < Rb^+ < NH_4^+ < Cs^+$

(Figure 2A). The estimation of permeability was carried out since the reversal

potential was very close to zero for all ions (Figure 2 B) and thus a selectivity

sequence cannot be obtained.

Although the single-channel conductance explains in part the larger macroscopic

conductance when the permeant ion has a larger atomic radius, i.e., larger single-

channel conductance corresponds to larger macroscopic conductance, the microscopic

current-amplitudes observed by measuring single-channels (Figure 2B and C) do not

fully correspond to what is observed in macroscopic currents (Figure 1C). Specifically,

the single-channel conductance with Rb^+ is greater than in K^+, Na^+ and Li^+, unlike the

macroscopic results, where the conductance with Rb^+ is almost the lowest, only

comparable with the conductance with Li^+.

This difference suggests that when Rb^+ is the permeant ion at positive voltages, the

channel might gate differently. To test this possibility, we measured the single-

channel open probability according to Eq. 2, with different permeant ions. Figure 3A

shows that in the presence of Rb^+, the channel opens in short bursts, unlike thezzz

openings in Na^+, which happen in long-lived bursts. Concurrently, the open probability

is significantly smaller with Rb^+, while it is close to maximum in all other ions tested

(Figure 3B). This result reconciles the observed effects of ions on microscopic and

macroscopic conductance and reveals that the permeant ion modulates gating of

TRPV1. In particular, Rb^+ makes TRPV1 gate with lower open probability.

TRPV1 single-channel currents show a complex temporal structure; the open state is

rather unstable, showing many rapid closures or flickers and an increased open

channel noise. We observed that current flicker seems to get more noticeable with

increasing ion size and voltage (Figure 2A), suggesting that this flickery gating is a

phenomenon that arises from the interaction of the permeant ion with the pore of the

channel.

Increased open channel noise can occur when channels undergo very rapid
transitions to closed states adjacent to the open state (Rauh et al., 2018). These
dynamics can be seen as excess noise in the power spectrum of the open channel
current or as a non-Gaussian distribution of the all-points current amplitude
histogram (Heinemann and Sigworth, 1991). It has been shown that for a two-state
process, such as a unimolecular fast open channel block, the kinetics of the process
can be characterized from all-points current amplitude histograms by means of fitting
a beta distribution, which gives the amplitude of open events derived from a filtered
two-state stochastic process (Fitzhugh, 1983; Yellen, 1984).

When the fast process involves more than two states, there is no exact analytic
solution; however, it has been shown that the amplitude histograms can still be
compared with theoretical distributions derived from a simulation of a theoretical,
infinite bandwidth process filtered with the same filter and frequency of experimental
data. In this way, different explicit Markov models can be used to extract rate
constants from extended beta distributions (Rauh et al., 2017; Schroeder, 2015).

Bursts of openings of TRPV1 contain both excess noise and apparently incomplete
closures (Figure 4A), these two phenomena give rise to "tails" of excess low amplitude
events in the all-points current amplitude histograms. Interestingly these tails become
more prominent when channels conduct larger ions.

Figure 4B compares histograms for all ions tested at 90 mV. It can be observed that
the smaller ions (Li$^+$ and Na$^+$) produce amplitude histograms that are roughly
Gaussian, while for the larger ions, the histograms deviate from Gaussian and are
asymmetric, with increased counts at the left arm of the histogram formed by
amplitude values nearer to the closed amplitude. This is made more evident in Figure
4C, which scales the histograms to the maximum open amplitude for all ions. To
extract rate constants for the fast processes that give rise to these tails, we employed
extended beta distributions derived from the following Scheme I:

$$C_S \underset{\alpha_c}{\overset{\alpha_o}{\rightleftarrows}} O \underset{\beta_c}{\overset{\beta_o}{\rightleftarrows}} C_M$$
$$O \underset{\gamma_c}{\overset{\gamma_o}{\rightleftarrows}} C_f$$

This scheme postulates the existence of three closed transitions adjacent to an open state, each transition representing a slow, intermediate and fast equilibrium given by alpha, beta and gamma rate constants, respectively. Although it is known that TRPV1 channels have many open and closed states, this simplified scheme can be applied to the study of bursts of openings at sub saturating capsaicin concentrations, since the effect of capsaicin is mainly to reduce the interburst interval (Hui et al., 2003).

This scheme generates gating behavior that is very similar to the experimentally observed busts (Figure 4D) and the derived all-points histograms provide a good fit to the experimental histograms (Figure 4E). We also tested two simpler schemes with one open and two closed states and show that these cannot provide an adequate fit to the data (Supplementary figure 1; green and blue traces, respectively).

The voltage dependence of the forward and backward rate constants obtained with the extended beta distribution method for all ionic conditions is shown in Figure 5. The slow opening transition (α_o) is determined mainly by longer lived closed states and shows a slight voltage dependence in all ions, except ammonium. Inversely, the closing transition (α_c) shows a higher voltage dependence. Intermediate opening and closing transitions (β_o and β_c) have less voltage dependence while the fast transitions (γ_o and γ_c) are again more voltage dependent. While the voltage-dependence of slow and intermediate rate constants can be fit to a single exponential function (Eq. 5), the fast-opening transitions show saturation at high positive voltages and can only be fitted with the subtraction of two exponential functions (Eq. 6), (see Table I for all fitted parameters). A very interesting finding derived from these measurements is that all rate constants are modulated by the permeant ion. In general, rate constants,

when compared at 0 mV, are faster with the larger ions such as Cs⁺ and slower with
smaller ions, although this is not a rule.

The interplay between voltage and the effect of the permeant cation can be quantified
by the total charge associated with each of the three transitions, $Z_i = (z_i^o + z_i^c)$, where
$i = \alpha, \beta, \gamma$ and o, c represent the opening or closing rate constant, respectively. This is
plotted in Figure 5B for all ions. It is readily apparent that the smaller ions Li⁺ and Na⁺
are associated with a larger Z than say, Cs⁺ or Rb⁺. The finding that these transitions,
which occur near the open state, are voltage dependent with a relatively high
associated valence that depends on the size of the ion suggest that the observed
voltage-dependence of the macroscopic conductance might arise in part from the
movement of ions in the pore's electric field. This is consistent with our previous
observation that the macroscopic voltage dependence at a saturating capsaicin
concentration is modulated by the permeant ion (Figure 1D). To compare the
observed voltage dependence of rate constants with the macroscopic conductance, we
measured conductance at the same concentration of capsaicin at which rate constants
were estimated (100 nM) and compared it to the prediction of Scheme I, using the
measured rates and charges.

The open probability predicted by Scheme I is given by:

$$Po = \frac{K_\alpha}{1+K_\alpha+K_\alpha K_\beta+K_\alpha K_\gamma} \qquad \text{Eq. 7}$$

With equilibrium constants K_i given by:

$$K_i = \frac{i_o(0)e^{\frac{q_{i_o}V}{KT}}}{i_c(0)e^{\frac{q_{i_c}V}{KT}}} \qquad \text{Eq. 8}$$

Figures 6A depicts macroscopic currents measured at saturating (10 μM) and
subsaturating (100 nM) capsaicin. Figure 6C and D show G-V curves at 100 nM
capsaicin for all ions and the prediction of Scheme I (Eq. 7) using values derived from
measurements in Figure 5 (Table I). The model shows good agreement with the data
and shows that the estimated apparent charge associated with channel activation is
modulated by the permeant ion (Figure 6E).

Discussion

TRPV1 is generally characterized as a cation permeable, non-selective channel with slightly higher permeability to Ca^{2+} ions (Caterina et al., 1997b). Surprisingly, little is known of the permeation mechanisms for cations in TRPV1. A study by (Samways and Egan, 2011) showed single-channel recordings at a negative voltage that suggested non-selective permeability of cations (also based in single-channel conductance) in the order: $Na^+ > Cs^+ > Li^+ > K^+$. In this study, we have employed single-channel recordings under bi-ionic conditions and measured a permeability sequence based on the conductance at positive voltages, where the single-channel conductance is higher and easier to measure. We observed a sequence $Cs^+ > Rb^+ > K^+ > Na^+ > Li^+$. Interestingly, this is consistent with an Eisenman sequence of type I (Eisenman, 1962), which is produced by a weak field-strength cation binding site that favors permeation of easily dehydrated larger ions. Molecular dynamics (MD) simulations show that the TRPV1 pore may contain up to four binding sites for monovalent cation permeation in which the free energy for Na^+ and K^+ is very similar (Jorgensen et al., 2016), consistent with the almost identical permeability ratio for Na^+ and K^+. However, no definitive permeation mechanism has been dilucidated for TRPV channels.

Our data suggest that while non-selective, the pore of TRPV1 channels can regulate the flux of monovalent cations based on the energetics of dehydration and can differentiate between the size of ions, which is borne by the permeability sequence obtained from single-channel records (Figure 2A).

While MD simulations have suggested a highly flexible selectivity filter in TRPV1 (Darré et al., 2015), our results indicated that filter dynamics can be modulated by the permeant ion. These results prompted us to explore the kinetics of transitions near the open state. To this end, we employ the extended beta distribution method to extract rate constants that arise from incomplete closures since simpler detection methods such as half-amplitude threshold crossing are not appropriate to detect rapid incomplete closings.

In single-channel recording experiments, we observed that the rapid transitions near
the open channel seems to speed up when large ions permeate, leading to increased
open channel noise. This observation could be explained if small ions like Li^+ and Na^+
are able to remain bound for longer times to the selectivity filter, due to their stronger
electrostatic electric field, and thus stabilize the open state, reducing fluctuations. This
is also consistent with the observation that smaller ions promote a higher open
probability (Fig. 3).

The rapid fluctuations we observe are consistent with the existence of transitions
from the open state to closed states that are adjacent to the open state. It is likely that
these represent rapid pore dynamics derived from structural transitions. Cryo-EM
experiments have revealed that many TRPV channels, including TRPV1, can exist in
different pore conformations, even in the presence of activators (Zhang et al., 2021).
Functional data have also provided evidence that TRPV channels can adopt different
open conformations in the presence of different ligands (Benítez-Angeles et al., 2023;
Canul-Sánchez et al., 2018).

Experiments employing single-molecule FRET in Kir channels have directly shown
dynamic changes of the structure of the selectivity filter as a function of the presence
of Na^+ or K^+ ions (Wang et al., 2019). Similar experiments employing solution nuclear
magnetic resonance have shown similar ion-dependent dynamics of the filter of non-
selective ion channel, which have a filter structure more closely related to TRPV1's
(Lewis et al., 2021) and the intrinsic pore dynamics of TRPV1 has been demonstrated
directly by measuring the fluorescence of genetically encoded non-canonical amino
acids (Steinberg et al., 2017).

An open question in TRPV channel biophysics, specially TRPV1, is the origin of the
mild voltage dependence of the macroscopic conductance. As mentioned before, the
VLSD does not support a canonical function as a voltage sensor (Palovcak et al.,
2015b). It has recently been proposed that a conformational change that rearranges
the position of negatively charged residues at the extracellular entrance of the
selectivity filter is responsible for the voltage dependence of TRPV1 gating, which is

equivalent to the displacement of ~0.55 elementary charges per channel. We find that

in the presence of saturating capsaicin, the apparent gating charge depends on the

identity of the permeant ion. The smallest alkali metal, Li^+, is associated with a steeper

G-V, with an apparent gating charge of 0.65 e_o as well as activation at more negative

voltages.

It is known that other types of ion channels obtain their voltage-dependence from the

concerted movements of permeant ions in the permeation pathway (Schewe et al.,

2016; Marchesi et al., 2012; Pusch et al., 1995) and that permeant blockers, also

moving in the pore's electric field, can produce highly voltage-dependent block

(Spassova and Lu, 1998). For example, in chloride ClC channels, voltage dependence is

a result of the interaction of Cl^- ions with a protopore gate (Richard and Miller, 1990)

and in Kir channels where permeating ions non-canonically regulate the voltage-

dependence of gating in these channels (Lu et al., 2001; Kurata et al., 2010).

A fascinating observation derived from our study is that at relatively low

concentration of capsaicin, were the open probability P_o ~0.4, the apparent gating

charge is largest for the larger permeant cations, while at saturating concentration,

with P_o = 0.98, the apparent gating charge is largest when the smaller ion (Li^+)

permeates.

Acknowledgments

We thank Gisela Rangel, Manuel Hernández and Elsa Evaristo for excellent technical

support. This work was supported by DGAPA-PAPIIT-UNAM grant No. IN215621. M.

G-A is a doctoral student from Programa de Doctorado en Ciencias Bioquímicas-UNAM

and was supported by a doctoral thesis scholarship from CONACyT No. 776082.

References

Benítez-Angeles, M., A.E.L. Romero, I. Llorente, I. Hernández-Araiza, A. Vergara-Jaque, F.H. Real,
Ó.E. Gutiérrez Castañeda, M. Arciniega, L.E. Morales-Buenrostro, F. Torres-Quiroz, R.
García-Villegas, L.B. Tovar-Y-Romo, W.B. Liedtke, L.D. Islas, and T. Rosenbaum. 2023.
Modes of action of lysophospholipids as endogenous activators of the TRPV4 ion channel.
J. Physiol. 601:1655–1673. doi:10.1113/JP284262.

Canul-Sánchez, J.A., I. Hernández-Araiza, E. Hernández-García, I. Llorente, S.L. Morales-Lázaro, L.D.
Islas, and T. Rosenbaum. 2018. Different agonists induce distinct single-channel
conductance states in TRPV1 channels. *J. Gen. Physiol.* 150:1735–1746.
doi:10.1085/jgp.201812141.

Cao, E. 2020. Structural mechanisms of transient receptor potential ion channels. *J. Gen. Physiol.*
152:e201811998. doi:10.1085/jgp.201811998.

Caterina, M.J., and D. Julius. 2001. The Vanilloid Receptor: A Molecular Gateway to the Pain
Pathway. *Annu. Rev. Neurosci.* 24:487–517. doi:10.1146/annurev.neuro.24.1.487.

Caterina, M.J., M.A. Schumacher, M. Tominaga, T.A. Rosen, J.D. Levine, and D. Julius. 1997a. The
capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature.* 389:816–
824. doi:10.1038/39807.

Caterina, M.J., M.A. Schumacher, M. Tominaga, T.A. Rosen, J.D. Levine, and D. Julius. 1997b. The
capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature.* 389:816–
824. doi:10.1038/39807.

Colquhoun, D., and F.J. Sigworth. 1995a. Fitting and Statistical Analysis of Single-Channel Records.
In Single-Channel Recording. B. Sakmann and E. Neher, editors. Springer US, Boston, MA.
483–587.

Colquhoun, D., and F.J. Sigworth. 1995b. Fitting and Statistical Analysis of Single-Channel Records.
In Single-Channel Recording. B. Sakmann and E. Neher, editors. Springer US, Boston, MA.
483–587.

Darré, L., S. Furini, and C. Domene. 2015. Permeation and dynamics of an open-activated TRPV1
channel. *J. Mol. Biol.* 427:537–549.

Deng, Z., N. Paknejad, G. Maksaev, M. Sala-Rabanal, C.G. Nichols, R.K. Hite, and P. Yuan. 2018.
Cryo-EM and X-ray structures of TRPV4 reveal insight into ion permeation and gating
mechanisms. *Nat. Struct. Mol. Biol.* 25:252–260. doi:10.1038/s41594-018-0037-5.

Eisenman, G. 1962. Cation selective glass electrodes and their mode of operation. *Biophys. J.*
2:259–323. doi:10.1016/s0006-3495(62)86959-8.

Fitzhugh, R. 1983. Statistical properties of the asymmetric random telegraph signal, with
applications to single-channel analysis. *Math. Biosci.* 64:75–89. doi:10.1016/0025-
5564(83)90028-7.

Geron, M., R. Kumar, W. Zhou, J.D. Faraldo-Gómez, V. Vásquez, and A. Priel. 2018. TRPV1 pore
turret dictates distinct DkTx and capsaicin gating. *Proc. Natl. Acad. Sci.* 115.
doi:10.1073/pnas.1809662115.

Grandl, J., S.E. Kim, V. Uzzell, B. Bursulaya, M. Petrus, M. Bandell, and A. Patapoutian. 2010.
Temperature-induced opening of TRPV1 ion channel is stabilized by the pore domain. *Nat.*
Neurosci. 13:708–714.

Hamill, O.P., A. Marty, E. Neher, B. Sakmann, and F.J. Sigworth. 1981. Improved patch-clamp
techniques for high-resolution current recording from cells and cell-free membrane
patches. *Pflugers Arch.* 391:85–100.

Heinemann, S.H., and F.J. Sigworth. 1991. Open channel noise. VI. Analysis of amplitude
histograms to determine rapid kinetic parameters. *Biophys. J.* 60:577–587.
doi:10.1016/S0006-3495(91)82087-2.

Hui, K., B. Liu, and F. Qin. 2003. Capsaicin activation of the pain receptor, VR1: multiple open
states from both partial and full binding. *Biophys. J.* 84:2957–2968. doi:10.1016/S0006-
3495(03)70022-8.

Jara-Oseguera, A., C. Bae, and K.J. Swartz. 2016. An external sodium ion binding site controls
allosteric gating in TRPV1 channels. *eLife*. 5:e13356. doi:10.7554/eLife.13356.

Jara-Oseguera, A., K.E. Huffer, and K.J. Swartz. 2019. The ion selectivity filter is not an activation
gate in TRPV1-3 channels. *Elife*. 8:e51212.

Jara-Oseguera, A., and K.J. Swartz. 2015. Unraveling Allosteric Coupling Mechanisms in the TRPV1
Channel. *Biophys. J.* 108:124a. doi:10.1016/j.bpj.2014.11.695.

Jordt, S.-E., M. Tominaga, and D. Julius. 2000. Acid potentiation of the capsaicin receptor
determined by a key extracellular site. *Proc. Natl. Acad. Sci.* 97:8134–8139.
doi:10.1073/pnas.100129497.

Jorgensen, C., S. Furini, and C. Domene. 2016. Energetics of Ion Permeation in an Open-Activated
TRPV1 Channel. *Biophys. J.* 111:1214–1222. doi:10.1016/j.bpj.2016.08.009.

Kurata, H.T., M. Rapedius, M.J. Kleinman, T. Baukrowitz, and C.G. Nichols. 2010. Voltage-
dependent gating in a "voltage sensor-less" ion channel. *PLoS Biol.* 8:e1000315.
doi:10.1371/journal.pbio.1000315.

Lewis, A., V. Kurauskas, M. Tonelli, and K. Henzler-Wildman. 2021. Ion-dependent structure,
dynamics, and allosteric coupling in a non-selective cation channel. *Nat. Commun.*
12:6225. doi:10.1038/s41467-021-26538-8.

Liao, M., E. Cao, D. Julius, and Y. Cheng. 2013. Structure of the TRPV1 ion channel determined by
electron cryo-microscopy. *Nature*. 504:107–112. doi:10.1038/nature12822.

Lu, T., L. Wu, J. Xiao, and J. Yang. 2001. Permeant ion-dependent changes in gating of Kir2.1
inward rectifier potassium channels. *J. Gen. Physiol.* 118:509–522.
doi:10.1085/jgp.118.5.509.

Marchesi, A., M. Mazzolini, and V. Torre. 2012. Gating of cyclic nucleotide-gated channels is
voltage dependent. *Nat. Commun.* 3:973. doi:10.1038/ncomms1972.

Matta, J.A., and G.P. Ahern. 2007. Voltage is a partial activator of rat thermosensitive TRP
channels: Voltage-independent gating of TRPV1 and TRPM8. *J. Physiol.* 585:469–482.
doi:10.1113/jphysiol.2007.144287.

Numazaki, M., T. Tominaga, K. Takeuchi, N. Murayama, H. Toyooka, and M. Tominaga. 2003.
Structural determinant of TRPV1 desensitization interacts with calmodulin. *Proc. Natl.*
Acad. Sci. U. S. A. 100:8002–8006. doi:10.1073/pnas.1337252100.

Palovcak, E., L. Delemotte, M.L. Klein, and V. Carnevale. 2015a. Comparative sequence analysis
suggests a conserved gating mechanism for TRP channels. *J. Gen. Physiol.* 146:37–50.

Palovcak, E., L. Delemotte, M.L. Klein, and V. Carnevale. 2015b. Comparative sequence analysis
suggests a conserved gating mechanism for TRP channels. *J. Gen. Physiol.* 146:37–50.

Pusch, M., U. Ludewig, A. Rehfeldt, and T.J. Jentsch. 1995. Gating of the voltage-dependent
chloride channel ClC-0 by the permeant anion. *Nature.* 373:527–531.
doi:10.1038/373527a0.

Rauh, O., U.-P. Hansen, S. Mach, A.J.W. Hartel, K.L. Shepard, G. Thiel, and I. Schroeder. 2017.
Extended beta distributions open the access to fast gating in bilayer experiments-assigning
the voltage-dependent gating to the selectivity filter. *FEBS Lett.* 591:3850–3860.
doi:10.1002/1873-3468.12898.

Rauh, O., U.P. Hansen, D.D. Scheub, G. Thiel, and I. Schroeder. 2018. Site-specific ion occupation in
the selectivity filter causes voltage-dependent gating in a viral K+ channel. *Sci. Rep.*
8:10406. doi:10.1038/s41598-018-28751-w.

Richard, E.A., and C. Miller. 1990. Steady-state coupling of ion-channel conformations to a
transmembrane ion gradient. *Science.* 247:1208–1210. doi:10.1126/science.2156338.

Rosenbaum, T., A. Gordon-Shaag, M. Munari, and S.E. Gordon. 2004. Ca2+/Calmodulin Modulates
TRPV1 Activation by Capsaicin. *J. Gen. Physiol.* 123:53–62. doi:10.1085/jgp.200308906.

Rosenbaum, T., S.L. Morales-Lázaro, and L.D. Islas. 2022. TRP channels: a journey towards a
molecular understanding of pain. *Nat. Rev. Neurosci.* 23:596–610. doi:10.1038/s41583-
022-00611-7.

Salazar, H., A. Jara-Oseguera, E. Hernandez-Garcia, I. Llorente, I.I. Arias-Olguin, M. Soriano-Garcia,
L.D. Islas, and T. Rosenbaum. 2009. Structural determinants of gating in the TRPV1
channel. *Nat. Struct. Mol. Biol.* 16:704–710.

Samways, D.S.K., and T.M. Egan. 2011. Calcium-dependent decrease in the single-channel
conductance of TRPV1. *Pflüg. Arch. - Eur. J. Physiol.* 462:681–691. doi:10.1007/s00424-
011-1013-7.

Schewe, M., E. Nematian-Ardestani, H. Sun, M. Musinszki, S. Cordeiro, G. Bucci, B.L. de Groot, S.J.
Tucker, M. Rapedius, and T. Baukrowitz. 2016. A Non-canonical Voltage-Sensing
Mechanism Controls Gating in K2P K+ Channels. *Cell*. 164:937–949.
doi:10.1016/j.cell.2016.02.002.

Schroeder, I. 2015. How to resolve microsecond current fluctuations in single ion channels: The
power of beta distributions. *Channels*. 9:262–280. doi:10.1080/19336950.2015.1083660.

Shimada, H., T. Kusakizako, T.H. Dung Nguyen, T. Nishizawa, T. Hino, M. Tominaga, and O. Nureki.
2020. The structure of lipid nanodisc-reconstituted TRPV3 reveals the gating mechanism.
Nat. Struct. Mol. Biol. 27:645–652. doi:10.1038/s41594-020-0439-z.

Spassova, M., and Z. Lu. 1998. Coupled Ion Movement Underlies Rectification in an Inward-
Rectifier K+ Channel. *J. Gen. Physiol.* 112:211–221. doi:10.1085/jgp.112.2.211.

Steinberg, X., M.A. Kasimova, D. Cabezas-Bratesco, J.D. Galpin, E. Ladron-de-Guevara, F. Villa, V.
Carnevale, L. Islas, C.A. Ahern, and S.E. Brauchi. 2017. Conformational dynamics in TRPV1
channels reported by an encoded coumarin amino acid. *eLife*. 6:e28626.
doi:10.7554/eLife.28626.

Wang, S., S.-J. Lee, G. Maksaev, X. Fang, C. Zuo, and C.G. Nichols. 2019. Potassium channel
selectivity filter dynamics revealed by single-molecule FRET. *Nat. Chem. Biol.* 15:377–383.
doi:10.1038/s41589-019-0240-7.

Yang, F., L. Xu, B.H. Lee, X. Xiao, V. Yarov-Yarovoy, and J. Zheng. 2020. An Unorthodox Mechanism
Underlying Voltage Sensitivity of TRPV1 Ion Channel. *Adv. Sci.* 7:2000575.
doi:10.1002/advs.202000575.

Yellen, G. 1984. Ionic permeation and blockade in Ca2+-activated K+ channels of bovine
chromaffin cells. *J. Gen. Physiol.* 84:157–186. doi:10.1085/jgp.84.2.157.

Yu, F.H., and W.A. Catterall. 2004. The VGL-Chanome: A Protein Superfamily Specialized for
Electrical Signaling and Ionic Homeostasis. *Sci. STKE*. 2004. doi:10.1126/stke.2532004re15.

Zhang, K., D. Julius, and Y. Cheng. 2021. Structural snapshots of TRPV1 reveal mechanism of
polymodal functionality. *Cell*. 184:5138-5150.e12. doi:10.1016/j.cell.2021.08.012.

Zubcevic, L., S. Le, H. Yang, and S.-Y. Lee. 2018. Conformational plasticity in the selectivity filter of
the TRPV2 ion channel. *Nat. Struct. Mol. Biol.* 25:405–415. doi:10.1038/s41594-018-0059-
z.

Figure 1. The voltage dependence of TRPV1 is influenced by the permeant ion.

A) Representative inside-out current-traces recorded in the same patch, elicited by 20 mV steps from -130 mV to 130 mV for a duration of 150 ms. The extracellular solution (pipette) contained 130 mM NaCl, this patch was first exposed to internal 130 mM

NaCl (black) followed by exposure to 130 mM NaCl + 10 μM capsaicin (red) and finally
to 130 mM LiCl + 10 μM capsaicin (blue). All experiments were performed following
the same experimental strategy to compare the permeation of different monovalent
ions (Li^+, K^+, Rb^+, NH_4^+, Cs^+) with respect to the permeation of Na^+ in the same patch.
B) Averages of current-voltage (I-V) relations for all ions tested. The number of
experiments, n, is in parenthesis. **C)** Conductance-voltage (G-V) relations for
monovalent ions, obtained from data as in B in the presence of 10 μM capsaicin,
normalized to data with a symmetrical concentration of Na^+ from each patch. G-V
relations were fitted to the Boltzmann function (lines). **D)** Scatter plots of the
apparent charge (Z; above) and $V_{1/2}$ (below) associated with channel opening, both
were estimated by fitting a Boltzmann function to G-V curves in B. The values of Z for
Li^+ were compared for statistical differences with all other ions, the p values are: *p =
0.00007 for Li^+ vs. Na^+; NS (not significant) p=0.213 for Li^+ vs. K^+; *p = 0.00034 for Li^+
vs. Rb^+; *p = 0.00005 for Li^+ vs. NH_4^+; *P = 0.01 for Li^+ vs. Cs^+. For $V_{1/2}$: *p = 0.0000009
for Li^+ vs. Na^+; *p = 0.00001 for Li^+ vs. K^+; *p = 0.00000001 for Li^+ vs. Rb^+; *P =
0.000002 for Li^+ vs. NH_4^+; *p = 0.02 for Li^+ vs. Cs^+. **E)** Scatter plot of the $\Delta\Delta G$ associated
to the ion effect in channel opening was estimated as: $\Delta\Delta G = \Delta G_{Ion} - \Delta G_{Na}$ + the data
was taken from the fit in B. *p = 0.00005 for Li^+ vs. K^+; *p = 0.0000002 for Li^+ vs. Rb^+;
*p = 0.000004 for Li^+ vs. NH_4^+; *p = 0.003 for Li^+ vs. Cs^+. Data are shown as mean ±
SEM. The significant difference was determined with a one-way ANOVA test followed
by a Tukey´s test post-hoc (*p < 0.05).

Figure 2

Figure 2. TRPV1 permeability to monovalent ions is size-dependent.

A) Representative current traces showing openings from longer single-channel recordings in the presence of varying permeant ions. Inside-out patches were first exposed to internal 130 mM Na$^+$ + 100 nM capsaicin and then to internal 130 mM (Li$^+$ (blue), Na$^+$ (red) K$^+$(mustard), Rb$^+$ (purple), NH$_4^+$ (black), Cs$^+$ (green)) + 100 nM capsaicin similarly to experiments in Figure 1A. The dashed line represents the closed state of the channel. B) Representative I-V relations from single-channel records. These were obtained from data as in A for each ion. C) Current carried by each ion relative to the single-channel current amplitude with Na$^+$. Ionic radii in Å are also shown for comparison. Data are mean ± SEM, n is shown in parentheses.

591
Figure 3. Rubidium ions strongly modulate the single-channel open probability
of TRPV1 channels.

A) Representative current traces from single-channel recordings at 90 mV when
intracellular Rb+ (purple) or Na+ (red) were used as permeant ions. B) Open
probability of TRPV1 channels at 90 mV in the presence of different permeant ions
and 100 nM of capsaicin, calculated from traces of current recordings as shown in A.
Data are mean ± s.e.m.

Figure 4. Fluctuations in the single-channel current during the open state can be fitted by a Markov model.

A) Representative current trace during a typical single TRPV1 opening burst recorded in the inside-out configuration at 90 mV, with internal 130 mM RbCl + 100 nM capsaicin. B) Normalized all-points histograms showing single-channel current amplitudes of openings elicited by 100 nM capsaicin at 90 mV, for each ion tested. The histograms are shown aligned to the closed state amplitude peak (left). C) Close-up of the tails of events of the all-points histograms in B aligned to the maximum current amplitude of the open state to highlight the asymmetric form of the histograms. D) Current trace of a single-channel at 90 mV from an inside-out patch exposed to internal 130 mM NH$_4$Cl + 100 nM capsaicin (black) and simulation with the model in Scheme I (blue). These simulated recordings are used to produce simulated all-points amplitude histograms of single-channel current, which are then fitted to experimental all-points amplitude histograms. E) Typical fit of a simulated histogram (blue) to experimental histogram from single current elicited by 110 mV with internal 130 mM NH$_4$Cl + 100 nM capsaicin (black) using the Markov model in Scheme I. Parameters obtained from the fit are: α_o = 1500; α_c = 27; β_o = 22500; β_c = 600; γ_o = 127500; γ_c = 30000; χ^2 = 0.00013873.

Figure 5. Voltage-dependent transitions near the open state of TRPV1 channels are modulated by the permeant ions.

A) Rate constants obtained from the Markov model fit to the histograms as in Figure 4D. The rate constants were classified as slow, medium, and fast according to their magnitude at 0 mV. The continuous curves represent the fit to Eq. 5 or 6.

625 B) Scatter plot of the sum of the apparent charge by transition ($Z^i_c + Z^i_o$ where I = α, β

626 or γ) obtained from the fit of equations 5 or 6 to rate constants in A. Black bars are

627 mean ± s.e.m.

Figure 6

Figure 6. The opening probability predicted by the Markov model explains the macroscopic experimental behavior.

A) Representative inside-out macroscopic current traces recorded in the same patch, elicited by 20 mV steps from -130 mV for 150 ms. The patch was exposed to internal 130 mM NaCl + 10 μM capsaicin (red) followed by 130 mM RbCl + 10 μM capsaicin (purple). B) Representative inside-out current traces like A but using 100 nM capsaicin. C) Normalized conductance-voltage (G-V) relations with monovalent ions, obtained from inside-out patches after treatment with internal 100 nM capsaicin. The G-V relation was fitted to the Boltzmann function (line). Data are shown as mean ± s.e.m., number of experiments n, in parenthesis. D) Normalized prediction of open probability by a Markov model for each monovalent ion. E) Bar graph of apparent charge associated with channel opening estimated by fitting a Boltzmann equation to curves in A and B. Data is shown as mean ± s.e.m.

Supplementary figure 1.

Supplementary Figure 1. Amplitude histogram fits to three different Markov models.

A) Markov models with 1 closed state (green), two closed states (blue) and three closed states (red). B) Fit of the simulated histograms from the models in A to experimental an all-points histogram of single channel current of TRPV1(black), elicited by a pulse to 90 mV from an inside-out patches exposed to 130 mM NaCl + 100 nM capsaicin. The fits parameters used for 1 close state were: α_o = 301.9535; α_c = 36.1664; χ^2 = 0.0697; 2 closed states α_o = 300; α_c = 37.5; γ_o = 28500; γ_c = 13500; χ^2 = 0.002342; three closed states α_o = 280.61; α_c = 49; β_o = 14724.475; β_c = 1133.16; γ_o = 52500; γ_c = 15396; χ^2 = 0.00017296.