Cation permeability and pore dynamics in TRPV1 ion channels.

Miriam García-Ávila[1], Javier Tello-Marmolejo[1,2], Tamara Rosenbaum[3] and León D.

Islas[1]*

From:

[1]Department of Physiology, School of Medicine, UNAM, Mexico City, 04510, Mexico.

[2]Current affiliation: Department of Physics, University of Gothenburg, 405 30
Gothenburg, Sweden.
[3]Department of Cognitive Neuroscience, Instituto de Fisiología Celular, UNAM, Mexico
City, 04510, Mexico.
11

*Correspondence to:

León D. Islas
Department of Physiology
School of Medicine
Universidad Nacional Autónoma de México (UNAM)
Circuito Escolar S/N, Ciudad Universitaria
Ciudad de México, 04510, México
leon.islas@gmail.com
20

21 ORCIDs

22 Javier Tello-Marmolejo: https://orcid.org/0000-0002-3665-793
23 L.D. Islas: https://orcid.org/0000-0002-7461-5214
24 Miriam García-Ávila: https://orcid.org/0000-0002-5326-7302
25 Tamara Rosenbaum: https://orcid.org/0000-0002-4791-3195
26

Abstract

The Transient Receptor Vanilloid 1 (TRPV1) is a non-selective ion channel, which is

activated by several chemical ligands and heat. We have previously shown that

activation of TRPV1 by different ligands result in single-channel openings with

different conductance, suggesting that the selectivity filter is highly dynamic. TRPV1 is

weakly voltage-dependent, here we sought to explore whether the permeation of

different monovalent ions could influence the voltage-dependence of this ion channel.

By using single-channel recordings, we show that TRPV1 channels undergo rapid

transitions to closed states that are directly connected to the open state, which may
result from structural fluctuations of their selectivity filters. Moreover, we
demonstrate that the rates of these transitions are ~~strongly~~ influenced by the
permeant ion, suggesting that ion permeation regulates the voltage dependence of
these channels.

Introduction

Transient receptor potential vanilloid (TRPV) channels are members of a superfamily
of voltage-gated-like ion channels (Yu and Catterall, 2004). The first member of the
TRPV family to be identified was TRPV1, which is a non-selective cationic channel
involved in thermal sensation and which participates in the physiology of pain and
inflammation (Rosenbaum et al., 2022). TRPV1 channels assemble as homotetramers,
and each subunit contains large intracellular N-and C-terminal domains of complex
architecture (Liao et al., 2013), and a transmembrane domain formed by six alpha
helical membrane-spanning segments. Within this transmembrane domain, the
transmembrane helices S1 to S4 form a helix bundle constituting a voltage sensor-like
domain (VSLD), that does not seem to function as the voltage-sensor domains (VSD)
present in canonical voltage-activated channels (Palovcak et al., 2015a), but which
contains multiple binding sites for activating and regulatory molecules (Cao, 2020).
The pore, which <u>encompasses</u> a selectivity filter and an activation gate, is formed by
the S5 segment, a pore loop and a helix motif and the S6 segment (Liao et al., 2013).

TRPV1 can be activated by varied stimuli such as capsaicin, external pH < 5.4, heat,
various ligands and voltage (Caterina and Julius, 2001; Jordt et al., 2000; Matta and
Ahern, 2007; Grandl et al., 2010; Rosenbaum et al., 2022). Interactions with these
stimuli lead to widening of the S6 activation gate, culminating in channel opening and
ion permeation (Salazar et al., 2009). Importantly, these activating factors are
allosterically coupled to the channel gate and also between them (Jara-Oseguera and
Swartz, 2015).

Several structures of TRPV1 and related channels (TRPV1-4) have been solved by X-ray crystallography and cryoEM methods (Liao et al., 2013; Zubcevic et al., 2018; Shimada et al., 2020; Deng et al., 2018). These structures show that in the presence of activators, the selectivity filter (SF) can adopt varying conformations, suggesting that in these channels the SF is a highly dynamic structure (Zubcevic et al., 2018). Functional experiments have shown that, while not an activation gate, the SF shows dynamic accessibility changes (Jara-Oseguera et al., 2019). It has also been suggested that voltage-induced conformational changes in this structure underlie the very small voltage dependence of activation (Yang et al., 2020).

The dynamic character of the selectivity filter of TRPV1 has not been studied functionally, although several lines of evidence point towards this characteristic. For example, different agonists of TRPV1 can open the channel to distinct single-channel conductance levels (Canul-Sánchez et al., 2018; Geron et al., 2018). Hence, we decided to characterize the effect of the permeant ions on the gating of TRPV1. We observed very rapid transitions to closed states that are directly connected to the open state and that might represent structural fluctuations of the SF. The rates of these transitions are strongly influenced by the permeant ion, in a manner that is consistent with ion transport being involved in the regulation of gating of these channels.

Methods

Cell culture and transfection.

We used the HEK293 cell line from American Type Culture Collection (ATCC; RRID: CVCL_0045) for heterologous expression of TRPV1. Cells were grown in Dulbecco′s Modified Eagle Medium (DMEM, Invitrogen) with the addition of 10 % fetal bovine serum (Invitrogen) and 1% of penicillin-streptomycin (Invitrogen) (referred to as supplemented DMEM). Cultures were incubated at 37°C, in a 5 % CO_2 atmosphere. Every 3 or 4 days, the cells were washed with PBS, followed by treatment with 1 ml of 0.05% trypsin-EDTA (Invitrogen) for 2 min, then 1 ml of supplemented DMEM was added. Subsequently, the cells were mechanically dislodged and 80 µl of the cell suspension were reseeded on 5 x 5 mm glass coverslips in 30 mm culture dishes with

2 ml of supplemented DMEM. After 1-2 days in culture, cells were cotransfected with

pcDNA3.1 plasmid with WT rat TRPV1 (rTRPV1) and pEGFP-N1 plasmid containing

the eGFP fluorescent protein using the jetPEI transfection reagent (Polyplus),

according to manufacturer's instructions. Single-channel recordings were done one

day after transfection, while macroscopic recordings were performed two or more

days after transfection.

Electrophysiological recordings.

The experiments were carried out in the inside-out configuration of the patch-clamp

recording technique (Hamill et al., 1981). Recordings were performed using a pipette

(extracellular) solution consisting of 130 mM NaCl, 10 mM HEPES and 5 mM EGTA, pH

7.4, and bath (intracellular) solutions with either 130 mM NaCl, KCl, RbCl, LiCl, NH$_4$Cl

or CsCl (depending on the experiment, as indicated in the figure legends), 10 mM

HEPES, and 5 mM EGTA, pH 7.4.

Currents were recorded with an EPC-10 patch-clamp amplifier (RRID:SCR_018399,

HEKA Elektronik) controlled by PatchMaster software (RRID:SCR_000034, HEKA

Elektronik). Borosilicate capillaries (Sutter instruments) were used to fabricate patch-

pipettes, which had a resistance after fire polishing of 4-6 MΩ for macroscopic

currents and 8-10 MΩ for single-channel recordings.

Macroscopic currents were filtered <u>with the build-in 4-pole Bessel filter of the EPC-10</u>

at 1 kHz <u>(-3dB) and sampled at 40 kHz while single-channel currents were filtered at</u>

<u>5 kHz and sampled at 50 kHz.</u>

Analysis of macroscopic current recordings.

The macroscopic conductance in each permeant ion condition was determined from

the current *I,* applying Ohm's law according to:

$$G = \frac{I}{V - V_{rev}}$$

where V is the test voltage and V_{rev} is the reversal potential. The conductance was

normalized to the maximum G_{max} and plotted as a function of voltage and was fit to a

Boltzmann function:

$$\frac{G}{Gmax} = \frac{1}{1 + e^{\frac{z(V - V_{1/2})}{K_B T}}}$$ Eq. 1

where z is the apparent valence of gating in e_o, $V_{1/2}$ is the potential where $G/G_{max}=0.5$,

K_B is Boltzmann's constant and T the temperature in Kelvin (296 K). The apparent free

energy change of activation by voltage was estimated as $\Delta G_{ion} = zV_{1/2}$ from values

obtained for each ion from fits to Eq. 1. $\Delta\Delta G$ was estimated as $\Delta G_{ion} - \Delta G_{Na}$

Analysis of single-channel recordings.

Single-channel recordings were analyzed to determine the open probability, P_o.

Openings were detected employing the 50% threshold crossing method (Colquhoun

and Sigworth, 1995a). Single-channel traces were interpolated with a cubic spline

function before event detection. Dwell times (d_o) of open events and dwell times of

closed events (d_c) were compiled and used to determine P_o as:

$P_o = \frac{d_o}{d_o + d_c}$ Eq. 2

~~The dwell times were also used to construct amplitude vs. open dwell time~~

~~relationships.~~

Single-channel kinetics were analyzed by constructing open and closed dwell time

distributions using log-transformed histograms according to (Sigworth and Sine,

1987). Events shorter than the filter death time T_d were discarded. $T_d = 0.179/f_c$. At

$f_c = 5$ kHz, $T_d = 36$ µs. Bursts were defined as groups of openings separated by a closed

time longer than a critical time, t_c. This critical time was calculated from the closed

time distribution from recordings in Li^+, Na^+, Rb^+ and Cs^+ according to (Colquhoun and

Sakmann, 1985). Closed, open and burst length distributions were fit to a sum of

exponential components. The number of components and their parameters are given

in the figure legends.

Analysis of fast single-channel current fluctuations.

Fast current fluctuations are apparent in single-channel records both as increases in

the open-channel noise and as incomplete closing transitions. Since the half-amplitude

crossing method is not appropriate to study these extremely fast events, we chose to

analyze these fast kinetics by studying the all-points amplitude histograms compiled

from bursts of channel activity, employing an extension of the Beta-distribution

analysis developed by (Schroeder, 2015), which has been shown to be applicable to

systems with more than two states. This analysis relies on fitting of the all-points

amplitude histograms to theoretical distributions derived from time series of

simulated channel activity from specific kinetic models. These fits yield forward and

backward rate constants that were analyzed over varying conditions of permeant ion

and voltages. Bursts of channel openings were selected manually from recordings and

included short periods of no channel activity in order to adjust the zero current level.

Experimental all-points histograms were constructed by accumulating the currents

from these selected bursts.

Model fitting.

A simulated time series of channel openings and closings from specific Markov models

of gating was first generated employing a Monte Carlo method. For exponentially

distributed dwell times, any particular random duration of value d_i, in any closed or

open state, i, is calculated from rate constants as:

$d_i = -\frac{1}{\sum_{i=1}^{j \neq i} k_{ij}} ln R_n$ Eq. 3

Here, k_{ij} are the rate constants leaving state i into state j and R_n is a uniformly

distributed random number from 0 to 1. Transition probabilities between sates i and j,

p_{ij} were calculated according to:

$p_{ij} = \frac{k_{ij}}{\sum_{i=1}^{j \neq i} k_{ij}}$

Once a time series is simulated, Gaussian noise with a variance similar to the variance

of the closed channel current level of each recording was added and the time series

was filtered by convolution with the impulse response of a Gaussian filter given by

(Colquhoun and Sigworth, 1995b):

$H(t) = \frac{1}{\sqrt{2\pi}\sigma_g} exp\left(\frac{-t^2}{2\sigma_g^2}\right)$ Eq. 4

σ_g is related to the filter corner frequency f_c by:

179 $\sigma_g = \frac{\sqrt{(ln2)}}{2\pi f_c}.$

180 Simulated and filtered data are converted to an all-points amplitude histogram which

181 is then fitted to the experimental histogram using a least-squares procedure and the

182 fit parameters used to simulate a new time series until the fit error is minimized. We

183 implemented a zero-order optimization algorithm with a number of iterations i = 100.

184 The merit function was $R_n(i+1) < R_n(i)$, where $R_n(i)$ is an average of n least squares

185 values between a simulation and the experimental data. The number of simulations

186 per iteration, was n = 10. It was essential to average these least squares values $R_n(i)$

187 since the simulation is a random process and non-optimal values could produce a low

188 R because of this randomness. Two types of steps were taken with equal probability in

189 the variation of the fitting parameters. A small step that looks for the minimum R

190 value and a large step. The large step was important to avoid getting stuck in local

191 minima.

192 Experimental amplitude histograms were fitted to simulated histograms resulting

193 from several models considering one open estate and one, two or more closed states

194 with appropriate voltage dependent rate constants.

195 *Fit to rate constants.*

196 The voltage-dependence of each transition was calculated by plotting the rate

197 constants as a function of voltage and fitting the data to the following function:

198 $k = k(0) \cdot e^{\left(\frac{zV}{K_BT}\right)}$    Eq. 5

199 Here, $k(0)$ is the rate constant at 0 mV, z is the apparent charge associated with the

200 transition, K_B is the Boltzmann constant, T is the temperature (296 K) and V is the

201 voltage applied.

202 ~~Fast transitions showed saturation at high voltages and were thus fitted using a~~

203 ~~subtraction of two exponential functions:~~

204 ~~$k = k(0)_1 \cdot e^{\left(\frac{z_1V}{K_BT}\right)} - k(0)_2 \cdot e^{\left(\frac{z_2V}{K_BT}\right)}$ Eq. 6~~

205 Data were analyzed with Patchmaster (HEKA) and custom-written programs in Igor

206 Pro 8 (RRID: SCR_000325, Wave-Metrics) and Python 3 using Jupyter Notebooks

207 (RRID:SCR_008394).

Statistical tests.

Statistical comparisons were made by using one-way ANOVA with significance

determined from Tukey´s post hoc test. *p < 0.05 indicates statistical significance.

Results

In these experiments, we focused on outward currents carried by varying permeant

ions, in the nominal absence of divalent ions and presence of a constant extracellular

concentration of NaCl. These conditions were chosen since it has been shown that Ca^{2+}

can permeate, block and produce Ca^{2+}-calmodulin mediated desensitization of TRPV1

(Rosenbaum et al., 2004; Numazaki et al., 2003) and that extracellular Na^+ ions are

required for normal gating in TRPV1 (Jara-Oseguera et al., 2016).

Permeation and voltage dependence of TRPV1.

Macroscopic currents through TRPV1 were activated by a saturating concentration of

capsaicin (CAP) of 10 μM and recorded in asymmetric ion concentrations, with Na^+

always present at 130 mM in the extracellular face of the channel. Outward currents

were carried by the test ion and were always preceded by recording of current in

symmetric Na^+ in the same patch, allowing the results to be compared over multiple

patches, taking as reference the condition in which the channel permeates Na^+ in both

directions. For example, red traces for only Na^+ vs. blue traces for Na^+/Li^+ (Figure 1A).

As can be seen from I-V curves normalized to the current in Na^+ at 130 mV, TRPV1

permeates monovalent ions (Figure 1B) (Caterina et al., 1997a).

Outward macroscopic currents are larger when the permeant ion has a larger ionic

radius like Cs^+ (1.6 $Å$). However, the reversal potential was not statistically different

from zero for each ion tested (Figure 1B), showing that the channel does not select

between monovalent ions. Conductance (G) vs. voltage (V) relationships, again,

normalized to G_{Na}, show that the magnitude of the maximal conductance at positive

voltages (G_{max}) depends on the permeant ion. These G-V curves were fitted to Eq. 1, to

estimate apparent charge, z and voltage of half activation $V_{1/2}$. Besides the effect on

G_{max}, the values of z and $V_{1/2}$, are different for each of the permeant ions studied;

specifically, there is a significant difference in the $\Delta\Delta G$ with Li^+ (Compared to Na^+,

Figure 1, D and E), which is brought about by a significant increase in z (increased

steepness of the G-V curve) and a negative shift of the $V_{1/2}$.

To understand why the macroscopic conductance is affected by the permeant ion, we

performed single-channel recordings in identical ionic conditions. Our single-channel

recordings in the inside-out configuration show a clear increase in the microscopic

conductance when larger ions permeate (Figure 2A). Since the reversal potential was

very close to zero for all ions (Figure 2B), to estimate the cation permeability, we used

the magnitude of the single-channel conductance, relative to the single-channel

conductance of Na^+ current in the same patch. We can estimate the permeability

sequence as: $Li^+ < Na^+ < K^+ < Rb^+ < NH_4^+ < Cs^+$ (Figure 2C).

Although the single-channel conductance explains in part the larger macroscopic

conductance when the permeant ion has a larger atomic radius, i.e., larger single-

channel conductance corresponds to larger macroscopic conductance, the microscopic

current-amplitudes observed by measuring single-channels (Figure 2B) do not fully

correspond to what is observed in macroscopic currents (Figure 1C). Specifically, the

single-channel conductance with Rb^+ is greater than in K^+, Na^+ and Li^+, unlike the

macroscopic results, where the conductance with Rb^+ is almost the lowest, only

comparable with the conductance with Li^+.

This difference suggests that when Rb^+ is the permeant ion at positive voltages, the

channel might gate differently. To test this possibility, we measured the single-

channel open probability according to Eq. 2, with different permeant ions. Figure 3A

shows that in the presence of Rb^+, the channel opens in short bursts, unlike the

openings in Na^+, which happen in long-lived bursts <u>at this concentration of capsaicin</u>.

Concurrently, the open probability is significantly smaller with Rb^+, while it is close to

the maximum <u>attained P_o at this capsaicin concentration (100 nM) and voltage in all</u>

<u>other ions tested (Figure 3B). Since the macroscopic current is proportional to the</u>

<u>product of P_o and single channel I-V,</u> this result reconciles the observed effects of ions

on microscopic and macroscopic conductance and reveals that the permeant ion

modulates gating of TRPV1. In particular, Rb^+ makes TRPV1 gate with lower open

probability.

To further explore the gating effects of permeant ions, we performed an analysis of

the kinetics of channel opening and closing. TRPV1 gating at the single-channel level is

complex (Hui et al., 2003), the open and closed dwell time histograms can be fitted by

a sum of three and four exponential components, respectively, indicating the presence

of multiple open and closed states (Supplementary figure 1 and Table I), with the

channel mostly opening in bursts. For comparison, we obtained the burst length

distributions in Li^+, Na^+, Cs^+ and Rb^+. As shown in Figure 4, burst length is also

multiexponential and the bursts become shorter when Rb^+ is the permeant ion. Burst

length distributions in Na^+, Li^+ and Cs^+ are very similar.

TRPV1 single-channel currents show a complex temporal structure, as mentioned

before. Moreover, the open state is rather unstable, showing many rapid closures or

flickers and an increased open channel noise. We observed that current flicker is more

noticeable with increasing ion size and voltage (Figure 2A), suggesting that this

flickery gating is a phenomenon that arises from the interaction of the permeant ion

with the pore of the channel.

Increased open channel noise can occur when channels undergo very rapid

transitions to closed states adjacent to the open state (Rauh et al., 2018). These

dynamics can be seen as excess noise in the power spectrum of the open channel

current or as a non-Gaussian distribution of the all-points current amplitude

histogram (Heinemann and Sigworth, 1991a). It has been shown that for a two-state

process, such as a unimolecular fast open channel block, the kinetics of the process

can be characterized from all-points current amplitude histograms by means of fitting

a beta distribution, which gives the amplitude of open events derived from a filtered

two-state stochastic process (Fitzhugh, 1983; Yellen, 1984).

When the fast process involves more than two states, there is no exact analytic

solution; however, it has been shown that the amplitude histograms can still be

compared with theoretical distributions derived from a simulation of an infinite

bandwidth process filtered with the same filter and frequency of experimental data. In

this way, different explicit Markov models can be used to extract rate constants from

extended beta distributions (Rauh et al., 2017; Schroeder, 2015). Importantly, this

<u>approach is able to estimate kinetic events that are faster than the filter rise time</u>

(Heinemann and Sigworth, 1991b).

Bursts of openings of TRPV1 contain both excess noise and apparently incomplete

closures (Figure 5A), these two phenomena <u>make the amplitude histograms deviate</u>

<u>from a Gaussian distribution and</u> give rise to "tails" of excess low amplitude events in

the all-points current amplitude histograms. Interestingly these tails become more

prominent when channels conduct larger ions.

Figure 5B compares histograms for all ions tested at 90 mV. It can be observed that

the smaller ions (Li^+ and Na^+) produce amplitude histograms that are roughly

Gaussian, while for the larger ions, the histograms deviate from Gaussian and are

asymmetric, with increased counts on the left arm of the <u>histogram contributed by</u>

<u>amplitude values that approach the closed amplitude (Supplementary Figure 2).</u> This

is made more evident in Figures 5C and D, which scale the histograms to the

maximum open amplitude for all ions <u>and between the closed and open amplitudes,</u>

<u>respectively. These figures show greater excess noise for the openings with larger</u>

<u>cations.</u>

To extract rate constants for the fast processes that give rise to these tails, we

employed extended beta distributions derived from the following Scheme I:

This scheme postulates the existence of three closed transitions adjacent to an open

state, each transition representing a slow, medium and fast equilibrium given by

alpha, beta and gamma rate constants, respectively. Although it is known that TRPV1

channels have many open and closed states, this simplified scheme can be applied to

the study of bursts of openings at sub saturating capsaicin concentrations, since the

effect of capsaicin is mainly to reduce the inter burst interval (Hui et al., 2003).

This scheme generates gating behavior that is very similar to the experimentally

observed currents (<u>Figure 5E</u>) and the derived all-points histograms provide a good

fit to the experimental histograms (<u>Figure 5F</u>). We also tested two simpler schemes

with one open and one or two closed states and show that these cannot provide an

adequate fit to the data (Supplementary figure 3; green and blue traces, respectively).

The voltage-dependence of the forward and backward rate constants obtained with

the extended beta distribution method for all ionic conditions is shown in Figure 6.

The slow opening transition (α_o) is determined mainly by longer lived closed states

and shows a slight voltage-dependence in all ions, except ammonium, <u>which tends to</u>

<u>have a higher valence</u>. Inversely, the closing transition (α_c) shows a higher voltage-

dependence. Intermediate opening and closing transitions (β_o and β_c) have less

voltage- dependence, while the fast transitions (γ_o and γ_c) are again more voltage

dependent. The voltage-dependence of these transition rate constants was fit to a

single exponential function of voltage (Eq. 5). ~~the fast-opening transitions show~~

~~saturation at high positive voltages and can only be fitted with the subtraction of two~~

~~exponential functions (Eq. 6),~~ (<u>see Table II for all fitted parameter</u>). An interesting

finding derived from these measurements is that the rate constants <u>for all three</u>

<u>transitions</u> are modulated by the permeant ion. <u>When plotting the values of the rate</u>

<u>constant at 0 mV (Figure 6B), the fast transition (γ_o) becomes faster for openings in</u>

<u>the presence of larger ions, although the change is not very large. In contrast, the</u>

<u>intermediate opening rate (β_o) clearly increase with the radius of the ion. The slow</u>

<u>opening transition (α_o) shows a dependence on the ionic radius that is rather</u>

<u>interesting, being fast for small and large ions and with a minimum for intermediate</u>

<u>size ions.</u> This slow transition mirrors the dependence of the open probability on the

ionic radius (Figure 3B). ~~In general, rate constants, when compared at 0 mV, are faster~~

~~with the larger ions such as Cs⁺ and slower with smaller ions, although this is not a~~

~~rule.~~

The ~~interplay between voltage and the effect of the permeant cation can be quantified~~

~~by the~~ total charge associated with each of the three transitions is given by: <u>$Z_i =$</u>

<u>$(z_i^o - z_i^c)$,</u> where the z_i's are the partial charges associated with each rate constant and

i = α, β, γ and o and c represent the opening or closing rate constant, respectively.

<u>There is no difference between the values in the different permeant ion conditions</u>

<u>(Data not shown).</u> ~~It is readily apparent that the smaller ions Li⁺ and Na⁺ are~~

~~associated with a larger Z than say, Cs⁺ or Rb⁺.~~

~~The finding that these transitions, which occur near the open state, are voltage~~

~~dependent with a relatively high associated valence, that depends on the size of the~~

~~ion suggest that the observed voltage-dependence of the macroscopic conductance~~

~~might arise in part from the movement of ions in the pore's electric field. This is~~

~~consistent with our previous observation that the macroscopic voltage-dependence at~~

~~a saturating capsaicin concentration is modulated by the permeant ion (Figure 1D),~~

~~with a tendency for smaller apparent charge for larger ions.~~ To compare the observed

voltage- dependence of rate constants with the macroscopic conductance, we

measured conductance at the same concentration of capsaicin at which rate constants

were estimated (100 nM) <u>from single-channel recordings</u> and contrasted it to the

prediction of Scheme I, using the measured rates and charges.

The open probability predicted by Scheme I is given by <u>(derivation in Appendix):</u>

$P_O = \dfrac{K_s}{1 + K_s + \frac{K_s}{K_m} + \frac{K_s}{K_f}}$ Eq. 6

With equilibrium constants K_i given by:

$K_i = \dfrac{i_o(0)e^{\frac{q_{i_oV}}{KT}}}{i_c(0)e^{\frac{q_{i_cV}}{KT}}}$, where i = α, β, γ <u>and α=s, β=m, γ=f</u>   Eq. 7

Figures 7A and B depict macroscopic currents measured at saturating (10 µM) and

subsaturating (100 nM) capsaicin. Figure 7C shows G-V curves at 100 nM capsaicin for

all ions and Figure 7D shows the prediction of Scheme I (Eq. 6 and 7), using values

derived from measurements in Figure 6 (Table I). <u>For all ions except ammonium</u>, the

model shows good agreement with the data and suggest that the estimated apparent

charge associated with channel activation is modestly modulated by the permeant ion

(Figure 7E).

Discussion

TRPV1 is generally characterized as a cation permeable, non-selective channel with
slightly higher permeability to Ca^{2+} ions (Caterina et al., 1997b). Surprisingly, little is
known of the permeation mechanisms for cations in TRPV1. A study by (Samways and
Egan, 2011) showed single-channel recordings at a negative voltages that suggested
non-selective permeability of cations (also based in single-channel conductance) in
the order: Na^+ > Cs^+ > Li^+ > K^+. In this study, we have employed single-channel
recordings under bi-ionic conditions and measured a permeability sequence based on
the conductance at positive voltages, where the single-channel conductance is higher
and easier to measure. We observed a sequence Cs^+ > Rb^+ > K^+ > Na^+ > Li^+ <u>for alkali</u>
<u>cations</u>. Interestingly, this is identical with an Eisenman sequence of type I (Eisenman,
1962), which is produced by a weak field-strength cation binding site that favors
permeation of easily dehydrated larger ions. Molecular dynamics (MD) simulations
show that the TRPV1 pore may contain up to four binding sites for monovalent cation
permeation in which the free energy for Na^+ and K^+ is very similar (Jorgensen et al.,
2016), consistent with the almost identical permeability ratio for Na^+ and K^+.
However, no definitive permeation mechanism has been dilucidated for TRPV
channels.

Our data suggest that while non-selective, the pore of TRPV1 channels can regulate
the flux of monovalent cations based on the energetics of dehydration and can
differentiate between the size of ions, which is borne by the permeability sequence
obtained from single-channel records (Figure 2C).

MD simulations support a highly flexible selectivity filter in TRPV1 (Darré et al., 2015)
and our results indicated that gating can also be dynamic and is modulated by the
permeant ion. These results prompted us to explore the <u>microscopic gating kinetics</u>
<u>and its possible dependence on the permeant ion</u>. <u>We observe that single-channel</u>
<u>kinetics are indeed modulated by the permeant ion, with Rb^+ reducing the burst</u>
<u>length and the open probability of the channels. The smaller ions Li^+ and Na^+ make it</u>
<u>more probable for the channel to open in long duration bursts and Cs^+ has an</u>

intermediate effect. On top of these effects on closed and open dwell times, we
observed that the noise while the channel is open also varied with the permeant ion.

To study in detail this behavior, we employ the extended beta distribution method to
extract rate constants that arise from all closure events, including fast closures that
appear incomplete due to filtering, since simpler detection methods such as half-
amplitude threshold crossing are not appropriate to detect the contribution of rapid
incomplete closings.

In single-channel recording experiments, the intermediate transitions near the open
state seem to speed up when large ions permeate, leading to increased open channel
noise. This observation could be explained if small ions like Li$^+$ and Na$^+$ are able to
remain bound to the selectivity filter for longer times, due to their stronger
electrostatic electric field, and thus stabilize the open state, reducing fluctuations. In
contrast, the slow transitions are faster for small and big ions, with a minimum for
intermedia radii ions. This behavior is also observed in the burst length distributions
estimated from half-amplitude threshold crossing. Interestingly, the open probability
shows a similar behavior, with a minimum for Rb$^+$ ions. This is to be expected since
the slow transitions will contribute longer dwell times (in this case, longer closed
times) and have a larger contribution to the measured Po (Fig. 3).

The rapid fluctuations we observe are consistent with the existence of transitions
from the open state to closed states that are adjacent to the open state. It is likely that
these represent rapid pore dynamics derived from structural transitions. Cryo-EM
experiments have revealed that many TRPV channels, including TRPV1, can exist in
different pore conformations, even in the presence of activators (Zhang et al., 2021).
Functional data have also provided evidence that TRPV channels can adopt different
open conformations in the presence of different ligands (Benítez-Angeles et al., 2023;
Canul-Sánchez et al., 2018).

Experiments employing single-molecule FRET in Kir channels have directly shown
dynamic changes of the structure of the selectivity filter as a function of the presence

of Na$^+$ or K$^+$ ions (Wang et al., 2019). Related experiments employing solution nuclear

magnetic resonance have shown similar ion-dependent dynamics of the filter of a non-

selective ion channel, which has a filter structure more closely related to TRPV1's

(Lewis et al., 2021) and the intrinsic pore dynamics of TRPV1 has been demonstrated

directly by measuring the fluorescence of genetically encoded non-canonical amino

acids (Steinberg et al., 2017). It has also been demonstrated that the selectivity filter

can undergo conformational changes, although it is not a gate for small ions *per se*

(Jara-Oseguera et al., 2016).

An open question in TRPV channel biophysics, specially TRPV1, is the origin of the

mild voltage-dependence of the macroscopic conductance. As mentioned before, the

VLSD does not support a canonical function as a voltage sensor (Palovcak et al.,

2015b). It has recently been proposed that a conformational change that rearranges

the position of negatively charged residues at the extracellular entrance of the

selectivity filter is responsible for the voltage-dependence of TRPV1 gating, which is

equivalent to the displacement of ~0.55 elementary charges per channel (Yang et al.,

2020). We find that in the presence of saturating capsaicin, the apparent gating charge

depends on the identity of the permeant ion. The smallest alkali metal, Li$^+$, is

associated with a steeper G-V, with an apparent gating charge of 0.65 e$_o$ as well as

activation at more negative voltages.

It is known that other types of ion channels obtain their voltage-dependence from the

concerted movements of permeant ions in the permeation pathway (Schewe et al.,

2016; Marchesi et al., 2012; Pusch et al., 1995) and that permeant blockers, also

moving in the pore's electric field, can produce highly voltage-dependent block

(Spassova and Lu, 1998). For example, in chloride ClC channels, voltage-dependence is

a result of the interaction of Cl$^-$ ions with a protopore gate (Richard and Miller, 1990)

and in Kir channels, where permeating ions non-canonically regulate the voltage-

dependence of gating in these channels (Lu et al., 2001; Kurata et al., 2010). CNG

channels also display a small voltage-dependence at low open probabilities, although

its origin has not been clarified (Benndorf et al., 1999).

~~A fascinating observation derived from our study is that at relatively low~~
~~concentration of capsaicin, were the open probability $P_o \sim 0.4$, the apparent gating~~
~~charge is largest for the larger permeant cations, while at saturating concentration,~~
~~with $P_o = 0.98$, the apparent gating charge is largest when the smaller ion (Li^+)~~
~~permeates.~~

Acknowledgments

We thank Gisela Rangel, Manuel Hernández and Elsa Evaristo for excellent technical
support. This work was supported by DGAPA-PAPIIT-UNAM grant No. IN215621 to
L.D.I. and CONACyT (A1-S-8760) and DGAPA-PAPIIT-UNAM (IN200423) to T.R.
M. G-A is a doctoral student from Programa de Doctorado en Ciencias Bioquímicas-
UNAM and was supported by a doctoral thesis scholarship from CONACyT No.
776082.

References

Benítez-Angeles, M., A.E.L. Romero, I. Llorente, I. Hernández-Araiza, A. Vergara-Jaque, F.H. Real,
Ó.E. Gutiérrez Castañeda, M. Arciniega, L.E. Morales-Buenrostro, F. Torres-Quiroz, R.
García-Villegas, L.B. Tovar-Y-Romo, W.B. Liedtke, L.D. Islas, and T. Rosenbaum. 2023.
Modes of action of lysophospholipids as endogenous activators of the TRPV4 ion channel.
J. Physiol. 601:1655–1673. doi:10.1113/JP284262.

Benndorf, K., R. Koopmann, E. Eismann, and U.B. Kaupp. 1999. Gating by cyclic GMP and voltage in
the alpha subunit of the cyclic GMP-gated channel from rod photoreceptors. *J Gen Physiol*.
114:477–90.

Canul-Sánchez, J.A., I. Hernández-Araiza, E. Hernández-García, I. Llorente, S.L. Morales-Lázaro, L.D.
Islas, and T. Rosenbaum. 2018. Different agonists induce distinct single-channel
conductance states in TRPV1 channels. *J. Gen. Physiol.* 150:1735–1746.
doi:10.1085/jgp.201812141.

Cao, E. 2020. Structural mechanisms of transient receptor potential ion channels. *J. Gen. Physiol.*
152:e201811998. doi:10.1085/jgp.201811998.

Caterina, M.J., and D. Julius. 2001. The Vanilloid Receptor: A Molecular Gateway to the Pain
Pathway. *Annu. Rev. Neurosci.* 24:487–517. doi:10.1146/annurev.neuro.24.1.487.

Caterina, M.J., M.A. Schumacher, M. Tominaga, T.A. Rosen, J.D. Levine, and D. Julius. 1997a. The
capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature*. 389:816–
824. doi:10.1038/39807.

Caterina, M.J., M.A. Schumacher, M. Tominaga, T.A. Rosen, J.D. Levine, and D. Julius. 1997b. The
502   capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature*. 389:816–
503   824. doi:10.1038/39807.

504 Colquhoun, D., and B. Sakmann. 1985. Fast events in single-channel currents activated by
505   acetylcholine and its analogues at the frog muscle end-plate. *J. Physiol.* 369:501–557.
506   doi:10.1113/jphysiol.1985.sp015912.

507 Colquhoun, D., and F.J. Sigworth. 1995a. Fitting and Statistical Analysis of Single-Channel Records.
508   *In* Single-Channel Recording. B. Sakmann and E. Neher, editors. Springer US, Boston, MA.
509   483–587.

510 Colquhoun, D., and F.J. Sigworth. 1995b. Fitting and Statistical Analysis of Single-Channel Records.
511   *In* Single-Channel Recording. B. Sakmann and E. Neher, editors. Springer US, Boston, MA.
512   483–587.

513 Darré, L., S. Furini, and C. Domene. 2015. Permeation and dynamics of an open-activated TRPV1
514   channel. *J. Mol. Biol.* 427:537–549.

515 Deng, Z., N. Paknejad, G. Maksaev, M. Sala-Rabanal, C.G. Nichols, R.K. Hite, and P. Yuan. 2018.
516   Cryo-EM and X-ray structures of TRPV4 reveal insight into ion permeation and gating
517   mechanisms. *Nat. Struct. Mol. Biol.* 25:252–260. doi:10.1038/s41594-018-0037-5.

518 Eisenman, G. 1962. Cation selective glass electrodes and their mode of operation. *Biophys. J.*
519   2:259–323. doi:10.1016/s0006-3495(62)86959-8.

520 Fitzhugh, R. 1983. Statistical properties of the asymmetric random telegraph signal, with
521   applications to single-channel analysis. *Math. Biosci.* 64:75–89. doi:10.1016/0025-
522   5564(83)90028-7.

523 Geron, M., R. Kumar, W. Zhou, J.D. Faraldo-Gómez, V. Vásquez, and A. Priel. 2018. TRPV1 pore
524   turret dictates distinct DkTx and capsaicin gating. *Proc. Natl. Acad. Sci.* 115.
525   doi:10.1073/pnas.1809662115.

526 Grandl, J., S.E. Kim, V. Uzzell, B. Bursulaya, M. Petrus, M. Bandell, and A. Patapoutian. 2010.
527   Temperature-induced opening of TRPV1 ion channel is stabilized by the pore domain. *Nat.*
528   *Neurosci.* 13:708–714.

529 Hamill, O.P., A. Marty, E. Neher, B. Sakmann, and F.J. Sigworth. 1981. Improved patch-clamp
530   techniques for high-resolution current recording from cells and cell-free membrane
531   patches. *Pflugers Arch*. 391:85–100.

532 Heinemann, S.H., and F.J. Sigworth. 1991a. Open channel noise. VI. Analysis of amplitude
533   histograms to determine rapid kinetic parameters. *Biophys. J.* 60:577–587.
534   doi:10.1016/S0006-3495(91)82087-2.

535 Heinemann, S.H., and F.J. Sigworth. 1991b. Open channel noise. VI. Analysis of amplitude
536   histograms to determine rapid kinetic parameters. *Biophys. J.* 60:577–587.
537   doi:10.1016/S0006-3495(91)82087-2.

Hui, K., B. Liu, and F. Qin. 2003. Capsaicin activation of the pain receptor, VR1: multiple open
states from both partial and full binding. *Biophys. J.* 84:2957–2968. doi:10.1016/S0006-
3495(03)70022-8.

Jara-Oseguera, A., C. Bae, and K.J. Swartz. 2016. An external sodium ion binding site controls
allosteric gating in TRPV1 channels. *eLife*. 5:e13356. doi:10.7554/eLife.13356.

Jara-Oseguera, A., K.E. Huffer, and K.J. Swartz. 2019. The ion selectivity filter is not an activation
gate in TRPV1-3 channels. *Elife*. 8:e51212.

Jara-Oseguera, A., and K.J. Swartz. 2015. Unraveling Allosteric Coupling Mechanisms in the TRPV1
Channel. *Biophys. J.* 108:124a. doi:10.1016/j.bpj.2014.11.695.

Jordt, S.-E., M. Tominaga, and D. Julius. 2000. Acid potentiation of the capsaicin receptor
determined by a key extracellular site. *Proc. Natl. Acad. Sci.* 97:8134–8139.
doi:10.1073/pnas.100129497.

Jorgensen, C., S. Furini, and C. Domene. 2016. Energetics of Ion Permeation in an Open-Activated
TRPV1 Channel. *Biophys. J.* 111:1214–1222. doi:10.1016/j.bpj.2016.08.009.

Kurata, H.T., M. Rapedius, M.J. Kleinman, T. Baukrowitz, and C.G. Nichols. 2010. Voltage-
dependent gating in a "voltage sensor-less" ion channel. *PLoS Biol.* 8:e1000315.
doi:10.1371/journal.pbio.1000315.

Lewis, A., V. Kurauskas, M. Tonelli, and K. Henzler-Wildman. 2021. Ion-dependent structure,
dynamics, and allosteric coupling in a non-selective cation channel. *Nat. Commun.*
12:6225. doi:10.1038/s41467-021-26538-8.

Liao, M., E. Cao, D. Julius, and Y. Cheng. 2013. Structure of the TRPV1 ion channel determined by
electron cryo-microscopy. *Nature*. 504:107–112. doi:10.1038/nature12822.

Lu, T., L. Wu, J. Xiao, and J. Yang. 2001. Permeant ion-dependent changes in gating of Kir2.1
inward rectifier potassium channels. *J. Gen. Physiol.* 118:509–522.
doi:10.1085/jgp.118.5.509.

Marchesi, A., M. Mazzolini, and V. Torre. 2012. Gating of cyclic nucleotide-gated channels is
voltage dependent. *Nat. Commun.* 3:973. doi:10.1038/ncomms1972.

Matta, J.A., and G.P. Ahern. 2007. Voltage is a partial activator of rat thermosensitive TRP
channels: Voltage-independent gating of TRPV1 and TRPM8. *J. Physiol.* 585:469–482.
doi:10.1113/jphysiol.2007.144287.

Numazaki, M., T. Tominaga, K. Takeuchi, N. Murayama, H. Toyooka, and M. Tominaga. 2003.
Structural determinant of TRPV1 desensitization interacts with calmodulin. *Proc. Natl.*
Acad. Sci. U. S. A. 100:8002–8006. doi:10.1073/pnas.1337252100.

Palovcak, E., L. Delemotte, M.L. Klein, and V. Carnevale. 2015a. Comparative sequence analysis
suggests a conserved gating mechanism for TRP channels. *J. Gen. Physiol.* 146:37–50.

Palovcak, E., L. Delemotte, M.L. Klein, and V. Carnevale. 2015b. Comparative sequence analysis
suggests a conserved gating mechanism for TRP channels. *J. Gen. Physiol.* 146:37–50.

Pusch, M., U. Ludewig, A. Rehfeldt, and T.J. Jentsch. 1995. Gating of the voltage-dependent
chloride channel ClC-0 by the permeant anion. *Nature*. 373:527–531.
doi:10.1038/373527a0.

Rauh, O., U.-P. Hansen, S. Mach, A.J.W. Hartel, K.L. Shepard, G. Thiel, and I. Schroeder. 2017.
Extended beta distributions open the access to fast gating in bilayer experiments-assigning
the voltage-dependent gating to the selectivity filter. *FEBS Lett.* 591:3850–3860.
doi:10.1002/1873-3468.12898.

Rauh, O., U.P. Hansen, D.D. Scheub, G. Thiel, and I. Schroeder. 2018. Site-specific ion occupation in
the selectivity filter causes voltage-dependent gating in a viral K+ channel. *Sci. Rep.*
8:10406. doi:10.1038/s41598-018-28751-w.

Richard, E.A., and C. Miller. 1990. Steady-state coupling of ion-channel conformations to a
transmembrane ion gradient. *Science*. 247:1208–1210. doi:10.1126/science.2156338.

Rosenbaum, T., A. Gordon-Shaag, M. Munari, and S.E. Gordon. 2004. Ca2+/Calmodulin Modulates
TRPV1 Activation by Capsaicin. *J. Gen. Physiol.* 123:53–62. doi:10.1085/jgp.200308906.

Rosenbaum, T., S.L. Morales-Lázaro, and L.D. Islas. 2022. TRP channels: a journey towards a
molecular understanding of pain. *Nat. Rev. Neurosci.* 23:596–610. doi:10.1038/s41583-
022-00611-7.

Salazar, H., A. Jara-Oseguera, E. Hernandez-Garcia, I. Llorente, I.I. Arias-Olguin, M. Soriano-Garcia,
L.D. Islas, and T. Rosenbaum. 2009. Structural determinants of gating in the TRPV1
channel. *Nat. Struct. Mol. Biol.* 16:704–710.

Samways, D.S.K., and T.M. Egan. 2011. Calcium-dependent decrease in the single-channel
conductance of TRPV1. *Pflüg. Arch. - Eur. J. Physiol.* 462:681–691. doi:10.1007/s00424-
011-1013-7.

Schewe, M., E. Nematian-Ardestani, H. Sun, M. Musinszki, S. Cordeiro, G. Bucci, B.L. de Groot, S.J.
Tucker, M. Rapedius, and T. Baukrowitz. 2016. A Non-canonical Voltage-Sensing
Mechanism Controls Gating in K2P K+ Channels. *Cell*. 164:937–949.
doi:10.1016/j.cell.2016.02.002.

Schroeder, I. 2015. How to resolve microsecond current fluctuations in single ion channels: The
power of beta distributions. *Channels*. 9:262–280. doi:10.1080/19336950.2015.1083660.

Shimada, H., T. Kusakizako, T.H. Dung Nguyen, T. Nishizawa, T. Hino, M. Tominaga, and O. Nureki.
2020. The structure of lipid nanodisc-reconstituted TRPV3 reveals the gating mechanism.
Nat. Struct. Mol. Biol. 27:645–652. doi:10.1038/s41594-020-0439-z.

Sigworth, F.J., and S.M. Sine. 1987. Data transformations for improved display and fitting of single-
channel dwell time histograms. *Biophys. J.* 52:1047–1054. doi:10.1016/S0006-
3495(87)83298-8.

Spassova, M., and Z. Lu. 1998. Coupled Ion Movement Underlies Rectification in an Inward-
Rectifier K+ Channel. *J. Gen. Physiol.* 112:211–221. doi:10.1085/jgp.112.2.211.

Steinberg, X., M.A. Kasimova, D. Cabezas-Bratesco, J.D. Galpin, E. Ladron-de-Guevara, F. Villa, V.
Carnevale, L. Islas, C.A. Ahern, and S.E. Brauchi. 2017. Conformational dynamics in TRPV1
channels reported by an encoded coumarin amino acid. *eLife*. 6:e28626.
doi:10.7554/eLife.28626.

Wang, S., S.-J. Lee, G. Maksaev, X. Fang, C. Zuo, and C.G. Nichols. 2019. Potassium channel
selectivity filter dynamics revealed by single-molecule FRET. *Nat. Chem. Biol.* 15:377–383.
doi:10.1038/s41589-019-0240-7.

Yang, F., L. Xu, B.H. Lee, X. Xiao, V. Yarov-Yarovoy, and J. Zheng. 2020. An Unorthodox Mechanism
Underlying Voltage Sensitivity of TRPV1 Ion Channel. *Adv. Sci.* 7:2000575.
doi:10.1002/advs.202000575.

Yellen, G. 1984. Ionic permeation and blockade in Ca2+-activated K+ channels of bovine
chromaffin cells. *J. Gen. Physiol.* 84:157–186. doi:10.1085/jgp.84.2.157.

Yu, F.H., and W.A. Catterall. 2004. The VGL-Chanome: A Protein Superfamily Specialized for
Electrical Signaling and Ionic Homeostasis. *Sci. STKE*. 2004. doi:10.1126/stke.2532004re15.

Zhang, K., D. Julius, and Y. Cheng. 2021. Structural snapshots of TRPV1 reveal mechanism of
polymodal functionality. *Cell*. 184:5138-5150.e12. doi:10.1016/j.cell.2021.08.012.

Zubcevic, L., S. Le, H. Yang, and S.-Y. Lee. 2018. Conformational plasticity in the selectivity filter of
the TRPV2 ion channel. *Nat. Struct. Mol. Biol.* 25:405–415. doi:10.1038/s41594-018-0059-
z.

633
634
635 **Figure legends.**

636
637

Figure 1. The voltage-dependence of TRPV1 is influenced by the permeant ion.
A) Representative inside-out current-traces recorded in the same patch, elicited by 20
mV steps from -130 mV to 130 mV for a duration of 150 ms. The extracellular solution
(pipette) contained 130 mM NaCl, this patch was first exposed to internal 130 mM
NaCl (black) followed by exposure to 130 mM NaCl + 10 µM capsaicin (red) and finally
to 130 mM LiCl + 10 µM capsaicin (blue). All experiments were performed following
the same experimental strategy to compare the permeation of different monovalent
ions (Li^+, K^+, Rb^+, NH_4^+, Cs^+) with respect to the permeation of Na^+ in the same patch.
B) Averages of current-voltage (I-V) relations for all ions tested. The number of
experiments, n, is in parenthesis and is the same for panels C-E. **C)** Conductance-
voltage (G-V) relations for monovalent ions, obtained from data as in B in the presence

of 10 μM capsaicin, normalized to data with a symmetrical concentration of Na^+ from
each patch. G-V relations were fitted to equation 1 (continuous lines). **D)** Scatter plots
of the apparent charge z (above) and $V_{1/2}$ (below) associated with channel opening,
both were estimated by fitting G-V curves to equation 1. The values of z for Li^+ were
compared for statistical differences with all other ions, the p values are: *p = 0.00007
for Li^+ vs. Na^+; NS (not significant) p=0.213 for Li^+ vs. K^+; p = 0.00034 for Li^+ vs. Rb^+; p
= 0.00005 for Li^+ vs. NH_4^+; P = 0.01 for Li^+ vs. Cs^+. For $V_{1/2}$: p = 0.0000009 for Li^+ vs. Na^+;
p = 0.00001 for Li^+ vs. K^+; p = 0.00000001 for Li^+ vs. Rb^+; P = 0.000002 for Li^+ vs. NH_4^+;
p = 0.02 for Li^+ vs. Cs^+. **E)** Scatter plot of the $\Delta\Delta G$ associated to the ion effect in channel
opening was estimated as: $\Delta\Delta G = \Delta G_{Ion} - \Delta G_{Na^+}$. The * symbol indicates statistical
differences, p = 0.00005 for Li^+ vs. K^+; p = 0.0000002 for Li^+ vs. Rb^+; p = 0.000004 for
Li^+ vs. NH_4^+; p = 0.003 for Li^+ vs. Cs^+. Data are shown as mean ± SEM. The significant
difference was determined with a one-way ANOVA test followed by a Tukey´s test
post-hoc.
663
Figure 2. TRPV1 permeability to monovalent ions is size-dependent.
A) Representative current traces showing openings from single-channel recordings in
the presence of varying permeant ions. Inside-out patches were first exposed to
internal 130 mM Na^+ + 100 nM capsaicin and then to internal 130 mM (Li^+ (blue), Na^+
(red) K^+(mustard), Rb^+ (purple), NH_4^+ (black), Cs^+ (green)) + 100 nM capsaicin,
similarly to experiments in Figure 1A. The dashed line represents the closed state of
the channel. Currents are shown filtered at 2.5 kHz, for clarity of illustration. **B)** I-V
relations from single-channel records. These were obtained from data as in A for each
ion. **C)** Current carried by each ion relative to the single-channel current amplitude
with Na^+. All currents measured at 90 mV. Ionic radii in Å are also shown for
comparison. Data are mean ± s.e.m., n is shown in parentheses and is the same for the
curves in **B**.
676
Figure 3. Rubidium ions strongly modulate the single-channel open probability
of TRPV1 channels.
A) Representative current traces from single-channel recordings at 70 mV when
intracellular Rb^+ (purple) or Na^+ (red) were used as permeant ions. Currents are
shown at full bandwidth (5 kHz). **B)** Open probability of TRPV1 channels at 70 mV in
the presence of different permeant ions and 100 nM of capsaicin, calculated from
traces of current recordings as shown in A. Data are mean ± s.e.m. Number of
experiments is: Li^+(4); Na^+(8); K^+(3); Rb^+(5); NH_4^+(4); Cs^+(4).
685
Figure 4. The permeant ion modulates single channel kinetics of TRPV1
channels. A) Representative single-channel traces with the indicated permeant ion.
Data is shown at full bandwidth (5 kHz) except Li^+ currents, which are filtered at 2.5
kHz. **B)** Burst length distributions corresponding to permeant ions in A. The
continuous curve is the sum of exponential components. The values of time constants
(τ, in s) and amplitudes (a_i) are: Cs^+; a_1=1.5474, τ_1 = 0.0013, a_2=5.2727, τ_2=0.04818,
a_3=4.3972, τ_3=0.5031. Na^+; a_1=0.3720, τ_1=0.0362, a_2=3.2643, τ_2=0.8346. Li^+;

a_1=2.9447, τ_1=0.0048, a_2=8.0241, τ_2=0.2362. Rb$^+$; a_1=5.0377, τ_1=7.244 x 10^{-5},

a_2=8.7838, τ_2=0.0021, a_3=8.4035, τ_3=0.0163, a_4=2.8719, τ_4=0.2094.

Figure 5. Fluctuations in the single-channel current during the open state can be

fitted by a Markov model.

A) Representative current trace during a typical single TRPV1 opening burst recorded

in the inside-out configuration at 90 mV, with internal 130 mM RbCl + 100 nM

capsaicin. **B)** Normalized all-points histograms showing single-channel current

amplitudes of openings elicited by 100 nM capsaicin at 90 mV, for each ion tested. The

histograms are shown aligned to the closed state amplitude peak. **C)** All-points

histograms aligned to the maximum current amplitude of the open state to highlight

the asymmetric form of the histograms. **D) The same histograms in C are shown**

normalized both in number of counts and current amplitude. This representation

emphasizes the ion-dependence of the open current asymmetry. **E)** Current trace of a

single-channel at 90 mV from an inside-out patch exposed to internal 130 mM NH$_4$Cl +

100 nM capsaicin (black) and simulation with the model in Scheme I (blue). These

simulated recordings are used to produce simulated all-points amplitude histograms

of single-channel current, which are then fitted to experimental all-points amplitude

histograms. **F)** Typical fit of a simulated histogram (blue) to experimental histogram

elicited by 110 mV with internal 130 mM NH$_4$Cl + 100 nM capsaicin (black) using the

Markov model in Scheme I. Parameters obtained from the fit are: α_o = 1500 s^{-1}; α_c = 27

s^{-1}; β_o = 22500 s^{-1}; β_c = 600 s^{-1}; γ_o = 127500 s^{-1}; γ_c = 30000 s^{-1}; χ^2 = 0.00013873.

Figure 6. Voltage-dependent transitions near the open state of TRPV1 channels

are modulated by the permeant ions.

A) Rate constants obtained from the Markov model fit to the histograms as in Figure

5F. The rate constants were classified as slow, medium, and fast according to their

magnitude at 0 mV. The continuous curves represent the fit to Eq. 5.

B) Plot of the value of the rate constants at zero mV obtained from the fit of equation 5

to rate constants in A. Data are mean ± s.e.m. n is: 3 (Li$^+$); 6 (Na$^+$); 2(K$^+$); 3 (Rb$^+$); 3

(NH$_4$$^+$) and 3 (Cs$^+$).

Figure 7. The opening probability predicted by the Markov partially model

explains the macroscopic experimental behavior.

A) Representative inside-out macroscopic current traces recorded in the same patch,

elicited by 20 mV steps from -130 mV to +150 mV for 150 ms. The patch was exposed

to internal 130 mM NaCl + 10 μM capsaicin (red) followed by 130 mM RbCl + 10 μM

capsaicin (purple). **B)** Representative inside-out current traces like A but using 100

nM capsaicin. **C)** Normalized conductance-voltage (G-V) relations with monovalent

ions, obtained from inside-out patches after treatment with internal 100 nM capsaicin.

The G-V relation was fitted to the Boltzmann function (line). Data are shown as mean

± s.e.m., number of experiments n, in parenthesis. **D)** Normalized prediction of open

probability by a Markov model for each monovalent ion. **E)** Bar graph of apparent

charge (q_{app})associated with channel opening estimated by fitting a Boltzmann

equation to curves in A and B. Data is shown as mean ± s.e.m.

739
740 **Supplementary figure legends.**
741
742
756
Supplementary Figure 3. Amplitude histogram fits to three different Markov
models. A) Markov models with one closed state (green), two closed states (blue) and
three closed states (red). **B)** Fit of the simulated histograms from the models in A to
experimental an all-points histogram of single channel current of TRPV1(black),
elicited by a pulse to 90 mV from an inside-out patches exposed to 130 mM NaCl + 100
nM capsaicin. The fits parameters used for one close state were: α_o = 301.9535 s^{-1}; α_c
= 36.1664 s^{-1}; χ^2 = 0.0697; two closed states: α_o = 300 s^{-1}; α_c = 37.5 s^{-1}; γ_o = 28500 s^{-1};
γ_c = 13500 s^{-1}; χ^2 = 0.002342; three closed states: α_o = 280.61 s^{-1}; α_c = 49 s^{-1}; β_o =
14724.475 s^{-1}; β_c = 1133.16 s^{-1}; γ_o = 52500 s^{-1}; γ_c = 15396 s^{-1}; χ^2 = 0.00017296.
766
767
768 **Appendix: Derivation of equation 6**
769 The equations describing the occupancy of each state in Scheme I are given by

$$\frac{dC_s}{dt} = \alpha_c O - \alpha_o C_s$$
770 $$\frac{dC_m}{dt} = \beta_c O - \beta_o C_m$$
771 $$\frac{dC_f}{dt} = \gamma_c O - \gamma_o C_f$$
772 $$\frac{dO}{dt} = \alpha_o C_s + \beta_o C_m + \gamma_o C_f - O(\alpha_c + \beta_c + \gamma_c)$$
773
774 These equations are subject to: $C_s + C_m + C_f + O = 1$ and the probability of being in the
775 open state, P_o is:

$$P_O = \frac{O}{C_s + C_m + C_f + O} = \frac{1}{1 + \frac{C_s}{O} + \frac{C_m}{O} + \frac{C_f}{O}}$$

In steady-state:

$0 = \begin{cases} \dfrac{dC_s}{dt} = \alpha_c O - \alpha_o C_s \\[2mm] \dfrac{dC_m}{dt} = \beta_c O - \beta_o C_m \\[2mm] \dfrac{dC_f}{dt} = \gamma_c O - \gamma_o C_f \\[2mm] \dfrac{dO}{dt} = \alpha_o C_s + \beta_o C_m + \gamma_o C_f - O(\alpha_c + \beta_c + \gamma_c) \end{cases}$

Therefore: $\dfrac{C_s}{O} = \dfrac{\alpha_c}{\alpha_o}; \dfrac{C_m}{O} = \dfrac{\beta_c}{\beta_o}; \dfrac{C_f}{O} = \dfrac{\gamma_c}{\gamma_o}$ and $P_O = \dfrac{1}{1 + \frac{\alpha_c}{\alpha_o} + \frac{\beta_c}{\beta_o} + \frac{\gamma_c}{\gamma_o}}$

$P_O = \dfrac{1}{1 + \frac{\alpha_c}{\alpha_o} + \frac{\beta_c}{\beta_o} + \frac{\gamma_c}{\gamma_o}} \cdot \left(\dfrac{\frac{\alpha_o}{\alpha_c}}{\frac{\alpha_o}{\alpha_c}}\right) = \dfrac{\frac{\alpha_o}{\alpha_c}}{1 + \frac{\alpha_o}{\alpha_c} + \frac{\alpha_o}{\alpha_c}\frac{\beta_c}{\beta_o} + \frac{\alpha_o}{\alpha_c}\frac{\gamma_c}{\gamma_o}}$. Defining $\dfrac{i_o}{i_c} = K_i \; for \; i = \alpha, \beta \; or \; \gamma$, we can

write:

$P_O = \dfrac{K_s}{1 + K_s + \frac{K_s}{K_m} + \frac{K_s}{K_f}}$, where $\alpha = s$ stands for slow, $\beta = m$ for medium and $\gamma = f$ for fast.

783
784
785

Figure 1

A

Li⁺ Na⁺ K⁺ Rb⁺ NH₄⁺ Cs⁺

10 pA
40 ms

110 mV C

50 mV C

-50 mV C

B

Cs⁺
NH₄⁺
Rb⁺
K⁺
Na⁺
Li⁺

Current (pA)

Membrane potential, (mV)

C

Ion	I_I/I_{Na^+}	Å
Cs⁺	1.58 ± 0.15 (4)	1.69
NH₄⁺	1.55 ± 0.19 (3)	1.48
Rb⁺	1.28 ± 0.08 (5)	1.48
K⁺	1.23 ± 0.20 (4)	1.33
Na⁺	1	0.96
Li⁺	0.48 ± 0.05 (4)	0.6

Figure 2

Figure 3

Figure 4

A

5 pA
50 ms

B

Normalized counts
1.0
0.5
0.0
0   5   10   15   20
Current, pA

C

Open
Cs⁺
NH₄⁺
Rb⁺
K⁺
Na⁺
Li⁺
12 14 16 18 20 22 24
Current, pA

D

Normalized counts
1
0.1
0.01
0   1
Normalized current

E

Simulated

Experimental

5 pA
40 ms

F

Counts
10^4
10^3
10^2
10^1
0   5   10   15   20
Current, pA

Figure 5

A

Slow
α_o

Medium
β_o

Fast
γ_o

α_c

β_c

γ_c

B

Figure 6

Figure 7

Permeant ion	Open time		Closed time	
	Time constant (s)	Amplitude	Time constant (s)	Amplitude
Li^+	$\tau_1 = 0.0002$ $\tau_2 = 0.004$ $\tau_3 = 0.02$	$a_1 = 2.3$ $a_2 = 3.4$ $a_3 = 4.8$	$\tau_1 = 0.0001$ $\tau_2 = 0.0005$ $\tau_3 = 0.005$ $\tau_4 = 0.06$	$a_1 = 7.5$ $a_2 = 4.2$ $a_3 = 2.5$ $a_4 = 1.1$
Na^+	$\tau_1 = 0.0001$ $\tau_2 = 0.0015$ $\tau_3 = 0.011$	$a_1 = 11.6$ $a_2 = 2.3$ $a_3 = 16.6$	$\tau_1 = 6.7 \times 10^{-5}$ $\tau_2 = 0.0005$ $\tau_3 = 0.0033$ $\tau_4 = 0.0305$	$a_1 = 21.4$ $a_2 = 11.8$ $a_3 = 3.7$ $a_4 = 0.02$
Rb^+	$\tau_1 = 8.71 \times 10^{-5}$ $\tau_2 = 0.0011$ $\tau_3 = 0.003$	$a_1 = 13.1$ $a_2 = 12.5$ $a_3 = 21.5$	$\tau_1 = 6.6 \times 10^{-5}$ $\tau_2 = 0.0007$ $\tau_3 = 0.007$ $\tau_4 = 0.21$	$a_1 = 27.1$ $a_2 = 15.9$ $a_3 = 10.2$ $a_4 = 4.1$
Cs^+	$\tau_1 = 0.0001$ $\tau_2 = 0.0071$ $\tau_3 = 0.0211$	$a_1 = 16.5$ $a_2 = 25$ $a_3 = 5.6$	$\tau_1 = 8.5 \times 10^{-5}$ $\tau_2 = 0.0009$ $\tau_3 = 0.0038$ $\tau_4 = 0.023$	$a_1 = 31.5$ $a_2 = 15.5$ $a_3 = 6.2$ $a_4 = 2$

Table 1. Time constants and amplitudes of individual exponential components fitted to the histograms in supplementary figure 1.

Ion	Rate constants	Rate constant at 0 mV (s^{-1})	$Z_1(e_0)$
Li$^+$	α_o	1422.3	0.02548
	α_c	198.6	-0.36236
	β_o	2042.7	0.4883
	β_c	60.337	0.6112
	γ_o	87664.2	0.028629
	γ_c	34744	-0.051124
Na$^+$	α_o	875.09	-0.13178
	α_c	163.28	-0.55262
	β_o	6302.1	0.15402
	β_c	191.277	0.30446
	γ_o	83222	0.021433
	γ_c	28800.5	-0.02206
K$^+$	α_o	291.38	-0.20251
	α_c	357.04	-0.5549
	β_o	1000.6	0.312546
	β_c	22.5	0.4867
	γ_o	62298.3	0.10147
	γ_c	28322	-0.01003
Rb$^+$	α_o	123.32	-0.23134
	α_c	213.72	-0.44403
	β_o	8294.0	0.042546
	β_c	210.2	0.24134
	γ_o	78027.2	0.044316
	γ_c	36391	-0.11627
NH$_4^+$	α_o	236.23	0.31361
	α_c	596.9	-0.50654
	β_o	3077.9	0.32154
	β_c	33.656	0.582
	γ_o	72723.3	0.08284
	γ_c	24770.1	-0.034484
Cs$^+$	α_o	1543.1	0.014812
	α_c	231.63	-0.4304
	β_o	21545.9	-0.055021
	β_c	521.0	0.21361
	γ_o	97846	0.037125
	γ_c	34519	0.011

Table 2. Parameters of the fits to equation 5 of the data in Figure 6.

Supplementary Figure 2

A

B

Supplementary figure 3