Evolution of the spider homeobox gene repertoire by tandem and whole genome duplication

Madeleine E. Aase-Remedios[1], Ralf Janssen[2]*, Daniel J. Leite[1], Lauren Sumner-Rooney[3], and Alistair P. McGregor[1]*

[1] Department of Biosciences, Durham University, Durham, DH1 3LE, UK.
[2] Department of Earth Sciences, Uppsala University, 752 36, Uppsala, Sweden.
[3] Museum für Naturkunde, Leibniz-Institut für Evolutions- und Biodiversitätsforschung, 10115 Berlin Germany.

*Corresponding authors: alistair.mcgregor@durham.ac.uk, ralf.janssen@geo.uu.se

ABSTRACT

Gene duplication generates new genetic material that can contribute to the evolution of

gene regulatory networks and phenotypes. Duplicated genes can undergo

subfunctionalisation to partition ancestral functions and/or neofunctionalisation to

assume a new function. We previously found there had been a whole genome

duplication (WGD) in an ancestor of arachnopulmonates, the lineage including spiders

and scorpions but excluding other arachnids like mites, ticks, and harvestmen. This

WGD was evidenced by many duplicated homeobox genes, including two Hox clusters,

in spiders. However, it was unclear which homeobox paralogues originated by WGD

versus smaller-scale events such as tandem duplications. Understanding this is key to

determining the contribution of the WGD to arachnopulmonate genome evolution. Here

we characterised the distribution of duplicated homeobox genes across eight

chromosome-level spider genomes. We found that most duplicated homeobox genes in

spiders are consistent with an origin by WGD. We also found two copies of conserved

homeobox gene clusters, including the Hox, NK, HRO, *Irx*, and SINE clusters, in all

eight species. Consistently, we observed one copy of each cluster was degenerated in

terms of gene content and organisation while the other remained more intact. Focussing

on the NK cluster, we found evidence for regulatory subfunctionalisation between the

duplicated NK genes in the spider *Parasteatoda tepidariorum* compared to their single-

copy orthologues in the harvestman *Phalangium opilio*. Our study provides new insights

into the relative contributions of multiple modes of duplication to the homeobox gene

repertoire during the evolution of spiders and the function of NK genes.

INTRODUCTION

Whole genome duplications (WGDs) have occurred several times during animal evolution (Ohno 1970; Putnam et al. 2008; Flot et al. 2013; Kenny et al. 2016; Schwager et al. 2017; Nong et al. 2021). Duplicated genes are thought to be released from selective constraints, allowing paralogues to subfunctionalise, specialise, or neofunctionalise either by changes to their cis-regulatory controls or protein sequence while collectively maintaining the robustness of the ancestral functionality (Force et al. 1999; Jiménez-Delgado et al. 2009; Tinti et al. 2012; Espinosa-Cantú et al. 2015; Marlétaz et al. 2018). The two rounds of WGD (2R WGD) in the gnathostome ancestor has been credited with facilitating the evolution of vertebrate novelties, however, the extent to which WGDs in other animal lineages have contributed to diversification is not well understood (Meyer and Schartl 1999; Shimeld and Holland 2000; Cañestro et al. 2013; Simakov et al. 2020; Aase-Remedios and Ferrier 2021).

Many retained ohnologues (genes duplicated via WGD) are transcription factors that regulate processes in development, which further implicates the role of duplication in the evolution of phenotypes and the underlying developmental regulatory networks (Brunet et al. 2017). Homeobox genes are a superclass of transcription factors that expanded by tandem duplication early in animal evolution into eleven recognised classes encompassing more than 100 bilaterian gene families with diverse and wide-reaching roles in development (Table 1) (Holland et al. 2007; Larroux et al. 2008; Zhong et al. 2008; Kumar 2009; Butts et al. 2010; Bürglin 2011; Zhong and Holland 2011; Holland 2013; Holland et al. 2017). Certain homeobox gene expansions may have coincided with innovations that occurred at the evolutionary origins of animals and subsequently bilaterians (Garcia-Fernàndez 2005; Holland 2013; Holland 2015). Tandem duplication of homeobox genes early in animal evolution resulted in clusters of related genes that have been conserved across animals (Pollard and Holland 2000; Larroux et al. 2007; Larroux et al. 2008; Ferrier 2016a). Homeobox gene clusters (Table 1) are thought to be maintained by functional constraint on their genomic organisation

exerted by complex and overlapping regulatory elements controlling the expression of

adjacent genes (Irimia et al. 2012; Ferrier 2016b).

Table 1 – Genes and *Drosophila melanogaster* synonyms in conserved bilaterian homeobox
clusters. Bilaterian homeobox gene **families** contain genes descendant from a single inferred gene in the
ancestor of all bilaterians, as previously defined by Ferrier (2016b) and the references therein. Families are
grouped into **classes,** which themselves originated by duplication earlier in animal evolution. **Clusters** are
composed of related genes that arose by tandem duplication and have been conserved across bilaterians.
The Hox cluster contains the Hox genes, while the ancestral bilaterian Super-Hox cluster includes the
related ANTP-class genes that are often found linked to the Hox cluster (Butts et al. 2010; Ferrier 2016b).
The ParaHox cluster is the sister of the Hox cluster and consists of three bilaterian families; dashes for
Xlox/Pdx indicate it is not found in arthropods (Brooke et al. 1998; Ferrier 2016a). The NK cluster also
belongs to the ANTP class. Some ANTP-class NK2 cluster genes were first described as part of a
'pharyngeal' cluster, containing *NK2.1*, *NK2.2*, and *Msxlx*, as well as two non-homeobox genes, *Pax1/9*
and *FoxA1/2* (Simakov and Kawashima 2017); linkage and sometimes clustering between *NK2.1*, *NK2.2*,
and *Msxlx* is evident across bilaterians, and here is referred to as the NK2 cluster (Simakov et al. 2015;
Ferrier 2016b). The PRD-class HRO cluster (Mazza et al. 2010) also includes the LIM-class gene *Isl* as
they are often found linked (Ferrier (2016b). The TALE-class *Irx* cluster likely underwent independent
duplications in different lineages; *ara* and *caup* are a fly-specific duplication of one *Irx* family, often clustered
with single arthropod *mirr* orthologues of a second *Irx* family (Irimia et al. 2008; Kerner et al. 2009). The
dash for *Irx3* indicates it is not found in *D. melanogaster*. The SINE-class cluster (Gallardo et al. (1999)
consists of three families and is conserved across bilaterians (Ferrier 2016b).

Cluster (Class)	Bilaterian family	Arthropod family	Synonyms	Cluster (Class)	Bilaterian family	Arthropod family	Synonyms
Super-Hox (ANTP)	*Hox1*	*lab*	*lab*	NK (ANTP)	*NK1*	*NK1*	*slou*
	Hox2	*pb*	*pb*		*NK3*	*NK3*	*bap*
	Hox3	*Hox3*	*zen, zen2, bcd*		*NK4*	*NK4*	*tin*
	Hox4	*Dfd*	*Dfd*		*NK5*	*NK5*	*Hmx*
	Hox5	*Scr*	*Scr*		*NK6*	*NK6*	*Hgtx*
	Hox6-8	*ftz*	*ftz*		*NK7*	*NK7*	*NK7.1*
		Antp	*Antp*		*Msx*	*Msx*	*Dr, msh*
		Ubx	*Ubx*		*Tlx*	*Tlx*	*C15, cll*
		abd-A	*abd-A*		*Lbx*	*Lbx*	*lbe, lbl*
	Hox9-15	*Abd-B*	*Abd-B*		*Hhex*	*Hhex*	*HHEX*
	Abox	*Abox*	*CG34031*		*Noto*	*Noto*	*Not*
	Dbx	*Dbx*	*Dbx*		*Emx*	*Emx*	*ems, E5*
	Dlx	*Dlx*	*Dll*	NK2 (ANTP)	*NK2.1*	*NK2.1*	*scro*
	En	*En*	*en, inv*		*NK2.2*	*NK2.2*	*vnd*
	Evx	*Evx*	*eve*		*Msxlx*	*Msxlx*	*CG15696*
	Gbx	*Gbx*	*unpg*		*Hlx*	*Hlx*	*H2.0*
	Meox	*Meox*	*btn*	IRX (TALE)	*Irx*	*Irx1*	*ara, caup*
	Mnx	*Mnx*	*exex*			*Irx2*	*mirr*
	Nedx	*Nedx*	*lms*			*Irx3*	*–*
	Ro	*Ro*	*ro*	HRO+Isl (PRD + LIM)	*Hbn*	*Hbn*	*hbn*
ParaHox (ANTP)	*Xlox/Pdx*	*–*	*–*		*Rax*	*Rax*	*Rx*
	Gsx	*Gsx*	*ind*		*Otp*	*Otp*	*otp*
	Cdx	*Cdx*	*cad*		*Isl*	*Isl*	*tup*
				SINE (SINE)	*Six1/2*	*Six1/2*	*so*
					Six3/6	*Six3/6*	*optix*
					Six4/5	*Six4/5*	*Six4*

The best known homeobox gene cluster, the Hox cluster, illustrates how the genomic

organisation of homeobox genes is important for their function. Most arthropod

genomes contain one Hox cluster consisting of ten ANTP-class genes (Table 1), which

are involved in patterning the antero-posterior axis of animals in the same relative order

as the genomic organisation of the genes within the cluster (Krumlauf 2018 and the

references therein). The Hox cluster is highly conserved across bilaterians, both in

genomic organisation and function, though differences in Hox gene expression are

known to underlie changes to animal body plans (Averof and Patel 1997; Abzhanov et

al. 1999; Janssen et al. 2014; Martin et al. 2016; Serano et al. 2016; Janssen and

Pechmann 2023). Hox clusters have also been retained in duplicate following WGDs, as

exhibited by the four vertebrate Hox clusters that arose via the 2R WGD (Holland and

Garcia-Fernàndez 1996; McLysaght et al. 2002; Holland et al. 2007; Putnam et al.

2008; Cañestro et al. 2013; Holland 2013; Holland and Ocampo Daza 2018). The 2R

WGD appears to have relaxed constraint on vertebrate Hox cluster organisation so that

while the complement of genes from *Hox1* to *Hox14* are present in at least one copy, no

cluster contains all 14 genes (Hoegg and Meyer 2005). It is still not well understood to

what extent the patterns of gene loss between duplicate clusters and the

subfunctionalisation observed for vertebrate homeobox gene clusters apply more widely

to other homeobox gene clusters (Table 1) and to other animal lineages that have

undergone WGD.

Besides gnathostomes, an independent WGD took place in the arachnopulmonate

ancestor approximately 450 MYA (Figure 1), resulting in nearly 60% of homeobox

genes retained in duplicate in spiders and scorpions and their relatives, including two

Hox clusters (Schwager et al. 2017; Leite et al. 2018; Harper et al. 2021). Within each

Hox gene family, one ohnologue exhibits different spatial and temporal expression

compared to the other, though overall, spatial and temporal collinearity is largely intact

for each cluster (Schwager et al. 2007; Schwager et al. 2017; Turetzek et al. 2022).

Several of the other retained homeobox genes exhibit divergent expression consistent

with both sub- and neofunctionalisation in *Parasteatoda tepidariorum*, showing that

duplication of these genes has played a role in the evolution of spider development

(Leite et al. 2018).

The lack of chromosomal-level genome assemblies for spiders has so far hindered a full

understanding of the relationships among duplicated homeobox genes and the relative

impact of the WGD

compared to tandem

duplication. It was not

possible to determine

whether clusters were fully

intact or had undergone

degradation, beyond one

of the Hox clusters

(Schwager et al. 2017).

Leite et al. (2018)

described clustering of a

few genes, each

belonging to several of the

other homeobox gene

clusters besides the Hox,

including fragments of the

NK, HRO, *Irx*, and SINE

clusters, some of which

were found in duplicate.

However, it remained

impossible to determine

whether these duplicates

were ohnologues,

indicating the retention of

ohnologous clusters, or if

they resulted from tandem

Figure 1 – Cladogram of arthropods used in this study. The red star indicates the timing of the whole genome duplication (WGD); pre-duplicate lineages have grey lines, duplicate lineages have black lines. Genome assembly level is indicated below the species binomial, in bold for the chromosome-level assemblies. Higher phylogenetic classifications are labelled on phylogeny nodes. The topology was drawn based on published molecular phylogenies (Ontano et al. 2021; Li et al. 2022). Silhouettes were taken from the PhyloPic database [https://www.phylopic.org/].

duplications occurring before or after the WGD. Furthermore, it could not be determined

if these duplicates were shared with other spider species because of the lack of

comparative genomic resources.

We surveyed the homeobox gene complement and synteny across eight recently

released chromosome-level genome assemblies available for spiders including *P.*

tepidariorum (Figure 1, see Materials and Methods). Our analysis allowed us to

distinguish WGD ohnologues from paralogues that arose via tandem duplication for the

first time among spiders. We found that many spider homeobox genes and clusters

were retained in duplicate following the WGD, within conserved patterns of whole-

genome synteny. We also characterised both ancient and recent tandem duplications

and determined the timing of these relative to the WGD and the divergence of major

lineages within arthropods. We found that the NK cluster was retained in duplicate and

largely conserved across spiders. Complementing the previous analysis of *Emx* and

Msx family genes (Leite et al. 2018), we described the expression of the rest of the NK

cluster genes during embryogenesis in the harvestman *P. opilio*, an outgroup to the

arachnopulmonates that did not undergo the WGD, and their orthologues and

ohnologues from the spider *P. tepidariorum*.

For these NK cluster genes, we detected no evidence of selection on coding sequences

in our comparisons between paralogues or between single copy *Phalangium opilio*

genes and their *P. tepidariorum* ohnologues, but for each family with retained

ohnologues in *P. tepidariorum*, we found distinct expression patterns for each

ohnologue during embryogenesis, providing evidence for regulatory

subfunctionalisation. Combined, these results provide new insight into both the patterns

of ohnologue retention and their consequences in spider evolution.

RESULTS

Homeobox gene repertoires in arthropods

We surveyed the homeobox genes of eight spider species with chromosome-level
genome assemblies, representing seven entelegyne and one synspermiatan species
and compared these to the scorpion *Centruroides sculpturatus*, the harvestman
Phalangium opilio, and the tick *Ixodes scapularis*, as well as three mandibulates
(Figures 1, 2 and Table 2). We found that spiders and scorpions have similar numbers
of homeobox genes, and on average, have 1.4 times more homeobox genes than
outgroup species that did not have an ancestral WGD (Table 2). The spiders and
scorpion had an average of 150 homeobox genes; the most are found in *Hylyphantes*
graminicola with 172 and the fewest in *Dysdera silvatica* with 140. Mandibulates and
non-arachnopulmonate arachnids had on average 107 homeobox genes, with the
fewest in the insects (103) and the most in *Strigamia maritima* (113). Previous surveys
found 96 and 69 homeobox genes in *I. scapularis* and *P. opilio*, respectively,145 in *P.*
tepidariorum, and 156 in *C. sculpturatus* (Schwager et al. 2017; Leite et al. 2018), but
we identified between 1-55% more in each of these species (Table 2). For *P. opilio*,
searching both the recent genome assembly (Gainett et al. 2021) as well as the
developmental transcriptome (Sharma et al. 2012) provided a more complete dataset
and enabled the identification of more homeobox genes than previously described.

In total, 90 described homeobox families were represented in our arthropod species set
(Table 2), though some were absent in different species, such that no species had a
gene from every family (Figure 2 A, Table S1). In addition, we also found several new
homeobox genes. A few of these may represent new lineage-specific unnamed ("un")
families, temporarily called *ANTP-un1*, *PRD-un1*, *SINE-un1*, and *TALE-un1* based on
the classes to which they likely belong. *ANTP-un1* is found only in the chelicerates,
while *PRD-un1* and *SINE-un1* were found only in the araneids, and *TALE-un1* only in
the *Trichonephila* spp. (Figure 2 A, Table S1). The '1' is simply a placeholder to indicate
that these genes have an orthologue in at least two species and to distinguish them

from other unnamed or unclassified homeodomain sequences that have no similarity to

described families and were found only in one species, sometimes in several copies.

Table 2 – Homeobox genes across arthropods are largely stable, except for an increase in the
arachnopulmonates. Homeobox families: the number of described bilaterian homeobox gene families with
at least one gene present in a genome (Holland et al. 2007; Zhong et al. 2008; Zhong and Holland 2011;
Ferrier 2016b). Tandem duplications are divided into ancient (predating the WGD) and recent (after the
WGD). Species to the right of the vertical line had an ancestral WGD. Species abbreviations: *Dmel*
Drosophila melanogaster, *Tcas Tribolium castaneum*, *Smar Strigamia maritima*, *Isca Ixodes scapularis*,
Popi Phalangium opilio, *Cscu Centruroides sculpturatus*, *Dsil Dysdera silvatica*, *Dpla Dolomedes*
plantarius, *Lele Latrodectus elegans*, *Ptep Parasteatoda tepidariorum*, *Hgra Hylyphantes graminicola*, *Abru*
Argiope bruennichi, *Tant Trichonephila antipodiana*, *Tcla Trichonephila clavata*.

	Dmel	Tcas	Smar	Isca	Popi	Cscu	Dsil	Dpla	Lele	Ptep	Hgra	Abru	Tant	Tcla
Homeobox genes	103	104	114	110	108	161	143	150	152	151	176	149	147	152
Homeobox families	83	85	90	89	91	90	79	87	89	91	86	88	87	89
Families with paralogues from ancient TD(s)	5	5	6	7	8	7	7	8	8	8	8	8	8	8
Families with paralogues from recent TD(s)	9	9	13	8	5	4	8	3	6	4	23	3	2	5
Families with only one gene	71	71	74	76	81	41	35	41	46	44	42	45	44	45
Families with retained ohnologues	–	–	–	–	–	46	41	44	41	44	37	40	40	40

Comparing this taxonomic breadth of species in concert with genomic synteny enabled

us to distinguish ancient and recent tandem duplicates from likely ohnologues and

determine the timing of duplications relative to major radiations in arthropod evolution.

We have defined ancient tandem duplicates as paralogues that are often linked or

clustered in most spider genomes and date to a duplication node in an ancestor before

the arachnopulmonate WGD, while recent tandem duplicates are also linked or

clustered, but have occurred either after the WGD or in the outgroups. Six homeobox

gene families, *Arx*, *Barhl*, *Irx*, *Pax3/7*, *Pax4/6*, and *Lhx2/9*, underwent ancient tandem

duplication events that are shared by other arthropods and may have ancestry even

deeper in the bilaterian phylogeny (Figure 2 B). Only three of these ancient tandem

paralogues have been subsequently lost; just one *Lhx2/9* gene is found in *D.*

melanogaster, and only one *Barhl* gene is found in *T. castaneum* and *D. silvatica*.

Within the chelicerates, but predating the arachnopulmonate WGD, two further tandem

duplications occurred. *Msx* likely duplicated in the common ancestor of ticks,

harvestmen, and arachnopulmonates, and *Emx* likely duplicated in the ancestor of

harvestmen and arachnopulmonates. *NK1* duplicated in tandem, likely just after the WGD, while the other ohnologue was probably lost. Besides these duplications, the majority of tandem duplications in spiders were more recent and lineage-specific (Figure 2 B). In particular, *H. graminicola* has the highest number of gene families with recent tandem duplications (23 families) (Figure 2 B), resulting in this species having the highest number of homeobox genes identified (172). This

Figure 2 – Many homeobox genes were retained in duplicate in scorpion and spider genomes. A) Heatmap of homeobox genes by family. The relationships among duplicates (except for species-specific recent tandem duplications or family/species-specific losses) are indicated to the right. O: ohnologue retained; SOL: spider ohnologue lost; EOL: entelegyne ohnologue lost; AT: ancient tandem duplication; RT: recent tandem duplication; SC: single copy retained. *The second *ftz* ohnologues in *D. plantarius* and *P. tepidariorum* are likely pseudogenes as there are stop codons in the homeodomains. B) The inferred timing of ancient and recent tandem duplications among the included arthropod lineages. Gene families are listed above branches (as well as below for the root and *H. graminicola*). Species abbreviations are as in Table 2.

could be an assembly artefact; however, overall the *H. graminicola* genome did not

exhibit an exceptionally large number of duplicated genes (Zhu et al. 2022).

In contrast to tandem duplicates, dispersed paralogues consistently found on different

chromosomes in surveyed spider genomes were inferred to be ohnologues. There were

between 37 and 46 families with retained ohnologues in arachnopulmonates, and the

families that were retained were largely consistent across species. All the Hox genes

besides *Hox3* and *ftz* were retained as ohnologues, as previously reported for two

spider species (*P. tepidariorum* and *Cupiennius salei*) (Schwager et al. 2017), as well as

a few Super-Hox genes, namely *Dbx*, *En*, and *Gbx* (Figure 2 A). Many of the NK cluster

genes were also retained as ohnologues, including *NK4*, *NK5*, *NK6*, *NK7*, *Lbx*, *Tlx*, as

well as the ohnologues of ancient tandem paralogues of *Emx* and *Msx* (Figure 2 A).

Ohnologues were likely lost from several families following the WGD (Figure 2 A).

Thirty-two families are found in single copies in scorpions and spiders (discounting

subsequent lineage-specific tandem duplications of single-copy genes), indicating that

an ohnologue was lost between the WGD and the divergence of scorpions and spiders.

These include many of the Super-Hox families (*Abox*, *Dlx*, *Evx*, *Meox*, *Mnx*, *Nedx*, and

Ro), the ParaHox family *Gsx*, most of the NK2 cluster families (*Hlx*, *Msxlx*, and *NK2.1*)

and the HRO cluster family *Hbn*. Six families, including the Hox family *ftz*, the ParaHox

family *Cdx* and the HRO cluster family *Otp*, were found in single copies in spiders, while

both ohnologues were retained in the scorpion, indicating the loss of an ohnologue in a

spider ancestor (Figure 2 A). Within spiders, an ohnologue was lost from four families in

the entelegyne ancestor: *Dlx*, *Six4/5*, *Msxlx*, and *NK3*. *D. silvatica*, the sole

synspermiatan representative, has lost the largest number of ohnologues from eighteen

families, though the extent to which this applies across synspermiatans is not known.

Only a few families underwent losses multiple times, namely *Hox3* in *C. sculpturatus*, *D.

plantarius*, *Latrodectus elegans*, and the araneids; *Tgif* in the theridiids, *Dolomedes

plantarius*, *D. silvatica* and *C. sculpturatus*; and *Barx* in both *H. graminicola* and *D.

silvatica*. Only one family, *Vax*, was potentially lost before the WGD, or both ohnologues

were lost in the arachnopulmonate ancestor (Figure 2 A). Though it was not reported

previously (Leite et al. 2018), we identified a single *Vax* gene in the harvestman

genome (Figure 2 A, Table S1).

Homeobox gene clusters in spider genomes

The Hox cluster

Each of the spider genomes studied here contains two Hox clusters (Figure 3),

consistent with previous descriptions for *P. tepidariorum* and other spiders (Schwager et

al. 2007; Schwager et al. 2017; Fan et al. 2021; Sheffer et al. 2021; Wang et al. 2022).

With the new *P. tepidariorum* chromosome-level assembly, we confirmed that each

cluster is found on a different chromosome. Cluster B is typically intact, organised, and

contains orthologues of all ten arthropod Hox genes, while cluster A often lacks *Hox3*

and *ftz*, and is sometimes rearranged (Figure 3). There are some exceptions to this,

however. For one, there have been several tandem duplications of *H. graminicola* Hox

genes resulting in four *lab* paralogues, and three *pb*, *Antp*, *Ubx*, and *Abd-B* paralogues

(Figure 3). While these duplications have occurred in both *H. graminicola* Hox clusters,

cluster B is ordered, and contains the single orthologues of *Hox3* and *ftz*, while cluster A

lacks those genes and has been rearranged (Figure 3). Furthermore, the Super-Hox

gene *Dlx-A* is between *pb-A-2* and *Dfd-A* separated by 124 kb from *pb-A-2* and 380 kb

from *Dlx-A* and *Dfd-A*. In *D. plantarius* there is a new gene, identified with highest

similarity in the homeodomain to *Scr*, but not linked to either Hox cluster, possibly

representing a divergent *Scr* paralogue (blue *'S'* in Figure 3). In *D. silvatica*, both Hox

clusters are rearranged, though one does contain both *Hox3* and *ftz*, while the other has

lost only *ftz* (Figure 3). Pseudogenes of *ftz* were found in *D. plantarius* and *P.*

tepidariorum, identified by the homeodomain disrupted by a stop codon. In *P.*

tepidariorum, the *ftz*-pseudogene is not linked to either Hox cluster, while in *D.*

plantarius, it is adjacent to *ftz*-B (Table S1, Figure S3 F).

Figure 3: Two ohnologous Hox clusters and a disintegrated arthropod ParaHox cluster are found in spiders. Genes are represented by coloured boxes; chromosomes are black lines. Gene orientation is indicated by arrows above or below genes, and distances are indicated by slashes between genes. Gene families are denoted by a one- or two-letter code, and by colour for Hox and ParaHox genes, or in grey for Hox-linked genes. ParaHox genes are coloured based on their relationship to Hox genes, *Gsx* with anterior genes and *Cdx* with posterior genes; *Xlox* was not found in any arthropod genome. Species abbreviations are as in Table 2.

The Super-Hox genes consist of the Hox cluster as well as several ANTP-class genes

consistently found linked to each other and/or the Hox cluster across bilaterians (Table

1) (Butts et al. 2008; Ferrier 2016b). While many Super-Hox genes are still linked to one

another and/or to the Hox cluster in the tick and insects, Super-Hox genes have

dispersed in spider genomes and many Super-Hox families have also returned to the

single copy (Figure 2 A, Figure 3). Amid the overall disintegration of the Super-Hox

cluster, *Abox* and *Meox* are consistently found clustered in all the entelegyne species.

Other remnants of linkage between Super-Hox genes include *Dbx* and *Gbx* linked to

Hox cluster B in *H. graminicola*, *P. tepidariorum*, and *D. plantarius* (Figure 3). *En* is also

often linked to *Mnx*, *Gbx* to *Nedx*, and *Dbx* to *Evx*, though these have more frequently

been disrupted and dispersed, unlike the consistency of *Abox* and *Meox*. Synteny may

be conserved by the constraint of regulatory elements or may result from the passive

maintenance of ancestral linkage. While the Hox cluster has largely been retained in

duplicate in spiders, most of the Super-Hox families have returned to the single copy

and linkage has dispersed following the WGD.

The ParaHox cluster has largely disintegrated in spiders. Neither ParaHox gene is

linked to the other, and both have returned to the single copy (Figure 3). The two *Cdx*

paralogues in *D. silvatica* likely arose in a recent lineage-specific tandem duplication, as

they form a two-gene cluster in only this species. In both insects, *Gsx* (*Ind*) and *Cdx*

(*Cad*) are unlinked, while in the tick they are on the same chromosome but separated

by a large distance (28 Mb) (Figure 3). This is consistent with the ancestral breakdown

of the ParaHox cluster in an arthropod ancestor, and the further separation of the genes

onto separate chromosomes in insects and spiders (Figure 3). It is not clear whether the

retained ParaHox genes belong to the same ohnology group or not (i.e., both A or both

B versus one A and one B), as there is only one ohnologue retained in any of the

species surveyed.

The NK cluster

We found two ohnologous NK clusters in all spider genomes (Figure 4). One consists of

Msx1-A, *NK4-A*, *NK3-A*, *Tlx-A*, *Lbx-A*, and *NK7-A* clustered on one chromosome, while

Msx1-B, *Tlx-B*, *Lbx-B*, and *NK7-B* are found on a different chromosome, also with *NK3-*

B and *NK4-B* in the species that have retained these genes (core NK clusters are

delineated with grey boxes in Figure 4). Cluster B is less organised; *NK3-B* has likely

been lost in entelegynes (estimated divergence 280-310 MYA) (Fernández, Kallal, et al.

2018; Shao and Li 2018) and *NK4-B* has likely been lost in *H. graminicola* and *P.*

tepidariorum, independently. NK core cluster A is conserved across spiders, except that

Msx1-A is not found in *L. elegans* or *P. tepidariorum*, which may indicate the loss of

Msx1-A in a theridiid ancestor (est. 120 MYA) (Fernández, Kallal, et al. 2018). While NK

cluster A is found in a conserved complement and order in almost all spider species

analysed, it is rarely linked to any other genes belonging to the larger ancestral

bilaterian NK complement (Table 1). In contrast, NK cluster B has a varying gene

complement, having undergone lineage-specific gene losses and more rearrangements,

Figure 4: Two ohnologous NK clusters are retained, and a disintegrated NK2 cluster is found in spiders. Genes are represented by coloured boxes; chromosomes are black lines. Gene orientation is indicated by arrows above or below genes, and distances are indicated by slashes between genes. Gene families are denoted by colour and a one- or two-letter code. The genes making up the spider core NK cluster are delineated by grey boxes. Species abbreviations are as in Table 2.

making it more degenerate and less conserved between species. This second cluster is,

however, often linked to other ancestral NK cluster genes *Hhex*, *Noto*, *Emx1-B*, *Emx2-*

B, *NK5-B*, and *NK1*, which has undergone a tandem duplication, and to *NK6-B* and

Msx2-B in all spiders (Figure 4). *NK6-A* is found linked to *Msx2-A* in all the entelegyne

species, and *NK6* and *Msx2* genes can be found in cluster A and/or B in all spiders,

suggesting this two-gene *NK6-Msx2* cluster predates the WGD and was retained in

duplicate some species, while others have lost either *Msx2* or *NK6* ohnologues.

The NK2 cluster has largely broken down in spiders, with successive gene losses: first

Hlx from chelicerates, and then *Msxlx* from entelegynes, though further lineage

sampling may change this pattern (Figure 4). In all spiders, *NK2.2* is retained in

duplicate on two separate chromosomes, indicating an ohnologous relationship, while

NK2.1 has returned to the single copy (Figure 4). *NK2.1* and *NK2.2* are unlinked in all

spiders except for *D. plantarius*, indicating either that the cluster has separated twice, or

alternatively, that different ohnologues have been lost in different lineages.

Phylogenetics were not able to determine which ohnologue of *NK2.1* was retained in

different species because no examined genome contains both ohnologues.

The SINE cluster

There are five SINE-class genes in two clusters in spiders, consistent with a duplication

in the WGD followed by the loss of one *Six4/5* ohnologue. Spider SINE clusters likely

underwent different rearrangements in different lineages (Figure 5). *D. sylvatica* has

retained only one copy of each gene, while the entelegynes have retained B

ohnologues of both *Six1/2* and *Six3/6* in a conserved cluster. In araneids, in cluster A all

three genes are linked, though *Six4/5-A* is separated by a larger distance than that

separating *Six1/2-A* from *Six3/6-A*, which remain tightly clustered. There has also been

an inversion which may have occurred ancestrally among the theridiids that brought

Six4/5-A closer to *Six3/6-A* than *Six1/2-A*. This indicates that the separation of *Six4/5-A*

from the other two genes in araneids occurred after the divergence of the theridiids and

that the three genes making up SINE cluster A were likely still tightly clustered in their

shared ancestor. Among the outgroups, all three SINE genes are linked in the tick, and

only two are linked in the fly, while none are linked in the beetle, indicating the spider

SINE clusters have been retained more intact relative to other arthropod lineages.

The HRO cluster

The PRD-class HRO cluster was so named because the order of the genes in the fly

genome is *hbn* (*Hbn*), *Rx* (*Rax*), then *otp* (*Otp*); this order is conserved in insects and in

the tick genome, indicating that it is likely the ancestral arthropod arrangement (Figure

5). In spiders, the HRO cluster is found intact, but reordered: *Rax*, *Hbn*, then *Otp*

(Figure 5). There have also been several inversions of the orientations of the genes

within the cluster, but the R-H-O order is conserved across spiders, suggesting this

rearrangement occurred once ancestrally. It likely did not predate the WGD but

occurred in a spider ancestor because the scorpion HRO cluster A order is *Hbn*, *Rax*,

then *Otp*, as in insects and the tick (Table S1). As well as one conserved intact HRO

cluster, spiders retained only *Rax* in duplicate, while there are two *Rax* and two *Otp*

ohnologues retained in the scorpion (Figures 2 and 5).

The LIM-class gene *Isl* is consistently found linked to the HRO cluster (Figure 5). Ferrier

(2016) suggested this pattern is evidence for the ancestral linkage between the LIM-

class and PRD-class genes in the Giga-cluster of homeobox progenitor genes in an

animal ancestor. The linkage of *Isl* to the HRO cluster has been observed across

species with a most recent common ancestor in the ancestor of all animals. The

consistent linkage pattern between these genes in spiders, the tick, and beetle

genomes supports this association. This broad phylogenetic distribution makes this

between-class linkage unlikely to be a convergent secondary association; rather it is

more likely to be a conserved ancestral trait.

The Irx cluster

Irx genes have duplicated independently many times (see Table 1), giving rise to six

genes in spiders. Each of the ohnologous *Irx* clusters contains one of each gene type,

Irx1, *Irx2*, and *Irx3*, though *Irx2-B* has been lost from one of the clusters in the

entelegynes (Figure 5). Consistent with the findings of Kerner et al. (2009), insect *Irx*

genes fall into two clades, one containing orthologues of *D. melanogaster mirr* and the other with orthologues of *ara* and *caup* (Table 1 and Figure S1). Spider and scorpion *Irx1-A* and *Irx1-B* group with *ara* and *caup*, while spider and scorpion *Irx2-A* and *Irx2-B* group with *mirr* (Figure S1). The exact relationships between *Irx3* and other arthropod genes remains somewhat unclear. There are three *Irx* genes in the centipede *S. maritima* and the millipede *Trigoniulus corallinus* genomes (Chipman et al. 2014; So et al. 2022), one of which is not contained within either the *Irx1/ara/caup* or *Irx2/mirr* clade, and is resolved as sister to the *Irx3* clade in the *Irx* gene phylogeny, albeit with very low bootstrap support (Figure S1). The potential *Irx3* is clustered with *Irx1* and *Irx2* in the *S. maritima* genome in the same relative position as spider *Irx3* (Figures 5 and S3 C). There is a clear orthologue of each of the three arachnopulmonate *Irx* gene families in both the harvestman and tick genomes, indicating the three gene types originated in or before the chelicerate ancestor, and possibly *Irx3* genes were lost

Figure 5: Two ohnologous HRO clusters, one of which is linked to one of the two ohnologous *Irx* clusters, and two ohnologous SINE clusters are retained in spiders. Genes are represented by coloured boxes; chromosomes are black lines. Gene orientation is indicated by arrows above or below genes, and distances are indicated by slashes between genes. Gene families are denoted by colour and a one- or two-letter code. Species abbreviations are as in Table 2.

from an insect ancestor after the divergence of Myriapoda (est. 520 MYA) (Fernández,

Edgecombe, et al. 2018). According to Kerner et al. (2009), *Irx* genes duplicated

independently in spiralians, ecdysozoans, and deuterostomes, suggesting that all

paralogues arose from a single ancestral bilaterian *Irx* gene. From our phylogeny

(Figure S1), we can tentatively infer that the duplication that gave rise to fly paralogues

is ancestral to the arthropods, and that there was a three-gene cluster in the ancestral

chelicerate that further duplicated in the arachnopulmonate WGD resulting in up to six

spider *Irx* paralogues in two ohnologous clusters.

In spiders, one of the two ohnologous *Irx* clusters is also linked to the HRO+*Isl* cluster

(Figure 5). Since this association occurs only to *Irx* cluster A, while *Irx* cluster B is

consistently not linked to the second *Rax* paralogue (Figure 5, Figure S3 E-L), a likely

explanation is that one *Irx* cluster translocated to be linked to HRO+*Isl* following the

WGD. We infer it occurred secondarily, i.e., on only one ancestral arachnopulmonate

chromosome, rather than deeper in animal ancestry due to the consistent lack of

linkage between HRO+*Isl* with *Irx* genes in any other species examined here, in contrast

to the linkage between HRO and *Isl* found across bilaterians.

Conserved macrosynteny of spider genomes

To contextualise our observations of homeobox cluster conservation and

rearrangements, we explored macrosynteny across spider chromosomes. We used *H.*

graminicola as the reference because the *D. silvatica* genome is highly rearranged

compared to all entelegynes in this study, and *H. graminicola* represents an early-

branching lineage realative to the majority of other entelegynes (Figure 6, Figure S4 A).

Across entelegynes, we detected clear patterns of orthology between chromosomes,

and consistency in the synteny blocks in which certain homeobox clusters are found,

but these patterns were not as evident when compared to the sole synspermiatan

representative *D. silvatica* (Figure 6, Figure S4 A). Depictions of locations of all

homeobox genes across genomes can be found in Figure S3.

Figure 6 – Conserved macrosynteny of spider chromosomes and associated locations of homeobox gene clusters. Vertical bars show stacks of genes on each chromosome representing a macrosynteny block, with *H. graminicola* as the reference genome. Chromosomes of other species were ordered to show the orthologous relationships to *H. graminicola*. Chromosome numbers are denoted below, with 'X' indicating sex chromosomes. Locations of HOX, NK, IRX, SINE and HRO clusters are shown with black arrows, along with the assignment for cluster A (red) versus cluster B (blue). The grey triangle in *T. antipodiana* indicates the partial translocation of Hox cluster A.

In each araneid analysed here,

there are 13 chromosomes;

however, in the theridiids, there

are 12 in *P. tepidariorum* and 14

in *L. elegans* (Figure 6, Figure

S4 A). In *P. tepidariorum*, the

synteny block corresponding to

chromosome 2 in *H.*

graminicola (coral pink in Figure

6) appears to have been mostly

lost or distributed throughout

the genome, and *Irx* cluster B

which belongs to this block has

been relocated (Figure 6,

Figure S4 A). In *L. elegans*

there appear to have been

large fission and fusion events

on chromosomes 2, 4, 5, 6, 8,

11, and 13. Such events in *L.*

elegans are consistent with Hox

cluster A and SINE cluster A

being found on separate chromosomes, as well as the rearrangement of *Irx* cluster B

(Figures 6 and S3 G, Figure S4 A).

Sex chromosomes have been identified in several spider species (Figure 6 and Figure

S4 A) (Sheffer et al. 2021; Zhu et al. 2022; Miller et al. 2023). Interestingly, Hox cluster

B and NK cluster B are both found on the same sex chromosome in *H. graminicola* (12)

and *D. silvatica* (X), but on separate sex chromosomes in other entelegynes, whereas

Hox cluster A is located on an autosome (Figure 6, Figure S4 A). Across entelegynes,

NK cluster A is found in the same conserved synteny block corresponding to a full

chromosome in all species except for *Trichonephila antipodiana*. In this species, a

rearrangement brought NK cluster A onto another chromosome, and this is the only

entelegyne species in which the NK cluster A is rearranged (Figure 4). Overall, this

analysis illustrates the conservation of macrosynteny blocks and chromosomes across

entelegyne genomes and shows several chromosomal rearrangements that have

affected homeobox gene clusters.

Analysis of selection on NK cluster genes

We next sought to understand how homeobox genes have evolved following WGD and

tandem duplications. In the NK cluster, we identified families with single-copy

orthologues, ancient tandem paralogues, and WGD ohnologues. To determine if genes

with different relationships were under different selective pressures, we compared the

aBSREL estimated ω values calculated across coding sequence phylogenies for seven

NK cluster families with different inferred duplication histories: *Msx*, *NK4*, *NK3*, *Tlx*, *Lbx*,

NK7, and *Emx*. While none of the focal ohnologues or tandem paralogues were found to

be under positive selection, certain genes from the pre-duplicate outgroups were. *NK7*

was found to be under positive selection in *D. melanogaster* and *I. scapularis*, *Tlx* also

in *I. scapularis*, and *NK3* in *T. castaneum*. However, certain ohnologues did have longer

branches separating them from other ohnologues or the single-copy outgroup genes in

the aBSREL-generated phylogenies, indicating a higher substitution rate. These include

Msx1 in arachnopulmonates and the harvestman, *NK3-A* in the entelegynes, and *Lbx-B*

in the scorpion. No consistent pattern was detected between A versus B ohnologues,

despite NK cluster A being more organised, nor was there a clear pattern between

ohnologues versus tandem duplicates. Supplementary material underpinning these

results is available on GitHub (see Materials and Methods).

In the one-to-one comparisons between *P. tepidariorum* and *P. opilio* genes, and

between *P. tepidariorum* ohnologues, we calculated dN/dS ratios with codeML to detect

positive selection. For all NK cluster genes, none approached 1; all were very low,

ranging from 0.0011 (between *P. opilio Msx2* and *P. tepidariorum Msx2-A*) to 0.1423

(between *P. tepidariorum Emx1-A* and *Emx1-B*) (Tables 3 and S3). There are some

patterns among gene families, e.g., lower dN/dS ratios for *Tlx* ohnologues compared to

a consistently higher rate for *NK7* ohnologues. The *Emx1* and *Emx2* and *Msx1* and

Msx2 ohnologues had consistently low dN/dS values when compared to their *P. opilio*

orthologues but varied when compared between ohnologues in *P. tepidariorum*. Overall,

these low dN/dS ratios likely reflect the highly conserved nature of homeobox genes.

Table 3 – The ratio of non-synonymous to synonymous substitution rates between *P. tepidariorum*
and *P. opilio* NK cluster genes shows no
evidence for positive selection among
paralogues. Pairwise dN/dS values were
calculated with codeML for each *P. opilio*
and *P. tepidariorum* NK cluster gene. The
bars at the right show the dN/dS value
relative to the black bar (length of 1) at the
top. The relationship between each gene
pair is indicated in the third column, where 1
to 1 indicates direct orthology (the loss of an
ohnologue), 1 to 2 indicates the retention of
two ohnologues in the spider, O is the
comparison between the two spider
ohnologues, and T is the comparison
between tandem paralogues. For families
with ancient tandem duplications (*Msx* and
Emx), gene names have been updated to
reflect the inferred relationships between
paralogues in *P. opilio* and subsequent
ohnologues in *P. tepidariorum*, such that,
e.g., *P. opilio Emx1* is the pro-orthologue of
P. tepidariorum Emx1-A and *Emx1-B*, and
the same for *P. opilio Emx2* and *P.*
tepidariorum Emx2-A and *Emx2-B*. Raw
substitutions (S and N) and substitution
rates (dS and dN) as calculated with
codeML can be found in Table S3.

Popi	Ptep		dN/dS	
NK4	NK4	1 to 1	0.0673	
NK3	NK3	1 to 1	0.0584	
Tlx	Tlx-B	1 to 2	0.0060	
	Tlx-A	1 to 2	0.0094	
	Tlx-A x Tlx-B	O	0.0042	
Lbx	Lbx-B	1 to 2	0.0421	
	Lbx-A	1 to 2	0.0299	
	Lbx-A x Lbx-B	O	0.0076	
NK7	NK7-B	1 to 2	0.0642	
	NK7-A	1 to 2	0.0430	
	NK7-A x NK7-B	O	0.0575	
Msx1	Msx1-B	1 to 1	0.0104	
Msx2	Msx2-A	1 to 2	0.0011	
	Msx2-B	1 to 2	0.0106	
	Msx2-A x Msx2-B	O	0.0051	
	Msx1-B x Msx2-B	T	0.0845	
Emx1	Emx1-A	1 to 2	0.0123	
	Emx1-B	1 to 2	0.0128	
Emx2	Emx2-A	1 to 2	0.0120	
	Emx2-B	1 to 2	0.0094	
	Emx1-A x Emx1-B	O	0.1423	
	Emx2-A x Emx2-B	O	0.0118	
	Emx1-A x Emx2-A	T	0.0188	
	Emx1-B x Emx2-B	T	0.0124	

Figure 7 – Expression of *P. tepidariorum* and *P. opilio NK3* and *NK4*. In all panels, anterior is to the left. Panels A-C, F, G, I, K, M, and N show ventral views. Panels D and H show anterior-dorsal views. Panels E, and J show dorsal views. Panels L and O show lateral views and dorsal up. Panel P shows a posterior-dorsal view. Embryos in panels A, F and K are flat-mounted. Panel B shows a magnification of the posterior region of a flat-mounted embryo. Panels A´-L´, O´ and P´ show SYBR green stained embryos as shown in panels A-L, O and P. Developmental stages of *P. tepidariorum* (after Mittmann and Wolff (2012)) and *P. opilio* (after Gainett et al. (2022)) are indicated in the upper right corner of each panel with a bright field image. Arrows in panels A, C-J, O and P mark expression of *NK3* and *NK4* in the developing heart of *P. tepidariorum*, and expression of *NK4* in the developing heart of *P. opilio*. The arrowhead in panel A points to expression in the labrum. The asterisk in panels A and D mark expression in the head. In panel B, expression in the mesoderm is marked (note absence of expression in the overlying ectoderm). Double arrowheads in panels F, G and I point to dot-like expression of *NK4* in the opisthosoma of the spider. Asterisks in panels F and H mark stripes of expression in the anterior of the head. Asterisks in panels K and L mark two domains of *NK3* expression in the head of the harvestman. Arrows in panels K-M mark short segmental stripes of *NK3* expression. The arrowhead in panel N marks expression in the labrum at late developmental stages. Abbreviations: ch, chelicera; ect, ectoderm; hl, head lobe; L1-L4, first to fourth walking leg; lr, labrum; mes, mesoderm; O, opisthosomal segment; pp, pedipalp; s, stomodaeum.

Developmental expression of NK cluster genes in P. tepidariorum *and* P. opilio

It was previously shown that several ohnologues of spider homeobox genes had

undergone sub- and/or neofunctionalisation with respect to their single-copy

orthologues in the harvestman (Leite et al. 2018). To better understand the roles of

homeobox genes following WGD in the arachnopulmonate ancestor, we assayed the

expression of NK cluster genes in the spider *P. tepidariorum* in comparison to the

harvestman *P. opilio*.

Single-copy genes

Only one copy each of *NK3* and *NK4* were found in both *P. opilio* and *P. tepidariorum*,

indicating the loss of an ohnologue of each from the spider following the WGD. The

single spider *NK3* is strongly expressed in the developing heart and is weakly

expressed around the stomodaeum and in the head (Figure 7 A-E). This is very similar

to the single spider *NK4*, which is also expressed strongly in the heart (Figure 7 F-J).

Unlike *NK3*, *NK4* is also expressed in a dot-like fashion in the ventral nervous system of

the opisthosoma, and at the anterior rim of the head lobes (Figure 7 asterisks and

double arrowheads in Figure 7 F-I). We did not detect any *NK3* expression in the

developing heart of the harvestman; instead *NK3* is expressed in a segmental fashion

ventral to the base of the appendages and in similar short transverse segmental stripes

in the opisthosomal segments (Figure 7 K-N). The expression of harvestman *NK3* is

thus more like the expression of its orthologue in *D. melanogaster* (Azpiazu and Frasch

1993) and the millipede *Glomeris marginata* (Figure S5) than that of the spider.

Expression of *NK4* in the harvestman is restricted to the developing dorsal vessel at late

developmental stages (Figure 7 O-P), a pattern that is conserved among panarthropods

(Azpiazu and Frasch 1993; Janssen and Damen 2008; Treffkorn et al. 2018).

Ancient tandem paralogues

Both *Emx* and *Msx* families have likely undergone an ancient tandem duplication,

resulting in two paralogues of each gene in the harvestman genome, with four *Emx* and

three *Msx* paralogues retained in *P. tepidariorum*. In previous transcriptome-based

analyses of *P. opilio*, the highly similar sequences for each *Emx* paralogue were

merged into a single transcript model, and only a partial sequence was found for *Msx1*,

split across two different transcript models (Sharma et al. 2012), but the genome

revealed distinct loci for each paralogue. In *P. tepidariorum*, Leite et al. (2018)

described *Emx1-A* and *Emx2-A* (previously designated *Emx4* and *Emx3*, respectively)

expression in the precheliceral region and in patches in each segment along the ventral

midline, and *Emx1-B* and *Emx2-B* expression (previously designated *Emx2* and *Emx1*)

in the anterior of each opisthosomal segment, with *Emx1-B* also expressed at the base

of the prosomal appendages. *P. opilio Emx2* (previously designated *Emx*) was detected

at the base of the appendages and in patches along the ventral midline in stage 10-11

embryos (Leite et al. 2018).

Figure 8 – Expression of *P. opilio Emx1* and *Msx2*. In panels A-E, anterior is to the left and ventral views (except panel C, lateral view and dorsal up). Panels F and G each show dissected head lobes and chelicerae, a walking leg and a pedipalp. The dissected heads are shown with anterior up, ventral views. The dissected legs and pedipalps present lateral views, dorsal to the right. Panels A´-E´ show SYBR green stained embryos as shown in panels A-E. Developmental stages after Gainett et al. (2022) are indicated in the upper right corner of each panel with a bright field image. Arrowheads in panels A and B mark the sharp anterior border of *Emx1* expression anterior to the cheliceral segment. The arrows in panels A and C mark sharp distal border of expression in the appendages. Asterisks in panel D mark expression in the ventral nervous system, ventral to the base of the appendages. Double arrowheads in panels A, C and E mark expression in the opisthosoma. Arrows in panels F and G point to expression at the base of the appendages. Arrowheads in panels F and G point to dot-like expression in the chelicerae. Asterisks in panels F and G mark a ring-like expression in the legs. Abbreviations: ch, chelicera; hl, head lobe; L1-L4, first to fourth walking leg; lr, labrum; pp, pedipalp.

Very similar to its paralogue *Emx2*, *P. opilio Emx1* is expressed in all posterior

segments including the cheliceral segment (Figure 8). The sharp anterior border of

expression is conserved among other arthropods, including spiders (Walldorf and

Gehring 1992; Simonnet et al. 2006; Schinko et al. 2008; Birkan et al. 2011; Janssen

2017; Leite et al. 2018) and an onychophoran (Janssen 2017). Indeed, the overall

expression pattern of *P. opilio Emx1* is identical with that of the previously reported

Emx2, with the exception that expression of *Emx1* is detected earlier.

According to Leite et al. (2018), *P. tepidariorum Msx1-B* (previously designated *Msx3*) is

expressed in the base of the prosomal appendages at stage 11, while *Msx2-B*

(previously designated *Msx2*) is expressed in the chelicerae at stage 12, and *Msx2-A*

(previously designated *Msx1*) is expressed in each segment along the ventral midline in

stages 8 and 10 (Akiyama-Oda and Oda 2020). In *P. opilio*, *Msx1* (previously

designated *Msx*) was detected in a stripe in each segment at stage 7 and along the

ventral midline in each segment at stage 13 (Leite et al. 2018). In contrast, we detected

P. opilio Msx2 expression only at late developmental stages and in a very restricted dot-

like patch at the base of the appendages, and a ring-like domain in the legs (Figure 8 F-

G). Thus, *P. opilio Msx1* expression in the segments and along the ventral midline is

similar to *P. tepidariorum Msx2-A* expression, while *P. opilio Msx2* expression at the

base of appendages is similar to *P. tepidariorum Msx1-B* expression. *P. opilio Msx2*

expression was also detected in a small region of the chelicerae, matching the

expression of *Msx2-B* in stage 12 *P. tepidariorum* that was observed by Leite et al.

(2018) and interpreted as a novel expression domain. Our description of a matching

expression pattern from a previously unknown harvestman gene, *Msx2*, means that the

inferred neofunctionalisation is more likely a case of subfunctionalisation between the

tandem *Msx* paralogues.

Ohnologues

Three other NK families, *NK7*, *Tlx*, and *Lbx*, are found in the single copy in *P. opilio* and

have two retained ohnologues in *P. tepidariorum* (Figure 4). The two ohnologues of *P.*

tepidariorum Tlx are both expressed in the distal region of the legs, but in

Figure 9 – Expression of *P. tepidariorum* and *P. opilio Tlx*, *Lbx* and *NK7* ohnologues. In all panels showing whole embryos, anterior is to the left. Dissected heads in panels m, n, and q, are anterior up. Dissected appendages are shown by lateral views, dorsal to the right. Panels A, E, T, V, and g show lateral views and dorsal up. Panels B, C, D, F, I, J, N, O, U, W, Y, b, c, d, h, and i show ventral views. Panels K and P show anterior views. Panel Q shows a posterior view. Panel X shows a dorsal view. The embryo in panel I is flat-mounted. Panels A´-F, I-K´, N´, O´, T´-Y´, b´-d´, and g´-i´ show SYBR green stained embryos as shown in panels with corresponding bright field panels. Developmental stages of *P. tepidariorum* (after Mittmann and Wolff (2012)) and *P. opilio* (after Gainett et al. (2022) are indicated in the upper right corner of each panel with a bright field image. Arrowheads in panels B, D, and E mark dorsal expression in the opisthosoma. The asterisks in panels C and E mark expression in the labrum. The arrows in panels E and F mark expression in the posterior spinneret, and the double arrowhead marks ventral expression in the opisthosoma. Asterisks in panels G and H mark two stripes of stronger expression within the distal region of *Tlx-A* expression in the legs. The asterisk in panel J marks expression of *Tlx-B* in the very posterior of the embryo. The double arrowhead in panel K marks a transverse stripe of expression in the head. The arrows in panels N and O mark expression of *Tlx* in the labrum of the harvestman. The asterisk in panel P marks expression in the head lobes. The arrow in panel U points to expression in the mesoderm of the leg. Arrowheads in panels V-Y point to dorsal expression in the opisthosoma and double arrowheads point to expression ventral to the base of the opisthosomal appendages. The asterisks in panels g-i mark expression in the very posterior of the embryos. The arrows in panels l, m, and q point to expression in the labrum. Asterisks and arrowheads in panels m and q mark expression of *NK7* in the head of the spider and the harvestman. Asterisks in panels o, p, r, and s mark expression in the appendages. Abbreviations: ch, chelicera; L1-L4, first to fourth walking leg; mes, mesoderm; pp, pedipalp; saz, segment addition zone.

complementary patterns (Figure 9 A-M). *Tlx-A* is first expressed in the complete distal

region of the appendages and restricts into two rings of stronger expression around

stage 10 (Figure 9 G-H). *Tlx-B* expression only begins after this point and is strongest in

the region between the two rings of *Tlx-A* expression (Figure 9 M). Apart from its

expression in the developing appendages, *Tlx-A* is also expressed in the dorsal tissue

of the opisthosoma in a pattern that suggests a role in heart development (Figure 9 B,

D-E), and at later stages, in the ventral opisthosoma (Figure 9 F). In contrast, *Tlx-B* is

only expressed in the opisthosoma at the very posterior at late developmental stages,

after segment addition is complete (Figure 9 J). Unlike *Tlx-A*, *Tlx-B* is expressed in the

form of a transverse stripe in the developing head at late developmental stages (double

arrowheads in Figure 9 K) at the anterior edge of non-neurogenic ectoderm that

overgrows the neurogenic ectoderm of the head lobe from the anterior.

The single *Tlx* gene of the harvestman shares many aspects with one or both spider

ohnologues, indicating the spider *Tlx* ohnologues have subfunctionalised. *P. opilio Tlx* is

expressed in the distal half of the developing appendages (Figure 9 R-S), reflecting the

combined expression of both spider *Tlx* ohnologues (Figure 9 G-H, L-M). Like *P.*

tepidariorum Tlx-A, *P. opilio Tlx* is expressed in dorsal tissue of the opisthosoma and in

the labrum (Figure 9 N-O), and like *P. tepidariorum Tlx-B*, *P. opilio Tlx* is expressed at

the anterior edges of the head lobes (asterisk in Figure 9 P).

Expression of *P. tepidariorum Lbx-A* first appears in the mesoderm of the outgrowing

prosomal limb buds (Figure 9 T-U). Later, expression in the appendages is also seen in

the ventral ectoderm, but the dorsal ectoderm remains free of expression (Figure 9 W-

Y). Additional expression of *Lbx-A* is found in the dorsal ectoderm of the opisthosoma

and in segmental patches in the opisthosoma (Figure 9 V-Y). This latter expression is

possibly associated with the ventral-most tissue of the opisthosomal limb buds.

Compared to *Lbx-A*, expression of *P. tepidariorum Lbx-B* appears later and is restricted

to the prosomal appendages (Figure 9 b-d). *Lbx-B* expression within the developing

appendages is similar, but not identical, to that of *Lbx-A*; *Lbx-B* is expressed in fewer

ventral cells than *Lbx-A*. (Figure 9 e-f). Expression of the single *P. opilio Lbx* gene is

similar to that of *P. tepidariorum Lbx-A*, appearing early during development in the

mesoderm of the prosomal appendages (Figure 9 g-h), and later extending into ventral

ectodermal cells of the limbs (Figure 9 j-k). Unlike the two *P. tepidariorum Lbx*

ohnologues, *P. opilio Lbx* is also expressed at the posterior of the developing embryo

(asterisks in Figure 9 h-i).

Expression of both ohnologues of *P. tepidariorum NK7* were only detected in relatively

late stages of embryogenesis. *NK7-A* is exclusively expressed in the stomodaeum and

in a few patches in the head lobes (Figure 9 l-m). *NK7-B* is expressed in three rings in

the developing walking legs (Figure 9 n-p). The expression of *P. opilio NK7* combines

the expression of both *P. tepidariorum NK7* genes as it is expressed in the head lobes

and in three ventral domains in the legs (Figure 9 q-s), again suggesting the

subfunctionalisation of the spider *NK7* genes.

DISCUSSION

Evolution of the repertoire and organisation of spider homeobox genes before and after

the WGD

We detected widespread conserved patterns of homeobox gene content and clusters

across spider genomes that were impacted both by tandem duplications and the WGD.

Leite et al. (2018) previously discovered that up to 59% of spider homeobox families

were duplicated in arachnopulmonates and that most of these were dispersed

duplicates rather than closely linked tandem duplicates. However, the few fragmented

spider genomes available at the time meant that whether these were ohnologues or

tandem duplicates could not reliably be determined. Here we showed that most

duplicated homeobox genes are likely ohnologues retained after WGD, and the majority

have been retained in all spiders surveyed (Figure 2 A, Table 2). The

arachnopulmonate WGD has therefore played a prominent role in shaping the

homeobox gene repertoires of spiders.

Duplicated homeobox gene clusters seem to have evolved asymmetrically following the

WGD. Between the two ohnologous Hox clusters as well as the two NK, *Irx*, and SINE

clusters, one cluster is more conserved in gene content, orientation, and order than the

other. This asymmetry could be a result of mechanisms of regulatory

subfunctionalisation (e.g., complementary degenerating mutations to regulatory

elements (Force et al. 1999)) acting not only at the level of individual genes, but also on

homeobox gene clusters.

Contextualising the organisation of the conserved duplicated homeobox gene clusters

across entire spider genomes, we found that synteny is largely conserved among the

entelegynes (Figure 6). We also observed a few rearrangements of homeobox clusters

associated with chromosomal rearrangements (Figure S4 B and C). In *H. graminicola*,

Hox cluster B and NK cluster B are both on chromosome 12, one of the two sex

chromosomes, while in the other entelegynes, the synteny blocks that make up the *H.*

graminicola sex chromosomes are found on two distinct chromosomes, one with Hox

cluster B and the other with NK cluster B. This could have arisen by a rearrangement

among the sex chromosomes in *H. graminicola* or independent rearrangements in

theridiids and araneids. Given the X_1X_2 male and $X_1X_1X_2X_2$ female sex determination in

most entelegynes including the species shown here (Maddison and Leduc-Robert

2013), differential rates of purifying selection on sex chromosomes versus autosomes

may impact the extent to which the synteny of these chromosomes is conserved, thus

affecting the homeobox clusters on them (Kořínková and Král 2013; Charlesworth 2017;
Cordellier et al. 2020).

There seems to be a disconnect between the relative organisation of spider Hox
clusters and their regulation. Hox genes exhibit spatial and temporal collinearity (Ferrier
2019 and the references therein). The regulatory elements responsible for temporal
collinearity constrain Hox gene organisation, thus it has been hypothesised that Hox
cluster integrity is essential for the mechanism of temporal collinearity (Duboule 2007;
Krumlauf 2018; Ferrier 2019). While spatial collinearity is exhibited by both spider Hox
clusters, temporal collinearity is more evident in cluster A, though this cluster is less
organised (Schwager et al. 2017; Turetzek et al. 2022). Hox cluster organisation and
regulation vary between animals, for example, in mammals, each Hox cluster is
condensed and is regulated globally, while in some invertebrates the Hox cluster is
organised and expressed in sub-clusters (Kmita and Duboule 2003; Deschamps and
Duboule 2017; Wang et al. 2017). This necessitates a better understanding of the
relationship between the organisation and regulation of the ohnologous spider Hox
clusters.

The conservation of other homeobox clusters indicates that they may similarly be
constrained by regulatory elements. ANTP-class Super-Hox genes arose with the Hox
cluster, and many are still linked to one another and to the Hox cluster across
bilaterians (Butts et al. 2008; Ferrier 2016b). However, in spiders, the only consistent
pattern among Super-Hox genes was the conserved clustering of *Abox* and *Meox*,
though these two genes were not linked to either Hox cluster. This suggests that amid
many rearrangements and gene losses, there may be a functional constraint on this
two-gene cluster in spiders that would not have been recognised without comparisons
across this taxonomic breadth.

Our comparison of NK cluster genes in spiders with other bilaterians revealed a pattern
of conserved sub-clusters. In spiders, we found a core NK cluster consisting of a subset
of the bilaterian NK complement largely conserved in gene order (Figure 4). Its
ohnologous cluster, NK cluster B, is less conserved but is linked to more of the

bilaterian NK complement (Figure 4). Some NK gene organisation patterns seen in

spiders are similar to those in other arthropods. While the NK genes in different

drosophilid species have undergone a series of rearrangements, fragmenting and then

reuniting the NK cluster, certain gene pairs or triplets are consistently clustered and

move as units referred to as "contiguities" (Chan et al. 2015). In spiders, *NK3* and *NK4*

are adjacent, as in *D. melanogaster*, *T. castaneum*, *Bombyx terrestris*, many butterflies,

and *Anopheles gambiae*, and *Tlx* and *Lbx* are adjacent, as in *D. melanogaster*, *T.*

castaneum, several butterflies, *A. gambiae* and *I. scapularis* (Figure 4) (Luke et al.

2003; Chan et al. 2015; Mulhair et al. 2022). Though *NK1* has undergone a recent

tandem duplication in spiders, *NK5* and *NK1* are also adjacent in *T. castaneum*, *B.*

terrestris, *A. gambiae*, and *I. scapularis*, but not *D. melanogaster* or butterflies (Figure 4)

(Chan et al. 2015; Mulhair et al. 2022).

These NK sub-clusters are even conserved across other bilaterians (Luke et al. 2003;

Larroux et al. 2007; Wotton et al. 2009; Hui et al. 2012; Mulhair et al. 2022). In the

annelid *Platynereis dumerilii* and the invertebrate chordate amphioxus *Branchiostoma*

floridae, *NK3* and *NK4* are clustered, as are *Lbx* and *Tlx* (Hui et al. 2012), while in

human, one *NK4* ohnologue clusters with *Tlx* and *Lbx* ohnologues (Luke et al. 2003;

Wotton et al. 2009). In contrast, the NK sub-clusters of tardigrades consist of different

subgroups (Treffkorn et al. 2018), which most likely is a result of rearrangements in this

lineage. Overall, these patterns suggest that while the entire NK cluster has undergone

many rearrangements in different animal lineages, these events have repeatedly

occurred between these sub-clusters, and far less often within them. This semi-flexible

organisation distinguishes the NK cluster from the Hox cluster, which has a more

consistent gene complement and order, but each of the NK sub-clusters may be

maintained by a conserved regulatory mechanism similar to the Hox cluster.

NK cluster genes show conserved and subfunctionalised expression

We observed that the genes in NK sub-clusters are expressed in similar patterns. NK

cluster genes are broadly involved in patterning the mesoderm (Jagla et al. 2001;

Saudemont et al. 2008; Treffkorn et al. 2018). The expression of *NK3* and *NK4* in both

arachnids is similar to the *D. melanogaster* orthologues *tin* and *bap* in the heart and

visceral mesoderm (Azpiazu and Frasch 1993). The fly *Tlx* orthologue, *C15*, and *Lbx*

paralogues *lbe* and *lbl* are also expressed in legs like their orthologues in both *P.*

tepidariorum and *P. opilio* (Jagla et al. 2001; Campbell 2005; Maqbool et al. 2006). This

suggests these genes share conserved cis-regulatory elements that may not only

preserve their clustering but also result in their co-expression. It would be interesting to

identify these regulatory elements and to investigate the extent of co-expression within

NK sub-clusters more broadly across bilaterians.

Though a few examples of potential neofunctionalisation have been described,

subfunctionalisation seems to be the more common fate for duplicated homeobox

genes (Schwager et al. 2017; Leite et al. 2018), and other genes (Schomburg et al.

2015; Janssen et al. 2021) in spiders. Using a more thorough annotation of NK genes

from the harvestman, we were able to determine the relative functional consequences

of tandem duplication versus the WGD, which was particularly important for families

with ancient tandem duplication events like *Emx* and *Msx*. Expression of *Emx* tandem

paralogues were similar while A and B ohnologues have undergone temporal

subfunctionalisation (Leite et al. 2018), despite the *Emx* tandem duplication now

estimated to predate the WGD. Furthermore, the two *Emx* paralogues in *P. opilio* are

expressed in near-identical patterns (Figure 8) (Leite et al. 2018), which suggests that

while the tandem duplication gave rise to two conserved paralogues, *Emx* genes did not

subfunctionalise until after the WGD. For *Msx*, we found that *P. opilio Msx2* exhibits a

similar expression pattern to *Msx2-B*, which now seems more likely to have undergone

subfunctionalisation instead of neofunctionalisation. This stresses the importance of

understanding the full evolutionary history of a gene family before inferring sub- and/or

neofunctionalisation. Furthermore, the roles of the ancestral single copies of *Msx* and

Emx are not yet known but could be determined with comparisons to earlier-branching

lineages of chelicerates or arachnids.

It is thought that duplicated genes encoding developmental transcription factors like

homeobox genes are more likely to undergo regulatory subfunctionalisation, driven by

the complementary loss of enhancers (Force et al. 1999; Jiménez-Delgado et al. 2009;

Espinosa-Cantú et al. 2015; Marlétaz et al. 2018). Indeed, the clear patterns of

subfunctionalisation observed for the *P. tepidariorum* NK ohnologues and single-copy

orthologues from *P. opilio* despite no NK cluster ohnologues being found to be under

positive selection suggests changes primarily occurred in their regulatory regions.

Comparisons to the outcomes of other WGD events

Our results suggest that there have been similar outcomes between the

arachnopulmonate WGD and the likely independent WGDs in horseshoe crabs (Nossa

et al. 2014; Kenny et al. 2016; Shingate et al. 2020; Nong et al. 2021). Here, many

homeobox gene families are expanded to four or more copies and there are multiple

duplicated homeobox clusters, including up to six Hox clusters, six NK clusters, and

seven SINE clusters as a result of up to three WGDs (Nong et al. 2021). Like spiders, in

horseshoe crabs some ohnologous clusters are more intact than others; for instance,

two of the six Hox clusters are intact and ordered, while the others have lost between

two and eight genes (Shingate et al. 2020). However, in the absence of additional

chromosome-level assemblies, the extent to which this applies to other horseshoe crabs

is not yet clear.

In comparison to vertebrate homeobox gene clusters resulting from the 2R WGD, the

ohnologous clusters of spiders are more conserved in terms of gene content. Across the

four human Hox clusters, complementary gene losses resulted in no cluster containing

all gene families, but in spiders, one Hox cluster is intact while the other has lost only

two genes (Figure 3). The four ohnologous human NK clusters are spread across five

chromosomes, and most genes are retained in two or maximum three copies (Luke et

al. 2003; Wotton et al. 2009). While in spiders the HRO cluster is intact, *Hbn* is not

found in chordates, and *Otp* is not linked to *Rax* or *Rax2* in the human genome (Mazza

et al. 2010). However, for SINE genes, the clusters have evolved similarly between

vertebrates and spiders following WGD (Kawakami et al. 2000; Ferrier 2016b). In both

of these lineages, linkage of all three SINE genes is conserved on one chromosome,

while the ohnologous cluster(s) have undergone losses and rearrangements. Therefore,

the pattern of one copy being retained intact while the other has degenerated, as seen

for most ohnologous spider homeobox gene clusters, is similar for only some human

homeobox clusters. Still, these few similarities illustrate how duplication can lead to

asymmetric degradation of otherwise conserved gene clusters.

Besides conserved clusters, some homeobox genes have returned to the single copy in

both humans and spiders or scorpions, namely *Hlx*, *Hhex*, *Mnx*, *Dmbx*, *Otp*, *Prop*, and

Mkx (Holland et al. 2007; Leite et al. 2018). Genes often found in the single copy in both

spiders and horseshoe crabs are *Bari*, *Barx*, *CG11294*, *Dmbx*, *Hhex*, *Meox*, *Mkx*, *Msxlx*,

and *NK4* (Nong et al. 2021). This suggests that certain genes may be constitutive

singletons, disadvantageous in multiple copies, or unable to subfunctionalise and thus

lost. However, these comparisons are only between three distantly related species;

wider sampling of WGD lineages may reveal clearer patterns of genes that tend to

return to the single copy and the underlying reasons.

WGDs are hypothesised to have had important evolutionary consequences, primarily

based on the finding that 2R WGD occurred along the vertebrate stem. However, the

impact of WGDs in other animal lineages remains unclear. We still do not know to what

extent the evolution of the morphological novelties and species diversity of spiders was

driven by the arachnopulmonate WGD, though our work contributes to a growing body

of evidence detailing widespread subfunctionalisation of developmental ohnologues.

The homeobox genes we have characterised here as well as other developmental

genes provide candidates for future investigations, which, in concert with future studies

of other chelicerate lineages, advance our aim to understand the impact of the

arachnopulmonate WGD and WGDs more widely across animals.

CONCLUSIONS

Analyses of WGDs in arachnopulmonates, horseshoe crabs, and vertebrates shows

that homeobox genes, as well as other regulatory genes are commonly retained in

duplicate (Cañestro et al. 2013; Berthelot et al. 2014; Leite et al. 2016; Harper et al.

2021; Janssen et al. 2021). Across eight spiders, representing three major clades we

found that the majority of duplicated homeobox genes likely originated in the

arachnopulmonate WGD and that often, cluster or sub-cluster integrity has been

maintained in at least one ohnologous copy. Comparisons of expression patterns of

selected homeobox genes support extensive subfunctionalisation among spider

ohnologues. Taken together, our results help us understand the impact of WGD and

tandem duplication on the evolution of homeobox genes and genomes in spiders and

provide an important comparison to WGD in other animals.

MATERIALS AND METHODS

Homeobox gene annotation

We classified the homeobox gene complement of the publicly available chromosome-

level genomes from eight spider species, *D. silvatica* (Sánchez-Herrero et al. 2019;

Escuer et al. 2022), *D. plantarius* (Blaxter et al. 2022), *L. elegans* (Wang et al. 2022), *P.*

tepidariorum (Zhu et al. 2023), *H. graminicola* (Zhu et al. 2022), *A. bruennichi* (Sheffer

et al. 2021), *T. antipodiana* (Fan et al. 2021), and *T. clavata* (Hu et al. 2022). We

examined the previous homeobox gene classification of the scorpion *C. sculpturatus*,

which shared the ancestral WGD, as an outgroup to the spiders (Schwager et al. 2017;

Leite et al. 2018). As non-WGD outgroups, we included two arachnids, the harvestman

P. opilio (Gainett et al. 2021), and the tick *I. scapularis* (De et al. 2023), as well as three

mandibulates: the centipede *S. maritima* (Chipman et al. 2014; So et al. 2022), the fruit

fly *D. melanogaster* (BDGP6.32), in which many of these homeobox genes and families

were first described (Zhong et al. 2008), and the beetle *T. castaneum* (Herndon et al.

2020), which was used by Butts et al. (2008) to describe the Super-Hox cluster. Only

chromosome-level assemblies were used for synteny analyses, but homeobox genes

were counted and classified in all genomes. Genome accessions are shown in Table 4.

Table 4 – Genome assemblies used in this study and accessions. NCBI accession numbers or
sequence database DOIs are given for genomes with homeobox genes annotated in this study for the first
time. Citations are given for genomes with published homeobox gene annotations. Genome size, N50, and
BUSCO scores are shown for each genome as calculated for the annotated chromosome-level assemblies
with the arthropoda_odb10 database using BUSCO v5.4.7 or were taken from NCBI and the genome
papers (cited in the text above). Abbreviations: C: completeness; S: single copy; D: duplicated. The first-
listed *A. bruennichi* genome, sequenced by the Darwin Tree of Life project, had a higher N50 and larger
size, so was used for homeobox annotation, however it has not been annotated, so an older version, the

second-listed, was used for the whole genome synteny comparison as it has a corresponding annotation.
A homeobox annotation was also conducted on this older version and can be found at the bottom of Table
S1. The older *P. tepidariorum* assembly was not to chromosome-level so homeobox genes were
reannotated on the newer chromosome-level genome despite existing annotations on the previous version.

Species	Accession or Citation	Size	N50	BUSCO
T. clavata	https://spider.bioinfotoolkits.net/base/download/-1	2.6 Gb	202.1 Mb	C:83.8% [S:80.8%, D:3.0%]
T. antipodiana	http://dx.doi.org/10.5524/100868	2.3 Gb	172.9 Mb	C:96.3% [S:92.5%, D:3.8%]
A. bruennichi (DToL)	GCF_947563725.1	1.8 Gb	139.1 Mb	unannotated
A. bruennichi (OLD)	GCA_015342795.1	1.7 Gb	124.2 Mb	C:87.0% [S:81.5%, D:5.5%
H. graminicola	GCA_023701765.1	936.1 Mb	77.1 Mb	C:96.0% [S:92.2%, D:3.8%]
P. tepidariorum (OLD)	GCF_000365465.2; annotated by Schwager et al. 2017, Leite et al. 2018	1.4 Gb	4.1 Mb	98% completeness reported
P. tepidariorum (NEW)	https://doi.org/10.11922/sciencedb.o00019.00014	1.1 Gb	93.9 Mb	C:95.2% [S:92.4%, D:2.8%]
L. elegans	http://dx.doi.org/10.5524/102210	1.6 Gb	114.3 Mb	C:58.3% [S:55.2%, D:3.1%]
D. plantarius	GCA_907164885.2	2.8 Gb	216.7 Mb	unannotated
D. silvatica	GCA_006491805.2	1.4 Gb	174.2 Mb	C:77.2% [S:74.7%, D:2.5%]
C. sculpturatus	GCA_000671375.2; annotated by Schwager et al. 2017, Leite et al. 2018	925.5 Mb	537.5 kb	96.8% completeness reported
P. opilio	GCA_019434445.1	576.9 Mb	211 kb	95.1% reported
I. scapularis	GCA_016920785.2; annotated by Schwager et al. 2017, Leite et al. 2018	2.2 Gb	132.1 Mb	C:98.8% [S:93.4%, D:5.4%]
S. maritima	GCA_000239455.1; annotated by Chipman et al. 2014, Leite et al. 2018	176.2 Mb	139.5 kb	93.8% completeness reported
T. castaneum	GCA_000002335.3; HomeoDB; Zhong et al. 2008, Zhong and Holland, 2011	165.9 Mb	4.5 Mb	C:98.4% [S:97.7%, D:0.7%]
D. melanogaster	GCA_000001215.4; HomeoDB; Zhong et al. 2008, Zhong and Holland, 2011	143.7 Mb	25.3 Mb	C:92.9% [S:92.3%, D:0.6%]

We used tBLASTn searches (Altschul et al. 1990) with the contents of HomeoDB

(accessed July 2020) (Zhong et al. 2008; Zhong and Holland 2011), as well as the

previous classification of *P. tepidariorum* homeobox genes (Schwager et al. 2017; Leite

et al. 2018) as the query set to retrieve homeodomain sequences from all eight spider

genomes, and re-annotated the homeobox genes for the harvestman, tick, and

centipede. Homeodomain sequences were then curated by tBLASTn searches against

annotated gene models or transcriptomes for all species except *D. plantarius*, for which

an annotation is not available, and classified based on sequence similarity and

diagnostic residues of the homeodomain for all families. For some gene families with

several duplication events, *e.g.*, *Irx*, *Pax4/6*, *Pax3/7*, as well as for the NK cluster genes,

phylogenies were constructed (see below). All homeodomain sequences can be found

in the supplementary information (Table S1).

Phylogenetics

To determine the timing of duplication of the arachnopulmonate *Irx* genes relative to

pre-duplicate paralogues, we made a full-length amino acid sequence alignment from a

from a subset of our spider species, the scorpion, and a transcriptome from another

synspermiatan spider, *Segestria senoculata* (Baudouin-Gonzalez unpublished). Full

sequences for genomes with poor annotations were curated from GenBank searches

and BLASTp (Altschul et al. 1990) searches against nr (Benson et al. 2018). These

sequences were aligned with MUSCLE set to the protein alignment defaults (Edgar

2004) and manually curated and trimmed to remove gaps. We inferred a Maximum-

Likelihood phylogeny with IQ-TREE (Nguyen et al. 2015), determined the best

substitution model with ModelFinder (Kalyaanamoorthy et al. 2017), and used 1,000

bootstrap replicates to optimise topology support. The phylogeny was visualised with

FigTree (FigTree 2018). The full and trimmed alignments and the phylogeny can be

found in the supplementary information on GitHub (Figure S1,

[https://github.com/madeleineaaseremedios/SpiderHomeoboxSequences]). Other family

phylogenies were made using only the homeodomain sequence and aligned manually,

but inferred as for the *Irx* family tree, except for the NK cluster and NK-linked genes

phylogeny, which used full protein sequences and was aligned with MUSCLE set to the

protein default (Edgar 2004) (Figure S2).

Whole genome synteny comparison

We performed whole genome synteny comparisons to explore chromosomal

rearrangements that might underlie trends in homeobox placement in spider genomes.

We used all spider species mentioned above, except for *D. plantarius* due to the lack of

genome annotation, and compared these to non-spider species *I. scapularis*, *T.*

castaneum, and *D. melanogaster*. Hierarchical orthogroups (HOGs) of all non-

homeobox genes were identified using Orthofinder (Emms and Kelly 2019) with default

settings, specifying the species tree:

((Dmel,Tcas)N1,(Isca,(Dsil,((Ptep,Lele)N5,(Hgra,(Abru,(Tcla,Tant)N8)N7)

N6)N4)N3)N2)N0);. These were combined with HOGs of our homeobox orthology and

used to reconstruct ancestral gene orders with Agora (Muffato et al. 2023), using the

basic pipeline. However, we used Agora to illustrate chromosomes with macrosynteny

and orthology relationships, rather than the ancestral synteny blocks, termed

Contiguous Ancestral Regions (CARs) detected by Agora, because it resolved a better

signal of chromosomal rearrangements across the whole genome compared to the

smaller coverage of CARs. We also used MCScanX (Wang et al. 2012) to capture

microsynteny relationships between spider genomes and used Circos (Krzywinski et al.

2009) to plot synteny relationships and homeobox positions in the genome.

Coding sequence evolutionary analyses

For core NK cluster genes, we curated full coding sequences for a subset of species

with the best annotations: *P. tepidariorum*, *A. bruennichi*, *D. silvatica*, *D. melanogaster*,

T. castaneum and *I. scapularis*, as well as four species not included in the synteny

comparison: the centipede *S. maritima*, the harvestman *P. opilio*, the scorpion *C.*

sculpturatus, and the onychophoran *Euperipatoides rowelli*, for which NK homeobox

genes have been described, but which lack chromosomal-level genome assemblies

(Chipman et al. 2014; Schwager et al. 2017; Leite et al. 2018; Treffkorn et al. 2018).

The NK cluster genes contained families with different duplication histories, including

WGD ohnologues in arachnopulmonates, tandem duplications both before and after the

WGD, and families with only single genes represented after the WGD, allowing us to

compare possible signatures of subfunctionalisation between the different evolutionary

trajectories and different types of duplicates.

We used aBSREL v2.3 (Smith et al. 2015) to detect selection on each branch of the

phylogeny for each of the spider NK core cluster gene families (*Msx*, *NK3*, *NK4*, *Tlx*,

Lbx, and *NK7*) and *Emx*. We used sequences from three spiders, *P. tepidariorum*, *D.*

silvatica, and *A. bruennichi*, and the scorpion, harvestman*,* tick, fly, and beetle, as

above, and rooted the trees with sequences from the onychophoran (Treffkorn et al.

2018). We created and trimmed codon alignments with PAL2NAL (Suyama et al. 2006)

from coding sequences and protein sequence alignments inferred with MUSCLE set to

the protein alignment default (Edgar 2004). We ran aBSREL using hyphy v2.5.46 and

the outputs were visualised with HyPhy Vision at [http://vision.hyphy.org/] (Pond et al.

2005; Kosakovsky Pond et al. 2020). We also conducted pairwise comparisons of

dN/dS ratios between *P. opilio* and *P. tepidariorum* for NK cluster genes calculated with

codeML (Goldman and Yang 1994) from codon alignments inferred as above for each

gene pair. Sequence data and aBSREL outputs can be found in the supplementary

information on GitHub at [https://github.com/madeleineaaseremedios/NKseqevoldata]

and visualised with HyPhy Vision as above.

Animal husbandry, embryo collection, PCR, riboprobe synthesis, and in situ

hybridization

Embryos of the spider *P. tepidariorum* were collected from the colony established in

Uppsala, Sweden, and maintained as described in Prpic et al. (2008). Embryos of the

harvestman *P. opilio* were collected from wild caught specimens in Uppsala, Sweden.

Adult specimens and embryos of *P. opilio* were treated as described in Janssen et al.

(2021). We applied the staging system of Mittmann and Wolff (2012) for *P.*

tepidariorum, and Gainett et al. (2022) for *P. opilio*. We investigated all developmental

stages of *P. tepidariorum* and *P. opilio* from the formation of the early germ band to

dorsal closure. Total RNA from *P. tepidariorum* and *P. opilio* was isolated from a mix of

embryonic stages using TRIzol (Invitrogen). mRNA was isolated from total RNA using

the Dynabeads mRNA Purification System (Invitrogen) and reverse transcribed into

cDNA using the SuperScriptII system (Invitrogen). PCRs were performed applying sets

of gene specific primers. For all investigated genes, reverse primers were ordered with

additional 5-prime T7-promotor sequences (gggTAATACGACTCACTATAG) for

subsequent antisense RNA probe synthesis (David and Wedlich 2001). *In situ*

hybridizations were performed as per Janssen et al. (2018). Primer sequences can be

found in Table S2.

DATA AVAILABILITY STATEMENT

All data underpinning this study is available through the links above or in the

supplementary information.

AUTHOR CONTRIBUTIONS

APM and LSR conceived of the project. MEAR annotated the homeobox genes,

conducted the phylogenetics and protein sequence analyses, and drafted the

manuscript. RJ characterised the developmental expression of NK genes. DJL

conducted the whole genome macrosynteny analysis. All authors contributed to writing

and approved the final version of the manuscript.

ACKNOWLEDGEMENTS

We would like to thank Julio Rozas and Paula Escuer for early access to the *D. silvatica*

genome. This project was funded in by a NERC grant (NE/T006854/1) to APM and

LSR.

REFERENCES

Aase-Remedios ME, Ferrier DEK. 2021. Improved Understanding of the Role of Gene
and Genome Duplications in Chordate Evolution With New Genome and
Transcriptome Sequences. *Front. Ecol. Evol.* 9:429.
https://doi.org/10.3389/fevo.2021.703163.

Abzhanov A, Popadic A, Kaufman TC. 1999. Chelicerate Hox genes and the homology
of arthropod segments. *Evol. Dev.* 1:77–89. https://doi.org/10.1046/j.1525-
142x.1999.99014.x.

Akiyama-Oda Y, Oda H. 2020. Hedgehog signaling controls segmentation dynamics
and diversity via msx1 in a spider embryo. *Sci. Adv.* 6:eaba7261.
https://doi.org/10.1126/sciadv.aba7261.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
search tool. *J. Mol. Biol.* 215:403–410. https://doi.org/10.1016/S0022-
2836(05)80360-2.

Averof M, Patel NH. 1997. Crustacean appendage evolution associated with changes in
Hox gene expression. *Nature* 388:682–686. https://doi.org/10.1038/41786.

Azpiazu N, Frasch M. 1993. tinman and bagpipe: two homeo box genes that determine
cell fates in the dorsal mesoderm of Drosophila. *Genes Dev.* 7:1325–1340.
https://doi.org/10.1101/gad.7.7b.1325.

Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Ostell J, Pruitt KD, Sayers EW.
2018. GenBank. *Nucleic Acids Res.* 46:D41–D47.
https://doi.org/10.1093/nar/gkx1094.

Berthelot C, Brunet F, Chalopin D, Juanchich A, Bernard M, Noël B, Bento P, Da Silva
C, Labadie K, Alberti A, et al. 2014. The rainbow trout genome provides novel
insights into evolution after whole-genome duplication in vertebrates. *Nat.*
Commun. 5:3657. https://doi.org/10.1038/ncomms4657.

Birkan M, Schaeper ND, Chipman AD. 2011. Early patterning and blastodermal fate
map of the head in the milkweed bug Oncopeltus fasciatus. *Evol. Dev.* 13:436–
447. https://doi.org/10.1111/j.1525-142X.2011.00497.x.

Blaxter M, Mieszkowska N, Di Palma F, Holland P, Durbin R, Richards T, Berriman M,
Kersey P, Hollingsworth P, Wilson W, et al. 2022. Sequence locally, think
globally: The Darwin Tree of Life Project. *Proc. Natl. Acad. Sci.* [Internet] 119.
https://doi.org/10.1073/pnas.2115642118. Available from:
http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2115642118/-
/DCSupplemental.

Brooke NM, Garcia-Fernàndez J, Holland PWH. 1998. The ParaHox gene cluster is an
evolutionary sister of the Hox gene cluster. *Nature* 392:920–922.
https://doi.org/10.1038/31933.

Brunet F, Lorin T, Bernard L, Haftek-Terreau Z, Galiana D, Schartl M, Volff JN. 2017.
Case studies of seven gene families with unusual high retention rate since the
vertebrate and teleost whole-genome duplications. In: Evolutionary Biology:
Self/Nonself Evolution, Species and Complex Traits Evolution, Methods and
Concepts. Springer International Publishing. p. 369–396.

Bürglin TR. 2011. Homeodomain Subtypes and Functional Diversity. In: Hughes TR,
editor. A Handbook of Transcription Factors. Subcellular Biochemistry.
Dordrecht: Springer Netherlands. p. 95–122. Available from:
https://doi.org/10.1007/978-90-481-9069-0_5

Butts T, Holland PWH, Ferrier DEK. 2008. The Urbilaterian Super-Hox cluster. *Trends*
Genet. 24:259–262. https://doi.org/10.1016/j.tig.2007.09.006.

Butts T, Holland PWH, Ferrier DEK. 2010. Ancient homeobox gene loss and the
evolution of chordate brain and pharynx development: deductions from
amphioxus gene expression. *Proc. R. Soc. B Biol. Sci.* 277:3381–3389.
https://doi.org/10.1098/rspb.2010.0647.

Campbell G. 2005. Regulation of gene expression in the distal region of the Drosophila
leg by the Hox11 homolog, C15. *Dev. Biol.* 278:607–618.
https://doi.org/10.1016/j.ydbio.2004.12.009.

Cañestro C, Albalat R, Irimia M, Garcia-Fernàndez J. 2013. Impact of gene gains,
losses and duplication modes on the origin and diversification of vertebrates.
Semin. Cell Dev. Biol. 24:83–94. https://doi.org/10.1016/j.semcdb.2012.12.008.

Chan C, Jayasekera S, Kao B, Páramo M, von Grotthuss M, Ranz JM. 2015.
Remodelling of a homeobox gene cluster by multiple independent gene reunions
in Drosophila. *Nat. Commun.* 6:6509. https://doi.org/10.1038/ncomms7509.

Charlesworth D. 2017. Evolution of recombination rates between sex chromosomes.
Philos. Trans. R. Soc. B Biol. Sci. 372:20160456.
https://doi.org/10.1098/rstb.2016.0456.

Chipman AD, Ferrier DEK, Brena C, Qu J, Hughes DST, Schröder R, Torres-Oliva M,
Znassi N, Jiang H, Almeida FC, et al. 2014. The First Myriapod Genome
Sequence Reveals Conservative Arthropod Gene Content and Genome
Organisation in the Centipede Strigamia maritima.Tyler-Smith C, editor. *PLoS*
Biol. 12:e1002005. https://doi.org/10.1371/journal.pbio.1002005.

Cordellier M, Schneider JM, Uhl G, Posnien N. 2020. Sex differences in spiders: from
phenotype to genomics. *Dev. Genes Evol.* 230:155–172.
https://doi.org/10.1007/s00427-020-00657-6.

David R, Wedlich D. 2001. PCR-Based RNA Probes: A Quick and Sensitive Method to
Improve Whole Mount Embryo In Situ Hybridizations. *BioTechniques* 30:768–
774. https://doi.org/10.2144/01304st02.

De S, Kingan SB, Kitsou C, Portik DM, Foor SD, Frederick JC, Rana VS, Paulat NS,
Ray DA, Wang Y, et al. 2023. A high-quality Ixodes scapularis genome advances
tick science. *Nat. Genet.* 55:301–311. https://doi.org/10.1038/s41588-022-01275-
w.

Deschamps J, Duboule D. 2017. Embryonic timing, axial stem cells, chromatin
dynamics, and the Hox clock. *Genes Dev.* 31:1406–1416.
https://doi.org/10.1101/gad.303123.117.

Duboule D. 2007. The rise and fall of Hox gene clusters. *Development* 134:2549–2560.
https://doi.org/10.1242/dev.001065.

Edgar RC. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* 32:1792–1797.
https://doi.org/10.1093/nar/gkh340.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biol.* 20:238. https://doi.org/10.1186/s13059-
019-1832-y.

Escuer P, Pisarenco VA, Fernández-Ruiz AA, Vizueta J, Sánchez-Herrero JF, Arnedo
MA, Sánchez-Gracia A, Rozas J. 2022. The chromosome-scale assembly of the
Canary Islands endemic spider Dysdera silvatica (Arachnida, Araneae) sheds
light on the origin and genome structure of chemoreceptor gene families in
chelicerates. *Mol. Ecol. Resour.* 22:375–390. https://doi.org/10.1111/1755-
0998.13471.

Espinosa-Cantú A, Ascencio D, Barona-Gómez F, De Luna A. 2015. Gene duplication
and the evolution of moonlighting proteins. *Front. Genet.* 6:227.
https://doi.org/10.3389/fgene.2015.00227.

Fan Z, Yuan T, Liu P, Wang LY, Jin JF, Zhang F, Zhang ZS. 2021. A chromosome-level
genome of the spider Trichonephila antipodiana reveals the genetic basis of its
polyphagy and evidence of an ancient whole-genome duplication event.
GigaScience 10:1–15. https://doi.org/10.1093/GIGASCIENCE/GIAB016.

Fernández R, Edgecombe GD, Giribet G. 2018. Phylogenomics illuminates the
backbone of the Myriapoda Tree of Life and reconciles morphological and
molecular phylogenies. *Sci. Rep.* 8:83. https://doi.org/10.1038/s41598-017-
18562-w.

Fernández R, Kallal RJ, Dimitrov D, Ballesteros JA, Arnedo MA, Giribet G, Hormiga G.
2018. Phylogenomics, Diversification Dynamics, and Comparative
Transcriptomics across the Spider Tree of Life. *Curr. Biol.* 28:1489-1497.e5.
https://doi.org/10.1016/j.cub.2018.03.064.

Ferrier DEK. 2016a. The origin of the Hox/ParaHox genes, the Ghost Locus hypothesis
and the complexity of the first animal. *Brief. Funct. Genomics* 15:333–341.
https://doi.org/10.1093/bfgp/elv056.

Ferrier DEK. 2016b. Evolution of Homeobox Gene Clusters in Animals: The Giga-
Cluster and Primary vs. Secondary Clustering. *Front. Ecol. Evol.* 4:1–13.
https://doi.org/10.3389/fevo.2016.00036.

Ferrier DEK. 2019. Space and time in Hox/ParaHox gene cluster evolution. In: Fusco G,
editor. Perspectives on Evolutionary and Developmental Biology: Essays for
Alessandro Minelli. Padova: Padova University Press. p. 245–274.

FigTree. 2018. FigTree v1.4.4. Available from: http://tree.bio.ed.ac.uk/software/figtree/

Flot J-F, Hespeels B, Li X, Noel B, Arkhipova I, Danchin EGJ, Hejnol A, Henrissat B,
Koszul R, Aury J-M, et al. 2013. Genomic evidence for ameiotic evolution in the

bdelloid rotifer Adineta vaga. *Nature* 500:453–457.
https://doi.org/10.1038/nature12326.

Force A, Lynch M, Pickett FB, Amores A, Yan Y, Postlethwait J. 1999. Preservation of
Duplicate Genes by Complementary, Degenerate Mutations. *Genetics*
151:1531–1545.

Gainett G, Crawford AR, Klementz BC, So C, Baker CM, Setton EVW, Sharma PP.
2022. Eggs to long-legs: embryonic staging of the harvestman Phalangium opilio
(Opiliones), an emerging model arachnid. *Front. Zool.* 19:11.
https://doi.org/10.1186/s12983-022-00454-z.

Gainett G, González VL, Ballesteros JA, Setton EVW, Baker CM, Barolo Gargiulo L,
Santibáñez-López CE, Coddington JA, Sharma PP. 2021. The genome of a
daddy-long-legs (Opiliones) illuminates the evolution of arachnid appendages.
Proc. R. Soc. B Biol. Sci. 288:20211168. https://doi.org/10.1098/rspb.2021.1168.

Gallardo ME, Lopez-Rios J, Fernaud-Espinosa I, Granadino B, Sanz R, Ramos C,
Ayuso C, Seller MJ, Brunner HG, Bovolenta P, et al. 1999. Genomic cloning and
characterization of the human homeobox gene SIX6 reveals a cluster of SIX
genes in chromosome 14 and associates SIX6 hemizygosity with bilateral
anophthalmia and pituitary anomalies. *Genomics* 61:82–91.
https://doi.org/10.1006/geno.1999.5916.

Garcia-Fernàndez J. 2005. The genesis and evolution of homeobox gene clusters. *Nat.*
Rev. Genet. 6:881–892. https://doi.org/10.1038/nrg1723.

Goldman N, Yang Z. 1994. A codon-based model of nucleotide substitution for protein-
coding DNA sequences. *Mol. Biol. Evol.* 11:725–736.
https://doi.org/10.1093/oxfordjournals.molbev.a040153.

Harper A, Baudouin Gonzalez L, Schönauer A, Janssen R, Seiter M, Holzem M, Arif S,
McGregor AP, Sumner-Rooney L. 2021. Widespread retention of ohnologs in key
developmental gene families following whole-genome duplication in
arachnopulmonates.Lott S, editor. *G3 GenesGenomesGenetics*
11:2020.07.10.177725. https://doi.org/10.1093/g3journal/jkab299.

Herndon N, Shelton J, Gerischer L, Ioannidis P, Ninova M, Dönitz J, Waterhouse RM,
Liang C, Damm C, Siemanowski J, et al. 2020. Enhanced genome assembly and
a new official gene set for Tribolium castaneum. *BMC Genomics* 21:47.
https://doi.org/10.1186/s12864-019-6394-6.

Hoegg S, Meyer A. 2005. Hox clusters as models for vertebrate genome evolution.
Trends Genet. 21:421–424. https://doi.org/10.1016/j.tig.2005.06.004.

Holland LZ, Ocampo Daza D. 2018. A new look at an old question: when did the second
whole genome duplication occur in vertebrate evolution? *Genome Biol.* 19:209.
https://doi.org/10.1186/s13059-018-1592-0.

Holland PWH. 2013. Evolution of homeobox genes. *Wiley Interdiscip. Rev. Dev. Biol.*
2:31–45. https://doi.org/10.1002/wdev.78.

Holland PWH. 2015. Did homeobox gene duplications contribute to the Cambrian
explosion? *Zool. Lett.* 1:1. https://doi.org/10.1186/s40851-014-0004-x.

Holland PWH, Booth HAF, Bruford EA. 2007. Classification and nomenclature of all
human homeobox genes. *BMC Biol.* 5:47. https://doi.org/10.1186/1741-7007-5-
47.

Holland PWH, Garcia-Fernàndez J. 1996. HoxGenes and Chordate Evolution. *Dev.*
Biol. 173:382–395. https://doi.org/10.1006/dbio.1996.0034.

Holland PWH, Marlétaz F, Maeso I, Dunwell TL, Paps J. 2017. New genes from old:
asymmetric divergence of gene duplicates and the evolution of development.
Philos. Trans. R. Soc. B Biol. Sci. 372:20150480.
https://doi.org/10.1098/rstb.2015.0480.

Hu W, Jia A, Ma S, Zhang G, Wei Z, Lu F, Luo Y, Zhang Z, Sun J, Yang T, et al. 2022.
Molecular atlas reveals the tri-sectional spinning mechanism of spider dragline
silk. :2022.06.21.496984. https://doi.org/10.1101/2022.06.21.496984. Available
from: https://www.biorxiv.org/content/10.1101/2022.06.21.496984v1

Hui JHL, McDougall C, Monteiro AS, Holland PWH, Arendt D, Balavoine G, Ferrier
DEK. 2012. Extensive Chordate and Annelid Macrosynteny Reveals Ancestral
Homeobox Gene Organization. *Mol. Biol. Evol.* 29:157–165.
https://doi.org/10.1093/molbev/msr175.

Irimia M, Maeso I, Garcia-Fernàndez J. 2008. Convergent Evolution of Clustering of
Iroquois Homeobox Genes across Metazoans. *Mol. Biol. Evol.* 25:1521–1525.
https://doi.org/10.1093/molbev/msn109.

Irimia M, Tena JJ, Alexis MS, Fernandez-Miñan A, Maeso I, Bogdanovic O, de la Calle-
Mustienes E, Roy SW, Gómez-Skarmeta JL, Fraser HB. 2012. Extensive
conservation of ancient microsynteny across metazoans due to cis-regulatory
constraints. *Genome Res.* 22:2356–2367. https://doi.org/10.1101/gr.139725.112.

Jagla K, Bellard M, Frasch M. 2001. A cluster of Drosophila homeobox genes involved
in mesoderm differentiation programs. *BioEssays* 23:125–133.
https://doi.org/10.1002/1521-1878.

Janssen R. 2017. Comparative analysis of gene expression patterns in the arthropod
labrum and the onychophoran frontal appendages, and its implications for the

1177  arthropod head problem. *EvoDevo* 8:1. https://doi.org/10.1186/s13227-016-0064-
1178  4.

1179 Janssen R, Andersson E, Betnér E, Bijl S, Fowler W, Höök L, Leyhr J, Mannelqvist A,
1180  Panara V, Smith K, et al. 2018. Embryonic expression patterns and phylogenetic
1181  analysis of panarthropod sox genes: insight into nervous system development,
1182  segmentation and gonadogenesis. *BMC Evol. Biol.* 18:88.
1183  https://doi.org/10.1186/s12862-018-1196-z.

1184 Janssen R, Damen WGM. 2008. Diverged and conserved aspects of heart formation in
1185  a spider. *Evol. Dev.* 10:155–165. https://doi.org/10.1111/j.1525-
1186  142X.2008.00223.x.

1187 Janssen R, Eriksson BJ, Tait NN, Budd GE. 2014. Onychophoran Hox genes and the
1188  evolution of arthropod Hox gene expression. *Front. Zool.* 11:22.
1189  https://doi.org/10.1186/1742-9994-11-22.

1190 Janssen R, Pechmann M. 2023. Expression of posterior Hox genes and opisthosomal
1191  appendage development in a mygalomorph spider. *Dev. Genes Evol.* [Internet].
1192  https://doi.org/10.1007/s00427-023-00707-9. Available from:
1193  https://doi.org/10.1007/s00427-023-00707-9

1194 Janssen R, Pechmann M, Turetzek N. 2021. A chelicerate Wnt gene expression atlas:
1195  novel insights into the complexity of arthropod Wnt-patterning. *EvoDevo* 12:12.
1196  https://doi.org/10.1186/s13227-021-00182-1.

1197 Jiménez-Delgado S, Pascual-Anaya J, Garcia-Fernàndez J. 2009. Implications of
1198  duplicated cis-regulatory elements in the evolution of metazoans: the DDI model
1199  or how simplicity begets novelty. *Brief. Funct. Genomic. Proteomic.* 8:266–275.
1200  https://doi.org/10.1093/bfgp/elp029.

1201 Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
1202  ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat.*
1203  *Methods* 14:587–589. https://doi.org/10.1038/nmeth.4285.

1204 Kawakami K, Sato S, Ozaki H, Ikeda K. 2000. Six family genes-structure and function
1205  as transcription factors and their roles in development. *BioEssays* 22:616–626.
1206  https://doi.org/10.1002/1521-1878(200007)22:7<616::AID-BIES4>3.0.CO;2-R.

1207 Kenny NJ, Chan KW, Nong W, Qu Z, Maeso I, Yip HY, Chan TF, Kwan HS, Holland
1208  PWH, Chu KH, et al. 2016. Ancestral whole-genome duplication in the marine
1209  chelicerate horseshoe crabs. *Heredity* 116:190–199.
1210  https://doi.org/10.1038/hdy.2015.89.

1211 Kerner P, Ikmi A, Coen D, Vervoort M. 2009. Evolutionary history of the iroquois/Irx
1212  genes in metazoans. *BMC Evol. Biol.* 9:74. https://doi.org/10.1186/1471-2148-9-
1213  74.

Kmita M, Duboule D. 2003. Organizing Axes in Time and Space; 25 Years of Colinear
Tinkering. *Science* 301:331–333. https://doi.org/10.1126/science.1085753.

Kořínková T, Král J. 2013. Karyotypes, Sex Chromosomes, and Meiotic Division in
Spiders. In: Nentwig W, editor. Spider Ecophysiology. Berlin, Heidelberg:
Springer. p. 159–171. Available from: https://doi.org/10.1007/978-3-642-33989-
9_12

Kosakovsky Pond SL, Poon AFY, Velazquez R, Weaver S, Hepler NL, Murrell B, Shank
SD, Magalis BR, Bouvier D, Nekrutenko A, et al. 2020. HyPhy 2.5—A
Customizable Platform for Evolutionary Hypothesis Testing Using Phylogenies.
Mol. Biol. Evol. 37:295–299. https://doi.org/10.1093/molbev/msz197.

Krumlauf R. 2018. Hox genes, clusters and collinearity. *Int. J. Dev. Biol.* 62:659–663.
https://doi.org/10.1387/ijdb.180330rr.

Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra
MA. 2009. Circos: an information aesthetic for comparative genomics. *Genome*
Res. 19:1639–1645. https://doi.org/10.1101/gr.092759.109.

Kumar JP. 2009. The sine oculis homeobox (SIX) family of transcription factors as
regulators of development and disease. *Cell. Mol. Life Sci.* 66:565–583.
https://doi.org/10.1007/s00018-008-8335-4.

Larroux C, Fahey B, Degnan SM, Adamski M, Rokhsar DS, Degnan BM. 2007. The NK
Homeobox Gene Cluster Predates the Origin of Hox Genes. *Curr. Biol.* 17:706–
710. https://doi.org/10.1016/j.cub.2007.03.008.

Larroux C, Luke GN, Koopman P, Rokhsar DS, Shimeld SM, Degnan BM. 2008.
Genesis and Expansion of Metazoan Transcription Factor Gene Classes. *Mol.*
Biol. Evol. 25:980–996. https://doi.org/10.1093/molbev/msn047.

Leite DJ, Baudouin-Gonzalez L, Iwasaki-Yokozawa S, Lozano-Fernandez J, Turetzek
N, Akiyama-Oda Y, Prpic N-M, Pisani D, Oda H, Sharma PP, et al. 2018.
Homeobox Gene Duplication and Divergence in Arachnids. *Mol. Biol. Evol.*
35:2240–2253. https://doi.org/10.1093/molbev/msy125.

Leite DJ, Ninova M, Hilbrant M, Arif S, Griffiths-Jones S, Ronshaugen M, McGregor AP.
2016. Pervasive microRNA Duplication in Chelicerates: Insights from the
Embryonic microRNA Repertoire of the Spider Parasteatoda tepidariorum.
Genome Biol. Evol. 8:2133–2144. https://doi.org/10.1093/gbe/evw143.

Luke GN, Castro LFC, McLay K, Bird C, Coulson A, Holland PWH. 2003. Dispersal of
NK homeobox gene clusters in amphioxus and humans. *Proc. Natl. Acad. Sci.*
100:5292–5295. https://doi.org/10.1073/pnas.0836141100.

Maddison WP, Leduc-Robert G. 2013. MULTIPLE ORIGINS OF SEX CHROMOSOME
FUSIONS CORRELATED WITH CHIASMA LOCALIZATION IN HABRONATTUS
JUMPING SPIDERS (ARANEAE: SALTICIDAE). *Evolution* 67:2258–2272.
https://doi.org/10.1111/evo.12109.

Maqbool T, Soler C, Jagla T, Daczewska M, Lodha N, Palliyil S, VijayRaghavan K,
Jagla K. 2006. Shaping Leg Muscles in Drosophila: Role of ladybird, a
Conserved Regulator of Appendicular Myogenesis. *PLOS ONE* 1:e122.
https://doi.org/10.1371/journal.pone.0000122.

Marlétaz F, Firbas PN, Maeso I, Tena JJ, Bogdanovic O, Perry M, Wyatt CDR, de la
Calle-Mustienes E, Bertrand S, Burguera D, et al. 2018. Amphioxus functional
genomics and the origins of vertebrate gene regulation. *Nature* 564:64–70.
https://doi.org/10.1038/s41586-018-0734-6.

Martin A, Serano JM, Jarvis E, Bruce HS, Wang J, Ray S, Barker CA, O'Connell LC,
Patel NH. 2016. CRISPR/Cas9 Mutagenesis Reveals Versatile Roles of Hox
Genes in Crustacean Limb Specification and Evolution. *Curr. Biol.* 26:14–26.
https://doi.org/10.1016/j.cub.2015.11.021.

Mazza ME, Pang K, Reitzel AM, Martindale MQ, Finnerty JR. 2010. A conserved cluster
of three PRD-class homeobox genes (homeobrain, rx and orthopedia) in the
Cnidaria and Protostomia. *EvoDevo* 1:3. https://doi.org/10.1186/2041-9139-1-3.

McLysaght A, Hokamp K, Wolfe KH. 2002. Extensive genomic duplication during early
chordate evolution. *Nat. Genet.* 31:200–204. https://doi.org/10.1038/ng884.

Meyer A, Schartl M. 1999. Gene and genome duplications in vertebrates: the one-to-
four (-to-eight in fish) rule and the evolution of novel gene functions. *Curr. Opin.*
Cell Biol. 11:699–704. https://doi.org/10.1016/S0955-0674(99)00039-3.

Miller J, Zimin AV, Gordus A. 2023. Chromosome-level genome and the identification of
sex chromosomes in Uloborus diversus. *GigaScience* 12:giad002.
https://doi.org/10.1093/gigascience/giad002.

Mittmann B, Wolff C. 2012. Embryonic development and staging of the cobweb spider
Parasteatoda tepidariorum C. L. Koch, 1841 (syn.: Achaearanea tepidariorum;
Araneomorphae; Theridiidae). *Dev. Genes Evol.* 222:189–216.
https://doi.org/10.1007/s00427-012-0401-0.

Muffato M, Louis A, Nguyen NTT, Lucas J, Berthelot C, Roest Crollius H. 2023.
Reconstruction of hundreds of reference ancestral genomes across the
eukaryotic kingdom. *Nat. Ecol. Evol.* 7:355–366. https://doi.org/10.1038/s41559-
022-01956-z.

Mulhair P, Crowley L, Boyes DH, Harper A, Lewis OT, Consortium DT of L, Holland
PWH. 2022. Diversity, duplication, and genomic organization of homeobox genes

1286    in Lepidoptera. *Genome Res.*:gr.277118.122.
1287    https://doi.org/10.1101/gr.277118.122.

1288 Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and
1289    Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
1290    *Mol. Biol. Evol.* 32:268–274. https://doi.org/10.1093/molbev/msu300.

1291 Nong W, Qu Z, Li Y, Barton-Owen T, Wong AYP, Yip HY, Lee HT, Narayana S, Baril T,
1292    Swale T, et al. 2021. Horseshoe crab genomes reveal the evolution of genes and
1293    microRNAs after three rounds of whole genome duplication. *Commun. Biol.* 4:83.
1294    https://doi.org/10.1038/s42003-020-01637-2.

1295 Nossa CW, Havlak P, Yue J-X, Lv J, Vincent KY, Brockmann HJ, Putnam NH. 2014.
1296    Joint assembly and genetic mapping of the Atlantic horseshoe crab genome
1297    reveals ancient whole genome duplication. *GigaScience* 3:2047-217X-3–9.
1298    https://doi.org/10.1186/2047-217X-3-9.

1299 Ohno S. 1970. Evolution by Gene Duplication. Berlin, Heidelberg: Springer Berlin
1300    Heidelberg Available from: http://link.springer.com/10.1007/978-3-642-86659-3

1301 Pollard SL, Holland PWH. 2000. Evidence for 14 homeobox gene clusters in human
1302    genome ancestry. *Curr. Biol.* 10:1059–1062. https://doi.org/10.1016/S0960-
1303    9822(00)00676-X.

1304 Pond SLK, Frost SDW, Muse SV. 2005. HyPhy: hypothesis testing using phylogenies.
1305    *Bioinformatics* 21:676–679. https://doi.org/10.1093/bioinformatics/bti079.

1306 Prpic N-M, Schoppmeier M, Damen WGM. 2008. Collection and Fixation of Spider
1307    Embryos. *Cold Spring Harb. Protoc.* 2008:pdb.prot5067.
1308    https://doi.org/10.1101/pdb.prot5067.

1309 Putnam NH, Butts T, Ferrier DEK, Furlong RF, Hellsten U, Kawashima T, Robinson-
1310    Rechavi M, Shoguchi E, Terry A, Yu J-K, et al. 2008. The amphioxus genome
1311    and the evolution of the chordate karyotype. *Nature* 453:1064–1071.
1312    https://doi.org/10.1038/nature06967.

1313 Sánchez-Herrero JF, Frías-López C, Escuer P, Hinojosa-Alvarez S, Arnedo MA,
1314    Sánchez-Gracia A, Rozas J. 2019. The draft genome sequence of the spider
1315    Dysdera silvatica (Araneae, Dysderidae): A valuable resource for functional and
1316    evolutionary genomic studies in chelicerates. *GigaScience* 8:giz099.
1317    https://doi.org/10.1093/gigascience/giz099.

1318 Saudemont A, Dray N, Hudry B, Le Gouar M, Vervoort M, Balavoine G. 2008.
1319    Complementary striped expression patterns of NK homeobox genes during
1320    segment formation in the annelid Platynereis. *Dev. Biol.* 317:430–443.
1321    https://doi.org/10.1016/j.ydbio.2008.02.013.

Schinko JB, Kreuzer N, Offen N, Posnien N, Wimmer EA, Bucher G. 2008. Divergent functions of orthodenticle, empty spiracles and buttonhead in early head patterning of the beetle Tribolium castaneum (Coleoptera). *Dev. Biol.* 317:600–613. https://doi.org/10.1016/j.ydbio.2008.03.005.

Schomburg C, Turetzek N, Schacht MI, Schneider J, Kirfel P, Prpic N-M, Posnien N. 2015. Molecular characterization and embryonic origin of the eyes in the common house spider Parasteatoda tepidariorum. *EvoDevo* 6:15. https://doi.org/10.1186/s13227-015-0011-9.

Schwager EE, Schoppmeier M, Pechmann M, Damen WGM. 2007. Duplicated Hox genes in the spider Cupiennius salei. *Front. Zool.* 4:10. https://doi.org/10.1186/1742-9994-4-10.

Schwager EE, Sharma PP, Clarke T, Leite DJ, Wierschin T, Pechmann M, Akiyama-Oda Y, Esposito L, Bechsgaard J, Bilde T, et al. 2017. The house spider genome reveals an ancient whole-genome duplication during arachnid evolution. *BMC Biol.* 15:62. https://doi.org/10.1186/s12915-017-0399-x.

Serano JM, Martin A, Liubicich DM, Jarvis E, Bruce HS, La K, Browne WE, Grimwood J, Patel NH. 2016. Comprehensive analysis of Hox gene expression in the amphipod crustacean Parhyale hawaiensis. *Dev. Biol.* 409:297–309. https://doi.org/10.1016/j.ydbio.2015.10.029.

Shao L, Li S. 2018. Early Cretaceous greenhouse pumped higher taxa diversification in spiders. *Mol. Phylogenet. Evol.* 127:146–155. https://doi.org/10.1016/j.ympev.2018.05.026.

Sharma PP, Schwager EE, Extavour CG, Giribet G. 2012. Hox gene expression in the harvestman Phalangium opilio reveals divergent patterning of the chelicerate opisthosoma. *Evol. Dev.* 14:450–463. https://doi.org/10.1111/j.1525-142X.2012.00565.x.

Sheffer MM, Hoppe A, Krehenwinkel H, Uhl G, Kuss AW, Jensen L, Jensen C, Gillespie RG, Hoff KJ, Prost S. 2021. Chromosome-level reference genome of the European wasp spider Argiope bruennichi: a resource for studies on range expansion and evolutionary adaptation. *GigaScience* 10:1–12. https://doi.org/10.1093/GIGASCIENCE/GIAA148.

Shimeld SM, Holland PWH. 2000. Vertebrate innovations. *Proc. Natl. Acad. Sci. U. S. A.* 97:4449–4452. https://doi.org/10.1073/pnas.97.9.4449.

Shingate P, Ravi V, Prasad A, Tay BH, Garg KM, Chattopadhyay B, Yap LM, Rheindt FE, Venkatesh B. 2020. Chromosome-level assembly of the horseshoe crab genome provides insights into its genome evolution. *Nat. Commun.* 11:1–13. https://doi.org/10.1038/s41467-020-16180-1.

Simakov O, Kawashima T. 2017. Independent evolution of genomic characters during
major metazoan transitions. *Dev. Biol.* 427:179–192.
https://doi.org/10.1016/j.ydbio.2016.11.012.

Simakov O, Kawashima T, Marlétaz F, Jenkins J, Koyanagi R, Mitros T, Hisata K,
Bredeson J, Shoguchi E, Gyoja F, et al. 2015. Hemichordate genomes and
deuterostome origins. *Nature* 527:459–465. https://doi.org/10.1038/nature16150.

Simakov O, Marlétaz F, Yue J, O'Connell B, Jenkins J, Brandt A, Calef R, Tung C,
Huang T, Schmutz J, et al. 2020. Deeply conserved synteny resolves early
events in vertebrate evolution. *Nat. Ecol. Evol.* 4:820–830.
https://doi.org/10.1038/s41559-020-1156-z.

Simonnet F, Célérier M-L, Quéinnec E. 2006. Orthodenticle and empty spiracles genes
are expressed in a segmental pattern in chelicerates. *Dev. Genes Evol.* 216:467–
480. https://doi.org/10.1007/s00427-006-0093-4.

Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, Kosakovsky Pond SL. 2015.
Less is more: an adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. *Mol. Biol. Evol.* 32:1342–1353.
https://doi.org/10.1093/molbev/msv022.

So WL, Nong W, Xie Y, Baril T, Ma H, Qu Z, Haimovitz J, Swale T, Gaitan-Espitia JD,
Lau KF, et al. 2022. Myriapod genomes reveal ancestral horizontal gene transfer
and hormonal gene loss in millipedes. *Nat. Commun.* 13:3010.
https://doi.org/10.1038/s41467-022-30690-0.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. *Nucleic Acids Res.*
34:W609–W612. https://doi.org/10.1093/nar/gkl315.

Tinti M, Johnson C, Toth R, Ferrier DEK, Mackintosh C. 2012. Evolution of signal
multiplexing by 14-3-3-binding 2R-ohnologue protein families in the vertebrates.
Open Biol. 2:120103. https://doi.org/10.1098/rsob.120103.

Treffkorn S, Kahnke L, Hering L, Mayer G. 2018. Expression of NK cluster genes in the
onychophoran Euperipatoides rowelli: implications for the evolution of NK family
genes in nephrozoans. *EvoDevo* 9:17. https://doi.org/10.1186/s13227-018-0105-
2.

Turetzek N, Pechmann M, Janssen R, Prpic N-M. 2022. Hox genes in spiders: Their
significance for development and evolution. *Semin. Cell Dev. Biol.*:in press.
https://doi.org/10.1016/j.semcdb.2022.11.014.

Walldorf U, Gehring W j. 1992. Empty spiracles, a gap gene containing a homeobox
involved in Drosophila head development. *EMBO J.* 11:2247–2259.
https://doi.org/10.1002/j.1460-2075.1992.tb05284.x.

Wang S, Zhang J, Jiao W, Li J, Xun X, Sun Y, Guo X, Huan P, Dong B, Zhang L, et al.
2017. Scallop genome provides insights into evolution of bilaterian karyotype and
development. *Nat. Ecol. Evol.* 1:1–12. https://doi.org/10.1038/s41559-017-0120.

Wang Y, Tang H, DeBarry JD, Tan X, Li J, Wang X, Lee T, Jin H, Marler B, Guo H, et
al. 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene
synteny and collinearity. *Nucleic Acids Res.* 40:e49.
https://doi.org/10.1093/nar/gkr1293.

Wang Z, Zhu K, Li H, Gao L, Huang H, Ren Y, Xiang H. 2022. Chromosome-level
genome assembly of the black widow spider Latrodectus elegans illuminates
composition and evolution of venom and silk proteins. *GigaScience* 11:1–11.
https://doi.org/10.1093/GIGASCIENCE/GIAC049.

Wotton KR, Weierud FK, Juárez-Morales JL, Alvares LE, Dietrich S, Lewis KE. 2009.
Conservation of gene linkage in dispersed vertebrate NK homeobox clusters.
Dev. Genes Evol. 219:481–496. https://doi.org/10.1007/s00427-009-0311-y.

Zhong YF, Butts T, Holland PWH. 2008. HomeoDB: A database of homeobox gene
diversity. *Evol. Dev.* 10:516–518. https://doi.org/10.1111/j.1525-
142X.2008.00266.x.

Zhong YF, Holland PWH. 2011. HomeoDB2: Functional expansion of a comparative
homeobox gene database for evolutionary developmental biology. *Evol. Dev.*
13:567–568. https://doi.org/10.1111/j.1525-142X.2011.00513.x.

Zhu B, Jin P, Hou Z, Li J, Wei S, Li S. 2022. Chromosomal-level genome of a sheet-
web spider provides insight into the composition and evolution of venom. *Mol.
Ecol. Resour.* 22:2333–2348. https://doi.org/10.1111/1755-0998.13601.

Zhu B, Jin P, Zhang Y, Shen Y, Wang W, Li S. 2023. Genomic and transcriptomic
analyses support a silk gland origin of spider venom glands. *BMC Biol.* 21:82.
https://doi.org/10.1186/s12915-023-01581-7.

1422