Convergent evolution of noxious heat sensing by TRPA5, a novel class of heat sensor in *Rhodnius prolixus*

Marjorie A. Liénard[1,2,3*], David Baez-Nieto[4*], Cheng-Chia Tsai[5], Wendy A. Valencia-Montoya[2], Balder

Werin[6], Urban Johanson[6], Jean-Marc Lassance[2,7], Jen Q. Pan[4], Nanfang Yu[5], Naomi E. Pierce[2]

[1]Department of Biology, Lund University, Lund, Sweden

[2]Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology, Harvard University,
Cambridge, MA 02138, USA

[3]Broad Institute, Cambridge, MA 02142, USA

[4]Stanley Center for Psychiatric Research, Broad Institute, Cambridge, MA 02142, USA

[5]Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027, USA

[6]Division of Biochemistry and Structural Biology, Department of Chemistry, Lund University, Lund, Sweden

[7]GIGA-Research, University of Liège, Liège, Belgium

[*]These authors contributed equally to the work

ORCIDs:

MAL:0000-0003-3193-3666

DBN:0000-0002-2238-8486

CCT:0000-0001-5662-6908

WAVM:0000-0001-9246-2330

BW:0009-0006-1188-3496

UJ: 0000-0001-8862-128X

JML: 0000-0002-3675-6956

JQP: 0000-0003-0767-086X

NY:0000-0002-9462-4724

NEP:0000-0003-3366-1625

Correspondence:

marjorie.lienard@biol.lu.se

Keywords: transient receptor potential channel, temperature, noxious heat, insect vector, patch-clamp

recordings, triatomine, Chagas disease, convergent evolution

ABSTRACT

Insects are ectotherms, and as such, rely on a diverse repertoire of thermoreceptors to monitor environmental temperature and control behavioral thermoregulation. Here, we use structural, phylogenetic, genetic, and functional analyses to show that TRPA5 genes, widespread across numerous insect orders, encode a novel class of noxious heat receptors. We show that in the triatomine bug *Rhodnius prolixus*, the primary vector of Chagas disease, Rp-TRPA5$_2$ differs biophysically and structurally from noxious thermoTRPAs previously described in insects. This includes key changes in the ankyrin repeat domain and the selectivity filter of the channel. *In vitro*, we find evidence that the homo-tetrameric channel is not activated by voltage, but displays high thermosensitivity with an enthalpy change (ΔH) of 72 kcal/mol associated with the channel activation, with a Q_{10} = 25 and $T°_{half}$ = 58.6°C. Structural analyses reveal parallels in the overall ion channel architecture between fruit fly TRPA1 and Rp-TRPA5$_2$; however, functional properties and expression patterns indicate that the role of Rp-TRPA5$_2$ is more similar to that of *Pyrexia* noxious heat receptors found in fruit flies. *Pyrexia* genes have been lost in true bugs, and our findings suggest that the rapidly evolving insect TRPA gene family has given rise to an independent evolutionary origin of a molecular transducer that is responsive to noxious thermal stimuli.

Introduction

Animal thermosensation is critical for performance in fluctuating environments. Changes in environmental temperature are transduced by the sensory system as part of physiological feedback controlling responses such as metabolic homeostasis, feeding, finding suitable habitats, and extreme-heat avoidance [1, 2]. At the molecular level, thermal perception is mediated by the temperature-dependent activation of specific cold- and heat-activated receptors [3, 4]. Although families such as ionotropic receptors (IRs) and gustatory receptors (GRs) have been linked to peripheral innocuous thermosensation in insects [3-6], the transient receptor potential (TRP) receptor family encodes the greatest diversity of thermosensitive channels. TRP receptors are remarkably diverse (TRPA, TRPC, TRPN, TRPM, TRPML and TRPV) and play salient roles as polymodal ion channels responding to chemical, mechanical, and thermal stimuli [7-12].

Mammalian TRP channels involved in temperature detection (thermoTRPs) belong to the TRPA, TRPV and TRPM subfamilies (Table 1) and are activated by temperatures from noxious cold to noxious heat [4, 9, 13-16]. In invertebrates, known thermoTRP channels have so far been restricted to the ankyrin TRPA subfamily of genes including *Painless*, *Pyrexia*, *TRPA1*, and *Hs-TRPA* (Fig. 1; Fig. S1) [11, 12]. In *Drosophila melanogaster, Painless*, *Pyrexia* and *Dm-TRPA1* isoforms A, B and D encode receptors that exhibit distinct biophysical properties, cellular expression patterns and temperature activation thresholds ranging from 19°C to 46°C [17-24]. TRPA1 is also a heat–activated TRP sensor in *Anopheles gambiae* (25-37°C), and other mosquitos [25, 26], playing a key role in tuning heat-seeking behavior. Outside the Diptera, TRPA1 has been characterized as a heat-sensitive channel in other insects as it is known to regulate the induction of embryonic diapause in *Bombyx mori* at temperatures above 21°C [27]. The subfamily *Waterwitch* includes receptors responding to stimuli in different modalities from ancestral hygrosensation found in fruit flies [20] to derived heat sensing exhibited by hymenopterans and mediated by the Hs-TRPA subfamily, which diverged following a duplication from *Waterwitch* [12]. Thus, despite the loss of TRPA1 in Hymenoptera, in honeybees, *Apis mellifera* Am-HsTRPA responds to temperatures around 34°C, in parasitoid wasps, *Nasonia*

vitripennis Nv-HsTRPA activates in response to small temperature differences in the range 8°C to

44°C regardless of initial temperatures, and in fire ants, *Solenopsis invicta,* Si-HsTRPA is activated in

the range 28-37°C [28, 29]. Notably, the insect TRP ankyrin family has an additional subfamily of

unknown function, TRPA5, which is seemingly absent from the fruit fly genome yet found across

several other orders of insects [11].

Here we deorphanize and characterize an ankyrin TRPA5 ion channel from the "kissing" bug,

Rhodnius prolixus. Long used as a model organism in studies of insect development and physiology

[30], *R. prolixus* (Hemiptera, Reduviidae: Triatominae) has become increasingly relevant for molecular

and functional studies. This is primarily explained by its long-term medical and societal impact as a

haematophagous vector of *Trypanosoma cruzi*, the causative agent of Chagas' disease [31]. Due to

the progressive adaptation of wild triatomine vector species to domestic environments, vector

transmission to human populations has increased in recent years [32, 33]. The disease currently

affects over 8 million people worldwide, with vector transmission causing around 30,000 new cases

yearly [32, 34, 35]. Extensive long-term efforts towards decoding the sensory ecology of triatomines

[36-38] have identified olfactory, thermal and environmentally-mediated cues as well as the

neuroethology underlying its complex host-seeking behavior [38-45]. Moreover, the annotated *R.*

prolixus genome [37] and recent transcriptomic studies [44-46] provide detailed expression profiles of

candidate sensory receptor genes, including olfactory, ionotropic, pickpocket, and transient receptor

potential receptors that can be used to probe the genetic basis of sensory traits [47, 48].

In this study, we leverage the rich genomic and transcriptomic resources available for *R.*

prolixus along with molecular, structural modeling and functional approaches to characterize a TRPA5

ion channel enriched in canonical sensory tissues. The biophysical properties of the ion channel

demonstrate that *TRPA5* encodes a novel ankyrin type of heat-activated TRP receptor responding to

noxious temperatures. Analyses of predicted structures reveal that the channel displays unique

features among the ankyrin family, potentially affecting its ion conduction properties despite sharing

conserved structural domains with other ankyrin thermoTRPs. These findings may facilitate future

studies of agonist responses of TRPA5 to assist in the development of novel genetic tools for vector

control efforts.

Table 1. Vertebrate and invertebrate TRP ion channels involved in thermal transduction.

Nomenclature	Other names	Organism	Expression	TRP family	Thermal treshold*	Activity range	Entropy change (ΔS) cal/mol*K	Activation enthalpy (ΔH) kcal/mol	Q10**	Reference
Vertebrate transient receptor potential (TRP) channels involved in thermosensation										
TRPV1	Vr1	*Rattus norvegicus* (rat), *Homo sapiens* (human)	nociceptor neurons	Ankyrin	≥ 42℃ (40-45)		260 (at 30mV)	90-100	16.8, 20*	Caterina et al 1997, Cheng et al 2012*, Yao et al 2010*
TRPV1(l)	TRPV1- long	*Desmodus rotundus* (vampire bat)	DRG and TG neurons	Ankyrin	40℃				12	Gracheva et al. 2011
TRPV1(s)	TRPV1- short	*Desmodus rotundus* (vampire bat)	TG neurons	Ankyrin	30℃				5	Gracheva et al. 2011
TRPV2	Vrl2	*Rattus norvegicus* (rat), *Homo sapiens* (human)	nociceptor neurons	Ankyrin	≥ 52℃ (50-53)		586	200	20.6	Caterina et al 1999
hTrpv3	Vrl3	*Homo sapiens* (human)	skin keratinocytes	Ankyrin	≥ 33℃ (34-38℃)	33℃~50℃ (Max 41-47℃)	n.d		6.62	Peier et al 2002; Smith et al 2002
hTrpv4	OTRPC4, VR-OAC, Trp12, Vrl2	*Homo sapiens* (human)	skin and dorsal root ganglion neurons	Ankyrin	~ 27-34℃		n.d		9.9+/-3.8	Güler et al 2002; Watanabe et al 2002
TRPM8	CMR1			Melastatin	≤ 23-25℃	0℃-25℃	-384	-112	24	McKemy et al 2002; Brauchi et al 2004;
TRPM3		*Mus musculus* (mouse)	DRG and TG sensory neurons	Melastatin	≥ 43℃				7.2	Vriens et al 2011
TRPC5		*Mus musculus* (mouse), *Homo sapiens* (human)	DRG neurons	Canonical	<37℃->25℃		n.d.	-40		Zimmermann et al 2011
Anktm1	TRPA1	*Mus musculus* (mouse), *Homo sapiens* (human)	DRG neurons	Ankyrin	≤ 17℃	0℃-17℃	-140	-40	6	Story et al 2003
Chicken TRPA1	TRPA1	*Gallus gallus domesticus* (chicken)	DRG neurons	Ankyrin	39.4℃					Saito et al 2014
xtTRPA1	TRPA1	*Xenopus tropicalis* (Western clawed frog)	DRG neurons	Ankyrin	39.7℃				59.24+/-18	Ohkita et al 2012
xlTRPA1	TRPA1	*Xenopus laevis* (African clawed frog)	DRG neurons	Ankyrin	36.2 +/- 0.4℃					Saito et al. 2016
snTRPA1	TRPA1	*Crotalus atrox* (rattlesnake)	TG neurons	Ankyrin	27.6℃				13.7	Gracheva et al. 2010
python TRPA1	TRPA1	*Python regius* (python)	TG neurons	Ankyrin	32.7℃				n.d.	Gracheva et al. 2010
boa TRPA1	TRPA1	*Corallus hortulanus* (boa)	TG neurons	Ankyrin	29.6℃				n.d.	Gracheva et al. 2010
ratsnake TRPA1	TRPA1	*Elaphe obsoleta lindheimeri* (rat snake)	TG neurons	Ankyrin	37.2℃				8.8	Gracheva et al. 2010
Anole TRPA1	TRPA1	*Anolis carolinensis* (green anole)	TG neurons	Ankyrin	33.9℃				45.71+/-6	Saito et al. 2012
A. allogus TRPA1	TRPA1	*Anolis allogus*	Dorsal skin	Ankyrin	33.5 +/- 0.7℃					Akashi et al. 2018
A. homolechis TRPA1	TRPA1	*Anolis homolechis*	Dorsal skin	Ankyrin	36.4 +/- 0.8℃					Akashi et al. 2018
A. sagrei TRPA1	TRPA1	*Anolis sagrei*	Dorsal skin	Ankyrin	33.5 +/- 0.7℃					Akashi et al. 2018
axTRPA1	TRPA1	*Ambystoma mexicanum* (axolotl)	Brain, lung, heart, stomach	Ankyrin	39.7 +/- 1.0℃					Oda et al. 2019
zTRPA1b***	TRPA1	*Danio rerio* (zebrafish)	Sensory neurons innervating skin and cranial sensory ganglia	Ankyrin	variable < 10℃ (cold); > 25℃ (heat)				8.2+/-0.6	Oda et al. 2016
olTRPA1	TRPA1	*Oryzias latipes* (medaka)		Ankyrin	variable					Oda et al. 2017
trTRPA1	TRPA1	*Takifugu rubripes* (pufferfish)		Ankyrin	variable < 8℃ (cold); > 25℃ (heat)					Oda et al. 2018

Invertebrate transient receptor potential (TRP) channels involved in thermosensation

Name	Synonym	Species	Location	Domain	Threshold				Q10	Reference
ceTRPA1****		*Caenorhabditis elegans* (nematode)	neurons, muscle, intestine, and epithelial cells		≤ 17℃					Reviewed in Laursen et al. 2015
Painless	dAnktm1	*Drosophila melanogaster* (fruit fly)		Ankyrin	~39-42℃					Tracey et al 2003; Sokabe et al. 2008
Pyrexia	Pyx-PA and Pyx-PB	*Drosophila melanogaster* (fruit fly)	embryos: peripheral nerves and central nerves, multidendritic neurons in larval epidermis;adults: sensory neurons around bristle eyes, britsle neurons along thorax, neurons in maxillary palps, proboscis, antennae	Ankyrin	≥ 40 ℃				Pyx-PA 18.145; Pyx-PB 15.329	Lee et al. 2005; Neely et al. 2011
dTRPA1(A)	TrpA1-RI (Prom B, ex10a), dTrpA1, dANKTM1, DmTRPA1	*Drosophila melanogaster* (fruit fly)	larval sensory neurons, adult proboscis	Ankyrin	27-29℃				9	Viswanath et al. 2003; Rosenzweig et al. 2005; Hamada et al 2008; Kang et al. 2012
dTRPA1(D)	TrpA1-RG (Prom A, ex10a), dTrpA1, dANKTM1, DmTRPA1	*Drosophila melanogaster* (fruit fly)	larval nociceptors	Ankyrin	≥ 46 ℃				116	Zhong et al. 2012
AgTRPA1		*Anopheles gambiae* (African malaria mosquito)	antennae, head	Ankyrin	25.2℃					Wang et al. 2009
AsTRPA1(A)		*Anopheles stephensi* (Asian malaria mosquito)	antennae, head	Ankyrin	30.3 +/- 0.9℃				14,5	Li et al. 2019
AaTRPA1(B)		*Aedes aegypti* (yellow fever mosquito)	antennae, head	Ankyrin	32 +/- 0.8℃				20,7	Li et al. 2019
CpTRPA1(A)		*Culex pipiens pallens* (northern house mosquito)	antennae, head	Ankyrin	21.8 +/- 0.7℃				61,2	Li et al. 2019
TRPA1		*Bombyx mori* (silk moth)		Ankyrin	21.6℃				20,5	Sato et al. 2014
hsTRPA	AmhsTRPA	*Apis mellifera* (honeybee)	antennae	Ankyrin	33.9 +/-0.6℃				17.2+/-4.0	Kohno et al. 2010
Rp-TRPA5₂	Rp-TRPA5B	*Rhodnius prolixus* (kissing bug)	head, rostrum, legs, antennae	Ankyrin	53℃; Thalf 58.6℃	53-68℃	274	72	25	*this study*

* Activation threshold temperature as determined in heterologous expression systems
** Q10 is the fold current increase over 10 ℃ increase as a measure of sensitivity, the higher Q10 value, the more sensitivity to heat the channel

Results and discussion

TRPA5 genes are ancient ankyrin receptors found across many insect orders but lost in Diptera

To start investigating the molecular basis of thermosensation in Rhodnius, we first reanalyzed the genome annotations (Version RproC3.3) complemented with available transcriptomic resources (*see Methods*) to gain insights into gene variation and genomic architecture within the TRP ankyrin family. Genomes of triatomines [37] and additional surveyed hemipteran species (Table S1) all lack an ortholog to *Pyrexia* (*Pyx*) TRP but possess one gene copy of the three canonical ankyrin TRP genes: *Waterwitch* (*Wtrw*), *TRPA1* and *Painless* (*Pain*) (Fig. 1A, Fig. S1). Rhodnius *TRPA1* and *Painless* exhibit a wide mRNA tissue distribution (Fig. 1B) that potentially indicates a canonical role in thermosensation, similarly to TRPAs expression patterns in other insect species [27-29, 49].

Figure 1. A. Phylogenetic reconstruction of the ankyrin TRP (TRPA) channel family in representative insect species. TRPA5 channels are present across insect Orders but absent from dipteran genomes (see also Fig. S1 and S2). Gene abbreviations: *Painless (Pain), Pyrexia (Pyx), Waterwitch (Wtrw)*, hymenopteran-specific TRPA (*hsTRPA*). Silkmoth, *Bombyx mori*; Hornworm moth, *Manduca sexta*; Mosquito, *Anopheles gambiae*; Fruit fly, *Drosophila melanogaster*; Flour beetle, *Tribolium castaneum*; Fire ant, *Solenopsis invicta*; Honeybee, *Apis mellifera*; Bed bug, *Cimex lectularis;* Kissing bug*, Rhodnius prolixus*; Termite, *Zootermopsis nevadensis*; Bluetail Damselfly, *Ischnura elegans.* **B.** Phylogenetic relationships of TRP genes in *R. prolixus* and their corresponding

expression levels across tissues in published transcriptomic data. Heat maps compare the expression levels

across tissues and developmental stages. Expression levels are represented as Log$_2$ FPKM +1 and depicted

with a gradient color scale. Gene models are based on genomic annotations [37], *de novo* transcriptome

assembly [44] and manual annotation of gene models not annotated in the two aforementioned studies (see

Methods).

Our scaffold-mapping analysis of three *TRPA5* isoforms previously annotated in *R. prolixus* [37] shows

that the predicted-isoforms A and B map to different genomic locations, and consist of two physically

close tandem-duplicate loci, whereas predicted-isoform C maps to a distinct scaffold. Therefore,

TRPA5A, TRPA5B and TRPA5C are hereafter referred to as three distinct loci, *TRPA5$_1$*, *TRPA5$_2$*, and

TRPA5$_3$. Intrigued by the finding of multiple *TRPA5* gene copies, we performed an extensive *TRPA5*

gene search across annotated genomic and transcriptomic datasets available for the insect Orders

Anoplura, Diptera, Coleoptera, Hemiptera, Hymenoptera, Isoptera, Lepidoptera, Odonata and

Thysanoptera. Our phylogenetic reconstruction shows that the TRPA5 ankyrin subfamily is completely

absent in all surveyed dipteran genomes (Fig. S2). TRPA5 orthologs were nonetheless found

spanning the orders Lepidoptera, Coleoptera, Hymenoptera, Hemiptera, Isoptera and Odonata (Fig.

S2), including remarkable group-specific expansions such as those in the fire ant, *Solenopsis invicta*

[29], the damselfly *Ischnura elegans*, the tobacco hornworm moth *Manduca sexta*, and several

hemipterans (Fig. S2). In addition to lineage-specific expansions through duplications, complex

alternative splicing also seems to play a role in TRPA5 functional diversification [50]. Altogether, our

large-scale phylogenetic analyses recapitulate that the insect TRPA ankyrin family comprises five

rapidly evolving clades consisting of *Waterwitch* (and HsTRPA1), *Pyrexia* (presumably lost in

Hemiptera), *TRPA5* (presumably lost in Diptera), *TRPA1* and *Painless* (Fig. 1A, SI File 1, Table S1).

Prior to interrogating a possible role for Rhodnius TRPA5 in thermosensation, we asked whether these

TRPs are expressed in sensory tissues. Whereas Rp-TRPA5$_1$ and Rp-TRPA5$_3$ mRNAs are expressed

at low or below detection thresholds across tissues (Fig. 1B), Rp-TRPA5$_2$ mRNA appears to be the

most abundant TRPA5 transcript in adult Rhodnius. Rp-TRPA5$_2$ is significantly enriched in adult male

and female heads (Fig. 1B, Fig. S3, Table S5). We further examined expression profiles of Rp-TRPA5$_2$

via quantitative PCR of additional canonical sensory tissues. Rp-TRPA5$_2$ is abundant in the rostrum

and legs and expressed at lower levels in antennae (Fig. 1B, Fig. S3), a first indication in line with a

possible role in thermosensation.

Rhodnius TRPA5$_2$ exhibits unique structural features in the Ankyrin Repeat Domain and the

selectivity filter

To visualize structural features of Rhodnius ankyrin TRP homologs, we used DeepMind's protein

structure prediction software AlphaFold 2.0 [51-53]. Monomeric structures of Rhodnius TRPA1,

Painless, Waterwitch, and TRPA5$_2$ were modeled, as well as a tetrameric model of the Rhodnius

TRPA5$_2$. We then performed pairwise comparisons of the Rhodnius orthologs to their Drosophila

correspondent monomeric structures for TRPA1, Painless and Waterwitch, and compared Rp-TRPA5$_2$

to Dm-Pyx separately (Fig. 2). Each monomeric prediction is presented using a coloring scheme

reflecting a structural reliability measure (Fig. 2A) next to a rainbow representation running from the

N- to the C-terminus (Fig. 2B).

All Rhodnius and Drosophila ankyrin TRP monomeric structures shared several expected features

with the cryo-EM structure of HsTRPA1 [54], including the N-terminal ankyrin repeat domain (ARD),

six transmembrane α-helices (S1-S6), and a region corresponding to the allosteric nexus of Hs-TRPA1

connecting the ARD and the transmembrane region [54]. The C-terminal region features at least one

α-helix, which together with the corresponding helices from the other subunits, most likely form a

coiled-coil in the tetramer as seen in the solved TRPA1 structures. Furthermore, the recently published

structure of Dm-TRPA1-A in state 1 confirms distinct predicted features in our model, including the

interfacial helix and the interaction between AR12 and the region C-terminal of the coiled-coil helix

[55]. However, without the constraints of the other monomers and the interactions that would force the

C-terminus into the coiled-coil, some of the helices are oriented in unrealistic directions while the

secondary structure remains plausible.

Figure 2. Predicted tetrameric structure of Rp-TRPA5$_2$ alongside monomers of all TRPA homologs in *Drosophila melanogaster* and *Rhodnius prolixus*. A. TRPA monomers colored by pLDDT score from the AlphaFold modeling. Blue represents a plDDT score of more than 90, reflecting a high confidence. **B.** TRPA monomers colored by chain bows, with the N-terminus in blue and the C-terminus in red. **C.** Tetrameric model of *R. prolixus* TRPA5$_2$, colored as chain bows as in B. The black box indicates the location of the selectivity filter shown in D. **D.** Top view of the selectivity filter of the pore of *R. prolixus* TRPA5$_2$ (top) and human TRPA1 (*H. sapiens*, pdb:6V9Y) (bottom). Three important residues identified by Zhao et al [56]– L913, G914 and D915 – are marked in TRPA1. The equivalent residues L913 and E914 are marked in TRPA5$_2$. The sequence alignment shows the selectivity filter. Note the glycine is absent in TRPA5$_2$ (see also Fig. S2).

Since dipteran and hemipteran insects lack *TRPA5* and *Pyrexia*, respectively, we asked if the two channels may occupy homologous structural and functional niches. From a pairwise overall comparison of the reliable monomeric models, despite the conserved regions mentioned above, Rp-TRPA5$_2$ appears to deviate from Dm-Pyrexia in three regions: the ARD, the transmembrane domain, and the pore helices that flank the selectivity filter (Fig. S4). First, Rp-TRPA5$_2$ possesses 14 AR

compared to 9 in Dm-Pyrexia, and displays longer loops, compared to Dm-Pyrexia and all other

channels, including between the third and the fourth ankyrin repeats, within the fifth ankyrin repeat,

and between the fifth and the sixth ankyrin repeats, counting from the N-terminus. This observation is

interesting as ankyrin repeats are 31-33 residue protein motifs consisting of two α-helices connected

by a ß-turn that occur in tandem arrangement to form ARDs critical for physiological processes [57],

including a previously suggested role in thermal activation sensitivity [58, 59].

Hence, in vertebrates, two regions of 6 ARs each in the snake TRPA1 (AR3-8; AR10-15) have been

shown to revert the channel thermal sensitivity by conferring heat-sensitivity to a chimeric AR human

TRPA1 (Hs-TRPA1) [60]. Furthermore, transfer of a part of the ARD from Dm-TRPA1 (AR10-15), a

region shown to control thermosensitivity in the fruitfly TRPA1, to Hs-TRPA1 also produced a heat-

sensitive Hs-TRPA1 [60]. The temperature-dependent dynamics of the ARD has also recently been

investigated in the TRPV1 channel, demonstrating that the ARD undergoes structural changes at

similar temperatures that lead to TRPV1 activation, which suggested a potential role in the

temperature-dependent structural changes leading to the channel opening [58]. The N-terminus region

of mosquito TRPA1 also seems to be quite critical for heat-sensitivity [19]; however, there have been

contradicting data for TRPA1, both from human and mosquito, arguing that additional regions

controlling thermosensitivity are located outside the ARD [26, 61]. Another interesting feature is the

disruption in the ankyrin repeat stacking between the fifth and the sixth ankyrin repeat in both Rhodnius

and Drosophila Painless, which is not seen in Dm-Pyrexia and Rp-TRPA5$_2$. This breaking point

coincides with the resolved N-terminal end of the recently reported structure of Dm-TRPA1-A in state

2, which is suggested to represent a temperature sensitized, pre-opened conformation of the channel

[55]. Although additional functional evidence that ARDs may play a general role in insect TRP

thermosensitivity is needed, these differences between Painless, Pyrexia and TRPA5 hint at a similar

activation mechanism in Painless as proposed for TRPA1, yet potentially distinct from the activation

mechanism of Dm-Pyrexia and Rp-TRPA5$_2$. In contrast, the shape of the ARD in Dm-Pyx, Dm-Wtrw

and Rp-Wtrw all exhibit remarkable similarities compared to that of Rp-TRPA5$_2$ (Fig. 2A) and the

predicted number and position of ankyrin repeats also appears remarkably conserved between *R.*

prolixus and *D. melanogaster* for TRPA1, Painless, and Wtrw (Table 2) suggestive of potentially

conserved interspecific functions for each Rhodnius ortholog of the latter three channels.

Second, looking at the selectivity filter and the upper gate of the Rp-TRPA5$_2$ tetramer model (Fig. 2C

and D), one conserved glycine (Gly914) is notoriously absent in Rp-TRPA5$_2$ (Fig 2D). This is striking

as it is conserved in most other TRPAs, and either conserved or substituted for Serine or Threonine

in most other non-hemipteran TRPA5 proteins (Fig. S2). Comparing it to a structure of Hs-TRPA1

published by Zhao et al. (PDB:6V9Y) [56], the Leu913 and Glu914 in Rp-TRPA5$_2$ seem to remain

largely in the same locations as Leu913 and Asp915 in Hs-TRPA1, despite the shorter pore loop. The

main difference therefore is that there are only two carbonyls in the pore loop of Rp-TRPA5$_2$, as

opposed to three carbonyls in Hs-TRPA1. Although the difference is smaller than anticipated, since

Gly914 is suggested to be important in gating, and lies in the location of the selectivity filter, this may

affect permeation properties of Rp-TRPA5$_2$.

Finally, a difference in the lower gate can be seen where the Hs-TRPA1 structure is much narrower

than that of Rp-TRPA5$_2$, and where the pore of the Rp-TRPA5$_2$ model is blocked by the position of the

sidechain of Glu914. Since AlphaFold is better at modeling backbone folding than individual

sidechains, the significance of this finding may be limited, but the Rp-TRPA5$_2$ model features a

narrower pore than Hs-TRPA1 resolved in a closed conformation in complex with an antagonist [56].

Our findings that Rp-TRPA5$_2$ shares several conserved features of ankyrin TRPs, as well as unique

structural novelties, raises the question as to whether Rp-TRPA5$_2$ may have undergone selection to

fill the same function as *Pyrexia*, or whether its distinctive structural features may underlie a new type

of thermosensitivity phenotype. If so, it would be interesting to determine whether the heat activation

of the channel may be affected.

Table 2. Number of Ankyrin repeats observed in each monomer structure of thermos TRPA of *Rhodnius prolixus*

and *Drosophila melanogaster.*

	Number of Ankyrin repeats	
Name	*Rhodnius prolixus*	*Drosophila melanogaster*
TRPA1	17	17
Painless	10	10
Waterwitch	9	10
Pyrexia		9
TRPA5$_2$	14	

TRPA5$_2$ encodes a novel class of thermosensitive insect receptor

In order to demonstrate a potential role of candidate TRPA5$_2$ as a thermosensitive ion channel, we

optimized an *in vitro* cell-based platform to record temperature-elicited currents from HEK293T cells

under whole-cell patch-clamp configuration. We transiently expressed a bicistronic T2A-fluorescent

marker cassette [62] together with the candidate TRP channel, which localized well to the plasma

membrane (Fig. S5). We delivered a fast heat stimulus by coupling an infrared laser diode fiber optic

[63] to a PID controller that conveyed millisecond current pulses (Fig. S6).

At the molecular level, non-denaturing SDS-Page analysis showed that Rp-TRPA5$_2$ assembles

similarly to other TRPs as a membrane-bound tetramer when expressed in HEK293T cells (Fig. S5).

To validate the infrared (IR) patch clamp system, we first transiently expressed two known

thermoTRPs, the rat TRPV1 (rTRPV1) and fruit fly TRPA1 isoform D (Dm-TRPA1-D) (Figs. S6, S7),

both of which formed expected homotetrameric structures (Fig. S5). First, we used the ionic current

increments through the open patch pipette (holding potential -2 mV), to calculate the temperature

changes associated with the different laser intensities. Typically, the current pulses were set to result

in temperature increments at the cell membrane in the range of 23.5-71.7°C. A similar laser stimulation

protocol led to marginal whole-cell current changes in non-transfected cells (Fig. 3A, Fig. S7A-B).

Compared to non-transfected cells, we then observed a strong increase in the current amplitude of

cells expressing rTRPV1 (Fig. 3B, Fig. S7C-D) with an enthalpy change associated with the activation

of 88.3±9.4 kcal/mol, which is comparable to the published enthalpy values obtained using millisecond

temperature jumps of ΔH = 85 kcal/mol, $T_{1/2}$ = 47.5°C for rTPRV1 (calculated from enthalpy and

entropy values for steady-state activation from Yao et al. 2010, see Table 1) [63-65]. A temperature-

induced activation response was also observed for the heat activated fruit fly channel, Dm-TRPA1-D,

for which in our more precise setup, at 46.3°C [22], the open probability (Po) of the channel is about

10% (Po=0.1), corroborating a noxious activation temperature > 42°C [22]. Assuming complete

activation by temperature of this channel (Po=1), which was not measured in previous studies due to

limitations in the maximum temperature to which the Dm-TRPA1-D channel could be subjected, the

activation process is characterized by an enthalpy change ΔH= 68.7±13.1 kcal/mol and $T_{1/2}$ = 53.5°C

(Fig. 3C, Fig. S7E-F). These results demonstrate highly consistent biophysical properties of known

thermoTRP channels evaluated with our laser-based delivery method, using the modified C-terminus

in our expression cassette.

Figure 3. Thermodynamics of Rp-TRPA5₂ temperature-activated currents. A. Activation time course of

temperature for the open pipette calibration (upper panel), and baseline current traces for a control cell (non-

transfected). During each voltage pulse, a temperature step of 700ms was presented from room temperature to

71°C. **A.** Open patch-clamp pipette current traces in response to increasing voltage pulses (10 - 23 mV at 1 mV

voltage inputs). **B.** Whole-cell currents evoked by temperature at -30mV of HEK293T cells expressing rat

TRPV1, the heat-activated mammalian vanilloid thermo-TRP channel. **C.** Whole-cell currents evoked by

HEK293T cells expressing Dm-TRPA1-D. **D**. Whole-cell currents evoked by HEK293T cells expressing Rp-

TRPA5₂ under whole-cell configuration patch-clamp, held at -30 mV (n=9). Negative currents indicate that Rp-

TRPA5₂ is a cationic channel. Data are presented as means ± standard errors. **E**. Current-Temperature

relationship for Rp-TRPA5₂ where current density is plotted after normalization by cell membrane capacitance

and scaling to the mean maximum. **F.** Fraction of Rp-TRPA5$_2$ channels in the open state (Open probability, P$_o$)

as a function of the temperature. The Po vs 1/T was fitted to a Boltzman function with the midpoint of activation

(T$_{1/2}$) reached at 58.6 °C. **G** Van't Hoff plot estimates of Rp-TRPA5$_2$ with an activation enthalpy of the

endothermic transition at 92 kcal/mol and an entropic change associated to the temperature activation process

at 274 cal/mol*K. at -30 mV as a function of the temperature [63]. **H.** Coupling between enthalpic (ΔH) and

entropic (ΔS) changes for each one of the experiments recorded **I.** ΔG vs. temperature plot for Rp-TRP5A$_2$

channels, temperature activation is associated with small ΔG (free energy) changes, as reported for other

families of mammalian thermo receptors. ΔG was calculated as -RT*ln(Keq) [66].

Finally, when holding the membrane potential at -30mV in patched mRuby2-expressing cells

transfected with Rp-TRPA5$_2$, whole cell currents were evoked by temperature steps from 53°C to 68°C

(Fig. 3D, 3E). The average temperature for the activation "threshold" was 53°C, defined as Po = 0.1

calculated from the Van't Hoff plots. The channel opening appeared to saturate at 68°C (Po=0.9) (Fig.

3F), with a T$_{1/2}$ = 58.6°C. The steady-state parameters of activation were calculated from the current

at the end of the 700 ms temperature pulse. The current density versus temperature relationship (Fig.

3E) indicates that the opening of Rp-TRPA5$_2$ involves an activation enthalpy of approximately 72.6±14

kcal/mol (Fig. 3G). Similar large enthalpy changes ranging from 60 to 200 kcal/mol are involved in the

opening of TRPM8, TRPV1, and TRPV2 [9], supporting that the TRPA5 ion channel activity is

extremely temperature-dependent with high enthalpy change associated with the channel opening.

The large entropy value further indicates that the channel transits between a highly ordered closed

state and a strongly disordered open configuration, similarly to TRPV1 (316 cal/mol·K at -60mV).

Altogether, these results demonstrate that Rp-TRPA5$_2$ acts as an insect thermoTRP receptor of

noxious heat *in vitro*.

Based on the open probability (Po), we calculated a Q$_{10}$ value of ~ 25, which is in the range of

characterized noxious vertebrate (rTRPV1 Q$_{10}$ = 16.8; rTRPV2 Q$_{10}$ = 20.6) and invertebrate

thermoTRPs (fruit fly Pyrexia Q$_{10}$ = 18.2) (Table 1). From a thermodynamic point of view, many TRP

ion channels are modulated by temperature [67], but our results clearly support that TRPA5$_2$ belongs

to a restricted category of thermoTRPs as it is directly activated by temperature as the sole stimulus

[8, 66]. Rp-TRPA5$_2$ appears to be activated in a higher noxious range compared to all known

invertebrate thermoTRPs thus far (Fig. 1A, Table 1, Fig. S2) including the fruit fly Painless and TRPA1

channels that mediate thermal nociceptive escape through larval mdIV neurons at temperatures above

40°C and 46°C, respectively [17], or Pyrexia channels that induce paralysis in adult flies upon

exposure to 40°C [18]. In mammals, only TRPV2 contributes to highly noxious (>52 °C) heat sensing

[64, 68, 69].

ThermoTRP channels can integrate voltage and temperature allosterically [67]. We aimed to establish

the temperature sensitivity of the channel directly from the Van't Hoff plot, and not from the influence

of temperature on the voltage activation process, allowing us to establish the thermodynamics of

temperature activation process independently from other stimuli. Using fast temperature jumps, we

could activate Rp-TRPA52 channels directly, validating that the channel is a thermoTRP in the noxious

temperature range. In addition, similarly to rTRPV1 and Dm-TRPA1-D, Rp-TRPA5$_2$ has large enthalpy

changes related to the channel activation, which is related to the high sensitivity of these channels to

temperature changes. In rTRPV1, thermal activation at +60 mV causes only a relatively small enthalpy

change (30 kcal/mol) compared to 100 kcal/mol when rTRPV1 is activated at -60 mV, representing a

three-time increase in temperature sensitivity at negative voltages [63]. Similarly to rTRPV1, Rp-

TRPA5$_2$ shows almost no heat-elicited activity at depolarized potentials (> 0mV), but a robust response

at negative voltages (-30mV). Although the temperature activation for Rp-TRPA5$_2$ appears very similar

to Dm-TRPA1-D and rTRPV1, the activation kinetics of Rp-TRPA5$_2$ is slower compared with these

channels, which could be due to intrinsic molecular interactions influencing the transition between

closed and open states [67], in light of our observations of several key defining structural differences.

This difference in the opening kinetics has been observed before for other thermoTRPs, like the

temperature-dependent HsTRPV3, which exhibits a high temperature sensitivity, comparable to

hTRPV1, but presents slower kinetics [65, 70].

TRPA5's physiological role as a noxious heat sensor

Sensory receptors in the same clades are often tuned to detect a stimulus over a discrete

window of intensities, enabling the recognition of physiologically relevant cues over a wide dynamic

range [1, 3, 71]. The TRP Ankyrin family is an excellent example of this pattern, as distinct, yet closely

related channels account for thermal responsiveness over a range from innocuous to noxious heat [4,

12]. In addition, orthologous thermoTRPs often have different activation temperatures, and this has

been postulated to reflect functional adaptive evolution to different optimal temperatures, coordinating

thermoregulatory behaviors such as host seeking, thermal avoidance, and tracking of optimal

temperatures [72]. Our findings on the temperature activation of Rp-TRPA5$_2$, a TRPA clade not

functionally characterized, clearly support a role for this previously orphan gene as a molecular sensor

in the noxious heat spectrum. Notably, a known noxious heat receptor in Diptera, *Pyrexia,* is missing

from *Rhodnius*, while a TRPA5 ortholog has not been found in flies and mosquitoes*,* raising the

interesting possibility that convergence and functional redundancy might account for the evolutionary

patterns of differential gain and retention of thermoTRPs in insects.

Consistent with a role as a noxious thermosensor, quantitative analyses show that Rp-TRPA5$_2$

is expressed broadly across tissues, with high levels detected in the head, rostrum, legs, thorax,

abdomen and to a lesser extent in the antennae (Fig. 1, Fig. S3). By qPCR, we detected the highest

expression in the head (Fig. S3), in agreement with previous transcriptomic analyses [37] and

remarkably similar to the expression pattern of *Pyrexia* in the fruit fly [18]. Noxious environmental

temperatures are extremely common in natural environments of small insects with low thermal

capacity, thus, detecting and avoiding heat is critical to prevent injury. Temperature distributions vary

widely for natural objects. For instance, dry and moderately gray-colored or dark objects such as tree

bark or rocks easily reach temperatures above 50°C [73]. If the humidity level is high, and radiative

cooling of the sky is not effective, the same objects can reach temperatures above 60°C in the full

sun. For example, temperatures of dry leaf substrates on the ground can exceed 50°C in full sun since

they do not undergo evaporative cooling, which would typically prevent a leaf's surface temperature

from going above 40°C. In lab simulated natural environments and in field thermal imaging studies,

insects can reach 60°C under full sun with high humidity in as little as 15 seconds [73]. Considering

that Rhodnius adults are about 3 cm in length and dark colored, with a small thermal capacity, and

that they typically inhabit tropical environments with high humidity, they are likely to rapidly reach

temperatures above 60°C if exposed to full sun, suggesting that TRPA5 may mediate noxious heat

avoidance. Although the physiological and behavioral role of Rp-TRPA5$_2$ will need to be examined in

detail, the discovery of the activation range of TRPA5$_2$, opens avenues for exploring the convergent

evolution of noxious heat sensing between TRPA5$_2$ in Rhodnius bugs and *Pyrexia* in Drosophila flies,

and for assessing whether it is also relevant for both inner temperature regulation and warm substrate

avoidance.

To better understand the mechanisms of Rp-TRPA5$_2$ function and heat stimulus integration,

future studies could examine co-expression with other TRPs. Rp-TRPA5$_2$ is also expressed in

antennae, albeit at lower levels (Fig. S3). Antennae in triatomine bugs function as a multimodal

sensory organ, notably harboring highly specialized thermosensory sensilla involved in the detection

of host signals such as heat and moisture [36]. Heat sensing in triatomines is driven by air temperature

gradients around the host (conductive heat) [39, 40] as well as infrared radiation [74]. Antennal

thermoreceptors are known to be required for orientation and distance estimation [75] as well as for

precisely locating warm skin blood vessels before initiating proboscis extension and biting [76].

Similarly to homologous noxious heat detectors in mosquito antennae, Rp-TRPA5$_2$, might contribute

to tuning host-selective thermotaxis to avoid stimuli exceeding host temperatures. An interesting next

step would be to map the cellular location of Rp-TRPA5$_2$ in peripheral sensory neurons to investigate

potential colocalization with other channels involved in innocuous heat detection.

TRPA5 as novel target to mitigate the impact *Rhodnius prolixus* in the transmission of Chagas

disease

ThermoTRPs are polymodal sensors of physical and chemical stimuli [71]. For example, channels in

the insect TRPA1 and HsTRPA clades are typically activated by AITC and various plant-derived

chemicals such as carvacrol and citronellal [23, 28, 29]. However, characterized receptors of noxious

heat in insects such as Pyrexia and Painless do not exhibit chemical sensitivity to electrophiles [17].

Since our phylogenetic analyses support that Rp-TRPA5$_2$ is the sister clade to TRPA1, it is possible

that Rp-TRPA5$_2$ also plays a chemosensory role. Indeed, live Rhodnius treated with capsaicin, the

vanilloid pungent extract of chili peppers, were recently shown to have impaired orientation towards a

thermal source [77]. Notably, this compound can directly activate the mammalian TRPV1 receptor

independent of temperature, and the mammalian noxious temperature receptor, TRPV2, when bearing

only four mutations [3, 15]. Other than capsaicin, both TRPV1 and TRPV2 are readily activated by

additional vanilloid compounds such as resiniferatoxin, an active compound from the cactus *Euphorbia*

resinifera used for medicinal purposes and other plant-derived compounds that act as chemical

agonists [78]. Findings of botanical compounds triggering chemical activation of TRPA5$_2$ combined

with *in vivo* behavioral exposure studies, may thus contribute to uncovering new classes of natural

repellents potentially co-mediating heat-avoidance. This would be significant, not only in Rhodnius but

also for other triatomines and hemipteran vectors sharing a close TRPA5$_2$ orthologue such as the bed

bug, *Cimex lectularius*. Together, our study deorphanizes and characterizes the first ankyrin TRPA5

ion channel acting as noxious heat sensor, consistent with independent evolutionary origins of the

molecular transduction of noxious stimuli in insects, while simultaneously opening the door for further

pharmacological studies of TRP receptors in triatomine vectors.

Material and Methods

Phylogenetic analyses. Amino acid sequences of insect TRPA channels from the Anoplura (sometimes

included under Psocodea or Phthiraptera), Coleoptera, Diptera, Hemiptera, Hymenoptera, Isoptera and

Lepidoptera insect orders were retrieved from the InsectBase repository [79], FlyBase version FB2020_03 [80],

VectorBase (https://www.vectorbase.org), BeeBase [81], NCBI-blast [82], EnsemblMetazoa

(https://metazoa.ensembl.org), the i5k Workspace@ NAL [83] and OrthoFinder [84]. The TRP sequences from

insect model systems including *Drosophila melanogaster*, *Tribolium castaneum*, *Bombyx mori*, *Apis mellifera*

and *Rhodnius prolixus* were used as templates to mine and curate orthologous TRP ORF sequences from

annotated insect genomes and transcriptomes. To classify the uncharacterized TRPs, amino acid sequences

were aligned using MAFFT [85], and Maximum-Likelihood phylogenetic trees were inferred in IQ-TREE v1.6.11

using ModelFinder (Ultrafast Bootstrap, 1000 replicates), using a best-fit model JTT+F+I+G4 measured by the

Bayesian information criterion (BIC) [86-88]. The phylogenetic trees were visualized, rooted at mid-point and

annotated in R V3.6.3 using the ggtree package [89] and Evolview [90]. The accession numbers are listed in

Table S1.

TRPA5 gene annotation and tissue expression. We collected Illumina read data from *R. prolixus* tissue

libraries published in the Sequence Read Archive (SRA) at NCBI under Bioproject accession numbers

PRJNA281760/SRA:SRP057515 (antennal library from larvae, female adult and male adult),

PRJEB13049/SRA:ERP014587 (head library), and PRJNA191820/SRA:SRP006783 (ovary and testes library).

We performed low-quality base trimming and adaptor removal using cutadapt version 1.16 [91] and aligned the

trimmed read pairs against the *R. prolixus* assembly version RproC3.0.3 (retrieved from VectorBase.org)

genome using HISAT2 version 2.2.0 [92]. The existing annotation was used to create a list of known splice sites

using a python script distributed with HISAT2. We used StringTie version 2.1.3b [93] with the *conservative*

transcript assembly setting to improve the annotation, reconstruct a non-redundant set of transcripts observed

in any of the RNA-Seq samples, and compute expression estimates.

We applied Trinotate version 3.2.1 [94] to generate a functional annotation of the transcriptome data. In

particular, the functional annotation of TRP genes for which the initial genome annotation was absent or

incomplete (i.e *TRPA5, Nan, Pain*) were localized in Trinotate annotation followed by validation using the Apollo

gene browser [95]. Rp-TRPA5-A, Rp-TRPA5-B and Rp-TRPA5-C are three independent loci. Therefore, we

opted to rename them *Rp-TRPA5$_1$*, *Rp-TRPA5$_2$,* and *Rp-TRPA5$_3$,* respectively, to avoid confusion with standard

gene isoform nomenclature, which applies in the case of TRPA1-A and TRPA1-B, two isoforms of the Rhodnius

TRPA1 locus. All TRP gene identifiers are presented in Table S2.

The alignment BAM files were used to estimate transcript abundance using StringTie together with our improved

annotation. The abundance tables from StringTie were imported into R using the *tximport* package [96], which

was used to compute gene-level abundance estimates reported as FPKM. We used the R package *pheatmap*

to visualize the expression level of TRP genes.

Monitoring of TRPA5$_2$ expression levels by quantitative PCR. Live adults of *R. prolixus* were obtained from

BEI Resources (USA). Female antennae, rostrum, legs, heads (minus antenna and rostrum), and bodies (thorax

minus legs + abdomen) were dissected and pooled from 15 individuals in DNA/RNA shield reagent (Zymo) and

stored at -20°C until further processing. Total RNA was isolated using the Monarch RNA extraction procedure

(New England Biolabs), including tissue grinding in liquid nitrogen and a DNAse I step. cDNAs were synthesized

using the GoScript cDNA synthesis procedure (Promega) prior to concentration assessment using the Qubit

High sensitivity DNA kit (Invitrogen). Two gene-specific primer (GSP) sets were designed for Rhodnius Actin

(Genbank acc. Nr. EU233794.1) and TRPA5$_2$ using Primer3 version 2.3.7 in Geneious [97] (Table S3). Each

primer set was initially validated by calculating standard curves from serial dilutions of template cDNA (2 ng/µL

to 0.25 ng/µL) and primer mix (5 to 0.25 µM) with choosing amplification efficiencies (E) between 95 and 100%.

qPCR amplification products from initial runs were additionally checked on 2% agarose gels to verify the correct

amplicon sizes and the absence of primer dimers. As a final validation, qPCR products were purified using Exo-

SAP (Fermentas) prior to Sanger sequencing to ensure product amplification specificity. Quantitative PCR

reactions were then run in three technical replicates on a CFX384 Real-Time PCR system (Bio-Rad) with

quantification and dissociation curves analyses performed for three independent experiments using the CFX

Maestro Software 2.3 (Bio-Rad). Each five-microliter reaction contained 2.5 µL 2x SsoAdvanced Universal SYBR

Green Supermix (Biorad), 0.25 ng cDNA and 0.125 µM primers. Cycling conditions were as follows: 95°C for 2

min, 39 cycles of 95°C for 10 s, 60°C for 10 s followed by a dissociation curve analysis from 65.5°C to 89.5°C

with gradual heating at 0.6°C/s. Relative log-fold expression levels were normalized per tissue type against the

reference gene and calibrated relative to Antennae (log fold expression = 1).

Alpha-fold modeling and DALI analyses. Monomer structures of Rhodnius TRPA1, Rhodnius Painless,

Rhodnius Waterwitch, Rhodnius TRPA5$_2$, Drosophila TRPA1, Drosophila Painless, Drosophila Waterwitch and

Drosophila Pyrexia were generated using AlphaFold2 with amber relaxation activated [51] on Colab's server

[53]. To model the Rhodnius TRPA5$_2$ tetramer, due to limitations in computational power, the transmembrane

region (residues 608-1078) was modeled first, and then used as a custom template to model a monomer of

residues 42-1078. The first 41 residues and the C-terminal of the monomers from residue 1079 were disordered

and truncated to avoid clashes when assembling the tetramer. A tetramer was assembled of four copies of the

monomer by aligning them to each of the chains of the truncated transmembrane tetramer in PyMOL [98]. The

monomer models were compared with pairwise structural alignment using the Dali server [99]. The PDB files

are provided as supplementary material.

Molecular cloning. Antennae from twenty Rhodnius adult individuals were obtained from a laboratory culture

(Orchard lab, University of Toronto Mississauga, Canada) and stored in DNA/RNA Shield™ reagent (Zymo

Research). Tissues were disrupted in Trizol using a Premium Multi-Gen 7XL Homogenizer (PRO Scientific) and

RNA was subsequently extracted using the Direct-zol RNA kit (Zymo Research), including a DNAse step to

remove genomic DNA contamination. cDNA was synthesized from 1ug Total RNA using the GoScript™ Reverse

Transcriptase kit (Promega) and random hexamers following the recommended manufacturers' protocol. RNA

and cDNA qualities were verified using a Nanodrop (Nanodrop 2000/2000c UV-vis spectrophotometer, Thermo

Scientific) and quantified using a Qubit Fluorometer (ThermoFisher). The coding regions of Rhodnius *Rp-*

TRPA5$_2$ was amplified from antennal cDNA using gene-specific primers designed based on Rhodnius full length

TRP sequences [37] and containing unique restriction sites (Table S3). PCR reactions were performed in a

Veriti™ Thermal Cycler (ThermoFisher) using the Advantage® 2 PCR Kit (Takara Bio) in a touchdown cycling

program as follows: 95°C for 2 min, 16 cycles of 95°C for 30 sec, 68°C for 1 min (-0.5°C/cycle), 68°C for 4 min

followed by 20 cycles of 95°C for 30 sec, 60°C for 1 min, 68°C for 4 min, and a final step at 68°C for 10 min.

Amplification products were analyzed by electrophoresis, and fragments of expected size were excised from the

gel, purified using the Monarch® DNA gel extraction kit (NEB) and subjected to Sanger Sequencing for ORF

sequence-verification prior to codon-optimization at Genscript and subcloning. For the rat rTRPV1 and the fruit

fly Dm-TRPA1-D, gene specific primers (Table S3) were used to amplify the ORF including suitable flanking

restriction sites prior to gel purification and double restriction digestion. The digested PCR products were gel

purified and ligated in an expression cassette containing the human cytomegalovirus (CMV) immediate early

promoter and engineered to include a C-terminal tag by the monoclonal antibody FLAG epitope sequence

(DYKDDDDK), followed by a Ser-Gly-Ser linker peptide, a T2A peptide sequence (EGRGSLLTCGDVEENPG)

and the coding region of the cytoplasmic fluorescent marker protein mRuby2 [62, 100]. The ligation mixtures

were used to transform *Stbl3* competent *E. coli* cells (ThermoFisher) using standard protocols. Plasmid DNAs

were purified using the Qiaprep spin Miniprep (Qiagen) and verified by Sanger sequencing using internal gene-

specific and vector primers to ensure overlapping sequence information in both forward and reverse directions.

High yield pure plasmid DNA preparations were subsequently obtained from 100 mL overnight LB broth cultures

using the endo-free ZymoPURE™ II Plasmid Midiprep Kit (Zymo Research, USA).

Transient HEK293T cell expression. Plasmid DNAs clones from TRP cDNAs were transiently expressed in

HEK293T cells to optimize expression conditions via mRuby2 visualization and western blot analysis prior to

whole cell patch clamp recordings. HEK293T cells were seeded at a density of 0.6×10^6 cells on day 0 in 60 mm

culture dishes (ref 25382-100, VWR) in DMEM High Glucose, GlutaMAX (Life Technologies) supplemented with

10% FBS (Seradigm Premium, VWR, USA). For each transfection, lipid complexes containing 2.5 µg DNA: 10

µL L2000 (Life Technologies) mixed in Opti-MEM I Reduced Serum (Life Technologies) were added dropwise

to the cells at 50% confluency (1.2×10^6 cells, day 1). The culture medium was exchanged with new DMEM/FBS

medium six-hours post-transfection. Cells were incubated at 37°C in a humidified HERAcell 150i incubator

(Thermo Scientific) with 5% CO_2.

Biochemistry. For whole-cell TRP expression analysis, cells were harvested 72h post-transfection; the medium

was decanted, cells were collected in 2mL cold D-PBS, centrifuged for 5 min at 4,000 rpm at 4°C and then the

supernatant was discarded. The cell pellet was gently suspended in 50 µL cold Ripa lysis buffer (Thermo

Scientific) supplemented with 1% Triton-X100 (Sigma-Aldrich) and complete EDTA-free protein inhibitors

(Sigma-Aldrich). Cell membranes were lysed for 1h at 4°C with gentle rotation on a sample homogenizer, and

cell debris were collected by centrifugation at 4°C for 15 min at 13,000 rpm. The crude protein lysate

concentration was quantified by bovine serum albumin (BSA) (Sigma-Aldrich) and 25 µg crude extract was

loaded on NuPAGE™ 3-8% Tris-Acetate gels (ThermoFisher) and transferred to a polyvinylidene difluoride

membrane on a TurboBlotTransfer system (Bio-Rad Laboratories). The membranes were blocked with 5% milk

(Bio-Rad) in Tris-buffered saline containing 0.1% Tween 20 (TBST, Bio-Rad) and incubated overnight with

aFLAG antibody 1:2,500 (GE Healthcare) on a gently rocking platform at 4°C. After washing with TBST the

membranes were incubated for 1h at ambient temperature in the dark with horseradish peroxidase (HRP) ECL

anti-mouse conjugated antibody (Amersham, USA) diluted in 5% milk in TBS-Tween at 1:2,500. Membranes

were rinsed in TBST and revealed using the SuperSignal West Femto (Thermo Scientific) and imaged on a

ChemiDoc system (Bio-Rad Laboratories).

For membrane surface expression, the plasma membrane expression of Rp-TRPA5$_2$ channels was assessed

using the Pierce Cell surface Protein isolation kit (Thermo Scientific). On day 0, four T75 cm^2 flasks were seeded

with 1 x 10^6 HEK293T cells. Forty hours later, each flask was transfected with lipid complexes containing 48 µg

endo-free plasmid DNA and 96 µl Lipofectamine 2000 diluted in Opti-MEM serum and incubated at 37°C. 72

hours post-transfection, cells were gently washed with ice-cold PBS, labeled with Sulfo-NHS-SS-Biotin, and

harvested following the manufacturer's protocol. Cells were lysed on ice for 30 min in the manufacturer's lysis

buffer supplemented with 0.5% Triton-X100 and complete EDTA-free protein inhibitors (Sigma-Aldrich), with

gentle 5s vortexing every 5 min, and two 5x-1s sonicating bursts on ice. Following centrifugation, the cell lysate

was bound to NeutrAvidin agarose resin and gently mixed for 60 min at ambient temperature on a platform

rotator. The membrane-bound fraction was eluted with 50mM Dithiothreitol in SDS-Sample buffer (62.5 mM

Tris/HCl pH6.8, 1% SDS, 10% Glycerol) and then placed on ice. For Western Blot analysis, 32 µl of the

membrane protein eluate fraction were mixed with Laemmli buffer (Bio-Rad) supplemented with 10% 2-

mercaptoethanol. Sixteen µl of the homogenized protein-loading buffer sample were loaded in duplicates on a

NuPAGE™ 3-8% Tris-Acetate gel (ThermoFisher) to be probed separately with FLAG and ATPase antibodies.

Proteins were separated by electrophoresis for 3h at 80V at 4°C, then transferred to a polyvinylidene difluoride

membrane on a TurboBlotTransfer system (Bio-Rad Laboratories). The membranes were blocked in parallel

with 5% milk (Bio-Rad) in Tris-buffered saline containing 0.1% Tween 20 (TBS-T, Bio-Rad) and incubated

overnight on a gently rocking platform at 4°C with aFLAG antibody 1:2,500 (GE Healthcare) or with Anti-Sodium

Potassium ATPase antibody 1:2,500 (ab76020, Abcam) diluted in 5% milk. After three washes with TBST, the

membranes were incubated for 1h at ambient temperature in the dark with HRP ECL anti-mouse conjugated

antibody (Amersham, USA) at a 1:2,500 dilution in 5% milk/TBST. Membranes were rinsed in TBST and revealed

using the SuperSignal West Femto (Thermo Scientific) and imaged on a ChemiDoc system (Bio-Rad

Laboratories).

Temperature control using a laser system. We used a manual patch-clamp station (Axopatch 200, Molecular

Devices) equipped with a fiber-delivered laser system to record temperature-activated currents under a precise

voltage-clamp control. The setup was modified after Yao et al (2009) [101] (Fig. S5) and takes advantage of

water's IR absorption band to generate rapid temperature jumps from RT to high temperatures. It combines an

infrared diode laser (λc =1460 nm (+/-20 nm), Output power = 4.8 watts) (Seminex Inc.) coupled with a 100-um

optical fiber with a striped tip (ThorLabs, Inc.) as the controllable heat source. Two independent

micromanipulators allowed us to precisely align the relative positions of the patch-clamp electrode and the fiber

on a single cell (Fig. S8). To calibrate the optic fiber position with respect to the patch pipette we used a visible

laser (Fig. S8). Marks on the computer screen were used to keep the position of the fiber and the pipette

consistent for the different experiments. Cells under whole-cell voltage-clamp control were held at -30mV during

the experiment. To program fast pseudo-transient temperature changes, the patch pipette current was used to

read the temperature changes in real-time as the feedback to the laser diode controller (LDC-37620, ILX

Lightwave) to perform proportional-integral-derivative (PID) control of the driving current of the laser diode (*see*

Extended Methods). This laser-heating setup provides a rapid and precise heating rate on the order of 50°C

within tens of milliseconds, essential to provide both adequate temporal resolution and controllable steady-state

temperatures in the range of 35°C to 70°C to analyze the channel activation. Fig. S7A shows constant

temperature steps were achieved with a rising time constant of 34.2±3.3 ms, independent of the laser power.

The temperature jump associated with successive current pulses is precisely calculated by running an open

pipette calibration following the same current sequence at the end of each run (Fig. S6).

Temperature calibration. We used the resistance of the open pipette to measure the temperature jump

magnitudes following the equation $T=\{1/T_0 - R/E_a \times \ln(I/I_0)\}$ - 1, where R is the gas constant, T_0 and I_0 are

respectively room temperature and the corresponding electrode current at room temperature. The activation

energy (E_a) of the system corresponds to 3.84 kcal/mol as was established by Yao et al. (2009) for the pair of

solutions used in the recordings [101]. The equation describes the change in ion motility as a function of

temperature changes in the system. The current change was used as a feedback signal for a laser-diode

controller software coded in Labview that uses a proportional-integral-derivative (PID) control algorithm. To

account for the variability in the diameter between the different patch pipettes used in different experiments, the

instrument was calibrated before each experiment to assure comparable temperature jumps in each experiment,

adjusting the diode power outputs to the desired temperature accordingly.

Whole cell patch-clamp recordings. Cells were seeded at low density in a 30 mm culture dish (VWR)

containing round glass coverslips 48h post-transfection (Table S5, Fig. S5). Cells were first rinsed with D-PBS

at room temperature, trypsinized with 0.5 mL Accutase (Stemcell Technologies) and suspended in 4.5 mL pre-

warmed DMEM-FBS medium. Two hundred microliters of this cell suspension were mixed with 1.8 mL pre-

warmed DMEM-FBS medium, dispensed drop wise in the culture dish, and incubated for 24h at 30°C. In a typical

experiment, one glass cover slip was gently retrieved from the culture dish using sterile forceps, rinsed with a

recording solution using a Pasteur pipette and placed in the recording chamber (Fig. S6). The fluorescence of

mRuby-expressing cells was monitored to select bright, healthy, isolated cells for whole-cell patch clamping.

Experiments under the whole-cell configuration were carried out 72h after induction. The electrodes were

fabricated with borosilicate capillaries, using a horizontal micropipette puller P-1000 (Sutter Instrument, Novato,

CA, USA), and polished using a microforge (Narishige, Japan) to a final diameter between 2-4 um. The internal

electrode was filled with the following solution in mM: 126 CsCl, 10 EGTA, 10 HEPES, 1 EDTA, and 4 MgATP.

The extracellular recording solution contained 2 mM $CaCl_2$, 10 mM HEPES, 140 mM NaCl, pH 7.2 (adjusted with

NaOH). The electrode resistance ranged between 2-4 MΩ, and the Vjp was estimated at ~18 mV for the

recordings. The current traces were amplified using a MultiClamp 700B amplifier (Molecular Devices, Sunnyvale,

CA, USA). The amplified electrical signals were acquired with a Digidata 1440A using the pClamp10 software

(Molecular Devices, Sunnyvale, CA, USA). Series Resistance (Rs) was compensated in 80%, as well as the fast

and slow capacitive components of the current. The current density was fitted to the following Boltzmann

function:

$$I\_total = I\_leak\ e^{(-(\Delta H\_leak)/RT)} + (I\_max\ e^{(-(\Delta H\_i)/RT)})/(1 + e^{(-(\Delta G\_T)/RT)})$$

Whereby the first term ΔH_leak is the enthalpy change of the leak current. The second term accounts for the

channel activity, with $\Delta G = \Delta H - T\Delta S$ is the free energy change involved in the closed-open reaction, and ΔH_i

accounts for the linear temperature dependence of the ionic conductivity and leakage current [101]. The

corrected temperature current density (*I*) was used to calculate the equilibrium constant from the relative fraction

of the channel in the open conformation (Po), assuming a two-state model, where Po = I/Imax.

$$Po = 1/(1 + Keq^{-1})$$

where $\ln(K_{eq})$ = -(ΔH/RT) + (ΔS/R). Thus from the Van't Hoff plots ln(Keq) vs 1/T, the enthalpy and entropy

associated with the channel opening can be obtained [66].

Funding

This work was supported by a Mind Brain Behavior Interfaculty grant to NEP and MAL, an Alice and

Knut Wallenberg fellowship at the Broad Institute of MIT and Harvard to MAL, National Science

Foundation PHY-1411123 and PHY-1411445 to NEP and NY respectively, funding from Formas 2017-

01463 to UJ and from the Carl Tryggers Foundation (CTS 21:1309) to MAL. The funders had no role

in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

The authors acknowledge Andrew Allen for helpful discussions on TRP biochemistry, Xi-Shi for early

advice with HEK cell culture, Rhiannon Macrae for her valuable guidance, Feng Zhang for his support

and for giving access to resources at the Broad, Ian Orchard for providing *Rhodnius prolixus* antennae,

BEI resources (http://www.beiresources.org/) and Ellen M. Dotson for providing Rhodnius adults,

Rachel Gaudet for providing a rTRPV1 plasmid template and Pengyu Gu for providing a Drosophila

Dm-TRPA1-D (A10a, flybase PG) plasmid template.

Data Availability Statement

All datasets generated and analyzed for this study will be published as supplementary files

accompanying the paper.

References

[1] McKemy DD. 2007 Temperature sensing across species. *Pflugers Archiv : European journal of*
physiology **454**, 777-791. (10.1007/s00424-006-0199-6).

[2] Vriens J, Nilius B, Voets T. 2014 Peripheral thermosensation in mammals. *Nature Reviews*
Neuroscience **15**, 573-589. (10.1038/nrn3784).

[3] Xiao R, XU S. 2021 Temperature sensation: from molecular themosensors to neural circuits and
coding principles. *Annual review of physiology* **83**, 205-230.

[4] Castillo K, Diaz-Franulic I, Canan J, Gonzalez-Nilo F, Latorre R. 2018 Thermally activated TRP
channels: molecular sensors for temperature detection. *Physical Biology* **15**, 021001.
(10.1088/1478-3975/aa9a6f).

[5] Garrity P, Goodman M, Samuel AD, Sengupta P. 2010 Running hot and cold: behavioral
strategies, neural circuits, and the molecular machinery for thermotaxis in *C. elegans* and
Drosophila. *Genes and Development* **24**, 2365-2382.

[6] Klein M, Afonso B, Vonner AJ, Hernandez-Nunez L, Berck M, Tabone CJ, Kane EA, Pieribone
VA, Nitabach MN, Cardona A, et al. 2015 Sensory determinants of behavioral dynamics in
Drosophila thermotaxis. *Proceedings of the National Academy of Sciences* **112**, E220-E229.

[7] Clapham DE. 2003 TRP channels as cellular sensors. *Nature* **426**, 517-524.
(10.1038/nature02196).

[8] Venkatachalam K, Montell C. 2007 TRP channels. *Annu Rev Biochem* **76**, 387-417.
(10.1146/annurev.biochem.75.103004.142819).

[9] Baez-Nieto D, Castillo J, Dragicevic C, Alvarez O, Latorre R. Thermo-TRP channels: biophysics
of polymodal receptors.

[10] Fowler M, Montell C. 2013 Drosophila TRP channels and animal behavior. *Life Sci* **92**, 394-403.

[11] Peng G, Shi X, Kadowaki T. 2015 Evolution of TRP channels inferred by their classification in
diverse animal species. *Molecular Phylogenetics and Evolution* **84**, 145-157.
(doi.org/10.1016/j.ympev.2014.06.016).

[12] Matsuura H, Sokabe T, Kohno K, Tominaga M, Kadowaki T. 2009 Evolutionary conservation
and changes in insect TRP channels. *BMC Evolutionary Biology* **9**, 228. (10.1186/1471-2148-9-228).

[13] Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, Julius D. 1997 The
capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* **389**, 816-824.
(10.1038/39807).

[14] Gracheva EO, JF C-M, Carcacia-González JA, Ingolia N, Manno C, Aranguren C, Weissman J,
Julius D. 2011 Ganglion-specific splicing of TRPV1 underlies infrared sensation in vampire bats.

[15] Caterina MJ, Rosen TA, Tominaga M, Brake AJ, Julius D. 1999 A capsaicin-receptor
homologue with a high threshold for noxious heat. *Nature* **398**, 436-441. (10.1038/18906).

[16] Vriens J, Owsianik G, Hofmann T, Philipp Stephan E, Stab J, Chen X, Benoit M, Xue F,
Janssens A, Kerselaers S, et al. 2011 TRPM3 Is a Nociceptor Channel Involved in the Detection of
Noxious Heat. *Neuron* **70**, 482-494. (doi.org/10.1016/j.neuron.2011.02.051).

[17] Tracey WD, Wilson RI, Laurent G, Benzer S. 2003 painless, a Drosophila Gene Essential for
Nociception. *Cell* **113**, 261-273. (10.1016/S0092-8674(03)00272-1).

[18] Lee Y, Lee Y, Lee J, Bang S, Hyun S, Kang J, Hong S-T, Bae E, Kaang B-K, Kim J. 2005
Pyrexia is a new thermal transient receptor potential channel endowing tolerance to high
temperatures in Drosophila melanogaster. *Nature Genetics* **37**, 305. (10.1038/ng1513).

[19] Kang K, Panzano V, Chang E, Ni L, Dainis A, Jenkins A, Regna K, Muskavitch M, Garrity P.
2012 Modulation of TRPA1 thermal sensitivity enables sensory discrimination in Drosophila. *Nature*
481, 76-81.

[20] Liu L, Li Y, Wang R, Yin C, Dong Q, Hing H, Kim C, Welsh MJ. 2007 Drosophila hygrosensation
requires the TRP channels water witch and nanchung. *Nature* **450**, 294-298. (10.1038/nature06223).

[21] Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, Garrity PA. 2008 An
internal thermal sensor controlling temperature preference in Drosophila. *Nature* **454**, 217-220.
(10.1038/nature07001).

[22] Zhong L, Bellemer A, Yan H, Honjo K, Robertson J, Hwang Richard Y, Pitt Geoffrey S, Tracey
WD. 2012 Thermosensory and Nonthermosensory Isoforms of *Drosophila melanogaster*TRPA1
Reveal Heat-Sensor Domains of a ThermoTRP Channel. *Cell Reports* **1**, 43-55.
(10.1016/j.celrep.2011.11.002).

[23] Guntur A, Gy P, Takle K, Chen J, Yang X, Yang C-H. 2015 Drosophila TRPA1 isoforms detect
UV light via photochemical production of H2O2. *Proc Natl Acad Sci* **E5753-E5761**.

[24] Gu P, Gong J, Shang Y, Wang F, Ruppell KT, Ma Z, Sheehan AE, Freeman MR, Xiang Y. 2019
Polymodal Nociception in Drosophila Requires Alternative Splicing of TrpA1. *Curr Biol* **29**, 3961-
3973.e3966. (10.1016/j.cub.2019.09.070).

[25] Wang G, Qiu YT, Lu T, Kwon H-W, Jason Pitts R, Van Loon JJA, Takken W, Zwiebel LJ. 2009
Anopheles gambiae TRPA1 is a heat-activated channel expressed in thermosensitive sensilla of
female antennae. *European Journal of Neuroscience* **30**, 967-974. (10.1111/j.1460-
9568.2009.06901.x).

[26] Survery S, Moparthi L, Kjellbom P, Hogestatt E, Zygmunt PM, Johanson U. 2016 The N-terminal
Ankyrin Repeat Domain Is Not Required for Electrophile and Heat Activation of the Purified Mosquito
TRPA1 Receptor.

[27] Sato A, Sokabe T, Kashio M, Yasukochi Y, Tominaga M, Shiomi K. 2014 Embryonic
thermosensitive TRPA1 determines transgenerational diapause phenotype of the silkworm, Bombyx
mori. *Proc Natl Acad Sci U S A* **111**, E1249-1255. (10.1073/pnas.1322134111).

[28] Kohno K, Sokabe T, Tominaga M, Kadowaki T. 2010 Honey Bee Thermal/Chemical Sensor,
AmHsTRPA, Reveals Neofunctionalization and Loss of Transient Receptor Potential Channel
Genes. *The Journal of Neuroscience* **30**, 12219.

[29] Wang X, Li T, Kashio M, Xu Y, Tominaga M, Kadowaki T. 2018 HsTRPA of the Red Imported
Fire Ant, *Solenopsis invicta,* Functions as a Nocisensor and Uncovers the Evolutionary Plasticity of
HsTRPA Channels. *eNeuro* **5**, ENEURO.0327-0317.2018. (10.1523/ENEURO.0327-17.2018).

[30] Wigglesworth V. 1939 *The Principles of Insect Physiology (7 hardback editions; paperback*
edition 1984). London, UK, Methuen and Co., Ltd.

[31] Coura JR, Viñas PA. 2010 Chagas disease: a new worldwide challenge. *Nature* **465**, S6-S7.
(10.1038/nature09221).

[32] 2015 Chagas disease in Latin America: an epidemiological update based on 2010 estimates.
Wkly Epidemiol Rec **90**, 33-43.

[33] WHO. 2002 Control of Chagas disease : second report of the World Health Organization (WHO)
expert committee. (World Health Organization.

[34] Clayton J. 2010 Chagas disease 101. *Nature* **465**, S4-S5. (10.1038/nature09220).

[35] Ceccarelli S, Balsalobre A, Medone P, Cano ME, Gurgel Gonçalves R, Feliciangeli D, Vezzani
D, Wisnivesky-Colli C, Gorla DE, Marti GA, et al. 2018 DataTri, a database of American triatomine
species occurrence. *Scientific data* **5**, 180071-180071. (10.1038/sdata.2018.71).

[36] Wigglesworth B, Gillett J. 1934 The function of the antennae in *Rhodnius prolixus* (Hemiptera)
and the mechanism of orientation to the host. *J. Exp. Biol* **11**, 120-138.

[37] Mesquita RD, Vionette-Amaral RJ, Lowenberger C, Rivera-Pomar R, Monteiro FA, Minx P,
Spieth J, Carvalho AB, Panzera F, Lawson D, et al. 2015 Genome of *Rhodnius prolixus*, an insect
vector of Chagas disease, reveals unique adaptations to hematophagy and parasite infection.
Proceedings of the National Academy of Sciences **112**, 14936-14941.

[38] Barrozo R, Reisenman C, Guerenstein P, Lazzari C, Lorenzo MG. 2016 An inside look at the
sensory biology of triatomines. *Journal of Insect Physiology* **97**, 3-19. (doi:
10.1016/j.jinsphys.2016.11.003).

[39] Lazzari CR, Núñez J. 1989 The response to radiant heat and the estimation of the temperature
of distant sources in Triatoma infestans. *Journal of Insect Physiology* **35**, 525-529.

[40] Lazzari CR, Nunez J. 1989 Blood temperature and feeding-behavior in triatoma infestans
(heteroptera: Reduviidae). *Entomologia Generalis* **14**, 183-188.

[41] Guerenstein PG, Lazzari CR. 2009 Host-seeking: How triatomines acquire and make use of
information to find blood. *Acta tropica* **110**, 148-158. (10.1016/j.actatropica.2008.09.019).

[42] Zopf L, Lazzari C, Tichy H. 2014 Differential effects of ambient temperature on warm cell
responses to infrared radiation in the bloodsucking bug Rhodnius prolixus. *Journal of*
Neurophysiology **111**, 1341-1349.

[43] Zopf L, Lazzari C, Tichy H. 2014 Infrared detection without specialized infrared receptors in the
bloodsucking bug *Rhodnius prolixus*. *Journal of Neurophysiology* **112**, 1606.

[44] Latorre-Estivalis JM, Robertson HM, Walden KKO, Ruiz J, Gonçalves LO, Guarneri AA, Lorenzo
MG. 2017 The molecular sensory machinery of a Chagas disease vector: expression changes
through imaginal moult and sexually dimorphic features. *Scientific Reports* **7**, 40049.
(10.1038/srep40049).

[45] Latorre-Estivalis JM, Lorenzo MG. 2019 Molecular bases of sensory processes in kissing bugs,
vectors of Chagas disease. *Current Opinion in Insect Science* **34**, 80-84.
(10.1016/j.cois.2019.03.010).

[46] Latorre-Estivalis JM, Sterkel M, Ons S, Lorenzo MG. 2020 Transcriptomics supports local
sensory regulation in the antenna of the kissing-bug Rhodnius prolixus. *BMC Genomics* **21**, 101.
(10.1186/s12864-020-6514-3).

[47] Chaverra-Rodriguez D, Macias VM, Hughes GL, Pujhari S, Suzuki Y, Peterson DR, Kim D,
McKeand S, Rasgon JL. 2018 Targeted delivery of CRISPR-Cas9 ribonucleoprotein into arthropod
ovaries for heritable germline gene editing. *Nature Communications* **9**, 3008. (10.1038/s41467-018-
05425-9).

[48] Pereira J, Diogo C, Fonseca A, Bomfim L, Cardoso P, Santos A, Dittz U, Miranda K, de Souza
W, Gioda A, et al. 2020 Silencing of RpATG8 impairs the biogenesis of maternal autophagosomes in
vitellogenic oocytes, but does not interrupt follicular atresia in the insect vector *Rhodnius prolixus*.
PLOS Neglected Tropical Diseases **14**, e0008012. (10.1371/journal.pntd.0008012).

[49] Neely GG, Keene AC, Duchek P, Chang EC, Wang Q-P, Aksoy YA, Rosenzweig M, Costigan
M, Woolf CJ, Garrity PA, et al. 2011 TrpA1 Regulates Thermal Nociception in Drosophila. *PLOS*
ONE **6**, e24343. (10.1371/journal.pone.0024343).

[50] Cattaneo AM, Bengtsson JM, Montagné N, Jacquin-Joly E, Rota-Stabelli O, Salvagnin U, Bassoli A, Witzgall P, Anfora G. 2016 TRPA5, an ankyrin subfamily insect TRP channel, is expressed in antennae of *Cydia pomonella* (Lepidoptera: Tortricidae) in multiple splice variants. *Journal of Insect Science* **16**, 83-83. (10.1093/jisesa/iew072).

[51] Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Zidek A, Potapenko A, et al. 2021 Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589. (doi:10.1038/s41586-021-03819-2).

[52] Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G, Laydon A, et al. 2021 AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Research* **50**, D439-D444. (10.1093/nar/gkab1061).

[53] Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022 ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682. (10.1038/s41592-022-01488-1).

[54] Paulsen CE, Armache JP, Gao Y, Cheng Y, Julius D. 2015 Structure of the TRPA1 ion channel suggests regulatory mechanisms. *Nature* **520**, 511-517. (10.1038/nature14367).

[55] Wang X, Li Y, Wei H, Yang Z, Luo R, Gao Y, Zhang W, Liu X, Sun L. 2023 Molecular architecture and gating mechanisms of the Drosophila TRPA1 channel. *Cell Discov* **9**, 36. (10.1038/s41421-023-00527-1).

[56] Zhao J, Lin King JV, Paulsen CE, Cheng Y, Julius D. 2020 Irritant-evoked activation and calcium modulation of the TRPA1 receptor. *Nature*. (10.1038/s41586-020-2480-9).

[57] Gaudet R. 2008 A primer on ankyrin repeat function in TRP channels and beyond. *Molecular bioSystems* **4**, 372-379. (10.1039/b801481g).

[58] Ladrón-de-Guevara E, Dominguez L, Rangel-Yescas GE, Fernández-Velasco DA, Torres-Larios A, Rosenbaum T, Islas LD. 2020 The Contribution of the Ankyrin Repeat Domain of TRPV1 as a Thermal Module. *Biophys J* **118**, 836-845. (10.1016/j.bpj.2019.10.041).

[59] Zheng W, Qin F. 2015 A combined coarse-grained and all-atom simulation of TRPV1 channel gating and heat activation. *J Gen Physiol* **145**, 443-456. (10.1085/jgp.201411335).

[60] Cordero-Morales JF, Gracheva EO, Julius D. 2011 Cytoplasmic ankyrin repeats of transient receptor potential A1 (TRPA1) dictate sensitivity to thermal and chemical stimuli. *Proceedings of the National Academy of Sciences* **108**, E1184.

[61] Moparthi L, Survery S, Kreir M, Simonsen C, Kjellbom P, Högestätt ED, Johanson U, Zygmunt PM. 2014 Human TRPA1 is intrinsically cold- and chemosensitive with and without its N-terminal ankyrin repeat domain. *Proc Natl Acad Sci U S A* **111**, 16901-16906. (10.1073/pnas.1412689111).

[62] Liénard M, Bernard GD, Allen A, Lassance J, Song S, Childers R, Yu N, Ye D, Stephenson A, Valencia-Montoya W, et al. 2021 The evolution of red colour vision is linked to coordinated rhodopsin tuning in lycaenid butterflies. *PNAS* **118** e2008986118. (doi.org/10.1073/pnas.2008986118).

[63] Yao J, Liu B, Qin F. 2010 Kinetic and energetic analysis of thermally activated TRPV1 channels.

[64] Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, Julius D. 1997 The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* **389**, 816-824. (10.1038/39807).

[65] Liu B, Qin F. 2017 Single-residue molecular switch for high-temperature dependence of vanilloid receptor TRPV3. *Proc Natl Acad Sci U S A* **114**, 1589-1594. (10.1073/pnas.1615304114).

[66] Latorre R, Brauchi S, Orta G, Zaelzer C, Vargas G. 2007 ThermoTRP channels as modular
proteins with allosteric gating. *Cell Calcium* **42**, 427-438.

[67] Baez D, Raddatz N, Ferreira G, Gonzalez C, Latorre R. 2014 Chapter Three - Gating of
Thermally Activated Channels. In *Current Topics in Membranes* (eds. L. D. Islas & F. Qin), pp. 51-
87, Academic Press.

[68] Caterina MJ, Julius D. 2001 The Vanilloid Receptor: A Molecular Gateway to the Pain Pathway.
Annual Review of Neuroscience **24**, 487-517. (10.1146/annurev.neuro.24.1.487).

[69] Güler A, Lee H, Lida T, Shimizu I, Tominaga M, Caterina M. 2002 Heat-evoked activation of the
ion channel, TRPV4.

[70] Smith GD, Gunthorpe MJ, Kelsell RE, Hayes PD, Reilly P, Facer P, Wright JE, Jerman JC,
Walhin JP, Ooi L, et al. 2002 TRPV3 is a temperature-sensitive vanilloid receptor-like protein. *Nature*
418, 186-190. (10.1038/nature00894).

[71] Diver MM, Lin King JV, Julius D, Cheng Y. 2022 Sensory TRP Channels in Three Dimensions.
Annu Rev Biochem **91**, 629-649. (10.1146/annurev-biochem-032620-105738).

[72] Laursen WJ, Anderson EO, Hoffstaetter LJ, Bagriantsev SN, Gracheva EO. 2015 Species-
specific temperature sensitivity of TRPA1.

[73] Tsai C-C, Childers RA, Nan Shi N, Ren C, Pelaez JN, Bernard GD, Pierce NE, Yu N. 2020
Physical and behavioral adaptations to prevent overheating of the living wings of butterflies. *Nature*
Communications **11**, 551. (10.1038/s41467-020-14408-8).

[74] Schmitz H, Trenner S, Hofmann MH, Bleckmann H. 2000 The ability of *Rhodnius prolixus*
(Hemiptera; Reduviidae) to approach a thermal source solely by its infrared radiation. *Journal of*
Insect Physiology **46**, 745-751.

[75] Flores GB, Lazzari CR. 1996 The role of the antennae in Triatoma infestans: Orientation
towards thermal sources. *Journal of Insect Physiology* **42**, 433-440. (doi.org/10.1016/0022-
1910(95)00137-9).

[76] Ferreira RA, Lazzari CR, Lorenzo MG, Pereira MH. 2007 Do Haematophagous Bugs Assess
Skin Surface Temperature to Detect Blood Vessels? *PLOS ONE* **2**, e932.
(10.1371/journal.pone.0000932).

[77] Zermoglio PF, Latorre-Estivalis JM, Crespo JE, Lorenzo MG, Lazzari CR. 2015
Thermosensation and the TRPV channel in Rhodnius prolixus. *Journal of Insect Physiology* **81**, 145-
156. (doi.org/10.1016/j.jinsphys.2015.07.014).

[78] Voets T, Talavera K, Owsianik G, Nilius B. 2005 Sensing with TRP channels. *Nature Chemical*
Biology **1**, 1-8.

[79] Yin C, Shen G, Guo D, Wang S, Ma X, Xiao H, Liu J, Zhang Z, Liu Y, Zhang Y, et al. 2015
InsectBase: a resource for insect genomes and transcriptomes. *Nucleic Acids Research* **44**, D801-
D807. (10.1093/nar/gkv1204).

[80] Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates L S, Marygold SJ, Matthews BB,
Millburn G, Antonazzo G, Trovisco V, et al. 2018 FlyBase 2.0: the next generation. *Nucleic Acids*
Research **47**, D759-D765. (10.1093/nar/gky1003).

[81] Elsik CG, Tayal A, Diesh CM, Unni DR, Emery ML, Nguyen HN, Hagen DE. 2015 Hymenoptera
Genome Database: integrating genome annotations in HymenopteraMine. *Nucleic Acids Research*
44, D793-D800. (10.1093/nar/gkv1208).

[82] Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997 Gapped
BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids*
Research **25**, 3389-3402.

[83] Poelchau M, Childers C, Moore G, Tsavatapalli V, Evans J, Lee C-Y, Lin H, Lin J-W, Hackett K.
2014 The i5k Workspace@NAL—enabling genomic data access, visualization and curation of
arthropod genomes. *Nucleic Acids Research* **43**, D714-D719. (10.1093/nar/gku983).

[84] Emms D, Kelly S. 2019 OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* **20**, 238. (doi:10.1186/s13059-019-1832-y).

[85] Katoh K, Standley D. 2013 MAFFT Multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780.
(doi:10.1093/molbev/mst010).

[86] Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015 IQ-TREE: A fast and effective
stochastic algorithm for estimating maximum likelihood phylogenies. *Mol Biol Evol* **32**, 268-274.

[87] Trifinopoulos J, Nguyen L-T, von Haeseler A, Minh BQ. 2016 W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research* **44**, W232-W235.
(10.1093/nar/gkw256).

[88] Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS. 2017 ModelFinder: Fast
model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587-589.

[89] Yu G. 2020 Using ggtree to Visualize Data on Tree-Like Structures. *Current Protocols in*
Bioinformatics **69**, e96. (10.1002/cpbi.96).

[90] Zhang H, Gao S, Lercher M, Hu S, Chen W-H. 2012 EvolView, an online tool for visualizing,
annotating and managing phylogenetic trees. *Nucleic Acids Res* **40**, W569-W572.

[91] Martin M. 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 **17**, 3. (10.14806/ej.17.1.200).

[92] Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019 Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**, 907-915. (10.1038/s41587-
019-0201-4).

[93] Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015 StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* **33**, 290-
295. (10.1038/nbt.3122).

[94] Bryant DM, Johnson K, Tia D, Tickle T, Couger MB, Payzin-Dogru D, Lee T, Leigh N, Kuo T, .,
et al. 2017 A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of Limb
Regeneration Factors. *Cell rep* **18**, 762-776.

[95] Lee E, Helt GA, Reese JT, Munoz-Torres MC, Childers CP, Buels RM, Stein L, Holmes IH, Elsik
CG, Lewis SE. 2013 Web Apollo: a web-based genomic annotation editing platform. *Genome*
Biology **14**, R93. (10.1186/gb-2013-14-8-r93).

[96] Soneson C, Matthes KL, Nowicka M, Law CW, Robinson MD. 2016 Isoform prefiltering
improves performance of count-based methods for analysis of differential transcript usage. *Genome*
Biology **17**, 12. (10.1186/s13059-015-0862-3).

[97] Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
Markowitz S, Duran C, et al. 2012 Geneious Basic: An integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647-1649.

[98] The PyMOL Molecular Graphics System, Version 1.8, Schrödinger, LLC. (

[99] Holm L. 2020 DALI and the persistence of protein shape. *Protein Sci* **29**, 128-140.
(10.1002/pro.3749).

[100] Liénard MA, Valencia-Montoya WA, Pierce NE. 2022 Molecular advances to study the
function, evolution and spectral tuning of arthropod visual opsins. *Philosophical Transactions of the*
Royal Society B: Biological Sciences **377**. (doi:10.1098/rstb.2021.0279).

[101] Yao J, Liu B, Qin F. 2009 Rapid Temperature Jump by Infrared Diode Laser Irradiation for
Patch-Clamp Studies. *Biophysical Journal* **96**, 3611-3619.

Extended data

Convergent evolution of noxious heat sensing by TRPA5, a novel class of heat sensor in *Rhodnius prolixus*

Marjorie A. Liénard[1,2,3*], David Baez-Nieto[4*], Cheng-Chia Tsai[5], Wendy A. Valencia-Montoya[2], Balder

Werin[6], Urban Johanson[6], Jean-Marc Lassance[2,7], Jen Q. Pan[4], Nanfang Yu[5], Naomi E. Pierce[2]

[1]Department of Biology, Lund University, Lund, Sweden

[2]Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology, Harvard University,
Cambridge, MA 02138, USA

[3]Broad Institute, Cambridge, MA 02142, USA

[4]Stanley Center for Psychiatric Research, Broad Institute, Cambridge, MA 02142, USA

[5]Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027, USA

[6]Division of Biochemistry and Structural Biology, Department of Chemistry, Lund University, Lund, Sweden

[7]GIGA-Research, University of Liège, Liège, Belgium

This document contains:

Supplementary Methods

Supplementary Figures S1 to S8

SI Tables S1 to S5 are in a separate file

Supplementary Methods

PID control

A PID (proportional-integral-derivative) control is the typical way to adjust the output according to the input

reading in real-time without knowing most of the environmental parameters. The idea is described in the following

function:

891
$$Output(t) = K_p Err(t) + K_i \int_0^t Err(\tau)d\tau + K_d \frac{dErr(t)}{dt}$$

892
$$Err(t) = Input(t) - Setpoint$$

with $Output(t)$ being the laser power, $Input(t)$ being the temperature, and $Setpoint$ being the desired value.

The $Output(t)$ is determined by $Err(t)$, which is the difference between $Input(t)$ and $Setpoint$.

There are 3 terms in this equation, the first term is the proportional term. This term varies linearly with $Err(t)$.

For instance, when the temperature reaches the setpoint, this term decreases, and when the temperature

exceeds the setpoint, this term becomes negative to bring the temperature back to the setpoint.

The second term is the integral term. This term provides a gradually increasing offset and this offset will stabilize

when the temperature stabilizes to setpoint, where $Err(t) \to 0$.

The third term is the derivative term. This term estimates the required change of output by watching the inertia

of $Err(t)$. For instance, when the temperature reaches the setpoint, the proportional term gives 0 and the integral

term gives a stabilized value, but if the temperature is still increasing, this term will decrease the output to prevent

the temperature from exceeding the setpoint in the next time interval.

Open pipette current measurements

The temperature is measured by monitoring the current through an open patch pipette, which means there are

no cells but only water, assuming the thermal property of the cell is the same as water. First, a series of open-

pipette measurements with different laser powers and power waveforms is performed and used to calculate the

temperature evolution from real-time current in the patch-clamp recording pipette. The patch-clamp experiment

is conducted by applying the same laser powers and waveforms to the cell (Fig. S4)

Fiber preparation

The attenuation coefficient of water at the wavelength of 1460 nm is ~3000 cm^{-1} which means the absorption

of the laser power through a $100 nm$ thick water layer is over 40%. Using the fiber above the water surface, the

laser would thus deliver most of its power in the upper water layer. The temperature of water above the targeted

cell would be much higher than the cell temperature itself, and the temperature change would not be confined

to a single cell due to conduction and convection in the water. To resolve this, we cut the fiber with a diamond

blade under a stereomicroscope and stripped the fiber which allowed us to place the fiber tip in the water layer

directly above the cell in the recording chamber (Fig. S5). The relative position of the fiber tip and cell was

adjusted using patch-clamp rig microscope.

Supplementary Figures

Figure S1. Phylogeny of insect TRPA channels

Figure S1. Phylogeny of insect TRPA channels. The Maximum-likelihood phylogeny of amino acid sequences
includes representative ion channel members of the TRPA1, Pain, Pyrexia, HsTRPA, TRPA5, Wtw subfamilies.
Accession numbers are listed in Table S1. The tree was inferred in IQ-TREE v1.6.11 using ModelFinder,
Ultrafast Bootstrap (UFboot), 1000 replicates, using a best-fit model JTT+F+I+G4 measured by the Bayesian
information criterion (BIC). Branches were assigned Shimodaira-Hasegawa approximate likelihood ratio test
(SH-aLRT) and UFboot supports. The tree was visualized in Rstudio (2021.09.2) using ggtree (Script provided).
Species represented are **Anoplura**: *Pediculus humanus* (Phu, head louse), **Coleoptera**: *Tribolium castaneum*
(Tca, red flour beetle), *Dendroctonus ponderosae* (Dpo, Mountain pine beetle), *Photynus pyralis* (Ppy, Common
Eastern Firefly), *Agrilus planipennis* (Apl, Emerald ash borer), *Anoplophora glabripennis* (All, Asian longhorned
beetle), *Aethina tumida* (Aut, Small hive beetle), *Onthophagus taurus* (Ota, bull dung beetle), **Diptera**:
Drosophila melanogaster (Dme, fruitfly), *Mayetiola destructor* (Mde, Hessian fly), *Culex quinquefasciatus* (Cqu,
Southern house mosquito), *Anopheles stephensi* (Ast), *Anopheles gambiae* (Aga), *Anopheles darlingi* (Ada),
Bactrocera cucurbitae (Bcu, Melon fly), *Ceratitis capitata* (Cca, Mediterranean fruit fly), *Musca domestica* (Mdo,
housefly), *Lutzomyia longipalpis (Llo, Sandfly), Phlebotomus papatasi* (Ppa, Sandfly), *Stomoxys calcitrans* (Sca,
Barn fly), **Hemiptera:** *Acyrthosiphon pisum* (Api, pea aphid)*, Nilaparvata lugens* (Nlu, Brown planthopper),
Diaphorina citri (Dci, Asiatic citrus psyllid), *Cimex lectularius* (Cle, Bed bug)*, Triatoma infestans* (Tin, Winchuka)
Rhodnius prolixus (Rp, Kissing bug)*, **Hymenoptera:** Apis mellifera* (Ame, Western honeybee) *, Bombus*
terrestris (Bte, Buff-tailed bumblebee)*, Megachile rotundata* (Mro, Alfalfa leaf cutting bee), *Nasonia vitripenis*
(Nvi, Jewel wasp), *Harpegnathos saltator* (Hsa, Jumping ant)*, Linepithema humile* (Lhu, Argentine ant),
Campanatus floridanus (Cfl, Florida carpenter ant), *Pogonomyrmex barbatus* (Pba, red harvester ant), *Atta*
cephalotes (Ace, Leafcutter ant), *Acromyrmex echinatior* (Aec, Panamian leafctutter ant), *Solenopsis invicta*
(Sin, Red imported fire ant)*, Vollenhovia emeryi* (Vem, ant), *Athalia rosae* (Aro, Turnip sawfly), *Cerapachys biroi*
(Cbi, Clonal raider ant), *Ceratosolen solmsi* (Cso, Fig wasp), *Wasmannia auropunctata* (Wau, Electric ant),
Isoptera: *Zootermopsis nevadensis* (Zne, Dampwood termite), **Lepidoptera**: *Bombyx mori* (Bmo, Silkmoth)*,*
Chilo suppressalis (Csu, Asiatic rice borer), *Danaus plexippus* (Dpl, Monarch butterfly), *Heliconius melpomene*
(Hme, Postman butterfly), *Plutella xylostella* (Diamondback moth), *Manduca sexta* (Mse, Tobacco hornworm)
Cydia pomonella (Cpo, Codling moth), **Odonata**: *Ischnura elegans* (Iel, Bluetail damselfly).

Figure S2. Phylogeny of insect TRPA5 channels

Figure S2. Phylogeny of insect TRPA5 channels. TRPA5 channels are present across insect Orders but
Diptera. The Maximum-likelihood phylogeny of amino acid sequences includes representative ion channel
members of the TRPA5 subfamily. The tree was inferred in IQ-TREE v1.6.11 using ModelFinder, Ultrafast
Bootstrap (UFboot), 1000 replicates, using a best-fit model JTT+F+I+G4 measured by the Bayesian information
criterion (BIC). Branches were assigned Shimodaira-Hasegawa approximate likelihood ratio test (SH-aLRT) and
UFboot supports. The tree was visualized in Rstudio (2021.09.2) using ggtree (script provided). Species
abbreviations are as follows: **Coleoptera:** *Tribolium castaneum* (Tca, red flour beetle), *Photynus pyralis* (Ppy,
Common Eastern Firefly), *Anoplophora glabripennis* (Agl, Asian longhorned beetle), *Aethina tumida* (Aut, Small
hive beetle), *Onthophagus taurus* (Ota, bull dung beetle); **Hemiptera**: *Bemisia tabaci* (Bta, Silverleaf Whitefly),
Cimex lectularius (Cle, Bed bug)*, Halyomorpha halys* (Hha, Brown marmorated stinkbug), *Nilaparvata lugens*
(Nlu, Brown planthopper), *Rhodnius prolixus* (Rp, Kissing bug); **Hymenoptera**: *Apis mellifera* (Ame, Western
honeybee) *, Athalia rosae* (Aro, Turnip sawfly), *Bombus terrestris* (Bte, Buff-tailed bumblebee)*, Megachile
rotundata* (Mro, Alfalfa leaf cutting bee), *Nasonia vitripennis* (Nvi, Jewel wasp), *Harpegnathos saltator* (Hsa,
Jumping ant)*, Linepithema humile* (Lhu, Argentine ant), *Campanatus floridanus* (Cfl, Florida carpenter ant),
Pogonomyrmex barbatus (Pba, red harvester ant), *Atta cephalotes* (Ace, Leafcutter ant), *Acromyrmex echinatior*
(Aec, Panamian leafcutter ant), *Solenopsis invicta* (Sin, Red imported fire ant)*, Vollenhovia emery* (Vem, ant),
Ooceraea biroi (Cbi, Clonal raider ant), *Wasmannia auropunctata* (Wau, Electric ant), *Polisted dominula* (Pdo,
European paper wasp), *Neodiprion lecontei* (Nle, Red-headed pine sawfly), *Cephus cinctus* (Cci, Wheat stem
sawfly); **Isoptera**: *Zootermopsis nevadensis* (Zne, Dampwood termite); **Lepidoptera:** *Bombyx mori* (Bmo,
Silkmoth)*, Chilo suppressalis* (Csu, Asiatic rice borer), *Danaus plexippus* (Dpl, Monarch butterfly), *Heliconius
melpomene* (Hme, Postman butterfly), *Plutella xylostella* (Pxy, Diamondback moth), *Manduca sexta* (Mse,
Tobacco hornworm), *Cydia pomonella* (Cpo, Codling moth), *Vanessa tameamea* (Vta, Kamehameha butterfly),
Trichoplusia ni (Tni, Cabbage looper), *Spodoptera litura* (Sli, Tobacco cutworm), *Galleria mellonella* (Gmo,
Greater wax moth), **Odonata**: *Ischnura elegans* (Iel, damselfly), **Thysanoptera**: *Frankliniella occidentalis* (Foc,
thrips). Capital letters above branches represent the amino acid residues at the ion channel selectivity filter (see
Fig. 2).

Figure S3. qCPR analysis

Figure S3. Differential expression pattern of Rp-TRPA5$_2$ transcripts isolated from *R. prolixus* female sensory
tissues and monitored by quantitative PCR. The mean relative expression scores were calculated from raw cycle
threshold (Ct) values (±SEM, n= 9) (Table S5) and boxplots were visualized in R (script provided). Log$_2$ fold
change expression values are shown relative to the *Rhodnius* housekeeping gene (Actin) and calibrated to
expression in Antenna.

Fig. S4. DALI Analysis

995
Figure S4. DALI Analysis. Pairwise structural alignment using the Dali server
(http://ekhidna2.biocenter.helsinki.fi/dali/), of different monomers against Rhodnius TRPA5$_2$. **A.** Rhodnius
TRPA1, **B**. Rhodnius Painless, **C.** Rhodnius Waterwitch, **D.** Drosophila TRPA1, **E.** Drosophila Painless, **F.**
Drosophila Waterwitch, **G.** Drosophila Pyrexia. The red circles in **G** indicate areas where the Pyrexia monomer
stands out in that the pore region and the voltage sensor domain are less similar to TRPA5$_2$, but the ARD is
more similar to TRPA5$_2$.

Figure S5. TRP whole cell and surface expression.

Figure S5. A-B. **Rp-TRPA5$_2$ anti-FLAG signals in whole cell lysates**. Untransfected HEK293T cells (lanes
1,4) and HEK293T cells transfected with 2.5 µg pcDNA-FLAG-T2A-mRuby plasmid DNAs for Rp-TRPA5$_2$ (A,
lanes 2,3), and fruitly Dmel-TRPA1-D (B, lane 5). Cells were collected 72 hrs after transfection. The protein
ladder image taken from the same membrane is juxtaposed to the left of the immunoblot. One and two asterisks
represent predicted dimeric and tetrameric TRP forms, respectively. The predicted monomeric MW is indicated
with a black arrowhead: Rp-TRPA5$_2$, 127.78 kDa; fruit fly Dmel-TRPA1-D 138.82 kDa. **C.** Surface expression
analysis of Rp-TRPA5$_2$. Biotinylated surface protein eluates were run in parallel wells on the same SDS-page
gel to probe TRP (left) and ATPase (right). Anti-FLAG levels in surface protein fraction are from non-transfected
HEK293T cells (lane 1), cells expressing rTRPV1 (lane 2), and cells expressing Rp-TRPA5$_2$ (lane 3). Lanes 4
to 6 are the corresponding anti-ATPase biotin-surface fraction from non-transfected HEK293T cells (lane 4),
cells expressing rTRPV1 (lane 5), and cells expressing Rp-TRPA5$_2$ (lane 6). One and two asterisks represent
predicted dimeric and tetrameric TRP forms, respectively. The predicted monomeric MW is indicated with a black
arrowhead (rTRPV1, 94.95 kDa; Rp-TRPA5$_2$, 127.78 kDa).

1016

Figure S6. Schematics of the functional assay workflow and optical fiber-patch clamp recording setup

Figure S6. Schematic drawing of the experimental workflow used to characterize *Rhodnius prolixus*

thermo TRPA5$_2$ ion channel. A. The full TRPA5$_2$ ORF sequence was amplified from *Rhodnius prolixus* antenna

and gene-specific primers, sequenced-verified, then codon-optimized for mammalian expression, and subcloned

in a custom-made pFRT-TO-FLAG-T2A-mRuby2 expression cassette under expression of the CMV promoter

(1) **(B)**. **C**. HEK293T cells seeded at low density were transiently transfected with plasmid DNA-lipid complexes

and incubated at 37°C for 48h to allow TRP surface membrane expression. Monomeric mRuby2 fluorescent

protein (mRuby2) was co-expressed as a cytoplasmic marker. 48h post-transfection, cells were prepared for

patch-clamp recording by seeding in a 30-mm^2 culture dish overlaid with round glass cover slips. Cells were

incubated at 30°C. **D**. Electrophysiology recordings took place after 24h to 48h using an optical fiber-based setup

adapted after Yao et al 2009 (2), designed to couple manual patch clamp recordings with fiber optics as a mean

to provide controllable optical and thermal stimulations to individual cells expressing candidate thermosensitive

receptor proteins. The setup consists of a fiber launch system combining a high-power optical fiber tuned to

near-infrared wavelengths (λc =1460 nm (+/-20 nm), Po= 4.8 watts), a visible alignment laser (red), and a laser

diode controller, forming a PID control loop using the patch clamp current as the feedback signal. **E**. During the

experiment, a laser spot is aligned with one single patched cell (see Fig. S8) stably expressing the membrane

receptor protein of interest in the cover slip placed in the recording chamber.

Figure S7. Validation with rTRPV1 and Dm-TRPA1-D

Figure S7. Validation with rTRPV1 and Dm-TRPA1-D. **A.** Time course of the ionic current through the open
pipette at -10 mV holding voltage when a current of 0.50A (green trace) and 0.63A (red trace) was applied to the
laser diode. The temperature jumps for those currents correspond to 32.1°C and 59.7°C, from 22.6°C,
respectively. Both temperature jumps can be fitted by a mono-exponential function with a time constant of ~35
ms (black line). **B.** Time course comparison between the current through the open pipette (magenta trace) and
a HEK293T cell expressing rTRPV1 channels under voltage clamp at -30 mV (orange trace), in response to a
temperature jump from 22.6°C to 59.7°C (0.63A). The current through rTRPV1 channels is three times slower
than the laser kinetics with a time constant of ~100 ms. **C.** Representative heat-activated current traces of a
HEK293T cell expressing rTRPV1 under voltage clamp (holding voltage of -30 mV). The currents were elicited
by temperature jumps from room temperature (22.6°C) up to 59.7°C with a duration of 700 ms. **D.** Fraction of
rTRPV1 channels in the open state (Open probability, Po) as a function of the temperature. The Po vs 1/T was
fitted to a Boltzman function (red line) with the midpoint of activation (T$_{0.5}$) reached at 51.6°C. The Van't Hoff plot

estimates for rTRPV1 provides an activation enthalpy of the endothermic transition at 88.3±9.4 kcal/mol and an

entropic change associated to the temperature activation process at 271±28 cal/mol*K. at -30 mV as a function

of the temperature. These values in our new expression cassette are very close to previously published values

of ΔH = 85 kcal/mol, $T_{0.5}$ = 45.6°C (2,3). **E.** Representative heat-activated current traces of a HEK293 cell

expressing Dm-TRPA1-D channels under voltage clamp (holding voltage of -30 mV). The currents were elicited

by temperature jumps from room temperature (19.4°C) up to 63.5°C with a duration of 700 ms. **F.** Fraction of

Dm-TRPA1-D channels in the open state (Open probability, Po) as a function of the temperature. The Po vs 1/T

was fitted to a Boltzman function (red line) with the midpoint of activation ($T_{0.5}$) reached at 53.5°C, the Van't Hoff

plot estimates for Dm-TRPA1-D an activation enthalpy of the endothermic transition at 68.7±13.1 kcal/mol and

an entropic change associated to the temperature activation process at 211±40 cal/mol·K at -30 mV. The

stationary current at the end of the temperature pulse (last 100 ms) was used to calculate all the thermodynamics

parameters. Data are presented as means ± standard deviation.

Figure S8.

1060

Figure S8. Open pipette temperature calibration and optic fiber positioning prior to each cell patch clamp

experiment A. Schematic representation of open pipette temperature calibration. First the open pipette is

positioned under the center of the laser using the reference marks on the screen. Then we proceed to record

the current through the open pipette elicited as a function of a series of near IR laser pulses. With these current

traces we can estimate the temperature jumps magnitude that will be applied later to the target cell. After this

the target cell is positioned in the center of the laser beam using the microscope stage translation (the optic fiber

stays fixed). Once the cell is in the right position, we use the same pipette used for the calibration to carry out

the electrophysiological recording, applying the same set of pulses used in the calibration **B.** Histogram of the

time constants from the exponential fit to the open pipette current traces from the temperature jumps used in the

rTRPV1 experiments (n=56), the mean time constant was 33.3±1.8 ms. **C.** View of HEK293T cells seeded at

low density on a glass cover in the recording chamber on the patch-clamp rig station. **D.** After finding a target

cell co-expressing mRuby2 as a fluorescent marker (see Fig. 1B), with the help of an automatized

micromanipulator, the optic fiber is placed in the recording solution using the reference marks on the computer

screen (not shown here) defined during the calibration with the visible laser. The depth in the solution is adjusted

so that the fiber is directly above a single cell in the recording medium. **E.** The field of view is changed to align

the patch clamp electrode to the reference marks. **F.** The target cell expressing mRuby2 is patched with the

recording electrode, now visible on the left side, whereas the fiber is directly above the target cell. The relative

position between the recording pipette and the fiber was established using a visible laser during the setup of the

system, and is constant for all the experiments.

Extended data References

(1) Liénard MA, Valencia-Montoya WA, Pierce NE. 2022 Molecular advances to study the function, evolution
and spectral tuning of arthropod visual opsins. *Philosophical Transactions of the Royal Society B: Biological*
Sciences 377. (doi:10.1098/rstb.2021.0279).

(2) Yao J, Liu B, Qin F. 2010 Kinetic and energetic analysis of thermally activated TRPV1 channels. *Biophys J*
99, 1743-1753. (10.1016/j.bpj.2010.07.022).

(3) Liu B, Hui K,Qin F. 2003 Thermodynamics of Heat Activation of Single Capsaicin Ion Channels VR1
Biophys J. 85(5): 2988–3006. (10.1016/S0006-3495(03)74719-5)