Title: Coupling and uncoupling of midline morphogenesis and cell flow in amniote gastrulation

Authors: Rieko Asai[1], Vivek N. Prakash[2], Shubham Sinha[2], Manu Prakash[3], Takashi Mikawa[1]*

Affiliations: [1]Cardiovascular Research Institute, University of California, San Francisco. San Francisco, California 94158, USA. [2]Department of Physics, University of Miami, Coral Gables, Florida 33146, USA. [3]Department of Bioengineering, Stanford University, Stanford, California 94305, USA.

*Correspondence to: takashi.mikawa@ucsf.edu

Summary: Large-scale cell flow characterizes gastrulation in animal development. In amniote gastrulation, particularly in avian gastrula, a bilateral vortex-like counter-rotating cell flow, called 'polonaise movements', appears along the midline. Here, through experimental manipulations, we addressed relationships between the polonaise movements and morphogenesis of the primitive streak, the earliest midline structure in amniotes. Suppression of the Wnt/planar cell polarity (PCP) signaling pathway maintains the polonaise movements along a deformed primitive streak. Mitotic arrest leads to diminished extension and development of the primitive streak and maintains the early phase of the polonaise movements. Ectopically induced Vg1, an axis-inducing morphogen, generates the polonaise movements, aligned to the induced midline, but disturbs the stereotypical cell flow pattern at the authentic midline. Despite the altered cell flow, induction and extension of the primitive streak are preserved along both authentic and induced midlines. Finally, we show that ectopic axis-inducing morphogen, Vg1, is capable of initiating the polonaise movements without concomitant PS extension under mitotic arrest conditions. These results are consistent with a model wherein primitive streak morphogenesis is required for the maintenance of the polonaise movements, but the polonaise movements are not necessarily responsible for primitive streak morphogenesis. Our data describe a previously

undefined relationship between the large-scale cell flow and midline morphogenesis in

gastrulation.

Introduction

Large-scale cell flow during gastrulation is an evolutionarily conserved biological phenomenon

in embryogenesis (1-3). During this period, animals with bilateral symmetry, called bilaterians,

initiate development of three germ layers (i.e. the ectoderm, mesoderm and endoderm) and

midline structures (e.g. the notochord and primitive streak) along the midline axis (4-6). The

large-scale cell flow has been linked to early embryonic morphogenesis, including axial

structures, through rearrangement of cells and/or transportation of signaling molecules (2, 3).

Coupling of the large-scale cell flow and body axis morphogenesis has been extensively studied

in invertebrate bilaterians, particularly insect models (7-9). However, in vertebrates, our

understanding of whether and how the large-scale cell flow and midline morphogenesis are

coupled is limited.

Vertebrates are classified into two groups: non-amniotes (e.g. fish and amphibians) and amniotes

(e.g. birds and mammals) (1). In non-amniotes, particularly amphibians, the large-scale cell flow

is coupled with formation of the notochord through Wnt/planar cell polarity (PCP) pathway-

regulated convergent extension, which does not depend on mitosis (10, 11). In amniotes,

particularly in avian gastrula (i.e. embryonic disc), a bilateral vortex-like counter-rotating cell

flow, termed 'polonaise movements', occurs within the epiblast along the midline axis, prior to

and during primitive streak (PS) formation (12-14). The PS is an evolutionarily unique

embryonic structure in amniotes, which is not only an organizing center for gastrulation but also

the earliest midline structure (6, 15-17). During amniote gastrulation, widespread mitosis has

been identified throughout the embryo, including the PS (18), and previous studies showed that

mitotic arrest leads to morphologically diminishing PS extension (19, 20). Further, the Wnt/PCP

pathway also plays an important role for proper PS extension and patterning (21). Since the chick

embryo is readily accessible and closely represents the human gastrula (22, 23), much of what is

known about the cell flow and PS development has come from studies using this model system

(14). The PS is first visible as a cell cluster at the posterior embryonic disc at the initial streak

stage [HH2; Hamburger and Hamilton staging (24)] formed through the influence of several axis

inducing factors [e.g. Vg1 and Chordin (25, 26)]; it then extends anteriorly along the midline

axis (6, 14, 15). During PS development, displacement of cells and/or tissue tension potentially

generates biomechanical forces, which lead to cell flows and/or PS morphogenesis (20, 21, 27,

28). Recent studies discuss that cell intercalation and ingression underlying PS formation,

leading to the polonaise movements through altering the neighbor tissue tension (21, 28, 29),

suggesting that the polonaise movements are likely a consequence of PS morphogenesis (16, 30,

31). Other studies demonstrate that the polonaise movements, generated by the myosin-cable-

mediated tissue tension with cell intercalation and ingression, lead to PS morphogenesis (20, 27,

31). Thus, the relationship between the polonaise movements and PS morphogenesis remains

controversial (31).

Here, we characterize previously undefined relationships between the polonaise movements and

PS morphogenesis. Suppression of the Wnt/PCP pathway generates a wider and shorter PS while

maintaining the polonaise movements. Mitotic arrest diminishes PS extension and maintains the

early phase of the polonaise movements. Moreover, experimentally manipulating the large-scale

cell flow by inducing a secondary midline axis indicates that the authentic PS forms despite

altered cell flows. Ultimately, a single axis-inducing morphogen, Vg1, is capable of initiating the

polonaise movements despite defective PS morphogenesis. These results suggest that PS

morphogenesis is not responsible for the initiation and early phase of the polonaise movements.

Further, the polonaise movements are not necessarily responsible for PS morphogenesis. This

study suggests that amniote gastrulation may be evolutionarily distinct from non-amniote

gastrulation in the mechanisms coupling large-scale cell flow and midline morphogenesis.

Results

Suppression of the Wnt/PCP pathway maintains the bilateral vortex-like counter-rotating

cell flow along a deformed PS

Throughout this work, the cell flows were recorded by tagging individual epiblast cells with

electroporated fluorescent tracers at the pre-streak stage (32). The tagged embryos were live-

imaged until HH3 with a modified New culture system, which we previously described (33, 34).

To reconstruct trajectories of the tagged cells and quantitatively analyze the cell flow pattern, the

recorded images were processed using tracking analysis tools (35), Particle Image Velocimetry

(PIV) (36-40), and subsequent visualization techniques (41, 42).

To address the relationship between the polonaise movements and PS morphogenesis, we

experimentally manipulated the shape of the PS while maintaining the population of PS cells

(Fig. 1A-B). The DEP domain of Dishevelled (Dsh; a transducer protein of Wnt signaling) is

responsible for the non-canonical Wnt/PCP pathway (43, 44), and misexpression of dominant-

negative Dsh lacking DEP [dnDsh(ΔDEP)] leads to deformation of the midline structures,

including the PS (21). Further, the Wnt/PCP pathway is involved in cellular polarity and

migration, while the canonical Wnt pathway regulates cell proliferation (45). We refer the

dnDsh(ΔDEP)-GFP construct that we generated, as ΔDEP-GFP, and tested its ability to alter

cellular polarity, resulting in PS deformation' (Fig. 1C-E, Fig. S1A-D). In the control-GFP-

introduced PS and epiblast cells, Pk1 (a PCP core component) exhibited a polarized localization

at the E-cadherin-based adherens junctions, particularly cell-cell junctions (Fig. S1A-B, n=3). In

contrast, this Pk1-localization pattern was not identified in the ΔDEP-GFP-misexpressing PS and

epiblast cells (Fig. S1C-D, n=3), implying that the construct efficiently disrupted the cellular

polarity within the misexpressing cells. The expression of *Brachyury* indicated that ΔDEP-GFP-

PS cells maintained their identity (Fig. 1E) but displayed a shorter and wider morphogenesis than

the control PS (Fig. 1C-E, n≥4 for each). The number of PS cells was not significantly different

from the control- and ΔDEP-GFP-PS (control $8.5 \pm 0.4 \times 10^3$ cells vs ΔDEP $8.8 \pm 1.2 \times 10^3$ cells,

p=0.68, n=3 for each, Fig. 1F). These results indicated that ΔDEP-misexpression was capable of

suppressing the Wnt/PCP pathway through disrupting the localization pattern of the PCP core

component while maintaining the number of the PS cells. Having confirmed the function of the

ΔDEP-GFP construct, we applied it as a molecular tool for manipulating the shape of the PS.

To address relationship(s) between the polonaise movements and the shape of the PS, we live-

imaged the control- and ΔDEP-GFP-embryos from pre-streak stage to HH3 and analyzed the

resulting cell flows (Fig. 1G-L''', and movie S1). In the control-GFP embryo, the cell flow

pattern, which was visualized by tracking tools and quantitatively analyzed by PIV with

subsequent flow-visualization techniques (41, 42), identified robust polonaise movements (Fig.

1G-G''', I-I'', K-K''', and movie S1, n=5). Surprisingly, these analyses indicated that

suppression of the Wnt/PCP pathway maintained the polonaise movements despite PS

deformation (n=5, Fig. 1C-L''' and movie S1). While the topology of the counter-rotating flow

pattern were maintained under suppression of the Wnt/PCP pathway (Fig. 1J-J'''), the distance

between the left-right rotating cell flows was wider in the ΔDEP embryos than the controls

(control 860.2±86.2μm vs ΔDEP 1008.4±183.8μm, p=0.026, n=4 for each, Fig. S1E-I), implying

that the shape of the PS and/or the Wnt/PCP signaling-mediated cell intercalation affected the

location of these rotations. These data demonstrated that the polonaise movements persisted even

though the PS as a midline structure was deformed under the condition of suppressing the

Wnt/PCP pathway.

Mitotic arrest diminishes PS formation but initially preserves the bilateral vortex-like

rotating cell flow

The above data showed that a deformed midline structure allowed the polonaise movements to

occur while the position of the LR rotations in the polonaise movements was affected (Fig. 1 and

Fig. S1). However, it remained unclear the relationship between the polonaise movements and

the population of the PS cells, particularly expansion of the PS precursor cells in PS

morphogenesis (Fig. 2A-B). To address this question, we applied a mitotic arrest approach using

aphidicolin (a specific DNA polymerase inhibitor), which has been shown to diminished PS

development (19, 20). In chick embryonic disc, active cell proliferation was identified in both the

epiblast and PS cells (Fig. 2C)(18). As verified by BrdU incorporation assay, the aphidicolin-

treated embryos exhibited little or no mitotic activity (Fig. 2D). Morphogenesis of the PS was

monitored by *Brachyury* (a marker for PS and mesodermal cells) and *Sox3* (an epiblast marker).

The expression domain and midline extension of cells, expressing *Brachyury* and negative for

Sox3, were considerably reduced in the aphidicolin-treated embryos, compared to the control

(Fig. 2E; n≥4 for each). The expression pattern of *Brachyury* and PS extension followed an

aphidicolin dose-dependent diminishment (Fig. S2A-D). The *Vg1*-expressing area was also

diminished but detectable in the aphidicolin-treated embryos (Fig. 2E, n=4), suggesting that the

axis-inducing morphogen was preserved in the embryonic disc, but expansion of the PS

precursor cells was suppressed by mitotic arrest. These results indicate that mitotic arrest,

induced by the aphidicolin-treatment, remarkably reduced PS extension and development,

consistent with the previous works (19, 20).

To address the relationship between the polonaise movements and the population of the PS cells,

we live-imaged the fluorescently tagged embryos under aphidicolin-treatment and analyzed the

resulting cell flow (Fig. 2F-K''' and movie S2). Tracking and PIV-based analyses visualized the

cell flow pattern in control and aphidicolin-treated embryos (Fig. 2F-K''' and movie S2). In

control embryos, the polonaise movements occurred from pre-streak until HH3 for more than

four hours (n=4, Fig. 2F-F''' and movie S2). The streamline and vorticity plots illustrated that the

polonaise movements became more robust in a time-dependent manner (Fig. 2H-H''' and J-J''').

Strikingly, in the aphidicolin-treated embryos (n=4), the polonaise movements initiated but were

maintained for a shorter period, 2.2±1.2 hours (control 6.5±1.2 hours, p=0.0038, Fig. G-G''', I-

I''', and movie S2). The streamline and vorticity plots also indicated the initiation and early

phase of the polonaise movements in the aphidicolin-treated embryos (Fig. 2J-J''' and K-K''').

Together, these data demonstrated that the typical pattern of the polonaise movements was

induced and initially maintained in the short-term, while there was no or little indication of

robust PS formation or extension.

Aphidicolin potentially induces apoptosis (46), which might indirectly lead to the cell flow

changes through tissue tension alterations. To test this possibility, we examined apoptosis in the

control and aphidicolin-treated embryos (Fig. S3A-B). TUNEL assay showed that the

aphidicolin-treatment slightly increased apoptotic rate but there was no significant difference

between these embryos and controls (control sham, 10.6±3.5%; aphidicolin, 12.1±3.7%, p=0.20,

n=4 for each, Fig. S3B), suggesting that the bilateral rotating cell flow in the aphidicolin-treated

embryos was not indirectly induced by apoptosis.

Previous studies have shown that a large-scale ring-like distribution of phospholyrated-myosin

(pMyosin) cables among the epiblast cell layer is required for tissue tension related to cell flow

(20, 27, 28). We examined whether the pMyosin cables were maintained in the treated embryos

over the course of the imaging (Fig. S3C-D"). In the aphidicolin-treated embryos, the large-scale

ring-like distribution pattern of the pMyosin cables remained in the epiblast (Fig. S3D-D"),

similar to the control embryos (Fig. S3C-C"), implying that the mitotic arrested-embryos

maintained tissue tension, consistent with the previous report (20). Thus, these results support a

model that mitosis is coupled with PS morphogenesis but is not necessarily required for the

initiation and early phase of the polonaise movements.

PS properly extends along the midline despite altering the large-scale cell flow

Given that PS morphogenesis is required for maintenance of the polonaise movements, we next

examined the converse of the relationship; that the polonaise movements may be required for PS

formation. To test this model, we experimentally reprogrammed the large-scale cell flow (Fig.

3A-B). Ectopically induced Vg1 leads to formation of a *Brachyury*-positive secondary PS and a

secondary midline axis (Fig. 3C-D)(47-49). COS cells expressing either a control-GFP (control-

GFP/COS) or Vg1 (Vg1/COS) construct were placed at the anterior marginal zone (AMZ) at pre-

streak stage, approximately 180° from the posterior marginal zone where the authentic PS forms,

and we analyzed the resulting cell flow during both authentic and ectopically induced PS

extension (Fig. 3B). If the bilateral rotating cell flow is required for PS formation, both the

authentic and induced PSs should each be accompanied by polonaise movements during the

extension. The trajectory analysis by Imaris and PIV-based analyses showed that the control-

COS implanted embryos exhibited stereotypical polonaise movements along the authentic

midline axis, indicating that the COS cell implantation maintained an authentic cell flow pattern

(Fig. 3E-E''', G-G''', I-I''', K, Fig. S4, and movie S3, n=4). In contrast, the analyses of the cell

flow in the Vg1/COS implanted embryos illustrated the disruption of the authentic polonaise

movements at the extending authentic PS (Fig. 3F-F''', H-H''', J-J''', Fig. S5, and movie S3,

n=4). At pre-streak stage, the epiblast cells initially converged to the center of the embryonic

disc along both the authentic and the induced axes, and the bilateral rotating cell flows began to

appear at both axes (Fig. 3F-F''', H-H''', Fig. S5, and movie S3). This cell flow pattern continued

for approximately one hour (Fig. 3H', Fig. S5, and movie S3). Before extension of the authentic

PS, the bilateral rotating cell flow along the induced axis became dominant, and eventually

overrode the original cell flow (Fig. 3H'-H''', Fig. S5, and movie S3). Despite global disruption

and reversed counter-rotating cell flows, the authentic PS extended properly along the authentic

midline (Fig. 3D, F''', H''', J''', K, Fig. S5, and movie S3; n=4). Secondary midline-axis

induction by ectopic Vg1 created the polonaise movements around the secondary PS and altered

the original cell flow pattern, generating a reprogrammed large-scale cell flow (Fig. 3F-F''', H-

H''', J-J''', K, Fig. S6, and movie S3). These results indicate that extension of the authentic PS is

preserved even under the defective polonaise movements.

Vg1 is capable of inducing the bilateral vortex-like rotating cell flow despite diminished PS

extension under mitotic arrest

The above data indicate the lack of a strong relationship between the polonaise movements and

PS morphogenesis; however, it remained unclear how the polonaise movements were induced.

Given that mitotic arrest maintained both Vg1-expression and the initiation of the polonaise

movements (Fig. 2 and movie S2) and that induction of the secondary midline axis by ectopic

Vg1 reprograms the epiblast cell flow (Fig. 3, Fig. S4, S5, and movie S3), we tested the

possibility that the axis-inducing morphogen, Vg1, would be capable of inducing the polonaise

movements in the absence of PS extension (Fig. 4A). We combined the secondary axis induction

by AMZ implantation of Vg1/COS cells with aphidicolin-treatment and analyzed the resulting

cell flow (Fig. 4B, Fig. S6, S7, and movie S4). The BrdU assay showed that the aphidicolin

treatment effectively blocked mitosis in Vg1/COS implanted embryos (control sham 88±9.0% vs

control-GFP/COS with aphidicolin 0.2±0.1% vs Vg1/COS with aphidicolin 0.19±0.2%, p=

0.0005×10^{-3}, n=3 for each, Fig. 4C-D) and diminished both the authentic and induced PSs (Fig.

4E-F, n=3 for each). The control-GFP/COS implanted embryo under mitotic arrest condition

initially displayed the polonaise movements along the authentic midline axis, however, the

vorticity decreased time-dependently, resulting in a defective rotating cell flow (n=3, Fig. 4G-

G''', I-I''', K-K''', Fig. S6, and movie S4), consistent with the non-COS cell implanted embryos

under aphidicolin-treatment (shown in Fig. 2). In contrast, the Vg1/COS implanted embryos

under mitotic arrest condition exhibited the polonaise movements predominantly at the induction

point of the secondary axis with disrupted authentic cell flow at the site of the authentic midline

axis (n=3, Fig. 4H-H''', J-J''', L-L''', and movie S4). These data suggest that the midline axis

induction by the single morphogen brings about the polonaise movements, even when the

midline structure (i.e. extending PS) is diminished. Taken together, our data show that Vg1 is

capable of inducing and reprogramming the large-scale cell flow during gastrulation even though

the midline structure is diminished.

Discussion

Here, we described a previously undefined relationship between the polonaise movements and

PS morphogenesis (Fig. 5). Our approaches to molecularly manipulating PS morphogenesis

revealed that the initiation and early phase of the polonaise movements occur despite diminished

PS formation (Fig. 5A). Conversely, the experimental manipulation of the cell flow

demonstrated that the authentic PS extends regardless of aberrant flow patterns (Fig. 5B).

Further, a single axis-inducing morphogen, Vg1, is capable of inducing the bilateral rotating cell

flow even under mitotic arrest, when the ectopic PS is reduced (Fig. 5B, C). Our data support a

model in which the PS population is required for maintaining the polonaise movements, as seen

in the ΔDEP embryos, while the polonaise movements are not necessarily responsible for PS

morphogenesis (Fig. 5D).

A similar mitotic arrest has been described as simultaneously diminishing PS extension and

disrupting the polonaise movements (19, 20), but it was not determined whether a lack of PS

extension and/or diminished cell population of the PS at the midline led to defective polonaise

movements. In this study, our live-imaging system combined with the quantitative analyses,

molecular tools for manipulating PS formation, and the gene expression analyses allowed us to

clarify the relationships among PS morphogenesis, the polonaise movements, mitosis, and the

Wnt/PCP pathway (Fig. 1, Fig. 2, and Fig. 5A, C). Our data demonstrated that the shape of the

PS as a midline structure is not necessarily responsible for initiation of the polonaise movements

while the population of the PS cells, expanded by mitosis, is required for sustained polonaise

movements (Fig. 1 and 2). Mesendoderm specification is required for large-scale cell flows

associated with later stage ingression and intercalation at the PS during gastrulation (27, 50).

Here, we have described the distinctive set of cell flows that precede gastrulation in the chick

embryo, and the relationship between early and late cell flow has yet to be examined.

The Wnt/PCP pathway has been assumed to be coupled with the polonaise movements, as an

evolutionarily conserved mechanism (21, 51). In contrast, another study showed that

misexpression of dnWnt11 or dnDsh-ΔPDZ [suppressing both the Wnt canonical and non-

canonical pathway, including the Wnt/PCP (43)] maintained the bilateral counter-rotating cell

flow along a defective PS (52). Thus, the role of the Wnt/PCP signaling pathway for the

polonaise movements has remained unresolved. In this study, our loss-of-function technique,

introducing ΔDEP to PS cells, demonstrated that the polonaise movements were preserved along

the deformed PS (Fig. 1, and movie S1). Our electroporation-mediated gene-transfection allowed

us to restrict gene transduction in area and number of cells (53). Despite the technical limitations,

the suppression level of the Wnt/PCP pathway through the ΔDEP-misexpression was sufficient

to lead to the PS deformation (Fig. 1 and Fig. S1). The Wnt/PCP pathway potentially works both

cell-autonomously and non-cell autonomously (45, 54). In the PS and non-PS epiblast cells, the

disruption of the Pk1-localization was identified not only in the ΔDEP-misexpressing cells, but

also in neighbors; however, this study is not designed to provide insight into questions of cell

autonomy. Further, the Wnt/PCP pathway is a major regulator of cell intercalation (51, 55).

Recent studies have shown that cell intercalation plays an important role in PS morphogenesis

and the initiation of the polonaise movements (20, 21, 27-29). Our data demonstrates that

suppression of the Wnt/PCP pathway preserves the initiation and maintenance of the polonaise

movements, but the lateral positioning of the rotations is enlarged when the PS is deformed (Fig.

1 and Fig. S1). These results suggest that the Wnt/PCP pathway may not be associated with

directional cell intercalation for driving the polonaise movements (Fig. 5C).

To investigate the role of the polonaise movements for PS morphogenesis, a technical

challenge/limitation is how to stop or disrupt the cell flow while preserving the cellular process

intact. For example, previous loss-of-function studies, using knockdown techniques (such as

siRNA and inhibitor treatments), identified the necessity of myosins, including pMyosin and

actomyosin, for both the large-scale cell flow and PS morphogenesis, through regulation of

tissue tension/biomechanical force (27, 28). Myosins are fundamental cellular components and

involved in several cell biological processes, such as tissue tension, cell division, and

intracellular transport (56). Therefore, it remains a possibility that myosin-related mechanisms

could be involved in the cell flow and PS morphogenesis. Recent studies using injury and

pressure models further demonstrated that the cell flow pattern can be altered by a physical cut to

an embryonic disc or removal of a large portion of an embryo disc through change of the tissue

tension or ablation (20, 57). While these models suggest that the cell flow pattern is dependent

on an intact embryonic architecture, it remains unclear as to involvement of other elements such

as signaling pathways and gene expression, which can be also molecularly and cell-biologically

changed as a result of surgeries. Thus, we developed an experimentally manipulation of the cell

flow system, using a Vg1-induced two-midline axis model, disturbing the stereotypical cell flow

pattern while maintaining embryonic development (Fig. 3 and Fig. 5B).

Our two-midline axis model provided an important insight into the axis-inducing morphogen for

induction and patterning of the polonaise movements (Fig. 3, Fig. 4, Fig. S4-S7, and movie S3,

S4). Our data demonstrated that the ectopic Vg1, mimicking the inducing activity of the posterior

marginal zone, generated the bilateral rotating cell flow aligned to the induced midline axis but

disrupted the cell flows at the authentic midline axis (Fig. 3 and Fig. 4). Despite the disturbed

cell flow, PS extension was preserved along both authentic and induced midline axes (Fig. 4).

While this model ultimately disturbed the stereotypical cell flow pattern at the authentic midline

axis, cell flow was initiated there. Therefore, our study does not rule out potential roles of the

cell flow in the initiating steps of PS extension.

It remains unclear how the induced polonaise movements along with the secondary axis

overcame the authentic cell flows despite mitotic arrest (Fig. 4 and Fig. 5). As for axis-induction

by other axis-inducing morphogens (58), Vg1 may work dose-dependently for induction and

patterning of the polonaise movements, and ectopic application may have over-ridden

endogenous levels (Fig. 3 and Fig. 4). Further studies will clarify the relationship between the

large-scale cell flows and the concentration of the morphogen; this will add new insight to the

regulation mechanisms that connect axis-induction and the cell flow patterning.

This study further suggests evolutional distinctions underlying cell biological mechanisms of the

large-scale cell flow and midline morphogenesis, between amniotes and non-amniotes. The first

distinction is the necessity of the Wnt/PCP pathway to the large-scale cell flow. Whereas the

Wnt/PCP pathway is coupled with both the large-scale cell flow and midline morphogenesis in

non-amniotes (55), our data revealed that the signaling pathway is required for proper PS

morphogenesis, but contributes little to the polonaise movements (Fig. 1). The second distinction

is mitosis-dependency/independency comparisons between model systems of gastrulation (Fig.

2). Non-amniotes, in particular amphibians, utilize a blastopore-mediated gastrulation and

undergo notochord formation with the large-scale cell flow, all independent of mitosis (10). In

amniotes, mitosis is required for PS morphogenesis, and expanding the mass of the PS cells

contributes to the maintenance of the polonaise movements (Fig. 2). Thus, the large-scale cell

flow and midline morphogenesis in amniote gastrulation is not tightly coupled, which may be

evolutionarily distinct from non-amniote gastrulation.

In summary, our study illustrates the relationships between midline morphogenesis, the large-

scale cell flow, and an axis-inducing factor in amniote gastrulation. These findings will add

another layer of our understanding to how the bilateral body plan is initiated and patterned along

the midline axis.

Acknowledgments

We thank past and present Mikawa lab and Prakash lab members, particularly Drs J. Hyer and L. Hua for their invaluable suggestions and/or assistance. Imaging data for this study were acquired at the Center for Advanced Light Microscopy-CVRI at UCSF on microscopes obtained using funding from the Research Evaluation and Allocation Committee, the Gross Fund, and the Heart Anonymous Fund. This work was supported in part by grants from NIH (R01HL122375, R37HL078921, R01HL132832, R01HL148125) to T.M.; Uehara Memorial Foundation Fellowship and JSPS Postdoctoral Fellowship for Research Abroad to R.A.; Univ Miami funds to N.P.V.; NSF CCC (DBI-1548297), CZI BioHub Investigator Program, and Howard Hughes Medical Institute to M.P. **Author contributions:** T.M., R.A., V.N.P., and M.P. conceptualized the project. T.M. and R.A. wrote the manuscript. R.A. performed all experiments. R.A. and V.N.P. carried out the PIV analysis of cell flows. S.S. technically supported PIV analysis. V.N.P., S.S., and M.P. participated in project discussions and commented on the manuscript.

Competing interests: Competing interests No competing interests declared.

Correspondence and requests for materials should be addressed to takashi.mikawa@ucsf.edu.

Fig. 1. 'Polonaise movements' persist under suppression of Wnt/PCP pathway.
(A) Testing model of relationships between the polonaise movements, primitive streak (PS)
morphogenesis, and the Wnt/PCP pathway. A-P and L-R; anterior-posterior and left-right body
axes, respectively. (B) Diagram of experimental set up. (C, D) In situ hybridizations for
Brachyury (Bra) in control- and ΔDEP-GFP-misexpressing embryos. (E) Overlay of embryos
hybridized for Bra (n=4 for each). (F) Cell number in PS [control $8.5\pm0.4\times10^3$ cells vs ΔDEP
$8.8\pm1.2\times10^3$ cells, p=0.68, n=3 for each (two-tailed Student's t test)]. (G-H''') Trace of cell flow
path of electroplated epiblast cells with Flowtrace. The initiation of the polonaise movements
was set as Time 0 (t=0). See also Movie S1. (I-J''') Streamlines, visualizing averaged cell flows
during each time period. (K-L''') Vorticity plots, displaying an averaged measure of the local
rotation during each time period. Blue, clockwise; red, counter-clockwise rotation.

371
Fig. 2. Early phase of 'polonaise movements' remains under global mitotic arrest.
(A) Testing model of relationships between the polonaise movements, primitive streak (PS)
morphogenesis, and mitosis. Axes as in Fig. 1. (B) Diagram of experimental set up.
(C, D) BrdU incorporation in control-sham operated and aphidicolin treated in embryos. White
boxes show enlarged areas. (C', C") High magnification of boxed areas in PS and non-PS,
respectively. (E) In situ hybridizations for *Brachyury* (*Bra*), *Vg1*, and *Sox3* in control and
aphidicolin-treated embryos. Arrow heads indicate diminished PSs. Black boxes, enlarged area
including the diminished PS. (F-G''') Trace of cell flow path with Flowtrace analysis (green is
trace, yellow indicates endpoint) of fluorescently-tagged epiblast cells. The initiation of the
polonaise movements was set as Time 0 (t=0). See also Movie S2. (H-I''') Streamlines,
visualizing averaged cell flows during each time period. Interpolated vectors are displayed in
orange. (J-K''') Vorticity plots, displaying averaged measure of the local rotation during each
time period. Blue, clockwise; red, counter-clockwise rotation. **Note**: Since vorticity is calculated
for all deviation from set point, slight curves and full rotations receive the same color indication,
as seen in K''' which is not maintaining a bilateral vortex-like-rotating cell flows. See Materials
and Methods section for a full discussion.

Fig. 3. Authentic PS extends under disruption of authentic 'polonaise movements'.
(A) Testing model of relationships between the polonaise movements, primitive streak (PS)
morphogenesis, using ectopic Vg1-inducing system. Axes as in Fig. 1. (B) Diagram of
experimental set up. (C, D) In situ hybridizations for Brachyury (Bra) in control and ectopic
Vg1-induced PS, respectively. (E-F''') Trace of cell flow path of electroplated epiblast cells by
using Imaris tracking analysis. See also Movie S3. White circles, COS cell implanted site. The
initiation of the polonaise movements was set as Time 0 (t=0). (G-H''') Streamlines, visualizing
averaged cell flows during each time period of the vector fields. (I-J''') Vorticity plots,
displaying an averaged measure of the local rotation during each time period in the vector fields.
Blue, clockwise; red, counter-clockwise rotation. **Note**: vorticity is identified in both a full-
rotation and curve, particularly in (J-J'''). The white asterisks indicate the polonaise movements
at the induced PS which are opposite direction to the authentic PS. (K) Summary of the cell flow
patterns in control-GFP/COS- and Vg1/COS-implanted embryos, shown in (E-J''').

Fig. 4. Induced 'polonaise movements' persist despite defective PS morphogenesis.

(A) Testing model of relationships between the polonaise movements, primitive streak (PS)

morphogenesis, and Vg1. Axes as in Fig. 1. (B) Diagram of experimental set up.

(C, D) BrdU incorporation in control-GFP/COS- and Vg1/COS-implanted embryos under

aphidicolin-treatment. White boxes showed enlarged areas. (E, F) In situ hybridizations for

Brachyury (Bra, n=3 for each). Black arrows indicate diminished PSs. (G-H''') Trace of cell

flow path of electroplated epiblast cells by using Imaris tracking analysis. See also Movie S5.

White circles, COS cell implanted site. The initiation of the polonaise movements was set as

Time 0 (t=0). (I-J''') Streamlines, visualizing averaged cell flows during each time period of the

vector fields. (K-L''') Vorticity plots, displaying an averaged measure of the local rotation

during each time period. Note: vorticity is identified in both a full-rotation and curve, particularly

in (L-L'''). Blue, clockwise; red, counter-clockwise rotation. White asterisks, the polonaise

movements at the induced axis which are opposite direction to the authentic midline axis. **Note:**

vorticity is identified in both a full-rotation and curve in (K''') and (L'''). See description in the

material and method section.

418
Fig. 5. Summary of the patterns of the cell flow and PS morphogenesis.
(A) Cell flow pattern under experimental manipulation of PS morphogenesis.
(B) Authentic PS morphogenesis under disrupting authentic polonaise movements.
(C) Vg1-induced cell flow pattern under defective PS morphogenesis.
(D) Summary of the relationships between polonaise movements, PS morphogenesis, mitosis, and Wnt/PCP pathway.

References

1.   S. F. Gilbert, M. J. F. Barresi, *Developmental Biology* (Oxford University Press, 2017).
2.   L. Solnica-Krezel, D. S. Sepich, Gastrulation: making and shaping germ layers. *Annu Rev*
Cell Dev Biol **28**, 687-717 (2012).
3.   M. Leptin, Gastrulation movements: the logic and the nuts and bolts. *Dev Cell* **8**, 305-320
(2005).
4.   L. Solnica-Krezel, Conserved patterns of cell movements during vertebrate gastrulation.
Curr Biol **15**, R213-228 (2005).
5.   R. Keller, L. Davidson, Cell movements of gastrulation. *Gastrulation: from cells to*
embryos (ed. C. Stern), 291-304 (2004).
6.   T. Mikawa, A. M. Poh, K. A. Kelly, Y. Ishii, D. E. Reese, Induction and patterning of the
primitive streak, an organizing center of gastrulation in the amniote. *Dev Dyn* **229**, 422-
432 (2004).
7.   R. Keller, Shaping the Vertebrate Body Plan by Polarized Embryonic Cell Movements.
Science **298**, 1950-1954 (2002).
8.   E. W. Gehrels, B. Chakrabortty, M. E. Perrin, M. Merkel, T. Lecuit, Curvature gradient
drives polarized tissue flow in the Drosophila embryo. *Proc Natl Acad Sci U S A* **120**,
e2214205120 (2023).
9.   C. Bertet, L. Sulak, T. Lecuit, Myosin-dependent junction remodelling controls planar
cell intercalation and axis elongation. *Nature* **429**, 667-671 (2004).
10. R. Keller *et al.*, Mechanisms of convergence and extension by cell intercalation. *Philos*
Trans R Soc Lond B Biol Sci **355**, 897-922 (2000).
11. M. Tada, M. L. Concha, C.-P. Heisenberg (2002) Non-canonical Wnt signalling and
regulation of gastrulation movements. in *Seminars in cell & developmental biology*
(Elsevier), pp 251-260.
12. R. Wetzel, Untersuchungen am Hühnchen. Die Entwicklung des Keims während der
ersten beiden Bruttage. *Wilhelm Roux Arch Entwickl Mech Org* **119**, 188-321 (1929).
13. L. Gräper, Die Primitiventwicklung des Hühnchens nach stereokinematographischen
Untersuchungen, kontrolliert durch vitale Farbmarkierung und verglichen mit der
Entwicklung anderer Wirbeltiere. *Wilhelm Roux Arch Entwickl Mech Org* **116**, 382-429
(1929).
14. C. D. Stern, *Gastrulation in the chick*. C. D. Stern, Ed., Gastrulation: from cells to
embryo (CSHL Press, New York, 2004), pp. 219-232.
15. G. C. Schoenwolf, Molecular Genetic Control of Axis Patterning during Early
Embryogenesis of Vertebrates. *Annals of the New York Academy of Sciences* **919**, 246-
260 (2000).
16. A. Raffaelli, C. D. Stern, Signaling events regulating embryonic polarity and formation
of the primitive streak in the chick embryo. *Curr Top Dev Biol* **136**, 85-111 (2020).
17. G. Sheng, A. Martinez Arias, A. Sutherland, The primitive streak and cellular principles
of building an amniote body through gastrulation. *Science* **374**, abg1727 (2021).
18. E. J. Sanders, M. Varedi, A. S. French, Cell proliferation in the gastrulating chick
embryo: a study using BrdU incorporation and PCNA localization. *Development* **118**,
389-399 (1993).
19. C. Cui, X. Yang, M. Chuai, J. A. Glazier, C. J. Weijer, Analysis of tissue flow patterns
during primitive streak formation in the chick embryo. *Dev Biol* **284**, 37-47 (2005).

20.   M. Saadaoui, D. Rocancourt, J. Roussel, F. Corson, J. Gros, A tensile ring drives tissue
flows to shape the gastrulating amniote embryo. *Science* **367**, 453-458 (2020).

21.   O. Voiculescu, F. Bertocchini, L. Wolpert, R. E. Keller, C. D. Stern, The amniote
primitive streak is defined by epithelial cell intercalation before gastrulation. *Nature* **449**,
1049-1052 (2007).

22.   R. O'Rahilly, F. Muller, Developmental stages in human embryos: revised and new
measurements. *Cells Tissues Organs* **192**, 73-84 (2010).

23.   R. Asai, M. Bressan, T. Mikawa, Avians as a Model System of Vascular Development.
Methods Mol Biol **2206**, 103-127 (2021).

24.   V. Hamburger, H. L. Hamilton, A series of normal stages in the development of the chick
embryo. 1951. *Dev Dyn* **195**, 231-272 (1992).

25.   E. A. Seleiro, D. J. Connolly, J. Cooke, Early developmental expression and experimental
axis determination by the chicken Vg1 gene. *Curr Biol* **6**, 1476-1486 (1996).

26.   A. Streit *et al.*, Chordin regulates primitive streak development and the stability of
induced neural cells, but is not sufficient for neural induction in the chick embryo.
Development **125**, 507-519 (1998).

27.   M. Chuai, G. Serrano Nájera, M. Serra, L. Mahadevan, C. J. Weijer, Reconstruction of
distinct vertebrate gastrulation modes via modulation of key cell behaviors in the chick
embryo. *Sci Adv* **9**, eabn5429 (2023).

28.   E. Rozbicki *et al.*, Myosin-II-mediated cell shape changes and cell intercalation
contribute to primitive streak formation. *Nat Cell Biol* **17**, 397-408 (2015).

29.   O. Voiculescu, L. Bodenstein, I. J. Lau, C. D. Stern, Local cell interactions and self-
amplifying individual cell ingression drive amniote gastrulation. *Elife* **3**, e01817 (2014).

30.   O. Voiculescu, Movements of chick gastrulation. *Curr Top Dev Biol* **136**, 409-428
(2020).

31.   J. X. Wang, M. D. White, Mechanical forces in avian embryo development. *Semin Cell
Dev Biol* **120**, 133-146 (2021).

32.   H. Eyal-Giladi, S. Kochav, From cleavage to primitive streak formation: a
complementary normal table and a new look at the first stages of the development of the
chick. I. General morphology. *Dev Biol* **49**, 321-337 (1976).

33.   D. A. T. New, A New Technique for the Cultivation of the Chick Embryo in vitro.
Journal of Embryology and Experimental Morphology **3**, 326-331 (1955).

34.   L. Maya-Ramos, T. Mikawa, Programmed cell death along the midline axis patterns
ipsilaterality in gastrulation. *Science* **367**, 197-200 (2020).

35.   W. Gilpin, V. N. Prakash, M. Prakash, Flowtrace: simple visualization of coherent
structures in biological fluid flows. *J Exp Biol* **220**, 3411-3418 (2017).

36.   W. Thielicke, E. J. Stamhuis, PIVlab – Towards User-friendly, Affordable and Accurate
Digital Particle Image Velocimetry in MATLAB. *Journal of Open Research Software* **2**
(2014).

37.   E. J. Stamhuis, Basics and principles of particle image velocimetry (PIV) for mapping
biogenic and biologically relevant flows. *Aquatic Ecology* **40**, 463-479 (2006).

38.   M. Raffel, C. E. Willert, J. Kompenhans, *Particle image velocimetry: a practical guide*
(Springer, 1998), vol. 2.

39.   C. K. Batchelor, G. K. Batchelor, *An introduction to fluid dynamics 2nd edition*
(Cambridge university press, 2012).

40. R. Aris, *Vectors, tensors and the basic equations of fluid mechanics* (Courier
Corporation, 2012).
41. M. Jiang, R. Machiraju, D. Thompson, Detection and visualization of Vortices. *The
visualization handbook* **295** (2005).
42. H. G. W. Christopher J. Greenshields, *Notes on Computational Fluid Dynamics: General
Principles* (CFD Direct Limited, 2022).
43. M. Sharma, I. Castro-Piedras, G. E. Simmons, Jr., K. Pruitt, Dishevelled: A masterful
conductor of complex Wnt signals. *Cell Signal* **47**, 52-64 (2018).
44. U. Rothbächer *et al.*, Dishevelled phosphorylation, subcellular localization and
multimerization regulate its role in early embryogenesis. *Embo j* **19**, 1010-1022 (2000).
45. Y. Yang, M. Mlodzik, Wnt-Frizzled/Planar Cell Polarity Signaling: Cellular Orientation
by Facing the Wind (Wnt). *Annual Review of Cell and Developmental Biology* **31**, 623-
646 (2015).
46. J. M. Glynn, T. G. Cotter, D. R. Green (1992) Apoptosis induced by actinomycin D,
camptothecin or aphidicolin can occur in all phases of the cell cycle. (Portland Press
Ltd.).
47. J. Cooke, Morphogenesis and regulation in spite of continued mitotic inhibition in
Xenopus embryos. *Nature* **242**, 55-57 (1973).
48. Y. Wei, T. Mikawa, Formation of the avian primitive streak from spatially restricted
blastoderm: evidence for polarized cell division in the elongating streak. *Development*
127, 87-96 (2000).
49. S. B. Shah *et al.*, Misexpression of chick Vg1 in the marginal zone induces primitive
streak formation. *Development* **124**, 5127-5138 (1997).
50. G. S. Nájera, C. J. Weijer, The evolution of gastrulation morphologies. *Development* **150**
(2023).
51. I. Roszko, A. Sawada, L. Solnica-Krezel, Regulation of convergence and extension
movements during vertebrate gastrulation by the Wnt/PCP pathway. *Semin Cell Dev Biol*
20, 986-997 (2009).
52. M. Chuai *et al.*, Cell movement during chick primitive streak formation. *Dev Biol* **296**,
137-149 (2006).
53. Y. Ishii, T. Mikawa, Somatic transgenesis in the avian model system. *Birth Defects Res C
Embryo Today* **75**, 19-27 (2005).
54. C. F. Davey, C. B. Moens, Planar cell polarity in moving cells: think globally, act locally.
Development **144**, 187-200 (2017).
55. M. Tada, C.-P. Heisenberg, Convergent extension: using collective cell migration and
cell intercalation to shape embryos. *Development* **139**, 3897-3904 (2012).
56. M. A. Hartman, J. A. Spudich, The myosin superfamily at a glance. *Journal of Cell
Science* **125**, 1627-1632 (2012).
57. P. Caldarelli, A. Chamolly, O. Alegria-Prévot, J. Gros, F. Corson, Self-organized tissue
mechanics underlie embryonic regulation. *bioRxiv* 10.1101/2021.10.08.463661,
2021.2010.2008.463661 (2021).
58. J. Green, Morphogen gradients, positional information, and Xenopus: Interplay of theory
and experiment. *Developmental Dynamics* **225**, 392-408 (2002).

Material and methos

Embryo isolation and culture conditions

Fertilized eggs of White Leghorn (*Gallus Gallus domesticus*) were obtained from Petaluma Farms (Petaluma, CA) and were incubated at 37°C in a humidified incubator to the appropriate embryonic stages. We electroporated unincubated eggs at pre-streak stages IX to XII for cell flow imaging. Embryos were isolated in Tyrode's solution (137mM NaCl, 2.7 mM KCL, 1mM $MgCl_2$, 1.8mM $CaCl_2$, 0.2 mM Ha_2HPO_4, 5.5mM D-glucose, pH 7.4). After manipulations (such as electroporation, inhibitor treatment, and implantation), the embryos were live-imaged on a vitelline membrane stretched around a glass ring according to the New culture method as previously described (33, 34).

Implantation of COS cells

$2x10^5$ of COS7 cells were plated in a 35mm cell culture dish (Thermo Fisher Scientific, 153066), 24 hours prior to transfection. The cells were transfected by Lipofectamine 3000 (Thermo Fisher Scientific, L3000015) with 6µg of pMT23-Vg1-myc-GDF1 plasmid DNA (a gift from Drs. Claudio D. Stern and Jane Dodd) for 5 hours. The next day, the transfected cells were trypsinized, and cultured using the hanging-drop method. Each hanging drop contained $10x10^6$ cells in 20 µl of culture media and were cultured overnight (O/N). The cell spheres were then rinsed in serum-free DMEM and implanted into the anterior marginal zone in the chick embryonic discs at pre-streak stages.

Plasmid generation

ΔDEP-GFP was cloned from a plasmid, XE124 XDsh delta DEP-GFP-CS2+ in pCS2+ (a gift from Dr. Randall Moon, Addgene, #16785), by PCR (Q5 High-Fidelity DNA Polymerase, New England Biolabs, M0491) with the following primers: 5'-TACCGCGGGCCCGGGATCCAGCCACCATGGCGGAGACT-3' and 5'- AGCCTGCACCTGAGGAGTGCTTATTTGTATAGTTCATCCATGCCATGTGTAATCC'. After PCR purification with QIAquick Gel Extraction Kit (Qiagen, #28704), the PCR product was inserted into a backbone vector, pCAG-GFP (a gift from Connie Cepko, Addgene, #1115) by using Gibson-assembly (New England Biolabs, E2611).

Electroporation

Embryos were transfected with expression vectors, and/or control-oligo DNA conjugated with

Carboxyfluorescein or Lissamine (GENE TOOLS) using an electroporator (Nepagene) with 3 pulses of

2.4-3.8 Volts, 50 milliseconds duration, 500 milliseconds interval, and platinum electrodes. The DNA

solution delivered to the epiblast contained 0.1% fast green (final 0.02%), 80% glucose (final 4%) and

5μg/μl of expression vectors, or 1mM control-oligo (5'-CCTCTTACCTCAGTTACAATTTATA-3').

Aphidicolin treatment

Aphidicolin (SIGMA, A0781), dissolved in DMSO (SIGMA, B23151), was added to Tyrode's solution to

a final concentration of 0.1 to 100μM. Embryos were isolated, and soaked in Tyrode's solution with

either 0.3% DMSO (control), or various concentrations of aphidicolin for 15 minutes at 37°C. Embryos

were then cultured using the New culture method at 37°C.

BrdU assay

Embryos were soaked in Tyrode's solution containing BrdU (final concentration; 0.1mM, Thermo Fisher,

B23151) with or without aphidicolin (SIGMA, A0781) at 37°C for 15 minutes, and cultured for 12 hours

at 37°C (BrdU was incorporated for 12 hours). Embryos were then fixed in 4% paraformaldehyde (PFA,

Electron Microscopy Sciences)/PBS for 30 minutes at room temperature (RT). Embryos were then

washed with PBS to remove PFA, and unincorporated BrdU, and incubated with 1M HCl for 1 hour at

RT to denature DNA. BrdU signal was detected by immunofluorescence staining with anti-BrdU

antibody (1□200, Millipore, MAB3424).

Tunel assay

The In Situ Cell Death Detection Kit, TMR red (Rosche, 12156792910) was used, and the kit instructions

followed with these additional modifications. Embryos were fixed in 4% PFA for 30 min at RT and

washed in PBS 3X for 10 minutes each. Permeabilization was performed in TBST (0.5% Triton-X) for 30

minutes at RT. Embryos were then incubated with the TUNEL reaction and DAPI staining for 3 hours at

37°C and washed with PBS five times for 10 minutes each.

Immunofluorescent staining

For chicken embryos:

Embryos were fixed in 4% PFA/PBS for 30 minutes at RT, or 4:1 Methanol/DMSO at 4°C O/N and

washed in PBS 3X for 30 minutes each. Embryos were then incubated in blocking reagent [1% bovine

serum albumin (BSA) and 0.1% triton in PBS] for 1 hour at RT. Embryos were then incubated with

primary antibodies at 4°C O/N. After washing in PBS 3 X for 30 minutes each, embryos were then

incubated with secondary antibodies for 2 hours at RT. Embryos were then washed in PBS 2X for 30

minutes each. For the BrdU assay, embryos had an additional incubation with Streptavidin, Alexa Fluor

647 conjugate (1:1000, Thermo Fisher Scientific) for 1 hour at RT, and washed in PBS 2X for 30 min

each. Embryos were incubated with DAPI for 1 hour, washed for 30 minutes, then mounted between

cover slips (VWR, 48366249), and slide glasses (Thermo Fisher Scientific, 12-544-7) with Aqua-

PolyMount (Polysciences, Inc., #18606-20).

Primary antibodies:

Anti-BrdU (1□200, Millipore, MAB3424), anti-E-Cadherin (1:500, BD Biosciences, 610181), anti-GFP

(1:1000, Rockland, 600-101-215), , anti-Prickle1 antibody (1:300, Proteintech, 22589-1-AP), and anti-

GFP-HRP antibody (1:200, Abcam, 600-101-215), anti-ZO1 antibody (1:300, Thermo Fisher Scientific,

33-9100).

Secondary antibodies:

Anti-mouse-IgG (H+L) Alexa Fluor 647 antibody (1:500, Thermo Fisher Scientific, A-31571), anti-goat

IgG (H+L) Alexa Fluor 488 (1:500, Thermo Fisher Scientific, A11055), anti-rabbit IgG (H+L) Alexa

Fluor 568 antibodies (1:500, Thermo Fisher Scientific, A10042), anti-goat IgG Alexa Fluor 488 (1:1000,

Thermo Fisher Scientific, A11055) and anti-mouse IgG Alexa Fluor 594 antibodies (1:1000, Thermo

Fisher Scientific, A21203).

Whole Mount In Situ Hybridization (WISH) and subsequent immunohistochemistry

Embryos were fixed in 4% PFA/PBS in PBS O/N at 4°C. WISH was performed as previously described

(39), using plasmids for *Brachyury* and *Sox3* (gift from Dr. Raymond B. Runyan), and *Vg1*. All WISH

were carried out with paralleled control embryos for managing the color development time. For

subsequent IHC, embryos were fixed in 4:1 methanol/DMSO at 4°C O/N.

The next day, embryos were incubated with 4:1:1 methanol/DMSO/30% Hydrogen peroxide at RT and

incubated for 2 hours. Then embryos were washed in Tris-Buffered Saline (pH7.6) with 0.1% Tween 20

(TBST) 3X for 10 minutes each, and incubated with blocking solution (1% sheep serum in TBST) for 1

hour at RT. Primary antibodies, anti-GFP antibody-HRP conjugated (1:200, Abcam, ab6663), were

applied for at 4°C O/N incubation. Embryos were then washed in TBST 5X for 1 hour each at RT,

followed by 10 minutes with 0.05M Tris-HCl (pH 7.6), and incubated for 30 minutes with DAB (3,3′-

Diaminobenzidine)/Tris-HCl (pH7.6) 0.0003% Hydrogen peroxide in the dark. After the reaction was

stopped, the embryos were washed in PBS, and placed in 4% PFA at 4°C for long term storage.

Imaging of chick embryos

For live-imaging, embryos were cultured in a 35 mm dish by the New culture method at 37°C. Time-lapse

images were recorded with Nikon TIRF/Spinning Disk microscope (Nikon Ti inverted fluorescent

Microscope with CSU-22 spinning disc confocal) supported by Prime 95B Scientific CMOS camera

(Photometrix), Nikon Widefield Epifluorescence inverted microscope (Nikon Ti inverted fluorescent

Microscope with CSU-W1 large field of view) supported by ANDOR iXon camera and Nikon Eclipse

TE2000-E supported by Hamamatsu ORCA-Flash 2.8 camera. The acquisition time was every 3 minutes

using Nikon Elements Advance Research software V4.00.07. All time-lapse images were recorded every

3 minutes.

Immunostained chick embryos were imaged by a confocal microscope (Leica TCS SPE). Fixed WISH

samples were imaged by Leica MZ16F microscope with Leica DFC300 Fx camera and Leica FireCam

V.3.4.1 software.

Quantification and Statistical Analysis

Cell number, mitotic rate, BrdU-, and Tunel-assay:

To measure frequency of BrdU-positive or Tunel-positive cells, 3D-surface was created to an

immunostained image of the embryonic disc by using the Surface tool in Imaris X64 9.2.0 software, and

then the center of the embryonic disc was set for the following measurement. 3x3 squares with 800μm of

each side length were then arranged on the embryonic disc. Numbers of DAPI and BrdU-positive nuclei

within each square were counted by Spots tool and averaged to get a frequency of BrdU- or Tunel-

positive nuclei (Fig. 2F, Fig. S1A, and Fig. S2C).

Cell flow analyses:

Cell flow analysis was performed with Imaris X64 9.2.0 software, Flowtrace and Particle Image

Velocimetry (PIV) analysis. Frowtrace stacks every 20-40 frames from original live-imaging data and

generates projection images of particle trajectories for visualizing cell movements. PIV analysis is

described in the following part.

PIV analysis and cell flow visualizing techniques:

For quantitatively analyzing the cell flow pattern, the recorded images were further processed by using

Particle Image Velocimetry (PIV) and subsequent visualization techniques. PIV analysis technique

[PIVlab package in MATLAB (36)], which is a common technique for mapping displacement of particles

(e.g. tagged cells) over a short time interval measured from a series of consecutive images to obtain the

velocity vector field is employed to quantify the cell flows in the experimental time-lapse datasets. PIV is

a standard technique used in the Fluid Physics and Engineering community to quantitatively measure and

characterize fluid flows in different systems. This analysis provides comprehensive and quantitative

information of a whole flow as a velocity vector field. The PIV analysis involves several steps that are

carried out on image sequences over time intervals of 3 minutes, which is also the imaging acquisition

speed in the experiments. First, we apply a high-pass filter during the image preprocessing step to

highlight the bright particles (fluorescently tagged cell) and minimize the background. Then the PIV

analysis was carried out by partitioning the image into smaller interrogation windows, typically of size 64

x 64 pixels with 50% overlap on the first pass, and a window size of 32 x 32 pixels with 50% overlap on

the second pass. After this, raw velocity vector fields over time were obtained, and then these vectors

were post-processed and validated by setting threshold limits, data smoothing and interpolation. A

calibration was applied to convert pixel/frame units into μm/s units.

The generated velocity vector field through PIV displayed the averaged cell pattern of the whole-flow-

field in the certain time period in this study. To highlight the more specific rotating cell flows,

mathematical/physical information (such as topology and vorticity) was extracted from these velocity

vector fields via computational visualization techniques (36, 39, 40, 42). Topology of the cell flow was

determined by streamlines, which are a group of curves that approximate to contours of the vectors in the

velocity vector field at a particular moment in time (39, 40, 42). Flowtrace visualized the trajectories of

the cell flow, whereas the streamlines illustrated the averaged flow pattern in the time period. Therefore,

the trajectories and streamlines show similar pattern when the cell flow is in stable-state; however, they

are not completely consistent with each other when the cell flow is unstable and time-dependently

changing (39, 40, 42). Vorticity, also termed as 'curl of vorticity', is a measure of a local rotation in the

larger flow and its magnitude is twice that of the local angular velocity (39, 40, 42). The vorticity plot

further displays directionality of the rotation. Since vorticity arises from any rotational movements,

including both a full- and partial rotation, the vorticity plot displayed the vorticities in both the vortex-like

rotating and curves of the cell flow (36, 39, 40, 42).

Measurement of the distance between the left-right rotations of the polonaise movements in control-

GFP and ΔDEP-embryos

The vector fields in control- and ΔDEP-GFP-embryos were generated from the recorded movies by using

PIVlab and averaged the flow pattern of the polonaise movements for 6 hours. The centers of the left-

right rotating cell flow of the polonaise movements in the averaged vector field were identified by

streamlines and measured the distance between them by using ImageJ.

Statistics:

Statistical analyses were performed using two-tailed Student's t tests, one-way ANOVA in R (Ver3.5.0)

and Excel 2022 (Microsoft). Statistical tests, individual p values, and sample numbers are described in

Figure panels and legends.

References for Material and methods:
732
733

1.   S. F. Gilbert, M. J. F. Barresi, *Developmental Biology* (Oxford University Press, 2017).

2.   L. Solnica-Krezel, D. S. Sepich, Gastrulation: making and shaping germ layers. *Annu Rev*
Cell Dev Biol **28**, 687-717 (2012).

3.   M. Leptin, Gastrulation movements: the logic and the nuts and bolts. *Dev Cell* **8**, 305-320
(2005).

4.   L. Solnica-Krezel, Conserved patterns of cell movements during vertebrate gastrulation.
Curr Biol **15**, R213-228 (2005).

5.   R. Keller, L. Davidson, Cell movements of gastrulation. *Gastrulation: from cells to*
embryos (ed. C. Stern), 291-304 (2004).

6.   T. Mikawa, A. M. Poh, K. A. Kelly, Y. Ishii, D. E. Reese, Induction and patterning of the
primitive streak, an organizing center of gastrulation in the amniote. *Dev Dyn* **229**, 422-
432 (2004).

7.   R. Keller, Shaping the Vertebrate Body Plan by Polarized Embryonic Cell Movements.
Science **298**, 1950-1954 (2002).

8.   E. W. Gehrels, B. Chakrabortty, M. E. Perrin, M. Merkel, T. Lecuit, Curvature gradient
drives polarized tissue flow in the Drosophila embryo. *Proc Natl Acad Sci U S A* **120**,
e2214205120 (2023).

9.   C. Bertet, L. Sulak, T. Lecuit, Myosin-dependent junction remodelling controls planar
cell intercalation and axis elongation. *Nature* **429**, 667-671 (2004).

10.   R. Keller *et al.*, Mechanisms of convergence and extension by cell intercalation. *Philos*
Trans R Soc Lond B Biol Sci **355**, 897-922 (2000).

11.   M. Tada, M. L. Concha, C.-P. Heisenberg (2002) Non-canonical Wnt signalling and
regulation of gastrulation movements. in *Seminars in cell & developmental biology*
(Elsevier), pp 251-260.

12.   R. Wetzel, Untersuchungen am Hühnchen. Die Entwicklung des Keims während der
ersten beiden Bruttage. *Wilhelm Roux Arch Entwickl Mech Org* **119**, 188-321 (1929).

13.   L. Gräper, Die Primitiventwicklung des Hühnchens nach stereokinematographischen
Untersuchungen, kontrolliert durch vitale Farbmarkierung und verglichen mit der
Entwicklung anderer Wirbeltiere. *Wilhelm Roux Arch Entwickl Mech Org* **116**, 382-429
(1929).

14.   C. D. Stern, *Gastrulation in the chick*. C. D. Stern, Ed., Gastrulation: from cells to
embryo (CSHL Press, New York, 2004), pp. 219-232.

15.   G. C. Schoenwolf, Molecular Genetic Control of Axis Patterning during Early
Embryogenesis of Vertebrates. *Annals of the New York Academy of Sciences* **919**, 246-
260 (2000).

16.   A. Raffaelli, C. D. Stern, Signaling events regulating embryonic polarity and formation
of the primitive streak in the chick embryo. *Curr Top Dev Biol* **136**, 85-111 (2020).

17.   G. Sheng, A. Martinez Arias, A. Sutherland, The primitive streak and cellular principles
of building an amniote body through gastrulation. *Science* **374**, abg1727 (2021).

18.   E. J. Sanders, M. Varedi, A. S. French, Cell proliferation in the gastrulating chick
embryo: a study using BrdU incorporation and PCNA localization. *Development* **118**,
389-399 (1993).

19.   C. Cui, X. Yang, M. Chuai, J. A. Glazier, C. J. Weijer, Analysis of tissue flow patterns
during primitive streak formation in the chick embryo. *Dev Biol* **284**, 37-47 (2005).

20.   M. Saadaoui, D. Rocancourt, J. Roussel, F. Corson, J. Gros, A tensile ring drives tissue
flows to shape the gastrulating amniote embryo. *Science* **367**, 453-458 (2020).

21.   O. Voiculescu, F. Bertocchini, L. Wolpert, R. E. Keller, C. D. Stern, The amniote
primitive streak is defined by epithelial cell intercalation before gastrulation. *Nature* **449**,
1049-1052 (2007).
22.   R. O'Rahilly, F. Muller, Developmental stages in human embryos: revised and new
measurements. *Cells Tissues Organs* **192**, 73-84 (2010).
23.   R. Asai, M. Bressan, T. Mikawa, Avians as a Model System of Vascular Development.
Methods Mol Biol **2206**, 103-127 (2021).
24.   V. Hamburger, H. L. Hamilton, A series of normal stages in the development of the chick
embryo. 1951. *Dev Dyn* **195**, 231-272 (1992).
25.   E. A. Seleiro, D. J. Connolly, J. Cooke, Early developmental expression and experimental
axis determination by the chicken Vg1 gene. *Curr Biol* **6**, 1476-1486 (1996).
26.   A. Streit *et al.*, Chordin regulates primitive streak development and the stability of
induced neural cells, but is not sufficient for neural induction in the chick embryo.
Development **125**, 507-519 (1998).
27.   M. Chuai, G. Serrano Nájera, M. Serra, L. Mahadevan, C. J. Weijer, Reconstruction of
distinct vertebrate gastrulation modes via modulation of key cell behaviors in the chick
embryo. *Sci Adv* **9**, eabn5429 (2023).
28.   E. Rozbicki *et al.*, Myosin-II-mediated cell shape changes and cell intercalation
contribute to primitive streak formation. *Nat Cell Biol* **17**, 397-408 (2015).
29.   O. Voiculescu, L. Bodenstein, I. J. Lau, C. D. Stern, Local cell interactions and self-
amplifying individual cell ingression drive amniote gastrulation. *Elife* **3**, e01817 (2014).
30.   O. Voiculescu, Movements of chick gastrulation. *Curr Top Dev Biol* **136**, 409-428
(2020).
31.   J. X. Wang, M. D. White, Mechanical forces in avian embryo development. *Semin Cell
Dev Biol* **120**, 133-146 (2021).
32.   H. Eyal-Giladi, S. Kochav, From cleavage to primitive streak formation: a
complementary normal table and a new look at the first stages of the development of the
chick. I. General morphology. *Dev Biol* **49**, 321-337 (1976).
33.   D. A. T. New, A New Technique for the Cultivation of the Chick Embryo in vitro.
Journal of Embryology and Experimental Morphology **3**, 326-331 (1955).
34.   L. Maya-Ramos, T. Mikawa, Programmed cell death along the midline axis patterns
ipsilaterality in gastrulation. *Science* **367**, 197-200 (2020).
35.   W. Gilpin, V. N. Prakash, M. Prakash, Flowtrace: simple visualization of coherent
structures in biological fluid flows. *J Exp Biol* **220**, 3411-3418 (2017).
36.   W. Thielicke, E. J. Stamhuis, PIVlab – Towards User-friendly, Affordable and Accurate
Digital Particle Image Velocimetry in MATLAB. *Journal of Open Research Software* **2**
(2014).
37.   E. J. Stamhuis, Basics and principles of particle image velocimetry (PIV) for mapping
biogenic and biologically relevant flows. *Aquatic Ecology* **40**, 463-479 (2006).
38.   M. Raffel, C. E. Willert, J. Kompenhans, *Particle image velocimetry: a practical guide*
(Springer, 1998), vol. 2.
39.   C. K. Batchelor, G. K. Batchelor, *An introduction to fluid dynamics 2nd edition*
(Cambridge university press, 2012).
40.   R. Aris, *Vectors, tensors and the basic equations of fluid mechanics* (Courier
Corporation, 2012).

41.   M. Jiang, R. Machiraju, D. Thompson, Detection and visualization of Vortices. *The*
visualization handbook **295** (2005).
42.   H. G. W. Christopher J. Greenshields, *Notes on Computational Fluid Dynamics: General*
Principles (CFD Direct Limited, 2022).
43.   M. Sharma, I. Castro-Piedras, G. E. Simmons, Jr., K. Pruitt, Dishevelled: A masterful
conductor of complex Wnt signals. *Cell Signal* **47**, 52-64 (2018).
44.   U. Rothbächer *et al.*, Dishevelled phosphorylation, subcellular localization and
multimerization regulate its role in early embryogenesis. *Embo j* **19**, 1010-1022 (2000).
45.   Y. Yang, M. Mlodzik, Wnt-Frizzled/Planar Cell Polarity Signaling: Cellular Orientation
by Facing the Wind (Wnt). *Annual Review of Cell and Developmental Biology* **31**, 623-
646 (2015).
46.   J. M. Glynn, T. G. Cotter, D. R. Green (1992) Apoptosis induced by actinomycin D,
camptothecin or aphidicolin can occur in all phases of the cell cycle. (Portland Press
Ltd.).
47.   J. Cooke, Morphogenesis and regulation in spite of continued mitotic inhibition in
Xenopus embryos. *Nature* **242**, 55-57 (1973).
48.   Y. Wei, T. Mikawa, Formation of the avian primitive streak from spatially restricted
blastoderm: evidence for polarized cell division in the elongating streak. *Development*
127, 87-96 (2000).
49.   S. B. Shah *et al.*, Misexpression of chick Vg1 in the marginal zone induces primitive
streak formation. *Development* **124**, 5127-5138 (1997).
50.   G. S. Nájera, C. J. Weijer, The evolution of gastrulation morphologies. *Development* **150**
(2023).
51.   I. Roszko, A. Sawada, L. Solnica-Krezel, Regulation of convergence and extension
movements during vertebrate gastrulation by the Wnt/PCP pathway. *Semin Cell Dev Biol*
20, 986-997 (2009).
52.   M. Chuai *et al.*, Cell movement during chick primitive streak formation. *Dev Biol* **296**,
137-149 (2006).
53.   Y. Ishii, T. Mikawa, Somatic transgenesis in the avian model system. *Birth Defects Res C*
Embryo Today **75**, 19-27 (2005).
54.   C. F. Davey, C. B. Moens, Planar cell polarity in moving cells: think globally, act locally.
Development **144**, 187-200 (2017).
55.   M. Tada, C.-P. Heisenberg, Convergent extension: using collective cell migration and
cell intercalation to shape embryos. *Development* **139**, 3897-3904 (2012).
56.   M. A. Hartman, J. A. Spudich, The myosin superfamily at a glance. *Journal of Cell*
Science **125**, 1627-1632 (2012).
57.   P. Caldarelli, A. Chamolly, O. Alegria-Prévot, J. Gros, F. Corson, Self-organized tissue
mechanics underlie embryonic regulation. *bioRxiv* 10.1101/2021.10.08.463661,
2021.2010.2008.463661 (2021).
58.   J. Green, Morphogen gradients, positional information, and Xenopus: Interplay of theory
and experiment. *Developmental Dynamics* **225**, 392-408 (2002).
866
867

Fig. S1. Distance between the LR rotations increases under suppression of the Wnt/PCP
pathway.
(A, C) Immunofluorescent staining images of control- and ΔDEP-GFP-misexpressing PSs at
HH3 with GFP (green) and DAPI (cyan). (B, D) Immunostaining images of epiblast and PS
cells in control- and ΔDEP-GFP-misexpressing embryos at HH3 with GFP (green), Prickle1
(Pk1; magenta), and DAPI (blue). The white asterisks indicate GFP-expressing cells. (E, G)
Averaged bilateral vortex-like-rotating cell flows (6 hours), visualized by vorticity plot, in
control- and ΔDEP-GFP-misexpressing embryos. Black lines, measurement of the distance
between the left-right (LR) rotations. Blue, clockwise; red, counter-clockwise rotation. (F, H)
Averaged bilateral vortex-like-rotating cell flows (6 hours), visualized by streamline plot, in
control- and ΔDEP-GFP-misexpressing embryos. (I) Distance between the LR rotations in
control- and ΔDEP-GFP-misexpressing embryos [control-GFP 860.2±86.2 μm vs ΔDEP-GFP
1008.4±183.8μm, p=0.026, n=4 for each (two-tailed Student's t test)].

Fig. S2. PS extension dose-dependently responds to aphidicolin.
(A) Dose-dependency of aphidicolin-induced mitotic arrest (PS, control 98.8±1.3% vs 10uM aphidicolin 10.3±4.3% vs 100uM aphidicolin 0±0%, p=$0.002e^{-4}$ in one-way ANOVA; embryo, control 87.9±9.0% vs 10uM aphidicolin 9.0±4.4% vs 100uM aphidicolin 0.02±0.01%; n=4 for each; p=$0.003e^{-5}$ in one-way ANOVA). (B-C) In situ hybridization for Brachyury (Bra) to different concentrations of aphidicolin treatment. (D) Length of PS at HH3 for different concentrations of aphidicolin treatment (control 1.3±0.2mm vs 0.1μM aphidicolin 1.3±0.3mm vs 1μM aphidicolin 0.8±0.4mm vs 10μM aphidicolin 0.3±0.1mm vs 100μM aphidicolin 0.1±0.1mm; n=6 for each, p=$6.2e^{-9}$ in one way ANOVA).

894
Fig. S3. Aphidicolin-treatment maintains apoptotic index and pMyosin cables.
(A) TUNEL staining at stage HH 3 in control and aphidicolin-treated embryos. (B) Apoptotic
index in control and aphidicolin-treated embryos (control 10.6±3.5% vs aphidicolin 12.1±3.7%,
p=0.20, n=4 for each). (C-D") Immunofluorescent staining images of control- and aphidicolin-
treated embryos at HH3 with ZO-1 (magenta) and pMyosin (cyan). White boxes show enlarged
areas.
901

902
Fig. S4. Time evolution of cell flow in control-GFP/COS-implanted embryo.
(A) Trace of cell flow path of electroplated epiblast cells by using Imaris tracking analysis. See
also Movie S3. A-P and L-R; anterior-posterior and left-right body axes, respectively. White
circles, COS cell implanted site. (B) Streamlines, visualizing averaged cell flows during each
time period. (C) Vorticity plots, displaying averaged measure of the local rotation during each
time period. Blue, clockwise; red, counter-clockwise rotation.
909

Fig. S5. Time evolution of cell flow in Vg1/COS-implanted embryo.
(A) Trace of cell flow path of electroplated epiblast cells by using Imaris tracking analysis. See
also Movie S3. Axes are shown in Fig S4. White circles, COS cell implanted site. (B)
Streamlines, visualizing averaged cell flows during each time period. (C) Vorticity plots,
displaying averaged measure of the local rotation during each time period. Blue, clockwise; red,
counter-clockwise rotation. White asterisks, the polonaise movements at the induced axis which
are opposite direction to the authentic axis.
918

919
Fig. S6. Time evolution of cell flow in control-GFP/COS-implanted embryo under
aphidicolin-treatment.
(A) Trace of cell flow path of electroplated epiblast cells by using Imaris tracking analysis. See
also Movie S5. Axes are shown in Fig S4. White circles, COS cell implanted site. (B)
Streamlines, visualizing averaged cell flows during each time period. (C) Vorticity plots,
displaying averaged measure of the local rotation during each time period. Blue, clockwise; red,
counter-clockwise rotation.
927

928
Fig. S7. Time evolution of cell flow in Vg1/COS-implanted embryo under aphidicolin-
treatment.
(A) Trace of cell flow path of electroplated epiblast cells by using Imaris tracking analysis. See
also Movie S5. Axes are shown in Fig S4. White circles, COS cell implanted site. (B)
Streamlines, visualizing averaged cell flows during each time period. (C) Vorticity plots,
displaying averaged measure of the local rotation during each time period. Blue, clockwise; red,
counter-clockwise rotation. White asterisks, the polonaise movements at the induced axis which
are opposite direction to the authentic axis.
937

Supplemental movies

Movie S1. 'Polonaise movements' continue under suppression of Wnt/PCP pathway. Time-lapse movie was analyzed by using Imaris tracking. Embryos were co-electroporated with lissamine-tagged oligo (red) and control-GFP or ΔDEP-GFP expression vectors (cyan) at pre-streak stage X-XII. Live imaging performed at 4x objective lens on a spinning disk confocal microscope until HH3. Related to Fig. 1 and Fig. S1. Scale bars, 300 μm.

Movie S2. Mitotic arrest maintains early phase of 'polonaise movements'. Time-lapse movie of fluorescently-tagged epiblast cells in control and aphidicolin-treated embryos from pre-streak to HH3, processed by using Flowtrace. The trajectories, from green to yellow, indicate movement of fluorescently tagged cells (30 min projections). Embryos were electroporated with fluorescein-tagged oligo (green) and treated with DMSO (control-sham operated) or 100 μM of aphidicolin. Live-imaging performed at 4x objective lends on a spinning disk confocal microscope for 10 hours. Related to Fig. 2, Fig. S2, and S3. Scale bars, 200 μm.

Movie S3. Authentic PS extends in absence of regular 'polonaise movements'. Embryos were electroporated with lissamine-tagged oligo (red) and implanted with COS cells expressing either a control-GFP (control-GFP/COS) or Vg1 (Vg1/COS) expression vector at anterior marginal zone at pre-streak stage IX-X. Live-imaging performed at a 2x objective lens on a wide-field epifluorescent microscope. Magenta dots; lissamine-tagged epiblast cells, cyan lines; trajectories of the tagged epiblast cells for 2 hours. Related to Fig. 3, Fig. S4, S5. Scale bars, 300 μm.

Movie S4. Ectopic Vg1 leads to 'polonaise movements' despite defective PS morphogenesis after mitotic arrest. Embryos were electroporated with lissamine-tagged oligo (red), implanted with either control-GFP/COS or Vg1/COS at the anterior marginal zone (AMZ), and treated with 100 μM of aphidicolin at pre-streak stage IX to X. Live imaging performed at a 2x objective lens with a wide-field epifluorescent microscope. Magenta dots; lissamine tagged epiblast cells, cyan lines; trajectories for 2.5 hours. Related to Fig. 4, Fig. S6, and S7. Scale bars, 300 μm.