FHY3/FAR1 **transposable elements generate adaptive genetic variation in the *Bassia scoparia* genome**

Nathan Hall[1], Jinyi Chen [1,2], Christopher Saski[3], Phil Westra[4], Todd Gaines[4], Eric Patterson [1]

[1] Michigan State University, Department of Plant, Soil, and Microbial Sciences
[2] Nanjing Agricultural University, College of Plant Protection
[3] Clemson University, Department of Plant and Environmental Sciences
[4] Colorado State University, Department of Agricultural Biology

Abstract

Transposable element activity has greatly impacted the evolution of *Kochia scoparia* syn. *Bassia scoparia* (kochia), an important agricultural weed. In this study, an improved kochia genome assembly of the glyphosate-susceptible line 7710 was analyzed in comparison to a glyphosate-resistant line, M32. A near complete genome assembly consisting of 14 pseudochromosomes with a total length of 969.6 Mb, and N50 scaffold length of 99.88 Mb, was generated with hybrid short-read (Illumina), and single-molecule (PacBio/Bionano genomics) data, in addition to chromatin interaction mapping (Hi-C). The 9 largest scaffolds corresponded to the 9 chromosomes of a close relative, *Beta vulgaris*. From this assembly, 54,387 protein-coding gene loci were annotated. After gene annotation it was revealed that 58% of the whole-genome consisted of transposable elements. We identified that gene families containing *FHY3* (FAR-Red Elongated Hypocotyl 3) and *FAR1* (FAR-Red Impaired Response 1) functional domains have undergone a massive *Bassia*-lineage specific expansion. *FHY3* and *FAR1* domains were first described as a "domesticated" (non-active) sub-type of Mutator Don-Robertson transposase (MuDR) that function as transcription factors; however, the domains may still function as transposases in the correct genomic context, but this has yet to be observed. Mutator elements are associated with large DNA segmental transposition within the genome including gene duplication events, which can greatly influence plant phenotypes and evolution. We discovered in our work that putative MuDR elements with detectable *FHY3*/*FAR1* domains were tightly associated with segmental duplications of 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*) that confer resistance to the herbicide glyphosate. Further, in a case study of MuDR activity, we characterized a new MuDR subtype, named here as "Muntjac", which contributes to the evolution of herbicide resistance in kochia through the process of transduplication. Collectively, our study provides insights into the de-domestication of a *FHY3*/*FAR1* transposon and provides new perspectives on the evolution of glyphosate resistance in kochia.

Keywords: Transposable Elements, *MuDR*, *FHY3*/*FAR1*, Glyphosate Resistance, Kochia, reference genome, genome resequencing, 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*), copy number variation

1. Introduction

Transposable elements (TEs) are engines of genetic change that can fundamentally alter an organisms genome and genetic diversity, even within relatively short evolutionary timescales[1]. TEs are active and mobile in the genome and propagate through replication cascades that are independent from the duplication of the background genome. While primarily thought of as detrimental as they can disrupt proper gene function and expression, there may be benefits from novel genetics that these TEs generate at the population and species level. The two major classes of TEs are defined by their propagation mechanism, known as Class I and II[2]. Class I TEs replicate via an RNA intermediate, a so-called "copy and paste" mechanism, while Class II TEs are without RNA intermediates and mobilize DNA through excision and re-insertion, via a "cut and paste" mechanism. When TEs insert into a new location in the genome, they may impart local structural changes to the DNA, such as methylation and gene expression[3,4]. At a chromosomal scale, TE insertions can facilitate larger-scale genomic rearrangements such as deletions, inversions, and segmental duplications[5,6]. At the gene scale, TEs can cause loss of gene function or gene shuffling through the process of transduplication[7–9]. Finally, and most rarely, TEs may contribute novel genes to the genome through a process of TE domestication, as in the case of the genes *FHY3* (Far-red elongated hypocotyl 3) and *FAR1* (far-red-impaired response). These

domesticated TEs were originally derived from a Class II TE known as a Mule (Mutator like) transposon in the MuDR (Mutator Don Robertson) TE clade but they have been hypothesized to have lost their ability to transpose and neofunctionalized as transcription factors that coordinate response to light signals[10].

All functioning, autonomous MuDR elements have a shared set of characteristics: 1) a DDE motif in the transposase domain, 2) terminal inverted repeats (TIR), and 3) target site duplications (TSD) that are created when the MuDR is inserted in a novel location[10]. When Mule elements move, including MuDR, their excision and reinsertion is known to pick up a 'payload' of genomic material composed of the genes found in the MuDR flanking sequence[11]. The transposase domain of MuDRs have been well characterized and is mechanistic in the complete excision of both the forward and reverse DNA strands from the genome, frequently leaving an excision footprint[12]. MuDR TIRs aid in the formation of their circularized double stranded DNA intermediate during transposition. TSDs form at the site of DNA insertion at a new location, which are typically short (8-11 bp) directed repeats adjacent to the TIRs[12]. *FHY3* and *FAR1* are notable in that as part of their domestication process they have lost their TIRs, resulting in the prediction that *FHY3* and *FAR1* cannot form the DNA intermediate needed for translocation of a Class II TE[13]. An undomesticated TE, named *Jittery,* is sister to the *FHY3* and *FAR1* genes and retains its TIRs and mobility[14].

Bassia scoparia (L.) Schrad. (kochia) is a common and economically important weed in crop and ruderal areas across the Canadian Prairies and the Great Plains of the USA[15]. Kochia is diploid with a somatic chromosome number of 2n=18 and an estimated haploid genome size of 0.7-1 Gb[16,17]. Glyphosate resistance in kochia is due to a tandemly arrayed copy number variation (CNV) of glyphosate's target, 5-enolpyruvylshikimate-3-phosphate synthase (*EPSPS*)[18,19]. In previous work, a mobile genetic element (MGE) that contained a gene annotated as having a *FHY3* and a *FAR1* protein domain was identified flanking the *EPSPS* CNV suggested that the origin of the EPSPS CNV was closely related with *FHY3* and *FAR1* mobility[17]. In this research, we investigate a massive expansion of the *FHY3/ FAR1*-like gene family in the kochia lineage and provide evidence that some or even most of these *FHY3/ FAR1* like genes are not domesticated but are still actively mobile, at least in the kochia genome. Furthermore, we specifically describe the *FHY3/FAR1* MGE involved in glyphosate resistance and characterize it in the broader context of the MuDR transposon clade. We rename this specific transposing *FHY3/FAR1* gene "Muntjac", for the eponymous ship imagined by Lev Grossman in The Magicians Trilogy in order to differentiate it from domesticated *FHY3/FAR1* genes[20,21]. We propose that the transposable element, like the fabled ship, is highly mobile, at least in the kochia genome and that it has exhibiting degrees of contraction and expansion even at recent timescales. Furthermore, we propose that the activity of Muntjac is a signature of the evolutionary trajectory of the kochia genome in response to human selection from the herbicide glyphosate.

2. Material and methods

2.1. Germplasm

Several collections of glyphosate susceptible and resistant kochia plants were used for these experiments. The herbicide-susceptible reference genome assembly was generated from the highly inbred lab strain known as '7710', originally collected from western Nebraska[22]. This line has been used extensively as a susceptible control for many herbicide studies. The 7710 line has a consistent phenotype and it is well-studied compared to many kochia collections[23,24]. The primary resistant line used for comparison was the inbred lab line known as 'M32', collected from north-eastern Colorado[25], which has been confirmed to have highly amplified copy numbers of *EPSPS* and consistent (non-segregating) inheritance of the glyphosate resistant phenotype[17,26]. Furthermore, the *EPSPS* locus in M32 has been extensively studied and assembled using BAC sequencing[17].

To discover the original locus of Muntjac we collected two glyphosate-susceptible populations, one located from Israel in the native range of kochia (referred to as Israel), and a second from northern Colorado (referred to as CO-S). Additionally, we used glyphosate resistant plants with high *EPSPS* copy number but no amplification of the Muntjac gene (referred to as Oregon)[27]. For all studies, seeds were sowed in SureMix® and kept in growth chamber (25 °C in the day and 20 °C at night, 14/10 h light period) and watered every other day.

2.2. Kochia genome sequencing and assembly

DNA was extracted from leaf tissue of a single glyphosate susceptible kochia plant from the line 7710 using a modified CTAB protocol[17,28,29]. DNA was sent for PAC-BIO Sequel sequencing at The University of Delaware DNA Sequencing & Genotyping Center. Expanding leaf tissue was sent to Phase genomics for HiC library prep and sequencing. An optical map was constructed at the McDonnell Genome Institute using the BioNano Genomics Saphyr system. Short, high-coverage, paired end reads were generated at the University of Illinois on an Illumina HiSeq. These Illumina reads were used to assess the genome size of the kochia glyphosate-susceptible line 7710. Paired-end Illumina reads (144.34 Gb) (with an insert size of 350 bp) were selected to generate 17-mer frequency data for k-mer analysis. The k-mer depth distribution was plotted against the k-mer frequency, with the highest peak occurring at a depth of 97. Based on the total number of k-mers (71,593,662,168 bp), the kochia genome size was calculated to be 969.61 Mb, with genome heterozygosity of 0.06%. Additional flow cytometry data confirmed this size estimate with individuals from 900-1050 Mb. Like most chenopods, kochia has a chromosome count of $n = 9$[30].

Long read sequences were initially assembled using MeCat2[31]. Short reads were then aligned to this assembly using HiSat2 v2.1.0[32]. The assembly and alignment were used in conjunction to polish the genome using Pilon[33]. Phased haplotigs were merged using HaploMerger2[34]. Merged contigs were scaffolded using HiC data and the SALSA pipeline[35]. After initial scaffolding, the assembly was further scaffolded using the optical map at the McDonnell Genome Institute and the BioNano Solve software[36]. This assembly was then gap filled using PBJelly[37]. Newly filled gaps were polished once again using Pilon. Finally, super-scaffolds were manually scaffolded into pseudo-chromosomes using the HiC data and Juicer[38].

Gene models in the assembly were predicted using the GenSAS pipeline (https://www.gensas.org/gensas)[39]. The genome assembly was uploaded to the GENSAS webserver along with RNA-seq data for annotation[23]. The RepeatMasker and RepeatModeler tools were used to identify the repeats, and mask the genome for gene model calling[40,41]. Publicly available RNAseq data (NCBI SRA: SRX3276189- SRX3276192) were aligned with the masked genome, using BLAST+, BLAT, HISAT2, TopHat, and PASA [42–45]. The genome and these alignments were run through the Augustus, BRAKER, GeneMark-ES, Genscan, and SNAP tools for gene model prediction with default parameters[46–49]. Next, we made consensus gene models using EvidenceModeler. Each predictive tool was uniquely weighted. The following final weights were assigned to each prediction: 10 for transcript alignments and BRAKER; 5 for Augustus and protein alignments; 1 for GeneMark-ES and SNAP. PASA refinement was conducted using BRAKER-hisat2 as OGS. For the functional annotation, predicted genes were aligned to the NCBI plant protein database using both BLASTp and Diamond. Targeting and subcellular localization was predicted using Pfam, SignalP and TargetP, with default parameters for plants. Additional functional annotation was performed with Multiloc2, Iprscan, and an MMseqs2 search of Uniref50 database[50–52]. The final annotation was solidified after running BUSCO using the embryophyta_odb10 lineage database on the predicted proteins to assess the completeness of the annotation[53] .

2.3. Gene Family Expansions and the FHY3/ FAR1-like gene family in kochia

To determine the history of Muntjac and if *FHY3/FAR1* domains were prevalent in gene expansions, orthogroup clustering was performed using Orthofinder v2.4.0[54]. All annotated genes in the kochia genome were subjected to orthologous gene clustering with all annotated genes from eight closely related species from the Caryophyllales clade (Chenopodiaceae and Amaranthaceae). The species tree and related gene trees were inferred by Orthofinder using representative proteins produced using CGAT v0.06[54,55]. A CAFÉ analysis was conducted to detect significant expansions within orthogroups[56]. Distances within species tree produced by Orthofinder were scaled by a factor of 100 with ETE3 and CAFÉ was run with default settings[57].

The *FHY3/FAR1*-like susceptible protein described by Patterson et al[17] was used to validate the annotation of Muntjac. The gene sequence was identified by protein blast to the assembly and reads from three different SRA runs (ERR364385, SRR6165247, and SRR1198330) were mapped to the gene with HISAT2 to update the annotation. These results were compared to HISAT2 junction files, and matching junctions were searched for in 31 SRAs. The manually produced transcript was searched for ORFs using NCBI ORF Finder (https://www.ncbi.nlm.nih.gov/orffinder/ last accessed 3/8/2021), searched for conserved domains using NCBI Conserved Domain Search

(https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi last accessed 3/8/2021) and using InterPro
online search portal (https://www.ebi.ac.uk/interpro/, last accessed 3/8/2021).
Past research has demonstrated that kochia genomes may exhibit appreciable structural variability.
To classify types of gene duplication and detect perturbations to genome structure, MCSCANX was
run according to its documentation. To confirm structural differences between the glyphosate
susceptible line 7710 and glyphosate resistant line M32, micro-assemblies were created for the genomic
regions surrounding an intact Muntjac (Bs.00g352450) and a fragmented Muntjac insertion
(Bs.00g145150). Resistant reads were mapped against the kochia genome with HiSat2, and mapped
reads from target genes and the surrounding sequence were extracted with SAMTools v1.10[58]. Extracted
reads were assembled with SPAdes v3.14.0 (default settings)[59]. Assembled regions from resistant lines
were compared back to each locus of origin to determine if an excision had occurred and search for
target site duplications and terminal inverted repeats (TIRs).
To determine the relationship of Muntjac to *FHY3*, *FAR1* and *Jittery*, a gene tree was produced
using protein sequences that represented several TE families including Jittery, Hop, and MuDRs.
Briefly, full protein records were found on NCBI by searching reported transposase domains against
NCBI using BLAST online search utility with searches including eight NCBI taxon IDs. Amino acid
sequences were aligned and curated. Curated alignment was analyzed using IQTree v1.6.12 for
Windows with flags (-alrt 1000, -bb 1000). For the purposes of visualization, the reduced set of proteins
was searched for Pfam domains using NCBI Conserved Domain Search
(https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi, last accessed 21/8/2021). Detected Pfam
regions were compared with a down sampled gene tree produced with ETE3 v3.1.2 and Python v6.5.3
(https://github.com/NDHall/kochia_scoparia_analysis). Figures prepared with Inkscape v0.92.4 (
https://inkscape.org/, last accessed 3/8/2021).
189
2.4. Assessing copy number differences and expression differences.
To determine genome wide copy number variation (CNV) in the previously characterized
glyphosate resistant line M32, an individual was selected for genomic Illumina resequencing[17]. Paired
end data were aligned to the susceptible reference genome assembly (i.e. 7710) using HiSat2 on
standard parameters. Additionally, Illumina paired end data from the susceptible plant was also aligned
to the reference to control for shared CNV events between the resistant and susceptible plants using
HiSat2 on standard parameters. The resulting alignments were scanned using CNVnator v0.4.1 and a
1,000 bp sliding window to detect read coverage changes outside of the normal read depth distribution[60].
Duplicated and deleted regions were identified using a read depth cutoff of 2x and 0.5x coverage for
duplications and deletions (respectively) and an adjusted p-value < 0.05. CNV regions shared between
the susceptible and resistant lines were removed from consideration as they most likely exist in the
background of the kochia genome and were not subject to glyphosate selection. The genes in the
resulting glyphosate-resistant CNVs were collected and the gene ontology terms associated with those
genes were analyzed for GO enrichment.
To measure the relationship of predicted gene copy number and expression, publicly available
RNAseq data (4-7710 and 4-M32 individuals) was aligned to the genome-predicted transcriptome using
HiSat2[17,32]. Counts were extracted from alignments using samtools idxstats and differential expression
was calculated using recommend settings in EdgeR[58,61]. Relative expression was correlated to relative
gene copy number differences between the S and R lines.
209
2.5. Muntjac and its MGE activity and transposition
PCR primers were designed to identify 1) the upstream and downstream boundaries of the MGE
containing Muntjac's insertion on chromosome 9 (chr9), 2) the excision of the MGE from chr9 in a
glyphosate resistant individual, 3) the insertion of the MGE at the *EPSPS* locus on chr1, and the 4) the
presence of the flanking genomic sequence at the chr9 position. For this assay, we used several
individuals from the five kochia collections described in section 2.1: 7710 (S), CO-S (S), Israel (S),
M32 (R), and Oregon (R). DNA was extracted from individual plants of all populations. There were
two replicates per population. Primers were designed based on kochia chr9 assembled sequences and
were blasted to the whole kochia genome to confirm their specificity. The five pairs of primers were
used for validation (Table 3). The PCR was conducted in a 25-μL volume, consisting of 1 μL of DNA,
1 μL of each primer (10 μM) and 12.5 μL of GoTaq Green Master mix (Promega®, USA). The PCR

was run in a thermocycler (BioRad, USA) with the following profile: 95 ℃ for 5 min, 35 cycles of 95
℃ for 30 s, 55 ℃ for 30 s, and 72 ℃ for 1 min, followed by a final extension step of 5 min at 72 ℃.
223
3. Results
3.1. The Kochia Genome
The genome was assembled using a combination of high coverage RSII PacBio single-molecule
real-time (SMRT) sequencing (130 Gb of raw data), deep-coverage Illumina shotgun sequencing data
(144 Gb of raw data), a Hi-C linkage dataset (87 Gb of raw data), and a BioNano Optical map. The
assembly consisted of 14 scaffolds with an N50 scaffold length of 99.88 Mb and an N50 contig length
of 0.91 Mb. Total sequenced length of the assembly was 837.60 Mb, with 132.0 Mb in gaps. The
scaffolds were ordered and numbered as pseudochromosomes against the *Beta vulgaris* genome,
kochia's phylogenetically nearest sequenced relative. Due to *Beta vulgaris* and kochia being
significantly phylogenetically divergent, several inversions, duplications and deletions are apparent in
the whole genome alignment. The 5 remaining, much smaller, scaffolds from kochia remain
unincorporated in the chromosome assembly due to a lack of information placing them in the larger
pseudomolecules. For the most part, they are not included in further analyses. The average length of the
9 pseudomolecules is 100.6 Mb, among which the shortest was 84.85 Mb (chr2), and the longest was
117.96 Mb (chr5) (Figure 1).
Structural and functional gene annotation revealed that a majority (95.0%) of the 1,614 BUSCOs
in the embryophyta_odb10 BUSCO dataset were either complete or fragmented, which indicated a high
degree of assembly and annotation completeness (Table 1). In total, 54,387 gene models were predicted,
45,372 received functional annotation and 8,333 unique Interpro annotations were assigned to 30,046
genes and 2,234 unique Gene Ontologies were assigned to 19,357 genes.
244
3.2. Gene Family Expansions and the FHY3/ FAR1-like gene family in kochia
Orthologous gene clustering using CAFÉ and all annotated genes from eight closely related species
from the Caryophyllales clade (Chenopodiaceae and Amaranthaceae) revealed 206 of gene families that
are significantly expanded with 0 families contracting in kochia relative to all other chenopods (Figure
2, Table 2). Of the expanded gene families, 10 gene families contain proteins annotated as containing
FHY3/FAR1 family and/or FAR1 DNA binding domain (IPR031052 and IPR004330 respectively)
accounting for 267 genes total. Furthermore, using the results from Orthofinder and MCSCANX we
see an especially high enrichment of *FHY3* and *FAR1* related genes annotated with IPR004330,
IPR031052, or IPR037766 interpro identifiers within segmental duplications. A total 80 of the 282
whole genome/segmental duplications contained at least one *FAR1* related gene that had undergone
duplication. *FAR1* related genes and their segmental duplications seem to be concentrated in gene rich
areas of the chromosomes rather than the centromere (Figure 1).
257
3.3. Copy number variation and expression differences
Previous work identified a CNV event that increases the copy number of the gene *EPSPS*, the target
of the herbicide glyphosate, which in turn increases mRNA and protein production. The increased
EPSPS protein production due to the *EPSPS* CNV is not inhibited by typical glyphosate application
rates and therefore confers high levels of glyphosate resistance[62–64]. This CNV is widespread in kochia
populations throughout North America[27,65]. Several different arrangements of the *EPSPS* CNV have
been suggested[27]; however, the *EPSPS* CNV event being described here is confirmed 1) to be in a
tandem arrangement (i.e., not dispersed) with two distinct repeat units of different sizes, 2) to include 7
genes (one of which was *EPSPS*), and 3) to be associated with the insertion of one or more 15 kb mobile
genetic element(s) (MGE) containing the complete *Muntjac* gene. This specific arrangement was
previously confirmed in the resistant line M32 by BAC sequencing[17]. This MGE is flanked by TIRs
suggesting that the MGE is composed of Muntjac plus additional genetic payload.
Shotgun Illumina resequencing from a glyphosate resistant line with an *EPSPS* CNV was aligned
to the susceptible genome assembly. CNV detection was then performed based on changes in read depth
using CNVnator. Dozens of small CNV events were predicted between the glyphosate susceptible and
glyphosate resistant lines but none so large or nearly as high copy number as the *EPSPS* CNV event.
The MGE is present as a single copy in the susceptible genome chr9, and this MGE sequence from the
susceptible genome is nearly identical to the MGE located next to *EPSPS* on chr1 in the resistant line

(>99.9% sequence similarity). Breaks in read mapping coverage indicate the presence of a small structural variant between the MGE at chr9 in susceptible and the MGE on chr1 in resistant, which is also detected by CNVnator (Figure 3). RNA-Seq data from the two populations show that slight changes in predicted genomic copy number do not correlate well with expression changes for several the 7 genes located in the *EPSPS* CNV. The additional copies of *EPSPS* resulted in higher levels of *EPSPS* expression in a linear correlation.

3.4. Muntjac activity and transposition

Muntjac and the MGE contain three key traits of an active autonomous transposase: 1) an intact Mutator domain, 2) target site duplication, and 3) TIRs. The results of CNVnator and read mapping indicate that Muntjac is internal of the MGE that is present in the CNV related to herbicide resistance in kochia. Interruptions and variations in the aligned reads are due to the MGE containing Muntjac that occurs at an entirely different locus within the glyphosate resistant line consistent with transposase activity (Figure 3). Micro-assemblies of the susceptible Chr9 and resistant Chr1 insertion sites of the entire MGE provide further support for Muntjac as an autonomous Mutator Transposase (Figure 4). First, a target site duplication was detected at both the insertion next to *EPSPS* on Chr1 (~bp 6,110,000) in the resistant line as well as at its location on Chr9 (~bp 98,280,000) in the assembled susceptible line. Second, a TIR of the appropriate length (88 bp) was identified adjacent to each TSD. A survey of protein domains within MuDR type demonstrates that Muntjac retains key conserved domains for transposase activity including a Mutator domain (MULE) and a SWIM zinc-finger (Figure 5). The detection of a putative YCF1 (Yeast Cadmium Factor-1) domain within the Muntjac predicted protein is unique and does not appear to be in any other *FHY3* or *FAR1* genes in the susceptible genome. Finally, phylogenetic analysis places Muntjac within the MuDR TEs, sister to *FHY3* and *FAR1*. This placement of Muntjac within the MuDR clade indicates that Muntjac, *FHY3,* and *FAR1* are more closely related than they are to Jittery (Figure 6). It should be noted that we also identified, through functional annotation, what might be another copy of Muntjac at a novel position on Chr1 (~bp 22,725,000); however, it was missing TIRs and had fused with another unrelated gene suggesting that this copy is now non-functional (Figure 4).

3.5. Molecular verification of Muntjac activity and transposition

Active MuDR elements are mobile and therefore are at different loci in different individuals. To determine if Muntjac exhibited mobility with the kochia genome, we designed five pairs of PCR primers to detect the absence/presence of Muntjac at Chr9 (~bp 98,280,000) (Table 3) in a variety of kochia plants from diverse locales and glyphosate resistance. For these assays, if the MGE at its position in Chr9, the primer pair F1/R1 will exhibit no bands since the amplicon is too long to be amplified with the applied settings. Additionally, if the MGE is present at the Chr9 locus, then primer pairs F1/R4 and F3/R3 should amplify since R4 and F3 are both located within the conserved Muntjac sequence in the MGE and F1 and R3 are located in the flanking region, effectively detecting both the upstream and downstream boundaries of the MGE. These primers were tested and validated using DNA from populations M32 and 7710 since we have genomic information about these lines. As predicted, the F1/R1 primer pair did not amplify in 7710 since the MGE fragment (~16,000 bp) was too long to be amplified under the PCR setup (Table 4). In contrast, M32 plants had the predicted ~900 bp target band, indicating the MGE was not in its Chr9 position in this resistant line. Both M32 and 7710 showed target length single bands with these two primer pairs. Primer pairs F1/R4 and F3/R3 amplified the junction between MGE and the insertion point, which further demonstrated the presence of the full MGE in Chr9. As expected, 7710 plants showed target band with both primers while M32 plants did not. Primer pairs F1/R2 and F2/R3 were used to verify that the assembled DNA sequence flanking the Chr9 insertion site was present and correct (Figure 7, Table 4).

We also screened novel kochia populations that had no genomic data, both with and without the *EPSPS* CNV and glyphosate resistance, to determine if the MGE containing Muntjac on Chr9 is the native location and the position on Chr1 in the resistant line M32 is the novel insertion site. We selected a resistant line from Oregon and plants from two susceptible populations CO-S and Israel. In these populations, plants Oregon-1, CO-S-1, Israel-1, and Israel-2 showed target length bands with primer F1/R1, suggesting the MGE is not present at the Chr 9 locus in these resistant and susceptible populations. As anticipated from this result the primer used to test the boundaries of the MGE, F1/R4

and F3/R3, also did not amplify (Table 4). Primer pairs F1/R2 and F2/R3 confirmed the DNA around the Chr9 insertion line was present and normal (Table 4). Therefore, in these plants, MGE is not inserted in the chromosome 9 locus as was found in the 7710 population. For plants Oregon-1 and CO-S-1, there were target bands under F1/R2 and F2/R3, but three primer pairs F1/R1, F1/R4 and F3/R3, used to detect the MGE presence/absence did not amplify and confirm its presence or absence. In Oregon-2, there was no MGE insertion at Chr9, as F1/R1 worked, and F1/R4 and F3/R3 did not. However, in this individual, multiple bands were identified with primers F1/R2 and F2/R3. Results in this plant suggest the presence of nucleotide variations in this region among different kochia individuals. The PCR product of F1/R1, F1/R2 and F2/R3 from each population was sequenced. All sequences were highly similar with the kochia genome template, although some indels were present. Overall, the primer pairs used for validation here can be applied to most kochia populations and demonstrated that MGE position varies with kochia plants/populations and that the MGE at the Chr9 position is not shared among all susceptible populations.

4. Discussion

4.1. The Kochia Genome

The genome of Kochia scoparia comprises approximately 1 Gb of genetic material, which is characterized by highly repetitive and challenging to assemble centromeres and telomeres, analogous to other chenopod species. Through an integrative approach that employed hybrid short and long read sequencing techniques, nucleosome contact mapping, and optical mapping, we achieved an assembly of approximately 83.8% of the total sequence data, covering 97% of the predicted genome length. Although numerous gaps remain in these scaffolds, the gaps are notably gene-poor and consist mainly of highly repetitive sequences. The assessment of gene space completeness using the embryophyte odb10 dataset via BUSCO estimates was found to be 95.0%. Although the development of newer sequencing and assembly technologies may allow for a more contiguous Kochia genome, such refinement was not necessary for the present study, given the assembly's comprehensive coverage of gene space and relevant EPSPS and Muntjac loci in our susceptible reference.i

Beta vulgaris represents the nearest sequenced species to kochia despite being phylogenetically distant. Like most chenopods, both species have nine chromosomes. Upon alignment of the two genomes, a significant degree of macro-synteny and collinearity is observed. However, owing to their evolutionary divergence, numerous inversions, gaps, and transpositions have occurred. Notably, the regions adjacent to the centromere are particularly divergent. It may be that misassembles of the kochia genome also account for some of these differences. Furthermore, available data indicates that the Beta vulgaris genome is estimated to encompass approximately 758 Mb, approximately 75% of Kochia's genome size. The basis of this discrepancy is not yet apparent; however, preliminary analysis indicates the presence of a high number of transposable elements (TEs) in Kochia. Notably, there is currently no evidence of any polyploidy events or significant genomic duplication in either species since their divergence.

4.2. FHY3/ FAR1-like gene family in kochia

When the annotated genes of kochia are compared to other sequenced chenopods and amaranths, there are several gene families with conserved gene functions that are significantly expanded in kochia according to our Café Analysis (Figure 2). Many of these enriched families contain functions related to TEs (*e.g.* IPR005162: Retrotransposon gag domain, IPR001878: Zinc finger, CCHC-type superfamily, and IPR00752: Zinc finger, SWIM-type); however, of all the enriched genes, members of the *FHY3/FAR1* family are particularly interesting (Figure 1). Past research indicated that one of these *FHY3/FAR1* genes was in a MGE that caused a CNV of the gene *EPSPS*. This CNV, in turn, endows resistance to the herbicide glyphosate and is the major mechanism by which kochia has adapted to this herbicide[18,26]. *FHY3/FAR1* are generally considered one of the rare cases of 'domesticated' transposons and have been though to lose their ability to readily translocate and self-duplicate[13,66]. The expansion of the *FHY3/FAR1* may indicate that the *FHY3/FAR1* genes have either retained their transposable element capabilities or they have re-acquired transposable element capabilities in the *Bassia scoparia* lineage.

The interaction between genomic rearrangement(s) and adaptation underscores the profound impact that genomic plasticity can have on plant evolution, particularly under sustained abiotic stresses. While

not a novel concept, previous research has shown that TE activity and CNVs are essential pathways for generating novel genetic diversity. Nevertheless, the potential contribution of structural variation as a driver of adaptive genetic variation is often underestimated as supported by several studies[67,68]. Structural variation has the unique advantage of changing gene expression without changing the protein structure or the regulation of the promotor region. This process is not stable and instead is dynamic and responsive to abiotic stress with CNV numbers being driven higher by higher levels of selection[69].

This study utilized deep-coverage, short read genomic sequences of a glyphosate resistant individual to examine the *EPSPS* locus and the *FHY3/FAR1* MGE described previously [17] for mapping the location of the MGE. A genome-wide survey was conducted to explore potential loci, beyond the EPSPS locus, that demonstrated high frequency within the resistant genome. As anticipated, given the sequence divergence between the two Kochia individuals, the analysis revealed hundreds of small candidate copy number variations (CNVs) at both lower and higher copy number than the EPSPS locus. Many of the CNVs at very high read depths contained genes annotated as TEs. Most CNVs had little correlation to gene expression, indicating that the plants were able to maintain some level of control over the expression of duplicated genes from these other CNVs, the exception being *EPSPS* in the *EPSPS* CNV.

4.3. Muntjac activity and transposition

Muntjac's locus is variable within the genomes of several kochia samples. We designed PCR markers that amplify if, and only if, the MGE is at its position on chr9 like the susceptible genome. Additionally, we developed markers that amplify only if the MGE is absent (Figure 7b). The former primers span the upstream and downstream boundaries of the MGE and the surrounding genome, while the latter span the insertion site. If the entire MGE is present, the amplicon is too large to amplify (~16 Kb) but if it is absent, it is only 900 bp. We ran these molecular markers on the susceptible kochia line from which the genome was developed as well as from the glyphosate resistant line from which we first identified Muntjac. We confirm that Muntjac-MGE is indeed missing from its position on chr9 in the glyphosate resistant line, M32, while it is present in the glyphosate susceptible line, 7710. From these data we hypothesized that the Muntjac-MGE transposed from its position on chr9 to the EPSPS locus on chr1 using a "cut-and paste" mechanism. To test whether chr9 is indeed the "native" location for Muntjac we used our markers on two unique susceptible lines (divergent to the genome line) and a glyphosate resistant line unrelated to the glyphosate resistant line we have been working with. Muntjac was not at the position on chr9 or the position on chr1 (underlying resistance in concert with the *EPSPS* CNV), in any of these populations or individuals, which indicates that a conserved, original, locus of Muntjac is still unknown if it exists at all, and that the location on chr9 in the susceptible genome and insertion in chr1 in the resistant line are due to independent and recent insertions at these loci. Recently, *EPSPS* duplication without Muntjac as part of the CNV has been identified in some kochia populations, indicating multiple unique paths to *EPSPS* copy number variation[27]. This suggests high levels of structural variation and therefore genetic diversity exists in the kochia species as a whole.

The *FHY3/ FAR1* gene in the MGE that caused the *EPSPS* CNV is complete, containing key protein domains for transposition. Close molecular examination of the MGE and Muntjac in R lines from Oregon show evidence of differential transposition when compared to resistant and susceptible lines from Colorado and Israel. This line of evidence demonstrates the variability of the insertion site described from M32 on chr9 and suggests that the insertion that caused *EPSPS* duplication in M32 occurred recently. The detection of TSD, TIRs, and the apparent current mobility of Muntjac provides strong lines of evidence supporting active transposition of Muntjac within kochia.

Conclusion

We assembled the genome of a glyphosate susceptible kochia plant to chromosome scale. By performing orthologous gene clustering of related chenopods and amaranths, we observed an expansion in genes containing *FHY3* and/or *FAR1* annotations in kochia. Resequencing of a glyphosate resistant plant using high coverage Illumina data revealed the location of a MGE that is involved in the formation of the *EPSPS* CNV. Within this element, a gene annotated as having *FHY3* and *FAR1* protein domains was identified. Furthermore, this element had signature target-site duplications, terminal inverted repeats, and a functional transposase domain in the *FHY3/FAR1* gene. We believe this element has all the necessary features to be an autonomous Class II transposon and represents a "de-domestication" of the *FHY3/FAR1* like genes. We therefore rename this element as "Muntjac" to distinguish it from other

Mules and *FHY3/FAR1* related genes. We verified using molecular markers that Muntjac exists in at least two locations: Chr9 (~bp 98,280,000) in the glyphosate susceptible individual we assembled and flanking the glyphosate target gene *EPSPS* on Chr1 (~bp 6,110,000) in a glyphosate resistant individual. Furthermore, we used these markers to attempt to decipher which one of these locations is Muntjac's "native" location; however, in other susceptible lines (namely plants from Israel, from kochia's native range), Muntjac is at neither location. We hypothesize that Muntjac is still very active within kochia and that its native location and recent activity may only be resolvable using a more in-depth pan-genomic approach. Regardless, we believe Muntjac's involvement with generating glyphosate resistance alleles is an strong demonstration of the critical role TEs play with the production of novel variation and therefore the adaptive potential of weedy species.

Data availability.
The assembled genomes, associated GFF annotation files, and all functional annotation information are publicly available through the International Weed Genomics Consortium online database, 'Weedpedia' (https://weedpedia.weedgenomics.org/).

Acknowledgement

This work was partially supported by USDA-NIFA Hatch project COL00783 at Colorado State University.

References

References

1.   Lisch, D. How important are transposons for plant evolution? *Nature Reviews Genetics* vol. 14
49–61 (2013).

2.   Finnegan, D. J. Eukaryotic transposable elements and genome evolution. *Trends Genet.* **5**,
103–107 (1989).

3.   Lisch, D. Epigenetic regulation of transposable elements in plants. *Annu. Rev. Plant Biol.* **60**,
43–66 (2009).

4.   Burgess, D., Li, H., Zhao, M., Kim, S. Y. & Lisch, D. Silencing of mutator elements in maize
involves distinct populations of small RNAs and distinct patterns of DNA methylation.
Genetics **215**, 379–391 (2020).

5.   Bennetzen, J. L. Transposable element contributions to plant gene and genome evolution.
Plant Mol. Biol. **42**, 251–269 (2000).

6.   Devos, K. M., Brown, J. K. M. & Bennetzen, J. L. Genome size reduction through illegitimate
recombination counteracts genome expansion in Arabidopsis. *Genome Res.* **12**, 1075–1079
(2002).

7.   Hsing, Y. I. *et al.* A rice gene activation/knockout mutant resource for high throughput
functional genomics. *Plant Mol. Biol.* **63**, 351–364 (2007).

8.   Candela, H. & Hake, S. The art and design of genetic screens: Maize. *Nat. Rev. Genet.* **9**, 192–
203 (2008).

9.   Juretic, N., Hoen, D. R., Huynh, M. L., Harrison, P. M. & Bureau, T. E. The evolutionary fate
of MULE-mediated duplications of host gene fragments in rice. *Genome Res.* **15**, 1292–1297
(2005).

10.   Dupeyron, M., Singh, K. S., Bass, C. & Hayward, A. Evolution of Mutator transposable
elements across eukaryotic diversity. *Mob. DNA* **10**, (2019).

11.   Lisch, D. Mutator and MULE Transposons . *Microbiol. Spectr.* **3**, (2015).

12.   Wells, J. N. & Feschotte, C. A Field Guide to Eukaryotic Transposable Elements. *Annu. Rev.*
Genet. **54**, 539–561 (2020).

13.   Lisch, D. R., Freeling, M., Langham, R. J. & Choy, M. Y. Mutator transposase is widespread
in the grasses. *Plant Physiol.* **125**, 1293–1303 (2001).

14.   Xu, Z. *et al.* Jittery, a Mutator distant relative with a paradoxical mobile behavior: Excision
without reinsertion. *Plant Cell* **16**, 1105–1114 (2004).

15.   Kumar, V., Jha, P., Jugulam, M., Yadav, R. & Stahlman, P. W. Herbicide-resistant kochia
(Bassia scoparia) in North America: A review. *Weed Sci.* **67**, 4–15 (2019).

16.   Friesen, L. F., Beckie, H. J., Warwick, S. I. & Van Acker, R. C. The biology of Canadian
weeds. 138. Kochia scoparia (l.) schrad. *Can. J. Plant Sci.* **89**, 141–167 (2009).

17.   Patterson, E. L. *et al.* The Draft Genome of Kochia scoparia and the Mechanism of Glyphosate
Resistance via Transposon-Mediated EPSPS Tandem Gene Duplication. *Genome Biol. Evol.*
11, 2927–2940 (2019).

18. Wiersma, A. T. *et al.* Gene amplification of 5-enol-pyruvylshikimate-3-phosphate synthase in
glyphosate-resistant Kochia scoparia. *Planta* **241**, 463–474 (2015).

19. Jugulam, M. *et al.* Tandem amplification of a chromosomal segment harboring 5-
Enolpyruvylshikimate-3-Phosphate synthase locus confers glyphosate resistance in Kochia
scoparia. *Plant Physiol.* **166**, 1200–1207 (2014).

20. Grossman, L. *The magician king*. (2012).

21. Grossman, L. *The Magician's Land*. (2015).

22. Preston, C., Belles, D. S., Westra, P. H., Nissen, S. J. & Ward, S. M. Inheritance of Resistance
to The Auxinic Herbicide Dicamba in Kochia (Kochia scoparia). *Weed Sci.* **57**, 43–47 (2009).

23. Pettinga, D. J. *et al.* Increased chalcone synthase (CHS) expression is associated with dicamba
resistance in Kochia scoparia. *Pest Manag. Sci.* **74**, 2306–2315 (2018).

24. Busi, R. *et al.* Weed resistance to synthetic auxin herbicides. *Pest Manag. Sci.* **74**, 2265–2276
(2018).

25. Westra, E. P., Nissen, S. J., Getts, T. J., Westra, P. & Gaines, T. A. Survey reveals frequency
of multiple resistance to glyphosate and dicamba in kochia (Bassia scoparia). *Weed Technol.*
33, 664–672 (2019).

26. Gaines, T. A. *et al.* EPSPS Gene Copy Number and Whole-Plant Glyphosate Resistance Level
in Kochia scoparia. *PLoS One* **11**, 1–11 (2016).

27. Ravet, K. *et al.* Genomic-based epidemiology reveals independent origins and gene flow of
glyphosate resistance in Bassia scoparia populations across North America. *Mol. Ecol.* **30**,
5343–5359 (2021).

28. Jinlu, L., Shuo, W., Jing, Y., Ling, W. & Shiliang, Z. A Modified CTAB Protocol for Plant
DNA Extraction. *Chinese Bull. Bot.* **48**, 72–78 (2013).

29. Doyle J J & Doyle J L. A rapid DNA isolation procedure for small quantities of fresh leaf
tissue. *Phytochem. Bull.* **19**, 11–15 (1987).

30. Cheng, Y. *et al.* Studies on genome size estimation, chromosome number, gametophyte
development and plant morphology of salt-tolerant halophyte Suaeda salsa. *BMC Plant Biol.*
19, (2019).

31. Xiao, C. Le *et al.* MECAT: Fast mapping, error correction, and de novo assembly for single-
molecule sequencing reads. *Nat. Methods* **14**, 1072–1074 (2017).

32. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

33. Walker, B. J. *et al.* Pilon: An integrated tool for comprehensive microbial variant detection and
genome assembly improvement. *PLoS One* **9**, (2014).

34. Huang, S., Kang, M. & Xu, A. HaploMerger2: Rebuilding both haploid sub-assemblies from
high-heterozygosity diploid genome assembly. *Bioinformatics* **33**, 2577–2579 (2017).

35. Ghurye, J., Pop, M., Koren, S., Bickhart, D. & Chin, C. S. Scaffolding of long read assemblies
using long range contact information. *BMC Genomics* **18**, (2017).

36.   Staňková, H. *et al.* BioNano genome mapping of individual chromosomes supports physical
mapping and sequence assembly in complex plant genomes. *Plant Biotechnol. J.* **14**, 1523–
1531 (2016).

37.   English, A. C. *et al.* Mind the Gap: Upgrading Genomes with Pacific Biosciences RS Long-
Read Sequencing Technology. *PLoS One* **7**, (2012).

38.   Forcato, M. *et al.* Comparison of computational methods for Hi-C data analysis. *Nat. Methods*
14, 679–685 (2017).

39.   Humann, J. L., Lee, T., Ficklin, S. & Main, D. Structural and functional annotation of
eukaryotic genomes with GenSAS. *Methods Mol. Biol.* **1962**, 29–51 (2019).

40.   Tempel, S. Using and understanding repeatMasker. *Methods Mol. Biol.* **859**, 29–51 (2012).

41.   Smit, A. & Hubley, R. RepeatModeler open-1.0. *Available at http://www.repeatmasker.org*
2008–2015 (2015).

42.   Cock, P. J. A., Chilton, J. M., Grüning, B., Johnson, J. E. & Soranzo, N. NCBI BLAST+
integrated into Galaxy. *Gigascience* **4**, (2015).

43.   D, K. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions,
deletions, and gene fusions. *Genome Biol.* **14**, R36 (2013).

44.   James, K. W. BLAT---The BLAST-Like Alignment Tool. *Genome Res.* **12**, 656–664 (2002).

45.   Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using EVidenceModeler and
the Program to Assemble Spliced Alignments. *Genome Biol.* **9**, (2008).

46.   Stanke, M., Steinkamp, R., Waack, S. & Morgenstern, B. AUGUSTUS: A web server for gene
finding in eukaryotes. *Nucleic Acids Res.* **32**, (2004).

47.   Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M. & Stanke, M. BRAKER1:
Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and AUGUSTUS.
Bioinformatics **32**, 767–769 (2016).

48.   Lomsadze, A., Ter-Hovhannisyan, V., Chernoff, Y. O. & Borodovsky, M. Gene identification
in novel eukaryotic genomes by self-training algorithm. *Nucleic Acids Res.* **33**, 6494–6506
(2005).

49.   Ter-Hovhannisyan, V., Lomsadze, A., Chernoff, Y. O. & Borodovsky, M. Gene prediction in
novel fungal genomes using an ab initio algorithm with unsupervised training. *Genome Res.*
18, 1979–1990 (2008).

50.   Cock, P. J. A., Grüning, B. A., Paszkiewicz, K. & Pritchard, L. Galaxy tools and workflows
for sequence analysis with applications in molecular plant pathology. *PeerJ* **2013**, (2013).

51.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the
analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

52.   Suzek, B. E., Huang, H., McGarvey, P., Mazumder, R. & Wu, C. H. UniRef: Comprehensive
and non-redundant UniProt reference clusters. *Bioinformatics* **23**, 1282–1288 (2007).

53.   Waterhouse, R. M. *et al.* BUSCO applications from quality assessments to gene prediction and
phylogenomics. *Mol. Biol. Evol.* **35**, 543–548 (2018).

54. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, (2019).

55. Sims, D. *et al.* CGAT: Computational genomics analysis toolkit. *Bioinformatics* **30**, 1290–
1291 (2014).

56. De Bie, T., Cristianini, N., Demuth, J. P. & Hahn, M. W. CAFE: A computational tool for the
study of gene family evolution. *Bioinformatics* **22**, 1269–1271 (2006).

57. Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and Visualization of
Phylogenomic Data. *Mol. Biol. Evol.* **33**, 1635–1638 (2016).

58. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).

59. Anton, B. *et al.* SPAdes: a new genome assembly algorithm and its applications to single-cell
sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

60. Abyzov, A., Urban, A. E., Snyder, M. & Gerstein, M. CNVnator: An approach to discover,
genotype, and characterize typical and atypical CNVs from family and population genome
sequencing. *Genome Res.* **21**, 974–984 (2011).

61. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2009).

62. Patterson, E. L., Pettinga, D. J., Ravet, K., Neve, P. & Gaines, T. A. Glyphosate resistance and
EPSPS gene duplication: Convergent evolution in multiple plant species. *J. Hered.* **109**, 117–
125 (2018).

63. Gaines, T. A., Patterson, E. L. & Neve, P. Molecular mechanisms of adaptive evolution
revealed by global selection for glyphosate resistance. *New Phytol.* **223**, 1770–1775 (2019).

64. Gaines, T. A. *et al.* Gene amplification confers glyphosate resistance in Amaranthus palmeri.
Proc. Natl. Acad. Sci. U. S. A. **107**, 1029–1034 (2010).

65. Martin, S. L. *et al.* High gene flow maintains genetic diversity following selection for high
EPSPS copy number in the weed kochia (Amaranthaceae). *Sci. Rep.* **10**, (2020).

66. Hudson, M. E., Lisch, D. R. & Quail, P. H. The FHY3 and FAR1 genes encode transposase-
related proteins involved in regulation of gene expression by the phytochrome A-signaling
pathway. *Plant J.* **34**, 453–471 (2003).

67. Hardigan, M., Crisovan, E., Hamilton, J. & Kim, J. Genome reduction uncovers a large
dispensable genome and adaptive role for copy number variation in asexually propagated
Solanum tuberosum. *The Plant* (2016).

68. Fuentes, R. R. *et al.* Structural variants in 3000 rice genomes. *Genome Res* **29**, 870–880
(2019).

69. Horváth, V., Merenciano, M. & González, J. Revisiting the Relationship between
Transposable Elements and the Eukaryotic Stress Response. *Trends Genet.* **33**, 832–841
(2017).

Figures

Figure 1: Circos plot of *Bassia scoparia* (kochia) genome assembly; a) chromosomes (chr1..chr9) and the two largest unplaced scaffolds (u1,u2) with gaps larger than 10 Mb indicated in gray; b) the placement of genes were part of a target set of genes containing some combination of the following Interpro designations *FHY1* (IPR037766), *FAR1* DNA binding domain (IPR004330), and *FHY3/ FAR1* family (IPR031052); c) repetitive sequence coverage per 1 Mb; d) gene coverage per 1 Mb; e) segmental duplications detected using MCScanX. Red lines indicate segmental duplications contains a target set gene. Green lines do not contain a target set gene.

Figure 2: A cladogram of *Bassia scoparia* and several other *Amaranthaceae* species that have
sequenced genomes. Significant gene family expansions for each clade are listed above the branch in
blue while significant gene family contractions are reported below the branch in orange as calculated
by Computational Analysis of gene Family Evolution (Cafe).

Figure 3: Genomic DNA Illumina reads from the resistant *Bassia scoparia* (kochia) line (M32) mapped to the susceptible genome. Putative copy number
variants predicted using CNVnator. Structural variants predicted using complete breaks in read mapping, and micro assembly.

Figure 4: Micro assemblies from the glyphosate resistant *Bassia scoparia* (kochia) line (M32) were
assembled for two Muntjac insertions in the susceptible line (7710). a) A completely intact Muntjac
insertion on Chr9, with detected target site duplication (TSD), and detectable terminal inverted repeats
(TIR). b) A fragmented insertion on Chr1. The Muntjac gene has been pseudogenized or incorporated
into another a gene. The TSD provides evidence of insertion, but the fragmentation of Muntjac and
the absence of TIRs suggest that this insertion on Chr1 is no longer active.
637

639 Figure 5: A reduced gene tree produced from an alignment of representative MuDR sequences searched against representative plant clades showing the
640 representative domains within the MuDR clade. *FHY3/ FAR1* domain was detected in all sequences but is not shown for clarity.
641

Figure 6: Simplified gene tree showing relationship of named MuDRs, *FHY3*, *FAR1*, and Muntjac.
Muntjac clade is sister to *FHY3/FAR1* clade. The Muntjac clade diverged from *FHY3/FAR1* at the
base of angiosperms, but it is more closely related to *FHY3/FAR1* than Jittery. Gene tree was
produced taking representative MuDR sequences and searching against representative plant clades.
Alignment was produced with COBALT, trimmed with GGBlocks and phylogeny estimated using
IQTree.
649

651 Figure 7: Diagram of the primer positions used to verify the presence/absence of MGE sequence on
652 Chr9 in glyphosate resistant and susceptible *Bassia scoparia* (kochia) individuals.

Kochia Chromosome 9

653
654

655 **Tables**
656 Table 1: BUSCO scores for the kochia transcriptome using the Embryophyta_ODB 10 database.
657

	Number of BUSCOs	Percentage of BUSCOs
Total BUSCO groups searched	1,614	
Complete BUSCOs (C)	1,505	93.3%
Complete and single-copy BUSCOs (S)	1,468	91.0%
Complete and duplicated BUSCOs (D)	37	2.3%
Fragmented BUSCOs (F)	27	1.7%
Missing BUSCOs (M)	82	5.0%

Table 2: A summary of the output from Computational Analysis of gene Family Evolution (Cafe) indicating where in the evolution of the Bassia scoparia gene family expansions and contractions occurred. Terminal branches are labeled with the species while internal branches are bolded and species included in those branches are made explicit using a 6 character Genus:species ID.

Branch Label	Expansions	Genes Gained	Contractions	Genes Lost	Average Expansion	Significant Expansions	Significant Contractions
Beta vulgaris	16	19	25,664	27,667	-1.008	0	4
AmaHyp, AmaHyb, AmaPal, AmaTub, CheQui, AtrHor, SpiOle, BasSco	142	175	0	0	0.006	0	0
All Amaranthus Sp.	656	794	8,019	8,050	-0.265	16	1
Amaranthus tuberculatus	3,750	6,811	5,063	5,235	0.057	44	0
AmaHyp, AmaHyb, AmaPal	462	550	993	1,010	-0.017	6	0
Amaranthus palmeri	3,989	8,704	5,312	5,480	0.118	109	2
AmaHyp, AmaHyb	926	1,029	1,960	2,111	-0.039	5	8
Amaranthus hybridus	1,423	2,588	3,385	3,560	-0.035	58	11
Amaranthus hypochondriacus	2,059	2,638	2,082	2,484	0.006	22	55
CheQui, AtrHor, SpiOle, BasSco	928	1,231	4,161	4,181	-0.108	20	1
Bassia scoparia	5,331	22,938	6,405	6,541	0.598	206	0
CheQui, AtrHor, SpiOle	837	1,081	3,002	3,147	-0.075	17	1
Spinacia oleracea	3,334	10,766	4,605	5,272	0.200	107	26
CheQui, AtrHor	1,857	3,344	2,399	2,513	0.030	110	6
Chenopodium quinoa	11,787	18,049	2,284	2,907	0.552	130	50
Atriplex hortensis	2,170	8,704	4,121	4,713	0.146	217	37

664 Table 3 Information of kochia MGE validation primers

Assay	Primer name	Primer sequence (5' to 3')	Target length (bp)	Annealing temperature (°C)	Function
1	F1	CATCTTCTCTTCTCAACCACTC	904		Test for the presence of MGE
	R1	ACCACACCCACTCATTCAC			
2	F1	CATCTTCTCTTCTCAACCACTC	403		Amplify upstream of MGE insertion site
	R2	CGGAAAACAACACTTAGAATGC			
3	F2	GTGAATGAGTGGGTGTGGT	222	55	Amplify downstream of MGE insertion site
	R3	TGAACAACTTAGCAGCAAGG			
4	F1	CATCTTCTCTTCTCAACCACTC	993		Amplify the upstream junction of MGE insertion
	R4	CTACAGTCAACAAAACCAAACC			
5	F3	ACCCAAATTCTTCAACAGCAC	931		Amplify the downstream junction of MGE insertion
	R3	TGAACAACTTAGCAGCAAGG			

666 Table 4 Results of MGE validation primer pairs.

Assay	Primer pair	Populations									
		Oregon		CO-S		Israel		M32		7710 (S mixture)	
		Oregon-1	Oregon-2	CO-S-1	CO-S-2	Israel-1	Israel-2	M32-1	M32-2	S-1	S-2
1	F1/R1	×	✓	×	✓	✓	✓	✓	✓	×	×
2	F1/R2	✓	× *	✓	✓	✓	✓	✓	✓	✓	✓
3	F2/R3	✓	× *	✓	✓	✓	✓	✓	✓	✓	✓
4	F1/R4	×	×	×	×	×	×	×	×	✓	✓
5	F3/R3	×	×	×	×	×	×	×	×	✓	✓

667 ✓: band of/close to target length amplified

668 ×: band of/close to target length not amplified

669 * Multiple bands amplified