Complexes of vertebrate TMC1/2 and CIB2/3 proteins form hair-cell

mechanotransduction cation channels

Arnaud P. J. Giese[1,#,*], Wei-Hsiang Weng[2,3,*], Katie S. Kindt[4], Hui Ho Vanessa Chang[5],

Jonathan S. Montgomery[2,6], Evan M. Ratzan[7], Alisha J. Beirl[4], Roberto Aponte Rivera[4],

Jeffrey M. Lotthammer[2], Sanket Walujkar[2], Mark P. Foster[2,3,6], Omid A. Zobeiri[8], Jeffrey R.

Holt[7], Saima Riazuddin[1], Kathleen E. Cullen[9], Marcos Sotomayor[2,3,6] , Zubair M.

Ahmed[1,10,11]

[1]Department of Otorhinolaryngology – Head & Neck Surgery, University of Maryland

School of Medicine, Baltimore, MD, USA.

[2]Department of Chemistry and Biochemistry, The Ohio State University, Columbus, OH,

USA.

[3]Biophysics Graduate Program, The Ohio State University, Columbus, OH, USA.

[4]Section on Sensory Cell Development and Function, National Institute on Deafness and

Other Communication Disorders, National Institutes of Health, Bethesda, MD, USA.

[5]Department of Physiology, McGill University, Montreal, QC, Canada.

[6]Ohio State Biochemistry Program, The Ohio State University, Columbus, OH, USA.

[7]Departments of Otolaryngology and Neurology, F.M. Kirby Neurobiology Center, Boston

Children's Hospital, Harvard Medical School, Boston, MA, USA.

[8]Department of Biomedical Engineering, McGill University, Montreal, QC, Canada.

[9]Departments of Biomedical Engineering, Neuroscience, and Otolaryngology and Head and

Neck Surgery, Johns Hopkins University School of Medicine, Baltimore, MD, USA.

[10]Department of Biochemistry and Molecular Biology, University of Maryland School of

Medicine, Baltimore, MD, USA.

[11]Department of Ophthalmology and Visual Sciences, University of Maryland School of

Medicine, Baltimore, MD, USA.

[#]Current address: SENSORION SA, Montpellier, France.

*Contributed equally.

Correspondance:

Emails: sotomayor.8@osu.edu; zmahmed@som.umaryland.edu

Keywords: CIB2, CIB3, mechanotransduction, TMC1, TMC2, Vestibular hair cells

Abstract

Calcium and integrin-binding protein 2 (CIB2) and CIB3 bind to transmembrane

channel-like 1 (TMC1) and TMC2, the pore-forming subunits of the inner-ear mechano-

electrical transduction (MET) apparatus. Whether these interactions are functionally relevant

across mechanosensory organs and vertebrate species is unclear. Here we show that both CIB2

and CIB3 can form heteromeric complexes with TMC1 and TMC2 and are integral for MET

function in mouse cochlea and vestibular end organs as well as in zebrafish inner ear and lateral

line. Our AlphaFold 2 models suggest that vertebrate CIB proteins can simultaneously interact

with at least two cytoplasmic domains of TMC1 and TMC2 as validated using nuclear magnetic

resonance spectroscopy of TMC1 fragments interacting with CIB2 and CIB3. Molecular

dynamics simulations of TMC1/2 complexes with CIB2/3 predict that TMCs are structurally

stabilized by CIB proteins to form cation channels. Overall, our work demonstrates that intact

CIB2/3 and TMC1/2 complexes are integral to hair-cell MET function in vertebrate

mechanosensory epithelia.

Introduction

The vertebrate senses of hearing and balance depend on the process of mechano-electrical transduction (MET) in inner-ear hair cells. This process is initiated when mechanical forces from sound waves and head movements deflect specialized mechanosensory projections at the apical surface of hair cells, known as stereocilia (reviewed in Fettiplace, 2017; Frolenkov et al., 2004; Gillespie and Müller, 2009). Deflections of stereocilia result in tensioning of the extracellular tip links that convey mechanical force to the MET channels. Mature tip links are heterotetrameric filaments composed of cadherin 23 (CDH23) and protocadherin 15 (PCDH15) (Ahmed et al., 2006; Kazmierczak et al., 2007; Siemens et al., 2004; Sotomayor et al., 2012). In mammalian auditory hair cells, MET channels are localized at the PCDH15 end of the tip link (Beurg et al., 2009). Significant progress in our understanding of MET has been recently made by identifying the transmembrane channel-like proteins 1 and 2 (TMC1 and TMC2) as the pore-forming subunits of the MET channel (Kawashima et al., 2011; Pan et al., 2018). In hair cells, two other transmembrane proteins, LHFPL5 and TMIE, are essential for sustained hearing. LHFPL5 binds to PCDH15 and may regulate force transmission to TMC1 and TMC2 (Qiu et al., 2023; Xiong et al., 2012). TMIE also interacts with TMC1/2 and regulates unitary conductance and ion selectivity of the MET channels (Cunningham et al., 2020; Naz et al., 2002). Interestingly, MET currents can still be recorded from hair cells lacking LHFPL5 or TMIE, suggesting that neither are essential for channel function (Fettiplace et al., 2022).

We previously showed that calcium (Ca^{2+}) and integrin-binding protein 2 (CIB2), and the closely related CIB3, interact with TMC1/2 (Giese et al., 2017). Furthermore, we also demonstrated that CIB2 is essential for MET function in auditory hair cells (Giese et al., 2017). Recessive variants of CIB2 predominantly cause nonsyndromic DFNB48 hearing loss in

humans (Booth et al., 2018; Michel et al., 2017; Riazuddin et al., 2012; Seco et al., 2016; Wang

et al., 2017). Similarly, recessive alleles of *Cib2* cause hearing loss in mice but no vestibular

deficits (Giese et al., 2017; Michel et al., 2017; Wang et al., 2017). CIB2 belongs to a family

of four closely related proteins (CIB1-4) that have similar structural domains and have partial

functional redundancy (Huang et al., 2012). Wang and co-authors investigated the function of

Cib1 in cochlear hair cells through generation of a knockout strain and concluded that although

Cib1 is expressed in the mouse cochlear hair cells, loss of CIB1 protein did not affect auditory

function (Wang et al., 2017). A recent study demonstrated that while CIB2 is essential for

proper localization of TMC1/2 at the tips of auditory hair cell stereocilia, CIB3 can restore

MET currents in cochlear hair cells lacking CIB2 (Liang et al., 2021). The same study reported

a crystal structure of a TMC1 intracellular linker in complex with CIB3, which suggests that

CIB2/3 proteins are likely to be auxiliary subunits of hair cell MET channels (Liang et al.,

2021).

Previous work on CIB proteins primarily focused on auditory hair cells, leaving open

the question of whether they play a role in vestibular function. This question was addressed in

part by another recent study that reported vestibular deficits in mice lacking both CIB2 and

CIB3 (Wang et al., 2023). However, whether CIB2/3 can function and interact with TMC1/2

proteins across sensory organs, hair-cell types, and species is still unclear. Here we provide

further evidence that CIB2 and CIB3 are indeed evolutionary conserved and functionally

redundant components of the vertebrate transduction apparatus necessary for MET channel

activity in mouse vestibular hair cells, as well as in zebrafish mechanosensory epithelia.

Moreover, based on our structural analyses and simulations, we propose that CIB2 and CIB3

are required for stabilization and channel function of the MET proteins TMC1 and TMC2.

Results

CIB2 and CIB3 are expressed in mouse vestibular hair cells

We and others previously demonstrated that CIB2 is essential for MET in auditory hair cells (Giese et al., 2017; Michel et al., 2017; Wang et al., 2017). In contrast to auditory hair cells, we found that the vestibular hair cells in $Cib2^{KO/KO}$ mice have intact MET. We assessed MET via uptake of FM 1-43 (Figure 1A), a styryl dye that permeates into hair cells through functional MET channels (Meyers et al., 2003), indicating that there may be another CIB protein playing a functionally redundant role. Bulk RNA sequencing revealed that all four members of the Cib family are expressed in mouse inner ear tissues (Figure 1B). However, when assessed by single cell RNA sequencing analysis, only $Cib2$ and $Cib3$ are expressed in immature hair cells in the mouse utricle (Figure 1C), a vestibular sensory organ required to detect linear acceleration (McInturff et al., 2018). Further, we find that the expression of $Cib2$ and $Cib3$ is maintained in mature type 1 and type 2 utricular hair cells (Figure 1C). We also examined the expression of $Tmc1$ and $Tmc2$ in utricular hair cells and found that in immature hair cells $Tmc1$ is highly expressed, while $Tmc2$ expression is low (Figure 1C). $Tmc2$ expression increases during development and becomes comparable to $Tmc1$ in both type 1 and type 2 hair cells upon maturation (Figure 1C). CIB2 and CIB3 have been reported to interact with TMC1/2, and CIB2 is essential for auditory hair cell MET (Giese et al., 2017; Liang et al., 2021). The co-expression of $Cib2$ and $Cib3$ in developing and mature vestibular hair bundles, and the ability of CIB2/3 to interact with TMC1 and TMC2, all make them attractive candidates to regulate MET function in vestibular hair cells.

CIB3 is not required for hearing function in mice

To evaluate whether CIB3 regulates MET function both in auditory and vestibular hair

cells, we generated *Cib3* knockout (*Cib3^{KO}*) mice using CRISPR/Cas9 technology (Figure 1D).

An indel in exon 4 of *Cib3* was introduced leading to a nonsense mutation (p.Asp54*), thus

causing a loss of allele function (Figure 1E-F). We first characterized hearing function in

Cib3^{KO/KO} and control littermate mice at P16 by measuring auditory-evoked brainstem

responses (ABRs). Normal ABR waveforms and thresholds were observed in wild-type and

Cib3^{KO/KO} mice, indicating normal hearing (Figure 2A-B). Next, we investigated if there was a

functional redundancy between CIB2 and CIB3 in auditory hair cells. For this analysis, the

Cib3^{KO} allele was introduced into the *Cib2^{KO}* background. Similar to what has been previously

reported in *Cib2^{KO/KO}* mice (Giese et al., 2017; Michel et al., 2017; Wang et al., 2017), the

Cib2:Cib3 double knockout (*Cib2^{KO/KO};Cib3^{KO/KO}*) mice were also profoundly deaf. The

Cib2^{KO/KO};Cib3^{KO/KO} mice did not respond to click or strong tone burst stimuli at 100 dB sound

pressure level (SPL) (Figure 2A-B). As reported previously for *Cib2^{KO/KO}* mice, we observed

that auditory hair cells in *Cib2^{KO/KO};Cib3^{KO/KO}* mice did not accumulate FM 1-43 dye (Figure

2C). Also, similar to *Cib2^{KO/KO}* mice (Giese et al., 2017; Michel et al., 2017; Wang et al., 2017),

we observed progressive degeneration of auditory hair cells in *Cib2^{KO/KO};Cib3^{KO/KO}* mice

(Figure 2D). Taken together with previous studies, our results confirm that CIB2 is

indispensable for hearing and MET function in the auditory hair cells. Furthermore,

endogenous CIB3 is unable to compensate for CIB2 loss in the auditory hair cells, perhaps due

to extremely low expression level of CIB3 in these cells and the lack of compensatory

overexpression of CIB3 in the cochlea of *Cib2^{KO/KO}* mice (Giese et al., 2017).

CIB2 and CIB3 are required for vestibular function in mice

After we assessed auditory function, we turned our analyses toward vestibular function.

Similar to *Cib2^{KO/KO}* mice (Giese et al., 2017), *Cib3^{KO/KO}* mice did not display any obvious

indications of vestibular dysfunction, such as circling, hyperactivity or head bobbing,

performed well when exploratory behavior was tested, and had normal vestibulo-ocular

reflexes (Figure 3). In contrast, visual observation of the $Cib2^{KO/KO};Cib3^{KO/KO}$ mice indicated

altered exploratory behavior in an open-field test, suggesting an impairment of the vestibular

system (Figure 3A). To assess the implications of the $Cib2^{KO/KO};Cib3^{KO/KO}$ mice on vestibular

function, we next performed a series of tests, including: the vestibulo-ocular reflex (VOR),

performance in a rotating rod, and head movement analysis (Figure 3B-H). The VOR is

mediated by a direct three neuron pathway linking the vestibular nerve to the extraocular

muscles and produces compensatory eye movements in the direction opposite of the head

movement to provide stable gaze (reviewed in Cullen, 2019). The dynamic response properties

(i.e., gain and phase) of the VOR can be precisely quantified in response to rotations applied

at different frequencies in darkness (VORd). In response to VORd testing, $Cib2^{+/+};Cib3^{KO/KO}$

mice generated robust compensatory eye movements (Figure 3B, black trace). In contrast,

$Cib2^{KO/KO};Cib3^{KO/KO}$ mice generated negligible eye movements (Figure 3B, red trace).

Consistent with this observation, quantification of the VORd across frequencies (Figure 3C)

revealed gains near zero in the $Cib2^{KO/KO};Cib3^{KO/KO}$ mice that were significantly reduced

($p<0.05$) compared to those measured in $Cib2^{+/+};Cib3^{KO/KO}$ mice. Interestingly, the VORd

gains of $Cib2^{KO/+};Cib3^{KO/KO}$ mice also showed attenuation with a significant decrease at 2 Hz

($p<0.01$) compared to $Cib2^{+/+};Cib3^{KO/KO}$ mice. On the other hand, the response phase of VORd

eye movements were comparable for $Cib2^{KO/+};Cib3^{KO/KO}$ and $Cib2^{+/+};Cib3^{KO/KO}$ mice at all

frequencies ($p>0.05$) (Figure 3C).

As a control for possible changes in visual system and/or ocular motor pathway, we

also quantified the optokinetic reflex (OKR), which generates eye movements in responses to

a moving visual scene and in light. The OKR normally works together with the VOR to

stabilize the visual field on the retina. Both $Cib2^{+/+};Cib3^{KO/KO}$ and $Cib2^{KO/+};Cib3^{KO/KO}$ showed

robust OKN responses which similarly decreased as a function of frequency (Figure 3D;

$p>0.05$). Likewise, OKR gain values in $Cib2^{KO/KO};Cib3^{KO/KO}$ were comparable with the single

exception of a significant gain decrease at 0.2 Hz ($p<0.05$). Analysis of OKN similarly revealed

comparable response phase for all 3 groups of mice at all frequencies ($p>0.05$) with the

exception that the $Cib2^{KO/KO};Cib3^{KO/KO}$ mice demonstrated less lag at 0.8 and 2 Hz (Figure 3D-

E). Overall, the OKR responses of mutant and control mice were generally comparable.

In addition to generating compensatory VOR eye movements to stabilize gaze, the

vestibular system also plays a key role in the maintenance of posture and control of balance.

Thus, we also tested all 3 groups of mice on several vestibular tasks that assessed balance and

postural defects. First, quantification of open-field testing experiments confirmed that mice

exhibited circling – a behavior that is commonly associated with loss of vestibular function in

mice (Figure 3F; $p<0.002$). We also completed rotarod testing in which performance was

quantified by measuring the amount of time the mouse remained on the rotating rod as its

acceleration increased. While $Cib2^{+/+};Cib3^{KO/KO}$ and $Cib2^{KO/+};Cib3^{KO/KO}$ mice exhibited

similar performance, $Cib2^{KO/KO};Cib3^{KO/KO}$ mice exhibited significantly poorer performance

across all time points from day 1 through day 4 (Figure 3G, $p<0.002$). Finally, it was also

possible to distinguish $Cib2^{KO/KO};Cib3^{KO/KO}$ mice from the other two groups in their home

cages since they commonly displayed head bobbing, a trait also generally associated with

vestibular dysfunction in mice. To quantify head movement in our mice, we attached a 6D

MEMS module consisting of three gyroscopes and three linear accelerometers to the mouse's

head implant (see Methods). Analysis of power spectra of head motion revealed significantly

higher power for $Cib2^{KO/KO};Cib3^{KO/KO}$ mice in all six dimensions of motion, as compared to

$Cib2^{KO/+};Cib3^{KO/KO}$ and $Cib2^{+/+};Cib3^{KO/KO}$ mice (Figure 3H, compare red vs. blue and black).

Further, there were no significant differences in the head movements power spectra of

Cib2^{KO/+};Cib3^{KO/KO} and *Cib2^{+/+};Cib3^{KO/KO}* mice (Figure 3H, blue vs. black). Taken together

our findings indicate that *Cib2^{KO/KO};Cib3^{KO/KO}* mice show marked deficits in vestibulomotor

function when compared to both *Cib2^{KO/+};Cib3^{KO/KO}* and *Cib2^{+/+};Cib3^{KO/KO}* mice.

CIB2 and CIB3 are functional subunits of the MET apparatus in mouse vestibular hair

cells

To understand the root cause of vestibulomotor function deficits described above for

Cib2^{KO/KO};Cib3^{KO/KO} mice, we first analyzed vestibular hair bundle morphology using confocal

microscopy. High-resolution confocal imaging did not reveal any obvious vestibular hair cell

loss and hair bundles looked indistinguishable from control in *Cib2^{KO/KO};Cib3^{KO/KO}* mice

(Figure 4A). Next, we visualized MET activity in vestibular hair cells of saccular and utricular

end organs of *Cib2^{KO/KO};Cib3^{KO/KO}* mice using the styryl dye FM 1-43. These vestibular end

organs are divided into two main physically and functionally distinct domains, a central striolar

domain and a peripheral extrastriolar domain. At P5, FM 1-43 labeling of *Cib2^{KO/KO}; Cib3^{KO/KO}*

tissue failed to label vestibular hair cells in appreciable numbers (Figure 4B). Similarly, fixable

FM 1-43FX intraperitoneal injections at P60 also showed no dye uptake in *Cib2^{KO/KO};*

Cib3^{KO/KO} mice (Figure 4C). *Cib2^{KO/KO}* and *Cib3^{KO/KO}* single mutant mice had dye uptake both

in striolar and extrastriolar hair cells, although a subtle reduction in FM 1-43FX was observed

in extrastriolar hair cells of both single mutants at P60 (Figure 4C). Taken together, these

results indicate that CIB2 and CIB3 have functionally redundant but compulsory roles in the

vestibular hair cell MET complex.

Cib2 and Cib3 function is evolutionary conserved in zebrafish

Numerous studies have shown that the same proteins are required for hair cell MET in

zebrafish and mammals (Chen et al., 2020; Erickson et al., 2017; Ernest et al., 2000; Gleason

et al., 2009; Gopal et al., 2015; Maeda et al., 2017, 2014; Phillips et al., 2011; Seiler et al.,

2004; Söllner et al., 2004). Unlike mammals, the zebrafish inner ear does not have a cochlea.

Instead, zebrafish rely on their saccule for hearing and utricle for balance. Zebrafish also utilize

hair cells in a specialized sensory system, the lateral line, to detect local water movement. The

hair cells within both the zebrafish inner ear and lateral line share many morphologically

similarities to mammalian vestibular hair cells (Nicolson, 2017). Given the zebrafish reliance

on the saccule for hearing and the similarities between zebrafish hair cells to mammalian

vestibular hair cells, we tested whether Cib2 and/or Cib3 were required for hair cell MET in

the zebrafish inner ear or lateral line.

To examine the impact of Cib2 and Cib3 loss in zebrafish, we created mutants using

CRISPR-Cas9 based methods to target each homologue (Varshney et al., 2016). We first

examined auditory function by testing the acoustic startle reflex in *cib2* and *cib3* mutants. For

our experiments we used a Zantiks behavioral system to quantify the probability of generating

a startle in response to a vibrational tap stimulus. This response is thought to rely primarily on

hair cells in the zebrafish inner ear and lateral-line system. We found that, compared to

controls, the probability of a startle response was significantly reduced in *cib2* mutants (Figure

5A, control: 0.71 ± 0.04; *cib2*: 0.28 ± 0.06, $p = 0.005$). In contrast, *cib3* mutants startle with a

similar frequency compared to controls (Figure 5A, $p = 0.36$). Interestingly, contrary to

mammals, where CIB2 is crucial for hearing, acoustic startle responses were not completely

eliminated in *cib2* mutants. One possibility for this partial loss of acoustic startle is that Cib3

may be able to compensate for Cib2 in the zebrafish. Therefore, we examined acoustic startle

responses in *cib2;cib3* mutants. In *cib2;cib3* mutant animals we observed a complete absence

of startle responses (Figure 5A). Furthermore, we observed that *cib2;cib3* mutants but not *cib2*

or *cib3* mutants exhibited spontaneous circling behavior, an additional indicator of inner ear

dysfunction in zebrafish (Nicolson et al., 1998). Together our behavioral results indicate that

Cib2 and Cib3 are both required to ensure normal acoustic startle responses in zebrafish.

Furthermore, just loss of Cib2, but not Cib3 can impair auditory function in zebrafish.

Cib2 and Cib3 are required for MET in the zebrafish lateral line

Our results demonstrate that both Cib2 and Cib3 are required for proper auditory

function in zebrafish. This raises the question of whether, similar to mice, these proteins are

critical for MET function in zebrafish hair cells. Therefore, we next turned our analysis to hair

cells in neuromast organs of the lateral line where methods to assess MET are well-documented

and straightforward (Lukasz and Kindt, 2018; Seiler et al., 2004).

First, to assess whether there were gross alterations to MET in lateral-line hair cells we

examined uptake of the FM 1-43. We found that, compared to controls, the percent of hair cells

per neuromast labeling with FM 1-43 was significantly reduced in *cib2* mutants (Figure

5B,D,E, control: 97.60 ± 1.17; *cib2*: 37.53 ± 6.92, $p < 0.0001$). In contrast, a similar percent of

hair cells per neuromast labeled with FM 1-43 in *cib3* mutants, compared to controls (Figure

5B,D,F, *cib3*: 95.67 ± 1.87, $p = 0.97$). We also quantified the average intensity of the FM 1-43

in hair cells (Figure 5C). We found that, compared to controls, the average FM 1-43 label was

reduced in *cib2* but not *cib3* mutants (Figure 5C). The residual FM 1-43 label in *cib2* mutant

hair cells led us to hypothesize that in lateral-line hair cells, similar to its role in zebrafish

auditory function, Cib3 may be able to substitute for Cib2 in MET. To assess whether Cib3

can substitute for Cib2, we examined FM 1-43 labeling in *cib2;cib3* mutants. In *cib2;cib3*

mutant animals we observed a complete absence of FM 1-43 labeling in lateral-line hair cells

(Figure 5B,D,G). Thus, taken together the results of our FM 1-43 labeling analysis are

consistent with a requirement for both Cib2 and Cib3 in hair cell MET in the zebrafish lateral

line.

To obtain a more quantitative assessment of MET we used Ca^{2+} imaging to perform

evoked measurements of hair cell function. For this assessment, we used a transgenic line that

expresses a membrane-localized GCaMP6s (memGCaMP6s) specifically in hair cells (Figure

5H-I'). This transgenic line has previously been used to measure Ca^{2+} influx into apical hair

bundles while stimulating lateral-line hair cells with a fluid-jet (Figure 5H-I') (Lukasz and

Kindt, 2018). Using this approach, we were able to stimulate and monitor stimulus-evoked

mechanosensitive Ca^{2+} responses in hair bundles within neuromast organs of each genotype

(Representative responses: Figure 5I-L). We quantified the number of hair bundles per

neuromast with mechanosensitive Ca^{2+} responses, and found that compared to controls,

significantly few hair bundles were mechanosensitive in *cib2* and *cib2*;*cib3* mutants (Figure 5-

figure supplement 1A, control: 92.16 ± 2.49; *cib2*: 49.88 ± 5.79, *cib2*;*cib3:* 18.99 ± 6.63, p >

0.0001). In contrast, in *cib3* mutants, the percent of mechanosensitive hair bundles per

neuromast was similar compared to controls (Figure 5-figure supplement 1A, *cib3*: 94.51 ±

1.82, p = 0.96). We also quantified the magnitude of the mechanosensitive Ca^{2+} signal in the

hair bundles, focusing on only the responsive hair cells for each genotype. This quantification

revealed that, while *cib3* mechanosensitive responses were not different compared to controls

(Figure 5-figure supplement 1B-C, GCaMP6s % ΔF/F control: 70.60 ± 5.87; *cib3*: 71.61 ±

5.23, p > 0.998), the magnitude of responses in *cib2* and *cib2*;*cib3* hair bundles was

significantly reduced (Figure 5-figure supplement 1B-C, GCaMP6s % ΔF/F control: 70.60 ±

5.87; *cib2*: 32.95 ± 5.77, *cib2*;*cib3:* 14.65 ± 1.85, p < 0.0001). Overall, our *in vivo* FM 1-43

and Ca^{2+}-imaging analyses revealed that Cib2 and Cib3 are both required to ensure normal

MET function in lateral-line hair cells. Furthermore, loss of Cib2 alone, but not Cib3 can impair

mechanosensory function in zebrafish lateral-line hair cells.

Cib2 is required in specific subsets of hair cells in the zebrafish lateral line and inner ear

Previous work has shown that within the zebrafish inner ear and lateral line there are distinct hair cell subtypes that rely on different complements of the mechanosensitive ion channel Tmc: Tmc1, 2a and 2b. In the primary posterior lateral line (pLL) hair cells are oriented to respond to either anterior or posterior fluid flow. The subset of hair cells that detects posterior flow relies on Tmc2b while the subset that detects anterior flow relies on both Tmc2b and Tmc2a (Figure 6A) (Chou et al., 2017). Further, in the zebrafish inner ear, work has shown that there are two morphologically distinct types of hair cells: short, teardrop-shaped cells with taller hair bundles and tall, gourd-shaped cells with shorter hair bundles (Smith et al., 2020; Zhu et al., 2021). Short cells rely on Tmc2a, while tall cells rely on either Tmc1 or Tmc2b for mechanosensitive function (Figure 6E). Based on this work, we hypothesized that Cib2 or Cib3 could have specific roles in zebrafish hair cell sensory organs linked to hair cell orientation in the lateral line or morphology type in the inner ear.

Our FM 1-43 labeling in neuromasts revealed no defect in *cib3* mutants, suggesting that Cib2 can compensate for Cib3 in all pLL hair cells. In contrast to *cib3* mutants, only a subset of hair cells per neuromast were labeled with FM 1-43 in *cib2* mutants (Figure 5E). Previous work has shown that in the pLL of *tmc2b* mutants, a subset of hair cells that respond to anterior flow label with FM 1-43 (Chou et al., 2017). This residual FM 1-43 label is dependent on Tmc2a. Therefore, we examined whether the remaining, FM 1-43 positive hair cells in *cib2* mutants have a specific hair cell orientation. For this analysis, we used a fixable form of FM 1-43, FM 1-43FX to label hair cells. After FM 1-43FX labeling, we fixed the larvae and labeled the apical hair bundles with phalloidin to reveal hair cell orientation. This analysis revealed that in sibling controls, a similar percentage of FM 1-43FX positive hair cells are oriented to respond to anterior and posterior flow (Figure 6B,B',D). In contrast, when we examined *cib2*

mutants, we found that significantly more of the FM 1-43 positive hair cells were oriented to

respond to posterior flow (Figure 6C-D). This indicates that in *cib2* mutants, the residual Cib3

present in pLL hair cells preferentially functions in hair cells that express primarily Tmc2b and

are oriented to respond to posterior flow (Figure 6A). On the other hand, Cib2 can function

with hair cells that express Tmc2b (respond to posterior flow) as well as hair cells that express

Tmc2b and Tmc2a (respond to anterior flow).

Next, we examined whether different hair-cell subtypes (short versus tall morphology)

in the zebrafish inner ear preferentially rely on Cib2 or Cib3 for mechanosensitive function.

To examine MET function, we injected FM 4-64 into the zebrafish ear in a transgenic

background that labels the hair cell membrane (memGCaMP6s). FM 4-64 is a redshifted

version of FM 1-43 and provides a similar readout of MET (Erickson et al., 2017; Smith et al.,

2020). After injection, we imaged FM 4-64 label in the medial cristae, an inner ear sensory

organ that allows for excellent delineation of short and tall cell types (Figure 6F-F'). We found

that similar to our FM 1-43 results in the pLL, no hair cells in the medial cristae were FM 4-

64 positive in *cib2;cib3* double mutants (Figure 6I-I'). In addition, similar to the pLL we

observed no defects in FM 4-64 labeling in the crista of *cib3* mutants compared to controls

(Figure 6G-G'). Interestingly we found that short hair cells in the crista of *cib2* mutants failed

to label with FM 4-64 (Figure 6H-H'). Previous work has shown that short hair cells rely on

Tmc2a for mechanosensitive function (Smith et al., 2020). Overall, our functional assessment

of MET based on hair cell subtypes within the zebrafish pLL and inner ear suggests that Cib2

is required for mechanosensitive function in hair cells that express Tmc2a.

CIB proteins simultaneously interact with at least two cytoplasmic domains of TMC1

To gain molecular insight into the role played by CIB proteins in hair-cell MET, we

generated AlphaFold 2 (AF2) (Jumper et al., 2021; Tunyasuvunakool et al., 2021) models of

the human TMC1 dimer, alone and in complex with human CIB2/3 (Figure 7A-B; Figure 7-

figure supplement 1; Figure 7-video 1). The AF2 model of TMC1 alone resembles previous

TMEM16/OSCA-based models of TMC1 and is also consistent with a recent cryo-EM

structure of the worm TMC-1 protein (Ballesteros et al., 2018; Jeong et al., 2022; Pan et al.,

2018; Walujkar et al., 2021). In these models, TMC1 has ten transmembrane domains (α1 to

α10; Figure 7-figure supplements 1-2) and a putative pore is formed by transmembrane helices

α4, α5, α6, and α7 at the periphery of each monomer (Pan et al., 2018; Walujkar et al., 2021),

away from the dimeric interface involving α10. Our new models feature an additional

amphipathic helix, which we denote α0, extending almost parallel to the expected plane of the

membrane bilayer without crossing towards the extracellular side (as observed for a mostly

hydrophobic α0 in OSCA channels) (Jojoa-Cruz et al., 2018; Liu et al., 2018; M. Zhang et al.,

2018). Extracellular and intracellular domains linking transmembrane helices, which were

either omitted or poorly predicted in previous work (Ballesteros et al., 2018; Pan et al., 2018;

Walujkar et al., 2021), have well-defined secondary structure and a high confidence score

(pLDDT > 70 for most regions; Figure 7-figure supplement 1A). The intracellular domain

linking helices α2 and α3, denoted here as IL1, adopts a helix-loop-helix with the two helices

running parallel to each other and differing in length. This is the same fold observed in its

crystal structure in complex with CIB3 (Figure 7-figure supplements 1-5) (Liang et al., 2021),

which validated the modeling approach.

Interestingly, AF2 models of TMC1 in complex with CIB2 suggest that the N-terminus

(NT) of TMC1 also interacts with CIB2 (Figure 7A-B). In our model, CIB2 is clamped between

TMC1-NT and TMC1-IL1 while the first helix of TMC1-IL1 runs parallel to the membrane

plane interacting with the base of α1 and the intracellular linkers between α4 and α5, α6 and

α7, and α8 and α9. Notably, the predicted interaction between TMC1-NT and CIB2 is

consistent with previous biochemical data (Giese et al., 2017), while the overall model resolves

seemingly contradictory results indicating that either the NT or IL1 of TMC1 were responsible

for interactions with CIB proteins (Giese et al., 2017; Liang et al., 2021). Additional AF2

models of TMC1 with CIB3 and of TMC2 with CIB2/3 (Figure 7-figure supplement 1)

predicted the same architecture for the TMC1/2 + CIB2/3 complexes where CIB proteins are

clamped between the IL1 and NT domains of TMCs. Predictions for fish complexes also show

the same general architecture (Figure 7-figure supplement 1A).

Nuclear magnetic resonance (NMR) experiments were used to validate the clamp

model for *Homo sapiens* (*hs*) TMC + CIB interactions. We produced uniformly ^{15}N labeled *hs*

CIB2 (22.7 kDa) and CIB3 (22.9 kDa) proteins ([U-^{15}N]-*hs* CIB2/3), alone or co-refolded with

unlabeled *hs* TMC1-IL1 (6.3 kDa). In parallel, we produced an unlabeled soluble TMC1-NT

fragment (6.9 kDa) and used these CIB and TMC1 samples to test protein-protein interactions

by monitoring their effects on 2D ^{1}H-^{15}N-correlation spectra of the labeled CIB2 and CIB3

proteins. All protein fragments were purified using size exclusion chromatography (SEC)

(Figure 7-figure supplement 6), revealing a single monodisperse peak for [U-^{15}N]-*hs* CIB2

(69.2 mL elution volume in S75 16/600 column) and a slightly displaced peak for [U-^{15}N]-*hs*

CIB2 + *hs* TMC1-IL1 (68.5 mL). In contrast, SEC experiments revealed two peaks for [U-

^{15}N]-*hs* CIB3 alone (potential dimer at 62.8 mL and potential monomer at 73.1 mL) and a

single peak for [U-^{15}N]-*hs* CIB3 + *hs* TMC1-IL1 (71.3 mL). ^{1}H-^{15}N-correlation spectra of [U-

^{15}N]-*hs* CIB2 alone were consistent with prior data indicating that many, but not all potential

peaks are resolved (Figure 7C, Figure 7-figure supplement 7A) (Huang et al., 2012). Spectra

for the [U-^{15}N]-*hs* CIB2 + *hs* TMC1-IL1 complex exhibit widespread perturbations, consistent

with extensive remodeling of the CIB2 structure. Moreover, addition of *hs* TMC1-NT induced

additional chemical shift perturbations in many, but not all peaks, demonstrating that the

TMC1-IL1 and TMC1-NT ligands bind simultaneously to [U-^{15}N]-*hs* CIB2 (Figure 7C, Figure

7-figure supplement 7B). We observed similar results when comparing spectra for [U-^{15}N]-*hs*

CIB3 alone, co-refolded with *hs* TMC1-IL1, and upon addition of *hs* TMC1-NT (Figure 7-

figure supplement 7C-D). A control experiment with the co-refolded mix of *hs* CIB3 + *hs*

TMC1-IL1 and [U-^{15}N]-*hs* CIB2 (Figure 7-figure supplement 7E) revealed no change in

chemical shifts indicating that direct heteromeric interactions between CIB2 and CIB3 are

unlikely. Thus, taken together, the NMR data strongly support a model in which human CIB2/3

proteins are clamped between TMC1-IL1 and TMC1-NT as suggested by AF2 and consistent

with prior biochemical data revealing interactions with either TMC1-NT or TMC1-IL1 (Giese

et al., 2017; Liang et al., 2021).

Simulations reveal CIB proteins stabilize TMC1 and TMC2

Our dimeric AF2 models of TMC1/2 alone and in complex with CIB2/3 are consistent

with previous experimental data that validated the location of the putative pore in similar

TMEM16/OSCA-based homology models of TMC1 (Pan et al., 2018). Our models are also

consistent with our NMR data supporting the clamp model for the TMC + CIB complex (Figure

7C). To improve the AF2 predictions and to gain mechanistic insights into TMC and CIB

function we turned to all-atom equilibrium molecular dynamics (MD) simulations (Karplus

and Petsko, 1990). We built ten different models that included dimeric TMC1 or TMC2 alone,

in complex with CIB2 or CIB3 with and without bound Ca^{2+}. These models were embedded in

either pure POPC or stereocilia-like mixed composition bilayers and solvated to generate a

total of twenty systems that were equilibrated for 100 ns each (Figure 7D, Figure 7-Table

supplement 1).

Equilibrium simulations of TMC1 and TMC2 alone revealed stable conformations for

the transmembrane domains of these proteins, but a highly mobile NT domain (Figure 7E,

Figure7-figure supplement 8). In simulations where CIB2 or CIB3 were present, with or

without bound Ca^{2+}, we observed more stable conformations of N-termini along with stable

salt bridges (Figure 7F). Buried surface area (BSA) for the TMC + CIB complexes (Figure 7-

figure supplement 9A,C,E, Table supplement 2) revealed a possible trend in which TMC1 and

TMC2 have more contacts with CIB3 than with CIB2. While additional sampling and longer

simulations might be needed for a robust prediction, these results suggest variable affinities for

TMC + CIB protein complexes that might be modulated by Ca^{2+} (Figure 7-figure supplement

9F).

422   Additional analysis of the putative pore in our simulations of TMC1 and TMC2

revealed hydration as observed in previous simulations of TMEM16/OSCA-based TMC1

models, as well as movement of lipids that lined up the predicted permeation pathway for

cations suggesting a membrane-based gating mechanism (Pan et al., 2018; Sukharev and

Corey, 2004; Walujkar et al., 2021). Longer, Anton 2, 960-ns long equilibrium trajectories of

TMC1 with CIB2 and bound Ca^{2+} in a mixed bilayer and TMC1 with CIB2 and (un)bound

Ca^{2+} in a POPC bilayer revealed how parts of the amphipathic α0 helices spontaneously

partitioned at the membrane interface with their hydrophobic sides pointing toward the lipid

bilayer core (Figure 7G). The N-terminal cytoplasmic domain of LHFPL5 has been shown to

interact with this TMC1 amphipathic helix, which might be essential for channel function in

mice (Qiu et al., 2023). The BSA for the dimeric TMC1 + CIB2 complex in these longer

simulations (Figure 7-figure supplement 9E) decreased as contacts between TMC1-α0 and

CIB2 were lost for one of the monomers in the mixed bilayer system. In the monomer in which

contacts were maintained, a single cholesterol was situated in the middle of the putative pore.

Similarly, TMC1-α0 and CIB2 contacts were lost for both monomers in the POPC bilayer

systems. Regardless, in all Anton 2 equilibrium trajectories contacts between CIB2 and the

TMC1-IL1 and -NT remained, thus confirming the stability of the complex over a microsecond

timescale.

TMC1 and TMC2 complexes with CIB2 and CIB3 are predicted to be cation channels

To further explore the properties of our models of TMC/CIB complexes we took

equilibrated TMC1 systems and applied a -0.5 V transmembrane voltage while monitoring ion

permeation through the putative pore during 100-ns long trajectories (Figure 7-figure

supplements 10-11, Figure 7-Table supplement 2, Figure 7-video 2). In systems with pure

POPC bilayers we observed permeation of K^+ in either one or both pores of the TMC1 dimer,

with or without CIB2 or CIB3 and with or without bound Ca^{2+}. Similar results were obtained

for four TMC1 systems simulated at -0.25 V. Variability of the conductance state was attributed

to partial blockage by lipids and poor sampling, but in all cases in which a clear conductive

state was monitored we observed almost exclusive passage of K^+ with conductance values that

reached up to ~138 pS for a single pore. While simulations tend to overestimate bulk KCl

conductivity and in some cases predictions need to be scaled down (by a factor of ~0.6)

(Walujkar et al., 2021), the conductance values monitored here are consistent with previous

simulations and with the expected conductance of the inner-ear transduction channel (Beurg et

al., 2021; Walujkar et al., 2021). In contrast, we observed smaller conductance values for

TMC1 systems with mixed lipid bilayers (at most ~22 pS for a single pore), regardless of the

presence of CIB2 or CIB3, with or without bound Ca^{2+} (Figure 7-figure supplement 11B).

Additional Anton 2 simulations for one of these TMC1 + CIB2 + Ca^{2+} systems with a mixed

membrane, equilibrated for 240 ns and then subjected to -0.43 V for 480 ns, revealed a low

conductance value of ~6.2 pS. The reduced conductance values observed for mixed-membrane

systems were attributed in part to pore blockage by cholesterol and lipids in the initial setup.

Interestingly, Anton 2 simulations of TMC1, TMC1 + CIB2, and TMC1 + CIB2 + Ca^{2+} in pure

POPC membranes subjected to -0.5 V, -0.25 V, and -0.125 V for 480 ns, 480 ns, and 960 ns,

respectively, revealed a trend in which TMC1 exhibits higher conductance with CIB2 bound

(Figure 7H, Table supplement 2). This may begin to explain why CIB2 is essential for normal

TMC1 function.

Simulations of TMC2 with an applied transmembrane voltage of -0.5 V revealed a more

closed state when compared to our TMC1 model, regardless of membrane composition and

presence or absence of CIB2, CIB3 and bound Ca^{2+} (Figure 7-figure supplement 11).

Conduction of K^+ was observed in most cases, but the maximum conductance did not reach

above ~42 pS, suggesting that the conformation simulated is less open than the ones explored

in simulations of TMC1. Regardless, our simulations unequivocally indicate that TMC1 and

TMC2 are cation channels and that these proteins can form stable complexes with CIB2 and

CIB3.

Discussion

Previous work on CIB proteins had primarily focused on mammalian auditory hair

cells, leaving open the question of whether they play a functional role in vestibular function

and across vertebrate species. Recently, Wang et al. have suggested that CIB2 and CIB3 are

important for stereocilia maintenance and MET in mouse vestibular hair cells (Wang et al.,

2023). This work demonstrated that similar to *Tmc1* and *Tmc2*, *Cib2* is highly expressed in the

striolar region while *Cib3* is expressed in the extrastriolar regions, and mice lacking both *Cib2*

and *Cib3* had no MET and deficits in vestibular hair cell bundle structure (Wang et al., 2023).

Taking a step forward, here, we now establish that CIB2 and CIB3 are highly conserved

regulators of MET in both mouse vestibular hair cells and zebrafish hair cells. We demonstrate

that vestibular hair cells in mice and zebrafish lacking CIB2 and CIB3 appear structurally

normal but have no detectable MET, assessed via FM 1-43 dye uptake, at time points when

MET function is well developed in wild-type hair cells. Similar structural-function phenotype

has been reported in auditory hair cells in *Tmc1/Tmc2* double mutant mice (Kawashima et al.,

2015). Besides the known *in vitro* interactions (Giese et al., 2017; Liang et al., 2021), our study

provides strong genetic evidence for the direct functional relationship between CIB and TMC

proteins in regulating MET both in auditory and vestibular hair cells.

Previous studies have shown that the loss of CIB2 leads to profound hearing loss from

birth in both human and mice (Giese et al., 2017; Liang et al., 2021; Seco et al., 2016). This

suggests that the endogenous levels of CIB3 present in auditory hair cells are not sufficient to

compensate for CIB2 in hearing. Prior work has also shown that over-expression of CIB3 in

mouse auditory outer hair cells can only partially compensate for the loss of CIB2 in MET

(Liang et al., 2021). However, whether overexpression of CIB3 can rescue impaired hearing

thresholds in *Cib2* mutant mice is not known. While CIB2 is also present in vestibular hair

cells, compared to auditory hair cells, the expression of CIB3 is several folds higher in

vestibular end organs. Indeed, our current data demonstrate that, unlike auditory hair cells,

endogenous levels of CIB3 can fully compensate for loss of CIB2 in vestibular hair cells in

mice. Together our work on the vestibular system in mice highlights that both CIB2 and CIB3

can play overlapping roles in MET function.

Interestingly, our corresponding work in zebrafish indicates that Cib3 can also

compensate for Cib2, but only in specific subsets of hair cells. Recent genetic, morphology and

scRNAseq data have revealed that within the zebrafish inner ear, each sensory epithelia

(utricle, saccule, cristae) have two distinct types of hair cells (Shi et al., 2023; Smith et al.,

2020). In these epithelia, short cells are present in the striolar or central region, while tall cells

are present in the extrastriolar or peripheral region (Figure 6A,E). Genetic studies have

revealed that these two types of hair cells rely on different Tmc combinations for MET–short

cells rely on Tmc2a while tall cells rely on Tmc2b and Tmc1 (Smith et al., 2020). Recent

scRNAseq work has validated this *Tmc* requirement and shown that *cib2* is expressed in both

short and tall cells, while *cib3* is specifically expressed in just tall cells (Shi et al., 2023). Our

work on MET in the zebrafish inner ear provides functional support for this *cib* RNAseq data,

as FM 4-64 labeling is absent from short hair cells when Cib2 is lost (Figure 6H). In contrast,

loss of Cib3 alone does not impact FM 4-64 labeling in the zebrafish inner ear, indicating that

Cib2 can compensate for loss of Cib3 in both short and tall cells (Figure 6G). Currently it is

not clear what role these two distinct types of hair cells play in zebrafish hearing and balance.

Behavioral work on zebrafish *tmc2a* mutants has shown that when MET is absent from just

short hair cells, the acoustic startle response is severely diminished (Smith et al., 2020). This

is consistent with our present findings showing that zebrafish *cib2* mutants with impaired MET

in short hair cells have a reduced acoustic startle response (Figure 5A). Together these

behavioral studies suggest that short hair cells play an important role in detecting acoustic

stimuli necessary to generate a robust escape behavior.

In addition to the inner ear, we likewise found that both Cib2 and Cib3 play an

important role in the zebrafish lateral line. Similar to the zebrafish inner ear and mouse

vestibular system, we show that in lateral-line hair cells, Cib2 can compensate for MET in all

hair cells when Cib3 is lost (Figure 5B,C,F,K, Figure 5-figure supplement 1A-C). However,

like in the zebrafish inner ear, when Cib2 is lost, MET is lost from a specific population of

lateral-line hair cells. In the lateral line, each neuromast has two cell populations that detect

flow in opposing directions (López-Schier and Hudspeth, 2006; Pujol-Martí and López-Schier,

2013). Work in the posterior lateral line has demonstrated that cells that detect anterior flow

rely on Tmc2a/2b for MET, while cells that detect posterior flow rely on just Tmc2b (Chou et

al., 2017). In our work we find that cells that detect anterior flow primarily rely on Cib2 for

MET (Figure 6C-D). Further, this work indicates that Cib3 cannot compensate for loss of Cib2

in these cells. Interestingly, our functional assessment in the zebrafish lateral line and inner ear

suggests that Cib2 is specifically required for mechanosensitive function in populations of hair

cells that express Tmc2a. Whether Cib3 is unable to pair with Tmc2a or is simply not expressed

along with Tmc2a remains an unanswered question. The significance of this pairing is also

unclear. Recent work has shown that the addition of Tmc2a to lateral line cells that detect

anterior flow significantly augments the mechanosensitive responses of these cells (Kindig et

al., 2023). Whether Cib2 is required along with Tmc2a for these augmented responses remains

unclear.

Our *in vivo* studies of mouse TMC + CIB and zebrafish Tmc + Cib complexes suggest

a conserved functional relationship between these two protein families across vertebrates.

Structural predictions using AF2 show conserved folds for human and zebrafish proteins, as

well as conserved architecture for their protein complexes. Predictions are consistent with

previous experimentally validated models for the TMC1 pore (Ballesteros et al., 2018; Pan et

al., 2018), with the structure of human CIB3 coupled to mouse TMC1-IL1 (Liang et al., 2021),

with the architecture of the nematode TMC-1 and CALM-1 complex (Jeong et al., 2022), and

with our NMR data validating the interaction between human TMC1 and CIB2/3 proteins.

Taken together, the modeling predictions, NMR experiments, and simulations

presented here provide a structural framework to interpret and understand seemingly

contradictory results. In a previous study (Giese et al., 2017), we showed that the N-termini of

TMC1 and TMC2 interact with CIB2, and subsequent work (Liang et al., 2021) presented

biochemical data validating the interaction between the TMC1-NT with CIB2. However, X-

ray crystallography data demonstrated that the main interaction between TMC1 and CIB3

involved IL1, leaving the role played by CIB2-TMC-NT interactions unaddressed (Liang et

al., 2021). Here we show using NMR that both TMC1-NT and TMC1-IL1 bind to CIB2 and

CIB3 non-competitively. These interactions are consistent with AF2 models for TMC + CIB

complexes where CIB proteins are clamped between TMC1-NT and IL1. Simulations of these

models indicate that these interactions are stable and last for microseconds, with some potential

preferential binding of TMC1 and TMC2 to CIB3 over CIB2 predicted from BSA calculations.

Interestingly, the behavior of CIB2 and CIB3 in solution (SEC experiments) is different

in the absence of TMC1-IL1. While CIB2 appears as a single peak, CIB3 shows two clear

peaks indicating the presence of monomeric and dimeric states, consistent with the crystal

structure of an apparent CIB3 dimer (Liang et al., 2021). In the presence of TMC1-IL1, the

CIB3 dimeric peak is absent, and the potentially monomeric peak is slightly shifted accounting

for the added TMC1-IL1, as observed for CIB2. This result suggests that CIB2, both in the

presence and in the absence of TMC1-IL1, is monomeric in solution (Figure 7-figure

supplement 6). This proposal is consistent with prior conclusions that CIB2 is a monomer (Dal

Cortivo et al., 2019), but is in disagreement with other reports indicating that CIB2 is a dimer

in solution (Vallone et al., 2018) and our own prior pull-down assays indicating that CIB2 co-

immunoprecipitated with differentially tagged CIB2 or CIB3 (Giese et al., 2017; Riazuddin et

al., 2012). Moreover, our NMR data indicates that TMC1-IL1 + CIB2 is unlikely to directly

interact with CIB3. The simplest explanation for these seemingly contradictory results comes

from the TMC1 and TMC2 dimeric states (Pan et al., 2018) and our modeling indicating that

each TMC monomer interacts with one CIB protein (Figure 7-figure supplement 12). Pull-

down assays might simply be reflecting the oligomeric state of TMC proteins. In such scenario,

and considering that TMC proteins have been suggested to form hetero-oligomers (Pan et al.,

2018), there are up to ten different combinations of TMC1/2 and CIB2/3 proteins (Figure 7-

figure supplement 12) that might have distinct properties. Such variety of complexes with

potentially different conductance values might explain tonotopic changes of conductance along

the cochlea, although other explanations exist, including a variable number of TMC proteins

around the tip link (Beurg et al., 2018) and the direct influence of membrane composition on

conductance (Effertz et al., 2017; Walujkar et al., 2021).

Our simulations in the presence of a biasing potential unequivocally demonstrate that

both TMC1 and TMC2, in the presence or the absence of CIB proteins, with or without bound

Ca^{2+}, and in two types of membranes, are cation channels with predicted single-channel

conductance values that can reach values that are consistent with those of the transduction

channel. Our models emphasize again the role played by the membrane in lining the pore of

TMC proteins (Pan et al., 2018; Walujkar et al., 2021), and also indicate that CIB proteins are

ideally positioned to connect various parts of cytosolic inter-helix linkers with the helices that

form the pore, specially α3 and α4, recently implicated in gating (Akyuz et al., 2022; Walujkar

et al., 2021). Last, our modeling also positions the myristoylated N-terminus of CIB2 and CIB3

close enough to the membrane bilayer, which might be yet another way to sense membrane

stretch and influence gating.

Overall, the biochemically supported atomic-resolution structural models and

simulations presented here along with our results obtained using animal models strongly

suggest that TMC1/2 proteins must function with CIB2/3 in normal hair-cell MET across

sensory organs, hair-cell types, and species. This is supported by structural data indicating

multiple contact points between TMC and CIB proteins, simulation data predicting that CIB

proteins stabilize TMC cytosolic domains and may enhance cation conduction, and *in vitro* and

in vivo functional data indicating that both TMC1/2 and CIB2/3 proteins are needed for hair-

cell MET.

Materials and methods

<u>**Key resources table**</u>

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 568	Thermo Fisher Scientific	Cat#A-11031; RRID: AB_144696
Goat anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor 488	Thermo Fisher Scientific	A32723
Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Thermo Fisher Scientific	Cat#A-11034; RRID: AB_2576217
Goat anti-rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 568	Thermo Fisher Scientific	A-11011
Mouse monoclonal anti-FLAG M2	Sigma-Aldrich	Cat#F3165; RRID: AB_259529
Mouse monoclonal anti-FLAG M2-HRP	Thermo Fisher Scientific	Cat#A8592; RRID: AB_439702
Mouse monoclonal anti-V5	Thermo Fisher Scientific	Cat# R960-25; RRID: AB_2556564
Mouse monoclonal anti-V5-HRP	Thermo Fisher Scientific	Cat#R961-25; RRID: AB_2556565
Mouse monoclonal GAPDH	Santa Cruz	sc-32233
Rabbit polyclonal anti-CIB3	Sigma	SAB2103524
Rabbit polyclonal anti-GFP	Invitrogen	A-11122
Rabbit polyclonal anti-GFP-HRP	Thermo Fisher Scientific	Cat#A10260; RRID: AB_2534022
Rabbit polyclonal anti-Myosin VIIa	Proteus Biosciences	#25-6790
Rabbit polyclonal anti-V5	Abcam	Cat#ab9116; RRID: AB_307024
Chemicals		
FM™ 1-43FX	Thermo Fisher Scientific	F35355
FM™ 1-43	Thermo Fisher Scientific	T3163
FM™ 4-64	Thermo Fisher Scientific	T13320
Alexa Fluor™ 488 Phalloidin	Thermo Fisher Scientific	A12379
Alexa Fluor™ 633 Phalloidin	Thermo Fisher Scientific	A22284
16% Paraformaldehyde	Electron Microscopy Sciences	50-980-487

Triton X-100	Polysciences	04605-250
DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride)	Invitrogen	D1306
Protease inhibitor mixture	Roche	11697498001
Protein A–Sepharose CL-4	Thermo Fisher Scientific	101042
Polyethylenimine	Polysciences	23966
Critical Commercial Assays		
4–20% Tris-Glycine gel	Novex	XV04205PK20
SMARTScribe Reverse Transcriptase kit	Clontech	639537
SYBRgreen technology	Qiagen	330503
Amersham™ ECL™ Prime Western Blotting Detection Reagent	GE Healthcare	RPN2236
QIAshredder	Qiagen	Cat#79656
Trans-Blot® Turbo™ Transfer System	Bio-Rad	Cat#1704156
Trans-Blot® Turbo™ Midi PVDF Transfer Packs	Bio-Rad	Cat#1704157
4–20% Mini-PROTEAN® TGX Stain-Free™ Protein Gels	Bio-Rad	Cat#4568094
4–20% Criterion™ TGX Stain-Free™ Protein Gel	Bio-Rad	Cat#5678094
GFP-Trap Magnetic Agarose	Chromotek	Cat#gtma-20
Anti-v5 agarose affinity gel	Sigma-Aldrich	Cat#A7345
Anti-FLAG® M2 Magnetic Beads	Sigma-Aldrich	Cat#M8823
Nunc™ Lab-Tek™ chambered coverglass	Thermo Fisher Scientific	Cat#155411PK
Lipofectamine™ 3000 Transfection Reagent	Thermo Fisher Scientific	Cat#L3000015
Experimental Models: Organisms/Strains		
$Cib2^{tm1a}$	EUCOMM	
$Cib3^{KO}$	This study	N/A
$cib2^{idc21}$ zebrafish	This study	N/A
$cib3^{idc22}$ zebrafish	This study	N/A
Tg(myo6b:GCaMP6s-CAAX)^{idc1} zebrafish	(Sheets et al., 2017)	ZDB-ALT-170113-3

Mouse strains and husbandry

$Cib2^{KO}$ mice: $Cib2^{\ tm1a(EUCOMM)Wtsi}$ mice were obtained from the EUCOMM repository and

maintained on C57BL/6 N background. A gene trap cassette containing lacZ and neomycin

resistance genes flanked by FRT sites was inserted downstream of exon 3, with exon 4 flanked

by loxP sites.

Cib3^{KO} mice: Cib3 knockout mice were generated using CRISPR/Cas9 technology. An

indel in exon 4 of *Cib3* was introduced leading to a nonsense mutation and a premature stop at

the protein level (p.Asp54*). Primers used to genotype Cib3 mutant mice were F:

GTAGGGGCATGGATAACATCTG and R: GTTGAATGCTTGTCTCCCCAGG. *Cib3^{KO}*

mice were backcrossed using C57Bl6/JN *Cib2^{KO}* mice for several generations. *Cib3* allele was

validated at the genomic, RNA messenger, and protein level.

Animals were group-housed with their littermates on a 12:12 h light: dark cycle at 20°C

with ad libitum access to food and water. All animal procedures were approved by the

Institutional Animal Care and Use Committees (IACUCs) at University of Maryland (protocol

#0420002) and at Harvard Medical School (protocol #00001240).

Western blot and qPCR in mice

Mouse heart tissues were homogenized in CHAPS buffer containing 1X protease inhibitors,

1 mM Na orthovanadate, 10 mM Na glycerophosphate, and 10 mM NaF. Equivalent protein

concentrations were fractionated on a 4–20% Bis–Tris gel (Invitrogen) and transferred to

polyvinylidene fluoride (PVDF) membrane (Millipore). Membranes were blocked and then

probed with the rabbit anti-CIB2 (1:500), CIB3 (1:500), and GAPDH (1:500) overnight at 4°C,

followed by three washes of 30 min each in 1X TBST. After washing, blots were incubated

with horseradish peroxidase-conjugated anti-rabbit antibody (1:1000, Sigma NA934V, Lot #

17640116) for 2 h at room temperature, followed by detection using the ECL Prime Western

Blotting System (Thermo Fisher Scientific 32,106).

RNAs from brain were extracted from *Cib2;Cib3* mutant and control mice using the

Ribopure kit (Ambion). SMARTScribe Reverse Transcriptase kit (Clontech) was used to

generate cDNAs, and SYBRgreen technology (Qiagen) was used to perform the qPCR. The

following primers were used to amplify *Cib2* and *Cib3*: Cib2_exon4_F:

CTCTGTGCTCTGCGAATCAG, Cib2_exon5_R: GGCCAGCGTCATCTCTAAGT,

Cib3_exon3_F: TGGTGCCTCTTGACTACACG, Cib3_exon5_R:

CAGCACCTTCTCACAGACCA. *hprt* amplification was used to normalize the samples:

hprt_exon8_F: TGTTGTTGGATATGCCCTTG, hprt_exon9_R:

GGCTTTGTATTTGGCTTTTCC.

Quantification of VOR and OKR in mice

Surgical techniques and experimental setup have been previously described (Beraneck and

Cullen, 2007). Eye movement data were collected using an infrared video system (ETL-200,

ISCAN system). The rotational velocity of the turntable (head velocity) was measured using a

MEMS sensor (MPU-9250, SparkFun Electronics). Eye movements during OKR were evoked

by sinusoidal rotations of a visual surround (vertical black and white stripes, visual angle width

of 5°) placed around the turntable at frequencies 0.2, 0.4, 0.8, 1, and 2 Hz with peak velocities

of ±16∘/s. To record VOR responses, the turntable was rotated at sinusoidal frequencies 0.2,

0.4, 0.8, 1, and 2 Hz with peak velocities of ±16∘/s in both light and dark. In the light condition,

the visual surround remained stationary, whereas in the dark condition, both the visual surround

and turntable rotated in phase. Head and eye movement signals were low-passed filtered at 125

Hz and sampled at 1 kHz. Eye position data were differentiated to obtain velocity traces. Cycles

of data with quick phases were excluded from the analysis. Least-square optimization

determined the VOR and OKN gains and phases (Beraneck et al., 2008) plotted as mean ±

standard deviation (SD) against all frequencies for all mice. $Cib2^{+/+}; Cib3^{KO/KO}$ ($n = 4$),

$Cib2^{KO/+}; Cib3^{KO/KO}$ ($n = 5$), and $Cib2^{KO/KO}; Cib3^{KO/KO}$ ($n = 4$) mice were tested for all

frequencies. Two-way ANOVAs and Bonferroni post hoc tests were used to test statistical

significance.

Mouse rotarod test

Cib2$^{+/+}$;Cib3$^{KO/KO}$ (n = 6), *Cib2$^{KO/+}$;Cib3$^{KO/KO}$* (n = 6), and *Cib2$^{KO/KO}$;Cib3$^{KO/KO}$* (n = 6) mice

were used. Motorized rotating rod (Rotarod, IITC Life Science) was programmed to accelerate

from 4 rpm to 40 rpm with a ramp of 300 s for each test trial. The time taken for a mouse to

fall off the rotarod was recorded. Three trials were performed each day for 5 consecutive days.

First day (day 0) was considered the training day.

Quantification of head motion in mice

Head motion was recorded in mice via a miniature motion sensor (MPU-9250, TDK

Corporation) attached to the top of the skull. The motion sensor recorded 3-axis linear

acceleration (i.e., fore-aft, lateral, and vertical) and 3-axis angular velocity (roll, pitch, and

yaw) at 100 Hz sampling rate. The power spectra of the motion signals in all six dimensions

were computed using Welch's averaged periodogram with nfft = 512 and a Bartlett window

(pwelch function, MATLAB, MathWorks).

Circling behavior in mice

Mice were placed in a cylindrical-shape container (27 cm in diameter, 30 cm in height) and

their behavior were recorded using a cell-phone camera. The number of rotations made during

120-second recording were counted offline by watching the recorded videos. A rotation is

defined as a completed 360-degree rotation of the mouse's heading.

Immunostaining in mice

The cochlear and vestibular sensory epithelia were isolated, fine dissected and permeabilized

in 0.25% Triton X-100 for 1 h and blocked with 10% normal goat serum in PBS for 1 h. The

tissue samples were probed with primary antibody overnight and after three washes were

incubated with the secondary antibody for 45 min at room temperature. Rhodamine phalloidin

or Alexa fluor phalloidin 488 were used at a 1:250 dilution for F-actin labeling. Nuclei were

stained with DAPI (Molecular Probes). Images were acquired using either LSM 700 laser

scanning confocal microscope (Zeiss, Germany) with a 63× 1.4 NA or 100× 1.4 NA oil

immersion objectives or Leica SP8 laser scanning confocal microscope with a 100× 1.44 NA

objective lens. Stacks of confocal images were acquired with a Z step of 0.05-0.5 μm and

processed using ImageJ software (National Institutes of Health). Experiments were repeated at

least 3 times, using at least three different animals.

FM 1-43 dye uptake experiments in mice

In the Ahmed lab, cochlear explants from *Cib2* and *Cib3* mutant mice were dissected at

postnatal day 5 (P5) and cultured in a glass-bottom petri dish (MatTek, Ashland, MA). They

were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% FBS

(Life Technologies) for 2 days at 37 °C and 5% CO_2. Explants were incubated for 10 s with

3 μM FM 1-43, washed three times with Hank's balanced salt solution, and imaged live using

a Zeiss LSM 700 scanning confocal microscope.

In the Holt Lab, P60 mice were injected intraperitoneally with 5 mg/g mouse weight of

FM 1-43FX fixable form as previously described (Meyers et al., 2003). Tissue was collected

after ~24 h and fixed for 1 h at 25 °C in 4%PFA/1X PBS protected from light. Tissue was then

transferred into 120 mM EDTA pH 7.4 for 48-72 h protected from light. Then tissue was

transferred to 1XPBS, dissected, incubated in 1:1000 dilution of Phalloidin (Invitrogen),

washed three times in 1XPBS and mounted under glass coverslips for microscopy. Imaging

was completed using a 20× objective on a Zeiss 800 confocal microscope with laser intensity,

detector gain, and post processing set equally for each sample.

ABR measurements in mice

Hearing thresholds of *Cib2* and *Cib3* mutant mice ($n = 5$–10 each genotype) were evaluated by

recording ABRs. All ABR recordings, including broadband clicks and tone-burst stimuli at

three frequencies (8, 16, and 32 kHz), were performed using an auditory-evoked potential RZ6-

based auditory workstation (Tucker-Davis Technologies) with high frequency transducer

RA4PA Medusa PreAmps. Maximum sound intensity tested was 100 dB SPL. TDT system III

hardware and BioSigRZ software (Tucker Davis Technology) were used for stimulus

presentation and response averaging.

Zebrafish strains and husbandry

Zebrafish (*Danio* rerio) were grown at 28°C using standard methods. Zebrafish were

maintained in a Tubingen or TAB background. Larvae were raised in E3 embryo medium

(5 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, and 0.33 mM $MgSO_4$, buffered in HEPES, pH

7.2). Zebrafish work at the National Institute of Health (NIH) was approved by the Animal Use

Committee at the NIH under animal study protocol #1362-13. All experiments were performed

on larvae at 5-6 days post fertilization (dpf). At this age sex determination is not possible.

Neuromasts (L1-L5) in the primary posterior lateral line were used for our analyses. For Ca^{2+}

imaging the following transgenic line was used: *Tg(myo6b:memGCaMP6s)[idc1]* (Jiang et al.,

2017).

Cib2[idc21] and *cib3[idc22]* mutants were generated in-house using CRISPR-Cas9 technology

as previously described (Varshney et al., 2016). Exon 3 and exon 6 were targeted for *cib2* and

cib3 respectively. Guides for *cib2* and *cib3* are as follows: 5'-

GGAGGAGCTTACACCAG(AGG)-3' and 5'-GGAAATCCTCCAGAGAAAGG(CGG)-3'.

Founder fish were identified using fragment analysis of fluorescent PCR products (Varshney

et al., 2016). Founder fish containing a 10 bp insertion in *cib2* 5'-

GCTTACACCA(<u>CCTTTGGTAG</u>)GAGGAGGTTA-3' and a 4 bp deletion in *cib3* 5'-

TGGCCGCCT(TTCT)CTGGAGG-3' were propagated and selected for analysis. After

behavior or imaging analyses, genotypes were confirmed using standard PCR and sequencing.

The following primers were used for genotyping: *cib2*_FWD 5'-

ACTGAAGCAACTCCTTTGTGG-3' and *cib2*_REV 5'-GAACCATGCATTTTTACAGC-

3' and *cib3*_FWD 5'-ATAGTCTCTCCATCTATAGACT-3' and *cib3*_REV 5'-

TTTTAGGGTGAAATAAAACAGA-3'. Crosses of *cib2$^{+/-}$;cib3$^{-/-}$* X *cib2$^{+/-}$;cib3$^{+/-}$* or *cib2$^{-/-}$*

;cib3$^{+/-}$ X *cib2$^{+/-}$;cib3$^{+/-}$* adults were used to propagate larvae for our analyses. All controls or

siblings are double heterozygous animals. Larvae were chosen at random, except for *cib2;cib3*

homozygous mutants which, similar to other zebrafish mutants with hearing and balance

defects, had no swim bladder and no acoustic startle response.

<u>Labeling zebrafish hair cells with FM 1-43, FM 4-64, and FM 1-43FX</u>

To label lateral-line hair cells with FM 1-43, larvae were immersed in 1 µM FM 1-43

(ThermoFisher, T3163) in E3 for 30 s, and then washed 3 times in E3. After washing, larvae

were mounted in 1% low melt agarose containing an anesthetic, 0.02% tricaine

methanesulfonate (Syndel, ANADA 200-226). FM 1-43 labeled hair cells were then imaged

on an A1 Nikon upright laser-scanning confocal microscope with a 60× 1.0 NA water

objective. FIJI was used to process image Z-stacks. To estimate the number of hair cells per

neuromast, laser scanning DIC images were used to count the number of hair bundles per

neuromast. To quantify FM 1-43 label, a 5 µm circular region of interest (ROI) was placed on

the soma of each hair cell with a hair bundle in the DIC image. These ROIs were used to

quantify the mean FM 1-43 intensity per hair cell. These intensity values were averaged to

calculate the average FM 1-43 intensity per neuromast.

FM 4-64 (ThermoFisher, T13320) was used to label hair cells in the crista within the

zebrafish inner ear. For this labeling, *Tg(myo6b:memGCaMP6s)*[idc1] larvae were mounted on

their side on Sylgard chamber in E3 embryo media containing 0.02% MESAB. Approximately

2-3 nL of 10 μM FM 4-64 in 0.1 M KCl were injected into the otic capsule as previously

described (Smith et al., 2020). FM 4-64 labeled hair cells of the medial crista were then imaged

on an A1 Nikon upright laser-scanning confocal microscope with a 25× 1.10 NA or 60× 1.0

NA water objective. FIJI was used to process image Z-stacks. To assess tall versus short hair

cells in each crista *memGCaMP6s* label was used.

To correlate hair bundle orientation in lateral line hair cells with intact

mechanotransduction, larvae were immersed in 3 μM FM 1-43FX (ThermoFisher, F25255) in

E3 for 30 s, and then washed 3 times in E3. After washing, larvae were fixed with 4%

paraformaldehyde in PBS at 4°C overnight. After fixation, larvae were washed 4 × 5 min in

0.1% Tween in PBS (PBST). Larvae were then placed in blocking solution (2% goat serum,

1% bovine serum albumin, 2% fish skin gelatin in PBS + 0.3% triton) for 2 h at room

temperature. After block Alexa Fluor 633 Phalloidin (ThermoFisher, A22284) was added at

1:100 in PBST. Larvae were incubated for 4 h at room temperature and then washed 4 × 5 min

in PBST. Larvae were mounted on glass slides with Prolong Gold (ThermoFisher Scientific)

using No. 1.5 coverslips. To image hair bundle orientation and FM 1-43FX label in lateral-line

hair cells, a Zeiss LSM780 confocal microscope with Airyscan was used. Hair cells were

imaged using a 63× 1.3 NA oil objective and processed with an Airyscan processing factor of

7.0. For analysis, Z-stacks were max-projected in FIJI. Images were background subtracted

using a rolling ball correction. Hair cells with a FM 1-43FX intensity more than twice the

background intensity were scored as FM 1-43FX positive.

Zebrafish Ca^{2+} imaging and behavior

For imaging of mechanosensitive Ca^{2+} signals in neuromast hair bundles*,*

Tg(myo6b:memGCaMP6s) animals were mounted on their side on Sylgard chamber in E3

embryo media containing 0.2% MESAB (Lukasz and Kindt, 2018). Larvae were then pinned

and a solution containing α-bungarotoxin (125 μM, Tocris) was injected into the heart cavity

for paralysis during imaging. After paralysis, larvae were immersed in extracellular imaging

solution (in mM: 140 NaCl, 2 KCl, 2 CaCl$_2$, 1 MgCl$_2$ and 10 HEPES, pH 7.3, OSM 310 +/-

10). To image Ca^{2+} signals in the hair bundles we used a Bruker Swept-Field confocal system

equipped with a Rolera EM-C2 CCD camera (QImaging, Surrey, Canada) and a Nikon CFI

Fluor 60×1.0 NA water immersion objective as previously described (Q. Zhang et al., 2018).

The system includes a band-pass 488/561 nm filter set (59904-ET, Chroma) and is controlled

using Prairie View software (Bruker Corporation). A piezoelectric motor (PICMA P-882.11-

888.11 series, PI Instruments) attached to the objective and used to acquire 5-plane Z-stacks

every 0.5 μm using a 35 μm slit at a 50-Hz frame rate for a 10-Hz volume rate. A fluid-jet

composed of a glass capillary attached to a pressure clamp system (HSPC-2-SB, ALA) was

used to deliver a 500-ms anterior and posterior step stimulation to deflect hair bundles along

their axis of sensitivity.

To quantify GCaMP6s fluorescent intensity changes during stimulation, we registered

the Z-stacks and then average projected the Z-stacks into a single plane. We then loaded the

projected images into FIJI and used the Time Series Analyzer V3 plugin to create circular ROIs

with a ~1.7 μm diameter. ROIs were placed on the center of each individual bundle. We then

measured and plotted change in the mean intensity ($\Delta F/F_0$) within the region during the

recording period. The mean intensity within each ROI was computed for each hair bundle. A

hair bundle with a signal magnitude (peak value of intensity change upon stimulation) above

10% $\Delta F/F_0$ was considered mechanosensitive. To create the AVG hair bundle GCaMP6s

($\Delta F/F_0$) per neuromast plot in Figure 5-figure supplement 1C, only traces from

mechanosensitive cells were averaged.

A Zantiks MWP behavioral system was used to examine startle responses. The Zantiks

system tracked and monitored behavioral responses via a built-in infrared camera at 30 frames

per second. A 12-well plate was used to house larvae during behavioral analysis. Each well

was filled with E3 and 1 larva. All fish were acclimated in the plate within the Zantiks chamber

in dark for 15 min before each test. To induce startle, an integrated stepper motor was used to

drive a vibration-induced startle response. A vibrational stimulus that triggered a maximal %

of animals startling in controls without any tracking artifacts (due to the vibration) was used

for our analyses. Each larva was presented with the vibrational stimulus 5 times with 100 s

between trials. During the tracking and stimulation, a Cisco router connected to the Zantiks

system was used to relay x, y coordinates of each larva every frame. To qualify as a startle

response, a distance above 4 pixels or ~1.9 mm was required within 2 frames after stimulus

onset. Animals were excluded from our analysis if no tracking data was recorded for the animal.

<u>Zebrafish Statistics</u>

All data shown are mean ± standard error of the mean (SEM). All experiments were performed

on at least 4 animals and 8 neuromasts and 4 cristae. Behavior analysis and Ca^{2+} imaging was

repeated at least twice on two independent days. All statistical analysis was performed using

GraphPad Prism 8.0. A D'Agostino and Pearson test was used to test data for normality. A

one-way ANOVA with a Dunnett's correction for multiple comparisons was used to compare

differences between siblings and controls regarding the % of mechanosensitive hair cells,

number of hair cells, % of FM 1-43 positive hair cells, average FM 1-43 label. For proportion

of animals startling and comparisons for FM 1-43FX label, a Kruskal-Wallis test with a

Dunnett's correction for multiple comparisons was used.

Expression and purification of bacterially expressed *hs* CIB2, CIB3, TMC1-IL1, and

TMC1-NT

DNA sequences encoding for *hs* CIB2, CIB3, TMC1-IL1, and TMC1-NT were subcloned into

NdeI and *XhoI* sites of the pET21a vector. All DNA constructs were sequence verified. Protein

fragments were expressed in *Escherichia coli* BL21 Rosetta (DE3) (Novagen) cells, which

were cultured in LB or TB media, induced at OD_{600} ~0.4-0.6 with 1 mM IPTG and grown at

20°C or 30°C for ~16 h. Protein fragments for NMR were produced using the same cells

cultured in M9 minimal media supplemented with MEM vitamin solution (Gibco) and 1g/L

$^{15}NH_4Cl$ (Cambridge Isotopes) to produce uniformly labeled [U-^{15}N] protein. Cells were lysed

by sonication in denaturing buffer (20 mM Tris HCl, pH 7.5, 6 M guanidine hydrochloride, 10

mM $CaCl_2$, and 20 mM imidazole) or non-denaturing buffer (20 mM Tris HCl, pH 7.5, and 20

mM imidazole for *hs* TMC1-NT). The cleared lysates were loaded onto Ni-Sepharose (GE

Healthcare) and eluted with denaturing buffer or non-denaturing buffer supplemented with 500

mM imidazole. Denatured proteins were refolded using MWCO 2,000 membranes

(Spectra/Por 7) in dialysis buffer (400 mM Arg-HCl, 20 mM Tris-HCl, pH 7.5, 150 mM KCl,

50 mM NaCl, 2 mM $CaCl_2$, 2 mM DTT) for ~16 hr. Before starting refolding reactions (when

applicable), elution solutions were diluted to ~0.5 mg/mL using the denaturing buffer, and then

reduced by using 2 mM DTT in the diluted sample. Refolded and soluble proteins were

concentrated using Vivaspin 20 or Amicon 15 centrifugal concentrators (10 kDa, 5 kDa, or 3

kDa molecular weight cutoff) and further purified on Superdex S75 columns (GE Healthcare)

in SEC buffer (50 mM HEPES, pH 7.5, 126 mM KCl, 10 mM NaCl, 3 mM CaCl$_2$, 5 mM DTT).

Proteins were concentrated to ~0.5 mL for nuclear magnetic resonance (NMR) experiments.

NMR

Purified NMR samples were at concentrations of ~1.4 mM, ~0.2 mM, ~0.8 mM, and ~0.2 mM

for CIB2, CIB3, CIB2 + TMC1-IL1, and CIB3 + TMC1-IL1, respectively, in SEC buffer (50

mM HEPES, pH 7.5, 126 mM KCl, 10 mM NaCl, 3 mM CaCl$_2$, 5 mM DTT), supplemented

with 10% (v/v) D$_2$O and 0.2 mM 2,2-dimethyl-2-silapentane-5-sulphonate (DSS) for field-

frequency locking and chemical shift referencing. Two-dimensional ^{1}H-^{15}N TROSY-HSQC

spectra (Nietlispach, 2005) were recorded for each sample at 25°C on a Bruker Avance III HD

800 MHz spectrometer equipped with a 5 mm TXI cryoprobe with z-gradients. Data were

collected using 20 to 24 scans, 2,048 x 280 datapoints, and spectral widths of 17482.518 x

4054.298 Hz in the ^{1}H and ^{15}N dimensions, respectively. Data were processed and visualized

using NMRFx (Norris et al., 2016).

Simulated Systems

Models for monomeric *hs* TMC1 (residues 84 to 760; NCBI ID: NP_619636.2) and *hs* TMC2

(148 to 863; NCBI ID: NP_542789.2) in complex with full-length *hs* CIB2 (NCBI ID:

NP_006374.1) and *hs* CIB3 (NCBI ID: NP_473454.1), as well as truncated dimeric models of

hs TMC1 (residues 260 to 760; NCBI ID: NP_619636.2) and *hs* TMC2 (residues 324 to 863;

NCBI ID: NP_542789.2) were generated using AF2 (Evans et al., 2021; Jumper et al., 2021).

The TMC2 C-terminal end (residues 818 to 863) was truncated to match the C-terminal end of

TMC1. Using VMD (Humphrey et al., 1996), a model of dimeric *hs* TMC1 + *hs* CIB3-Ca^{2+}

was constructed by aligning the monomeric *hs* TMC1 + *hs* CIB3 complex to the

transmembrane region of the truncated dimeric *hs* TMC1 model. Coordinates for Ca^{2+} ions

were obtained by aligning the crystal structure of Mg^{2+}-bound *hs* CIB3 in complex with a *hs*

TMC1-IL1 peptide (PDB code: 6WUD) (Liang et al., 2021) to the AF2 model of monomeric

hs TMC1 + *hs* CIB3, then manually changing Mg^{2+} for Ca^{2+}. This model was minimized for

1,000 steps in vacuum using NAMD (Phillips et al., 2005) and the CHARMM36 force field

(Best et al., 2012; Huang and MacKerell, 2014; Vanommeslaeghe and MacKerell, 2015) and

served as a reference model for aligning other combinations of *hs* TMC1 or *hs* TMC2 with *hs*

CIB2 or *hs* CIB3. All models were either embedded into pure POPC or mixed membrane lipid

bilayers using CHARMM-GUI (Jo et al., 2009, 2008, 2007; Lee et al., 2019) as summarized

in Table supplement 1. The mixed membrane mimicked stereocilia membrane composition

(Zhao et al., 2012). The protein-lipid system was then solvated using TIP3P explicit water and

neutralized with 0.150 M KCl, representative of endolymph ionic concentration (Bosher and

Warren, 1978).

<u>Simulations using NAMD</u>

All-atom MD simulations were performed and analyzed using NAMD 2.13 (Phillips et al.,

2005) and the CHARMM36 force field with CMAP corrections (Best et al., 2012; Huang and

MacKerell, 2014; Vanommeslaeghe and MacKerell, 2015). For all systems, equilibrium

simulations consisted of 2,000 steps of minimization, 0.5 ns of dynamics with everything

excluding lipid tails fixed/constrained, 0.5 ns of dynamics with harmonic constraints applied

to the protein (1 kcal mol^{-1} $Å^{-2}$), 1 ns of free dynamics in the *NpT* ensemble ($\gamma = 1$ ps^{-1}), and up

to 100 ns of free dynamics in the *NpT* ensemble ($\gamma = 0.1$ ps^{-1}) prior to non-equilibrium

production runs. A piston period of 200 fs and a damping timescale of 50 fs was used for

pressure control and atomic coordinates were saved every 5 ps. All simulations were performed

at 310 K and 1 atmosphere of pressure by using the Langevin thermostat and Nosé-Hoover

Langevin pressure control. A 12-Å cutoff distance was employed with a force-based switching

function starting at 10 Å. Periodic boundary conditions and the PME method were used to

calculate long-range electrostatic interactions with a grid density greater than 1 Å^{-3}. An

integration timestep of 2 fs with electrostatic interactions computed every other timestep was

utilized for every simulation in tandem with the SHAKE algorithm for the constraint of

hydrogen atoms. A constant electric field was applied to all atoms in non-equilibrium

simulations with transmembrane potential (Gumbart et al., 2012).

<u>Simulations using Anton 2</u>

We used Anton 2, a massively parallel special purpose machine for MD simulations (Shaw et

al., 2014), to run simulations of *hs* TMC1 + *hs* CIB2-Ca^{2+} in POPC (simulations S1d-g; Table

supplement 1), *hs* TMC1 + *hs* CIB2 in POPC (simulations S2d-2g; Table supplement 1), *hs*

TMC1 in POPC (simulations S3d-g; Table supplement 1), and *hs* TMC1 + *hs* CIB2 in a mixed

membrane (simulations S4c-d; Table supplement 1). Coordinates of the pre-equilibrated

systems were converted to an Anton 2 compatible dms file format using the NAMDtoDMS.py

script provided by the Pittsburgh Supercomputing Center. Simulations used CHARMM36

forcefields and were performed in the *NpT* ensemble (310 K, 1 atm) using the multigrator

integration framework (Lippert et al., 2013) with a 2.5 fs integration timestep and saved every

120 ps. The MTK barostat and Nosé-Hoover thermostat was updated every 480 and 24 steps,

respectively. Semi-isotropic pressure coupling was used. Long range electrostatic interactions

were calculated using u-series electrostatics with a 64 × 64 × 64 grid. Van der Waals

interactions were calculated with a cutoff of 12 Å. Transmembrane voltages were generated by

applying a constant electric field to all atoms (Gumbart et al., 2012).

<u>Simulation analysis procedures and tools</u>

BSA was computed in VMD by measuring SASA for both TMC and CIB monomers, then

subtracting SASA for the complex. BSA for interacting molecules A and B were computed as

$BSA_{AB} = 1/2(SASA_A + SASA_B - SASA_{AB})$. RMSD values were computed in VMD using Cα

atoms and initial models as references, with alignments carried out using the corresponding

subdomains when indicated. VMD was used to analyze trajectories, render molecular images,

and create videos. Xmgrace and gnuplot were used to generate plots. Sequences for CIB and

TMC isoforms from various species were obtained from the NCBI protein database. Sequences

were aligned utilizing the MUSCLE algorithm within Geneious (Kearse et al., 2012). Aligned

sequences were loaded into JalView (Waterhouse et al., 2009) and colored based on a sequence

identity threshold of 45%. Ionic currents were computed by counting the number of crossing,

with conductance values estimated as in Walujkar et al., 2021.

<u>Resource Availability</u>

Lead Contact

Further information and requests for resources and reagents should be directed to the Lead

Contacts, Drs. Marcos Sotomayor (sotomayor.8@osu.edu) and Zubair M. Ahmed

(zmahmed@som.umaryland.edu).

Materials availability

Materials generated in this study, including strains, plasmids and clones, are freely available

from the Lead Contact upon request.

Data and code availability

This study did not generate any unique datasets or code.

Acknowledgements

We acknowledge Sakina Rehman for assistance with animal work, Neal Taliwal for assistance with molecular cloning, Chunhua Yuan for assistance with initial nuclear magnetic resonance experiments, and Collin R. Nisler and Harsha Mandayam Bharathi for assistance with simulations analyses. Simulations were carried out at the Ohio Supercomputer Center (OSC grants PAS1037 and PAA0217) and the Pittsburgh Supercomputer Center (PSC Anton 2 grant MCB150024P). Nuclear magnetic resonance data were acquired at the Ohio State University Campus Chemical Instrument Center. This study was supported by NIH/NIDCD grants R01DC015271 to M.S. and R01DC012564 to Z.M.A. and G.I.F., an Intramural Research Program Grant 1ZIADC000085-01 to K.S.K., and NIH grants UF1NS111695, R01DC018304, R01DC002390, R01DC018061 to K.E.C.

Additional information

Author contributions

Z.M.A., K.S.K., and M.S. conceived the studies. A.P.J.G. carried out all audiometric recordings in mice, immunolocalization studies, cochlear and vestibular explant cultures, FM 1-43 uptake experiment on mouse tissues, RT-qPCRs, exploratory testing, *Cib2* and *Cib3* mouse phenotyping and genotyping, cloning of CIB2 and CIB3 in pET21a vectors. K.S.K. conducted zebrafish FM 1-43, FM 4-64, FM 1-43FX labeling, Ca^{2+} imaging and immunostaining. A.J.B. created the *cib2* and *cib3* zebrafish mutants. R.A.R. did the zebrafish behavioral studies. W-H.W. did the cloning and mutagenesis of CIB and TMC constructs, protein expression and purification for NMR experiments, as well as model building and molecular dynamics simulations. J.S.M. did NMR experiments and NMR data processing. E.M.R. performed FM 1-43 uptake experiments in mice. J.M.L. and S.W. did model building. M.P.F. supervised NMR experiments and project. M.S. supervised project and assisted with

References

Ahmed ZM, Goodyear R, Riazuddin Saima, Lagziel A, Legan PK, Behra M, Burgess SM,

Lilley KS, Wilcox ER, Riazuddin Sheikh, Griffith AJ, Frolenkov GI, Belyantseva IA,

Richardson GP, Friedman TB. 2006. The tip-link antigen, a protein associated with the

transduction complex of sensory hair cells, is protocadherin-15. *J Neurosci* **26**:7022–

7034. doi:10.1523/JNEUROSCI.1163-06.2006

Akyuz N, Karavitaki KD, Pan B, Tamvakologos PI, Brock KP, Li Y, Marks DS, Corey DP.

2022. Mechanical gating of the auditory transduction channel TMC1 involves the fourth

and sixth transmembrane helices. *Sci Adv* **8**:eabo1126. doi:10.1126/sciadv.abo1126

Ballesteros A, Fenollar-Ferrer C, Swartz KJ. 2018. Structural relationship between the

putative hair cell mechanotransduction channel TMC1 and TMEM16 proteins. *Elife* **7**.

doi:10.7554/eLife.38433

Beraneck M, Cullen KE. 2007. Activity of vestibular nuclei neurons during vestibular and

optokinetic stimulation in the alert mouse. *J Neurophysiol* **98**:1549–1565.

doi:10.1152/jn.00590.2007

Beraneck M, McKee JL, Aleisa M, Cullen KE. 2008. Asymmetric recovery in cerebellar-

deficient mice following unilateral labyrinthectomy. *J Neurophysiol* **100**:945–958.

doi:10.1152/jn.90319.2008

Best RB, Zhu X, Shim J, Lopes PE, Mittal J, Feig M, Mackerell Jr. AD. 2012. Optimization

of the additive CHARMM all-atom protein force field targeting improved sampling of

the backbone phi, psi and side-chain chi(1) and chi(2) dihedral angles. *J Chem Theory*

Comput **8**:3257–3273. doi:10.1021/ct300400x

Beurg M, Cui R, Goldring AC, Ebrahim S, Fettiplace R, Kachar B. 2018. Variable number of

TMC1-dependent mechanotransducer channels underlie tonotopic conductance

gradients in the cochlea. *Nat Commun* **9**:2185. doi:10.1038/s41467-018-04589-8

Beurg M, Fettiplace R, Nam J-H, Ricci AJ. 2009. Localization of inner hair cell

mechanotransducer channels using high-speed calcium imaging. *Nat Neurosci* **12**:553–

558. doi:10.1038/nn.2295

Beurg M, Nam J-H, Fettiplace R. 2021. The speed of the hair cell mechanotransducer

channel revealed by fluctuation analysis. *J Gen Physiol* **153**. doi:10.1085/jgp.202112959

Booth KT, Kahrizi K, Babanejad M, Daghagh H, Bademci G, Arzhangi S, Zareabdollahi D,

Duman D, El-Amraoui A, Tekin M, Najmabadi H, Azaiez H, Smith RJ. 2018. Variants

in CIB2 cause DFNB48 and not USH1J. *Clin Genet* **93**:812–821. doi:10.1111/cge.13170

Bosher SK, Warren RL. 1978. Very low calcium content of cochlear endolymph, an

extracellular fluid. *Nature* **273**:377–378. doi:10.1038/273377a0

Chen Z, Zhu S, Kindig K, Wang S, Chou SW, Davis RW, Dercoli MR, Weaver H, Stepanyan

R, McDermott BM. 2020. Tmc proteins are essential for zebrafish hearing where Tmc1

is not obligatory. *Hum Mol Genet* **29**:2004–2021. doi:10.1093/HMG/DDAA045

Chou SW, Chen Z, Zhu S, Davis RW, Hu J, Liu L, Fernando CA, Kindig K, Brown WC,

Stepanyan R, McDermott BM. 2017. A molecular basis for water motion detection by

the mechanosensory lateral line of zebrafish. *Nat Commun* **8**. doi:10.1038/s41467-017-

01604-2

Cullen KE. 2019. Vestibular processing during natural self-motion: implications for

perception and action. *Nat Rev Neurosci* **20**:346–363. doi:10.1038/s41583-019-0153-1

Cunningham CL, Qiu X, Wu Z, Zhao B, Peng G, Kim YH, Lauer A, Müller U. 2020. TMIE

Defines Pore and Gating Properties of the Mechanotransduction Channel of Mammalian

Cochlear Hair Cells. *Neuron* **107**:126-143.e8. doi:10.1016/j.neuron.2020.03.033

Dal Cortivo G, Marino V, Iacobucci C, Vallone R, Arlt C, Rehkamp A, Sinz A, Dell'Orco D. 2019. Oligomeric state, hydrodynamic properties and target recognition of human Calcium and Integrin Binding protein 2 (CIB2). *Sci Rep* **9**. doi:10.1038/s41598-019-51573-3

Effertz T, Becker L, Peng AW, Ricci AJ. 2017. Phosphoinositol-4,5-bisphosphate regulates auditory hair-cell mechanotransduction-channel pore properties and fast adaptation. *J Neurosci* **37**:11632–11646. doi:10.1523/JNEUROSCI.1351-17.2017

Erickson T, Morgan CP, Olt J, Hardy K, Busch-Nentwich E, Maeda R, Clemens R, Krey JF, Nechiporuk A, Barr-Gillespie PG, Marcotti W, Nicolson T. 2017. Integration of Tmc1/2 into the mechanotransduction complex in zebrafish hair cells is regulated by Transmembrane O-methyltransferase (Tomt). *Elife* **6**. doi:10.7554/eLife.28474

Ernest S, Rauch GJ, Haffter P, Geisler R, Petit C, Nicolson T. 2000. Mariner is defective in myosin VIIA: A zebrafish model for human hereditary deafness. *Hum Mol Genet* **9**:2189–2196. doi:10.1093/hmg/9.14.2189

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, Ronneberger O, Bodenstein S, Zielinski M, Bridgland A, Potapenko A, Cowie A, Tunyasuvunakool K, Jain R, Clancy E, Kohli P, Jumper J, Hassabis D. 2021. Protein complex prediction with AlphaFold-Multimer. *bioRxiv* 2021.10.04.463034. doi:10.1101/2021.10.04.463034

Fettiplace R. 2017. Hair cell transduction, tuning, and synaptic transmission in the mammalian cochlea. *Compr Physiol* **7**:1197–1227. doi:10.1002/cphy.c160049

Fettiplace R, Furness DN, Beurg M. 2022. The conductance and organization of the TMC1-

containing mechanotransducer channel complex in auditory hair cells. *Proc Natl Acad*

Sci U S A **119**:e2210849119. doi:10.1073/pnas.2210849119

Frolenkov GI, Belyantseva IA, Friedman TB, Griffith AJ. 2004. Genetic insights into the

morphogenesis of inner ear hair cells. *Nat Rev Genet* **5**:489–498. doi:10.1038/nrg1377

Giese APJ, Tang YQ, Sinha GP, Bowl MR, Goldring AC, Parker A, Freeman MJ, Brown

SDM, Riazuddin S, Fettiplace R, Schafer WR, Frolenkov GI, Ahmed ZM. 2017. CIB2

interacts with TMC1 and TMC2 and is essential for mechanotransduction in auditory

hair cells. *Nat Commun* **8**. doi:10.1038/s41467-017-00061-1

Gillespie PG, Müller U. 2009. Mechanotransduction by hair cells: models, molecules, and

mechanisms. *Cell* **139**:33–44. doi:10.1016/j.cell.2009.09.010

Gleason MR, Nagiel A, Jamet S, Vologodskaia M, López-Schier H, Hudspeth AJ. 2009. The

transmembrane inner ear (Tmie) protein is essential for normal hearing and balance in

the zebrafish. *Proc Natl Acad Sci U S A* **106**:21347–52. doi:10.1073/pnas.0911632106

Gopal SR, Chen DHC, Chou SW, Zang J, Neuhauss SCF, Stepanyan R, McDermott BM,

Alagramam KN. 2015. Zebrafish models for the mechanosensory hair cell dysfunction

in usher syndrome 3 reveal that Clarin-1 is an essential hair bundle protein. *J Neurosci*

35:10188–10201. doi:10.1523/JNEUROSCI.1096-15.2015

Gumbart J, Khalili-Araghi F, Sotomayor M, Roux B. 2012. Constant electric field

simulations of the membrane potential illustrated with simple systems. *Biochim Biophys*

Acta - Biomembr **1818**:294–302. doi:10.1016/j.bbamem.2011.09.030

Huang H, Bogstie JN, Vogel HJ. 2012. Biophysical and structural studies of the human

calcium-and integrin-binding protein family: Understanding their functional similarities

and differences. *Biochem Cell Biol* **90**:646–656. doi:10.1139/o2012-021

Huang J, MacKerell ADJ. 2014. CHARMM36 all-atom additive protein force field:

Validation based on comparison to NMR data. *J Comput Chem* **34**:2135–2145.

doi:10.1002/jcc.23354.CHARMM36

Humphrey W, Dalke A, Schulten K. 1996. VMD: Visual molecular dynamics. *J Mol Graph*

14:33–38. doi:10.1016/0263-7855(96)00018-5

Jeong H, Clark S, Goehring A, Dehghani-Ghahnaviyeh S, Rasouli A, Tajkhorshid E, Gouaux

E. 2022. Structures of the TMC-1 complex illuminate mechanosensory transduction.

Nature **610**:796–803. doi:10.1038/s41586-022-05314-8

Jiang T, Kindt K, Wu DK. 2017. Transcription factor emx2 controls stereociliary bundle

orientation of sensory hair cells. *Elife* **6**:1–26. doi:10.7554/eLife.23661

Jo S, Kim T, Im W. 2007. Automated builder and database of protein/membrane complexes

for molecular dynamics simulations. *PLoS One* **2**. doi:10.1371/journal.pone.0000880

Jo S, Kim T, Iyer VG, Im W. 2008. CHARMM-GUI: A web-based graphical user interface

for CHARMM. *J Comput Chem* **29**:1859–1865. doi:10.1002/jcc.20945

Jo S, Lim JB, Klauda JB, Im W. 2009. CHARMM-GUI membrane builder for mixed bilayers

and its application to yeast membranes. *Biophys J* **97**:50–58.

doi:10.1016/j.bpj.2009.04.013

Jojoa-Cruz S, Saotome K, Murthy SE, Tsui CCA, Sansom MS, Patapoutian A, Ward AB.

2018. Cryo-EM structure of the mechanically activated ion channel OSCA1.2. *Elife* **7**.

doi:10.7554/eLife.41845

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,

Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie

A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D,

Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S,

1105   Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly

1106   accurate protein structure prediction with AlphaFold. *Nature* 1–11. doi:10.1038/s41586-

1107   021-03819-2

1108   Karplus M, Petsko GA. 1990. Molecular dynamics simulations in biology. *Nature* **347**:631–

1109   9. doi:10.1038/347631a0

1110   Kawashima Y, Géléoc GSG, Kurima K, Labay V, Lelli A, Asai Y, Makishima T, Wu DK,

1111   Della Santina CC, Holt JR, Griffith AJ. 2011. Mechanotransduction in mouse inner ear

1112   hair cells requires transmembrane channel-like genes. *J Clin Invest* **121**:4796–809.

1113   doi:10.1172/JCI60405

1114   Kawashima Y, Kurima K, Pan B, Griffith AJ, Holt JR. 2015. Transmembrane channel-like

1115   (TMC) genes are required for auditory and vestibular mechanosensation. *Pflügers Arch -*

1116   *Eur J Physiol* **467**:85–94. doi:10.1007/s00424-014-1582-3

1117   Kazmierczak P, Sakaguchi H, Tokita J, Wilson-Kubalek EM, Milligan RA, Müller U, Kachar

1118   B. 2007. Cadherin 23 and protocadherin 15 interact to form tip-link filaments in sensory

1119   hair cells. *Nature* **449**:87–91. doi:10.1038/nature06091

1120   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,

1121   Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious

1122   Basic: an integrated and extendable desktop software platform for the organization and

1123   analysis of sequence data. *Bioinformatics* **28**:1647–9. doi:10.1093/bioinformatics/bts199

1124   Kindig K, Stepanyan R, Kindt KS, McDermott BM. 2023. Asymmetric mechanotransduction

1125   by hair cells of the zebrafish lateral line. *Curr Biol* **33**:1295-1307.e3.

1126   doi:10.1016/j.cub.2023.02.033

Lee Jumin, Patel DS, Ståhle J, Park SJ, Kern NR, Kim S, Lee Joonseong, Cheng X, Valvano

MA, Holst O, Knirel YA, Qi Y, Jo S, Klauda JB, Widmalm G, Im W. 2019. CHARMM-

GUI Membrane Builder for Complex Biological Membrane Simulations with

Glycolipids and Lipoglycans. *J Chem Theory Comput* **15**:775–786.

doi:10.1021/acs.jctc.8b01066

Liang X, Qiu X, Dionne G, Cunningham CL, Pucak ML, Peng G, Kim YH, Lauer A, Shapiro

L, Müller U. 2021. CIB2 and CIB3 are auxiliary subunits of the mechanotransduction

channel of hair cells. *Neuron*. doi:10.1016/j.neuron.2021.05.007

Lippert R a, Predescu C, Ierardi DJ, Mackenzie KM, Eastwood MP, Dror RO, Shaw DE.

2013. Accurate and efficient integration for molecular dynamics simulations at constant

temperature and pressure. *J Chem Phys* **139**:164106. doi:10.1063/1.4825247

Liu X, Wang J, Sun L. 2018. Structure of the hyperosmolality-gated calcium-permeable

channel OSCA1.2. *Nat Commun* **9**:5060. doi:10.1038/s41467-018-07564-5

López-Schier H, Hudspeth AJ. 2006. A two-step mechanism underlies the planar polarization

of regenerating sensory hair cells. *Proc Natl Acad Sci U S A* **103**:18615–18620.

doi:10.1073/pnas.0608536103

Lukasz D, Kindt KS. 2018. In vivo calcium imaging of lateral-line hair cells in larval

zebrafish. *J Vis Exp* **2018**:1–14. doi:10.3791/58794

Maeda R, Kindt KS, Mo W, Morgan CP, Erickson T, Zhao H, Clemens-Grisham R, Barr-

Gillespie PG, Nicolson T. 2014. Tip-link protein protocadherin 15 interacts with

transmembrane channel-like proteins TMC1 and TMC2. *Proc Natl Acad Sci U S A*

111:12907–12. doi:10.1073/pnas.1402152111

Maeda R, Pacentine I V., Erickson T, Nicolson T. 2017. Functional analysis of the

transmembrane and cytoplasmic domains of Pcdh15a in zebrafish hair cells. *J Neurosci*

2216–16. doi:10.1523/JNEUROSCI.2216-16.2017

McInturff S, Burns JC, Kelley MW. 2018. Characterization of spatial and temporal

development of Type I and Type II hair cells in the mouse utricle using new cell-type-

specific markers. *Biol Open* **7**:1–13. doi:10.1242/bio.038083

Meyers JR, MacDonald RB, Duggan A, Lenzi D, Standaert DG, Corwin JT, Corey DP. 2003.

Lighting up the Senses: FM1-43 Loading of Sensory Cells through Nonselective Ion

Channels. *J Neurosci* **23**:4054–4065. doi:10.1523/JNEUROSCI.23-10-04054.2003

Michel V, Booth KT, Patni P, Cortese M, Azaiez H, Bahloul A, Kahrizi K, Labbé M, Emptoz

A, Lelli A, Dégardin J, Dupont T, Aghaie A, Oficjalska-Pham D, Picaud S, Najmabadi

H, Smith RJ, Bowl MR, Brown SD, Avan P, Petit C, El-Amraoui A. 2017. CIB2,

defective in isolated deafness, is key for auditory hair cell mechanotransduction and

survival. *EMBO Mol Med* **9**:1711–1731. doi:10.15252/emmm.201708087

Naz S, Giguere CM, Kohrman DC, Mitchem KL, Riazuddin Saima, Morell RJ, Ramesh A,

Srisailpathy S, Deshmukh D, Riazuddin Sheikh, Griffith AJ, Friedman TB, Smith RJH,

Wilcox ER. 2002. Mutations in a novel gene, TMIE, are associated with hearing loss

linked to the DFNB6 locus. *Am J Hum Genet* **71**:632–636. doi:10.1086/342193

Nicolson T. 2017. The genetics of hair-cell function in zebrafish. *J Neurogenet*.

doi:10.1080/01677063.2017.1342246

Nicolson T, Rüsch A, Friedrich RW, Granato M, Ruppersberg JP, Nüsslein-Volhard C. 1998.

Genetic analysis of vertebrate sensory hair cell mechanosensation: The zebrafish circler

mutants. *Neuron* **20**:271–283. doi:10.1016/S0896-6273(00)80455-9

Nietlispach D. 2005. Suppression of anti-TROSY lines in a sensitivity enhanced gradient

selection TROSY scheme. *J Biomol NMR* **31**:161–6. doi:10.1007/s10858-004-8195-7

Norris M, Fetler B, Marchant J, Johnson BA. 2016. NMRFx Processor: a cross-platform

NMR data processing program. *J Biomol NMR* **65**:205–216. doi:10.1007/s10858-016-

0049-6

Pan B, Akyuz N, Liu X-P, Asai Y, Nist-Lund C, Kurima K, Derfler BH, György B,

Limapichat W, Walujkar S, Wimalasena LN, Sotomayor M, Corey DP, Holt JR. 2018.

TMC1 Forms the Pore of Mechanosensory Transduction Channels in Vertebrate Inner

Ear Hair Cells. *Neuron* **99**:736-753.e6. doi:10.1016/j.neuron.2018.07.033

Phillips JB, Blanco-Sanchez B, Lentz JJ, Tallafuss A, Khanobdee K, Sampath S, Jacobs ZG,

Han PF, Mishra M, Titus TA, Williams DS, Keats BJ, Washbourne P, Westerfield M.

2011. Harmonin (Ush1c) is required in zebrafish Müller glial cells for photoreceptor

synaptic development and function. *DMM Dis Model Mech* **4**:786–800.

doi:10.1242/dmm.006429

Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kalé

L, Schulten K. 2005. Scalable molecular dynamics with NAMD. *J Comput Chem*

26:1781–1802. doi:10.1002/jcc.20289

Pujol-Martí J, López-Schier H. 2013. Developmental and architectural principles of the

lateral-line neural map. *Front Neural Circuits* **7**:1–9. doi:10.3389/fncir.2013.00047

Qiu X, Liang X, Llongueras JP, Cunningham C, Müller U. 2023. The tetraspan LHFPL5 is

critical to establish maximal force sensitivity of the mechanotransduction channel of

cochlear hair cells. *Cell Rep* **42**:112245. doi:10.1016/j.celrep.2023.112245

Riazuddin Saima, Belyantseva IA, Giese APJ, Lee K, Indzhykulian AA, Nandamuri SP,

Yousaf R, Sinha GP, Lee S, Terrell D, Hegde RS, Ali RA, Anwar S, Andrade-Elizondo

PB, Sirmaci A, Parise L V., Basit S, Wali A, Ayub M, Ansar M, Ahmad W, Khan SN, Akram J, Tekin M, Riazuddin Sheikh, Cook T, Buschbeck EK, Frolenkov GI, Leal SM, Friedman TB, Ahmed ZM. 2012. Alterations of the CIB2 calcium-and integrin-binding protein cause Usher syndrome type 1J and nonsyndromic deafness DFNB48. *Nat Genet* **44**:1265–1271. doi:10.1038/ng.2426

Seco CZ, Giese AP, Shafique S, Schraders M, Oonk AMM, Grossheim M, Oostrik J, Strom T, Hegde R, Van Wijk E, Frolenkov GI, Azam M, Yntema HG, Free RH, Riazuddin S, Verheij JBGM, Admiraal RJ, Qamar R, Ahmed ZM, Kremer H. 2016. Novel and recurrent CIB2 variants, associated with nonsyndromic deafness, do not affect calcium buffering and localization in hair cells. *Eur J Hum Genet* **24**:542–549. doi:10.1038/ejhg.2015.157

Seiler C, Ben-David O, Sidi S, Hendrich O, Rusch A, Burnside B, Avraham KB, Nicolson T. 2004. Myosin VI is required for structural integrity of the apical surface of sensory hair cells in zebrafish. *Dev Biol* **272**:328–338. doi:10.1016/j.ydbio.2004.05.004

Shaw DE, Grossman JP, Bank JA, Batson B, Butts JA, Chao JC, Deneroff MM, Dror RO, Even A, Fenton CH, Forte A, Gagliardo J, Gill G, Greskamp B, Ho CR, Ierardi DJ, Iserovich L, Kuskin JS, Larson RH, Layman T, Lee L-S, Lerer AK, Li C, Killebrew D, Mackenzie KM, Mok SY-H, Moraes MA, Mueller R, Nociolo LJ, Peticolas JL, Quan T, Ramot D, Salmon JK, Scarpazza DP, Schafer U Ben, Siddique N, Snyder CW, Spengler J, Tang PTP, Theobald M, Toma H, Towles B, Vitale B, Wang SC, Young C. 2014. Anton 2: Raising the Bar for Performance and Programmability in a Special-Purpose Molecular Dynamics SupercomputerSC14: International Conference for High Performance Computing, Networking, Storage and Analysis. IEEE. pp. 41–53. doi:10.1109/SC.2014.9

Sheets L, He XJ, Olt J, Schreck M, Petralia RS, Wang YX, Zhang Q, Beirl A, Nicolson T,

Marcotti W, Trapani JG, Kindt KS. 2017. Enlargement of ribbons in zebrafish hair cells

increases calcium currents but disrupts afferent spontaneous activity and timing of

stimulus onset. *J Neurosci* **37**:6299–6313. doi:10.1523/JNEUROSCI.2878-16.2017

Shi T, Beaulieu MO, Saunders LM, Fabian P, Trapnell C, Segil N, Crump JG, Raible DW.

2023. Single-cell transcriptomic profiling of the zebrafish inner ear reveals molecularly

distinct hair cell and supporting cell subtypes. *Elife* **12**:1–24. doi:10.7554/eLife.82978

Siemens J, Lillo C, Dumont R a, Reynolds A, Williams DS, Gillespie PG, Müller U. 2004.

Cadherin 23 is a component of the tip link in hair-cell stereocilia. *Nature* **428**:950–955.

doi:10.1038/nature02483

Smith ET, Pacentine I, Shipman A, Hill M, Nicolson T. 2020. Disruption of tmc1/2a/2b

Genes in Zebrafish Reveals Subunit Requirements in Subtypes of Inner Ear Hair Cells. *J*

Neurosci **40**:4457–4468. doi:10.1523/JNEUROSCI.0163-20.2020

Söllner C, Rauch G-J, Siemens J, Geisler R, Schuster SC, the Tübingen 2000 Screen

Consortium, Müller U, Nicolson T. 2004. Mutations in cadherin 23 affect tip links in

zebrafish sensory hair cells. *Nature* **428**:955–959. doi:10.1038/nature02484

Sotomayor M, Weihofen WA, Gaudet R, Corey DP. 2012. Structure of a force-conveying

cadherin bond essential for inner-ear mechanotransduction. *Nature* **492**:128–32.

doi:10.1038/nature11590

Sukharev S, Corey DP. 2004. Mechanosensitive channels: multiplicity of families and gating

paradigms. *Sci STKE* **2004**:re4. doi:10.1126/stke.2192004re4

Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Žídek A, Bridgland A, Cowie A,

Meyer C, Laydon A, Velankar S, Kleywegt GJ, Bateman A, Evans R, Pritzel A,

Figurnov M, Ronneberger O, Bates R, Kohl SAA, Potapenko A, Ballard AJ, Romera-

Paredes B, Nikolov S, Jain R, Clancy E, Reiman D, Petersen S, Senior AW,

Kavukcuoglu K, Birney E, Kohli P, Jumper J, Hassabis D. 2021. Highly accurate protein

structure prediction for the human proteome. *Nature* 1–9. doi:10.1038/s41586-021-

03828-1

Vallone R, Dal Cortivo G, D'Onofrio M, Dell'Orco D. 2018. Preferential Binding of Mg2+

Over Ca2+ to CIB2 Triggers an Allosteric Switch Impaired in Usher Syndrome Type 1J.

Front Mol Neurosci **11**. doi:10.3389/fnmol.2018.00274

Vanommeslaeghe K, MacKerell AD. 2015. CHARMM additive and polarizable force fields

for biophysics and computer-aided drug design. *Biochim Biophys Acta* **1850**:861–871.

doi:10.1016/j.bbagen.2014.08.004

Varshney GK, Carrington B, Pei W, Bishop K, Chen Z, Fan C, Xu L, Jones M, LaFave MC,

Ledin J, Sood R, Burgess SM. 2016. A high-throughput functional genomics workflow

based on CRISPR/Cas9-mediated targeted mutagenesis in zebrafish. *Nat Protoc*

11:2357–2375. doi:10.1038/nprot.2016.141

Walujkar S, Lotthammer JM, Nisler CR, Sudar JC, Ballesteros A, Sotomayor M. 2021. In

Silico Electrophysiology of Inner-Ear Mechanotransduction Channel TMC1 Models.

bioRxiv 2021.09.17.460860. doi:10.1101/2021.09.17.460860

Wang X, Liu S, Cheng Q, Qu C, Ren R, Du H, Li N, Yan K, Wang Y, Xiong W, Xu Z. 2023.

CIB2 and CIB3 regulate stereocilia maintenance and mechanoelectrical transduction in

mouse vestibular hair cells. *J Neurosci*. doi:10.1523/JNEUROSCI.1807-22.2023

Wang Y, Li J, Yao X, Li W, Du H, Tang M, Xiong W, Chai R, Xu Z. 2017. Loss of CIB2

causes profound hearing loss and abolishes mechanoelectrical transduction in mice.

Front Mol Neurosci **10**. doi:10.3389/fnmol.2017.00401

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2--a

multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**:1189–

1191. doi:10.1093/bioinformatics/btp033

Xiong W, Grillet N, Elledge HM, Wagner TFJ, Zhao B, Johnson KR, Kazmierczak P, Müller

U. 2012. TMHS is an integral component of the mechanotransduction machinery of

cochlear hair cells. *Cell* **151**:1283–95. doi:10.1016/j.cell.2012.10.041

Zhang M, Wang D, Kang Y, Wu J-X, Yao F, Pan C, Yan Z, Song C, Chen L. 2018. Structure

of the mechanosensitive OSCA channels. *Nat Struct Mol Biol* **25**:850–858.

doi:10.1038/s41594-018-0117-6

Zhang Q, Li S, Wong HTC, He XJ, Beirl A, Petralia RS, Wang YX, Kindt KS. 2018.

Synaptically silent sensory hair cells in zebrafish are recruited after damage. *Nat*

Commun **9**:1–16. doi:10.1038/s41467-018-03806-8

Zhao H, Williams DE, Shin J-B, Brügger B, Gillespie PG. 2012. Large membrane domains in

hair bundles specify spatially constricted radixin activation. *J Neurosci* **32**:4600–9.

doi:10.1523/JNEUROSCI.6184-11.2012

Zhu J, Xu H, Song H, Li X, Wang N, Zhao J, Zheng X, Kim KY, Zhang H, Mao Q, Xia H.

2021. CRISPR/Cas9-mediated grna gene knockout leads to neurodevelopmental defects

and motor behavior changes in zebrafish. *J Neurochem* **157**:520–531.

doi:10.1111/jnc.15307

1286

1287

 Figures and figure legends

Figure 1. Expression of *Cib* genes in the mouse inner ear and generation of *Cib3* knockout mice. (**A**) Maximum intensity projections of Z-stacks of confocal fluorescent images and corresponding DIC images of *Cib2*$^{KO/+}$ and *Cib2*$^{KO/KO}$ cultured vestibular end organ explants imaged after exposure to 3 µM of FM 1-43 for 10 s. The samples were dissected at P5 and kept 2 days *in vitro* (P5 + 2div). Scale bar: 20 µm. (**B**) Real-time quantitative RT–PCR analysis of mRNA levels in the cochlear and vestibular sensory epithelia at P12 revealed different expression of *Cib1*, *Cib2*, *Cib3,* and *Cib4* in the auditory and vestibular end organs. (**C**) Violin plots generated by gEAR portal showing the expression levels of *Cib2*, *Cib3*, *Tmc1,* and *Tmc2* by scRNA-seq in immature, type 1, and type 2 utricular vestibular hair cells. TPM: transcript per million. (**D**) Gene structure of the wild-type and *Cib3*^{KO} alleles. Exons are shown in yellow. Indel in exon 4 leads to a nonsense mutation and a premature stop at the protein level (p.

Asp54*) (**E**) Real-time quantitative RT–PCR analysis using cDNA from brain tissue of

Cib2;Cib3 mutant mice was used to analyze the expression of *Cib2* and *Cib3*. Contrary to *Cib2*

mRNA, *Cib3* mRNA appears to be stable in the *Cib2$^{KO/KO}$;Cib3$^{KO/KO}$* mice and does not go

through nonsense mediated decay process. The relative expression of *Cib2* and *Cib3* members

were normalized against *hprt*. Asterisks indicate statistical significance: ***, $p<0.001$

(Student's t-test). n.s.: not significant. Error bars represent SEM. (**F**) Western Blot analysis on

Cib2;Cib3 mouse heart tissue was performed using CIB3 antibody. No CIB3 protein was

detected in the *Cib2$^{KO/KO}$;Cib3$^{KO/KO}$* mice, validating our mouse model. GAPDH detection was

used as a loading control.

1310

1311

1312

Figure 2. CIB2/3 double mutant mice have profound hearing loss. (**A**) ABR thresholds to

broadband clicks and tone-pips with frequencies of 8 kHz, 16 kHz, 24 kHz and 32 kHz in

Cib2$^{+/+}$;Cib3$^{KO/KO}$ (black; $n = 5$) and *Cib2$^{KO/KO}$;Cib3$^{KO/KO}$* (red, $n = 5$) mice at P16. The same

animals were tested with clicks and tone pips. (**B**) ABR pure tone traces of P32

Cib2$^{+/+}$;Cib3$^{KO/KO}$ (black) and *Cib2$^{KO/KO}$;Cib3$^{KO/KO}$* (red) mice measured at 32 kHz. Hearing

thresholds are in Decibels (dB SPL). (**C**) Maximum intensity projections of Z-stacks of

confocal fluorescent images (right) and corresponding DIC images (left) of *Cib2$^{KO/+}$;Cib3$^{+/+}$*,

Cib2$^{KO/KO}$;Cib3$^{+/+}$, *Cib2$^{+/+}$;Cib3$^{KO/KO}$* and *Cib2$^{KO/KO}$;Cib3$^{KO/KO}$* cultured organ of Corti

explants imaged after exposure to 3 µM of FM 1-43 for 10 s. The samples were dissected at P5

and kept 2 days in vitro (P5 + 2div). Scale bar: 20 µm. (**D**) Maximum intensity projections of

confocal Z-stacks of apical, medial and basal turns of organ of Corti of control and

Cib2$^{KO/KO}$;Cib3$^{KO/KO}$ mice immunostained with an anti-myosin VIIa antibody (green) and

counterstained with phalloidin (red) at P18. Asterisks indicate hair cell loss. Scale bar: 10 µm.

1326

1327

1328

Figure 3. CIB2/3 double mutant mice have vestibular dysfunction. (**A**) Traces showing the

open-field exploratory behavior of P60 $Cib2^{+/+};Cib3^{+/+}$, $Cib2^{KO/+};Cib3^{+/+}$, $Cib2^{+/+};Cib3^{KO/KO}$,

$Cib2^{KO/+};Cib3^{KO/KO}$ and $Cib2^{KO/KO};Cib3^{KO/KO}$ mutant mice. (**B**) Examples of head-velocity

(grey) and resultant eye-velocity traces evoked during VORd testing. Note, the eye movement

is compensatory, and the trace has been inverted to facilitate comparison with head velocity.

(**C**) VORd Gain and phase (mean ± SD) plotted as a function of frequency for

$Cib2^{+/+};Cib3^{KO/KO}$ ($n = 4$), $Cib2^{KO/+};Cib3^{KO/KO}$ ($n = 5$), $Cib2^{KO/KO};Cib3^{KO/KO}$ ($n = 4$). (**D**) OKN

gain and phase (mean ± SD) plotted as a function of frequency for $Cib2^{+/+};Cib3^{KO/KO}$ ($n = 4$),

$Cib2^{KO/+};Cib3^{KO/KO}$ ($n = 5$), $Cib2^{KO/KO};Cib3^{KO/KO}$ ($n = 4$). (**E**) VORl gain and phase (mean ±

SD) plotted as a function of frequency for $Cib2^{+/+};Cib3^{KO/KO}$ ($n = 4$), $Cib2^{KO/+};Cib3^{KO/KO}$ ($n = $

5), $Cib2^{KO/KO};Cib3^{KO/KO}$ ($n = 4$). Comparisons made with two-way ANOVA followed by

Bonferroni post-hoc test for (A-D); *p < 0.05. (**F**) Quantification of the number of rotations in

120 seconds (mean ± SEM), showing that, unlike $Cib2^{+/+};Cib3^{KO/KO}$, and $Cib2^{KO/+};Cib3^{KO/KO}$

mice, $Cib2^{KO/KO};Cib3^{KO/KO}$ mutant mice display a circling behavior and a vestibular defect (*t*-

test) ($n = 4$ for each genotype). (**G**) Quantification of the time mice remained on the rotating

rod with increasing acceleration (mean ± SEM). Comparisons made with two-way ANOVA

followed by Bonferroni post-hoc test for (F-K); *P<0.002. ($Cib2^{+/+};Cib3^{KO/KO}$ ($n = 6$),

$Cib2^{KO/+};Cib3^{KO/KO}$ ($n = 6$), $Cib2^{KO/KO};Cib3^{KO/KO}$ ($n = 6$)). (**H**) Comparison of power spectra

density of head movements in translational axes (top) and rotational axes (bottom) between

wild type ($Cib2^{+/+};Cib3^{KO/KO}$, black), heterozygous mutants ($Cib2^{KO/+};Cib3^{KO/KO}$, blue), and

homozygous mutants ($Cib2^{KO/KO};Cib3^{KO/KO}$, red). The double knockout $Cib2^{KO/KO};Cib3^{KO/KO}$

exhibit significantly higher power than $Cib2^{+/+};Cib3^{KO/KO}$ and $Cib2^{KO/+};Cib3^{KO/KO}$ across all

frequencies (0–30 Hz), in all six translational and rotational axes.

1352

1353

Figure 4. Vestibular hair cells do not have functional MET channels at rest in CIB2/3

double mutant mice. (**A**) Maximum intensity projections of confocal Z-stacks of the

vestibular end organs of P18 *Cib2^{KO/+};Cib3^{KO/KO}* and *Cib2^{KO/KO};Cib3^{KO/KO}* mutant mice

immunostained with myosin VIIa antibody (green) and counterstained with phalloidin (red)

and DAPI (blue). Scale bar = 50 μm. (**B**) Adult vestibular end organs of *Cib2^{+/+};Cib3^{KO/KO},*

Cib2^{KO/+};Cib3^{KO/KO} and *Cib2^{KO/KO};Cib3^{KO/KO}* mutant mice imaged after exposure to 3 μM of

FM 1-43 for 10 s. Vestibular hair cell bodies (green doted lines), and stereocilia bundles (red

1361   arrows) are shown. Scale bar: 100 μm. (**C**) FM 1-43FX (fixable form) labeling of saccules and

1362   utricles by IP injection at P60. Obvious reduction in signal in $Cib2^{KO/KO};Cib3^{KO/KO}$, and subtle

1363   extrastriolar reduction in $Cib2^{+/+};Cib3^{KO/KO}$ mice was observed. Scale bar = 100 μm.

1364

1365

1366

Figure 5. Both Cib2 and Cib3 are required in zebrafish for acoustic startle and MET function. (**A**) Compared to controls (siblings) *cib3* mutants have a normal acoustic startle response. *Cib2;cib3* double mutants completely lack an acoustic startle response, while *cib2* mutants have a reduced probability to startle compared to controls (*n* = 43 sibling, 20 *cib2*, 19 *cib3* and 7 *cib2;cib3* animals). (**B**) Compared to controls *cib3* mutants have a normal % of hair

cells that label with FM 1-43 per neuromast. In *cib2;cib3* mutants no hair cells label with FM

1-43, and in *cib2* mutants a significantly reduced % of hair cells label with FM 1-43 per

neuromast. (**C**) The average intensity of FM 1-43 labeling in *cib3* mutant hair cells is similar

to controls. In contrast, in *cib2* mutants, hair cells that label with FM 1-43, have a significantly

reduced intensity compared to controls ($n = 8$ neuromasts per genotype in B and C). (**D-G**)

Representative neuromasts labeled with FM 1-43 for each genotype. FM 1-43 label is overlaid

onto laser scanning DIC images. (**H**) Schematic showing the localization of membrane-

localized GCaMP6s (memGCaMP6s); the indicator used to measure Ca^{2+} influx into lateral-

line hair bundles. The plane used for Ca^{2+} imaging and to determine if a hair cell is

mechanosensitive is indicated by the dashed arrow. (**I-I'**) Representative example of a hair

bundle imaging plane (I), along with the resulting Ca^{2+} responses (I') from hair bundles from

a control neuromast. The ROIs used to measure Ca^{2+} signals in each hair bundle are indicated

in (I). (**J-L**) Representative examples of Ca^{2+} responses from neuromasts in *cib2* (J), *cib3* (K)

and *cib2;cib3* mutants (L). FM 1-43 imaging and behavior were performed at 5 dpf. Ca^{2+}

imaging was performed at 5 or 6 dpf. A Kruskal-Wallis test was used in A; A one-way ANOVA

was used in B and C. The scale bar in D and I = 5 μm. The online version of this article includes

source data and the following figure supplement(s) for Figure 5.

1389

1390   **Figure supplement 1.** Summary of mechanosensitive Ca^{2+} responses in zebrafish.

1391

1392

1393

Figure 6. Cib2 is required in specific subsets of hair cells in the zebrafish pLL and inner

ear. (**A**) Overview of Tmc requirements in pLL hair cells. Hair cells that sense posterior flow

(orange) rely on Tmc2b, while hair cells that sense anterior flow (blue) can rely on Tmc2b

and/or Tmc2a for mechanosensitive function. (**B-C'**) FM 1-43FX labeling in pLL neuromasts

from sibling control (B) and *cib2* mutants (C). The residual cells in *cib2* mutants rely on Cib3

for mechanosensitive function. Phalloidin label can be used to link FM 1-43FX label to the

orientation of pLL hair cells (C,C'; see colored asterisks). In B', orange and blue arrows rest

above hair cells that are oriented to respond to anterior and posterior flow, respectively. (**D**)

Quantification reveals that a similar percentage of anterior and posterior responsive pLL hair

cells label with FM 1-43FX in sibling controls (posterior flow: 49.7 %, anterior flow: 49.2 %,

n = 9 neuromasts). In contrast, in *cib2* mutants, a significantly higher percentage of posterior

responsive pLL hair cells label with FM 1-43FX in *cib2* mutants (posterior flow: 60.5 %,

anterior flow: 39.5 %, *n* = 26 neuromasts). (**E**) Overview of Tmc and Cib requirements in the

hair cells of zebrafish cristae. Short teardrop-shaped cells rely primarily on Tmc2a, while tall

gourd-shaped cells rely primarily on Tmc2b and Tmc1. (**F-I'**) Medial cristae for each genotype.

MemGCaMP6s labels all hair cells, while FM 4-64 labels hair cells with intact

mechanosensitive function. In siblings and *cib3* mutants, both teardrop and gourd cells label

with FM 4-64 (F-F', G-G'). In *cib2;cib3* double mutants, no hair cells label with FM 4-64

(asterisks) (I-I'). In *cib2* mutants many short cells fail to label with FM 4-64 (asterisks) (H-H'),

(*n* = 13 double het sibling, 7 *cib3*, 14 *cib2* and 4 double mutant crista). All images were acquired

at 5 dpf. A Kruskal-Wallis test was used in D; $P > 0.01$. The scale bars in B, B', and F' = 5

µm.

1416

1417

1418

Figure 7. AF2 predictions, NMR data, and simulations support a clamp-like model for the human TMC1 and CIB2 complex. (**A**) AF2 model of *hs* TMC1 in complex with *hs* CIB2. Side view of monomeric *hs* TMC1 (light purple) in complex with *hs* CIB2 (light sea green). N-terminal residues of TMC1 known to interact with CIB2 are in red. (**B**) Top left panel shows residues at the *hs* TMC1-NT fragment (115-117) shown to weaken interaction with CIB2 upon mutation (Liang et al., 2021) at the interface with CIB2. Additional panels show details of contacts between TMC1 and CIB2 with formation of K124:D14 and K112:E80 salt bridges, as well as hydrophobic interactions. (**C**) Overlay of ^{1}H-^{15}N TROSY-HSQC spectra of *hs* [U-^{15}N]-CIB2 (black) alone and bound to either only *hs* TMC1-IL (red) or both *hs* TMC1-IL and *hs* TMC1-NT (blue). (**D**) A complete simulation system of *hs* TMC1 in complex with *hs* CIB2-Ca^{2+} is shown in surface representation. *hs* TMC1 is shown in light (monomer A) and dark (monomer B) purple, *hs* CIB2 in light sea green, water in transparent light blue, lipid membrane

in gray lines, and K$^+$ and Cl$^-$ ions are shown as pink and green spheres, respectively. (**E**) The

hs TMC1-NT domain explores diverse conformational space in the absence of CIB during

equilibrium simulations (S3a). (**F**) Salt bridges K124:D14 and K112:E80 remain throughout

equilibrium simulations. (**G**) snapshot of amphipathic helix α0 inserting into the lipid bilayer

(S4c). (**H**) Number of K$^+$ crossings as a function of time for pores A and B of *hs* TMC1, *hs*

TMC1 + *hs* CIB2 + Ca^{2+}, and *hs* TMC1 + *hs* CIB2 in a POPC bilayer at -0.5 V, -0.25 V, and -

0.125 V. Insets show top views of pore region with TMC1 (dark purple) and lipids (light gray)

in surface representation (protein and lipids partially excluded for clarity). Representative

images of K$^+$ permeating the pore region between α4 and α6 of *hs* TMC1 are shown with

protein in surface representation and ions as pink spheres. Conductance values calculated for

K$^+$ crossings.

The online version of this article includes source data and the following figure supplement(s)

for Figure 7.

Figure supplement 1. Overview of TMC1/2 and CIB2/3 AF2 models.

Figure supplement 2. TMC sequence alignment.

Figure supplement 3. CIB2 sequence alignment.

Figure supplement 4. CIB3 sequence alignment.

Figure supplement 5. CIB sequence alignment.

Figure supplement 6. SEC of *hs* CIB2 and *hs* CIB3 either refolded alone or co-refolded with *hs*

TMC1-IL1, and *hs* TMC1-NT

Figure supplement 7. ^{1}H-^{15}N TROSY-HSQC spectra.

Figure supplement 8. Stability of TMC/CIB complexes, monomers, and subdomains.

Figure supplement 9. BSA in *hs* TMC1/2 and *hs* CIB2/3 complexes.

Figure supplement 10. Number of K$^+$ crossings as a function of time for pores A and B of *hs* TMC1

and *hs* TMC2.

Figure supplement 11. Conduction events in long timescale simulations and summary of predicted

conductance values.

Figure supplement 12. Possible combinations of TMC1, TMC2, CIB2, and CIB3 complexes.

Figure 7-video 1. Overview of *hs* TMC1 (light and dark purple) in complex with *hs* CIB2
(light sea green)-Ca^{2+} (green spheres). K$^+$ ions (pink) were shown to be permeating the pore
region between α4 and α6. Water molecules, lipids, and protein side chains were omitted for
clarity.
1463

1464   **Figure 7-video 2.** Side view of the *hs* TMC1 channel pore during a trajectory at -0.5 V
1465   (monomer B). K$^+$ (pink) are seen permeating the pore from one side (S1b); *hs* TMC1 (monomer
1466   B) is shown in dark purple and *hs* CIB2 is shown in light sea green. Water molecules, lipids,
1467   *hs* TMC1 (monomer A), and protein side chains were omitted for clarity.
1468

1469