ADORE: A Benchmark Dataset for Machine Learning in Ecotoxicology

Christoph Schür[1,†,*], Lilian Gasser[2,†], Fernando Perez-Cruz[2,4], Kristin Schirmer[1,3,5], and Marco Baity-Jesi[1]

[1]Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland
[2]Swiss Data Science Center (SDSC), Zürich, Switzerland
[3]ETH Zürich: Department of Environmental Systems Science, Zürich, Switzerland
[4]ETH Zürich: Department of Computer Science, Zürich, Switzerland
[5]EPF Lausanne, School of Architecture, Civil and Environmental Engineering, Lausanne, Switzerland
[*]corresponding author: Christoph Schür (christoph.schuer@eawag.ch)
[†]these authors contributed equally to this work

ABSTRACT

The use of machine learning for predicting ecotoxicological outcomes is promising, but underutilized. The curation of data with informative features requires both expertise in machine learning as well as a strong biological and ecotoxicological background, which we consider a barrier of entry for this kind of research. Additionally, model performances can only be compared across studies when the same dataset, cleaning, and splittings were used. Therefore, we provide *ADORE*, an extensive and well-described dataset on acute aquatic toxicity in three relevant taxonomic groups (fish, crustaceans, and algae). The core dataset describes ecotoxicological experiments and is expanded with phylogenetic and species-specific data on the species as well as chemical properties and molecular representations. Apart from challenging other researchers to try and achieve the best model performances across the whole dataset, we propose specific relevant challenges on subsets of the data and include datasets and splittings corresponding to each of these challenge as well as in-depth characterization and discussion of train-test splitting approaches.

Background & Summary

Regulation of chemicals aims to protect both human health and the environment, with ecotoxicological research focusing on the latter. Regulatory hazard assessment relies on extensive animal testing, *e.g.,* in the European Union, acute fish toxicity testing for chemicals of a manufacturing or import volume > 10 tons p.a. is required by the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) legislation[1]. A recent pre-print estimates the global annual usage of fish and birds to range between 440,000 and 2.2 million individuals at a cost upwards of $39 million p.a.[2]. With over 200 million substances in the archive of the Chemical Abstracts Service (CAS, www.cas.org) and more than 350,000 chemicals and mixtures currently registered on the market worldwide[3], chemical hazard assessment is a Herculean task.

These ethical and financial considerations are incentives for alternatives to animal testing, which include computational (*in silico*) methods. A long established *in silico* method is Quantitative structure-activity relationship (QSAR) modeling. It aims to predict biological and chemical properties of compounds based on linear or nonlinear relationships of their molecular structure and experimentally determined activities and properties[4]. The increase of computational capacities and accessibility of educational resources and software led to the adoption of machine learning (ML, computational methods that continually improve their prediction without additional outside intervention) in ecotoxicology over the past decade. Published studies on predicting ecotoxicity are for the most part limited to aquatic systems. They are using animal-derived (*in vivo*) data and encompass supervised learning approaches: classification of chemicals into toxicity brackets, and regression-based prediction of concrete toxicity values[5–7]. The latter more closely resembles the outcome of ecotoxicity experiments, where specimens are exposed to multiple concentrations of the same substance and an effect is recorded at certain time intervals. A non-linear regression model is then fit to the data to derive toxicity values.

Biological and ecotoxicological expertise is required to guide informative feature selection and data cleaning prior to ML modeling, and to help interpret results[8]. In ML-based research, insufficient reporting and data availability hinders reproducibility and comparison across studies and, thus, objective judgment of model performances[9]. Additionally, the way a dataset is split into training and test data immensely affects model performance and can inflate it through data leakage (*i.e.,* the model is trained and tested on similar data)[10].

Ultimately, realistically assessing and comparing model performance among different works is only feasible when training and testing were performed on the same dataset, using the same data cleaning and splitting strategy. This is fundamental to ML-based research and demonstrated by the plethora of publications associated with long-standing benchmark datasets, such as CIFAR and ImageNet and many others (for example, in the NeurIPS benchmark data track). Compared to NeurIPS, environmental sciences are slowly progressing towards adopting the best practices in the field of ML, including the availability of standard datasets, with a notable pioneer being the CAMELS dataset from hydrology[11] with its recent evolution to CARAVAN[12]. Indeed, to serve as benchmark requires a well-characterized, expert-curated and -described, freely available dataset, which we provide with this work. The focus of our dataset is mortality and related endpoints for the three taxonomic groups fish, crustaceans, and algae, extracted from the ECOTOX database[13]. The dataset is supplemented by chemical, phylogenetic, and species-specific features that can be used in modeling toxicity outcomes or in addressing other questions.

Here, we describe the data sources and the reasoning behind data selection and processing steps. We provide defined splits of the data, based on chemical occurrence and molecular scaffolds. We propose concrete challenges to the community, including extrapolation across taxonomic groups, to learn more about the potential and limitations of ML in ecotoxicology.

Methods

The dataset is assembled from different sources. The core contains ecotoxicology data, *e.g.,* toxicity values and experimental conditions. These are expanded with species-related and chemical data (Fig. 2). Data sources, filtering and other processing steps are described in the following subsections. The section "Data records" is focused on the different files that contain subsets of the data, whereas data characteristics are shown in the Section "Technical validation". A glossary of all features is given in the Annex Table 8.

Ecotoxicology data
Core dataset on acute mortality

The main source of our dataset is the ECOTOX database (https://cfpub.epa.gov/ecotox/) from the United States Environmental Protection Agency's (US EPA). ECOTOX is updated on a quarterly basis. Our dataset is based on the release from September 2022, which contains over 1.1 million entries of more than 12,000 chemicals and close to 14,000 species[13]. The EnviroTox database[14] is similar to our dataset, but without the additional chemical- and species-specific features which we curated with ML modeling in mind, and without a standardization that allows straightforward comparability between studies. We had to evaluate a trade-off between a more noisy dataset encompassing more chemical and organismal diversity, and a substantially smaller, but cleaner dataset that is not as chemically and taxonomically diverse.

We include entries for three distinct taxonomic and trophic levels of importance in aquatic ecotoxicology: fish, crustaceans, and algae. These taxonomic groups represent some of the biggest data subsets available in ECOTOX by covering 41% of all the entries.

ECOTOX distinguishes between effects and endpoints. Effects are observed responses resulting from chemical exposure, ranging from mortality (MOR), *i.e.,* death of the specimen, to more subtle effects like behavioral changes, and gene regulation. Endpoints are the targets which can be numerically determined, *e.g.,*, LC50 and EC50. The use of the lethal concentration 50 (LC50), the concentration at which 50% of specimen died after a certain time period, is very common and allows for comparison of toxicity across chemicals and species. In aquatic ecotoxicology, the LC50 refers to the concentration of a substance in the test medium water surrounding a species[15]. For effects that are not lethality-based, toxicity is commonly described as the effective concentration 50 (EC50), that analogously describes the induction of a 50% effect level, often compared to a negative and/or positive control treatment. Both can be expressed in a mass-based (*e.g., mg/L*) and a mol-based (*e.g., mol/L*) manner. The former is easier to correlate to environmental concentrations, while the latter is more biologically informative, assuming that a molecule is the entity that induces the biological response on a molecular level.

Our dataset focuses on short-term lethal (acute) mortality and we included other endpoints that are comparable to mortality (Table 1). Effects manifest differently in different taxa. Mortality was the sole effect chosen in fish, based on standardized test guidelines such as the Organisation for Economic Co-operation and Development (OECD) Test guideline 203[16]. For crustaceans, mortality and immobilization, the latter categorized in ECOTOX under intoxication (ITX), are used interchangeably in the respective test guidelines, *e.g.,* OECD Test guideline 202[17]. For algae, the focus is rarely on the individual, but often related to population health/size as a proxy for toxicity. Population growth is the most commonly used approach to testing effects in algae, *e.g.,* OECD guideline 201[18]), which led to the inclusion of data related to effects encoded as mortality, growth (GRO), population (POP), and physiology (PHY) (Table 1).

The standard observational period for fish acute mortality is 96 hours according to OECD guideline 203[16], where mortality is recorded every 24 hours. For crustaceans, the typical experimental period is 48 hours, *e.g.,* given in OECD guideline 202[17], while algae are commonly exposed over 72 hours[18]. Only experimental periods up to 96 hours were included.

Since, ultimately, the goal would be to replace animal testing, we excluded *in vitro* endpoints, *i.e.,* assays based on isolated cells[19], and assays based on early organismal life stages, such as eggs and embryos[20]. They comprise alternative test methods to animal testing[21]. Removing eggs and embryo tests is done under the limitation that most of the entries have uninformative values for the organism life stage and that life stages for the three taxonomic groups are hardly comparable (see Figure 10E).

Processing of the ECOTOX database

The core dataset was assembled from the downloadable, pipe-delimited ASCII version of the ECOTOX database. It is made available as a set of .txt files. We used the files for species, tests, results and media that contain information on the taxonomy of species, experimental-design-related factors, experimental outcomes, and media characteristics, respectively. The entries of these files can be matched to each other using different unique keys, *e.g.,* the test species is encoded with the key *species_number* that can be used to match the taxonomic information with the test table. Each data point is uniquely identified by the *result_id*.

Chemicals are identified through different descriptors, most prominently the CAS number, the DSSTox Substance ID (DTXSID), associated with the CompTox Chemicals Dashboard by the US EPA, and the IUPAC International Chemical Identifier (InChI) and the associated InChIKey by the International Union of Pure and Applied Chemistry[22]. These identifiers are retained in the dataset to ensure high compatibility with other data sources and retraceability of all data points. If possible, the InChIKey was used for matching, followed by DTXSID and CAS. Additionally, the dataset contains Simplified Molecular Input Line Entry System (SMILES) codes that represent the chemical structure in a string format[23]. Like others[14, 24, 25], we use the canonical SMILES, from PubChem[26, 27], which do not capture isomerism.

In the first step of the processing pipeline (Fig. 2), the ECOTOX files were separately harmonized and pre-filtered.

Species file The species files contains information on the test species. We removed all data points which had missing values for the species group (column `ecotox_group`) or the taxonomic columns class, order, family, species and genus. We then retained only entries for the three taxonomic groups of interest. This was done by filtering the species group entries to contain either "Fish", "Crusta", or "Algae".

Tests file The tests file contains information on the different descriptors of experimental conditions. *Exposure type* (`exposure_type`) refers to the method by which contact between the organism and the chemical is achieved. We only retain the exposure types that are most common and ensure consistent experimental designs[1]. *Media type* (`media_type`) is the liquid, in which the experiment was carried out. This feature was reduced to only contain fresh water and salt water experiments,

---

[1]Static (S, "Toxicity tests with aquatic organisms in which no flow of test solution occurs; solutions may remain unchanged throughout the duration of the test"), flow through (F, "continuous or very frequent passage of fresh test solution through a test chamber with no recycling"), renewal (R, "A test without continuous flow of solution, but with occasional renewal of test solutions after prolonged periods, e.g., 24 hours"), and not reported aquatic experiments (NR). These definitions are from the ECOTOX term appendix).

also with the intention to remove tests conducted in environmental water samples that are basically non-reproducible and could have an effect by themselves.

The tests table contained other features which we retain but do not deem informative for modeling purposes. For *control type* (`control_type`), many of the data points are either not encoded or it is nontransparently reported what the encoding entails. The *test location* (`location`) was "laboratory" for more than 98% of the data points. *Application frequency unit* (`application_freq_unit`), describing how often the chemical was applied/renewed in the system, had such a high variability that it was not possible to normalize to a common unit. Most of the *organism life stage* (`organism_lifestage`) levels are uninformative.

Results file   The results file contains the information related to each result, *e.g.,* endpoint, effect, and concentration.

Mostly, *concentrations* (`conc1_mean`) are given as specific EC50 values. For limit tests or if the EC50 was outside the tested concentration range, the values are only given as above or below a concentration range (`conc1_mean_op` is either $<$, $\leq$, $\geq$, $>$). These values were removed. Other entries do not contain a mean effect value but a minimum and a maximum value (`conc1_mean_min` and `conc1_mean_max`, respectively). We averaged these two values if they were within one order of magnitude.

Effect concentrations are given in a variety of units (`conc1_mean_unit`). We only kept the units related to a mass or molar concentration and unified them to *mg/L*, from which we calculated the molar concentration in *mol/L*.

Similarly, *experimental durations* (`obs_duration_mean`) are given in a number of different units. We converted them to hours and retained 24, 48, 72, and 96 hours experimental periods to cover highly standardized short-term toxicity scenarios (Figure 10B). Generally, during a 96 hour experiment effects are also recorded at the previous time points. So, often, later time points are an extension of previous ones, *i.e.,* repeated measurements in the same test. We checked several repeated tests and could not definitively determine that with similar experimental properties later measurements are consistently more or less sensitive than previous ones, even though this may be disguised in the general data variability. Thus, we included all such data points. If needed, one can filter to the latest data point for an experiment which is uniquely identified by the reference number.

For the *concentration type* (`conc1_type`), we retained all the entries with the levels "active ingredient" (A), "formulation" (F), and "total" (T). Chemicals, *i.e.,* the active ingredient that is of interest in toxicity testing, are often combined with other substances depending on their use case to delay or enhance their activity or facilitate transport to the target area/organ, *e.g.,* into a plant or adhering to a surface. We could not conclusively deduce that either of the concentration types is consistently more toxic than another and thus decided to include this information into our model dataset (see Table 8).

Media characteristics file   The media characteristics file contained temperature (`media_temperature_mean`) and pH (`media_ph_mean`) values for some of the experiments. We removed or corrected unrealistic values (see next paragraph), calculated the average values if only minimum and maximum value were available, and converted all temperatures to °C.

Manual filtering of unrealistic values   The original dataset contains several values that required manual evaluation and cleaning/correcting. The manual filtering is documented in the code repository https://renkulab.io/gitlab/mltox/adore in the file *src/preprocessing.py* using the functions *remove_and_correct_entries* and *correct_ph_temp_entries*.

Some toxicity values were found to be either very high or very low, *i.e.,* EC50/LC50 $\geq 10^5$ *mg/L* and $\leq 10^{-5}$ *mg/L*. These values were manually evaluated by checking against the original source. If the original source was not accessible, the values were checked against comparable data in the ECOTOX database, ideally from the same chemical and the same species. If that was not available, they were checked against the same chemical and a species from the same taxonomic group. If that was also not available or the values differed by several orders of magnitude, the data point in question was removed. Some data points were excluded because the toxicity values were obtained in multi-stressor scenarios, *e.g.,* testing chemical sensitivity in concert with salinity tolerance of a species or under non-representative conditions, *e.g.,* application of a chemical in orange juice as carrier medium[28]. We concluded that these toxicity values are not representative of the chemical's toxic potential. We are aware that experimental setups like these likely make up part of the rest of the dataset as well and that they contribute to the high variability of effective concentrations (Figure 1), but manually checking all values is not feasible and would negate the use of an already assembled dataset, such as the ECOTOX database.

Similarly, we found unrealistic values for the experimental properties pH of the medium and temperature, *i.e.,* pH values smaller than 4 and larger than 10, and temperatures exceeding 50 °C. Again, we checked the values manually against the original literature and excluded cases where (waste)water samples with extreme pH were the test medium and where the pH was strongly influenced by the test substance. For temperature, we employed expert judgment and concluded that the values were mostly conversion errors, *e.g.,* unit given as °C, when actually °F, or decimal errors, *e.g.,* 250 °C when actually 25.0 °C, and adjusted the values accordingly.

Theoretically, it should be impossible to derive an EC50 value larger than the chemical's water solubility. Nonetheless, a number of data points contain EC50 values that lie above the given water solubility of the compound. This can partly be explained by most water solubility values being based on predictions or derived from experiments based on ultrapure water,

while exposure media already contain several micronutrients and salts. One option on how to deal with these entries is to remove all data points for which the effective concentration is higher than the chemical's water solubility[29]. Alternatively, as was done for the EnviroTox database, entries where the toxicity value exceeds the water solubility by a factor of 5 are specifically marked, but not removed[14]. The EC50 values in our dataset are calculated from the nonlinear regression on the concentration-response-curves and therefore themselves come with inherent uncertainty. Accordingly, all data points are uncertain to one degree or another, which in this case only becomes apparent for the data points where the EC50 exceeds water solubility. Therefore, we decided against applying this filtering step.

Species-related data

We include two kinds of species-related data, species-specific ecological, life-history, and "pseudo"-data related to the DEB modeling framework and phylogenetic distances between species.

Add my Pet

Species-specific data was acquired from the Add my Pet collection (AmP), a manually curated database linked to the dynamic energy budget (DEB) theory framework[30], that contains information for more than 3,400 animals (August 2022). The DEB theory is based on heterotrophs, *e.g.,* animals like fish and crustaceans, and does not cover autotrophs such as algae. Therefore, comparable data on algae is not included in our dataset. We downloaded the AmP collection (dated November 25th, 2022) and extracted information for the species present in the core dataset. This includes three types of information: Ecological data, life history data, and pseudo-data. See Table 2 and Figure 11 for a feature overview.

Ecology   This data category includes information about ecological peculiarities and lifestyle of a species. We selected climate zone and ecozone, *i.e.,* the climatic and geographical area, respectively, where the species occurs (Tables 10 and 11). Additionally, we selected migratory behavior (Table 12, *e.g.,* anadromous/catadromous species that migrate up-/downstream for reproduction, and food type, *e.g.,*, carnivorous, herbivorous, parasitic, *etc.* (Table 13).

For three features, we reduced the complexity of the original encoding. The AmP collection provides 51 different levels for climate zone and 56 for ecozone, respectively. Similarly, the food encoding has 77 different levels. These features follow a hierarchical encoding, which was reduced by retaining only the top level. Distributions for the species-specific features can be found in Figure 11. Further information on the encodings can be found in the Annex (Tables 10-13) and on the AmP homepage.

Life history   The AmP collection also provides life history data, which are usually taken from literature, but sometimes estimated or extrapolated from related species. As a consequence, some features are not available for each species and the resulting dataset contains missing values. We included the features which had the highest coverage of number of data points with the core dataset: life span, body length at birth, body length at puberty, ultimate body length, and maximum reproductive rate (see Table 2 and Figure 11). Life span and ultimate body length are available for all species.

Pseudo-data   Pseudo-data are species-specific parameters related to the modeling framework that the AmP collection forms the basis of. We included the three pseudo-data features showing reasonable variability across species: allocation fraction to soma[31], energy conductance, and volume-specific somatic maintenance cost (Table 2; Figure 11). These features are available for every species included in the AmP database. Definitions are given on the DEB-portal and in Kooijman *et al.* (2008)[32].

Phylogenetic distances from TimeTree

Since closely related species often are similarly sensitive to the same chemicals compared to distantly-related species[33], we include the phylogenetic distance between species as possible modeling features. The phylogenetic distance represents how closely related a pair of species is to each other by the time since they diverged from the last common ancestor[34]. We used TimeTree Version 5 to produce a phylogenetic tree of the species in our data. The tree, visualized in Figure 6D, was then transformed into a phylogenetic distance matrix using the Python package dendropy[35].

Chemical data

We include several chemical features of which some are intended as modeling features, namely chemical properties and molecular representations. The others should only be used to help interpret modeling results, namely the ClassyFire ontology and the functional use categories.

Chemical properties

We added basic chemical properties from DSSTox[36], the PubChem database[26] and through the Python package RDKit[37]. The properties are molecular weight (MW in g/mol), water solubility (WS in mg/L), melting point (MP in °C), number of heavy atoms (*i.e.,* all atoms excluding hydrogen), number of bonds, double bonds, and triple bonds, number of ring structures, and the number of OH groups. Additionally, we included the octanol-water partition coefficient (LogP, sometimes referred to as

$LogK_{ow}$, the logarithm of the relative solubility of a compound in the solvents octanol and water[38], positive/higher values, indicate a higher lipophilicity of a compound), and the acid dissociation constant (pKa).

The pKa regulates at which pH a substance dissociates. Since, for some compounds, the pH of the medium influences the toxicity[39], it is reasonable to include the pKa as a feature to be used together with the medium's pH. The pKa values were obtained through the OECD QSAR toolbox (V4.5)[40] via the SMILES. When several predictions were available, we used their median. Median values above 30 were excluded.

Molecular representations

Presenting chemical structures to a ML model such that it can be used in an informative manner poses a challenge. Standard molecular formats such as connection tables and SMILES are not valid input for most ML models. Chemicals need to be provided in a form "understandable by an algorithm" which Staker *et al.*[41] refers to as a molecular representation. We used three well-established molecular fingerprints[42]: MACCS[43], PubChem[44] and Morgan[45] fingerprints as well as the mol2vec embedding[46].

Molecular representations can be divided into non-learned and learned representations[41]. Besides computable features *i.e.,* chemical properties, which we introduced in the previous section, molecular fingerprints are widely used non-learned representations. Fingerprints are fixed-length, binary vectors where each bit indicates presence ("1") or absence ("0") of a substructure. MACCS, consisting of 166 publicly available bits, and PubChem, consisting of 881 bits, are substructure keys-based, *i.e.,* each bit corresponds to a predefined substructure[2]. The Morgan fingerprint is a circular fingerprint based on all fragments that can be extracted within a bond radius around each atom of the molecule. These fragments are then hashed to create fingerprints comparable across chemicals of different size. We used RDKit[37] to calculate the MACCS and Morgan fingerprints for all chemicals in the dataset. The Morgan fingerprint was calculated with a radius of 2 bonds and hashed to 2048 bits. The PubChem fingerprint was extracted as CACTVS fingerprint using PubChemPy[27].

The fourth representation is learned. Jaeger *et al.* developed the molecular embedding mol2vec, which applies the word embedding word2vec[47] on substructures derived from the Morgan algorithm. This embedding treats substructures as if they were words, and molecules as if they were sentences[46]. Jaeger *et al.* trained a 300 dimensional embedding on 19.9 million compounds, which is available as a Python package via https://github.com/samoturk/mol2vec and was used to calculate 300-dimensional feature vectors for all chemicals in the dataset.

Chemical ontology from ClassyFire

Chemical structures can be organized into a taxonomical system through the standardized classification system ClassyFire[48]. This information can either be retrieved using the ClassyFire API or the Python package pybatchclassyfire based on InChIkeys. We consider this feature more useful for the interpretation of modeling results than for the modeling itself.

Functional use categories

The US EPA provides a Chemicals and Products Database (CPDat) containing product applications and functional uses for various chemical compounds (December 2020 Release). The database provides functional uses in two variants. The OECD function is an internationally harmonized functional use category[49], which is suitable for most chemicals in our dataset except for the category "biocide". Biocides can be subdivided in insecticides, herbicides, fungicides, antimicrobial agents, and others, which is relevant for our dataset. This information is contained in the reported functional uses, which are unharmonized data entries. Please refer to the official documentation[49] for definitions of the OECD functions.

We processed the data by combining closely related OECD functions as indicated by the description[49], *e.g.,* dye and pigment are renamed to colorant. The reported functional uses are harmonized in several steps: All entries are set to lower case and unnecessary white space is removed. Non-informative entries such as "not reported", "proprietary", or "impurity" are removed. The other entries are unified where possible, *e.g.,* entries starting with "herbicide" are renamed to only "herbicide", or entries containing the term "catalyst" are renamed to "chemical reaction regulator" as is the name of the corresponding OECD function. Then, reported functional uses longer than 30 characters or occurring less than four times are removed. This results in 43 OECD functions and 174 reported functional uses. For most chemicals, there are several OECD functions and reported functional uses. We provide all of them as well as only the most prevalent ones, *i.e.,* the uses making up more than 20% of the entries for a compound.

The functional use categories should *not* be included in modeling, but only be used to enable a better understanding of modeling results. In fact, its inclusion would constitute a form of data leakage, since, *e.g.,* labeling a substance group as biocides would correlate well with higher toxicity towards certain species, but the information was not transmitted through a chemical feature but through this label.

---

[2]See these links for the definition of each MACCS bit and PubChem bit.

Further data processing

Overlap between data categories

The data categories, described in the previous sections, did not necessarily contain a feature value for every data point, *e.g.,* the ECOTOX data contains 1,690 species and 4,775 chemicals after processing but only data for 665 and 270 of those species are available in TimeTree and AmP, respectively.

This leads to a trade-off between retaining more data points and expanding the feature space. The longest version of the dataset contains more data points but has fewer features, whereas the dataset with the highest number of features retains a far smaller number of data points. Since the bottleneck for environmental risk assessment models is usually the scarcity of data, we prefer a bigger dataset to a larger feature space and therefore provide a long core dataset.

Apart from using ML to predict toxicity, we envision our dataset being used for the exploration of other ecotoxicologically relevant research questions. Therefore, we provide incomplete data tables, *i.e.,* containing missing values for some features, which are intended for future studies that are not tied to the challenges proposed and discussed in this paper.

Filter by taxonomic group and effect group

After all input data was harmonized and pre-filtered, and taxonomic information as well as chemical properties were added, we filtered the data to only contain acute mortality experiments for the three taxonomic groups fish, crustaceans and algae. For each taxonomic group, this encompasses the corresponding effect groups (Table 1). The endpoints were filtered to only include LC50 and EC50.

Final filtering

The processed data on acute mortality was then filtered to obtain a dataset suitable for modeling. To employ the mol2vec embedding and compare it to the other molecular representations, we only keep compounds which are compatible with it[46]. Also, we only keep compounds for which water solubility is available as we deem it a potentially important modeling feature. Additionally, we remove data points without phylogenetic and AmP information. These filtering steps remove around 50% of the processed data and leaves us with a dataset comprised of predominantly organic chemicals.

Data splittings

Data splitting, *i.e.,* the generation of training and test data subsets, and of cross-validations folds, can greatly affect model performance. Possible causes are the inherent variability of the data itself and (non-obvious) data leakage. The latter describes a case where information about the target is introduced which should not legitimately be used for modeling[50]. Data leakage has also been described as a "spurious relationship between the independent variables and the target variable"[10] and leads to overestimated model performance[50].

We discuss different data splitting schemes, of which most account for some sources of data leakage. Ideally, a combination of chemical and species would only appear in either the training or the test set. Since our dataset is skewed both with respect to chemicals and species, splits by chemical *and* species would lead to highly imbalanced subsets and are not feasible. We can only reasonably apply a split by either chemical or species, *i.e.,* all data points belonging to a chemical or species, respectively, are found in either the training or the test set. We focus on splits by chemical, due to the higher variability in chemicals compared to species and because extrapolating across chemicals allows us to address the problem of predicting toxicity of untested or yet unknown chemicals.

Split totally at random   The simplest train-test-split can be achieved by random sampling of data points, provided by us as *totally random*, which has been the main approach in previous work applying ML to ecotoxicology, and generally suffices for a well-balanced dataset without repeated experiments[29,51,52]. For our dataset with repeated experiments, *i.e.,* data points coinciding in chemical, species, and experimental variables, see Fig. 1, the *totally random* approach has a high risk of data leakage and the associated overestimated model performances. With this splitting, the same chemical is likely to appear in both training and test set.

Thus, we present stratified splits as alternatives which ensure that chemicals are not shared between training and test set, and not between cross-validation folds. We provide splits for 5-fold cross-validation.

Splits by chemical compound   The straightforward approach is to split by chemical compound, ensuring that a chemical is put into either the training or the test set. We provide two variants here, where compounds are split at *random by chemical compound* or *by occurrence of chemical compounds*. For the latter, compounds are sorted by the number of experiments performed on them, *i.e.,* those with most experiments at the top. Then, the first five compounds are put into the training set and the sixth is put into the test set. This is repeated with the subsequent compounds until all are distributed. The five cross-validation folds are filled accordingly, *e.g.,* the most common compound goes to fold 1, the second most common to fold

2, and so on. With both variants, it is likely that similar chemicals are shared between the training and test set, and between the cross-validation folds.

Splits by molecular scaffold   Molecular scaffolds can be used to categorize molecules based on their core structure, *i.e.,* the molecular backbone, notwithstanding side chains and functional groups. Scaffold splitting has been used in drug discovery[53] and was deemed to lead to more optimistic modeling outcomes than relying on random splitting[54]. We utilize two scaffold types: the Murcko scaffold and the generic scaffold. The Murcko scaffold reduces a molecular structure to ring systems and the structures connecting these ring systems[55]. The generic scaffold goes one step further by replacing all heteroatoms in the Murcko scaffold with carbon atoms and all bonds with single bonds. Both scaffold types were calculated using RDKit[37].

For both scaffold types, we provide several data splits. Scaffolds are put at random into either training or test set (*scaffold-murcko* and *scaffold-generic*). Here, we follow the procedure described by Yang *et al.* and put molecules whose scaffold is more common than half the test size into the training set[53]. The others are distributed at random between training and test set. Additionally, we provide two leave-one-out (LOO) splits where we put either the most common scaffold (*scaffold-murcko-loo-0* and *scaffold-generic-loo-0*) or the second most common scaffold (*scaffold-murcko-loo-1* and *scaffold-generic-loo-1*) in the test set and the others in the training set. Also, we include a "leave-last-out" (LLO) alternative, where we put the least common scaffolds in the test set, until the test size is reached, and the remaining ones in the training set (*scaffold-murcko-llo* and *scaffold-generic-llo*). The last splitting scheme represents a generalization to new compounds with more exotic structures. We provide several splits by both scaffold types since this is an area of active research and it is not yet known when and if splits by scaffold should be used.

We provide 11 splits, including cross-validation folds if applicable, for the data subsets where we train and test on the same taxa. For the data subsets where we train on crustaceans and algae and test on fish, we do not provide splits by molecular scaffold. Cross-validation is applicable for *totally random*, *random by chemical compound*, *by occurrence of chemical compounds*, *scaffold-murcko* and *scaffold-generic* but not for the LOO and LLO options. Since the split *by occurrence of chemical compounds* puts one sixth, *i.e.,* 17%, of the data points in the test set, we maintain the associated ratio of 83:17 for all train-test-splits to have comparable sizes across the data subsets.

Data Records

The dataset will be made available on ERIC, the institutional data repository of Eawag under https://doi.org/10.25678/0008C9. For the reviewing process, it is temporarily available on https://figshare.com/s/3a3e4dc26cca88e8ea11. In the final dataset, there are three folders (*processed*, *chemicals*, *taxonomy*). All files are provided as csv. All data files, including the raw files, are also part of the code repository https://renkulab.io/gitlab/mltox/adore.

Intermediate files as well as the final files, one for each prediction challenge, can be found in the folder *processed*. We include the intermediate files such that the user can follow the processing steps. They encompass the processed separate files from ECOTOX for species (*ecotox_species.csv*), tests (*ecotox_tests.csv*), results (*ecotox_results.csv*), as well as compiled chemical properties (*ecotox_properties.csv*) in the subfolder *before_aggregation*. These files are then combined to get *ecotox_mortality_processed.csv* and then filtered to get *ecotox_mortality_filtered.csv*.

For each challenge, we provide a data subset which starts with the name of the challenge as given in Tables 3-5 and ends with *_mortality.csv*. These datasets contain all information needed for modeling and further columns to uniquely identify and retrace each entry (Table 8).

Additionally, we provide two separate files in the folder *chemicals*, the chemical ontology (*classyfire_output.csv*) and the functional use categories (*functional_uses_output.csv*). They can be matched with the data in the challenges files using the InChIkey or the DTXSID, respectively. The file *tax_pdm_species* in the folder *taxonomy* contains phylogenetic distance information which can be used for modeling.

Technical Validation

Here, we describe the dataset and its features. Information on how the dataset was compiled can be found in the "Methods" section. We provide several data subsets related to the proposed prediction challenges presented in the "Usage Notes" section. The figures and numbers given in this section refer to the comprehensive file *a-FCA2FCA_mortality.csv*.

Ecotoxicological Data
Toxicity
The outcome of ecotoxicological experiments is a continuous value, in our case an EC50 value, or LC50 for mortality. The distributions of the different effects for the three taxonomic groups are shown in Figure 4A. The overall spread of EC50 values is comparable across effect groups, especially for fish and crustaceans. Physiology, only recorded in algae, has a bias towards

low EC50 values, which is reflected in Figure 4C, where the distribution across toxicity classes is comparable to that across taxonomic groups (see Table 7), while few compounds were non-toxic to algae. This could entail a bias in the compounds tested on algae, either with the expectation of high toxicity or because algae are more sensitive than the other taxonomic groups. Similarly, since the database has grown over time, it could hint to historical shifts related to the inclusion of algae as model organisms. Figure 9 confirms that the first reported tests on algae was done in 1974, 28 years and 10 years later than for fish and crustaceans, respectively. Figures 4 B and C visualize the distribution of data points in our dataset across the three taxonomic groups for either a binary or a multi-class classification, respectively.

Repeated experiments and reproducibility

As we are dealing with highly variable *in vivo* data, experiments with the same experimental conditions can lead to differing results. The dataset contains a large fraction of "repeated experiments", *i.e.,* data points that overlap in species, chemical, and the experimental parameters effect type, media type, observation duration, exposure type and concentration type. As a consequence, our modeling target EC50 rather resembles a probability distribution than a singular ground truth. We show this in Figure 1A with a histogram of the number of data points sharing a parameter combination. Most experiments have only been done once or repeated a few times. There are three experimental settings that have been repeated around 80 times each: the pesticide Trichlorfon (CAS number 52-68-6) has been tested on the rainbow trout (*Oncorhynchus mykiss*) 83 times while the insecticide Methoxychlor (72-43-5) has been tested on the brook trout (*Salvelinus fontinalis*) over an experimental duration of 24 and 96 hours, respectively, 79 and 82 times. Some of the experimental outcomes vary up to several orders of magnitude, even within this restrictive set of parameter values, as can be seen in Figure 1B, which shows experiments repeated at least 25 times. This variability needs to be considered when evaluating model predictions originating from this data.

Experimental features

Experimental features like media type and observation duration may influence the experimental outcome. See Figure 10 for an overview of the features and their levels.

Species-related data

In total, there are 203 species across all taxononomic groups of which fish make up around two third (Table 7). A few species have been tested much more often than most, causing the dataset to be skewed with respect to species. For fish, those are rainbow trout (*Oncorhynchus mykiss*), fathead minnow (*Pimephales promelas*), and the bluegill sunfish (*Lepomis macrochirus*) (Figure 6A). By far the most tested crustacean species is the waterflea *Daphnia magna*, followed by another water flea species, *Daphnia pulex* (Figure 6B). For algae, the microalgae *Chlorella vulgaris* and the green algae *Chlamydomonas reinhardtii* are the most abundant species in the dataset (Figure 6C).

Generally, apart from the dominance of a number of model species, the diversity of organisms included in the dataset is high and many taxonomic sublevels are present. A dendrogram of the species visualizing the phylogenetic distances between species are shown in Figure 6D. The length of lines after branching indicates the distance between species. This is most striking in the early branching of the algae from the two animalia groups fish and crustaceans (black line). Even though we consider it a good indicator of species relatedness, the used needs to consider that the phylogenetic distance matrix cannot straightforwardly be used with every type of model.

Other species-related data are focused on ecology, life-history and pseudo-data from the DEB theory framework, originating from the AmP database. The distributions of these species-specific features are shown in the Annex in Figure 11. They are only available for fish and crustaceans. Many species are found in several climate and ecozones, or accept different kinds of food. Therefore, we included all these combinations of levels in the data. The level abbreviations are divided by an underscore and are meant to be reformatted into multi-hot-encoded matrices for modeling use. Definitions for the encoding levels are provided in the Tables 10-13. The life history data and the pseudo-data are given in Figure 11B. The histograms are dominated by the variability in fish species whereas crustaceans cover smaller ranges. This is more prominent for the life history features, body length at three stages and reproductive rate, than for the pseudo-data features. The dataset contains many fish species of which some grow large, whereas the covered crustaceans all belong to fairly small-bodied taxonomic orders.

The importance of these parameters for modeling is difficult to foresee but we are positive that information distinguishing species is valuable to at least partly explain variation in the response of animals.

Chemical data

In total, the dataset encompasses ecotoxicological tests on 2,411 chemicals. Substances are routinely tested on several species and taxonomic groups, which is reflected in the Venn diagram in Figure 3A. 111 chemicals have been tested in all three taxonomic groups, while 1,909 and 1,444 have been tested on fish, and crustaceans, respectively, while only 174 chemicals have been tested on algae. Likewise, Figure 3B visalizes the imbalance between the number of chemicals tested on different pairings of taxonomic groups and the number of data points for these tests. As an example, the 111 chemicals tested in all of

fish, crustaceans, and algae, represent 10,568 data points, while 452 chemicals tested just on crustaceans only make up 825 data points.

As with species, some chemicals have been tested much more often than others. Figures 5 (A-C) show the number of data points by taxonomic group for the top 15 chemicals with the most associated data points. Primarily, chemicals are tested on fish, then on crustaceans, and lastly on algae. The insecticide Carbaryl (63-25-2) has been tested most often on fish with almost 600 data points, whereas the pesticide Sodium Pentachlorophenate (131-52-2) has been tested most often on crustaceans. For algae, the herbizide Atrazine (1912-24-9) makes up around 15% of the data points.

To better understand the chemical space of our dataset, we look at pairwise similarities between chemicals. The Tanimoto similarity was shown to be an appropriate choice for similarities based on molecular fingerprints[56] and is given as a value between 0 (dissimmilar) and 1 (equal). We calculated the similarity using the MACCS fingerprint with the BulkTanimotoSimilarity function from the Python package RDKit. Figure 12 depicts a histogram of all pairwise comparisons of the 2,411 chemicals in our dataset, which shows that the majority of chemicals are fairly dissimilar, with a mean similarity of 0.085 as indicated by the dotted vertical line.

Understanding the chemical space can also be approached using the chemical ontology ClassyFire[48], whose hierarchical structure for our dataset is shown in the icicle chart in Figure 14. 2,250 of the 2,411 chemicals in the dataset could be categorized with ClassyFire. Our dataset mainly consists of organic compounds, which is a consequence of the final filtering step which only includes water soluble and mol2vec compatible compounds. The largest superclass of the 17 superclasses, benzenoids, in total makes up more than a third of the chemicals, followed by organoheterocyclic compounds, and organic acids and derivatives. The chemicals are further divided into 140 classes and 251 subclasses, thus indicating a wide variety of compounds in the dataset.

Benzenoids being the most prevalent chemical superclass coincides with the findings from the Murcko scaffolds. There are 525 different Murcko scaffolds in our dataset of which a benzene ring and "no scaffold", *i.e.,* the molecule does not contain a ring system, are by far most common for all taxonomic groups (Figure 5) (D-C). For algae, 1,3,5-Triazine (*c1ncncn1*) is more common than "no scaffold". The latter is a very heterogenous group, since the only commonality among them is the lack of rings in their structure. The generic scaffold can be considered more reductionist by creating even broader scaffold categories. Here, the most common scaffolds are "no scaffold", and cyclohexane.

Chemicals associated with different scaffolds produce different toxicity ranges which can be seen by the different toxicity ranges covered by the top 15 Murcko scaffolds (Figure 7). The two main scaffolds cover a fairly similar toxicity range, whereas less-represented scaffolds either mostly represent a narrower or sometimes a broader part of the toxicity range. Some scaffold are particularly toxic or non-toxic for certain taxonomic groups, *e.g.,* the *O=C(OCc1cccc(Oc2ccccc2)c1)C1CC1* scaffold is less toxic to algae than to fish and crustaceans. It is the backbone of pyrethroid insecticides, most notably Permethrin, Deltamethrin, and Cypermethrin. On the other hand, chemicals based on the 1,3,5-Triazine scaffold are more toxic to algae and include primarily herbicides such as Atrazine, Simazine, and Terbutryn. These findings, apparent from the raw data, underline the informational value contained in the grouping according to scaffolds.

Additional information on the functional use categories is given in Figure 15 as an overview of the most prevalent OECD functions and their non-harmonized counterpart. For 570 of the 2,411 chemicals in our dataset, we could determine a functional use. Biocides are by far the most common, making up 43%, followed by fragrances (11%) and solvents (9%). For the biocides, insecticides are most common (21% of all chemicals with a functional use category), followed by fungicides (13%, and therefore more common than fragrances and solvents), and herbicides (3%). This result is not surprising as an ecotoxicological dataset is inherently biased towards toxic compounds. The information on the most common functional use categories has to be taken with a grain of salt as no function use information is available for most of the data.

Data splittings

We have a closer look at three splitting schemes representative for the three splitting categories in Figure 8: *totally random* (A) where data leakage is expected, *by occurrence of chemical compounds* (B), and *by Murcko scaffold* (C). For each splitting, we compare the distributions for toxicity (as EC50), molecular weight, and logP for the training and test sets. *Totally random* splitting results in coinciding distributions of the three parameters across both training and test set. Splitting *by occurrence* and *by Murcko scaffold* produces EC50 distributions that cover a similar range but peaking at different EC50 values, which is more pronounced for the splitting *by Murcko scaffold*. The distributions for molecular weight and logP are less similar, but generally cover similar ranges. Therefore, predictions based on these splits originate from the same the toxicity range they were trained on. Intuitively, the model performance should be better when the toxicity range for the test set is included in the range that was contained in the training set. Exceptions can arise from activity cliffs, structurally similar compounds that differ strongly in biological activity[57,58]).

Intuitively, splitting by scaffold (with the second largest compound group having "no scaffold" due to a lack of ring structures), presents a very challenging generalization task. During training, the model will only be presented with a rather

heterogenous subset of chemicals that are very different from the chemicals in the other scaffold bins.

Usage Notes

Prediction Challenges

Here, we propose several tasks to be addressed with our dataset, so, ultimately, results can be compared between modeling studies based on this dataset. For each data subset, we provide the training and test sets as described in the corresponding "Methods" subsection. We also provide cross-validation folds if the splitting scheme is suitable for it. Model selection should be performed on the training set, and the test set should only be used once, at the end of the study, to assess the generalization performance of the model. The test set *must* not be used for model training[10].

For all challenges, we suggest to split by occurrence of chemical compounds which prevents data leakage through the same chemicals being both in training and test sets. Splits by scaffolds are worth exploring to gain a better understanding of how they affect model outcome and whether they are suitable for splitting schemes. We strongly advise against the use of the *totally random* split as it will lead to overestimated model performances.

Using the whole dataset

Ultimately, the goal of *in silico* methods is to replace *in vivo* experiments, at least with vertebrates. We propose two ways to approach this. The first option is a generalization to species, *i.e.,* train on crustaceans and algae, and test the models on fish (see Table 3, a-CA2F-same and a-CA2F-diff). If it is possible to learn from invertebrate and algae *in vivo* data, which are considered less ethically problematic and more cost efficient, the number of tests on fish may be reduced in the future. This generalization to species can be performed either on the same chemicals (a-CA2F-same) or on different chemicals (a-CA2F-diff). The latter corresponds to a generalization to both species and chemicals, which presumably is the hardest challenge we propose. The second option is to train and test on all data of the three taxonomic groups to improve predictions for the species and chemicals available in the dataset (a-FCA2FCA).

Within a taxonomic group

The three taxonomic groups might be too different to be modelled together. Alternatively, we propose prediction challenges within each taxonomic group (see Table 4). These challenges may help to gain a better understanding of mortality within each taxonomic group and to determine whether the inclusion of species-related features helps to find explainable models with good prediction quality.

On single species

Since most previous studies focused on single species, we also provide challenges for the most prominent species for each taxonomic group in our dataset (see Table 5). For these challenges, species-specific features are not informative and need to be ignored.

Limitations

We would like to address some limitations that concern the field of predictive ecotoxicology in general and our dataset in particular.

In vivo data is, by nature, highly variable and our core dataset spans several decades of experimental work (Figure 9). During that time, several standards and test guidelines, like those from the ISO and OECD, were instituted to achieve higher reproducibility of experiments. Similarly, not all information desirable for modeling is contained in the original sources. Generally, the abundance of features comes with the caveat that not all parameters were recorded in all experiments, resulting in a fair amount of "Not recorded" (NR, NC) values. For some parameters we remove those entries, for others they make up the majority of entries and could therefor not be removed, *e.g.* the organism life stage as shown in Figure 10 (E). The inclusion of an appropriate control treatment is under most circumstances a requirement of the ECOTOX selection process[13]. Nonetheless, the encoding in the database includes the factors "inappropriate", "unsatisfactory", "not reported", and "unspecified". These encodings indicate for which entries the desirable control types, *i.e.,* negative, positive, and solvent control, were not included in the original study[59]). Nonetheless, certain data points are included by ECOTOX, when the use of controls can be deduced from the original source. Here, we decided to trust the selection criteria for the ECOTOX database, but advise against using control type as modeling feature (see Table 8).

Furthermore, the toxicity of different compounds, especially metals, is highly dependent on the medium composition and medium properties. This behavior is very complex and hard to predict, but to a degree interlinked with the acid dissociation constant pKa[60]. Therefore, we included this parameter such that its impact on model performance can be studied. Concurrently, isomerism and 3d-structures of molecules are not captured by the molecular representations used here.

Ideally, chemical properties, molecular representations, and species-related information would fully explain all toxicity outcomes by covering every physical, chemical, and biological dimension of an experiment. This, however is not the case

with the features currently available. Mortality in itself is a highly unspecific endpoint that can be caused through a plethora of different mechanisms, particularly when considering the differing biology of the studied organism types fish, crustaceans, and algae. Cheminformatics use cases following a more mechanistic focus, *e.g.,* modeling the interaction of a molecule with a specific molecular targets such as triggering a molecular event/receptor response/gene expression change;[61], likely have a higher benefit from molecular representations, since there is an immediate cause-and-effect-relationship between the molecular structure and the molecular target at hand. With predictive ecotoxicology still in its infancy, this dataset hopefully fosters exploration and a better understanding of what kind of data is helpful or lacking to further progress the field. Accordingly, our goal is to help the field of predictive ecotoxicology progress by enabling reproducibility and comparability across studies based on our dataset.

Code availability

The code used to load and process the input data and generate the output dataset was created and run in Python 3.8 and is made available on https://renkulab.io/gitlab/mltox/adore. The repository contains code on how to load the data, prepare it for modeling, *e.g.,* create one-hot and multi-hot-encodings for categorical features, and apply the train-test-split for 5-fold cross-validation. A good starting point are the files in the folder *scripts* for random forests (*14_analysis_regression_rf.py* and *34_analysis_regression_test_rf.py*).

References

1. EC – European Commission. Regulation (EC) No 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), establishing a European Chemicals Agency, amending Directive 1999/45/EC and repealing Council Regulation (EEC) No 793/93 and Commission Regulation (EC) No 1488/94 as well as Council Directive 76/769/EEC and Commission Directives 91/155/EEC, 93/67/EEC, 93/105/EC and 2000/21/EC (2006).

2. Mittal, K. *et al.* Resource requirements for ecotoxicity testing: A comparison of traditional and new approach methods. Preprint, Pharmacology and Toxicology (2022). 10.1101/2022.02.24.481630.

3. Wang, Z., Walker, G. W., Muir, D. C. G. & Nagatani-Yoshida, K. Toward a Global Understanding of Chemical Pollution: A First Comprehensive Analysis of National and Regional Chemical Inventories. *Environ. Sci. & Technol.* **54**, 2575–2584, 10.1021/acs.est.9b06379 (2020).

4. Muratov, E. N. *et al.* QSAR without borders. *Chem. Soc. Rev.* **49**, 3525–3564, 10.1039/D0CS00098A (2020).

5. Wu, J. *et al.* Predicting chemical hazard across taxa through machine learning. *Environ. Int.* **163**, 107184, 10.1016/j.envint.2022.107184 (2022).

6. Tuulaikhuu, B.-A., Guasch, H. & García-Berthou, E. Examining predictors of chemical toxicity in freshwater fish using the random forest technique. *Environ. Sci. Pollut. Res.* **24**, 10172–10181, 10.1007/s11356-017-8667-4 (2017).

7. Li, F. *et al. In Silico* prediction of pesticide aquatic toxicity with chemical category approaches. *Toxicol. Res.* **6**, 831–842, 10.1039/C7TX00144D (2017).

8. Wu, X., Zhou, Q., Mu, L. & Hu, X. Machine learning in the identification, prediction and exploration of environmental toxicology: Challenges and perspectives. *J. Hazard. Mater.* **438**, 129487, 10.1016/j.jhazmat.2022.129487 (2022).

9. Alves, V. M. *et al.* Oy Vey! A Comment on "Machine Learning of Toxicological Big Data Enables Read-Across Structure Activity Relationships Outperforming Animal Test Reproducibility". *Toxicol. Sci.* **167**, 3–4, 10.1093/toxsci/kfy286 (2019).

10. Kapoor, S. & Narayanan, A. Leakage and the Reproducibility Crisis in ML-based Science (2022). 2207.07048.

11. Addor, N., Newman, A. J., Mizukami, N. & Clark, M. P. The CAMELS data set: Catchment attributes and meteorology for large-sample studies. *Hydrol. Earth Syst. Sci.* **21**, 5293–5313, 10.5194/hess-21-5293-2017 (2017).

12. Kratzert, F. *et al.* Caravan - A global community dataset for large-sample hydrology. *Sci. Data* **10**, 61, 10.1038/s41597-023-01975-w (2023).

13. Olker, J. H. *et al.* The ECOTOXicology Knowledgebase: A Curated Database of Ecologically Relevant Toxicity Tests to Support Environmental Research and Risk Assessment. *Environ. Toxicol. Chem.* **41**, 1520–1539, 10.1002/etc.5324 (2022).

14. Connors, K. A. *et al.* Creation of a Curated Aquatic Toxicology Database: EnviroTox. *Environ. Toxicol. Chem.* **38**, 1062–1073, 10.1002/etc.4382 (2019).

15. Newman, M. C. *Fundamentals of Ecotoxicology* (CRC Press, 2014).

16. OECD. *Test No. 203: Fish, Acute Toxicity Test* (Organisation for Economic Co-operation and Development, Paris, 2019).

17. OECD. *Test No. 202:* Daphnia *Sp. Acute Immobilisation Test* (Organisation for Economic Co-operation and Development, Paris, 2004).

18. OECD. *Test No. 201: Alga, Growth Inhibition Test*. OECD Guidelines for the Testing of Chemicals, Section 2: Effects on Biotic Systems (OECD Publishing, 2006).

19. OECD. *Test No. 249: Fish Cell Line Acute Toxicity: The RTgill-W1 Cell Line Assay* (OECD, 2021).

20. OECD. *Test No. 236: Fish Embryo Acute Toxicity (FET) Test*. OECD Guidelines for the Testing of Chemicals, Section 2 (OECD Publishing, 2013).

21. Fischer, M. *et al.* Repeatability and Reproducibility of the RTgill-W1 Cell Line Assay for Predicting Fish Acute Toxicity. *Toxicol. Sci.* **169**, 353–364, 10.1093/toxsci/kfz057 (2019).

22. Heller, S. R., McNaught, A., Pletnev, I., Stein, S. & Tchekhovskoi, D. InChI, the IUPAC International Chemical Identifier. *J. Cheminformatics* **7**, 23, 10.1186/s13321-015-0068-4 (2015).

23. Weininger, D. SMILES, a chemical language and information system. 1. Introduction to methodology and encoding rules. *J. Chem. Inf. Comput. Sci.* **28**, 31–36, 10.1021/ci00057a005 (1988).

24. Gómez-Bombarelli, R. *et al.* Automatic Chemical Design Using a Data-Driven Continuous Representation of Molecules. *ACS Cent. Sci.* **4**, 268–276, 10.1021/acscentsci.7b00572 (2018).

25. Belfield, S. J., Cronin, M. T., Enoch, S. J. & Firman, J. W. Guidance for good practice in the application of machine learning in development of toxicological quantitative structure-activity relationships (QSARs). *PLOS ONE* **18**, e0282924, 10.1371/journal.pone.0282924 (2023).

26. Kim, S. *et al.* PubChem 2023 update. *Nucleic Acids Res.* **51**, D1373–D1380, 10.1093/nar/gkac956 (2023).

27. Matt Swain *et al.* PubChemPy (2014).

28. Lumor, S. E., Diez-Gonzalez, F. & Labuza, T. P. Detection of Warfare Agents in Liquid Foods Using the Brine Shrimp Lethality Assay. *J. Food Sci.* **76**, T16–T19, 10.1111/j.1750-3841.2010.01966.x (2011).

29. Toma, C. *et al.* New Models to Predict the Acute and Chronic Toxicities of Representative Species of the Main Trophic Levels of Aquatic Environments. *Molecules* **26**, 6983, 10.3390/molecules26226983 (2021).

30. Kooijman, B. *Dynamic Energy Budget Theory for Metabolic Organisation* (Cambridge University Press, Cambridge, 2009), third edn.

31. Kooijman, S. A. L. M. & Lika, K. Resource allocation to reproduction in animals: Resource allocation to reproduction in animals. *Biol. Rev.* **89**, 849–859, 10.1111/brv.12082 (2014).

32. Kooijman, S. A. L. M., Sousa, T., Pecquerie, L., Van Der Meer, J. & Jager, T. From food-dependent statistics to metabolic parameters, a practical guide to the use of dynamic energy budget theory. *Biol. Rev.* **83**, 533–552, 10.1111/j.1469-185X.2008.00053.x (2008).

33. Moore, D. R., Priest, C. D., Galic, N., Brain, R. A. & Rodney, S. I. Correcting for Phylogenetic Autocorrelation in Species Sensitivity Distributions. *Integr. Environ. Assess. Manag.* **16**, 53–65, 10.1002/ieam.4207 (2020).

34. Kumar, S. *et al.* TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol. Biol. Evol.* 10.1093/molbev/msac174 (2022).

35. Sukumaran, J. & Holder, M. T. DendroPy: A Python library for phylogenetic computing. *Bioinformatics* **26**, 1569–1571, 10.1093/bioinformatics/btq228 (2010).

36. Williams, A. J. *et al.* The CompTox Chemistry Dashboard: A community data resource for environmental chemistry. *J. Cheminformatics* **9**, 61, 10.1186/s13321-017-0247-6 (2017).

37. Landrum, G. Rdkit/rdkit: 2022_09_2 (Q3 2022) Release. Zenodo, 10.5281/ZENODO.591637 (2022).

38. Mannhold, R., Poda, G. I., Ostermann, C. & Tetko, I. V. Calculation of Molecular Lipophilicity: State-of-the-Art and Comparison of LogP Methods on more than 96,000 Compounds. *J. Pharm. Sci.* **98**, 861–893, 10.1002/jps.21494 (2009).

39. Wang, Z., Meador, J. P. & Leung, K. M. Y. Metal toxicity to freshwater organisms as a function of pH: A meta-analysis. *Chemosphere* **144**, 1544–1552, 10.1016/j.chemosphere.2015.10.032 (2016).

40. Dimitrov, S. D. *et al.* QSAR Toolbox – workflow and major functionalities. *SAR QSAR Environ. Res.* **27**, 203–219, 10.1080/1062936X.2015.1136680 (2016).

41. Cartwright, H. M. (ed.) *Machine Learning in Chemistry: The Impact of Artificial Intelligence*. Theoretical and Computational Chemistry Series (Royal Society of Chemistry, Cambridge, 2020).

42. Cereto-Massagué, A. *et al.* Molecular fingerprint similarity search in virtual screening. *Methods* **71**, 58–63, 10.1016/j.ymeth.2014.08.005 (2015).

43. Durant, J. L., Leland, B. A., Henry, D. R. & Nourse, J. G. Reoptimization of MDL Keys for Use in Drug Discovery. *J. Chem. Inf. Comput. Sci.* **42**, 1273–1280, 10.1021/ci010132r (2002).

44. Bolton, E. E., Wang, Y., Thiessen, P. A. & Bryant, S. H. Chapter 12 - PubChem: Integrated Platform of Small Molecules and Biological Activities. In Wheeler, R. A. & Spellmeyer, D. C. (eds.) *Annual Reports in Computational Chemistry*, vol. 4, 217–241, 10.1016/S1574-1400(08)00012-1 (Elsevier, 2008).

45. Rogers, D. & Hahn, M. Extended-Connectivity Fingerprints. *J. Chem. Inf. Model.* **50**, 742–754, 10.1021/ci100050t (2010).

46. Jaeger, S., Fulle, S. & Turk, S. Mol2vec: Unsupervised Machine Learning Approach with Chemical Intuition. *J. Chem. Inf. Model.* **58**, 27–35, 10.1021/acs.jcim.7b00616 (2018).

47. Mikolov, T., Chen, K., Corrado, G. & Dean, J. Efficient Estimation of Word Representations in Vector Space (2013). 1301.3781.

48. Djoumbou Feunang, Y. *et al.* ClassyFire: Automated chemical classification with a comprehensive, computable taxonomy. *J. Cheminformatics* **8**, 61, 10.1186/s13321-016-0174-y (2016).

49. OECD. Internationally Harmonised Functional, Product and Article Use Categories (2017).

50. Kaufman, S., Rosset, S., Perlich, C. & Stitelman, O. Leakage in data mining: Formulation, detection, and avoidance. *ACM Transactions on Knowl. Discov. from Data* **6**, 1–21, 10.1145/2382577.2382579 (2012).

51. Wu, K. & Wei, G.-W. Quantitative Toxicity Prediction Using Topology Based Multitask Deep Neural Networks. *J. Chem. Inf. Model.* **58**, 520–531, 10.1021/acs.jcim.7b00558 (2018).

52. Song, R. *et al.* Accelerating the pace of ecotoxicological assessment using artificial intelligence. *Ambio* **51**, 598–610, 10.1007/s13280-021-01598-8 (2022).

53. Yang, K. *et al.* Analyzing Learned Molecular Representations for Property Prediction. *J. Chem. Inf. Model.* **59**, 3370–3388, 10.1021/acs.jcim.9b00237 (2019).

54. Simm, J. *et al.* Splitting chemical structure data sets for federated privacy-preserving machine learning. *J. Cheminformatics* **13**, 96, 10.1186/s13321-021-00576-2 (2021).

55. Bemis, G. W. & Murcko, M. A. The Properties of Known Drugs. 1. Molecular Frameworks. *J. Medicinal Chem.* **39**, 2887–2893, 10.1021/jm9602928 (1996).

56. Bajusz, D. Why is Tanimoto index an appropriate choice for fingerprint-based similarity calculations? *J. Cheminformatics* 13 (2015).

57. Stumpfe, D., Hu, H. & Bajorath, J. Evolving Concept of Activity Cliffs. *ACS Omega* **4**, 14360–14368, 10.1021/acsomega.9b02221 (2019).

58. Stumpfe, D., Hu, H. & Bajorath, J. Advances in exploring activity cliffs. *J. Comput. Mol. Des.* **34**, 929–942, 10.1007/s10822-020-00315-z (2020).

59. Moermond, C. T., Kase, R., Korkaric, M. & Ågerstrand, M. CRED: Criteria for reporting and evaluating ecotoxicity data. *Environ. Toxicol. Chem.* **35**, 1297–1309, 10.1002/etc.3259 (2016).

60. Mansouri, K. *et al.* Open-source QSAR models for pKa prediction using multiple machine learning approaches. *J. Cheminformatics* **11**, 60, 10.1186/s13321-019-0384-1 (2019).

61. Xu, T., Xia, M. & Huang, R. Modeling Tox21 Data for Toxicity Prediction and Mechanism Deconvolution. In Hong, H. (ed.) *Machine Learning and Deep Learning in Computational Toxicology*, Computational Methods in Engineering & the Sciences, 463–477, 10.1007/978-3-031-20730-3_19 (Springer International Publishing, Cham, 2023).

62. OECD. OECD QSAR Toolbox (2023).

Acknowledgements

This work was made possible through the SDSC grant "Enhancing Toxicological Testing through Machine Learning" (project NoC20-04) and partly carried out in the framework of the European Partnership for the Assessment of Risks from Chemicals (PARC) and has received funding from the European Union's Horizon Europe research and innovation programme under Grant

Author contributions statement

L.G. and C.S.: Conceptualization, Software, Validation, Formal analysis, Investigation, Data curation, Visualization, Writing - Original Draft, Writing - Review & Editing. F.P.C, K.S., M.B.J.: Conceptualization, Supervision, Funding acquisition, Writing - Review & Editing. All authors reviewed and approved the manuscript.

Competing interests

The authors declare no competing financial and/or non-financial interests.

Figures & Tables

Table 1. Effect groups for different taxonomic groups. Definitions given in the Annex B

Fish	Crustaceans	Algae
Mortality (MOR)	Mortality (MOR)	Mortality (MOR)
-	Intoxication (ITX; immobilization)	Growth (GRO)
-	-	Population (POP)
-	-	Physiology (PHY)

Table 2. Overview of the Add my Pet features and their encoding. More detailed descriptions of the discrete levels and continuous values for these features can be found in Figure 11 and the Annex.

type	feature	code	unit
ecology	ecozone		
	climate		
	migration		
	food		
life history	life span	am	d
	body length at birth	Lb	cm
	body length at puberty	Lp	cm
	ultimate body length	Li	cm
	reproductive rate	Ri	$\#/d$
pseudo-data	allocation fraction to soma	kap	-
	energy conductance	v	cm/d
	volume-specific somatic maintenance cost	p_M	$J/d \cdot cm^3$

Figure 1. A: Histogram of the number of data points associated with a combination of chemical, species, and observation duration, concentration type, exposure type, media type, and effect type in our dataset. B: Boxplot for the distribution of toxicity values for experimental conditions with at least 25 values.

Table 3. Prediction challenges using the whole dataset. The names of these challenges start with "a".

Name	Train on	Test on	Generalization to	Data split
a-CA2F-same	crustaceans and algae	fish	species	by occurrence of chemical compounds
a-CA2F-diff	crustaceans and algae	fish	species and chemicals	by occurrence of chemical compounds
a-FCA2FCA	all data	all data	neither	by occurrence of chemical compounds

Table 4. Prediction challenges within a taxonomic group. The names of these challenges start with "t".

Name	Train on	Test on	Data split
t-F2F	fish	fish	by occurrence of chemical compounds
t-C2C	crustaceans	crustaceans	by occurrence of chemical compounds
t-A2A	algae	algae	by occurrence of chemical compounds

Table 5. Single species prediction challenges. The names of these challenges start with "s".

Name	Train on	Test on	Data split
s-F2F	*Oncorhynchus mykiss* (Rainbow Trout)	*Oncorhynchus mykiss*	by occurrence of chemical compounds
s-C2C	*Daphnia magna* (Water Flea)	*Daphnia magna*	by occurrence of chemical compounds
s-A2C	*Chlorella vulgaris* (Microalga)	*Chlorella vulgaris*	by occurrence of chemical compounds

Figure 2. Conceptual overview of the data sources and processing steps.

Figure 3. A: Venn diagram of the 2,411 tested chemicals. As reading example, 917 chemicals were tested only on fish, 868 chemicals were tested on fish and crustaceans, and 111 chemicals were tested on all taxonomic groups. B: Overview of the number of chemicals tested (left) in the different combinations of taxonomic groups and the related number of data points (right), indicating that many tested chemicals do not in all cases translate to many data points.

Figure 4. A: Overview of the toxicity distributions across effect groups by taxonomic groups. B, C: Toxicity classes across taxonomic groups in a binary and 5-level classification system, respectively.

Figure 5. Top chemicals (A-C) and chemical scaffolds (D-F) for the three taxonomic groups. Table 14 gives an overview of the structures of each scaffold.

Figure 6. A-C: Overview of the top 15 species per taxonomic group. D: The phylogenetic distances as circular dendrogram. Line lengths after branching indicate phylogenetic distances. The species labels are only representative and do not necessarily cover all included species.

Figure 7. Overview of the effective concentration 50 (EC50, in mg/L) values for the top 15 Murcko scaffolds.

Figure 8. Overview of EC50 (log10-transformation of the mass concentration in mg/L), molecular weight (in g/mol), and logP for three data splittings: A: *Totally random*, B: *by occurrence of chemical compound*, and C: *by Murcko scaffold*.

Table 6. Overview of the chemical properties and their sources. CPDat: Chemicals and Product Database

Chemical property	Abbreviation	Unit	Source(s)
Molecular weight	MW	g/mol	DSSTox[36]
Octanol-water partition coefficient	LogP, $LogK_{ow}$	-	RDKit[37]
Water solubility	WS	mg/L	DSSTox[36]
Melting point	MP	°C	DSSTox[36]
Number of heavy atoms		-	PubChem[26]
Number of bonds		-	manually calculated from PubChem[26] data
Number of double bonds		-	manually calculated from PubChem[26] data
Number of triple bonds		-	manually calculated from PubChem[26] data
Number of ring structures		-	RDKit[37]
Number of OH groups		-	RDKit[37]
Acid dissociation constant	pKa	-	QSAR toolbox[62]
Chemical ontology			ClassyFire[48]
Functional use category			CPDat[49]

Table 7. Total number data points after processing. N: number of data points

Taxonomic group	Number of species	N	%
Fish	140	26,163	78.1
Crustaceans	17	6,631	19.8
Algae	46	704	2.1
All	**203**	**33,498**	**100.0**

A Glossary

The following table 8 includes descriptions and definitions for all columns of the dataset.

Table 8. A glossary of all columns in the dataset. The prefix of the column name identifies the feature category or source file, *e.g.,* column names starting with "test" originate from the ECOTOX tests file. The dataset contains modeling features as well as additional information. Only the features labelled "suitable for modeling" should be used for modeling. The others *must not* be used. Effective concentrations are labelled as targets. Only one target should be used and the others *must not* be used as modeling feature.

Column name	Description	Suitable for modeling
test_id	Unique identifier for an experiment (links with ECOTOX)	no
reference_number	Unique identifier for the original source (links with ECOTOX)	no
test_cas	CAS number of the chemical compound	no
test_location	Test location	no
test_exposure_type	Exposure type	yes
test_control_type	Control type	no
test_media_type	Media type	yes
test_application_freq_unit	Application frequency unit	no
test_organism_lifestage	Organism life stage	no
result_id	Unique identifier for each data point (links with ECOTOX)	no
result_effect	Effect group	no
result_endpoint	Endpoint	no
result_obs_duration_mean	Observation duration	yes
result_conc1_type	Exposure concentration type	yes
result_conc1_mean_op	Effective concentration operator	no
result_conc1_mean	Effective concentration value (in mg/L)	target
result_conc1_mean_mol	Effective concentration value (in mol/L)	target
media_ph_mean	Measured medium pH	yes
media_temperature_mean	Measured medium temperature (in °C)	yes
tax_all	Combination of all taxonomic levels	no
tax_name	Common species name	no
tax_class	Taxonomic class	no
tax_order	Taxonomic order	no
tax_family	Taxonomic family	no
tax_genus	Taxonomic genus	no
tax_species	Taxonomic species	no
tax_gs	Taxonomic genus and species	no
species_number	Species identifier (links with ECOTOX)	no
tax_group	Taxonomic group	no
tax_pdm_available	Boolean whether phylogenetic distance is available	no
tax_eco_climate	Ecology, climate zone	yes
tax_eco_ecozone	Ecology, ecozone	yes
tax_eco_food	Ecology, food class	yes
tax_eco_migrate5	Ecology, migratory behavior (5-level encoding)	yes
tax_eco_migrate2	Ecology, migratory behavior (2-level encoding)	yes
tax_lh_amd	Life history, life span (in d)	yes
tax_lh_lbcm	Life history, body length at birth (in cm)	yes
tax_lh_lpcm	Life history, body length at puberty (in cm)	yes
tax_lh_licm	Life history, ultimate body length (in cm)	yes
tax_lh_ri#/d	Life history, reproductive rate (in $\#/d$)	yes
tax_ps_ampv	Pseudo-data, energy conductance (in cm/d)	yes
tax_ps_ampkap	Pseudo-data, allocation fraction to soma	yes

Column name	Description	Suitable for modeling
tax_ps_amppm	Pseudo-data, volume-specific somatic maintenance cost (in $J/d \cdot cm^3$)	yes
result_conc1_mean_binary	Effective concentration category (more toxic/less toxic)	target
result_conc1_mean_log	Effective mass concentration value after a log10 transformation	target
result_conc1_mean_mol_log	Effective molar concentration value after a log10 transformation	target
chem_dtxsid	DSSTOX substance ID	no
chem_name	Name of chemical compound	no
test_cas_name	Concatenation of CAS number and name of chemical compound	no
chem_sf	Molecular formula	no
chem_mw	Molecular weight (in g/mol)	yes
chem_mp	Melting point (in °C)	yes
chem_ws	Water solubility (in mg/L)	yes
chem_ws_binary	Boolean whether water solubility is available	no
chem_dtxcid	DSSTOX compound ID	no
chem_inchi	InChI from DSSTox	no
chem_inchikey	InChIkey from DSSTox	no
chem_pcp_cid	PubChem compound ID	no
chem_pcp_inchi	InChI from PubChem	no
chem_pcp_inchikey	InChIkey from PubChem	no
chem_pcp_iupac_name	IUPAC name	no
chem_pcp_can_smiles	Canonical SMILES from PubChem	no
chem_pcp_fp	Collapsed PubChem fingerprint	yes
chem_pcp_heavy_atom_count	Number of heavy atoms, *i.e.,* not hydrogen	yes
chem_rdkit_clogp	Octanol-water partition coefficient	yes
chem_rdkit_can_smiles	Canonical SMILES from RDKit	no
chem_pcp_bonds_count	Number of bonds	yes
chem_pcp_doublebonds_count	Number of double bonds	yes
chem_pcp_triplebonds_count	Number of triple bonds	yes
chem_rings_count	Number of rings	yes
chem_OH_count	Number of OH groups	yes
chem_mol2vec_allowed	Boolean: compatible with mol2vec	no
chem_pka_median	Acid dissociation constant	yes
chem_MACCS_fp	Collapsed MACCS fingerprint	yes
chem_Morgan_fp	Collapsed Morgan fingerprint	yes
chem_mol2vec[000-299]	300-dimensional mol2vec embedding	yes
split_totallyrandom	Split data points totally at random (incl. cross-validation folds)	no
split_random	Split chemicals at random (incl. cross-validation folds)	no
split_occurrence	Split chemicals by occurrence (incl. cross-validation folds)	no
split_scaffold-murcko	Split chemicals by Murcko scaffold (incl. cross-validation folds)	no
split_scaffold-murcko-loo-0	Split chemicals by Murcko scaffold (only training and test set)	no
split_scaffold-murcko-loo-1	Split chemicals by Murcko scaffold (only training and test set)	no
split_scaffold-murcko-llo	Split chemicals by Murcko scaffold (only training and test set)	no
split_scaffold-generic	Split chemicals by generic scaffold (incl. cross-validation folds)	no
split_scaffold-generic-loo-0	Split chemicals by generic scaffold (only training and test set)	no
split_scaffold-generic-loo-1	Split chemicals by generic scaffold (only training and test set)	no
split_scaffold-generic-llo	Split chemicals by generic scaffold (only training and test set)	no

B ECOTOX data

Definitions for ECOTOX effect groups as given in the ECOTOX term appendix:

- **Mortality (MOR):** Measurements and endpoints where the cause of death is by direct action of the chemical.

- **Physiology (PHY)/Intoxication (ITX):** Measurements and endpoints regarding basic activity in cells and tissues of plants or animals; includes four effect groups - injury, immunity, intoxication and general physiological response.

- **Growth (GRO):** Category encompasses measures of weight and length, and includes effects on development, growth and morphology.

- **Population (POP):** Measurements and endpoints relating to a group of organisms or plants of the same species occupying the same area at a given time.

Toxicity categories The toxicity intervals given in Table 9 and shown for the chemicals in our dataset in Figure 4 are in accordance with EPA.

Table 9. EC50 intervals for the binary and multi-class toxicity classification used in Figure 4.

EC50 interval (mg/L)	Binary classification	Multi-class classification
$(-\infty, 10^{-1}]$	more toxic	very highly toxic
$(10^{-1}, 10^{0}]$	more toxic	highly toxic
$(10^{0}, 10^{1}]$	less toxic	moderately toxic
$(10^{1}, 10^{2}]$	less toxic	slightly toxic
$(10^{2}, +\infty)$	less toxic	non-toxic

Figure 9. Chronology of the references related to the different taxonomic groups added to ECOTOX over time. The publication year refers to the year the original study was published. It is noteworthy that some references are not singular scientific studies, but entail whole databases, such as the US EPA "Pesticide Ecotoxicity Database (Formerly: Environmental Effects Database (EEDB))", added in 1992.

Reference chronology

 B.1 Experimental properties

Figure 10. Overview of the most common values for experimental properties parameters. A: Media type; B: Observation duration; B: Exposure type; D: Control type; D: Organism life stage (only the top 15 of 48 total, less-abundant life stages are combined into "other"); F: Test location. The number of entries are shown on a logarithmic scale.

C Taxonomic data

C.1 Overview

Figure 11. A: Overview of the taxonomic properties for ecology. B: Overview for life history and pseudo-data parameter values. The encodings for the different levels are given in Tables 10-13.

C.2 Ecological data encoding

Here, we provide the encodings for the ecological data. For some parameters, we had to reduce the complexity of the encodings. More detailed descriptions of the parameters and the original encoding can be found in the Add my Pet collection

 ### C.2.1 Climate zone

Table 10. Climate zone definitions and encodings.

Encoding	Definition
A	Tropical (megathermal) climates: every month of the year with an average temperature of 18 °C or higher, with significant precipitation
B	Dry (arid and semi-arid) climates: little precipitation
C	Temperate (mesothermal) climates: the coldest month averaging between 0 and 18 °C and at least one month averaging above 10 °C
D	Continental (microthermal) climates: at least one month averaging below -3 °C and at least one month averaging above 10 °C
E	Polar and alpine (montane) climates: every month of the year with an average temperature below 10 °C
M	Marine climates

 ### C.2.2 Ecozone

Table 11. Ecozone definitions and encodings.

Encoding	Definition
MA	Atlantic ocean
MC	Circumglobal oceans
MI	Indian ocean
MN	Arctic ocean
MP	Pacific ocean
MS	Southern ocean
TA	Australasia (including New Guinea, New Zealand)
TH	Holarctic
TN	Neotropic (including Central America, and the Caribbean)
TP	Paleotropic

 ### C.2.3 Migratory behavior

Table 12. Migratory definitions and encodings for 2 and 5 levels.

2-level encoding	5-level encoding	Definition
no_migration	no_migration	Do not migrate
migration	amphidromous	Migrate from fresh water to the sea, or vice versa, but not for breeding
migration	diadromous	Migrate between the sea and fresh water
migration	oceanodromous	Live and migrate wholly in the sea
migration	potamodromous	Live and migrate wholly within fresh water

 ### C.2.4 Food

Table 13. Food definitions and encodings.

Encoding	Definition
B	Bacterivore (including micro-organisms)
C	Carnivore (living animals)
D	Detrivore (bacteria, small fungi, organic matter)
H	Herbivore (plants)
O	Omnivore (plants/animals/fungi)
P	Planktivore (small aquatic organisms, macro-movement controlled by flow, not by swimming)
S	Scavenger (dead animals)
T	Parasitic (animal tissue)

D Chemical data

D.1 Tanimoto similarity

Figure 12. Histogram of the Tanimoto similarities for the chemicals in our dataset, ranging from 0 (dissimmilar) to 1 (equal). The dashed vertical line indicates the mean similarity of 0.085.

 D.2 Chemical properties

Figure 13. Overview of the distributions of the chemical properties represented in our dataset. See Table 6 for more details.

Figure 14. Icicle chart representing the chemical ontology for 2,250 of the 2,411 chemicals in our dataset, with the hierarchical levels kingdom, superclass, class, and subclass.

 D.4 Functional use categories

Figure 15. Prevalent functional use categories for 570 of the 2,411 chemicals in our dataset. Most chemicals have several reported uses of which we select the prevalent ones, *i.e.,* those reported in at least 20% of the cases for each chemical. A: Internationally harmonized OECD functional uses, B: non-harmonized functional uses distinguish between the different biocides.

 D.5 Molecular scaffolds

Table 14. Canonical SMILES and chemical structures of the most common Murcko scaffolds. We do not include "no scaffold" here.

Canonical SMILES	Structure
c1ccccc1	
c1ccc(Cc2ccccc2)cc1	
c1ccncc1	

Canonical SMILES	Structure
c1ccc2ccccc2c1	
O=C(OCc1cccc(Oc2ccccc2)c1)C1CC1	
C1=CC2CC1C1C3CC(C4OC34)C21	
C1CCCCC1	
O=C(Nc1ccccc1)c1ccccc1	
O=c1[nH]nnc2ccccc12	
O=S1OCC2C3C=CC(C3)C2CO1	
c1ncncn1	

Canonical SMILES	Structure
c1cncnc1	
O=C1c2ccc3c(c2OC2COc4ccccc4C12)CCO3	
O=C(Cc1ccccc1)OCc1cccc(Oc2ccccc2)c1	
c1ccc2[nH]cnc2c1	
C1=NCCN1Cc1cccnc1	
c1ccc(Oc2ccccc2)cc1	
c1cc(-c2cc[nH+]cc2)cc[nH+]1	
O=c1cccnn1-c1ccccc1	

Canonical SMILES	Structure
c1ncnc(NC2CC2)n1	
C(=Cc1ccccc1)C(C=Cc1ccccc1)=NNC1=NCCCN1	
O=C(Nc1ncncn1)NS(=O)(=O)c1ccccc1	
c1ccc(C2OC2(Cn2cncn2)c2ccccc2)cc1	
c1cc[n+]2c(c1)-c1cccc[n+]1CC2	
c1ccc(C2(Cn3cncn3)OCCO2)cc1	