Biochemical, biomarker, and behavioral characterization of the Grn^{R493X} mouse model of frontotemporal dementia

Denise M. Smith[1,2,3], Geetika Aggarwal[1,2,3], Michael L. Niehoff[1,4], Spencer A. Jones[1,2,3], Subhashis Banerjee[1,2,3], Susan A. Farr[1,2,3,4], Andrew D. Nguyen[1,2,3,*]

[1]Saint Louis University School of Medicine, Department of Internal Medicine, Division of Geriatric Medicine, United States of America

[2]Saint Louis University School of Medicine, Department of Pharmacology and Physiology, United States of America

[3]Saint Louis University, Institute for Translational Neuroscience, United States of America

[4]Veterans Affairs Medical Center, United States of America

*Correspondence: andy.d.nguyen@health.slu.edu

ABSTRACT

Heterozygous loss-of-function mutations in the progranulin gene (*GRN*) are a major cause of frontotemporal dementia due to progranulin haploinsufficiency; complete deficiency of progranulin causes neuronal ceroid lipofuscinosis. Several progranulin-deficient mouse models have been generated, including both knockout mice and knockin mice harboring a common patient mutation (R493X). However, the *Grn*^R493X mouse model has not been characterized completely. Additionally, while homozygous *Grn* mice have been extensively studied, data from heterozygous mice is still limited. Here, we performed more in depth characterization of heterozygous and homozygous *Grn*^R493X knockin mice, which includes neuropathological assessment, behavioral studies, and analysis of fluid biomarkers. In the brains of homozygous *Grn*^R493X mice, we found increased expression of lysosomal genes, markers of microgliosis and astrogliosis, pro-inflammatory cytokines, and complement factors. Heterozygous *Grn*^R493X mice exhibited more limited increases in lysosomal and inflammatory gene expression. Behavioral studies found social and emotional deficits in *Grn*^R493X mice that mirror those observed in *Grn* mouse models, as well as impairment in memory and executive function. Overall, the *Grn*^R493X knockin mouse model closely phenocopies *Grn* knockout models. Lastly, in contrast to homozygous knockin mice, heterozygous *Grn*^R493X mice do not have elevated levels of fluid biomarkers previously identified in humans, including neurofilament light chain (NfL) and glial fibrillary acidic protein (GFAP) in both plasma and CSF. These results may help to inform pre-clinical studies that use this and other *Grn* mouse models.

Keywords: progranulin, frontotemporal dementia, neuronal ceroid lipofuscinosis, mouse model, biomarkers

INTRODUCTION

Progranulin is a widely expressed protein with pleiotropic effects, including growth factor-like properties, anti-inflammatory effects, promoting neurite outgrowth, and promoting autophagy.[1-3] However, the precise molecular function of progranulin is unclear. Within cells, progranulin localizes to lysosomes;[4] progranulin is also secreted from cells and found in blood and cerebrospinal fluid.[5]

Heterozygous loss-of-function mutations in the progranulin gene (*GRN*) are a major cause of frontotemporal dementia (FTD) due to progranulin haploinsufficiency.[6-8] FTD involves atrophy of the frontal and temporal lobes, with onset usually occurring between 40 to 55 years of age. *GRN* mutations cause behavioral variant of FTD (bvFTD), which often presents with decline in social conduct, apathy, emotional blunting, and loss of insight.[9] Neuropathologically, FTD-*GRN* is characterized by cytoplasmic ubiquitin-positive inclusions of TAR DNA-binding protein 43 (TDP-43), neuroinflammation, and no tau pathology.[6,7,10,11] Several fluid biomarkers have been identified in the CSF and blood of individuals with FTD, including neurofilament light chain (NfL), glial fibrillary acidic protein (GFAP), TDP-43, and phosphorylated TDP-43.[12-18] NfL is a particularly promising biomarker for FTD-*GRN*, because its levels correlate with disease severity.[12,13] Additionally, in asymptomatic *GRN* mutation carriers, high plasma NfL is predictive of disease conversion and progression during the subsequent two-year period.[19]

In contrast to heterozygous *GRN* mutations, complete progranulin deficiency causes neuronal ceroid lipofuscinosis (NCL), CLN11, with disease onset usually in the teenage to midlife years, although this can vary widely.[20-22] NCL/CLN11 presents with vision loss, seizures, and ataxia. NCL is a form of lysosomal storage disease, with characteristic lipofuscin accumulation, pointing to an important function of progranulin for proper lysosome function and/or homeostasis. Moreover, FTD-*GRN* and NCL highlight the gene dosage effect of progranulin deficiency which manifests as distinct diseases.

Several progranulin-deficient mouse models have been generated by different labs, including four *Grn* knockout mouse lines[23-26] and two *Grn* knockin mouse lines that harbor a common FTD-associated nonsense mutation.[27,28] Overall, similar phenotypes have been described for all these mouse lines, although most reports have focused on homozygous mice. Homozygous *Grn* knockout and knockin mice exhibit signs of lysosomal impairment, including lipofuscin accumulation, NCL-like lysosomal storage material by electron microscopy, and upregulation of lysosomal genes in the brain which is likely to compensate for the lysosomal impairment.[28-34] Homozygous *Grn* knockout and knockin mice also show age-dependent neuroinflammation and cytoplasmic accumulation of phosphorylated TDP-43.[25,27,28,32,34-37] Additionally, *Grn* knockout and *Grn*[R493X] knockin mice exhibit deficits in social and emotional behavior, including decreased sociability, age-dependent changes in social dominance, increased anxiety, and increased obsessive–compulsive behaviors.[28,35-40] Lastly, memory impairment has also been reported in *Grn* knockout mice, including in the Morris water maze and the novel object recognition test.[24,36,40]

There are currently several important gaps in our knowledge about *Grn* mouse models. Most notably, characterization of heterozygous *Grn* and *Grn*[R493X] is not complete. For example, do these mice develop lysosomal dysfunction, neuroinflammation, or memory impairment, similar

to the homozygous mice? Additionally, are the biomarkers identified in human *GRN* mutation carriers similarly increased in *Grn* mouse models? In the current study, we performed more in depth characterization of heterozygous and homozygous Grn^{R493X} knockin mice, which includes neuropathological assessment, behavioral studies, and analysis of fluid biomarkers. These results may help to inform pre-clinical studies that use this and other *Grn* mouse models.

MATERIAL AND METHODS

Mice. Grn^{R493X} knockin[28] and $Grn^{-/-}$ knockout mice[23] were on the C57BL/6J background (backcrossed more than 8 generations) and were genotyped by real-time PCR (Transnetyx). Mice were housed in a pathogen-free barrier facility with a 12-h light/12-h dark cycle and allowed food and water *ad libitum*. Experimental cohorts were generated by crossing heterozygous mice, and wild-type littermates were used for comparisons. All experimental procedures were approved by the Institutional Animal Care and Use Committee at Saint Louis University or the Harvard Medical Area Standing Committee on Animals.

Experimental design. At the indicated ages, tissues were isolated, flash frozen in liquid nitrogen, and stored at -80 °C until analyzed. Blood was collected in EDTA-coated tubes (Sarstedt, Microvette CB 300 K2 EDTA, 16.444.100); tubes were centrifuged at 2000 × *g* for 5 min and the plasma was removed and stored at -80 °C. For the experiments of Fig. 3A and Fig. S2A-B, blood was collected from the submandibular vein. In all other studies, trunk blood was collected. CSF was collected by capillary action from the fourth ventricle of anesthetized mice using a 30-gauge needle attached to Intramedic polyethylene tubing (Becton Dickinson, PE-10).

Behavioral studies. A cohort of 11-month-old male wild-type, heterozygous, and homozygous Grn^{R493X} mice underwent a test battery of behavioral tests to assess behavioral and cognitive function. Behavioral assessment was performed between 7am and 12pm. Mice were individually housed one week prior to the start of behavioral testing until the end of the study to ensure equal levels of arousal during behavioral testing. Each mouse received only one test per day. Following the behavioral studies, tissues were collected as described above at 12 months of age. For the tube test, an additional cohort of 8- to 11-month-old male mice tested by the NeuroBehavior Laboratory at the Harvard NeuroDiscovery Center was included to achieve appropriate statistical power.

Open field test. The open field test was used to assess exploratory and locomotor activity. For this test, each mouse was allowed to freely roam an empty arena (53 cm × 63 cm) for 5 min. The distance traveled and the time spent in the center of the open field were determined using the ANY-maze video tracking system (San Diego Instruments).

Forced swim test. The forced swim test was used to assess depressive behavior. Each mouse was placed in an inescapable transparent tank (30 cm height × 20 cm diameter, with a water level of 15 cm) for 6 min. The time spent moving versus the time spent immobile was recorded using the ANY-maze tracking system. Analysis was performed by comparing the time spent immobile during the last 4 min between groups. The greater the amount of time spent immobile indicates increased depressive behavior. The water temperature was monitored and maintained at 25 °C during testing by exchanging cold water in the maze with warm tap water as needed.

During testing, the mouse was continually monitored and rescued from drowning should it stop swimming or become compromised. After each trial, the mouse was placed in a cage with a warmed absorbent surface under a warming lamp until dry.

Nesting behavior. Nest building was used to assess general well-being and motor systems.[41] At the start of the study, one 4-gram Bed-r'Nest white shredded paper disc (Andersons, BRN4WSR) was placed in the front of a fresh shoebox cage that contained the equivalent of 200 ml of corn cob bedding in the bottom, along with a fresh wire top containing food and water. Each mouse was placed in the cage in the morning at 9am and left for 24 h, and then the nest was photographed. Nests were scored on a 5-point scale: Score 1 = shredded paper was scattered throughout the cage or disc was untouched. Score 2 = some shredded material was used to construct the nest, but >50% of material was scattered and untouched. Score 3 = a noticeable nest was constructed, but several pieces of shredded paper were still scattered. Score 4 = almost all material used for nest, and few pieces remained scattered or near the nest. Score 5 = all material used to make an identifiable nest.

Three-chamber sociability test. The three-chamber test[42] was used to assess the sociability of a mouse when given the choice between interacting with another mouse versus an inanimate object. For this test, each mouse was allowed to freely explore the empty three-chambered box (20.5 cm × 40.5 cm × 22 cm high) made of Plexiglas containing two empty wire cups in the outer chambers. After 10 min, a stranger mouse was placed under one of the wire cups and an inanimate object (a block) was placed under the other wire cup. During the next 10 min, the times spent interacting with the novel mouse and with the inanimate object were determined using the ANY-maze tracking system. A sociability ratio was calculated for each mouse by dividing the time spent interacting with the novel mouse by the time spent interacting with the inanimate object.

Tube test. The tube test was used to assess social dominance.[38] For this test, a wild-type mouse and a heterozygous Grn^{R493X} mouse from separate home cages were paired; each mouse was tested against three mice of the other genotype. Mice were tested without habituation to the tube. Mice were placed, head first, into opposing sides of a clear vinyl tube (3.8 cm internal diameter × 30.5 cm length) and released simultaneously. The first mouse to place two paws outside of the tube was considered to be the less dominant mouse. If the mice crossed each other or no mouse left the tube after 2 min, the trial was aborted and later repeated at the end of the session.

Elevated plus maze. The elevated plus maze was used to assess anxiety and disinhibition-like behavior. This test used a plus-shaped apparatus with two open arms and two enclosed arms, which is elevated 50 cm above the floor. Each arm is 35.5 cm in length; two opposite arms are open while the other two opposite arms are enclosed, as previously described.[43] For this test, each mouse was placed on the central platform facing an enclosed arm and allowed to freely explore the maze for 5 min. The time spent in open arms was monitored using the ANY-maze tracking system.

T-maze. The T-maze was used to assess long-term spatial memory,[44] and involves the mouse learning to traverse the maze and turning in the correct direction to find the goal box. The maze consists of a black plastic alley with a start box at one end and two goal boxes at the other. The

start box is separated from the alley by a plastic guillotine door that prevents movement down the alley until raised at the onset of training. An electrifiable floor of stainless-steel rods run throughout the maze to deliver a mild scrambled foot-shock 5 sec after the guillotine door is raised and a cue buzzer (door-bell type sounded at 55 dB). For this test, each mouse was trained to one avoidance (of a mild 0.35 mA foot shock). One week later, memory retention was tested, with a criterion for retention defined as 5 avoidances in 6 consecutive trials.

Y-maze. The Y-maze was used to assess working memory.[45] This test relies on the innate curiosity of rodents to explore previously unvisited areas, and it measures the number of visits to consecutive arms without revisiting the arm just visited. For this test, each mouse was placed in a Y-shaped maze for 5 min and the sequence of arm visits was monitored using the ANY-maze tracking system. The percentage of arm visits was determined as % alternations = (number of alternations / total number of arm entries) x 100.

Novel object recognition. The novel object recognition test was used to assess short-term memory.[46] This test relies on the innate curiosity mice to explore novel things in their environment, and it measures the ability of the mice to recognize the previously explored object. For this test, each mouse was allowed to explore a maze with two like objects placed on one wall for 5 min. One hour after the initial phase, the mouse was placed in the maze with one like object in its original place and one new object on the opposite wall for 5 min. The time spent exploring the old object and the new object was recorded, and the discrimination index (DI) was calculated as $DI = (T_{new} - T_{old}) / (T_{old} + T_{new})$.

Puzzle box. The puzzle box was used to assess complex problem solving and memory.[47,48] This test relies on the innate desire of mice to avoid open spaces, and it measures latency to find the dark side of the maze by traversing a tunnel with increasingly difficult obstacles placed at the opening of the tunnel. The puzzle box is an arena consisting of a Plexiglas white box divided by a removable barrier into two compartments: a brightly-lit start zone (58 cm × 28 cm) and a smaller, dark goal zone (15 cm × 28 cm). For this test, each mouse was introduced into the start zone and trained to move into the goal zone through a narrow underpass (~ 4 cm wide) located under the barrier. Each mouse underwent a total of 11 trials over 4 consecutive days, during which they were challenged with obstructions of increasing difficulty placed at the tunnel entrance: completely opened (Trials 1-2), partially blocked by a guillotine (Trials 3-4), blocked by nesting paper (Trials 5-7), or blocked by a T-shaped cardboard plug (Trials 8-11). Mice that took > 4 min to complete the first 3 trials were excluded from the data analysis. This experimental paradigm allows for assessment of problem solving ability, learning/short-term memory, and long-term memory.[47]

qPCR. Total RNA was isolated from flash frozen tissues using the RNeasy Mini kit (Qiagen, 74106) with on-column DNase digestion (Qiagen, 79256). RNA was reverse-transcribed to obtain cDNA using the iScript cDNA synthesis kit (Bio-Rad, 1708891), and qPCR was performed using PowerUp SYBR Green Master Mix (ThermoFisher, A25777) with a Bio-Rad CFX384 Real-Time System. Primers sequences are provided in Table S1. Results for qPCR were normalized to the average of two housekeeping genes (*36B4* and *Cyclo*), and evaluated by the comparative C_T method.

<u>Biomarkers.</u> NfL and GFAP levels were measured using the Quanterix Simoa HD-1 platform with the Simoa NF-L Advantage Kit or the Neurology 2-Plex B Advantage Kit, respectively. Plasma and CSF samples were diluted 40-fold and 200-fold, respectively. All samples were assayed in duplicate wells. For a subset of plasma samples, NfL levels were also measured using the Sigma Single Molecule Counting (SMC) platform with the SMC NF-L High Sensitivity Immunoassay Kit, with plasma samples diluted 16-fold.

<u>Statistical analyses.</u> Data are presented as mean ± SEM and were analyzed with GraphPad Prism software. Most comparisons were analyzed by one-way ANOVA with Dunnett post hoc test. In the experiment of Fig. 4B, the distribution of wins per mouse was analyzed by two-way binomial test, with the expected distribution set at 50% wins for each genotype, similar to what has been previously described.[38] P values < 0.05 were considered significant.

RESULTS

Neuropathology in the Grn^{R493X} mouse model

Previous studies of *Grn* knockout mouse models have documented increased expression of lysosomal proteins in the brain,[29-32] likely to compensate for impaired lysosomal function. To investigate if Grn^{R493X} mice exhibit similar changes, we analyzed gene expression by qPCR in cortex and thalamus tissues from Grn^{R493X} mice at 6 months and 12 months of age. As expected, heterozygous and homozygous knockin mice had decreased *Grn* mRNA levels (Fig. 1). Heterozygous and homozygous Grn^{R493X} mice both showed modest increases in expression of lysosomal proteins (including TFEB, CtsD, PSAP, and LAMP1) at 6 months and 12 months of age (Fig. 1). Similar changes were also observed in the thalamus of Grn^{R493X} mice (Fig. S1). These results indicate that, Grn^{R493X} mice exhibit a broad increase in lysosomal gene expression in the brain, similar to *Grn* mice.

Age-dependent neuroinflammation is a major phenotype that has been observed in multiple *Grn* knockout mouse models,[23,25,29,32,35] although examination in heterozygous mice has been more limited.[30,34,35] For the Grn^{R493X} mouse model, we previously reported age-dependent microgliosis in the thalamus of 12-month-old homozygous mice, as determined by quantification of Iba1[+] cells after immunohistochemistry.[28] Additionally, Frew and Nygaard reported increased microgliosis and astrogliosis in 18-month-old homozygous mice by immunostaining.[37] Here, we report increased expression of the microglia marker Iba1 and astrocyte marker GFAP in the cortex that is apparent in homozygous mice by 12 months of age (Fig. 2). In contrast, heterozygous Grn^{R493X} mice did not show a significant increase. Overall, these observations in Grn^{R493X} mice are consistent with a previous report in 12-month-old *Grn* mice.[35]

We found modest increases in expression of cytokines (including TNFα, IL-1β, and MCP1) in the cortex of homozygous Grn^{R493X} mice at 12 months of age (Fig. 2). Heterozygous mice had intermediate cytokine levels, which were not statistically different than the wild-type mice. Notably, we observed a marked increase in the expression of complement factor C3 in 12-month-old homozygous mice; trends toward increased levels of C1qa and C3 were also seen in heterozygous Grn^{R493X} mice, although the magnitude of the increases was smaller than in

homozygous mice. Together, these results show that *Grn^R493X* mice develop age-dependent neuroinflammation that is similar to previous reports in *Grn* mouse models.

Biomarker levels in the Grn^R493X mouse model

Previous human studies have identified several biomarkers that are increased in the blood or CSF of individuals with FTD, including those with heterozygous *GRN* mutations. These biomarkers include NfL,[12-14] GFAP,[14,18] total TDP-43,[16] and phosphorylated TDP-43.[15,17] In particular, NfL is a neuronal cytoskeletal protein that is released into CSF and blood upon neuronal damage or death. CSF and blood NfL levels have been proposed as an early indicators of neurodegeneration,[49] and their levels reflect FTD disease severity.[12,13] It is currently not known if levels of NfL and other biomarkers are similarly increased in *Grn* and *Grn^R493X* mouse models. To address this, we measured plasma NfL levels in a cohort of *Grn^R493X* mice between 2 months and 18 months of age using the Single Molecule Array (Simoa) platform by Quanterix. As shown in Fig. 3A, we observed an age-dependent increase in plasma NfL levels with all genotypes. We detected elevated levels in homozygous *Grn^R493X* mice beginning at 10-12 months of age. In contrast, plasma NfL levels did not differ between wild-type and heterozygous *Grn^R493X* mice. These trends were also observed when male and female mice were analyzed separately (Figs. S2A and S2B). These trends were also observed with NfL measured using the Single Molecule Counting (SMC) platform by Sigma (Fig. S2C); there was a strong correlation between the measurements from two assays in the 14-month-old plasma samples (r^2 = 0.8541, n=23, p < 0.0001 by two-tailed t test). We measured plasma NfL levels in additional cohorts of 12-month-old and 18-month-old mice. While homozygous *Grn^R493X* mice had increased NfL levels at both ages (Fig. 3B), we again found no differences between wild-type and heterozygous *Grn^R493X* mice at 12 months of age. In 18-month-old mice, we observed a trend toward increased plasma NfL levels in heterozygous *Grn^R493X* mice, although this did not reach statistical significance (p = 0.1105). For comparison, we measured plasma NfL levels in the *Grn* knockout mouse model[23] at 12 and 18 months of age; similarly, we found increased NfL levels in homozygous *Grn* knockout mice but not in heterozygous mice (Fig. S2D). We also measured NfL levels in a limited number of CSF samples; these results show elevated NfL levels in homozygous *Grn^R493X* mice but not in heterozygous mice at 12 months and 18 months of age (Fig. 3C).

We also measured levels of another biomarker, GFAP, in the plasma and CSF of *Grn^R493X* mice. In plasma, GFAP levels were not increased in heterozygous or homozygous *Grn^R493X* mice at 12 months of age (Fig. S3A). In CSF, we observed a trend toward increased GFAP levels in homozygous *Grn^R493X* mice but no increase in heterozygous mice at 12 months of age (Fig. S3B). Collectively, these results suggest that the FTD fluid biomarkers identified in humans are not similarly elevated in the blood and CSF of heterozygous *Grn* and *Grn^R493X* mouse models. Additionally, the NfL results in particular suggest that, compared to their wild-type littermate controls, heterozygous *Grn^R493X* mice do not exhibit increased neuronal death that is detectable by plasma or CSF NfL through 18 months of age.

Behavioral deficits in the Grn^R493X mouse model

Limited behavioral studies have been performed on the Grn^{R493X} mouse model, and no data is currently available for heterozygous mice. Specifically, we have reported excessive grooming behavior, which is a type of obsessive compulsive-like behavior, in the homozygous Grn^{R493X} mice,[28] and Frew and Nygaard have reported increased anxiety in homozygous male mice.[37] Here, we performed additional behavioral studies in 11-month-old male mice to further characterize Grn^{R493X} mice. Similar to previous reports in Grn knockout mouse models,[35,36] we observed decreased sociability in homozygous Grn^{R493X} mice in the three-chamber test (Fig. 4A). A similar trend was observed in heterozygous mice, although the difference was not statistically significant (p = 0.0869). In the tube test of social dominance, we observed decreased social dominance in heterozygous Grn^{R493X} mice, as reflected in the lower percentage of trials won by the heterozygous mice (Fig. 4B); these results mirror previous reports with similarly aged $Grn^{+/-}$ mice.[38,50] In the elevated plus maze, homozygous Grn^{R493X} mice spent less time spent in the open arms, suggestive of increased anxiety (Fig. 4C). This finding is consistent with a previous study in homozygous Grn mice using the elevated plus maze,[51] as well as reports of increased anxiety in homozygous Grn^{R493X} male mice[37] and in homozygous Grn mice[24,26] based on time spent in the center of the open field. We observed no differences between genotypes in other behavioral tests, including open field activity, time spent in the center of the open field, forced swim test, and nest building (Fig. S4). Together, these studies show that Grn^{R493X} mice exhibit social and emotional deficits similar to those previously reported in Grn knockout mouse models.

Lastly, we assesed cognitive function in the Grn^{R493X} mice. Using the T-maze to assess long-term spatial memory, we found impairment in heterozygous and homozygous Grn^{R493X} mice (Fig. 5A). This impairment was refected in the increased number of trials needed to reach the criterion of choosing the correct side of the T-maze 5 out of 6 consecutive times. Using the novel object recognition test (with 1 h retention interval), heterozygous and homozygous Grn^{R493X} mice exhibited impairment in short-term memory, as reflected in the lower discrimation index which is a calculation of the time spent interacting with the novel object (Fig. 5B). Grn^{R493X} mice showed no differences in working memory, as assessed by the Y-maze (Fig. 5C). In the puzzle box test, we found impaired executive function in homozygous Grn^{R493X} mice, but not in the heterozygous mice (Fig. 5D). Together, these results indicate that Grn^{R493X} mice exhibit cognitive impairment in several specific domains. These results are consistent with previous reports of memory impairment in Grn knockout mice as assessed by the Morris water maze,[36] novel object recognition test,[24] and fear conditioning test.[35]

DISCUSSION

In the current study, we have further characterized the Grn^{R493X} knockin mouse model of FTD, including analyses of neuropathology, fluid biomarkers, and behavior. The R493X mutation introduces a premature termination codon, which leads to degradation of the mutant Grn mRNA through the nonsense-mediated RNA decay pathway, and therefore Grn^{R493X} mice have markedly reduced progranulin protein levels.[28] Overall, our results show that Grn^{R493X} knockin mice phenocopy their Grn knockout counterparts, with a mild behavioral phenotype and modest neuropathological changes.

In the brain, heterozygous and homozygous Grn^{R493X} mice had increased expression of lysosomal genes by 6 months of age (Figs. 1 and S1), likely to compensate for impaired lysosomal function and/or homeostasis. Similar increases in lysosomal gene expression have been reported in Grn knockout mice.[29-32] Additionally, lipofuscin has been observed in the brains of both homozygous Grn and Grn^{R493X} mice,[24,28,29,34,37,50,52] further indicating that progranulin deficiency leads to lysosomal dysfunction in these mouse models. It is unclear why the upregulation of some lysosomal genes (i.e., LAMP1, Gba, HexA, PSAP) was diminished at 12 months of age. Whereas the changes in lysosomal gene expression were evident at 6 months of age, increased expression of inflammation-related genes was generally not observed in Grn^{R493X} mice until 12 months of age (Fig. 2). These genes include markers of microgliosis and astrogliosis, pro-inflammatory cytokines, and complement factors. These changes in inflammation-related genes were largely absent in heterozygous Grn^{R493X} mice. Overall, this pattern of neuroinflammation is consistent with previous reports in Grn mouse models.[25,29,34-36] Together, these results suggest that neuroinflammation may be secondary to lysosomal dysfunction in the Grn^{R493X} mouse model.

Our behavioral studies found social and emotional deficits in Grn^{R493X} mice which mirror those found in Grn mouse models. Decreased sociability has been reported in both heterozygous and homozygous Grn knockout mice,[35,36] and we similarly observed this in Grn^{R493X} mice. Decreased social dominance has also been reported in 9- to 16-month-old heterozygous Grn mice,[38,50] and we found Grn^{R493X} mice also exhibit this phenotype in 11-month-old $Grn^{+/R493X}$ mice. A male-specific increase in anxiety has been reported in homozygous Grn knockout mice and homozygous Grn^{R493X} mice.[26,37,51] In the current study, we found evidence of increased anxiety in homozygous Grn^{R493X} male mice in the elevated plus maze (Fig. 4C). Additionally, similar to Grn mice,[39] we previously reported that homozygous Grn^{R493X} mice have increased grooming behavior, which is a type of obssesive compulsive-like behavior in mice.[28] Together, our studies show that Grn^{R493X} mice phenocopy Grn mice with respect to behavior and that they mirror symptoms of FTD-GRN including diminished social interest, apathy, disinhibition, obsessive-compulsive behavior, and anxiety.[9,53]

Memory is typically spared in bvFTD until later the stages of disease, although there are reports of memory impairment.[9,54] Three studies have reported memory impairment in homozygous Grn mice using the Morris water maze, novel object recognition test, and fear conditioning test.[24,35,36] In the current study, we found that both homozygous Grn^{R493X} mice as well as heterozygous mice show impariments in certain types of memory. Specifically, short-term memory and long-term spatial memory are impaired, but working memory was unaffected. In bvFTD, there is often impairment of executive functions which correlates with atrophy in the dorsolateral and medial prefrontal cortex.[55,56] Therefore, we assessed executive function in Grn^{R493X} mice using the puzzle box test, which relies heavily on the prefrontal cortex as well as the hippocampus.[47] Overall, we found that homozygous, but not heterozygous, Grn^{R493X} mice exhibit impaired executive function. These results show that 11-month-old heterozygous and homozygous Grn^{R493X} mice have impairments in specific types of memory, and homozygous mice also have impaired executive function. This supports a previous study finding impaired associative learning and memory in the heterozygous Grn mice.[35]

We sought to test if fluid biomarkers identified in individuals with FTD-*GRN* are similarly elevated in *Grn* and *Grn^R493X* mouse models. We largely focused on NfL, which is a neuronal cytoskeletal protein that has emerged as a promising biomarker that could be used to monitor the progression of neurodegenerative diseases and the effects of disease-modifying treatments.[49] While we observed increased NfL levels in the plasma and CSF of homozygous *Grn^R493X* mice by 10-12 months of age, we did not find increased levels in heterozygous *Grn^R493X* mice through 18 months of age (Figs. 4 and S2A, B). We also measured plasma NfL levels in the *Grn* knockout mouse model and similarly found elevated levels in homozygous but not heterozygous mice (Fig. S2D). Examining levels of another FTD-*GRN* biomarker, we similarly found increased GFAP levels in the CSF of homozygous *Grn^R493X* mice at 12 months of age, but no increase in heterozygous *Grn^R493X* mice (Fig. S3C). Overall, these biomarker results suggest the absence of overt neuronal death in heterozygous *Grn* and *Grn^R493X* mice. This contrasts the the severe atrophy of the frontal and temporal lobes observed in humans with heterozygous *GRN* mutations, which is accompanied by increases in NfL and GFAP in the CSF and plasma. The reasons for these differences between humans and mouse models are unclear. Nonetheless, they suggest that blood NfL and GFAP levels are not useful as biomarkers for pre-clinical studies using the heterozygous *Grn* and *Grn^R493X* mouse models. Thus, it will be worthwhile to validate alternative blood biomarkers, such as compelement C1qa and C3[32,57] or possibly specific lipid species such as gangliosides[58,59] in these mouse models.

We found that heterozygous *Grn^R493X* mice phenocopy heterozygous *Grn* mice reported in two previous studies.[30,35] Notably, they do not exhibit key features of human FTD-*GRN* including neuroinflammation and increased NfL levels in plasma or CSF. Nonetheless, like *Grn* mice, heterozygous *Grn^R493X* mice have social and emotional deficits that recapitulate core symptoms of FTD-*GRN*. The presence of these behavioral deficits without observed increases in NfL levels suggests that in mice the social and emotional deficits may develop independently of overt neuronal death that is detectable by this sensitive biomarker. Taken together, although *Grn* gene dosage effects are also observed in mice, the heterozygous states manifest quite differently in humans and mice; the reasons for this species difference are unclear.

Overall, our findings in homozygous *Grn^R493X* mouse brains are consistent with the phenotypes previously reported in homozygous *Grn* mice. Key features of homozygous *Grn^R493X* mouse from this study and from previous studies include neuroinflammation, microgliosis, astrogliosis, lipofuscin accumulation, vacuolation, upregulation of lysosomal genes, and TDP-43 pathology. Notably, all of these phenotypes have also been observed in mouse models of NCL,[30,60-63] raising the possibility that these observed phenotypes in homozygous *Grn* and *Grn^R493X* mice may reflect NCL rather than FTD-*GRN*. Our results support that, despite their more subtle phenotype, heterozygous *Grn* and *Grn^R493X* mice are likely to be more appropriate models of FTD-*GRN* than the homozygous mice, as previously suggested.[35,64]

Our study has limitations. Our analysis of neuroinflammation and lysosomal dysfunction relied on transcriptional changes; studies assessing protein levels are warranted. For our biomarker studies, it should be noted that the sample sizes were relatively small for measurements in CSF and in 18-month-old mice; therefore, these findings must be interpreted with caution. Our behavioral studies were limited to male mice at 11 months of age; additional studies in female

mice are necessary to address possible sex-specific phenotypes, which have been noted in the *Grn* and *Grn^{R493X}* mouse models.[24,26,37,51]

In conclusion, we have performed in depth characterization of the *Grn^{R493X}* mouse model of FTD, including both heterozygous and homozygous mice. Overall, we found that heterozygous *Grn^{R493X}* and homozygous knockin mice phenocopy their *Grn* counterparts with respect to neuropathology, levels of a plasma biomarker of neuronal damage, and behavioral and cognitive deficits. In contrast to human FTD-*GRN*, heterozygous *Grn^{R493X}* and *Grn* mice do not show elevated levels of plasma NfL through 18 months of age, suggesting they do not have increased neuronal death compared to wild-type mice. These results may help to inform pre-clinical studies using this and other *Grn* mouse models.

ACKNOWLEDGEMENTS

We thank Robert Farese, Jr., Tobias Walther, John Morley, and Thi Nguyen for helpful discussions, the Bluefield Project to Cure FTD for *Grn* mouse plasma, the NeuroBehavior Laboratory at the Harvard NeuroDiscovery Center, the Quanterix Accelerator Lab, and Millipore Sigma for plasma NfL measurements using the SMC platform. This work was supported by grants from the National Institutes of Health (AG047339 and AG064069) and the Bluefield Project to Cure FTD to ADN. Research reported in this publication was also supported by the Washington University Institute of Clinical and Translational Sciences grant UL1TR002345 from the National Center for Advancing Translational Sciences (NCATS) of the National Institutes of Health.

FIGURE LEGENDS

Figure 1. *Grn^{R493X}* mice have increased expression of multiple lysosomal genes in the cortex. Quantification of mRNA levels by qPCR. Bars represent mean ± SEM (n = 5-10 per group, male and female mice); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$ compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn^{+/R493X}* heterozygous mice; KI, *Grn^{R493X/R493X}* knockin mice; n.s., not significant.

Figure 2. *Grn^{R493X}* mice exhibit age-dependent neuroinflammation in the cortex. Quantification of mRNA levels by qPCR. Bars represent mean ± SEM (n = 5-10 per group, male and female mice); * $p < 0.05$ and ** $p < 0.01$ compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn^{+/R493X}* heterozygous mice; KI, *Grn^{R493X/R493X}* knockin mice; n.s., not significant.

Figure 3. Homozygous *Grn^{R493X}* mice, but not heterozygous mice, have increased NfL levels in blood and CSF. A) Plasma NfL levels in an aging cohort of mice (n = 6-10 per group, male and female mice). B) Plasma NfL levels in additional cohorts of mice (n = 15-17 per group at 12 months old, n = 12-20 per group at 18 months old, male and female mice). C) CSF NfL levels (n = 3-8 per group at 12 months old, male and female mice presented separately; n = 1-5 per group at 18 months old, male and female mice). Bars represent mean ± SEM; * $p < 0.05$, **

p < 0.01, and *** p < 0.001 compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice; n.s., not significant.

Figure 4. *Grn*^{R493X} mice exhibit social and emotional deficits. A) Three-chamber test of sociability (n = 9 per group, male mice only, 11 months old). B) Tube test of social dominance (n = 15 per group, male mice only, 8-11 months old). Percent of trials won is shown. C) Elevated plus maze (n = 11 per group, male mice only, 11 months old). Bars represent mean ± SEM; * p < 0.05 compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test (in A and C). Results were analyzed by two-way binomial test (in B). WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice.

Figure 5. *Grn*^{R493X} mice exhibit memory impairment at 11 months of age. A) T-maze retention (n = 11-12 per group, male mice only). B) Novel object recognition with 1 h retention interval (n = 8 per group, male mice only). C) Y-maze (n = 11-12 per group, male mice only). D) Puzzle box (n = 10-12 per group, male mice only). Bars represent mean ± SEM; ** p < 0.01 and *** p < 0.001 compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice.

Figure S1. *Grn* mice have increased expression of several lysosomal genes in the thalamus. Quantification of mRNA levels by RT-qPCR. Bars represent mean ± SEM (n = 5-8 per group, male and female mice); * p < 0.05, ** p < 0.01, and **** p < 0.0001 compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice.

Figure S2. Homozygous *Grn*^{R493X} mice, but not heterozygous mice, have increased plasma NfL levels. A-B) Plasma NfL levels at different ages and separated by sex measured using the Quanterix Simoa platform (n = 3-7 per group for male mice, n = 4-5 per group for female mice). C) Plasma NfL levels at different ages and separated by sex measured using the Sigma SMC platform (n = 3-6 per group for male mice, n = 3-4 per group for female mice). D) Plasma NfL levels in *Grn* knockout mouse model measured using the Quanterix Simoa platform (n = 4-6 per group at 12 months old, n = 6-8 per group at 18 months old, male and female mice). Bars represent mean ± SEM; * p < 0.05 and ** p < 0.01 compared to wild-type group, as determined by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice.

Figure S3. Plasma and CSF levels of GFAP in *Grn*^{R493X} mice. A) Plasma GFAP levels (n = 6-7 per group at 12 months old, male and female mice; n = 4 per group at 18 months old, female mice only). B) CSF GFAP levels (n = 3-4 per group, female mice only). Bars represent mean ± SEM; statistical analysis was performed by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice.

Figure S4. *Grn*^{R493X} mice show no differences in other behavioral tests performed. Bars represent mean ± SEM (n= 8-12 per group, male mice only, 11 months old); statistical analysis was performed by one-way ANOVA with Dunnett post hoc test. WT, wild-type; Het, *Grn*^{+/R493X} heterozygous mice; KI, *Grn*^{R493X/R493X} knockin mice.

REFERENCES

1. Petkau TL, Leavitt BR. Progranulin in neurodegenerative disease. *Trends Neurosci.* 2014;37(7):388-398.
2. Bateman A, Cheung ST, Bennett HPJ. A brief overview of progranulin in health and disease. *Methods Mol Biol.* 2018;1806:3-15.
3. Cenik B, Sephton CF, Kutluk Cenik B, Herz J, Yu G. Progranulin: a proteolytically processed protein at the crossroads of inflammation and neurodegeneration. *J Biol Chem.* 2012;287(39):32298-32306.
4. Kao AW, McKay A, Singh PP, Brunet A, Huang EJ. Progranulin, lysosomal regulation and neurodegenerative disease. *Nat Rev Neurosci.* 2017;18(6):325-333.
5. Nguyen AD, Nguyen TA, Martens LH, Mitic LL, Farese RV, Jr. Progranulin: at the interface of neurodegenerative and metabolic diseases. *Trends Endocrinol Metab.* 2013;24(12):597-606.
6. Baker M, Mackenzie IR, Pickering-Brown SM, Gass J, Rademakers R, Lindholm C, Snowden J, Adamson J, Sadovnick AD, Rollinson S, Cannon A, Dwosh E, Neary D, Melquist S, Richardson A, Dickson D, Berger Z, Eriksen J, Robinson T, Zehr C, Dickey CA, Crook R, McGowan E, Mann D, Boeve B, Feldman H, Hutton M. Mutations in progranulin cause tau-negative frontotemporal dementia linked to chromosome 17. *Nature.* 2006;442(7105):916-919.
7. Cruts M, Gijselinck I, van der Zee J, Engelborghs S, Wils H, Pirici D, Rademakers R, Vandenberghe R, Dermaut B, Martin JJ, van Duijn C, Peeters K, Sciot R, Santens P, De Pooter T, Mattheijssens M, Van den Broeck M, Cuijt I, Vennekens K, De Deyn PP, Kumar-Singh S, Van Broeckhoven C. Null mutations in progranulin cause ubiquitin-positive frontotemporal dementia linked to chromosome 17q21. *Nature.* 2006;442(7105):920-924.
8. Gass J, Cannon A, Mackenzie IR, Boeve B, Baker M, Adamson J, Crook R, Melquist S, Kuntz K, Petersen R, Josephs K, Pickering-Brown SM, Graff-Radford N, Uitti R, Dickson D, Wszolek Z, Gonzalez J, Beach TG, Bigio E, Johnson N, Weintraub S, Mesulam M, White CL, 3rd, Woodruff B, Caselli R, Hsiung GY, Feldman H, Knopman D, Hutton M, Rademakers R. Mutations in progranulin are a major cause of ubiquitin-positive frontotemporal lobar degeneration. *Hum Mol Genet.* 2006;15(20):2988-3001.
9. Miller B, Llibre Guerra JJ. Frontotemporal dementia. *Handb Clin Neurol.* 2019;165:33-45.
10. Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, Bruce J, Schuck T, Grossman M, Clark CM, McCluskey LF, Miller BL, Masliah E, Mackenzie IR, Feldman H, Feiden W, Kretzschmar HA, Trojanowski JQ, Lee VM. Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science.* 2006;314(5796):130-133.
11. Bevan-Jones WR, Cope TE, Jones PS, Kaalund SS, Passamonti L, Allinson K, Green O, Hong YT, Fryer TD, Arnold R, Coles JP, Aigbirhio FI, Larner AJ, Patterson K, O'Brien JT, Rowe JB. Neuroinflammation and protein aggregation co-localize across the frontotemporal dementia spectrum. *Brain.* 2020;143(3):1010-1026.
12. Rohrer JD, Woollacott IOC, Dick KM, Brotherhood E, Gordon E, Fellows A, Toombs J, Druyeh R, Cardoso MJ, Ourselin S, Nicholas JM, Norgren N, Mead S, Andreasson U, Blennow K, Schott JM, Fox NC, Warren JD, Zetterberg H. Serum neurofilament light chain protein is a measure of disease intensity in frontotemporal dementia. *Neurology.* 2016;87(13):1329–1336.
13. Meeter LH, Dopper EG, Jiskoot LC, Sanchez-Valle R, Graff C, Benussi L, Ghidoni R, Pijnenburg YA, Borroni B, Galimberti D, Laforce R, Jr, Masellis M, Vandenberghe R, Ber IL, Otto M, Minkelen R, Papma JM, Rombouts SA, Balasa M, Öijerstedt L, Jelic V, Dick

KM, Cash DM, Harding SR, Jorge Cardoso M, Ourselin S, Rossor MN, Padovani A, Scarpini E, Fenoglio C, Tartaglia MC, Lamari F, Barro C, Kuhle J, Rohrer JD, Teunissen CE, Swieten JC. Neurofilament light chain: a biomarker for genetic frontotemporal dementia. *Ann Clin Transl Neurol.* 2016;3(8):623-636.

14.	Zhu N, Santos-Santos M, Illán-Gala I, Montal V, Estellés T, Barroeta I, Altuna M, Arranz J, Muñoz L, Belbin O, Sala I, Sánchez-Saudinós MB, Subirana A, Videla L, Pegueroles J, Blesa R, Clarimón J, Carmona-Iragui M, Fortea J, Lleó A, Alcolea D. Plasma glial fibrillary acidic protein and neurofilament light chain for the diagnostic and prognostic evaluation of frontotemporal dementia. *Transl Neurodegener.* 2021;10(1):50.

15.	Suárez-Calvet M, Dols-Icardo O, Lladó A, Sánchez-Valle R, Hernández I, Amer G, Antón-Aguirre S, Alcolea D, Fortea J, Ferrer I, van der Zee J, Dillen L, Van Broeckhoven C, Molinuevo JL, Blesa R, Clarimón J, Lleó A. Plasma phosphorylated TDP-43 levels are elevated in patients with frontotemporal dementia carrying a C9orf72 repeat expansion or a GRN mutation. *J Neurol Neurosurg Psychiatry.* 2014;85(6):684-691.

16.	Foulds P, McAuley E, Gibbons L, Davidson Y, Pickering-Brown SM, Neary D, Snowden JS, Allsop D, Mann DM. TDP-43 protein in plasma may index TDP-43 brain pathology in Alzheimer's disease and frontotemporal lobar degeneration. *Acta Neuropathol.* 2008;116(2):141-146.

17.	Foulds PG, Davidson Y, Mishra M, Hobson DJ, Humphreys KM, Taylor M, Johnson N, Weintraub S, Akiyama H, Arai T, Hasegawa M, Bigio EH, Benson FE, Allsop D, Mann DM. Plasma phosphorylated-TDP-43 protein levels correlate with brain pathology in frontotemporal lobar degeneration. *Acta Neuropathol.* 2009;118(5):647-658.

18.	Heller C, Foiani MS, Moore K, Convery R, Bocchetta M, Neason M, Cash DM, Thomas D, Greaves CV, Woollacott IO, Shafei R, Van Swieten JC, Moreno F, Sanchez-Valle R, Borroni B, Laforce R, Jr., Masellis M, Tartaglia MC, Graff C, Galimberti D, Rowe JB, Finger E, Synofzik M, Vandenberghe R, de Mendonca A, Tagliavini F, Santana I, Ducharme S, Butler CR, Gerhard A, Levin J, Danek A, Frisoni G, Sorbi S, Otto M, Heslegrave AJ, Zetterberg H, Rohrer JD. Plasma glial fibrillary acidic protein is raised in progranulin-associated frontotemporal dementia. *J Neurol Neurosurg Psychiatry.* 2020;91(3):263-270.

19.	Rojas JC, Wang P, Staffaroni AM, Heller C, Cobigo Y, Wolf A, Goh SM, Ljubenkov PA, Heuer HW, Fong JC, Taylor JB, Veras E, Song L, Jeromin A, Hanlon D, Yu L, Khinikar A, Sivasankaran R, Kieloch A, Valentin MA, Karydas AM, Mitic LL, Pearlman R, Kornak J, Kramer JH, Miller BL, Kantarci K, Knopman DS, Graff-Radford N, Petrucelli L, Rademakers R, Irwin DJ, Grossman M, Ramos EM, Coppola G, Mendez MF, Bordelon Y, Dickerson BC, Ghoshal N, Huey ED, Mackenzie IR, Appleby BS, Domoto-Reilly K, Hsiung GR, Toga AW, Weintraub S, Kaufer DI, Kerwin D, Litvan I, Onyike CU, Pantelyat A, Roberson ED, Tartaglia MC, Foroud T, Chen W, Czerkowicz J, Graham DL, van Swieten JC, Borroni B, Sanchez-Valle R, Moreno F, Laforce R, Graff C, Synofzik M, Galimberti D, Rowe JB, Masellis M, Finger E, Vandenberghe R, de Mendonça A, Tagliavini F, Santana I, Ducharme S, Butler CR, Gerhard A, Levin J, Danek A, Otto M, Sorbi S, Cash DM, Convery RS, Bocchetta M, Foiani M, Greaves CV, Peakman G, Russell L, Swift I, Todd E, Rohrer JD, Boeve BF, Rosen HJ, Boxer AL. Plasma neurofilament light for prediction of disease progression in familial frontotemporal lobar degeneration. *Neurology.* 2021;96(18):e2296-e2312.

20.	Smith KR, Damiano J, Franceschetti S, Carpenter S, Canafoglia L, Morbin M, Rossi G, Pareyson D, Mole SE, Staropoli JF, Sims KB, Lewis J, Lin WL, Dickson DW, Dahl HH, Bahlo M, Berkovic SF. Strikingly different clinicopathological phenotypes determined by progranulin-mutation dosage. *Am J Hum Genet.* 2012;90(6):1102-1107.

21.	Almeida M, Macario M, Ramos L, Baldeiras I, Ribeiro M, Santana I. Portuguese family with the co-occurrence of frontotemporal lobar degeneration and neuronal ceroid

lipofuscinosis phenotypes due to progranulin gene mutation. *Neurobiol Aging.* 2016;41:200 e201-205.

22. Huin V, Barbier M, Bottani A, Lobrinus JA, Clot F, Lamari F, Chat L, Rucheton B, Fluchère F, Auvin S, Myers P, Gelot A, Camuzat A, Caillaud C, Jornéa L, Forlani S, Saracino D, Duyckaerts C, Brice A, Durr A, Le Ber I. Homozygous GRN mutations: new phenotypes and new insights into pathological and molecular mechanisms. *Brain.* 2020;143(1):303-319.

23. Martens LH, Zhang J, Barmada SJ, Zhou P, Kamiya S, Sun B, Min SW, Gan L, Finkbeiner S, Huang EJ, Farese RV, Jr. Progranulin deficiency promotes neuroinflammation and neuron loss following toxin-induced injury. *J Clin Invest.* 2012;122(11):3955-3959.

24. Petkau TL, Neal SJ, Milnerwood A, Mew A, Hill AM, Orban P, Gregg J, Lu G, Feldman HH, Mackenzie IR, Raymond LA, Leavitt BR. Synaptic dysfunction in progranulin-deficient mice. *Neurobiol Dis.* 2012;45(2):711-722.

25. Yin F, Banerjee R, Thomas B, Zhou P, Qian L, Jia T, Ma X, Ma Y, Iadecola C, Beal MF, Nathan C, Ding A. Exaggerated inflammation, impaired host defense, and neuropathology in progranulin-deficient mice. *J Exp Med.* 2010;207(1):117-128.

26. Kayasuga Y, Chiba S, Suzuki M, Kikusui T, Matsuwaki T, Yamanouchi K, Kotaki H, Horai R, Iwakura Y, Nishihara M. Alteration of behavioural phenotype in mice by targeted disruption of the progranulin gene. *Behav Brain Res.* 2007;185(2):110-118.

27. Fujita K, Chen X, Homma H, Tagawa K, Amano M, Saito A, Imoto S, Akatsu H, Hashizume Y, Kaibuchi K, Miyano S, Okazawa H. Targeting Tyro3 ameliorates a model of PGRN-mutant FTLD-TDP via tau-mediated synaptic pathology. *Nat Commun.* 2018;9(1):433.

28. Nguyen AD, Nguyen TA, Zhang J, Devireddy S, Zhou P, Xu X, Karydas AM, Miller BL, Rigo F, Ferguson SM, Huang EJ, Walther TC, Farese RV, Jr. Murine knockin model for progranulin-deficient frontotemporal dementia with nonsense-mediated mRNA decay. *Proc Natl Acad Sci U S A.* 2018;115(12):E2849-E2858.

29. Tanaka Y, Chambers JK, Matsuwaki T, Yamanouchi K, Nishihara M. Possible involvement of lysosomal dysfunction in pathological changes of the brain in aged progranulin-deficient mice. *Acta Neuropathol Commun.* 2014;2:78.

30. Götzl JK, Mori K, Damme M, Fellerer K, Tahirovic S, Kleinberger G, Janssens J, van der Zee J, Lang CM, Kremmer E, Martin JJ, Engelborghs S, Kretzschmar HA, Arzberger T, Van Broeckhoven C, Haass C, Capell A. Common pathobiochemical hallmarks of progranulin-associated frontotemporal lobar degeneration and neuronal ceroid lipofuscinosis. *Acta Neuropathol.* 2014;127(6):845-860.

31. Chang MC, Srinivasan K, Friedman BA, Suto E, Modrusan Z, Lee WP, Kaminker JS, Hansen DV, Sheng M. Progranulin deficiency causes impairment of autophagy and TDP-43 accumulation. *J Exp Med.* 2017;214(9):2611-2628.

32. Lui H, Zhang J, Makinson S, Cahill M, Kelley K, Huang H, Shang Y, Oldham M, Martens L, Gao F, Coppola G, Sloan S, Hsieh C, Kim C, Bigio E, Weintraub S, Mesulam M, Rademakers R, Mackenzie I, Seeley W, Karydas A, Miller B, Borroni B, Ghidoni R, Farese R, Jr, Paz J, Barres B, Huang E. Progranulin deficiency promotes circuit-specific synaptic pruning by microglia via complement activation. *Cell.* 2016;165(4):921-935.

33. Ward ME, Chen R, Huang HY, Ludwig C, Telpoukhovskaia M, Taubes A, Boudin H, Minami SS, Reichert M, Albrecht P, Gelfand JM, Cruz-Herranz A, Cordano C, Alavi MV, Leslie S, Seeley WW, Miller BL, Bigio E, Mesulam MM, Bogyo MS, Mackenzie IR, Staropoli JF, Cotman SL, Huang EJ, Gan L, Green AJ. Individuals with progranulin haploinsufficiency exhibit features of neuronal ceroid lipofuscinosis. *Sci Transl Med.* 2017;9(385).

34. Ahmed Z, Sheng H, Xu YF, Lin WL, Innes AE, Gass J, Yu X, Wuertzer CA, Hou H, Chiba S, Yamanouchi K, Leissring M, Petrucelli L, Nishihara M, Hutton ML, McGowan E, Dickson DW, Lewis J. Accelerated lipofuscinosis and ubiquitination in granulin knockout mice suggest a role for progranulin in successful aging. *Am J Pathol.* 2010;177(1):311-324.

35. Filiano AJ, Martens LH, Young AH, Warmus BA, Zhou P, Diaz-Ramirez G, Jiao J, Zhang Z, Huang EJ, Gao FB, Farese RV, Jr., Roberson ED. Dissociation of frontotemporal dementia-related deficits and neuroinflammation in progranulin haploinsufficient mice. *J Neurosci.* 2013;33(12):5352-5361.

36. Yin F, Dumont M, Banerjee R, Ma Y, Li H, Lin MT, Beal MF, Nathan C, Thomas B, Ding A. Behavioral deficits and progressive neuropathology in progranulin-deficient mice: a mouse model of frontotemporal dementia. *FASEB J.* 2010;24(12):4639-4647.

37. Frew J, Nygaard HB. Neuropathological and behavioral characterization of aged Grn R493X progranulin-deficient frontotemporal dementia knockin mice. *Acta Neuropathol Commun.* 2021;9(1):57.

38. Arrant A, Filiano A, Warmus B, Hall A, Roberson E. Progranulin haploinsufficiency causes biphasic social dominance abnormalities in the tube test. *Genes Brain Behav.* 2016;15(6):588-603.

39. Krabbe G, Minami SS, Etchegaray JI, Taneja P, Djukic B, Davalos D, Le D, Lo I, Zhan L, Reichert MC, Sayed F, Merlini M, Ward ME, Perry DC, Lee SE, Sias A, Parkhurst CN, Gan WB, Akassoglou K, Miller BL, Farese RV, Jr, Gan L. Microglial NFκB-TNFα hyperactivation induces obsessive-compulsive behavior in mouse models of progranulin-deficient frontotemporal dementia. *Proc Natl Acad Sci U S A.* 2017;114(19):5029-5034.

40. Ghoshal N, Dearborn JT, Wozniak DF, Cairns NJ. Core features of frontotemporal dementia recapitulated in progranulin knockout mice. *Neurobiol Dis.* 2012;45(1):395-408.

41. Deacon R. Assessing burrowing, nest construction, and hoarding in mice. *J Vis Exp.* 2012;59:e2607.

42. Moy SS, Nadler JJ, Perez A, Barbaro RP, Johns JM, Magnuson TR, Piven J, Crawley JN. Sociability and preference for social novelty in five inbred strains: an approach to assess autistic-like behavior in mice. *Genes, Brain and Behavior.* 2004;3(5):287-302.

43. Farr SA, Niehoff ML, Kumar VB, Roby DA, Morley JE. Inhibition of glycogen synthase kinase 3beta as a treatment for the prevention of cognitive deficits after a traumatic brain injury. *J Neurotrauma.* 2019;36(11):1869-1875.

44. Farr SA, Banks WA, La Scola ME, Flood JF, Morley JE. Permanent and temporary inactivation of the hippocampus impairs T-maze footshock avoidance acquisition and retention. *Brain Res.* 2000;872(1-2):242-249.

45. Kraeuter AK, Guest PC, Sarnyai Z. The Y-maze for assessment of spatial working and reference memory in mice. *Methods Mol Biol.* 2019;1916:105-111.

46. Hammond RS, Tull LE, Stackman RW. On the delay-dependent involvement of the hippocampus in object recognition memory. *Neurobiol Learn Mem.* 2004;82(1):26-34.

47. Ben Abdallah NM, Fuss J, Trusel M, Galsworthy MJ, Bobsin K, Colacicco G, Deacon RM, Riva MA, Kellendonk C, Sprengel R, Lipp HP, Gass P. The puzzle box as a simple and efficient behavioral test for exploring impairments of general cognition and executive functions in mouse models of schizophrenia. *Exp Neurol.* 2011;227(1):42-52.

48. O'Connor AM, Burton TJ, Leamey CA, Sawatari A. The use of the puzzle box as a means of assessing the efficacy of environmental enrichment. *J Vis Exp.* 2014(94).

49. Khalil M, Teunissen CE, Otto M, Piehl F, Sormani MP, Gattringer T, Barro C, Kappos L, Comabella M, Fazekas F, Petzold A, Blennow K, Zetterberg H, Kuhle J. Neurofilaments as biomarkers in neurological disorders. *Nat Rev Neurol.* 2018;14(10):577-589.

50.   Arrant AE, Filiano AJ, Unger DE, Young AH, Roberson ED. Restoring neuronal progranulin reverses deficits in a mouse model of frontotemporal dementia. *Brain.* 2017;140(5):1447-1465.
51.   Chiba S, Matsuwaki T, Yamanouchi K, Nishihara M. Alteration in anxiety with relation to the volume of the locus ceruleus in progranulin-deficient mice. *J Reprod Dev.* 2009;55(5):518-522.
52.   Petkau TL, Hill A, Leavitt BR. Core neuropathological abnormalities in progranulin-deficient mice are penetrant on multiple genetic backgrounds. *Neuroscience.* 2016;315:175-195.
53.   Benussi A, Premi E, Gazzina S, Brattini C, Bonomi E, Alberici A, Jiskoot L, van Swieten JC, Sanchez-Valle R, Moreno F, Laforce R, Graff C, Synofzik M, Galimberti D, Masellis M, Tartaglia C, Rowe JB, Finger E, Vandenberghe R, de Mendonça A, Tagliavini F, Santana I, Ducharme S, Butler CR, Gerhard A, Levin J, Danek A, Otto M, Frisoni G, Ghidoni R, Sorbi S, Le Ber I, Pasquier F, Peakman G, Todd E, Bocchetta M, Rohrer JD, Borroni B. Progression of behavioral disturbances and neuropsychiatric symptoms in patients with genetic frontotemporal dementia. *JAMA Netw Open.* 2021;4(1):e2030194.
54.   Hodges JR, Davies RR, Xuereb JH, Casey B, Broe M, Bak TH, Kril JJ, Halliday GM. Clinicopathological correlates in frontotemporal dementia. *Ann Neurol.* 2004;56(3):399-406.
55.   Possin KL, Feigenbaum D, Rankin KP, Smith GE, Boxer AL, Wood K, Hanna SM, Miller BL, Kramer JH. Dissociable executive functions in behavioral variant frontotemporal and Alzheimer dementias. *Neurology.* 2013;80(24):2180-2185.
56.   Rosen HJ, Allison SC, Schauer GF, Gorno-Tempini ML, Weiner MW, Miller BL. Neuroanatomical correlates of behavioural disorders in dementia. *Brain.* 2005;128(Pt 11):2612-2625.
57.   van der Ende EL, Heller C, Sogorb-Esteve A, Swift IJ, McFall D, Peakman G, Bouzigues A, Poos JM, Jiskoot LC, Panman JL, Papma JM, Meeter LH, Dopper EGP, Bocchetta M, Todd E, Cash D, Graff C, Synofzik M, Moreno F, Finger E, Sanchez-Valle R, Vandenberghe R, Laforce R, Jr., Masellis M, Tartaglia MC, Rowe JB, Butler C, Ducharme S, Gerhard A, Danek A, Levin J, Pijnenburg YAL, Otto M, Borroni B, Tagliavini F, de Mendonça A, Santana I, Galimberti D, Sorbi S, Zetterberg H, Huang E, van Swieten JC, Rohrer JD, Seelaar H. Elevated CSF and plasma complement proteins in genetic frontotemporal dementia: results from the GENFI study. *J Neuroinflammation.* 2022;19(1):217.
58.   Boland S, Swarup S, Ambaw YA, Malia PC, Richards RC, Fischer AW, Singh S, Aggarwal G, Spina S, Nana AL, Grinberg LT, Seeley WW, Surma MA, Klose C, Paulo JA, Nguyen AD, Harper JW, Walther TC, Farese RV, Jr. Deficiency of the frontotemporal dementia gene GRN results in gangliosidosis. *Nat Commun.* 2022;13(1):5924.
59.   Logan T, Simon MJ, Rana A, Cherf GM, Srivastava A, Davis SS, Low RLY, Chiu CL, Fang M, Huang F, Bhalla A, Llapashtica C, Prorok R, Pizzo ME, Calvert MEK, Sun EW, Hsiao-Nakamoto J, Rajendra Y, Lexa KW, Srivastava DB, van Lengerich B, Wang J, Robles-Colmenares Y, Kim DJ, Duque J, Lenser M, Earr TK, Nguyen H, Chau R, Tsogtbaatar B, Ravi R, Skuja LL, Solanoy H, Rosen HJ, Boeve BF, Boxer AL, Heuer HW, Dennis MS, Kariolis MS, Monroe KM, Przybyla L, Sanchez PE, Meisner R, Diaz D, Henne KR, Watts RJ, Henry AG, Gunasekaran K, Astarita G, Suh JH, Lewcock JW, DeVos SL, Di Paolo G. Rescue of a lysosomal storage disorder caused by Grn loss of function with a brain penetrant progranulin biologic. *Cell.* 2021;184(18):4651-4668.e4625.
60.   Chien HF, Rodriguez RD, Bonifati V, Nitrini R, Pasqualucci CA, Gelpi E, Barbosa ER. Neuropathologic findings in a patient with juvenile-onset levodopa-responsive parkinsonism due to ATP13A2 mutation. *Neurology.* 2021;97(16):763-766.

61. Takahashi K, Nelvagal HR, Lange J, Cooper JD. Glial dysfunction and its contribution to the pathogenesis of the neuronal ceroid lipofuscinoses. *Front Neurol.* 2022;13:886567.

62. Nakanishi H, Zhang J, Koike M, Nishioku T, Okamoto Y, Kominami E, von Figura K, Peters C, Yamamoto K, Saftig P, Uchiyama Y. Involvement of nitric oxide released from microglia-macrophages in pathological changes of cathepsin D-deficient mice. *J Neurosci.* 2001;21(19):7526-7533.

63. Partanen S, Haapanen A, Kielar C, Pontikis C, Alexander N, Inkinen T, Saftig P, Gillingwater TH, Cooper JD, Tyynelä J. Synaptic changes in the thalamocortical system of cathepsin D-deficient mice: a model of human congenital neuronal ceroid-lipofuscinosis. *J Neuropathol Exp Neurol.* 2008;67(1):16-29.

64. Kashyap SN, Boyle NR, Roberson ED. Preclinical interventions in mouse models of frontotemporal dementia due to progranulin mutations. *Neurotherapeutics.* 2023;20(1):140-153.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5