Title: **Pseudouridine guides germline small RNA transport and epigenetic inheritance.**

Authors: Rowan P Herridge[1, ‡*], Jakub Dolata[1,4*], Valentina Migliori[2,3*], Cristiane de Santis Alves[1*], Filipe Borges[1], Frédéric Van Ex[1], Ann Lin[1], Mateusz Bajczyk[4], Tommaso Leonardi[2,5], Alan Hendrick[6], Andrea J Schorn[7], Tony Kouzarides[2**], and Robert A Martienssen[1**]

Affiliations:

[1]Howard Hughes Medical Institute, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 11724, USA

[2]The Gurdon Institute, University of Cambridge, Tennis Court Road, Cambridge CB2 1QN, UK

[3]present address: Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, UK

[4] Department of Gene Expression, Institute of Molecular Biology and Biotechnology, Adam Mickiewicz University, Poznan, Poland

[5]Center for Genomic Science of IIT@SEMM, Instituto Italiano di Tecnologia (IIT), 20139 Milan, Italy

[6]Storm Therapeutics, Ltd., Moneta Building (B280), Babraham Research Campus, Cambridge CB22 3AT, UK

[7]Cold Spring Harbor Laboratory, Cold Spring Harbor NY11724

‡Present address: Department of Biochemistry, University of Otago, Dunedin, NZ

*these authors contributed equally

**For correspondence: t.kouzarides@gurdon.cam.ac.uk, martiens@cshl.edu

Summary:

Epigenetic modifications that arise during plant and animal development, such as DNA and

histone modification, are mostly reset during gamete formation, but some are inherited from the

germline including those marking imprinted genes[1,2]. Small RNAs guide these epigenetic

modifications, and some are also inherited by the next generation[3,4]. In *C. elegans*, inherited

small RNA precursors have poly (UG) tails[5], but how inherited small RNAs are distinguished in

other animals and plants is unknown. Pseudouridine (Ψ) is the most abundant RNA modification

but has not been explored in small RNAs. Here, we develop novel assays to detect Ψ in short

RNA sequences, demonstrating its presence in mouse and *Arabidopsis* microRNAs and their

precursors. We also detect substantial enrichment in germline small RNAs, namely

epigenetically activated siRNAs (easiRNAs) in *Arabidopsis* pollen, and piwi-interacting piRNAs

in mouse testis. In pollen, pseudouridylated easiRNAs are localized to sperm cells, and we found

that *PAUSED/HEN5 (PSD)*, the plant homolog of Exportin-t, interacts genetically with Ψ and is

required for transport of easiRNAs into sperm cells from the vegetative nucleus. We further

show that Exportin-t is required for the triploid block: chromosome dosage-dependent seed

lethality that is epigenetically inherited from pollen. Thus, Ψ has a conserved role in marking

inherited small RNAs in the germline.

One-Sentence Summary: Pseudouridine marks germline small RNAs in plants and mammals,

impacting epigenetic inheritance via nuclear transport.

The germlines of both plants and metazoans have abundant small RNA, many of which are cell-type specific. In the mammalian germline, spermatids accumulate 26-28 nt PIWI-interacting RNAs (piRNAs) which silence TEs, while spermatocytes accumulate 29-30nt piRNA that also target genes[6]. In plants, the male germline terminates in mature pollen, which contains two cell types, the pollen grain (or vegetative cell), and two sperm cells enclosed within its cytoplasm. Plants do not have piRNAs, but instead 21-22 nt epigenetically activated siRNAs (easiRNAs) are produced from TEs in the vegetative nucleus (VN) and translocated to the sperm cells[7-9]. Furthermore, 24 nt siRNAs from some DNA transposons are translocated into developing meiocytes in the anther from the surrounding layer of somatic cells (the tapetum)[8]. Both types of small RNA are produced by the plant-specific RNA polymerase, Pol IV[8,10,11] and mediate epigenetic inheritance after fertilization [10,12]. In maize, 21 nt and 24 nt "phasiRNAs" are also transported from epidermal and tapetal cells into the meiocytes, demonstrating that these mechanisms may be conserved[13]. Both miRNAs and siRNAs in plants are modified by 2'-O-methylation, while in animals only piRNAs have this modification[14].

We set out to determine whether pseudouridine (Ψ) was present in small RNAs. Ψ is the most common non-canonical base in RNA but has not been explored in small RNAs due to difficulties in detection. High-throughput techniques to detect pseudouridylation in longer RNAs specifically label Ψ with N-cyclohexyl-N'-(2-morpholinoethyl)-carbodiimide-metho-*p*-toluenesulfonate (CMCT) to block reverse transcriptase (RT) during RNA-seq library preparation[15-18]. However, in the case of small RNA, RT blocks would result in truncated sequences too short to map back to the genome. More recently, anti-Ψ antibody binding[19], as

well as bisulphite treatment at neutral pH[20] achieve similar results. In mouse, long primary

miRNA precursor transcripts (pri-miRNAs) have been shown to contain Ψ [21,22], so we initially

tested for the presence of Ψ in small RNAs from mouse NIH/3T3 cells. Following

immunoprecipitation with a Ψ-specific antibody, we could robustly detect Ψ using a combination

of mass spectrometry (Extended Data Fig.1a) and dotblots (Extended Data Fig. 1b) using *in vitro*

synthesized pseudouridylated transcripts as a positive control. Using this antibody, we then

performed pseudouridine immunoprecipitation (Ψ-IP) and sequencing of small RNAs from

NIH/3T3 cells before and after knockdown of the Ψ synthase PUS1 (Extended Data Fig. 1c-e;

Supplementary Dataset S3). We detected Ψ in several miRNAs (Extended Data Fig. 1c, d) four

of which were strongly impacted in the *PUS1* knockdown (Fig. 1e). Members of the let-7 family

were heavily pseudouridylated (Extended Data Fig. 1d), but unaffected by *PUS1* knockdown,

consistent with let-7 precursor interaction with DKC1 and TruB1 instead[21,22].

In search of factors that might modify small RNAs in plants, we performed a yeast two-hybrid

screen for interactors of *Arabidopsis* ARGONAUTE 3 (AGO3) using a cDNA library of

transcripts from *Arabidopsis* flowers[23]. The screen (Methods) identified a potential interactor of

AGO3, namely H/ACA RIBONUCLEOPROTEIN COMPLEX SUBUNIT 2 (NHP2), and we

confirmed the interaction by bimolecular fluorescence complementation in phase-separated

nucleolar bodies (Fig. 1a). NHP2 plays a key role in RNA pseudouridylation via the

pseudouridine synthase Dyskerin (DKC1), often, though not always, via snoRNA guides [24-26].

NHP2 was also in close proximity with other AGO proteins, notably AGO4, which has been

found in nucleolar bodies and in related Cajal bodies (Extended Data Fig. 1f)[27].

Immunoprecipitation of GFP-tagged NHP2 followed by mass spectrometry revealed interactions

with other subunits of the Dyskerin complex (DKC1 and NOP10) as well as components of the

RNAi machinery, including AGO1, 2, 4, 5, 9 and 10 (Fig1b). Immunoprecipitation of GFP-

tagged NHP2 followed by RNA sequencing revealed snoRNAs, snRNAs and several pri-

miRNAs that co-immunoprecipitated with NHP2 either directly or as part of a larger complex

(Supplementary Dataset S2; Supp Extended Data Fig. 1g). In order to determine whether pri-

miRNAs were pseudouridylated in plants, as they are in animals, we then performed Ψ-IP and

sequencing on mRNAs from *Arabidopsis* flower buds (collectively called the inflorescence).

Several pri-miRNA transcripts were enriched, and there was a significant (p<0.01) overlap with

those enriched by immunoprecipitation with GFP-tagged NHP2 (Extended Data Fig. 1h;

Supplementary Dataset S4).

Encouraged by these results, we tested three approaches to detect Ψ directly in small RNAs in

Arabidopsis (Methods; Extended Data Fig.2 a): (i) Ψ-IP followed by small RNA sequencing, (ii)

CMC treatment of small RNA libraries following 5′-adapter ligation, to block RT and deplete

small RNAs that contain Ψ, and (iii) reverse transcription of CMC-treated small RNAs in the

presence of Mn^{2+}, which causes deletions and mutations at Ψ residues instead of a reverse

transcription stop[28]. To determine the efficacy of these techniques, we examined coverage at

known pseudouridylation sites in rRNA and tRNA. Ψ at UNΨAR motifs is directed by PUS7

and was detected by CMC-depletion in WT but not in *pus7* mutants (Extended Data Fig. 2b, c).

Ψ at position 55 in tRNA depends on PUS10 (Extended Data Fig 2b). tRNA fragments

containing Ψ-55 were IP-enriched and CMC-depleted, contained mismatches in CMC/Mn^{2+}-

treated libraries, and were ablated in *pus10* mutants (Extended Data Fig 2b, d, e). Predicted sites

of rRNA pseudouridylation were also enriched by Ψ-IP, depleted by CMC treatment, and

mismatched in CMC/Mn^{2+} libraries (Extended Data Fig2f-l). Thus, all three methods robustly

detected Ψ in small RNA fragments derived from tRNA and rRNA.

We then applied our assays to detect Ψ in mature small RNAs from *Arabidopsis* inflorescence

and detected more than forty pseudouridylated mature miRNAs (Fig. 1d), of which four

(miR159b-3p, miR403-5p, miR447a-3p and miR833a-5p) were detected using all three assays

(Fig. 1e; Supplementary Datasets S5-7). Examination of *pus7* and *pus10* mutant datasets

(Extended Data Fig.2 c, d) revealed that these tRNA modification enzymes had no detectable

effect on Ψ-IP enrichment or CMC-depletion of miRNA. Instead, we expected that DYSKERIN

(DKC1) might be responsible, given the enrichment for pri-miRNA in pulldowns with the

Dyskerin co-factor NHP2 (Extended Data Fig 1b). As mutants of NHP2 and other members of

the DKC complex are lethal[29], we rescued *dkc1* knockout mutations with an embryo-specific

DKC1 cassette (Extended Data Fig. 3a), which restored about 10% of wild-type *DKC1*

expression in the leaves (Extended Data Fig. 3b) and allowed seeds to germinate into viable but

infertile plants (Extended Data Fig. 3c). Several (44) miRNAs were significantly enriched by Ψ-

IP in WT plants of which 12 lost Ψ enrichment in *dkc1-2* leaves and another 15 required *DKC1*

to a lesser extent (<75% Ψ enrichment). Overall, Ψ levels in miRNA were significantly

decreased in *dkc1-2* (P<0.05; Extended Data Fig. 3d). We conclude that just as some miRNAs

are pseudouridylated by PUS1 in the mouse (Extended Data Fig. 1h), some are pseudouridylated

by DKC1 in Arabidopsis, but in both cases multiple pseudouridine synthases must be responsible

for miRNA pseudouridylation, just as they are for mRNA pseudouridylation in plants, yeast and

mammals[30].

Ψ-detection by mismatch sequencing revealed that pseudouridylation in *Arabidopsis* miRNAs

(Fig. 1g) and in siRNAs (Extended Data Fig. 4 a, b) was prevalent at the 5′ nucleotide, which is

frequently uridine in miRNAs and contributes to AGO loading preference[31-33]. As pri-miRNAs

associate with NHP2 and contain Ψ (Extended Data Fig. 1b), we predicted that Ψ in mature

miRNAs was deposited prior to processing of pri-miRNAs into small RNAs by the endonuclease

DICER-LIKE1 (DCL1). cDNA reverse transcribed from pri-miR159b in the presence of CMC

and Mn^{2+} revealed extensive mismatches (pseudouridylation) at the 5′ dicing site (25%Ψ), which

was exacerbated in a *hyponastic leaves1 (hyl1)* mutant (75%Ψ) that accumulates pri-miRNAs

due to loss of the microprocessor component R2D2 (Extended Data Fig. 4c). Ψ was also detected

at 5′ and 3′ sites in mature miR159b, suggesting that pseudouridylated pri-miRNAs are processed

into mature miRNAs (Extended Data Fig. 4c). We confirmed the presence of Ψ in mature

miR159b on Northern blots, where CMCT-treatment induces a mobility shift (Fig. 1h).

Intriguingly, it has recently been shown that miR159b is inherited from sperm cells by the

fertilized endosperm, where it impacts seed development[4]. As miR159b is heavily

pseudouridylated (Fig.1h), this suggests that pseudouridylated small RNA may be inherited from

pollen.

We therefore applied Ψ-IP, CMC and CMC/Mn^{2+} treatments to small RNA isolated from pollen.

We found that known sites of pseudouridylation in 18S rRNA were robustly detected in rRNA

fragments in pollen by all three techniques (Extended Data Fig. 5a-d). In both CMC and Ψ-IP

datasets, many more miRNAs in pollen were pseudouridylated than in flower buds, including

miR159b (Fig. 2a; Supplementary Datasets S7-9). This effect was even more dramatic for

siRNAs, which were heavily pseudouridylated in pollen (Fig.2b). Pseudouridylated siRNA (Ψ-

siRNA) were associated with most annotated TE families, although there were notable

exceptions (Copia41, Copia 36, AtGP1 and AtGP2). Next, we tested whether Ψ-miRNA and Ψ-

siRNA were enriched in sperm cells or associated with AGO proteins. We purified sperm cells

from total pollen by FACS and immunoprecipitated AGO proteins from total pollen, before

making small RNA sequencing libraries (Fig.2c). To determine whether pseudouridylation of

small RNA correlates with their localization, we compared pseudouridylation of pollen miRNAs

to sperm cell localization, revealing some association (Fig. 2d)[34]. Pseudouridylation of siRNA,

on the other hand, was strongly associated with sperm cell localization ($R^2 = 0.88$). AGO 2, 5, 6,

7 and 9 are all expressed in mature pollen, where they are mostly restricted to sperm cells[35],

including AGO5 and AGO9 which are localized to the perinuclear space (Fig. 1e,f). We

performed small RNA sequencing after immunoprecipitation (RIP-seq) with anti-AGO9

antibodies (Methods) and compared the abundance of siRNA in each TE family by sequencing

following Ψ-IP, along with published RIP-seq data from AGO1 and AGO5[36]. In contrast to Ψ-

miRNA, which were relatively weakly associated (Fig.2 g-i), Ψ-siRNA were strongly associated

with sperm cell localization, and with small RNA in AGO5, AGO9 and AGO1 pulldowns (Fig.

2j-l).

Pseudouridylation has been implicated in cytoplasmic localization of mRNA and tRNA

suggesting a role for Ψ in small RNA transport[17,37]. Pol IV-dependent 21-22 nt easiRNAs in

pollen are also mobile, and move into sperm cells from the VN[7-9]. To further investigate a role

for Ψ in inter-cellular mobility of small RNAs, we probed the function of Ψ in sperm cell

localization. Since our *dkc1* mutant was completely sterile (Extended Data Fig. 3c), we could not

use this to investigate the function of Ψ in pollen. Additionally, other PUS enzymes likely have

considerable redundancy, as twenty exist in Arabidopsis[30,38]. Instead, we took advantage of the

fact that pseudouridylation of tRNAs by Pus1[+] contributes to nuclear export in yeast, and

genetically interacts with the tRNA export factor Loss of Suppression 1 (Los1[+]), which encodes

Exportin-t[37]. Exportin-t binds the upper minihelix of L-shaped tRNA, consisting of the acceptor-

stem and the TψC-loop, and terminating at Ψ55 and C56[39]. This minihelix includes the 22nt 3'

tRNA fragment with a 5' Ψ and 3'CCA, resembling a 21-22nt small RNA duplex[40].

Intriguingly, the *Arabidopsis* homolog of *Los1[+]*, which is known as *PAUSED/HEN5 (PSD)*, was

previously identified in genetic screens for mobile small RNA phenotypes, along with mutants in

ARGONAUTE7 (*zippy*) and *HASTY (hst)*. *HST* encodes Exportin-5 and has recently been shown

to be required for long-range transport of artificial miRNA in Arabidopsis[41-44]. We therefore

investigated whether *psd* mutants could be used as a proxy for the function of Ψ in siRNA, as

mutants are viable and fertile.

First, we demonstrated that the *psd* mutation in *Arabidopsis* is synthetically lethal with *pus7* in a

manner similar to *los1[-]* and *pus1[-]* from yeast. We found that *pus7-/-; psd+/-* double mutants

were semi-sterile (Extended Data Fig. 6a). In contrast, *pus7+/-; psd-/-* plants had fertile pollen,

but had 25% aborted seed compared with less than 5% in *psd* alone (Extended Data Fig. 6b, c).

Viable double homozygous seeds were not recovered from either genotype upon self-

fertilization. These results indicated that tRNA require *PUS7* for modification in diploid cells

before meiosis (*pus7-/-*), and subsequently require *PSD* for nuclear export in haploid pollen

(from *psd+/-* parents). Next, we performed Ψ-IP and CMC depletion on small RNA from WT

and *psd* mutant pollen as well as WT and mutant inflorescence (flower buds) and found robust

detection of pseudouridylation sites in rRNA (Extended Data Fig.5 a-b, e-f). Of those miRNAs

that were significantly pseudouridylated in the inflorescence, only miR403 lost Ψ in the mutant

(Extended Data Fig. 5g), while miRNAs overall were unaffected (Extended Data Fig. 5h).

Similarly, siRNAs from TEs were mostly unmodified in the inflorescence, and were unaffected

by *psd* (Extended Data Fig. 5i). In pollen, however, miRNAs were significantly depleted of Ψ in

psd relative to WT (Fig. 3a). easiRNA accumulation depends on both PolIV and miR845

(miR845a and miR845b), the most abundant pollen miRNAs, which target hundreds of LTR

retrotransposons at their reverse transcription primer binding site[10,36]. siRNAs from miR845

targets were heavily pseudouridylated, which depended on *psd* (Fig. 3a), as were Pol IV-

dependent 21-22 nt and 23-24 nt siRNAs (Fig. 3b).

In pollen, RdRM components are prominently expressed in the VN[45], but easiRNAs accumulate

in sperm cells[7-9]. To determine if Ψ affected siRNA transport from VN into sperm cells, we

isolated sperm cells from mutant *psd-13* (Exportin-t), mutant *hst-6* (Exportin-5), and wild-type

pollen by FACS and compared the abundance of siRNAs and miRNAs (Fig. 3c, d). Analysis

revealed that 21-22nt easiRNAs were reduced in *psd* and elevated in *hst* (Fig. 3c). In contrast,

miRNAs were sharply reduced in *hst* sperm cells as expected, but unaffected by *psd* (Fig.3d)

consistent with the dependence of miRNA but not siRNA accumulation on Exportin-5[44,46].

Strikingly, 21-22nt easiRNA that depend on PolIV, were strongly depleted from sperm cells in

psd but not in *hst* mutants (Fig.3e, Extended Data Fig. 7a-f), consistent with transport being

dependent on Ψ. While overall levels of miRNA in sperm cells were unaffected by *psd* (Fig. 3d),

individual miRNA including miR159 and miR403 significantly lost Ψ in pollen (Fig. 3f). A

likely explanation is that miRNA that lose Ψ in *psd* mutants can use the *HST* pathway instead to

accumulate in sperm cells. One prediction of this model is that Ψ-miRNA might be able to use

the PSD pathway in *hst* mutants. This prediction was confirmed, as miRNA without 5' terminal

Ψ were lost in sperm cells from *hst* mutants, while those with terminal Ψ were retained,

suggesting PSD was sufficient for their transport (Fig. 3g).

easiRNAs from pollen target imprinted genes in Arabidopsis[10,12]. Imprinted genes in plants are

expressed in the seed from either the maternal or the paternal genome and are known as

maternally expressed and paternally expressed genes (MEGs and PEGs), respectively.

DEMETER (*DME*) encodes a DNA glycosylase that is expressed in the endosperm and in the

pollen VN, where it removes DNA methylation from TEs upstream and downstream of

imprinted genes resulting in parent-of-origin expression[47]. We therefore examined Ψ in siRNAs

from DME targets and imprinted genes. We detected Ψ in 21-22 nt easiRNAs and 24 nt siRNAs

from DME targets (Fig.4a), and from TEs surrounding maternally- and paternally-expressed

imprinted genes (MEGs and PEGs) (Extended Data Fig. 7g, h). Pseudouridylation of siRNAs

from upstream and downstream of DME targets was lost in *psd-13* mutant pollen (Fig. 4a), and

there was a striking depletion of easiRNAs from the promoters of DME target genes in *psd*

sperm cells relative to WT (Fig.4b-c). Ψ loss was more significant in MEGs rather than PEGs

(p<0.05; Extended Data Fig. 7a) and also affected sperm cell localization of siRNAs from both

MEGs and PEGs, with 20-22 nt siRNAs depleted and 23-25 nt siRNAs enriched in *psd-13* sperm

cells compared to wild-type (p<0.001; Extended Data Fig. 7b).

Pol IV-dependent 21-22 nt easiRNAs and miR845 mediate the triploid block, a dose-dependent

epigenetic phenomenon whereby *Arabidopsis* seeds abort during development, if they inherit an

excess of paternal genomes from diploid instead of haploid pollen [10-12]. As 21-22 nt Pol IV-

dependent easiRNAs were lost from sperm cells in *psd-13* pollen, this suggested that *psd-13*

might rescue triploid seed abortion[10]. As expected, we observed ~90% seed abortion when WT

Columbia plants were crossed with *omission of second division 1 (osd1)* mutants, which have

diploid pollen (Fig. 4d-g). However, when *osd1*; *psd-13* diploid pollen was applied, this resulted

in only ~10% seed abortion, similar to when *osd1*; *nrpd1a* diploid pollen was applied, which is

deficient in the large subunit of Pol IV (NRPD1). Importantly, levels of miR845a and miR845b

were not affected in *psd-13* mutant pollen, and were not responsible for rescuing the triploid

block (Extended Data Fig. 7i)[10]. Exportin-t is therefore essential for the triploid block and for

pseudouridylation of easiRNAs.

Thus *PAUSED/EXPORTIN-T* is required for transport of pseudouridylated easiRNAs, and of

pseudouridylated miRNAs into sperm cells from the VN, while *HASTY/EXPORTIN-5* transports

unmodified miRNAs (Fig. 4h). Pseudouridylated easiRNAs lose Ψ in *psd* mutants, presumably

because they fail to bind AGO proteins if they are not exported to sperm cells (Fig. 2e-h; Fig. 4i)

and are degraded. Alternatively, nucleo-cytoplasmic transport might be required for

pseudouridylation. This novel function for *PAUSED/EXPORTIN-T* is pollen-specific, as

pseudouridylated miRNAs and easiRNAs are mostly restricted to the male germline (Extended

Data Fig. 5g-i). It is noteworthy that tRNA have been found to carry a strong signal for

intercellular transport in plants, so that mRNA-tRNA fusions are mobile[48]. It is tempting to

speculate that Ψ and *PSD* may be involved there as well. Cytoplasmic connections between VN

and sperm cells are thought to be involved in transport [7,9], consistent with a role for

nucleocytoplasmic exportin. AGO5 is found in these "male germ units"[34], and AGO5 is also

required at least in part for the triploid block[36]. AGO9 has a similar localization (Fig. 2e, f) and

has previously been associated with siRNA movement in pollen[49,50]. We therefore examined

AGO9 localization using an mCherry N-terminal fusion and found that AGO9 was first localized

in the VN at the bicellular stage, but then moved to the sperm cells by the tricellular stage (Fig.

4i). This localization is consistent with a role for AGO9 before, and after siRNA transport as

previously proposed[49,51].

TE-derived siRNAs in the plant germline have similar functions to piRNAs in mammalian

germline, and so we asked if high levels of Ψ were conserved in piRNAs. We used Ψ-IP as well

as CMC depletion to detect Ψ in small RNAs from 8-week *post-partum* mouse testis. As

expected, 3'-tRNA fragments (tRFs) were enriched by Ψ-IP (Extended Data Fig. 8a) and

depleted by CMC (Extended Data Fig. 8b). Importantly many piRNAs from TEs, including from

LINE and LTR retroelements, were strongly enriched for Ψ (Extended Data Fig. 8c, d). Up to

3% of piRNAs were differentially enriched for Ψ, though many more were immunoprecipitated

and hence potentially pseudouridylated (Extended Data Fig. 8d). No individual piRNA cluster

was significantly enriched or depleted, suggesting sequence- rather than locus- specificity

(Extended Data Fig. 8e, f). 3'-tRF that were depleted by CMC treatment were 22nt in size as

expected for 3'tRF containing Ψ-55 from mature tRNA (Extended Data Fig.8e) while piRNA

were mostly 30 nt in size as expected (Extended Data Fig.8f). The same small RNA sizes and

classes were detected by Ψ-IP (Extended Data Fig.8g, h). Thus, not only 3'tRFs, but also

piRNAs and miRNAs in mammals are pseudouridylated. It remains to be seen if they also

depend on Ψ for transport by Exportin-t. Intriguingly, 5'tRFs are also pseudouridylated, and

were reported to be inherited suggesting a similar mechanism in mammalian sperm[52,53]. Why are

germline small RNAs so heavily modified in both plants and mammals? An intriguing possibility

is that modifications of RNA in the germline may avoid viral surveillance systems after

fertilization, which could otherwise recognize inherited small RNA as "non-self"[54].

Materials and methods

Cell lines and tissues

NIH/3T3 (ATCC, CRL-1658) were cultured in DMEM (Invitrogen), supplemented with 10%

fetal bovine serum (FBS) and 1% penicillin/streptomycin/glutamine. Cells were obtained from

ATCC and tested negative for mycoplasma contamination. Testes were harvested from

C57Bl/6 mice at 8 weeks *post-partum* and flash frozen in liquid nitrogen.

Plant growth conditions

Arabidopsis plants were grown under long-day conditions at 22 °C. Seeds were always surface-

sterilized with sodium hypochlorite, sown on Murashige and Skoog medium and stratified for 3

d at 4 °C. Seedlings were transplanted to soil 2 weeks after germination and grown under long-

day conditions at 22 °C. The following mutants were used: pus7-1 (SALK_098773), pus10¬-1

(SALK_091360), nrpd1a-3 (SALK_128428). Homozygous mutants were identified using PCR

with primers specific to the predicted site of T-DNA integration (Supplementary Table S1). psd-

13 and hst-6 alleles were previously identified[41,55]. The osd1-1 mutant was kindly provided by

Raphael Mercier.

Yeast two hybrid screening.

The two-hybrid screen was performed by Hybrigenics (Paris, France) using an Arabidopsis

thaliana Col-0 library (ATFO) prepared from young flowers, using an in-house cDNA clone of

AGO3 as bait. 7 candidate proteins were tested by bimolecular fluorescence complementation,

leading to the identification of AT5G08180, which encodes the H/ACA ribonucleoprotein

complex subunit 2 (NHP2), as a high confidence interactor. This was subsequently confirmed by

Mass Spectrometry.

Bi-molecular fluorescence complementation

BiFC was performed using pBiFC-2in1 vectors[56]. AGO3 and NHP2 were fused in all

orientations (NN, CN, NC, CC; only NN and CC complemented), while other AGO proteins

were fused only in NN configuration. Tobacco leaves were infiltrated essentially as previously

described[57]. Briefly, agrobacterium cultures harboring pBiFC vectors were grown overnight

before subculturing in 50 mL of LB broth with antibiotics and 20 μM acetosyringone. Cultures

were pelleted by centrifugation, and resuspended in infiltration buffer (10 mM MES, 10 mM

MgCl2, 100 μM acetosyringone). Leaves of ~3 weeks old Nicotiana benthamiana plants were

infiltrated on the abaxial surface using a syringe without a needle. Three days post-infiltration

leaf discs were removed and imaged using a Zeiss LSM780 confocal microscope. Primers for

cloning AGO proteins and NHP2 are detailed in Supplementary Table S1.

Construction of viable *DYSKERIN* (*dkc1*) mutants in Arabidopsis

Dyskerin (*dkc1*) viable mutants were generated as follows. *AtNAP57 (DKC1)* genomic sequence

was amplified and cloned into the pENTR vector. The *ABI3* promoter sequence was amplified

and inserted using the NotI restriction site. *pABI3:AtNAP57* cassette was transferred to the

pGWB501 vector using the Gateway cloning system and used for Agrobacterium-mediated,

floral dip Arabidopsis transformation of *nap57*/+ (SALK_031065) heterozygous plants.

Homozygous transformant lines were obtained by three rounds of Hygromycin resistance

selection followed by verification of *nap57/nap57* genetic background by PCR.

Co-immunoprecipitation and mass-spectrometry analyses

For each co-IP, 0.6 g of tissue from GFP-NHP2 plants was used. The powder was extracted for

30 min in 1.5 ml of lysis buffer (50 mM Tris HCl pH 8.0, 50 mM NaCl, 1% Triton X-100 and 1×

cOmplete™ ethylenediaminetetraacetic acid (EDTA)-free protease inhibitor (Roche). After

removal of cell debris by centrifugation (twice 10 min, 16 000 g, 4°C) the cleared supernatants

were incubated for 30 min with anti-GFP antibodies coupled to magnetic microbeads (μMACS

GFP isolation Kits, Miltenyi). Beads were loaded on magnetized MACS separation columns

equilibrated with lysis buffer and washed four times with 300 μl washing buffer (50 mMTris

HCl pH 7.5, 0.1% Triton X-100). Samples were eluted in 100 μl of pre-warmed elution buffer

(Milteny). Control IPs were performed using Col-0 plants and GFP antibodies. Eluted proteins

were digested with sequencing-grade trypsin (Promega). Peptide mixtures were separated using

Nano-Ultra Performance Liquid Chromatography coupled to Orbitrap Velos mass spectrometer

(Thermo) at IBB PAS, Warsaw. Data were searched against the TAIR10 database. Peptides were

identified with the Mascot algorithm (Matrix Science, London, UK). The total number of

MS/MS fragmentation spectra was used to quantify each protein from three replicates. The

statistical analysis based on spectral counts was performed using a homemade R package that

calculates fold change and p values using the quasi-likelihood negative binomial generalized log-

linear model implemented in the edgeR package[58]. P-value was adjusted using the Benjamini–

Hochberg method from the stats R package. The size factor used to scale samples was calculated

according to the DESeq2 normalization method[59].

miRNA precursor pull-down with NHP2-GFP

Anti-GFP antibodies were used to immunoprecipitate NHP2-GFP from three independent

transgenic lines expressing GFP-NHP2 under native promoter and WT plants as a negative

control. RNA was converted into cDNA libraries using NEBNext Ultra II Directional RNA

Library Prep Kit for Illumina. Libraries were sequenced with Nextseq HO 100bp paired end.

Adapters and unpaired reads were removed using Trimmomatic and files were analyzed using

Salmon software. Obtained TPM values for each transcript were adjusted with 0.1 or 0.01 for

snRNA/snoRNA and pri-/pre-miRNA respectively to avoid dividing by zero in the case of non-

detected transcripts. IP values were normalized against input, and Fold Change was calculated

for GFP-NHP2 vs. negative control (GFP IP in WT plants). Reproducibly enriched RNAs in

NHP2 IP had FC $\geq$ 1.2 in at least two lines.

RNA extraction of mammalian cells

Cells were lysed in Qiazol (QIAGEN) and total RNA was purified using the miRNEasy mini kit

(QIAGEN) according to the manufacturer's instructions. Tissues were lysed in Qiazol

(QIAGEN) and total RNA was purified using Tissuelyser II (QIAGEN), as per the

manufacturer's instructions. Small RNAs (<200nt) size-fractionated using RNA Clean &

Concentrator 5 column kits (Zymo), as per the manufacturer's instructions. Testes of 8 weeks old

mice were grinded under liquid nitrogen and RNA was extracted using Trizol (Thermo Fisher

Scientific) according to the manufacturer's instructions but using 80% EtOH during the wash

step.

Dot-blot

For dot-blot analysis, input RNA or RNA immunoprecipitated with either anti-Ψ or isotypic non-

specific antibodies was spotted onto a nitrocellulose membrane and UV cross-linked at 254 nm

(120 mJ/cm2). The membranes were blocked in Denhart's solution (1% Ficoll, 1%

polyvinylpyrrolidone, 1% bovine serum albumin; Thermo Fisher) for 1 h at room temperature

and incubated with Ψ antibody for 1 h at room temperature. Signal was detected using HRP

conjugated secondary antibodies and ECL (GE Healthcare) and developed on a Chemidoc MP

machine (BioRad).

Pseudouridine Immunoprecipitation

Small RNAs were isolated from 0.01-0.1 mg total RNA, first by separating <200 nt RNAs with

the Zymo clean and concentrator kit, and subsequently gel extracting RNA 20-30 nt on a 15%

polyacrylamide gel. Small RNAs were eluted from the crushed gel slice overnight in 500 μL

RNA elution buffer (0.3M NaCl, 0.5 mM EDTA). RNA was resuspended in 50 μL water,

typically containing ~4-40 ng/μL RNA. Resuspended RNA was mixed with 200 μL IPP buffer

(150 mM NaCl, 0.1% NP-40, and 10 mM Tris-HCl [pH 7.4] + 10 U/mL RNAse inhibitor) and 5

μg of mouse monoclonal pseudouridine antibody (D347-3, MBL intl.). Samples were incubated

for 2 hours at 4C with rotation before adding 50 μL sheep anti-mouse IgG dynabeads washed in

IPP buffer (Invitrogen). The unbound fraction was retained, and the beads were washed 4 times

with IPP buffer. To prepare RNA from the unbound fraction, 400 μL binding buffer from the

Zymo clean and concentrator kit was added, and RNA was run through the column according to

manufacturer's instructions. To elute bound plant siRNA from the beads, 150 μL Trizol was

added and incubated for 2 minutes before placing tubes on a magnetic rack. Trizol was

transferred to a new tube and 30 μL chloroform was added and mixed thoroughly. Samples were

centrifuged for 10 minutes at maximum speed and the aqueous phase was pipetted into a new

tube with an equal volume of 100% ethanol. Mouse small RNAs were eluted using 6.7 mM Ψ

triphosphate in IPP buffer with RNase-OUT (Thermo Fisher Scientific) for 30 minutes at 37C.

Samples were then run through a Zymo Clean and Concentrator kit according to manufacturer's

instructions, and eluted in 8 μL. Libraries were prepared using NEBNext small RNA sequencing

kit (New England Biolabs) or NextFlex small RNA sequencing kit (Perkin Elmer) according to

manufacturer's instructions, with the same temperature modifications as the CMC-treated

samples. Libraries were pooled and run on NextSeq 500, HiSeq 4000 or MiSeq high-throughput

sequencing systems (Illumina).

Nucleoside analysis of RNA

RNA samples were incubated overnight with a 2x nuclease mix containing 62.5 units of

Benzonase (Sigma Aldrich), 5 units of Antarctic Phosphatase (NEB) and 10 mU/μL of

phosphodiesterase I (PDEI) from Crotalus adamanteus venom (Sigma Aldrich) made up in a 5X

digest buffer composed of 20 mM Tris-HCl (pH 8), 20 mM MgCl2 and 100 mM NaCl. Samples

were then further purified to remove protein by filtration at 14,000 rpm for 30 minutes through

Amicon® Ultra-0.5 spin columns (Sigma Aldrich) which have a 30 kDa molecular weight cutoff

(56). Samples were then analysed using a Q Exactive HF Orbitrap mass spectrometer (Thermo

Scientific). Briefly, 2ul of digested sample was loaded onto a 100Å, 1.8 μm, 2.1 mm X 100 mm

ACQUITY UPLC HSS T3 Column (Waters) in an aqueous buffer consisting of 0.1% MS-grade

formic acid in water: acetonitrile 98:2 at a constant flow of 0.3ml/min. A gradient of organic

solvent was used to facilitate separation and elution as follows: from 2 to 8 minutes the

percentage of organic solvent (0.1% formic acid in acetonitrile) increased from 2% to 10%.

From 8 to 13 minutes the gradient increased from 10% to 98% organic solvent, where it was held

for 1 minute before returning to 2% organic to re-equilibrate the column for the next sample.

Samples were read on a full scan positive mode from 100-600 m/z and data were analysed using

Xcalibur software (Thermo Scientific). All nucleosides were compared to exogenous standards

which allowed the interpolation of concentrations of sample nucleosides. Isobaric nucleosides

were differentiated on the basis of their retention times as compared with pure commercial

standards. As an example, Ψ has a retention time of 1.55 minutes in this protocol, whereas

uridine elutes at 3.05 minutes.

RT-qPCR

Cells were lysed in Qiazol (QIAGEN) and total RNA was purified using the miRNEasy mini kit

(QIAGEN) according to the manufacturer's instructions. For mRNA detection, 1 μg of purified

total RNA was reverse transcribed using the high-capacity cDNA reverse transcription kit

(Applied Biosystems). For specific miRNA quantification, we size-fractionated small RNAs (<

200nt, containing pre-miRNAs) using RNA Clean & Concentrator 5 column kits (Zymo), as per

the manufacturer's instructions. miRNAs were reverse transcribed with miScript II RT kit

(QIAGEN). Primers were designed to anneal on the 5p or 3p arm of the miRNA of interest

(Supplementary Table S1). miRNAs were quantified using Fast SybrGreen PCR mastermix

(Applied Biosystems) according to the manufacturer's instructions.

CMC-treatment and small RNA sequencing

For each pair of libraries 4 µg of Arabidopsis inflorescence RNA, prepared using Zymo

Quickzol RNA kit, was ligated to the 3′ adapter supplied in the NEB NextSeq kit in a 40 µL

reaction. After ligation samples were divided in two and subjected to CMC or mock treatment.

80 µL of 0.5M CMC in BEU buffer (7 M urea, 4 mM EDTA, 50 mM bicine pH 8) was added to

20 µL sample and incubated at 37C for 30 minutes. Mock treated samples were incubated with

80 µL of BEU buffer. After CMC/mock treatment 25 µL 3M sodium acetate, 3 µL glycogen and

1 mL 100% ethanol was added and samples were incubated for 1 hour at -80C. Samples were

centrifuged at 4ºC for 30 minutes and the resulting pellets were washed with 80% ethanol,

followed by another 1 hour at -80C. Samples were centrifuged at 4C for 30 minutes and the

ethanol removed; pellets were allowed to dry for 5-10 minutes before addition of 50 µL freshly

prepared 0.05 M carbonate buffer (pH 10.4) containing 4 mM EDTA. Samples were incubated at

37C for 3 hours and subsequently cleaned using Zymo RNA clean and concentrator kit, eluting

in a final volume of 25.5 µL. After cleaning, libraries were prepared according to instructions in

the NEBNext small RNA sequencing kit (New England Biolabs) or Nextflex Small RNA-Seq

Kit v3 (Perkin Elmer), beginning with annealing of the RT primer, with some modifications.

Primer annealing was performed at 65C for 5 minutes, followed by 37C for 15 minutes and 25ºC

for 15 minutes. cDNA synthesis was carried out at 42C. After PCR amplification libraries were

cleaned using Qiagen PCR cleanup kit and run on a 6% acrylamide gel. Bands of ~140 nt were

cut from the gel, crushed, and eluted for 2 hours in DNA elution buffer followed by ethanol

precipitation as outlined in the manufacturer's instructions. Libraries were pooled and run on

NextSeq or MiSeq high-throughput sequencing systems (Illumina) at the Cold Spring Harbor

Laboratory sequencing facility.

CMC/Mn2+ treatment and sequencing

For high-throughput siRNA mismatch analysis 2 µg total Arabidopsis RNA was treated as

described above for CMC-siRNA sequencing libraries with some modifications. Nextflex Small

RNA-Seq Kit v3 (Perkin Elmer) was used for all samples. The RT step was modified to utilize

SuperScript II (Thermo Fisher) and a custom 5X First-Strand Buffer containing $MnCl_2$ (250 mM

Tris-HCl, pH 8.3; 375 mM KCl; 30 mM $MnCl_2$). For pri-miRNA mismatch analysis, 10 µg of

total RNA was treated in the same way as for siRNA mismatch analysis. Obtained cDNA was

used for amplification with pri-miRNA specific primers. PCR products were gel-purified and

cloned into pGEM-T Easy (Promega) for Sanger sequencing. The same number of clones was

sequenced for corresponding CMC+ and CMC- samples.

Northern Blotting

50 µg total RNA was treated with 0.5 M CMC in BEU buffer (typically 25 µL RNA in 200 uL

buffer). Samples were cleaned and prepared in the same way as for small RNA sequencing,

except that RNA <200 nt was prepared using Zymo RNA Clean and Concentrator to enrich small

RNA in a very low volume. Samples were briefly denatured at 65C for two minutes (higher

temperatures should be avoided as these can remove the CMC adducts) before loading on a 15%

denaturing polyacrylamide TBE gel. CMC+/CMC- were treated identically, except for addition

of CMC. Untreated RNA was loaded directly on the gel after denaturation. Gels were run until

the bromophenol blue band reached the bottom. RNA was transferred to Hybond NX membrane

using a semi-dry electroblotter for 30 minutes. Crosslinking was performed with EDC[60].

Membranes were blocked using DIG easy-hyb buffer (Roche) followed by incubation with 10

pmol/mL 5′/3′ DIG labeled LNA probe (Exiqon) in easy-hyb buffer at 65C overnight. Probes

were detected using Roche DIG wash and block buffer set, anti-DIG antibody and CDP star

reagent according to manufacturer's instructions.

Immunoprecipitation of AGO9

Immunoprecipitation of AGO9 from Arabidopsis pollen was performed as previously described

(35). Pollen was collected from open Arabidopsis flowers as described above, prior to grinding

in liquid nitrogen and homogenizing in 5 ml g–1 of extraction buffer for 1 h at 4C (50mM Tris-

HCl pH 7.5, 150mM NaCl, 10% Glycerol, 0.2% NP-40, 5mM MgCl2, 5mM DTT, containing

one tablet per 10 ml of protease inhibitor cocktail, Roche). Cell debris was removed by

centrifugation at 12 Krpm at 4C for 20 min and the supernatant was recovered. Protein extract

was pre-cleared by incubation with 10 μl Protein A Dynabeads (Invitrogen) at 4C for 30 min.

Supernatant was recovered by centrifugation at 12 Krpm at 4C for 1min. Precleared extracts

were incubated with AGO9 antibody (1:100 dilution) and 30 μl of Protein A Dynabeads at 4C

overnight. The immunoprecipitate was pelleted and washed three times (15 min each in rotator

shaker) in extraction buffer. Small RNAs from the purified AGO9 complex were isolated using

Ambion columns. Small RNA was resolved on a 12.5% denaturing PAGE 7M-Urea gel, and

stained with SYBR-gold (Invitrogen). For cloning, gel slices within the range of 18–30 nt were

excised, and the RNAs were eluted and purified as described above. Small RNAs were

sequenced by Fasteris (Switzerland).

Flow cytometry and staining of Arabidopsis pollen

Flow cytometry and Alexander staining was performed as previously reported[34]. Pollen was

collected in 1.5mL eppendorf tubes by vortexing open flowers in pollen extraction buffer (PEB,

10 mM CaCl2, 2 mM MES, 1 mM KCl, 1% H3BO3, 10%) for 3 min 47, followed by filtration

through a 30um mesh (Partec/Sysmex) and centrifugation at 5,000g for 1 min. Pollen was

suspended in 50ul of PEB, immediately frozen in liquid nitrogen and stored at -80C until RNA

extraction. Open flowers were collected into a 50 ml falcon tube, frozen in liquid nitrogen, and

stored at -80C for storage. The pollen enriched fraction was isolated with the procedure

described in Borges et al. (2012) using Sperm Extraction Buffer (SEB). Intact pollen grains were

sorted with fluorescence-activated cell sorting (FACS) using a S3e Cell Sorter Bio-Rad. To the

obtained pollen sample in 1.5 ml tube, 100 μl of acid-washed glass beads (425–600 μm, Sigma)

was added and vortexed continuously at medium speed for 1 minute in order to break mature

pollen grains. The sample was transferred to a new tube and the volume of the sample was made

up to 1 ml with SEB. 1 μl of Sybr Green was added for staining. The sample was ready for

FACS for sperm cells sorting according to Borges, et al. [34]. RNA from the sperm cells fraction

was isolated with Trizol, treated with DNase I, and used for small RNA library preparation

(NextFlex Small RNA Kit, Perkin Elmer). Pollen staining was done following Alexander's

staining protocol[61]. Only viable pollen grains take up the stain.

Bioinformatic analysis of rRNA coverage

Adapters were trimmed from reads, converted to fasta and collapsed using fastx toolkit.

Trimmed reads between 15-30nt in length were mapped to the Arabidopsis genome (TAIR10)

using bowtie[62] allowing mismatches and multi-mapping. Coverage across the rRNA locus on

chromosome 2 (position 3706-9521) was calculated using bedtools [63]. Coverage within a sample

was normalized to the total amount of coverage across the locus, and the log2 fold-change

between paired samples (IP/Unbound, CMC/Mock) was calculated and averaged between

technical replicates. Sites of predicted pseudouridylation were taken from published datasets and

the average log2 fold change surrounding these sites was calculated[64,65].

Bioinformatic analysis of CMC/Mn2+-treated Arabidopsis samples

For adapter trimming and size selection of reads (18-42 nt) Cutadapt was used[66]. Terminal

deletions in CMC+ samples were detected using DeAnnIso[67] or miRMOD[68]. For internal

deletions detection reads were aligned to the Arabidopsis genome (TAIR10) using BWA[69] and

further processed using SAMtools[70] bam-readcount and AWK homemade scripts.

Bioinformatic analysis of Arabidopsis siRNA/miRNA

Adapters were trimmed from reads, converted to fasta and collapsed using fastx toolkit.

Trimmed reads between 20-25nt in length were mapped to the Arabidopsis genome (TAIR10)

using bowtie allowing mismatches and discarding unmapped reads. Individual sequences with an

average of >10 reads per sample were selected and analysed using DESeq2[59], treating

CMC+/Mock or IP/Unbound fractions as pairs. Mature miRNA sequences from miRBase (v21)

were extracted from these datasets. Sequences mapping to tRNA, snoRNA, snRNA or rRNA

were removed for comparison with AGO-bound sequences from published datasets. UNUAR-

containing and 3′CCA tRFs were identified by extracting reads mapping to tRNAs and selecting

those containing the UNUAR motif or 3′CCA; these sequences were extracted from the DESeq2

output for plotting. Significance of the overlap between CMC, IP and CMC/Mn2+ results was

calculated by simulating 1x10e6 outcomes and calculating the probability of the observed

overlap with a custom script.

Bioinformatic analysis of mouse miRNA

The mouse small RNA Sequencing data from NIH/3T3 cells were analysed with the SeqImp

pipeline configured with the mm10 assembly of the Mus musculus genome, the transcriptome

annotation from Ensembl (v89) and the miRbase annotation of microRNAs (v21). The pipeline

was executed with the default mouse miRNA configuration and default parameters, specifying

the sequence of the 3' adapter used for library preparation (AGATCGGAAGAGCACACGTC).

The miRNA quantifications obtained with SeqImp were further analyzed in R/Bioconductor with

the DESeq2 package[59]. Briefly, counts were library-size normalized using the median ratio

method as implemented in the estimateSizeFactors function. Variance was estimated with the

estimateDispersions function and statistical testing for enrichment performed with the Wald test

as implemented in the nbinomWaldTest function. P-values were corrected for multiple

hypothesis testing with the Benjamini-Hochberg procedure.

Analysis of TE-derived siRNA

Collapsed reads 18-30 nt long were mapped to the TAIR10 genome allowing multimapping, but

no mismatches using Bowtie. Quantification of siRNA mapping to each TE was performed as

described previously[10]. Reads mapping to TE families were aggregated and the average log2

fold-change between paired samples was calculated. The mean of all technical replicates

(generated by separate CMC/mock or IP/unbound library preparations on the same RNA) for any

given biological replicate was taken prior to calculating the mean of all biological replicates for

each tissue (inflorescence/pollen) and treatment (CMC or IP) combination.

Metaplots

Uncollapsed reads 20-25nt in length were aligned to the Arabidopsis (TAIR10) genome using

Bowtie allowing mismatches but only a single mapping event per read. Mapping files from the

same treatment (ie. CMC+, CMC-, IP, Unbound) were merged after normalization (as above) to

enable the production of a single representative plot. Coverage was calculated using bedtools

(60). Metaplots were produced using the Versatile Aggregate Profiler[71] using a window size of

25, and a smoothing window of 6 for TE analysis, and a window size of 50 and smoothing

window of 12 for DME target analysis. For comparison between psd-13 and wild-type, metaplots

of three biological replicates were created and the mean enrichment was plotted.

Bioinformatic analysis of mouse testis small RNA

Reads were clipped for Illumina Truseq (AGATCGGAAGAGCACACGTCTGAAC) or Perkin

Elmer Nextflex Small RNA adapter sequences (N(4)TGGAATTCTCGGGTGCCAAGG) using

Cutadapt. Reads were required to be 14-44nt and have a quality score of at least Q20 over 90%

of their length. tRNA fragment sequences (tRFs) were analyzed as described[72] using all tRNA

sequences included in the current UCSC repeat annotation

(ftp://hgdownload.soe.ucsc.edu/goldenPath/mm10/database/rmskOutCurrent.txt.gz). Reads that

were not tRFs were mapped to the UCSC mm10 genome sequence using Bowtie 2, filtered for 0-

3 mismatches (to accommodate mismatches due to RNA modifications) using SAMtools and

BAMtools, and intersected with the UCSC repeat annotation using BEDtools. Repeat and non-

repeat reads were further intersected with mm10 mouse piRNA clusters

(https://www.smallrnagroup.uni-mainz.de/piCdb/) to result in piRNA reads that overlap (i)

transposon sequences, (ii) structural RNAs or (iii) neither. Raw counts of individual piRNA

sequences were analyzed for pseudouridine enrichment (enriched in IP, depleted in CMC-treated

libraries) using DESeq2[59]. Similarly, raw counts per piRNA cluster were analyzed using

DESeq2, but each cluster was required to have at least an average of 100 raw reads (per IP or per

CMC-untreated sample). Read counts that were used as input for DESeq2 analysis were from

three replicates of each condition (input, IP; CMC-untreated, CMC-treated). Log2 fold changes

were plotted against -log10(adjusted p-value) using R.

Acknowledgments: The authors would like to thank J.S. Parent and M. Donoghue for

discussions and observations, Joe Simorowski, Uma Ramu and Alex Oruci for technical

assistance, and Tim Mulligan for plant care.

Funding: This work was supported by the Howard Hughes Medical Institute, and by grants from

the National Institutes of Health (GM067014 and GM76396), the National Science Foundation

Plant Genome Research Program, and the Robertson Research Foundation (to R.A.M.). Work in

the Kouzarides laboratory is supported by a grant from Cancer Research UK (grant reference

RG96894), in addition to benefiting from core support from the Welcome Trust (WT203144)

and Cancer Research UK (grant reference C6946/A24843). V.M. was funded by a Kay Kendall

Leukemia Fund project grant (grant reference RG88664) and Cancer Research UK (grant

reference RG96894). J.D was funded by a grant from the Polish National Science Centre

(2020/39/D/NZ1/01918). The authors acknowledge assistance from the Cold Spring Harbor

Laboratory Shared Resources, which are funded in part by the Cancer Center (Support Grant

5PP30CA045508).

Author contributions: R.P.H., J.D., C.S.A., V.M., T.K. and R.A.M. conceived and designed

experiments. R.P.H., V.M., J.D., C.S.A., A.L, F.V., and A.H. performed the experiments. R.P.H.,

J.D., T.L. and A.J.S. performed bioinformatic analysis. F.B. provided code and advice for

transposon analysis. R.A.M. and R.P.H. wrote the manuscript with input from V.M., J.D.,

C.S.A., A.J.S., T.L. and T.K.

Competing interests: T.K. is a co-founder of Abcam Plc and Storm Therapeutics Ltd,

Cambridge, UK. A.H. is an employee of Storm Therapeutics Ltd, Cambridge, UK.

Data and materials availability: Data has been deposited in GEO under accession numbers

GSE229986; GSE230359;GSE230228; GSE222751.

Supplementary Materials

Supplemental Table S1

Supplemental Data S1 to S9

References

1   Kohler, C., Wolff, P. & Spillane, C. Epigenetic mechanisms underlying genomic
imprinting in plants. *Annu Rev Plant Biol* **63**, 331-352, doi:10.1146/annurev-arplant-
042811-105514 (2012).

2   Skvortsova, K., Iovino, N. & Bogdanovic, O. Functions and mechanisms of epigenetic
inheritance in animals. *Nat Rev Mol Cell Biol* **19**, 774-790, doi:10.1038/s41580-018-
0074-2 (2018).

3   Zhang, Y. *et al.* Dnmt2 mediates intergenerational transmission of paternally acquired
metabolic disorders through sperm small non-coding RNAs. *Nat Cell Biol* **20**, 535-540,
doi:10.1038/s41556-018-0087-2 (2018).

4   Zhao, Y. *et al.* Clearance of maternal barriers by paternal miR159 to initiate endosperm
nuclear division in Arabidopsis. *Nat Commun* **9**, 5011, doi:10.1038/s41467-018-07429-x
(2018).

5   Shukla, A. *et al.* poly(UG)-tailed RNAs in genome protection and epigenetic inheritance.
Nature **582**, 283-288, doi:10.1038/s41586-020-2323-8 (2020).

6   Girard, A., Sachidanandam, R., Hannon, G. J. & Carmell, M. A. A germline-specific
class of small RNAs binds mammalian Piwi proteins. *Nature* **442**, 199-202,
doi:10.1038/nature04917 (2006).

7   Slotkin, R. K. *et al.* Epigenetic reprogramming and small RNA silencing of transposable
elements in pollen. *Cell* **136**, 461-472, doi:10.1016/j.cell.2008.12.038 (2009).

8   Long, J. *et al.* Nurse cell–derived small RNAs define paternal epigenetic inheritance in
Arabidopsis. **373**, eabh0556 (2021).

9   Martinez, G., Panda, K., Köhler, C. & Slotkin, R. K. Silencing in sperm cells is directed
by RNA movement from the surrounding nurse cell. *Nat Plants* **2**, 1-8 (2016).

10   Borges, F. *et al.* Transposon-derived small RNAs triggered by miR845 mediate genome
dosage response in Arabidopsis. *Nat Genet* **50**, 186-192, doi:10.1038/s41588-017-0032-5
(2018).

11   Panda, K., McCue, A. D. & Slotkin, R. K. Arabidopsis RNA Polymerase IV generates
21-22 nucleotide small RNAs that can participate in RNA-directed DNA methylation and

may regulate genes. *Philos Trans R Soc Lond B Biol Sci* **375**, 20190417,
doi:10.1098/rstb.2019.0417 (2020).

12 Martinez, G. *et al.* Paternal easiRNAs regulate parental genome dosage in Arabidopsis.
Nat Genet **50**, 193-198, doi:10.1038/s41588-017-0033-4 (2018).

13 Zhai, J. *et al.* Spatiotemporally dynamic, cell-type-dependent premeiotic and meiotic
phasiRNAs in maize anthers. *Proc Natl Acad Sci U S A* **112**, 3146-3151,
doi:10.1073/pnas.1418918112 (2015).

14 Gainetdinov, I. *et al.* Terminal modification, sequence, length, and PIWI-protein identity
determine piRNA stability. *Mol Cell* **81**, 4826-4842 e4828,
doi:10.1016/j.molcel.2021.09.012 (2021).

15 Carlile, T. M. *et al.* Pseudouridine profiling reveals regulated mRNA pseudouridylation
in yeast and human cells. *Nature* **515**, 143-146, doi:10.1038/nature13802 (2014).

16 Li, X. *et al.* Chemical pulldown reveals dynamic pseudouridylation of the mammalian
transcriptome. *Nat Chem Biol* **11**, 592-597, doi:10.1038/nchembio.1836 (2015).

17 Schwartz, S. *et al.* Transcriptome-wide mapping reveals widespread dynamic-regulated
pseudouridylation of ncRNA and mRNA. *Cell* **159**, 148-162,
doi:10.1016/j.cell.2014.08.028 (2014).

18 Lovejoy, A. F., Riordan, D. P. & Brown, P. O. Transcriptome-wide mapping of
pseudouridines: pseudouridine synthases modify specific mRNAs in S. cerevisiae. *PLoS*
One **9**, e110799, doi:10.1371/journal.pone.0110799 (2014).

19 Martinez Campos, C. *et al.* Mapping of pseudouridine residues on cellular and viral
transcripts using a novel antibody-based technique. *RNA* **27**, 1400-1411,
doi:10.1261/rna.078940.121 (2021).

20 Dai, Q. *et al.* Quantitative sequencing using BID-seq uncovers abundant pseudouridines
in mammalian mRNA at base resolution. *Nat Biotechnol* **41**, 344-354,
doi:10.1038/s41587-022-01505-w (2023).

21 Kurimoto, R. *et al.* The tRNA pseudouridine synthase TruB1 regulates the maturation of
let-7 miRNA. **39**, e104708 (2020).

22 Viswanathan, S. R., Daley, G. Q. & Gregory, R. I. Selective blockade of microRNA
processing by Lin28. *Science* **320**, 97-100, doi:10.1126/science.1154040 (2008).

23 Chen, C. *et al.* Meiosis-specific gene discovery in plants: RNA-Seq applied to isolated
Arabidopsis male meiocytes. *BMC plant biology* **10**, 1-14 (2010).

24 Gurha, P., Joardar, A., Chaurasia, P. & Gupta, R. Differential roles of archaeal box
H/ACA proteins in guide RNA-dependent and independent pseudouridine formation.
RNA biology **4**, 101-109 (2007).

25 Lafontaine, D. L., Bousquet-Antonelli, C., Henry, Y., Caizergues-Ferrer, M. & Tollervey,
D. The box H+ ACA snoRNAs carry Cbf5p, the putative rRNA pseudouridine synthase.
Genes and development **12**, 527-537 (1998).

26 Watkins, N. J. *et al.* Cbf5p, a potential pseudouridine synthase, and Nhp2p, a putative
RNA-binding protein, are present together with Gar1p in all H BOX/ACA-motif
snoRNPs and constitute a common bipartite structure. *RNA* **4**, 1549-1568 (1998).

27 Li, C. F. *et al.* An ARGONAUTE4-containing nuclear processing center colocalized with
Cajal bodies in Arabidopsis thaliana. *Cell* **126**, 93-106, doi:10.1016/j.cell.2006.05.032
(2006).

28 Zhou, K. I. *et al.* Pseudouridines have context-dependent mutation and stop rates in high-
throughput sequencing. *RNA Biol* **15**, 892-900, doi:10.1080/15476286.2018.1462654
(2018).

29 Kannan, K., Nelson, A. D. & Shippen, D. E. Dyskerin is a component of the Arabidopsis
telomerase RNP required for telomere maintenance. *Mol Cell Biol* **28**, 2332-2341,
doi:10.1128/MCB.01490-07 (2008).

30 Sun, L. *et al.* Transcriptome-wide analysis of pseudouridylation of mRNA and non-
coding RNAs in Arabidopsis. *J Exp Bot* **70**, 5089-5600, doi:10.1093/jxb/erz273 (2019).

31 Havecker, E. R. *et al.* The Arabidopsis RNA-directed DNA methylation argonautes
functionally diverge based on their expression and interaction with target loci. *Plant Cell*
22, 321-334, doi:10.1105/tpc.109.072199 (2010).

32 Mi, S. *et al.* Sorting of small RNAs into Arabidopsis argonaute complexes is directed by
the 5' terminal nucleotide. *Cell* **133**, 116-127, doi:10.1016/j.cell.2008.02.034 (2008).

33 Montgomery, T. A. *et al.* Specificity of ARGONAUTE7-miR390 interaction and dual
functionality in TAS3 trans-acting siRNA formation. *Cell* **133**, 128-141,
doi:10.1016/j.cell.2008.02.033 (2008).

34 Borges, F. *et al.* FACS-based purification of Arabidopsis microspores, sperm cells and
vegetative nuclei. *Plant Methods* **8**, 1-9 (2012).

35 Jullien, P. E., Schroder, J. A., Bonnet, D. M. V., Pumplin, N. & Voinnet, O. Asymmetric
expression of Argonautes in reproductive tissues. *Plant Physiol* **188**, 38-43,
doi:10.1093/plphys/kiab474 (2022).

36 Oliver, C. *et al.* The miRNome function transitions from regulating developmental genes
to transposable elements during pollen maturation. *Plant Cell* **34**, 784-801,
doi:10.1093/plcell/koab280 (2022).

37 Grosshans, H., Lecointe, F., Grosjean, H., Hurt, E. & Simos, G. Pus1p-dependent tRNA
pseudouridinylation becomes essential when tRNA biogenesis is compromised in yeast. *J*
Biol Chem **276**, 46333-46339, doi:10.1074/jbc.M107141200 (2001).

38 Xie, Y. *et al.* Genome-Wide Identification and Expression Analysis of Pseudouridine
Synthase Family in Arabidopsis and Maize. *Int J Mol Sci* **23**, doi:10.3390/ijms23052680
(2022).

39 Cook, A. G., Fukuhara, N., Jinek, M. & Conti, E. Structures of the tRNA export factor in
the nuclear and cytosolic states. *Nature* **461**, 60-65, doi:10.1038/nature08394 (2009).

40 Schorn, A. J. & Martienssen, R. Tie-Break: Host and Retrotransposons Play tRNA.
Trends Cell Biol **28**, 793-806, doi:10.1016/j.tcb.2018.05.006 (2018).

41 Hunter, C. A., Aukerman, M. J., Sun, H., Fokina, M. & Poethig, R. S. PAUSED encodes
the Arabidopsis exportin-t ortholog. *Plant Physiol* **132**, 2135-2143 (2003).

42 Li, J. & Chen, X. PAUSED, a putative exportin-t, acts pleiotropically in Arabidopsis
development but is dispensable for viability. *Plant Physiol* **132**, 1913-1924 (2003).

43 Garcia, D., Collier, S. A., Byrne, M. E. & Martienssen, R. A. Specification of leaf
polarity in Arabidopsis via the trans-acting siRNA pathway. *Curr Biol* **16**, 933-938,
doi:10.1016/j.cub.2006.03.064 (2006).

44 Brioudes, F. *et al.* HASTY, the Arabidopsis EXPORTIN5 ortholog, regulates cell-to-cell
and vascular microRNA movement. *The EMBO Journal*, e107455 (2021).

45 Calarco, J. P. *et al.* Reprogramming of DNA methylation in pollen guides epigenetic
inheritance via small RNA. *Cell* **151**, 194-205, doi:10.1016/j.cell.2012.09.001 (2012).

46 Park, M. Y., Wu, G., Gonzalez-Sulser, A., Vaucheret, H. & Poethig, R. S. Nuclear
processing and export of microRNAs in Arabidopsis. *Proc Natl Acad Sci U S A* **102**,
3691-3696, doi:10.1073/pnas.0405570102 (2005).

47 Khouider, S. *et al.* Male fertility in Arabidopsis requires active DNA demethylation of
genes that control pollen tube function. *Nat Commun* **12**, 410, doi:10.1038/s41467-020-
20606-1 (2021).

48 Zhang, W. *et al.* tRNA-Related Sequences Trigger Systemic mRNA Transport in Plants.
Plant Cell **28**, 1237-1249, doi:10.1105/tpc.15.01056 (2016).

49 Wu, W. *et al.* Heterochromatic silencing is reinforced by ARID1-mediated small RNA
movement in Arabidopsis pollen. *New Phytol* **229**, 3269-3280, doi:10.1111/nph.16871
(2021).

50 Olmedo-Monfil, V. *et al.* Control of female gamete formation by a small RNA pathway
in Arabidopsis. *Nature* **464**, 628-632, doi:10.1038/nature08828 (2010).

51 Yelina, N. E. *et al.* Putative Arabidopsis THO/TREX mRNA export complex is involved
in transgene and endogenous siRNA biosynthesis. *Proc Natl Acad Sci U S A* **107**, 13948-
13953, doi:10.1073/pnas.0911341107 (2010).

52 Sharma, U. *et al.* Biogenesis and function of tRNA fragments during sperm maturation
and fertilization in mammals. *Science* **351**, 391-396, doi:10.1126/science.aad6780 (2016).

53 Chen, Q. *et al.* Sperm tsRNAs contribute to intergenerational inheritance of an acquired
metabolic disorder. *Science* **351**, 397-400, doi:10.1126/science.aad7977 (2016).

54 Lin, T. Y., Mehta, R. & Glatt, S. Pseudouridines in RNAs: switching atoms means
shifting paradigms. *FEBS Lett* **595**, 2310-2322, doi:10.1002/1873-3468.14188 (2021).

55 Bollman, K. M. *et al.* HASTY, the Arabidopsis ortholog of exportin 5/MSN5, regulates
phase change and morphogenesis. *Development* **130**, 1493-1504, doi:10.1242/dev.00362
(2003).

56 Grefen, C. & Blatt, M. R. A 2in1 cloning system enables ratiometric bimolecular
fluorescence complementation (rBiFC). *Biotechniques* **53**, 311-314,
doi:10.2144/000113941 (2012).

57 Li, X. Infiltration of Nicotiana benthamiana protocol for transient expression via
Agrobacterium. *Bio-protocol*, e95-e95 (2011).

58 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140, doi:10.1093/bioinformatics/btp616 (2010).

59 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8
(2014).

60 Pall, G. S. & Hamilton, A. J. Improved northern blot method for enhanced detection of
small RNA. *Nat Protoc* **3**, 1077-1084, doi:10.1038/nprot.2008.67 (2008).

61 Hedhly, A., Vogler, H., Eichenberger, C. & Grossniklaus, U. Whole-mount Clearing and
Staining of Arabidopsis Flower Organs and Siliques. *J Vis Exp*, doi:10.3791/56441
(2018).

62 Langmead, B. Aligning short sequencing reads with Bowtie. *Curr Protoc Bioinformatics*
Chapter 11, Unit 11 17, doi:10.1002/0471250953.bi1107s32 (2010).

63 Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033
(2010).

64 Marker, C. *et al.* Experimental RNomics: identification of 140 candidates for small non-
messenger RNAs in the plant Arabidopsis thaliana. *Current Biology* **12** (2002).

65 Piekna-Przybylska, D., Decatur, W. A. & Fournier, M. J. The 3D rRNA modification
maps database: with interactive tools for ribosome analysis. *Nucleic acids research* **36**,
D178-D183 (2007).

66 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet journal **17**, 3 doi:10.14806/ej.17.1.200 (2011).

67 Zhang, Y. *et al.* DeAnnIso: a tool for online detection and annotation of isomiRs from
small RNA sequencing data. *Nucleic Acids Res* **44**, W166-175, doi:10.1093/nar/gkw427
(2016).

68 Kaushik, A., Saraf, S., Mukherjee, S. K. & Gupta, D. miRMOD: a tool for identification
and analysis of 5' and 3' miRNA modifications in Next Generation Sequencing small
RNA data. *PeerJ* **3**, e1332, doi:10.7717/peerj.1332 (2015).

69 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

70 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079, doi:10.1093/bioinformatics/btp352 (2009).

71 Coulombe, C. *et al.* VAP: a versatile aggregate profiler for efficient genome-wide data
representation and discovery. *Nucleic Acids Res* **42**, W485-493, doi:10.1093/nar/gku302
(2014).

72   Schorn, A. J., Gutbrod, M. J., LeBlanc, C. & Martienssen, R. LTR-Retrotransposon
Control by tRNA-Derived Small RNAs. *Cell* **170**, 61-71 e11,
doi:10.1016/j.cell.2017.06.013 (2017).
823

Figure 1

Figure 1. microRNAs are pseudouridylated in plants and mammals. (**a**) Bimolecular

fluorescence complementation confirming interaction of AGO3 with the Dyskerin co-factor NHP2

in Arabidopsis (Methods). Split YFP was fused to the C- and N-, or N- and N-terminals of AGO3

and NHP2. Reconstituted YFP signifies interaction. p35S::RFP acts as a transformation control.

(**b**) Volcano plot showing proteins co-purified with GFP-NHP2 as compared to IPs performed in

wild-type plants. H/ACA ribonucleoprotein complex subunits and AGO family proteins are

depicted in orange (p-value adjusted by Benjamini-Hochberg correction). (**c**) Structure of uridine

and pseudouridine (Ψ), enzymes involved in catalysis, and methods of detection with specific

antibodies and chemical modification with CMCT (Methods; Supplemental Fig. 2). (**d**) Volcano

plot of miRNAs enriched by Ψ-IP compared to unbound fractions from *Arabidopsis* flower buds

(DESeq2). Blue, $p<0.01$; orange, both Ψ-IP-enriched and depleted from libraries following CMC-

treatment ($p<0.01$); dark red, not significant. (**e**) Venn diagram showing overlap of miRNAs

detected in flower buds by each technique. Predicted Ψ sites in miRNAs detected by all 3

techniques. (**f**) miRNAs enriched by Ψ-IP in WT and *dkc1* mutants in Arabidopsis flower buds (**g**)

Rate of deletion in CMC/Mn^{2+} treated libraries at each site of miRNAs aligned at the 5′ (left panel)

and 3′ (right panel) ends. (**h**) Northern blots of synthetic miR159 RNA pseudouridylated at position

10 and 21 (upper) and flower bud small RNA (lower) treated (+) or mock treated (-) with CMC,

and probed with miR159b to reveal mobility shifts.

Figure 2

Fig. 2 Pollen small interfering RNAs are highly pseudouridylated and loaded into

ARGONAUTE proteins in sperm cells. (**a**) Heatmaps of individual miRNA Ψ-enrichment in

flower buds and pollen by Ψ-IP (Log$_2$ IP/Unbound) and by CMC-mediated depletion from small

RNA libraries (Log$_2$ CMC+/CMC-) assessed by DESeq2. (**b**) Heatmaps of siRNA Ψ-enrichment

from individual TE families in flower buds and pollen by Ψ-IP and CMC-depletion as in (a). Small

RNAs from most TE families were enriched for Ψ in pollen, except for those highlighted. (**c**)

Small RNAs were sequenced from sperm cells isolated from pollen grains by FACS, and by RNA

immunoprecipitation (RIP-seq) with antibodies to ARGONAUTE (AGO) proteins. (**d**)

Association of pseudouridylation determined by Ψ-IP in pollen with sperm-cell abundance by

FACS for miRNAs and siRNAs from TE families. (**e**) GFP::AGO5 fusion protein expressed from

AGO5 promoter (*pAGO5*) in mature pollen is found in the perinuclear space around sperm cells

(SC). DAPI was used to stain the vegetative nucleus (VN). Reproduced, with permission, from

(*12*) (**f**) AGO9::GFP fusion protein expressed from the *AGO9* promoter (*pAGO9*) in mature pollen.

(**g-i**) Association of pseudouridylation determined by Ψ-IP with AGO1 (**g, j**), AGO5 (**h, k**), and

AGO9 (**i, l**) binding measured by RIPseq for miRNAs (**g-i**) and siRNAs from TE families (**j-l**).

Pearson R^2 correlations are indicated.

Figure 3

Fig. 3 Transport of pseudouridylated siRNAs into sperm cells depends on *PAUSED*

(Exportin-t). (**a**) Ψ-enrichment of miRNAs and some of their targets were determined by Ψ-IP

and small RNA sequencing in *psd-13* mutant pollen relative to wild-type. Ψ-enrichment of the

targets of miR845 were strongly dependent on *PSD* (paired t-test, miRNAs; one-way ANOVA,

miR845 targets) (**b**) Ψ-enrichment of Pol IV-dependent siRNA from "PolIV TEs" and other TEs

(non-PolIV) were determined by Ψ-IP and small RNA sequencing in *psd-13* mutant pollen relative

to wild-type (p<0.01, one-way ANOVA based on ranks). siRNA from PolIV TEs was strongly

dependent on *PSD*. (**c**) siRNA abundance was determined by sequencing small RNA from FACS

sorted sperm cells, and was reduced in *psd*, but increased in *hst* mutants relative to WT. Significant

differences between 20-22nt and 23-25nt siRNAs were observed (one-way ANOVA). (**d**) miRNA

abundance from FACS sorted sperm cells was reduced in *hst*, but not in *psd* mutant sperm cells

relative to wild type (one-way ANOVA). (**e**) Metaplots of 20-22 nt (red) and 23-25 nt (black)

siRNA abundance across "PolIV-TEs" from wild-type, *psd-13* and *hst-6* sperm cells sorted by

FACS. 250 bp up- and down-stream regions as well as 500 bp from each end of the TE are shown.

(**f**) Histogram of miRNA Ψ-enrichment in wild-type and *psd-13* pollen for those miRNA

dependent on *PSD* for pseudouridylation (student's t-test p<0.05; n = 3 biological replicates). (**g**)

Abundance of miRNA with and without terminal Ψ in *psd* and *hst* mutants relative to wildtype in

sperm cells sorted by FACS. miRNA without terminal Ψ were lost in *hst*, but not in *psd* sperm

cells (p<0.001). Asterisks indicate p<0.05 (*), 0.01 (**), 0.001 (***) and 0.0001(****).

Figure 4

Fig. 4. Pseudouridine and Exportin-t mediate epigenetic inheritance and the triploid block.
(**a**) Ψ-enrichment (Log$_2$ Ψ-IP/unbound) of 20-22 nt siRNAs and 23-25 nt siRNAs surrounding imprinted genes targeted by *DEMETER (DME),* in wild-type (blue) and *psd-13* (orange) pollen Asterisks indicate student's t-test; $p<0.05$ (*), 0.01 (**), 0.001 (***) and 0.0001(****). (**b, c**) sperm cell abundance of 20-22 nt siRNAs (red) and 23-25 nt siRNAs (black) surrounding imprinted genes targeted by *DEMETER (DME),* in wild-type (**b**) and *psd-13* (**c**) pollen. (**d, e, f**) Images of representative seeds from crosses between wild-type Col-0 female and (**d**) *osd1* diploid pollen, (**e**) double mutant *osd1*; *psd-13* diploid pollen or (**f**) double mutant *osd1; nrpd1a-1* diploid pollen. *NRPD1*a encodes the large subunit of Pol IV which is required for the triploid block (**g**) Frequency of aborted seeds from the same crosses. (**h**) Model for transport of small RNA during pollen development. As the pollen matures 21/22 nt easiRNAs and 24nt siRNAs are produced by Pol IV in the vegetative nucleus and pseudouridylated along with some miRNAs. pseudouridylated miRNAs and easiRNAs are exported via PSD (Exportin-t) into sperm cells, where they are loaded onto AGO1, AGO5 and AGO9, while unmodified miRNAs are transported via HST (Exportin-5) and loaded onto AGO5 and AGO1. Pseudouridylated sperm cell easiRNAs mediate dose-dependent lethality (the triploid block) by targeting maternally expressed imprinted genes in the seed. (**i**) mCherry::AGO9 fusion protein was expressed from the *AGO9* promoter (*pAGO9*) in developing pollen. DAPI was used to stain nucleic acids (blue). AGO9 is present in the VN at the binuclear stage, but is enriched in the sperm cells in mature pollen.

897   **Extended Data.**

Extended Data Figure 1

Extended Data Figure 1. miRNA from plants and mammals are pseudouridylated.

(**a**) Immunoprecipitation with anti-Ψ antibody enriches for Ψ-containing RNAs made by in vitro

transcription (IVT) as determined by mass spectrometry (n = 3 biological replicates, error bars ±

SD). (**b**) Immuno-dot blot of 100ng total input RNA or 100ng RNA immunoprecipitated with

anti-Ψ antibody or control immunoglobulin G (IgG). 100ng of 100% Ψ IVT RNA was used as

control. (**c**) Volcano plot of miRNAs enriched by small RNA immunoprecipitation using a Ψ-

specific antibody (Ψ-IP) in mouse fibroblast NIH/3T3 cells compared to input fractions.

Significantly enriched ($p<0.01$) miRNAs are highlighted in blue. (**d**) Validation of enriched

miRNAs using qRT-PCR. (**e**) Ψ-enrichment of miRNAs affected by PUS1 knockdown (KD) 3

days and 6 days after PUS1 shRNA transfection. (**f**) AGO proteins and Dyskerin subunit NHP2

were fused at the N-terminus to N- and C- terminally split YFP, respectively. Positive interaction

is shown by YFP signal, with 35S::RFP as an expression control. (**g**) Ψ-enrichment of precursor

pri-miRNAs in Arabidopsis flower buds plotted against pri-miRNAs bound to NHP2 (blue) and

detected by RIP-seq in 2 or more replicates (orange). (**h**) Violin plot showing significant

difference between NHP2-enriched and non-enriched pri-miRNA and Ψ enrichment of the

mature miRNA by Ψ -IP (Log_2 IP/IgG; (one-way ANOVA; **$p<0.01$).

Extended Data Figure 2

Extended Data Figure 2. Strategies for robust detection of Ψ in small RNAs.

(**a**) Three strategies to detect Ψ in small RNAs and their predicted effect on pseudouridylated

sequences. Details are in Methods (**b**) Known sites of pseudouridylation in tRNAs mediated by

PUS7 (targets UNUAR) and PUS10 (targets Ψ-55). Ψ-55 is 22 nt "upstream" of the 3′ end of -

CCA tailed tRNA fragments. (**c**) Heatmap showing individual tRNA fragments (tRFs) with and

without the UNUAR motif in wild-type and *pus7*-/- libraries treated with CMC. (**d**) 3′CCA tRFs

$\geq$22 nt contain canonical Ψ-55 and are enriched by anti-Ψ antibody and depleted by CMC

treatment, whereas shorter fragments do not contain this nucleoside and are not enriched/depleted.

Ψ-55 is ablated in mutants of PUS10. (**e**) Deletions in 3′ CCA tRFs based on position in

CMC/Mn^{2+}- and mock-treated libraries. (**f, g**) Metaplots showing coverage depletion/enrichment

in CMC (orange)/IP (blue) libraries at rRNA pseudouridylation sites (f) compared to mismatches

from Mn2+/CMC+ libraries (g). Average across 42 predicted pseudouridylated sites in 18S and

25S rRNA and from 17 predicted sites in 18S rRNA (CMC mismatch/Mock mismatch). (**h, i**)

Enrichment/depletion/mismatches across 18S rRNA in CMC/IP (h) and CMC/Mn^{2+} (i) libraries.

Predicted Ψs are marked with dotted lines.

a

b

d

Extended Data Figure 3

Extended Data Figure 3. Arabidopsis mutants in Dyskerin/ NAP57.

(**a**) Generation of *dkc1-2* transgenic plants. *nap57* (SALK_031065) was complemented with the

pABI3:AtNAP57 transgene driven by embryo specific *ABI3* promoter (**b**) *AtNAP57* transcript

level in *dkc1-2* transgenic lines analyzed by qRT-PCR (n=3) (**c**) *dkc1-2* transgenic plants show

severe developmental defects including dwarfism, abnormal leaf shape, deformed and infertile

flowers. (**d**) Ψ-enrichment of pseudouridylated miRNAs in WT and *dkc1-2* mutants (P<0.01 one

way ANOVA).

Extended Data Figure 4

Extended Data Figure 4. Mapping of miRNA pseudouridylation sites in Arabidopsis by

CMC labeling and reverse transcription in the presence of Mn^{2+} (CMC/Mn^{2+}).

(**a**, **b**) Bar chart showing proportions of 5′ and 3′ nucleotides detected by CMC/Mn2+ mismatch

analysis in (**a**) IP-enriched and (**b**) CMC-depleted siRNAs from inflorescence compared to

background. IP-enrichment and CMC-depletion show biases towards 5′ and 3′ U. (**c**) Sites of

predicted Ψ in precursor pri-miR159b amplified from seedlings, leaves and inflorescence of wild-

type (Col-0) and *hyl1-2* microprocessor mutants. had insertions and deletions at varying

frequencies (number of clones after PCR sequencing) corresponding to 25% (Col-0) and 75%

(*hyl1-2)* overall. Terminal sites were also found in mature miR159b in inflorescence (lower panel).

Extended Data Figure 5

Extended Data Figure 5. Pseudouridine in miRNAs and TE-derived siRNAs in pollen from

WT and *paused (psd-13)/*exportin-t mutants.

(**a**) Metaplots of read coverage at 42 known pseudouridylated sites in 18S and 25S rRNA, relative

to untreated controls, in CMC-treated WT (magenta) and Ψ-IP WT (blue) and *psd-13* (orange)

small RNA sequencing libraries from pollen. (**b**) browser of read coverage from the same libraries

at 17 rRNA pseudouridylation sites in 18S rRNA. (**c, d**) metaplot and browser analysis of

predicted sites in rRNA from CMC/Mn^{2+} analysis relative to mock (**e, f**) Metaplots of read

coverage relative to untreated controls in CMC-treated WT (magenta) and WT (blue) and *psd-13*

(orange) Ψ-IP small RNA sequencing libraries from inflorescence. Known positions of Ψ in **b, c**

and **f** are marked with dotted lines. (**g**) Ψ-enrichment of individual miRNA detected by Ψ-IP and

small RNA sequencing in WT and *psd* mutant inflorescence (n=2 biological replicates) (**h**) Ψ-

enrichment of miRNA overall in WT and *psd* mutant inflorescence (one-way ANOVA, p>0.05)

(**i**) Violin plot of Ψ-enrichment of siRNA from individual TEs based on IP-enrichment (Log$_2$

IP/Unbound) in flower buds (inflorescence) and pollen from wt and *psd* mutants. Significant Ψ

enrichment was observed in pollen, but not in inflorescence, and depended on *PSD* (p<0.001, one-

way ANOVA).

	Col-0	*pus7*	*psd-13*	*psd13-/-; pus7+/-*	*psd13+/-; pus7-/-*
a					
b					

c

Extended Data Figure 6

Extended data Figure 6. Mutation of *psd* and *pus7* is synthetically lethal in Arabidopsis.

(**a**) Alexander staining of pollen from Col-0, *pus7*, *psd-13*, *psd-13*-/-; *pus7*+/-, and *psd-13*+/-;

pus7-/- parents. *psd-13*+/; *pus7*-/- had semisterile pollen abortion (loss of Alexander staining,

arrows). (**b**) Mature seeds from each parent. Aborted seeds in *psd-13* are highlighted with arrows.

(**c**) Quantification of approximately 500 seeds from each genotype revealed *psd*-13-/-; *pus7*+/-

have 25% seed abortion, significantly more than *psd-13* alone (<1%; P<0.001). Double

homozygous mutants could not be obtained (n>150 plants). These results suggest that PSD acts in

haploid pollen, while PUS7-mediated pseudouridylation of tRNA acts in diploid pre-meiotic

tissue. Double homozygous embryos are lethal.

Extended Data Figure 7

Extended Data Figure 7. Characterization of pseudouridylation of siRNA in *psd-13* mutants.

(a-c) Metaplots of 20-22 nt (red) and 23-25 nt (black) siRNA abundance across transposable

elements transcribed by Pol IV ("Pol IV-TEs") from biological replicates of **(a)** wild-type, **(b)** *psd-*

13 and **(c)** *hst-6* sperm cells sorted by FACS. 250 bp up- and down-stream regions as well as 500

bp from each end of the TE are shown. **(d-f)** Scatterplots of small RNA sequencing data from two

biological replicates mapped to TE families in **(d)** wild-type, **(e)** *psd-13* and **(f)** *hst-6* sperm cells

isolated by FACS. **(g)** Ψ-enrichment of siRNAs from TEs surrounding maternally expressed

imprinted genes (MEGs) and paternally expressed imprinted genes (PEGs) in wild type (blue) and

psd-13 (orange) pollen. * = p<0.05, ANOVA. **(h)** abundance of 20-22 nt (red) and 23-25 nt (black)

siRNA abundance surrounding imprinted genes in wild-type (blue) and *psd-13* (orange) sperm

cells**.** siRNA in sperm cells matching MEGs and PEGs was lower in *psd-13* mutants for 21-22nt

siRNA (P<0.05) but relatively higher for 23-25nt siRNA (P<0.01). **(i)** levels of miR845a and

miR845b were unchanged or higher in *psd-13* mutant pollen.

Extended Data Figure 8

Extended Data Fig. 8. Pseudouridine in piRNAs and tRNA fragments (tRFs) from mouse testis.

(a, b) Ψ enrichment by (a) Ψ-IP and (b) CMC-depletion of small RNA from testis from 8 weeks-old male mice. Each dot corresponds to tRFs from a single tRNA anticodon isotype. Statistically significant (p-adj. <0.01, log2 fold change > |1|) enriched/depleted tRFs are in black, 3'-tRFs are colored blue; grey dots mark tRFs without significant enrichment/depletion. **(c)** Volcano plot of individual piRNA sequences in testis (8 weeks old males) by CMC-depletion and sequencing. Black: significant fold changes (p-adj. <0.01, log2 fold change > |1|); grey: not significant; blue: significantly enriched/depleted piRNAs overlapping LTR and LINE transposon sequences. **(d)** Volcano plot of individual piRNA sequences. Black: significant fold changes (p-adj. <0.01, log2 fold change > |1|); grey: not significant; blue: significantly enriched/depleted piRNAs overlapping LTR and LINE transposon sequences. **(e, f)** Volcano plot of read counts per piRNA cluster using **(e)** Ψ-IP and **(f)** CMC-detection (I). **(g, h)** Size distribution of (g) 3'-tRFs and (h) piRNA sequences (without structural RNAs) using CMC-depletion and sequencing. **(i, j)** Size distribution of (i) 3'-tRFs and (j) piRNA sequences (G, without structural RNAs) using Ψ-IP and sequencing.