Identification and quantification of transposable element transcripts using Long-Read RNA-seq in *Drosophila* germline tissues

Rita Rebollo[1], Eric Cumunel[2,3], Arnaud Mary[2,3], François Sabot[4], Nelly Burlet[2], Benjamin Gillet[5], Sandrine Hughes[5], Daniel S. Oliveira[2,6], Clément Goubert[7], Marie Fablet[2,11], Cristina Vieira*[2,3] and Vincent Lacroix*[2,3]

[1]Univ Lyon, INRAE, INSA-Lyon, BF2I, UMR 203, 69621 Villeurbanne, France.

[2]Laboratoire de Biométrie et Biologie Evolutive, Université Lyon 1, CNRS, UMR5558, Villeurbanne, Rhone-Alpes, 69100, France

[3]ERABLE team, Inria, Grenoble Rhône-Alpes, Villeurbanne, France

[4]DIADE unit, Univ Montpellier, Cirad, IRD, F-34394 Montpellier Cedex 5, France

[5]Institut de Génomique Fonctionnelle de Lyon (IGFL), UMR5242 CNRS, Ecole Normale Supérieure de Lyon, Université Claude Bernard Lyon 1 , F-69007 Lyon, France.

[6]São Paulo State University (Unesp),Institute of Biosciences, Humanities and Exact Sciences, São José do Rio Preto, SP, Brazil.

[8]Human Genetics, McGill University, Montreal, QC, Canada

[9]Canadian Centre for Computational Genomics, McGill University, Montreal, QC, Canada

[10]McGill Genome Centre, Montreal, QC, Canada.

[11]Institut Universitaire de France (IUF), Paris, Île-de-France , F-75231, France.

*Corresponding authors: vincent.lacroix@univ-lyon1.fr and cristina.vieira@univ-lyon1.fr

Abstract

Transposable elements (TEs) are repeated DNA sequences potentially able to move throughout the genome. In addition to their inherent mutagenic effects, TEs can disrupt nearby genes by donating their intrinsic regulatory sequences, for instance, promoting the ectopic expression of a cellular gene. TE transcription is therefore not only necessary for TE transposition per se but can also be associated with TE-gene fusion transcripts, and in some cases, be the product of pervasive transcription. Hence, correctly determining the transcription state of a TE copy is essential to apprehend the impact of the TE in the host genome. Methods to identify and quantify TE transcription have mostly relied on short RNAseq reads to estimate TE expression at the family level while using specific algorithms to discriminate copy-specific transcription. However, assigning short reads to their correct genomic location, and genomic feature is not trivial. Here we retrieved full-length cDNA (TeloPrime, Lexogen) of *Drosophila melanogaster* gonads and sequenced them using Oxford Nanopore Technologies. We show that long-read RNAseq can be used to identify and quantify TEs at the copy level. In particular, TE insertions overlapping annotated genes are better estimated using long reads than short reads. Nevertheless, long TE transcripts (> 5kb) are not well captured. Most expressed TE insertions correspond to copies that have lost their ability to transpose, and within a family, only a few copies are indeed expressed. Long-read sequencing also allowed the identification of spliced transcripts for

43 around 190 TE copies. Overall, this first comparison of TEs between testes and ovaries uncovers
44 differences in their transcriptional landscape, at the subclass and insertion level.
45
46 **Keywords**: long-read sequencing, ONT, transposable elements, regulation, RNAseq, full-length cDNA
47

Introduction

Transposable elements (TEs) are widespread DNA sequences that have the ability to move around
genomes in a process called transposition (Bourque et al., 2018). TEs can transpose either using an
RNA intermediate, in a copy-and-paste mechanism, *i.e.* retrotransposons, or directly through a DNA
molecule using different cut-and-paste strategies, *i.e.* DNA transposons. In both cases, the synthesis
of a messenger RNA is a fundamental step allowing the production of the transposition machinery,
and hence promoting TE replication in the host genome. TE transposition is *per se* a mutational
process, and several host mechanisms are in place in order to avoid novel TE insertions, including
chromatin remodelling factors, DNA methylation, and small RNAs (Slotkin & Martienssen, 2007). For
instance, in *Drosophila melanogaster* ovaries, TEs are the target of piwi-interacting RNAs (piRNAs) that
promote TE transcript cleavage, but also deposition of repressive chromatin marks within the TE
insertion, blocking any further transcription (Fabry et al., 2021).
In order to appreciate the dynamics of TE regulation, an accurate measure of TE expression is
required, including copy-specific information (Lanciano & Cristofari, 2020). While such analyses may
be easily obtained in genomes composed of mostly ancient TE copies, discrimination of young TE
families, such as LINE-1, AluY and SVA in humans, or the study of genomes composed of mostly active
copies as seen in many insects, remains a complex feat. Indeed, TE copies belonging to the same TE
family have, by definition, more than 80 % of sequence identity, hampering the study of TE regulation
and consequently TE expression in a copy-specific manner (Lanciano & Cristofari, 2020). Most
genome-wide analyses interested in TE expression, and even their regulation, focus on TE family-level
analysis, where short reads are mapped either against TE consensus sequences or to the genome/TE
copy sequences followed by grouping of read counts at the family level (TEcount from the TEtools
package (Lerat et al., 2017), TEtranscripts (Jin et al., 2015)). In the past years, many methods have
surfaced to take advantage of short-read sequencing datasets and circumvent the multi-mapping
problem in order to develop copy-level analysis (for a review see (Lanciano & Cristofari, 2020)). These
methods are based on different algorithms that are able to statistically reassign multi-mapped reads
to unique locations, for instance with the expectation-maximization algorithm used in TEtranscripts
(Jin et al., 2015), SQuIRE (Yang et al., 2019) and Telescope (Bendall et al., 2019).
In the past years, long-read sequencing has become an attractive alternative to study TE biology.
Such reads are able to refine TE annotation (Jiang et al., 2019; Panda & Slotkin, 2020), pinpoint new
TE insertions (Mohamed et al., 2020; Rech et al., 2022), determine TE DNA methylation rates at the
copy level (Ewing et al., 2020), estimate TE expression (Berrens et al., 2022), and finally, detect TE-
gene fusion transcripts (Panda & Slotkin, 2020; Dai et al., 2021; Babarinde et al., 2021). Furthermore,
long-read RNA sequencing can not only determine which TE copies are expressed but also discriminate
between isoforms of a single TE copy produced by alternative splicing. Indeed, TE alternative
transcripts have been described in the very first studies of TEs, using techniques such as northern blot
(Belancio et al., 2006), but concomitantly with accessible short-read genome-wide analysis, low
interest has been given to TE transcript integrity. Nonetheless, TE isoforms have been shown to
participate in TE regulation, as observed for the P element in *D. melanogaster*, where a specific

germline isoform encodes a functional transposase protein, while in somatic tissues, another isoform
acts as a P element repressor (Laski et al., 1986). The regulation of such tissue-specific splicing has
recently been attributed to piRNA-directed heterochromatinization of P element copies (Teixeira et
al., 2017). The retrotransposon *Gypsy* also produces two isoforms, including an envelope-encoding
infectious germline isoform, also controlled by piRNA-guided repressive chromatin marks (Pélisson et
al., 1994; Teixeira et al., 2017). Recently, Panda and Slotkin produced long-read RNA sequencing of
Arabidopsis thaliana lines with defects in TE regulatory mechanisms (Panda & Slotkin, 2020), and were
able to annotate TE transcripts, pinpoint TE splicing isoforms, and most importantly, demonstrate that
properly spliced TE transcripts are protected from small RNA degradation.
D. melanogaster harbours around 12-20% of TE content, and recent studies have suggested that
24 TE superfamilies are potentially active (Adrion et al., 2017). Nevertheless, no indication of which
copies are active has been documented. Here, we describe a bioinformatics procedure using long-
read RNA sequencing, which enables the efficient identification of TE-expressed loci and variation in
TE transcript structure and splicing. Furthermore, our procedure is powerful enough to uncover tissue-
specific differences, as illustrated by comparing testes and ovaries data.
102

Methods

Reference genome and annotation

The dmgoth101 genome assembly was produced from Oxford Nanopore Technologies (ONT) long-
read DNA sequencing and described in (Mohamed et al., 2020). Genome assembly has been deposited
in the European Nucleotide Archive (ENA) under accession number PRJEB50024. Gene annotation was
performed as described in (Fablet et al., 2022). Briefly, we used LiftOff (Shumate & Salzberg, 2021) on
FlyBase *D. melanogaster* annotation (dmel-all-r6.46.gtf.gz), with the option -flank 0.2. In order to
locate and count the reads aligned against TE insertions, we produced a GTF file with the position of
each TE insertion. We have used RepeatMasker with DFAM dataset from *D. melanogaster* TE copies
and then used OneCodeToFindThemAll to merge LTR and internal parts of TE in one unique feature.
Visualization of alignments of TEcopies to their consensus sequences were performed in SnapGene.

Drosophila *rearing and total RNA extraction*

D. melanogaster dmgoth101 strain was previously described by (Mohamed et al., 2020). Briefly,
an isofemale line was derived from a wild-type female *D. melanogaster* from Gotheron, France,
sampled in 2014, and sib-mated for 30 generations. Flies were maintained in 12-hour light cycles, and
24º C, in vials with nutritive medium, in small-mass cultures with approximately 50 pairs at each
generation. For long-read sequencing, forty-five pairs of ovaries and 62 pairs of testes were dissected
in cold PBS 1X from 4 to 8-day-old adults. Total RNA was extracted using the QiagenRNeasy Plus Kit
(Qiagen, reference 74104) after homogenization (using a pellet pestle motor) of the tissues. DNA
contamination was controlled and removed by DNAse treatment for 30 minutes at 37°C (Ambion).
Total RNA was visualized in agarose gel to check DNA contamination and RNA integrity before storing
at -80°C. For short-read sequencing, two replicates per strain were produced, each with 30 pairs of
ovaries and 70 pairs of testes. RNA extraction and DNAse treatment were carried out as for long-read
sequencing. Quality control was performed using an Agilent Bioanalyzer. Libraries were constructed

from mRNA using the Illumina TruSeq RNA Sample Prep Kit following manufacturer's
recommendations. Libraries were sequenced on Illumina HiSeq 3000 with paired-end 150 nt reads.

Long-read RNA-seq and analysis

The two RNA extracts were quantified with RNA BR reagents on Qubit 4.0 (Thermo Fisher
Scientific) and qualified with RNA ScreenTape on Tapestation 4150 instrument (Agilent Technologies),
the results showing no limited quantity and a high quality of the RNA molecules (RIN >9.8). We then
took advantage of the TeloPrime Full-Length cDNA Amplification kit V2 (Lexogen) in order to enrich
ovary and testis total RNA in full-length cDNAs (Figure S1). This protocol is highly selective for mRNAs
that are both capped and polyadenylated and allows their amplification. We determined the optimal
PCR cycle number for each sample by quantitative PCR. The quantity and quality of the cDNA produced
were checked again with Qubit (dsDNA BR) and Tapestation (D5000 DNA ScreenTape) to confirm the
correct amplification of the cDNA and absence of degradation in cDNA fragment length profiles. It is
important to note that we do not have replicates for the long-read dataset. Our primary goal for this
experiment was to evaluate the potential of this technique to identify the largest number of expressed
TE copies and isoforms. Enriched full-length cDNAs generated from ovaries and testes were then
processed into libraries using the SQK-LSK109 ligation kit (ONT) using 3 µg as starting material. The
two libraries were sequenced separately in two flow cells R10 (FLO-MIN110) with a MinION device
(Guppy version 2.3.6 and fast basecalling). We obtained 1 236 000 reads for ovaries and 2 925 554 for
testes that passed the default quality filter (>Q7). Quality control was performed with NanoPlot
v1.39.0 (De Coster et al., 2018). The median read length was 1.4 Kb, and the median quality was 8.4
(*i.e.* one error every 8 bases, Figure S2). Data are available online at the BioProject PRJNA956863.
Reads were mapped to the dmgoth101 genome using minimap2 (version 2.17-r974-dirty) (Li, 2018)
using the splice preset parameter: "minimap2 -ax splice FC30.fastq dmgoth101_assembl_chrom.fasta
-o FC30.bam". Ninety percent of reads were mapped to the genome (Table S1). The vast majority
(99%) mapped to a unique location (i.e. had no secondary alignment reported by minimap2). Out of
the reads with several locations reported, 90% had a unique best location, i.e. the alignment score (AS
flag in the SAM format) was higher for one of the locations. Indeed, if a read has several alignments
with the same alignment score, then this means the read stems from exact repeats in the genome and
they cannot be told apart, hence, we cannot know which copy is transcribed. However, if a read has
several alignments with distinct alignment scores, then it means that the read stems from inexact
repeats. The presence of this read in the dataset means that one of the copies is transcribed and we
consider that it is the one with the highest alignment score. While it could be possible that the read
actually stems from the copy with suboptimal alignment, this is highly unlikely because it would mean
that there is a sequencing error at the position of the divergence between the two copies of the
repeat. A sequencing error in any other position of the read would cause a decrease in the alignment
score of both locations. An example of a read that maps to several locations, one with an alignment
score larger than the others is given in Figure S3.

We also noticed that some reads were only partially mapped to the genome. In practice the query
coverage distribution is bimodal (Figure S4), 80% of reads have a query coverage centered on 90 %,
while the remaining 20 % have a query coverage centered on 50 %. A thorough inspection of the
unmapped regions of these partially mapped reads reveals that they stem from transcripts located
elsewhere on the genome. Given that the transcripts covered by the read are themselves fully covered

(both the primary locus and the secondary locus), we think that these chimeras are artifactual and
were probably generated during ligation steps as previously described (White et al., 2017). Here, we
chose to focus on the locus corresponding to the primary alignment and discard the secondary loci. In
practice, this corresponds to the longest of the two transcripts. We also ran the same analyses after
completely discarding those 20 % chimeric reads, but the quantification of TEs is essentially the same
(R=0.992, Figure S5). In the remainder of the paper, unless stated otherwise, for each read, we focused
on the alignment with the best AS score. We discuss the few cases of ties when required.

Once a read is assigned to a genomic location, it does not yet mean that it is assigned to a genomic
feature. In order to decide which reads could correspond to a TE, we applied the following filters. First,
we selected all reads where the mapping location overlaps the annotation of a TE, for at least one
base. Then, we discarded all reads that covered less than 10 % of the annotated TE. Finally, in the case
where a read mapped to a genomic location where there are several annotated features (a TE and a
gene, or two TEs), we assign the read to the feature whose genomic interval has the smallest
symmetric difference with the one of the read. The rationale for introducing this filter is best explained
with examples. Figure S6 corresponds to a TE annotation overlapping a gene annotation. All reads
map to both features, but the gene is fully covered while the TE is only partially covered. We conclude
that the gene is expressed, not the TE. Figure S7 corresponds to a genomic location where there are
five annotated TEs and a gene, which all overlap. The features that are best covered are Jockey and
the gene.

Most commonly, several features may be partially covered by a read and a read may extend
beyond each of these features. Both the subject and the query coverage need to be modeled. To
address this, we consider genomic intervals as sets of positions. For each pair read-feature, we
compute the symmetric difference between the two intervals, and we assign the read to the feature
with the smallest symmetric difference (Figure S8). This situation occurs frequently and ignoring it
yields an overestimation of TE expression. The impact of each filter is given in Figure S9. After all filters
are applied, there are 1 322 reads in ovaries and 8 219 reads in testes that are assigned to TE copies
(Table S2-S4).

Short-read RNA-seq and analysis

RNA extraction and short-read sequencing were retrieved either from (Fablet et al., 2022), at the
NCBI BioProject database PRJNA795668, or performed here and available at BioProject PRJNA981353.
Briefly, RNA was extracted from 70 testes and 30 ovaries from adults aged three to five days, using
RNeasy Plus (Qiagen) kit following the manufacturer's instructions. After DNAse treatment (Ambion),
libraries were constructed from mRNA using the Illumina TruSeq RNA Sample Prep Kit following the
manufacturer's recommendations. Libraries were sequenced on Illumina HiSeq 3000 with paired-end
150 nt reads. Short-read analysis was performed as in (Fablet et al., 2022), and SQUIRE (Yang et al.,
2019) was used for mapping and counting TE copy-specific expression. A detailed protocol on SQUIRE
usage in non-model species can be found here https://hackmd.io/@unleash/squireNonModel.

Results and discussion

Transposable element transcripts are successfully detected with long-read RNA-seq

In order to understand the TE copy transcriptional activity and transcript isoforms in gonads of *D. melanogaster*, we extracted total RNA from ovaries and testes of dmgoth101 adults, a French wild-derived strain previously described (Mohamed et al., 2020). Prior to long-read sequencing, we enriched the total RNA fraction into both capped and polyadenylated mRNAs in order to select mature mRNAs potentially associated with TE activity (see material and methods for the details on the TeloPrime approach, Figure 1A). Sequencing yielded between ~1 to ~3 million reads per tissue, ranging from 104 to 12 584 bp (median read length ~1.4 Kb, Table S1). Reads were subsequently mapped to the strain-specific genome assembly (Mohamed et al., 2020) using the LR aligner Minimap2 (version 2.17-r974-dirty) (Li, 2018). Ninety percent of reads could be mapped to the genome (Table S1). The vast majority (99%) mapped to a unique location (*i.e.* had no secondary alignment), and 99.9% mapped to a unique best location (see Methods).

In order to validate the long-read RNAseq approach, we first determined the read coverage of all expressed genes, and show that the majority (70.2% in ovaries and 71.8% in testes) harbour at least one read covering more than 80% of their sequence, regardless of their longest transcript length. Besides, few reads correspond to partial transcripts, as most reads (68.9% in ovaries, 78.6% in testes) correspond to well-covered transcripts (>80% coverage) (Figure 1B). This shows that the TeloPrime protocol was successful in capturing full-length transcripts. Besides, the transcriptomes obtained are enriched in typical germline ontology terms, such as "sperm individualization and competition" for testes, and "oogenesis" for ovaries (Figure 1C). Finally, while the first version of the TeloPrime protocol could not be used for quantification (Sessegolo et al., 2019), the quantifications obtained here correlate well with available short-read sequencing for genes harbouring a single isoform (rho=0.78, R=0.74, Figure 1D). These correlation coefficients are in agreement with those obtained when comparing direct RNAseq *vs* Illumina Truseq sequencing (Sessegolo et al., 2019).

Although most long reads map to a unique location on the genome, ensuring that a read stems from the TE copy is not straightforward. First, a read may overlap with a TE copy only for a few nucleotides, suggesting the read is not a consequence of TE transcription. To rule out these cases, all reads covering less than 10% of the annotated TE were discarded. Second, in many instances, the read overlapped both a gene and one or several TEs. In these cases, the feature for which the coverage was best (see Methods and Figure S6-7) was selected. Overall, after applying these filters, 1 322 reads in ovaries and 8 219 reads in testes were assigned to TE copies (Table S2). Out of these, 37-48% overlapped only one TE, 8-18% overlapped with multiple TEs, 16-21% were located in the intron of a gene, and 18-34% overlapped with a gene exon (but were assigned to the TE, not the gene) (Figure 1E).

To ensure this long-read dataset is able to recover transcripts encompassing all TE copy lengths present in the genome, we compared the length distribution of all TE insertions with the length of all mapped reads (Figure 1F). While TE copies range from a few base pairs to ~15 Kb, 75% of annotated copies are smaller than 2 Kb. The average length of reads mapping to TEs encompasses the majority

of TE copies, but does not cover TE transcripts longer than 2 Kb. On the other hand, reads mapping to
genes may reach 5 Kb. The absence of very long reads indicates that either very long mRNAs are absent
from the sample, or the TeloPrime technique is not well tailored for capturing very long transcripts. In
order to clarify this point, we compared the quantification obtained by Illumina and Nanopore
TeloPrime for short (<3 Kb), long (3 Kb-5 Kb) and very long transcripts (>5 Kb), and obtained the
following Spearman correlations of 0.79 (n=3 605), 0.54 (n=378) and 0.38 (n=130), respectively (Figure
1D). We concluded that indeed long and very long transcripts are a minority (9% and 3% of all
transcripts, respectively) but the TeloPrime protocol underestimates their abundance.

Figure 1: Long-read RNAseq of *Drosophila melanogaster* ovaries and testes. A. Experimental layout of
TeloPrime approach. B. Transcript coverage by long-read RNAseq in ovaries and testes. The majority of

transcripts recovered are full-length. C. Gene Ontology enrichment analysis of long-read transcriptomes of
testes and ovaries. In red are enriched terms related to male and female gonadal development. D. Gene
expression quantification using Illumina and ONT sequencing. Each dot is a gene with a single annotated
isoform. Transcripts longer than 5 kb tend to be undersampled using TeloPrime. E. Number of reads mapping
to TEs separated into four categories whether they overlap a TE only, multiple TEs, a TE and an exon, or a TE
and an intron. F. TE copy and read length distribution. Reads mapping to TEs encompass most TE copy length
but lack transcripts longer than 5 Kb, as also observed for reads mapping to genes.

TE mRNA landscape is sex-specific

Taking into account all the filtering steps, only 0.28% (8 219/2 925 554) and 0.11% (1 322/1 236
000) of long reads aligned to TE copies in testes and ovaries respectively, in agreement with the
previous observations using short-read sequencing (Fablet et al., 2022). We identified 130 TE families
capable of producing capped-polyadenylated mRNAs (Table S2), of which 69 belong to Long-terminal
repeat (LTR) elements (retrotransposons that possess LTR sequences surrounding a retroviral-like
machinery). Despite the high number of shared transcribed TE families (99/130), the transcriptional
landscape between ovaries and testes is quite different (Figure 2A). While LTRs dominate the
transcriptional landscape in both tissues, LINE elements are the second most transcribed TE subclass
in testes, while in ovaries, DNA families harbor more read counts (Figure 2A). The transcriptional
landscape within TE subclasses between males and females is very similar for retrotransposons, with
Gypsy and I-Jockey superfamilies being the most expressed LTR and LINE elements respectively.
However, the DNA subclass transcriptional landscape is quite different between males and females:
TcMar-Pogo is the most expressed DNA superfamily in females, while RC elements are abundantly
transcribed in males.

Globally, TE families show higher long-read counts in males compared to females (Figure 2B), both
because male samples were more deeply sequenced (2.3 times more), and because the proportion of
reads that map to TEs is higher in males (0.28% *vs* 0.1%). *HETA* (I-Jockey) and *DNAREP1_D* (RC) are the
top two families in male TE transcript counts, with 734 and 722 long read transcripts respectively. In
females, *pogo* (TcMar-pogo) is the TE family amounting the most transcripts with 224 long reads
(while only 34 in males), followed by *DNAREP1_D* (RC) with 142, and *Copia* (Copia) with 136 long reads
(but both families with higher transcript counts in males). There are only three TE families that are
specific to ovaries, *BARI_Dm* (TcMar-Tc1 - DNA), *Gypsy7* (Gypsy - LTR), and *Helena* (I-Jockey - LINE),
but they all show only one single long read suggesting their expression is low. There are 28 families
specifically transcribed in testes compared to ovaries, five DNA elements (*Transib-N1_DM* (CMC-
Transib), *NOF_FB* (MULE-NOF), *Looper1_Dm* (PiggyBac), and two TcMar-Tc1 (*Bari1* and *Minos*)), 12
LINE elements (nine I-Jockey, two R1, and one R1-LOA, see Table S2 for details), and 11 LTR families
(one Copia, four Gypsy, two Pao and four unknown families), ranging from one to 76 long reads per
TE family. Finally, nine TE families show no long-read mapping in either tissue. Collectively, long-read
sequencing is able to discriminate between female and male TE transcriptional landscapes and
suggests retrotransposons might be strongly and specifically expressed in males compared to females.

Short-read RNA sequencing of ovaries and testes, followed by estimation of TE family expression
with TEcount ((Lerat et al., 2017) - TE expression estimation *per* TE family, see material and methods
for more information)) recapitulates the long-read RNA sequencing profiles (Figure 2C). Despite the
fact that TE transcripts are overall poorly expressed, the estimation of their expression level is
reproducible across technologies. The correlation is higher for testes (r=0.48, rho=0.79) than for

ovaries (r=0.36, rho=0.59), where the coverage is weaker. Indeed, as previously stated, the total
contribution of TE to the transcriptome is weaker for ovaries and the sequencing shallower.

Long-read sequencing successfully retrieves single-copy transcripts

The main objective of using long-read sequencing after the TeloPrime full-length cDNA enrichment
protocol is to recover copy-specific mature TE transcripts and potential isoforms. We uncovered 1 115
and 448 TE copies harbouring at least one long-read transcript unambiguously in testes and ovaries
respectively. Within a TE family, the contribution of each TE copy to the family transcriptional activity
is variable. In general, only a few insertions produce transcripts (Figure 3A). However, *HETA* (I-Jockey)
copies are nearly all expressed in testes (14/15 annotated copies), while in ovaries, *pogo* harbors 17

copies producing transcripts out of 25. *DNArep1* (RC) is the most abundant TE family in the *D.*
melanogaster genome and is also the family harbouring the most transcribed copies in both females
and males (73 and 170 respectively). Finally, in testis, out of the 1 115 insertions with at least one
mapped read, 165 had in fact more than 10 mapped reads. In ovaries, out of the 448 insertions with
at least one mapped read, 26 had more than 10 mapped reads.

In addition, while many TE copies within a family are indeed able to produce transcripts, there are
significant differences in copy expression rate. For instance, in ovaries, where *pogo* has the highest
number of long reads (224), an insertion of 1 352 bp in the X chromosome (X_RaGOO:21,861,530-
21,866,880) accumulates nearly 30% of the family total read count (Figure 3B and C). This specific
pogo insertion is located in an intron of a gene (CG12061), at 370 bp from the transcription start site,
and anti-sense to the gene. CG12061 is a potential calcium exchange transmembrane protein and has
been previously shown to be highly expressed in the male germline (Li et al., 2022). Indeed, using long-
read sequencing, CG12061 is highly expressed in testes compared to ovaries, and curiously, the
intronic *pogo* insertion is only expressed when the gene is silent (in ovaries) (Figure 3D). The same
pattern is observed for the second most expressed *pogo* insertion, also located in the intron of a gene,
expressed in testes and not in ovaries (Figure S10). The other expressed insertions of *pogo* are located
in intergenic regions (Figure S11).

Finally, using short-read sequencing and a tool developed to estimate single-copy expression
(Squire (Yang et al., 2019)), we compared the overall TE copy transcriptional landscape between short
and long reads (Figure S12). There was a poor correlation with the ONT estimations (rho=0.22, r=0.17
for ovaries and rho=0.38, r=0.33 for testes). Examination of instances where the two techniques
differed most, shows that Squire tends to overestimate the expression of TE insertions overlapping
genes (Figure S13). At the family level, the quantifications obtained by Squire were comparable to the
ones obtained with TEcount.

Figure 3: Transcription of transposable element copies. A. Frequency of transcribed copies (read > 1) within TE subfamilies in testes and ovaries, along with genomic copy number (color bar, 1 to 200 (LINE/LTR) or 300 (DNA) copies). All TE families harbouring more than 200 (LINE) or 300 (DNA) copies are depicted in pink. For DNA elements, *DNAREP1_DM* harbours 2 472 copies, and the LINE family *DMCR1A* has 534 copies. Most TE subfamilies have only a couple of copies producing transcripts, while the majority of HETA copies are expressed in testes for instance (middle panel). B. Distribution of read count per copy for *pogo*, *HETA*, *Copia*, *GYPSY_10* and *GYPSY_12* showing the overall expression of specific copies within a TE family (Table S3 and S4). Copies are represented by different colors within a bar graph. C. Heatmap of copy read count per TE subfamily in testes and ovaries. *Pogo*, *HETA* and *Copia* are depicted. D. Simplified IGV screenshot of a *pogo* copy (X_RaGOO:21,861,530-21,866,88, in pink). In green, testis coverage and an excerpt of mapped reads, in purple

354   the same information for ovaries. Dmgoth101 repeat and gene tracks are also shown and more information on
355   the annotation can be seen in the material and methods section.

Full-length transposable element copies are rarely transcribed

Many insertions produce transcripts that are shorter than the annotated TE and are likely unable
to participate in TE transpositional activity. Furthermore, even in the case where the transcript fully
covers the insertion, the copy itself might have accumulated mutations, insertions and deletions
making it unable to transpose. To assess this, we computed the query coverage of the reads w.r.t. the
insertion they correspond to. We find that one-third of the insertions have at least 80% query
coverage (Figure 4A). However, out of these insertions, only a few of them are close in length to a
functional full-length sequence. Indeed, a close analysis of *pogo*, one of the most expressed TE
subfamilies in ovaries, shows only two potential complete copies expressed, and both at low levels (6
to 4 reads in ovaries, and 2 to 3 in testes, Figure 4B). Instead, the most expressed *pogo* copies in
ovaries (73 and 76 long reads), and in testes (18 long reads) are internally deleted. In order to search
for potentially functional, expressed copies, we filtered copies that produced at least five long reads
in ovaries, and ten in testes (due to coverage differences), and covered at least 80% of their consensus
sequences. In ovaries these filters correspond to seven insertions: one *pogo*, five *Copia* insertions and
one *MAX*. In testes, there are nine potentially functional insertions: five *Copia* insertions, three
Mariner2, and one *HETA* (Figure 4B). Collectively, long-read sequencing shows that despite the
presence of potentially functional, full-length copies in the *D. melanogaster* genome, only a few of
these copies are indeed transcribed in testes and ovaries.

Figure 4. TE transcripts stem mostly from deleted or truncated copies. A. Coverage of ONT reads on TE insertions. One-third of copies are covered for at least 80% of their length. B. Alignment of transcribed copies against their consensus for three major subfamilies: *pogo*, *Mariner2* and *Copia*, all harboring potentially full-length expressed copies. Read count is shown for each copy and full-length copies (> 80% coverage, reads > 5 in ovaries and 10 in testis) and their long-read count are depicted in red.

Long-read sequencing unveils novel spliced TE isoforms

While most TEs do not harbor introns, there are a couple of exemptions previously described in *D. melanogaster.* Indeed, P elements are known to be regulated tissue-specifically by alternative splicing mechanisms, involving piRNA targeting (Laski et al., 1986; Teixeira et al., 2017). *Gypsy* copies are able to produce ENV proteins through mRNA alternative splicing, also regulated by piRNAs (Pélisson et al., 1994; Teixeira et al., 2017). Finally, *Copia* elements produce two isoforms, a 5 Kb and a 2.1 Kb (which is a spliced product of the 5 Kb mRNA) (Miller et al., 1989; Yoshioka et al., 1990). The 2.1 Kb encodes the GAG protein and is produced at higher levels than the other proteins (Brierley & Flavell, 1990). While the shorter transcript can be processed by *Copia* reverse transcriptase, the 5 Kb full-length isoform is clearly preferred (Yoshioka et al., 1990). Most of these discoveries were obtained through RT-PCR sequencing of amplicons, or recently, through short-read mapping. Nevertheless, systematic analysis of TE alternative splicing in *D. melanogaster* is lacking, due to the difficulty of detecting such isoforms from short-read data. Here we used long-read sequencing to mine for such splicing isoforms. We searched for reads harboring a gap compared to the reference sequence (presence of N's in the CIGAR string). In order to ensure that those gaps corresponded to introns, we searched for flanking GT-AG splice sites. This was the case for more than 90% of the gaps (155/172 in testes, 37/40 in ovaries) The remaining cases likely correspond to genomic deletions. While most TE copies harbor intronless transcripts (visible in Figure 5, as circles located at 0 in the X-axis), there is a significant number of TE transcripts, both in testes and ovaries, that harbor spliced transcripts (circles deviating from 0 in the X-axis).

Figure 5. TE spliced transcripts are ubiquitous. Each circle depicts a TE insertion and their size is proportional to the expression level of the insertion. The TE family name is written for highly expressed insertions (>30 reads in testis, >10 reads in ovaries). The X-axis represents the proportion of reads that align with a gap (presence of N's in the CIGAR string), while the Y-axis represents the proportion of the insertion covered by reads. Unfilled circles correspond to gaps that do not exhibit GT-AG sites.

While the proportion of spliced transcripts stemming from a TE copy can vary, there are a couple of copies that only produce spliced transcripts, as POGO$2R_RaGOO$7201268$7202754 for instance. *Pogo* is the most expressed TE family in ovaries, with 17 out of 25 copies producing capped poly-A

transcripts corresponding to 224 long reads, while only 7 expressed copies with a total of 34 long reads
are observed in testes, despite the higher coverage. While we previously noted that only two full-
length copies are transcribed in ovaries (and in testes also but albeit with a lower number of reads),
there are many truncated or deleted copies that are transcribed (Figure 4B).
POGO$2R_RaGOO$7201268$7202754 is one of the internally deleted copies, and it produces a spliced
transcript present in both testis and ovaries (Figure 6A). The splicing of this short intron (55 nt) has
been previously reported (Tudor et al., 1992) and enables the splicing of the two ORFs of *pogo* into a
single continuous ORF. This particular copy (POGO$2R_RaGOO$7201268$7202754) is however non-
functional since it contains a large genomic deletion located in the ORF near the intron (Figure 6B).
POGOX_RaGOO21863530$21864880 (Figure 3D) also contains a large genomic deletion,
encompassing the intron, explaining why there are no spliced transcripts for this copy (Figure 6B).

Figure 6. *Pogo* splicing in testes and ovaries. A. Simplified IGV screenshot shows spliced transcripts using long-
read datasets. In green, testis coverage and an excerpt of mapped reads, in purple the same information for
ovaries. Dmgoth101 repeat and gene tracks are also shown and more information on the annotation can be
seen in the material and methods section.B. Visualization of *pogo* splicing against the consensus sequence.
Both *pogo* copies (spliced and unspliced) are depicted, along with POGO$2R_RaGOO$7201268$7202754
spliced transcript. In pink, the ORF1 encoded by *pogo*.

Despite the presence of full-length *Copia* insertions in the genome, only spliced transcripts were
uncovered in the long-read sequencing (Figure 5 and 7). In contrast, with Illumina short reads, we see
both spliced and unspliced transcripts (Figure 7A). The full-length transcripts are 5 Kb, and are less
abundant than the spliced transcripts (~10 times less). The lack of such a full-length transcript in the
long-read sequencing data might be explained by the lower expression level and the length of the
transcript. One can not discard the possibility that deeper long-read coverage might uncover full-
length, unspliced, *Copia* transcripts. It is important to stress that by using only short reads it is nearly
impossible to determine which *Copia* sequence is being expressed as the vast majority of short reads
map to multiple locations with the exact same alignment score. With short reads, at least one full-
length *Copia* insertion is expressed but its specific location remains unknown. Furthermore, if we
restrict the analysis to primary alignments (i.e. a randomly chosen alignment in the case of multiple

mapping), then the coverage of the intronic sequence decreases and it is no longer clear if the
insertion produces both spliced and unspliced transcripts (Figure S13). Overall, for *Copia*, the long
reads enable the identification of which insertion is being transcribed, and the short reads enable the
detection of the presence of the two splice variants. Finally, spliced transcripts are unable to produce
the complete transposition machinery as they lack the reverse transcriptase enzyme and are only able
to produce the gag protein (Figure 7B).

Figure 7. Example of *Copia* splicing. A. Simplified IGV screenshot shows spliced transcripts using long-read
datasets. In green, testis coverage and an excerpt of mapped reads, in purple the same information for
ovaries. Short-read data is also present in addition to ONT reads. In the excerpt of mapped reads, white
rectangles correspond to multi-mapping reads. Dmgoth101 repeat and gene tracks are also shown and more
information on the annotation can be seen in the material and methods section. B. Visualization of *Copia*
splicing against the consensus sequence. Copia_LTR$3L_RaGOO$10022428$10027467 genomic sequence and
its spliced transcripts are depicted. In pink, the ORF encoded by *Copia*.

While other examples of spliced transcripts are observed for different families that we have
manually curated (*FB4, TART-A, TAHRE* and others), one particular case involves the *1731*
retrotransposon (Figure 8A). Indeed, the copy DM1731_LTR$3L_RaGOO$24209333$24213898,
encoding a GAG and POL open reading frames, produces a 1.8 kb internal transcript, corresponding to
truncated integrase protein and reverse transcriptase, and a spliced 1.3 kb transcript, spanning the
full 4.5 kb genomic copy but splicing a 3.2 kb intron, and producing the RNAseH protein (Figure 8B).
Both transcripts are present in testes, while no expression of this copy is observed in ovaries.

Figure 8. Example of *1731* splicing. A. Simplified IGV screenshot shows spliced transcripts using long-read datasets. In green, testis coverage and an excerpt of mapped reads, in purple the same information for ovaries. Dmgoth101 repeat and gene tracks are also shown and more information on the annotation can be seen in the material and methods section. B. Visualization of *1731* splicing against the consensus sequence. DM1731_LTR$3L_RaGOO$24209333$24213898 genomic sequence and its spliced transcripts are depicted. In pink, both ORFs encoded by *1731*.

Conclusion

Long-read sequencing remains a major progress in the study of repeat transcription. Here we demonstrated the feasibility of assigning long reads to specific copies, which remains the biggest issue in TE expression analysis. In addition, quantification of TE expression with long-read sequencing is similar to short-read analysis, suggesting not only one could recover copy-specific information but also perform quantitative and differential expression analysis.

The genome of *D. melanogaster* contains many functional full-length copies but only a couple of such copies produce full-length transcripts in gonads. Given TEs are tightly controlled in the germline, one can wonder how many full-length copies might be expressed in somatic tissues. It is also important to stress that, to our knowledge, this is the first comparison of the expression of TEs between testes and ovaries, and we demonstrate a different TE transcriptional landscape regarding TE subclasses. Interestingly, some insertions like POGOX_RaGOO21863530$21864880, located in the intron of a gene, are expressed only in ovaries and seem to have a silencing effect on their host gene. Furthermore, in many instances, we see that TEs are spliced, independently of their structure or class. While some of these introns had been reported in the literature 30 years ago, the relevance and prevalence of these spliced transcripts have not always been investigated. Long-read sequencing could facilitate the exhaustive inventory of all spliceforms, in particular for recent TEs, where short reads are harder to use due to multiple mapping.

Finally, it is important to note that we did not recover TE transcripts longer than 2 Kb, despite gene transcripts up to 5 Kb. While the sequencing coverage might indeed play a role in the detection of rare transcripts, it would be important to verify if such long transcripts necessitate different RNA extraction methods. For instance, the distribution of cDNA used here for ONT library construction reflects the distribution of reads, with a low number of cDNAs longer than 3.5 Kb (Figure S1).

Acknowledgements

This work was performed using the computing facilities of the LBBE/PRABI.

Data, scripts, code, and supplementary information availability

Data are available online at the BioProject PRJNA956863 and PRJNA981353.

Conflict of interest disclosure

The authors declare that they have no financial conflicts of interest in relation to the content of the article.

Funding

This work was supported by French National research agency (ANR project ANR-16-CE23-0001 'ASTER and ANR-20-CE02-0015 LongevitY), along with UDL-FAPESP2020.

References

Adrion JR, Song MJ, Schrider DR, Hahn MW, Schaack S (2017) Genome-Wide Estimates of Transposable Element Insertion and Deletion Rates in Drosophila Melanogaster. *Genome Biology and Evolution*, **9**, 1329–1340. https://doi.org/10.1093/gbe/evx050

Babarinde IA, Ma G, Li Y, Deng B, Luo Z, Liu H, Abdul MM, Ward C, Chen M, Fu X, Shi L, Duttlinger M, He J, Sun L, Li W, Zhuang Q, Tong G, Frampton J, Cazier J-B, Chen J, Jauch R, Esteban MA, Hutchins AP (2021) Transposable element sequence fragments incorporated into coding and noncoding transcripts modulate the transcriptome of human pluripotent stem cells. *Nucleic Acids Research*, **49**, 9132–9153. https://doi.org/10.1093/nar/gkab710

Belancio VP, Hedges DJ, Deininger P (2006) LINE-1 RNA splicing and influences on mammalian gene expression. *Nucleic Acids Research*, **34**, 1512–1521. https://doi.org/10.1093/nar/gkl027

Bendall ML, Mulder M de, Iñiguez LP, Lecanda-Sánchez A, Pérez-Losada M, Ostrowski MA, Jones RB, Mulder LCF, Reyes-Terán G, Crandall KA, Ormsby CE, Nixon DF (2019) Telescope: Characterization of the retrotranscriptome by accurate estimation of transposable element expression. *PLOS Computational Biology*, **15**, e1006453. https://doi.org/10.1371/journal.pcbi.1006453

Berrens RV, Yang A, Laumer CE, Lun ATL, Bieberich F, Law C-T, Lan G, Imaz M, Bowness JS, Brockdorff N, Gaffney DJ, Marioni JC (2022) Locus-specific expression of transposable elements in single cells with CELLO-seq. *Nature Biotechnology*, **40**, 546–554. https://doi.org/10.1038/s41587-021-01093-1

Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvák Z, Levin HL, Macfarlan TS, Mager DL, Feschotte C (2018) Ten things you should know about transposable elements. *Genome Biology*, **19**, 199. https://doi.org/10.1186/s13059-018-1577-z

Brierley C, Flavell AJ (1990) The retrotransposon copia controls the relative levels of its gene products post-transcriptionally by differential expression from its two major mRNAs. *Nucleic Acids Research*, **18**, 2947–2951. https://doi.org/10.1093/nar/18.10.2947

Dai Z, Ren J, Tong X, Hu H, Lu K, Dai F, Han M-J (2021) The Landscapes of Full-Length Transcripts and Splice Isoforms as Well as Transposons Exonization in the Lepidopteran Model System, Bombyx mori. *Frontiers in Genetics*, **12**, 704162.

532    https://doi.org/10.3389/fgene.2021.704162

533 De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C (2018) NanoPack: visualizing and
534    processing long-read sequencing data. *Bioinformatics*, **34**, 2666–2669.
535    https://doi.org/10.1093/bioinformatics/bty149

536 Ewing AD, Smits N, Sanchez-Luque FJ, Faivre J, Brennan PM, Richardson SR, Cheetham SW,
537    Faulkner GJ (2020) Nanopore Sequencing Enables Comprehensive Transposable Element
538    Epigenomic Profiling. *Molecular Cell*, **80**, 915-928.e5.
539    https://doi.org/10.1016/j.molcel.2020.10.024

540 Fablet M, Salces-Ortiz J, Jacquet A, Menezes BF, Dechaud C, Veber P, Noûs C, Rebollo R, Vieira C
541    (2022) A quantitative, genome-wide analysis in Drosophila reveals transposable elements'
542    influence on gene expression is species-specific. https://doi.org/10.1101/2022.01.20.477049

543 Fabry MH, Falconio FA, Joud F, Lythgoe EK, Czech B, Hannon GJ (2021) Maternally inherited
544    piRNAs direct transient heterochromatin formation at active transposons during early
545    Drosophila embryogenesis. *eLife*, **10**, e68573. https://doi.org/10.7554/eLife.68573

546 Jiang F, Zhang J, Liu Q, Liu X, Wang H, He J, Kang L (2019) Long-read direct RNA sequencing by
547    5'-Cap capturing reveals the impact of Piwi on the widespread exonization of transposable
548    elements in locusts. *RNA biology*, **16**, 950–959.
549    https://doi.org/10.1080/15476286.2019.1602437

550 Jin Y, Tam OH, Paniagua E, Hammell M (2015) TEtranscripts: a package for including transposable
551    elements in differential expression analysis of RNA-seq datasets. *Bioinformatics (Oxford,*
552    *England)*, **31**, 3593–3599. https://doi.org/10.1093/bioinformatics/btv422

553 Krupovic M, Blomberg J, Coffin JM, Dasgupta I, Fan H, Geering AD, Gifford R, Harrach B, Hull R,
554    Johnson W, Kreuze JF, Lindemann D, Llorens C, Lockhart B, Mayer J, Muller E, Olszewski
555    NE, Pappu HR, Pooggin MM, Richert-Pöggeler KR, Sabanadzovic S, Sanfaçon H, Schoelz
556    JE, Seal S, Stavolone L, Stoye JP, Teycheney P-Y, Tristem M, Koonin EV, Kuhn JH (2018)
557    Ortervirales: New Virus Order Unifying Five Families of Reverse-Transcribing Viruses.
558    *Journal of Virology*, **92**, e00515-18. https://doi.org/10.1128/JVI.00515-18

559 Lanciano S, Cristofari G (2020) Measuring and interpreting transposable element expression. *Nature*
560    *Reviews. Genetics*, **21**, 721–736. https://doi.org/10.1038/s41576-020-0251-y

561 Laski FA, Rio DC, Rubin GM (1986) Tissue specificity of Drosophila P element transposition is
562    regulated at the level of mRNA splicing. *Cell*, **44**, 7–19. https://doi.org/10.1016/0092-
563    8674(86)90480-0

564 Lerat E, Fablet M, Modolo L, Lopez-Maestre H, Vieira C (2017) TEtools facilitates big data
565    expression analysis of transposable elements and reveals an antagonism between their activity
566    and that of piRNA genes. *Nucleic Acids Research*, **45**, e17.
567    https://doi.org/10.1093/nar/gkw953

568 Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*, **34**, 3094–3100.
569    https://doi.org/10.1093/bioinformatics/bty191

570 Li H, Janssens J, De Waegeneer M, Kolluru SS, Davie K, Gardeux V, Saelens W, David FPA, Brbić
571    M, Spanier K, Leskovec J, McLaughlin CN, Xie Q, Jones RC, Brueckner K, Shim J, Tattikota
572    SG, Schnorrer F, Rust K, Nystul TG, Carvalho-Santos Z, Ribeiro C, Pal S, Mahadevaraju S,
573    Przytycka TM, Allen AM, Goodwin SF, Berry CW, Fuller MT, White-Cooper H, Matunis
574    EL, DiNardo S, Galenza A, O'Brien LE, Dow JAT, FCA Consortium, Jasper H, Oliver B,
575    Perrimon N, Deplancke B, Quake SR, Luo L, Aerts S (2022) Fly Cell Atlas: A single-nucleus
576    transcriptomic atlas of the adult fruit fly. *Science*, **375**, eabk2432.
577    https://doi.org/10.1126/science.abk2432

578 Miller K, Rosenbaum J, Zbrzezna V, Pogo AO (1989) The nucleotide sequence of Drosophila
579    melanogaster copia-specific 2.1-kb mRNA. *Nucleic Acids Research*, **17**, 2134.
580    https://doi.org/10.1093/nar/17.5.2134

581 Mohamed M, Dang NT-M, Ogyama Y, Burlet N, Mugat B, Boulesteix M, Mérel V, Veber P, Salces-
582    Ortiz J, Severac D, Pélisson A, Vieira C, Sabot F, Fablet M, Chambeyron S (2020) A
583    Transposon Story: From TE Content to TE Dynamic Invasion of Drosophila Genomes Using
584    the Single-Molecule Sequencing Technology from Oxford Nanopore. *Cells*, **9**, 1776.
585    https://doi.org/10.3390/cells9081776

586 Panda K, Slotkin RK (2020) Long-Read cDNA Sequencing Enables a "Gene-Like" Transcript

Annotation of Transposable Elements. *The Plant Cell*, **32**, 2687–2698.
https://doi.org/10.1105/tpc.20.00115
Pélisson A, Song SU, Prud'homme N, Smith PA, Bucheton A, Corces VG (1994) Gypsy transposition
correlates with the production of a retroviral envelope-like protein under the tissue-specific
control of the Drosophila flamenco gene. *The EMBO journal*, **13**, 4401–4411.
https://doi.org/10.1002/j.1460-2075.1994.tb06760.x
Rech GE, Radío S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H, Jamilloux V,
Quesneville H, González J (2022) Population-scale long-read sequencing uncovers
transposable elements associated with gene expression variation and adaptive signatures in
Drosophila. *Nature Communications*, **13**, 1948. https://doi.org/10.1038/s41467-022-29518-8
Sessegolo C, Cruaud C, Da Silva C, Cologne A, Dubarry M, Derrien T, Lacroix V, Aury J-M (2019)
Transcriptome profiling of mouse samples using nanopore sequencing of cDNA and RNA
molecules. *Scientific Reports*, **9**, 14908. https://doi.org/10.1038/s41598-019-51470-9
Shumate A, Salzberg SL (2021) Liftoff: accurate mapping of gene annotations. *Bioinformatics*
(Oxford, England), **37**, 1639–1643. https://doi.org/10.1093/bioinformatics/btaa1016
Slotkin RK, Martienssen R (2007) Transposable elements and the epigenetic regulation of the
genome. *Nature Reviews Genetics*, **8**, 272–285. https://doi.org/10.1038/nrg2072
Teixeira FK, Okuniewska M, Malone CD, Coux R-X, Rio DC, Lehmann R (2017) piRNA-mediated
regulation of transposon alternative splicing in the soma and germ line. *Nature*, **552**, 268–
272. https://doi.org/10.1038/nature25018
Tudor M, Lobocka M, Goodell M, Pettitt J, O'Hare K (1992) The pogo transposable element family
of Drosophila melanogaster. *Molecular & general genetics: MGG*, **232**, 126–134.
https://doi.org/10.1007/BF00299145
Wei K, Aldaimalani R, Mai D, Zinshteyn D, Prv S, Blumenstiel JP, Kelleher ES, Brooks E (2022)
Rethinking the "gypsy" retrotransposon: A roadmap for community-driven reconsideration of
problematic gene names. https://doi.org/10.31219/osf.io/fma57
White R, Pellefigues C, Ronchese F, Lamiable O, Eccles D (2017) Investigation of chimeric reads
using the MinION. *F1000Research*, **6**, 631. https://doi.org/10.12688/f1000research.11547.2
Yang WR, Ardeljan D, Pacyna CN, Payer LM, Burns KH (2019) SQuIRE reveals locus-specific
regulation of interspersed repeat expression. *Nucleic Acids Research*, **47**, e27.
https://doi.org/10.1093/nar/gky1301
Yoshioka K, Honma H, Zushi M, Kondo S, Togashi S, Miyake T, Shiba T (1990) Virus-like particle
formation of Drosophila copia through autocatalytic processing. *The EMBO journal*, **9**, 535–
541. https://doi.org/10.1002/j.1460-2075.1990.tb08140.x
621