Ripening and synchronization of biomolecular condensates in a heterogeneous elastic medium

Lingyu Meng,[1] Sheng Mao,[2] and Jie Lin[1,3]

[1]*Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies,*
Peking University, Beijing 100871, China
[2]*Department of Mechanics and Engineering Science, BIC-ESAT, College of Engineering,*
Peking University, Beijing 100871, China
[3]*Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies,*
Peking University, Beijing 100871, China

(Dated: May 27, 2023)

Biomolecular condensates play a crucial role in regulating gene expression, but their behavior in chromatin remains poorly understood. Classical theories of phase separation are limited to thermal equilibrium, and traditional methods can only simulate a limited number of condensates. In this paper, we introduce a novel mean-field-like method that allows us to simulate millions of condensates in a heterogeneous elastic medium to model the dynamics of transcriptional condensates in chromatin. Using this method, we uncover an elastic ripening process in which the average condensate radius exhibits a unique temporal power-law scaling, $\langle R \rangle \sim t^{1/5}$ and derive the exponent based on energy conservation and scale invariance. We also introduce active dissolution to model the degradation of transcriptional condensates upon RNA accumulation. Surprisingly, three different kinetics of condensate growth emerge, corresponding to constitutively expressed, transcriptional-bursting, and silenced genes. Furthermore, the timing of growth initiation can be synchronized among bursting condensates, with power-law scaling between the synchronization period and dissolution rate. Our results shed light on the complex interplay between biomolecular condensates and the elastic medium, with important implications for gene expression regulation.

I. INTRODUCTION

Biomolecular condensates are membraneless cellular compartments with many crucial physiological functions, e.g., stress adaptation, accelerating biochemical reactions, reducing noise [1–6]. Specifically, transcription-dependent condensates comprised of RNA polymerases (RNAPs) are observed in both prokaryotes and eukaryotes [7–12], suggesting an essential role of RNAP condensates in gene expression regulation. Biomolecular condensates are often liquid droplets forming via liquid-liquid phase separation (LLPS) [2, 13], supported by their fluid-like behaviors [1, 14–16]. Classical LLPS theories focus on liquid droplets in a liquid environment at or evolving towards thermal equilibrium [13, 17, 18]. According to classical LLPS theories, molecules flow from small to big condensates to reduce the total surface energy, called Ostwald ripening. The outcome is a single large condensate because this configuration minimizes the surface energy. However, in many cases, the surrounding environments of biomolecular condensates are not simple viscous liquids, e.g., the nucleoplasm is filled with chromatin, and the cytoplasm contains cytoskeleton. It has been found that the elastic medium can interact with condensates and affect their growth and coarsening [19–24]. For example, light-activated condensates *in vivo* were observed to be constrained by the chromatin as the condensates could only stay in the chromatin-sparse region [22]. Soft matter experiments also found that the droplet-forming molecules flow from droplets in stiff medium to droplets in soft medium, suggesting a new driving force of coarsening due to elasticity beyond the classical LLPS theories [25, 26]. It is still unclear whether the elastic driving force generates any novel universality class of the ripening process beyond the Ostwald ripening.

Furthermore, numerous out-of-equilibrium processes consume energy in natural biological systems. In particular, the effects of active chemical reactions on the formation and morphologies of condensates have been intensely studied recently [27–29]. One notable function of active chemical reactions is to generate multiple stable coexisting condensates beyond the classical Ostwald ripening [1, 30–33]. While active chemical reactions consume energies, such as ATP, alternative mechanisms to generate coexisting condensates have been proposed, e.g., through the nonlinear elasticity of surrounding network [20, 21, 24], which do not consume energies. In the meantime, other out-of-equilibrium processes also play crucial roles in condensate formation and dissolution. For example, recent experiments found that RNAP condensates can dissolve simply due to the accumulation of transcribed mRNAs, which is also an active process consuming energies [34].

In this work, we study the out-of-equilibrium dynamics of biomolecular condensates in a heterogeneous elastic medium. Regarding the theoretical tools, we introduce a novel mean-field-like model, which allows us to simulate millions of condensates simultaneously and significantly exceeds the computation capacity of traditional phase-field simulations [13]. We first study the case of a neo-Hookean elastic medium in which each condensate is confined by a constant local elastic pressure that can

FIG. 1. Biomolecular condensates grow in the interspace of a heterogeneous elastic medium. Each condensate has a fixed inside concentration c_{in} and a shared outside concentration c_{out}. Condensates' growth is suppressed by the surrounding elastic medium through a confining pressure, which can be different among condensates.

be different among condensates. Surprisingly, we find a
new dynamical scaling of the average condensate radius
$\langle R \rangle \sim t^{1/5}$ during the ripening phase induced by hetero-
geneous elasticity, which we denote as elastic ripening.
The 1/5 exponent is beyond the 1/3 exponent in the clas-
sical Ostwald ripening, and we derive its value based on
energy conservation and scale invariance.

We also introduce nonlinear elasticity beyond the neo-
Hookean model in which the ripening of condensates
can be suppressed, and multiple condensates can coex-
ist. However, within this nonlinear model, the system
quickly reaches an equilibrium state without any tem-
poral changes. To incorporate biological activity, we as-
sume that condensates dissolve at a rate proportional
to their volume inspired by experiments in which RNAs
can dissolve transcription-related condensates [34]. In-
terestingly, depending on the local stiffness around con-
densates, the condensates can grow immediately after
dissolution, grow intermittently, or be suppressed en-
tirely, which we propose to correspond to constitutively
expressed, transcriptional-bursting, and silenced genes.
Surprisingly, a subset of the intermittently growing con-
densates can be synchronized to start growing simulta-
neously: a subset of intermittently growing condensates
with similar local stiffness always grow together. We in-
vestigate the fraction of synchronized condensates and
find a power-law scaling between the synchronization pe-
riod with the dissolution rate. Our work reveals a new
universality class for the ripening process of condensates
in a heterogeneous elastic medium and uncovers the po-
tential roles of chromatin elasticity in gene expression
regulation.

II. THE CONDENSATE GROWTH MODEL

We propose a mean-field-like model to describe the
growth dynamics of condensates in a heterogeneous elas-
tic medium (Figure 1). We assume a fixed concentra-
tion c_{in} inside the condensates and an outside concen-
tration c_{out} shared by all condensates. This assumption
is justified by typical phase-field simulations in which an
almost uniform outside concentration is observed (Sup-
plementary Material and Figure S3). We assume near-
equilibrium dynamics so that the changing rate of a con-
densate's volume is proportional to the derivative of free
energy with its volume,

$$\frac{\partial V}{\partial t} = -kV^{\frac{1}{3}}\frac{\partial F}{\partial V} = kV^{\frac{1}{3}}(g - p_c), \qquad (1)$$

130 where g is the reduced free energy for forming conden-
131 sates per unit volume, which we call condensing affinity
132 in the following, and

$$g = c_{\text{in}}k_B T \ln\left(\frac{c_{\text{out}}}{c_0}\right) - k_B T(c_{\text{out}} - c_0). \qquad (2)$$

Here, c_0 is the saturated concentration of phase sepa-
ration and $g = 0$ when $c_{\text{out}} = c_0$ as expected (Ap-
pendix A). The confining pressure $p_c = p_s + p_{\text{el}}$. Here,
$p_s = 2\gamma(\frac{4\pi}{3V})^{1/3}$ is the Laplace pressure and γ is the sur-
face tension. p_{el} is the elastic pressure due to the sur-
rounding medium, and we will explain its value later.
The $V^{1/3}$ factor on the right side of Eq. (1) comes
from the spherical geometry. The particle flux enter-
ing an absorbing sphere is proportional to its radius in
three dimensions: $J \propto 1/R \times R^2 \sim R$ where $1/R$ is
for the concentration gradient and R^2 is for the surface
area. As we show later, this $V^{1/3}$ factor is important
to obtain the correct scaling of the Ostwald ripening:
$\langle R \rangle \sim t^{1/3}$ where $\langle R \rangle$ is the average condensate radius
and t is the time since initiation. By selecting the en-
ergy unit $\epsilon_0 = k_B T$, the length unit $l_0 = c_0^{-1/3}$ and the
time unit $t_0 = 1/(kc_0^2 l_0 k_B T)$, we non-dimensionalize the
equation so that all the variables become dimensionless,
$\partial V/\partial t = V^{\frac{1}{3}}(g - p_c)$ with $g = \frac{c_{\text{in}}}{c_0}\ln\left(\frac{c_{\text{out}}}{c_0}\right) - \left(\frac{c_{\text{out}}}{c_0} - 1\right)$.

In this work, we mainly focus on the nucleation regime
of phase separation, which is the typical scenario for bi-
ological systems [35, 36]. We assign each condensate a
nucleation radius R_n as its minimum size, which can be
related to the lengths of some specific DNA sequences
such as the promoters [12]. The condensing affinity must
overcome the Laplace and elastic pressures for the con-
densate to grow. Therefore, the larger R_n, the easier the
condensate to initiate growth. In the following numerical
simulations, we set $c_{\text{in}} = 10$, the initial outside concen-
tration $c_{\text{out, ini}} = 2$ and $\gamma = 0.1$. c_{out} is calculated as
$(c_{\text{out,ini}}V_{\text{tot}} - \sum_{i=1}^{N}c_{\text{in}}V_i)/(V_{\text{tot}} - \sum_{i=1}^{N}V_i)$, where N is
the total number of nucleation sites. V_i is a condensate's
volume and V_{tot} is the total system volume, which we set

FIG. 2. Elastic ripening in a heterogeneous neo-Hookean elastic medium. (a) Visualization of simulation results based on the condensate growth model. The systems undergo ripening until only one condensate survives. Note that the positions of these condensates are generated randomly. (b) During the elastic ripening, the average radius, weighted by the condensates' volumes, exhibits a power law scaling with time, $\langle R \rangle \sim t^{1/5}$. The elastic pressure $p_{\text{el}} = E$ obeys a uniform random distribution in the range $[0, 2\bar{E}]$. At a later time, the elastic ripening crossovers to the Ostwald ripening in which $\langle R \rangle \sim t^{1/3}$. (c) The average local elastic pressure $\langle E \rangle$, weighted by the condensates' volumes, also exhibits a power law scaling with time, $\langle E \rangle \sim t^{-3/5}$. (d) Distributions of the normalized radii $R/\langle R \rangle$ from $t = 1000$ to $t = 4000$ with a fixed interval converge to a universal distribution, exhibiting scale invariance. The gray dashed line is the theoretical prediction (Supplementary Material). Inset: the raw distributions of R at different times. (e) Distributions of the normalized radii $E/\langle E \rangle$ from $t = 1000$ to $t = 4000$ with a fixed interval converge to a universal distribution, exhibiting scale invariance. Inset: the raw distributions of E at different times. In (a), (d), and (e), $\bar{E} = 1$. In all figures, the radii of nucleation sites $R_n = 1.5R_c$ where $R_c = 2\gamma/g_{\text{ini}}$. For the case of $\bar{E} = 0$, we add randomness to R_n so that it is uniformly distributed from R_c to $2R_c$ to avoid deterministic dynamics. In (b) to (e), the total number of nucleation sites is 5×10^5.

as $V_{\text{tot}} = 10^3 N$. Neither V_i nor V_{tot} include the nucleation site volumes, which are typically very small. We define $R_c \equiv 2\gamma/g_{\text{ini}}$, where g_{ini} is the initial condensing affinity so that R_c is the minimum nucleation radius for the condensate to grow at the beginning of the simulation in the absence of elastic pressure.

III. ELASTIC RIPENING

In the following, we use the neo-Hookean elasticity to model the elastic medium in which the elastic energy cost is proportional to the condensate's volume $F_{\text{el}} = EV$ where E is the local elastic pressure [24]. To mimic the heterogeneity of local stiffness in the nucleus, e.g., due to the spatial organization of euchromatin and heterochromatin, we assign each condensate a random E. In the case of Ostwald ripening, corresponding to the case of homogeneous E, small condensates shrink while big con-

densates grow as this ripening process reduces the overall surface energy. During the Ostwald ripening, the average and standard deviation of those survived condensates' radii exhibit power scaling with time, $\langle R \rangle \sim \sigma_R \sim (\gamma t)^{1/3}$ [13, 17, 18].

For the case of a heterogeneous elastic medium, we expect a similar ripening process in which condensates with large E shrink and condensates with small E grow since this can reduce the overall elastic energy. We simulate the condensate growth model and choose a uniform distribution of E from 0 to $2\bar{E}$ where $\bar{E}$ is the average (Appendix B). Indeed, the condensates with large E shrink during ripening (Figure S4). We note that nucleation sites with large E may not initiate condensate growth throughout the simulations due to their strong elastic pressures.

Surprisingly, the system undergoing elastic ripening exhibits a novel power-law scaling between the average condensate radius and time, which crossovers to the Ostwald ripening at a later time (Figure 2a, b and Movie

S1). Also, the standard deviation of the condensate radius exhibits a similar power-law scaling (Figure S5a). Moreover, the average and the standard deviation of the local elastic pressures also exhibit power-law scaling (Figure 2c and Figure S5b).To summarize,

$$\langle R \rangle \sim \sigma_R \sim At^{\alpha}, \tag{3}$$
$$\langle E \rangle \sim \sigma_E \sim Bt^{-\beta}, \tag{4}$$

Here, $\sigma_R^2 = \langle R^2 \rangle - \langle R \rangle^2$ and $\sigma_E^2 = \langle E^2 \rangle - \langle E \rangle^2$. The average variable $\langle ... \rangle$ is averaged over condensates weighted by their volumes, so the contribution of dissolved condensates is negligible. We also test the non-weighted average by excluding dissolved condensates explicitly and obtain similar results (Figure S6). α and β are the two power-law exponents. As we show later, A and B are the prefactors depending on the initial conditions.

In the following, we theoretically derive the two exponents. We use the following relation

$$\langle \dot{R} \rangle^2 \sim \langle \dot{E} \rangle \Rightarrow \beta = 1 - 2\alpha \tag{5}$$
$$\langle ER^3 \rangle \sim \sigma_{E,0} \Rightarrow \beta = 3\alpha \tag{6}$$

Eq. (5) comes from energy conservation: the dissipation power per unit volume $W = -\frac{\partial F}{\partial V}\dot{V}/V \sim \dot{V}^2/V^{\frac{4}{3}} \sim \dot{R}^2$ must be equal to the elastic energy changing rate per unit volume, which is $\dot{E}$. Eq. (6) comes from the scale invariance of the average elastic energy per condensate. $\sigma_{E,0}$ is the initial standard deviation of E with all condensates contributing equally (excluding condensates that never grow during the simulation). We remark that $\sigma_{E,0}$ is the only scale of the initial E distribution (shifting the E distribution by a constant does not affect the dynamics of condensate growth). Based on Eqs. (5, 6), we obtain that $\alpha = 1/5$ and $\beta = 3/5$. We can also obtain the expressions of the factor so that $A = \sigma_{E,0}^{1/5}$ and $B = \sigma_{E,0}^{2/5}$ using Eqs. (5, 6). Therefore,

$$\langle R \rangle \sim \sigma_R \sim \sigma_{E,0}^{1/5} t^{1/5}, \tag{7}$$
$$\langle E \rangle \sim \sigma_E \sim \sigma_{E,0}^{2/5} t^{-3/5}. \tag{8}$$

Our theoretical predictions are nicely confirmed for the average radius $\langle R \rangle$ (Figure 2b), the standard deviation of radius (Figure S5a), the average local elastic pressure $\langle E \rangle$ (Figure 2c), and the standard deviation of E (Figure S5b). We also verify the invariance of average condensate energy in Eq. (6) (Figure S7) and the expressions of the factors A and B (Figure S8). Our results do not depend on the distribution of E (Figure S9).

We have used the scale-invariance assumption in deriving Eq. (6), and the same power law scaling of $\langle R \rangle$ and σ_R supports this idea. To explicitly test the scale-invariance assumption, we plot the distributions of the normalized condensates' radii $R/\langle R \rangle$ at different times, and they indeed overlap, which means that the only length scale is the average radius $\langle R \rangle$ (Figure 2d). Notably, the distribution of $R/\langle R \rangle$ can be calculated semi-analytically, the gray dashed line in Figure 2d (Supplementary Material), although it depends on the distribution of E (Figure S10). Scale invariance is also observed for the distributions of $E/\langle E \rangle$ for the surviving condensates (Figure 2e). During the elastic ripening, the heterogeneity of local elastic pressures for the surviving condensates decreases while the heterogeneity of the radii for the surviving condensates increases. As a result, the Ostwald ripening eventually takes over the elastic ripening (Figure 2b).

FIG. 3. Elastic ripening is suppressed in an elastic medium beyond neo-Hookean elasticity. (a) Visualization of simulation results based on the condensate growth model. In this modified model, the elastic pressure increases with the condensate radius so that $p_{el} = E + YR$ where E obeys a uniform random distribution in the range $[0, 2\bar{E}]$ and Y is a constant. Here $Y = 0.1$. (b) The average radius $\langle R \rangle$ first exhibits the elastic ripening scaling and then reaches a plateau. In all panels, $\bar{E} = 1$ and the radii of nucleation sites $R_n = 1.5R_c$. In (b), the total number of nucleation sites is 5×10^5.

IV. BEYOND NEO-HOOKEAN

In a neo-Hookean medium, the elastic pressure is constant for each condensate, and the system undergoes ripening until only one condensate is left. It has been shown that a nonlinear elastic medium beyond neo-Hookean can suppress ripening [20, 21, 24]. To incorporate nonlinearity, we modify the elastic pressure so that

it increases with the condensate radius $p_{el} = E + YR$ where Y is a constant, in agreement with recent molecular dynamical simulations [21]. We find that the average radius first follows the 1/5 elastic ripening scaling and then saturates to a plateau R_{max} (Figure 3 and Movie S2). From Eq. (1), it is easy to find R_{max} at equilibrium, $R_{max} = \frac{g - \langle E \rangle + \sqrt{(g - \langle E \rangle)^2 - 8\gamma Y}}{2Y}$.

FIG. 4. Dynamics of condensate growth and dissolution and its mapping to transcriptional kinetics. (a) Condensates with weak local stiffness, i.e., small E, grow immediately after dissolution and can be associated with constitutively expressed genes. (b) Condensates with intermediate local stiffness grow intermittently with a delay since the last dissolution and can be associated with transcriptional bursting genes. (c) Condensates with strong local stiffness never grow, representing silenced genes that are not expressed. In (a) to (c), $R_n = 8R_c$. (d) Phase diagram of condensate growth dynamics as a function of R_n normalized by R_c and E. The blue, orange, and yellow regions represent condensates associated with constitutively expressed, transcriptional bursting, and silenced genes. The dark orange area represents the synchronized bursting condensates. In all panels, $\bar{E} = 1$, $k_{dis} = 10^{-5}$ and the number of nucleation sites $N = 2000$.

V. NONEQUILIBRIUM DYNAMICS OF ACTIVELY-DISSOLVING CONDENSATES

So far, we have analyzed the ripening dynamics of condensates as the system approaches thermal equilibrium, governed by the gradient of free energy. However, biological systems are often out of equilibrium due to energy-consuming processes. For example, transcriptional condensates can dissolve in the nucleus because of the ac-

cumulation of transcribed RNA [34], and this active dissolution process makes the system depart from thermal equilibrium. To investigate the effects of active dissolution on the dynamics of transcriptional condensates, we introduce an active dissolution rate to each condensate so that they have a constant rate to dissolve per unit volume k_{dis}, that is, the probability for condensate to dissolve within a short time window dt is $k_{dis}V dt$.

We first study the case of a heterogeneous neo-Hookean medium. Interestingly, we find that the growth dynamics of condensates can be categorized into three cases. Condensates with small E grow immediately after dissolution (Figure 4a); condensates with intermediate E grow after a delay since the last dissolution (Figure 4b); condensates with large E never grow (Figure 4c). Intriguingly, these three cases can be mapped to three different gene expression kinetics: constitutively expressed genes, transcriptional-bursting genes, and silenced genes (Figure 4d). Understanding the kinetics of transcription has been a central challenge in the study of stochastic gene expression. Multiple mechanisms have been proposed for the conditions of constitutive expression and transcriptional bursting, such as histone modifications and transcription factor availability [37–40]. Our results suggest that the local stiffness of chromatin provides a mechanical way to regulate transcription kinetics.

Furthermore, we observe a collective behavior of those bursting condensates: the initiation timings of a subset of those bursting condensates are synchronized. When the nucleation radius R_n is small, the complete set of bursting condensates are synchronized (Figure 4d, Figure 5a, and Movie S3). As R_n increases, a subset of the bursting condensates remain synchronized. Eventually, the initiation timings of all condensates become completely uncorrelated (Figure 5b, c and Movie S4). Intuitively, when the nucleation radius R_n is small, the condensate has to overcome a large Laplace pressure to initiate growth. Therefore, condensates must wait until other condensates dissolve so that the outside concentration c_{out} is high enough to overcome the Laplace pressure. While for large R_n, condensates can grow without the help of other condensates. We confirm that this phenomenon is generated by surface tension because the synchronized growth disappears in an artificial system without surface tension (Figure S11). This emerging synchronized dynamics of condensate growth suggests a mechanism of synchronized gene expression based on LLPS.

We define an order parameter f_{sync} to represent the fraction of synchronized condensates excluding condensates that never grow. When the fraction f_{sync} equals 1, all the condensates that can grow are synchronized. We obtain a heatmap of f_{sync} as a function of R_n and k_{dis} (Figure 5d). f_{sync} decreases gradually as R_n increases, indicating a continuous transition from fully synchronized to uncorrelated growth. Also, the f_{sync} decreases as the dissolution rate k_{dis} increases. This is because

FIG. 5. Synchronized growth of condensates in a neo-Hookean medium. (a) The temporal trajectories of condensate volumes V under a small R_n, in which the initiation timings of all condensates are fully synchronized (excluding condensates that never grow during the simulation). Here, $R_n = 1.2R_c$. (b) Here, $R_n = 4R_c$. In this case, a subset of the bursting condensates are synchronized (inset). (c) Here, $R_n = 10R_c$. In this case, the initiation timings of all condensates are completely uncorrelated. (d) The fraction of synchronized condensates as a function of R_n/R_c and $-\log_{10}(k_{dis})$. (e) The period T exhibits piecewise power-law scaling with k_{dis}. Inset: During one period of synchronization, the average volume over synchronized condensates $\langle V \rangle \sim t^{3/2}$ during the growth phase, then reaches the plateau phase, and finally undergoes ripening with $\langle V \rangle \sim t$. Here $k_{dis} = 10^{-7}$ and $\bar{E} = 0.01$. In (d), $R_n = 1.01R_c$ so that $f_{sync} = 1$ for the entire range of simulated k_{dis}. In (a) to (d), $\bar{E} = 1$. In all panels, the number of nucleation sites $N = 2000$.

condensates release molecules to the environment more frequently when k_{dis} is large; therefore, the outside concentration c_{out} is high so that the coupling between condensates is weakened.

For the fully synchronized case ($f_{sync} = 1$), we predict that the period T, which is the time interval between two successive synchronized growth, should exhibit piecewise power-law scalings with the dissolution rate k_{dis}. We estimate the period using $k_{dis}\langle V \rangle T \sim 1$ where $\langle V \rangle$ is the average volume without active dissolution. When the dissolution rate k_{dis} is large, the condensates dissolve during the initial growth phase. In this regime, $\langle V \rangle \sim t^{3/2}$ because $g - p_c$ is essentially constant [see Eq. (1) and the inset of Figure 5e]; therefore, $T \sim k_{dis}^{-0.4}$. As k_{dis} decreases, the condensates can grow larger before they dissolve, and the average condensate size can reach the plateau phase before ripening in which $\langle V \rangle \sim const$; therefore, $T \sim k_{dis}^{-1}$. As k_{dis} further decreases, the system should undergo ripening. If the ripening is driven by elastic ripening, $\langle V \rangle \sim t^{3/5}$, and the period $T \sim k_{dis}^{-0.625}$.

If the ripening is driven by Ostwald ripening, $V \sim t$, and the period $T \sim k_{dis}^{-0.5}$.

To test our predictions, we simulate the condensate growth model by choosing $R_n = 1.01R_c$ and a small $\bar{E}$ to ensure that $f_{sync} = 1$ in the entire range of simulated k_{dis}. Our predictions are nicely confirmed (Figure 5e). We do not see a clear signature of the scaling $T \sim k_{dis}^{-0.625}$ generated by elastic ripening in Figure 5e. We think that this is because the small range of E makes the regime of elastic ripening too short to observe (inset of Figure 5e). To test the existence of the scaling $T \sim k_{dis}^{-0.625}$, we increase the range of E and find a scaling behavior consistent with elastic ripening (Figure S12a). Our conclusions regarding the power-law scaling between T and the dissolution rate for the fully synchronized case are independent of the choice of R_n and E. Nevertheless, we note that the k_{dis}^{-1} and $k_{dis}^{-0.5}$ scalings occur only when $f_{sync} = 1$ (Figure S12b). This is because the plateau and ripening phases only apply to a closed system of synchronized condensates; that is, the outside concentration c_{out}

a

b

FIG. 6. Synchronized growth of condensates in a medium beyond neo-Hookean. (a) The fraction of synchronized condensates as a function of R_n given different Y's. Here $\bar{E} = 1$ and $k_{\text{dis}} = 10^{-5}$. (b) The synchronization period T exhibits piecewise power-law scaling with k_{dis}. Inset: during one synchronization period, the average volume over synchronized condensates $\langle V \rangle \sim t^{3/2}$ during the growth phase, then reaches the plateau phase without the ripening phase. Here $k_{\text{dis}} = 10^{-7}$ and $\bar{E} = 0.01$. $\langle V \rangle$ increases at the end of the synchronization period because of the increasing outside concentration due to the early dissolution of some condensates, which nevertheless does not affect the scaling relation. In (b), $Y = 0.1$, $R_n = 1.01 R_c$ so that $f_{\text{sync}} = 1$ for the entire range of simulated k_{dis}. In all panels, the number of nucleation sites $N = 2000$.

374 is set by synchronized condensates only.

VI. SYNCHRONIZED GROWTH BEYOND NEO-HOOKEAN

In the medium beyond neo-Hookean elasticity in which $p_{\text{el}} = E + YR$, we find that the phase space of bursting condensates becomes smaller under a finite Y (Figure S13). We also calculate the synchronized fraction f_{sync} under different Y's and find that f_{sync} decreases significantly when Y increases (Figure 6a). Because the increasing confining pressure limits the condensate growth, condensates tend to be smaller than the limiting case $Y = 0$, which means a higher outside concentration c_{out}. Therefore, it is easier for condensate to initiate growth without waiting for other condensates' dissolution. The synchronization period T also exhibits power-law scalings with the dissolution rate in the fully synchronized regime (Figure 6b). The main difference compared with the neo-Hookean model is that the $T \sim k_{\text{dis}}^{-0.5}$ scaling disappears since ripening is suppressed by nonlinear elasticity. Further, the $T \sim k_{\text{dis}}^{-1}$ scaling does not require the condition $f_{\text{sync}} = 1$ anymore since the condensate volume will reach a plateau value in any case due to the increasing elastic pressure $p_{\text{el}} = E + YR$ (Figure 3 and Figure S14).

VII. DISCUSSION

In this work, we have systematically investigated the dynamical behaviors of biomolecular condensates in a heterogeneous elastic medium. We introduce a mean-field-like model to investigate the condensate growth dynamics, allowing us to track millions of condensates simultaneously. Using this novel numerical method, we find a new dynamical scaling for the elastic ripening, $\langle R \rangle \sim t^{1/5}$, and theoretically derive the origin of the 1/5 exponent based on energy conservation and scale invariance. Furthermore, the heterogeneity of local elastic pressure decreases over time and exhibits power-law scaling, $\sigma_E \sim t^{-3/5}$. Ripening is suppressed in an elastic medium beyond neo-Hookean elasticity, and multiple condensates can coexist at equilibrium. Our theoretical predictions and numerical simulations nicely agree with each other.

We introduce a constant dissolution rate per unit volume to each condensate to model the dissolution of transcriptional condensates *in vivo* due to RNA accumulation. This active dissolution process drives the system out of equilibrium. Intriguingly, the temporal growth patterns of condensates resemble gene expression dynamics. Condensates in soft regions with weak local stiffness keep growing and continue to grow immediately after dissolution, corresponding to constitutively expressed genes. Meanwhile, condensates with stronger local stiffness initiate growth after a delay since the last dissolution, corresponding to transcriptional-bursting genes, which switch between active and inactive states and initiate transcription only in the active state. If the local stiffness is too strong, condensates can never grow, corresponding to silenced genes that are not expressed. Our results suggest that the local mechanical properties of chromatin play a key role in regulating gene expression kinetics, which can

be another layer of regulation on top of the compartmentalization of euchromatin and heterochromatin.

Notably, the timing of the growth initiation of bursting condensates can be synchronized. We remark that this is entirely a nonequilibrium effect due to the active dissolution process and finite surface tension. The fraction of synchronized condensates depends on the nucleation radius and dissolution rate. For the fully-synchronized cases, the period of synchronized growth exhibits piecewise power-law scalings with the dissolution rate. We theoretically derive the power-law exponents and show that it is related to the power-law scalings of average condensate sizes with time. In an elastic medium beyond neo-Hookean, synchronized growth is suppressed, and the results are qualitatively similar.

Some questions remain regarding the connection of our results to condensates in natural biological systems. In the experiments by Cho et al., stable and large condensates that virtually do not dissolve within the experimental time window coexist with smaller condensates that dissolve frequently [9]. This indicates that the nonequilibrium dissolution process of biomolecular condensates can be more complex than our simplified assumptions. The coalescence of condensates is also not included in our model. Recently, Lee et al. found a new dynamical scaling of ripening generated by the coalescence of subdiffusive condensates [22]. In the future, it will be interesting to explore the interference of elastic ripening, driven by the gradient of local stiffness and the subdiffusion of condensates themselves.

ACKNOWLEDGMENTS

We thank Fanlong Meng, Zhi Qi, and Yiyang Ye for useful discussions related to this work. The research was funded by National Key R&D Program of China (2021YFF1200500) and supported by grants from Peking-Tsinghua Center for Life Sciences.

L.M. conceived, designed, and carried out the theoretical and numerical part of this work. S.M. conceived and designed the theoretical part of this work. J.L. conceived, designed, and carried out the theoretical and numerical part of this work. All the authors contributed to the preparation of the manuscript.

APPENDIX A: DETAILED DERIVATION OF THE CONDENSATE GROWTH MODEL

We compute the condensing affinity g in the condensate growth model by considering a small change of the condensate volume V with its inside concentration fixed at c_{in}:

$$\begin{aligned} g &= -c_{in}\frac{\partial F}{\partial N} - \frac{\partial F}{\partial V} \\ &= c_{in}(\mu_{out} - \mu_{in}) + (p_{in} - p_{out}). \end{aligned} \tag{9}$$

Here, μ_{in} and μ_{out} are the chemical potentials of the condensate-forming molecules inside and outside the condensate. p_{in} and p_{out} are the pressures inside and outside the condensate. Next, we use the Gibbs-Duhem equation, $Nd\mu = -SdT + Vdp$ [41]. Therefore, $\mu_{in} = \mu_0 + (p_{in} - p_0)/c_{in}$ where μ_0 is the chemical potential of the condensate-forming molecules and p_0 is the pressure at equilibrium in the thermodynamic limit. Meanwhile, $\mu_{out} = \mu_0 + k_B T \ln(c_{out}/c_0)$ where c_0 is the saturation concentration at equilibrium in the thermodynamic limit. Furthermore, the outside pressure p_{out} has a correction due to the deviation of c_{out} from c_0, which we approximate as $p_{out} = p_0 + k_B T(c_{out} - c_0)$. Here, we have used the dilute assumption of the background solution for the calculation of μ_{out} and p_{out}. Coming all the above results, we obtain

$$g = c_{in}k_B T \ln\left(\frac{c_{out}}{c_0}\right) - k_B T(c_{out} - c_0). \tag{10}$$

APPENDIX B: DETAILS OF NUMERICAL SIMULATIONS AND DATA ANALYSIS

We perform numerical simulations by solving Eq. (1) using the explicit Euler method on MATLAB. In Figure 4d, a condensate is mapped to a constitutively expressed gene if over half of its growth can initiate within dimensionless time 100 after dissolution. Meanwhile, a condensate is mapped to a silenced gene if it never grows during the simulation. Finally, the rest condensates correspond to transcriptional-bursting genes.

In Figure 4d, Figure 5, and Figure 6, we label the condensate that can grow with the largest E as the first synchronized condensate. We then scan the condensates that can grow from the largest E to the smallest E, and the condensate is labeled as synchronized if over 80% of its initiation timings overlap with the growth timing of the latest synchronized condensate. After all the synchronized condensates are identified, we calculate the period by finding the average interval between the peaks of condensate growth timing distribution.

In Figure 5e and 6b, in the case of $\bar{E} = 0$, we add a small noise $\pm 0.001R_n$ to the nucleation radius to avoid deterministic dynamics. In the inset of Figure 5e, we exclude the $\langle V \rangle$ data when the survived condensates' number is smaller than 200 below which finite size effects are significant.

Movie S1: Condensates undergo elastic ripening in a heterogeneous neo-Hookean elastic medium until only one condensate exists. The simulation result is the same as Figure 2a in the main text.

Movie S2: Elastic ripening is suppressed in an elastic medium beyond neo-Hookean elasticity. The simulation result is the same as Figure 3a in the main text.

Movie S3: All actively-dissolving condensates are synchronized when R_n is small. Simulation parameters are the same as in Figure 5a in the main text.

Movie S4: All actively-dissolving condensates grow randomly when R_n is large. Simulation parameters are the same as in Figure 5c in the main text.

---

[1] C. P. Brangwynne, C. R. Eckmann, D. S. Courson, A. Rybarska, C. Hoege, J. Gharakhani, F. Jülicher, and A. A. Hyman, Germline p granules are liquid droplets that localize by controlled dissolution/condensation, Science **324**, 1729 (2009).

[2] A. A. Hyman, C. A. Weber, and F. Jülicher, Liquid-liquid phase separation in biology, Annual review of cell and developmental biology **30**, 39 (2014).

[3] S. F. Banani, H. O. Lee, A. A. Hyman, and M. K. Rosen, Biomolecular condensates: organizers of cellular biochemistry, Nature reviews Molecular cell biology **18**, 285 (2017).

[4] S. Alberti, A. Gladfelter, and T. Mittag, Considerations and challenges in studying liquid-liquid phase separation and biomolecular condensates, Cell **176**, 419 (2019).

[5] A. Klosin, F. Oltsch, T. Harmon, A. Honigmann, F. Jülicher, A. A. Hyman, and C. Zechner, Phase separation provides a mechanism to reduce noise in cells, Science **367**, 464 (2020).

[6] S. Alberti and A. A. Hyman, Biomolecular condensates at the nexus of cellular stress, protein aggregation disease and ageing, Nature Reviews Molecular Cell Biology **22**, 196 (2021).

[7] X. Weng, C. H. Bohrer, K. Bettridge, A. C. Lagda, C. Cagliero, D. J. Jin, and J. Xiao, Spatial organization of rna polymerase and its relationship with transcription in escherichia coli, Proceedings of the National Academy of Sciences **116**, 20115 (2019).

[8] A.-M. Ladouceur, B. S. Parmar, S. Biedzinski, J. Wall, S. G. Tope, D. Cohn, A. Kim, N. Soubry, R. Reyes-Lamothe, and S. C. Weber, Clusters of bacterial rna polymerase are biomolecular condensates that assemble through liquid–liquid phase separation, Proceedings of the National Academy of Sciences **117**, 18540 (2020).

[9] W.-K. Cho, J.-H. Spille, M. Hecht, C. Lee, C. Li, V. Grube, and I. I. Cisse, Mediator and rna polymerase ii clusters associate in transcription-dependent condensates, Science **361**, 412 (2018).

[10] B. R. Sabari, A. Dall'Agnese, A. Boija, I. A. Klein, E. L. Coffey, K. Shrinivas, B. J. Abraham, N. M. Hannett, A. V. Zamudio, J. C. Manteiga, *et al.*, Coactivator condensation at super-enhancers links phase separation and gene control, Science **361**, eaar3958 (2018).

[11] Y. E. Guo, J. C. Manteiga, J. E. Henninger, B. R. Sabari, A. Dall'Agnese, N. M. Hannett, J.-H. Spille, L. K. Afeyan, A. V. Zamudio, K. Shrinivas, *et al.*, Pol ii phosphorylation regulates a switch between transcriptional and splicing condensates, Nature **572**, 543 (2019).

[12] J. A. Morin, S. Wittmann, S. Choubey, A. Klosin, S. Golfier, A. A. Hyman, F. Jülicher, and S. W. Grill, Sequence-dependent surface condensation of a pioneer transcription factor on dna, Nature Physics **18**, 271 (2022).

[13] J. Berry, C. P. Brangwynne, and M. Haataja, Physical principles of intracellular organization via active and passive phase transitions, Reports on Progress in Physics **81**, 046601 (2018).

[14] C. P. Brangwynne, T. J. Mitchison, and A. A. Hyman, Active liquid-like behavior of nucleoli determines their size and shape in xenopus laevis oocytes, Proceedings of the National Academy of Sciences **108**, 4334 (2011).

[15] A. A. Hyman and C. P. Brangwynne, Beyond stereospecificity: liquids and mesoscale organization of cytoplasm, Developmental cell **21**, 14 (2011).

[16] A. Patel, H. O. Lee, L. Jawerth, S. Maharana, M. Jahnel, M. Y. Hein, S. Stoynov, J. Mahamid, S. Saha, T. M. Franzmann, *et al.*, A liquid-to-solid phase transition of the als protein fus accelerated by disease mutation, Cell **162**, 1066 (2015).

[17] I. M. Lifshitz and V. V. Slyozov, The kinetics of precipitation from supersaturated solid solutions, Journal of physics and chemistry of solids **19**, 35 (1961).

[18] M. Doi, *Soft matter physics* (Oxford University Press, 2013).

[19] Y. Shin, Y.-C. Chang, D. S. Lee, J. Berry, D. W. Sanders, P. Ronceray, N. S. Wingreen, M. Haataja, and C. P. Brangwynne, Liquid nuclear condensates mechanically sense and restructure the genome, Cell **175**, 1481 (2018).

[20] X. Wei, J. Zhou, Y. Wang, and F. Meng, Modeling elastically mediated liquid-liquid phase separation, Phys. Rev. Lett. **125**, 268001 (2020).

[21] Y. Zhang, D. S. W. Lee, Y. Meir, C. P. Brangwynne, and N. S. Wingreen, Mechanical frustration of phase separation in the cell nucleus by chromatin, Phys. Rev. Lett. **126**, 258102 (2021).

[22] D. S. Lee, N. S. Wingreen, and C. P. Brangwynne, Chromatin mechanics dictates subdiffusion and coarsening dynamics of embedded condensates, Nature Physics **17**, 531 (2021).

[23] E. Vidal-Henriquez and D. Zwicker, Cavitation controls droplet sizes in elastic media, Proceedings of the National Academy of Sciences **118**, e2102014118 (2021).

[24] P. Ronceray, S. Mao, A. Košmrlj, and M. P. Haataja, Liquid demixing in elastic networks: Cavitation, permeation, or size selection?, Europhysics Letters **137**, 67001 (2022).

[25] R. W. Style, T. Sai, N. Fanelli, M. Ijavi, K. Smith-Mannschott, Q. Xu, L. A. Wilen, and E. R. Dufresne, Liquid-liquid phase separation in an elastic network, Phys. Rev. X **8**, 011028 (2018).

[26] K. A. Rosowski, T. Sai, E. Vidal-Henriquez, D. Zwicker, R. W. Style, and E. R. Dufresne, Elastic ripening and inhibition of liquid–liquid phase separation, Nature Physics **16**, 422 (2020).

[27] D. Zwicker, R. Seyboldt, C. A. Weber, A. A. Hyman, and F. Jülicher, Growth and division of active droplets provides a model for protocells, Nature Physics **13**, 408 (2017).

[28] C. Weber, D. Zwicker, F. Jülicher, and C. Lee, Physics of active emulsions, Reports on Progress in Physics **82** (2019).

[29] D. Zwicker, The intertwined physics of active chemical reactions and phase separation, Current Opinion in Col-

loid & Interface Science , 101606 (2022).

[30] D. Zwicker, A. A. Hyman, and F. Jülicher, Suppression of ostwald ripening in active emulsions, Phys. Rev. E **92**, 012317 (2015).

[31] C. M. Caragine, S. C. Haley, and A. Zidovska, Nucleolar dynamics and interactions with nucleoplasm in living cells, eLife **8**, e47533 (2019).

[32] J. Söding, D. Zwicker, S. Sohrabi-Jahromi, M. Boehning, and J. Kirschbaum, Mechanisms for active regulation of biomolecular condensates, Trends in Cell Biology **30**, 4 (2020).

[33] D. S. Lee, C.-H. Choi, D. W. Sanders, L. Beckers, J. A. Riback, C. P. Brangwynne, and N. S. Wingreen, Size distributions of intracellular condensates reflect competition between coalescence and nucleation, Nature Physics , 1 (2023).

[34] J. E. Henninger, O. Oksuz, K. Shrinivas, I. Sagi, G. LeRoy, M. M. Zheng, J. O. Andrews, A. V. Zamudio, C. Lazaris, N. M. Hannett, *et al.*, Rna-mediated feedback control of transcriptional condensates, Cell **184**, 207 (2021).

[35] Y. Shin, J. Berry, N. Pannucci, M. P. Haataja, J. E. Toettcher, and C. P. Brangwynne, Spatiotemporal control of intracellular phase transitions using light-activated optodroplets, Cell **168**, 159 (2017).

[36] S. F. Shimobayashi, P. Ronceray, D. W. Sanders, M. P. Haataja, and C. P. Brangwynne, Nucleation landscape of biomolecular condensates, Nature **599**, 503 (2021).

[37] A. Sanchez and I. Golding, Genetic determinants and cellular constraints in noisy gene expression, Science **342**, 1188 (2013).

[38] D. Nicolas, N. E. Phillips, and F. Naef, What shapes eukaryotic transcriptional bursting?, Mol. BioSyst. **13**, 1280 (2017).

[39] A. J. Larsson, P. Johnsson, M. Hagemann-Jensen, L. Hartmanis, O. R. Faridani, B. Reinius, Å. Segerstolpe, C. M. Rivera, B. Ren, and R. Sandberg, Genomic encoding of transcriptional burst kinetics, Nature **565**, 251 (2019).

[40] E. Tunnacliffe and J. R. Chubb, What is a transcriptional burst?, Trends in Genetics **36**, 288 (2020).

[41] D. Kondepudi and I. Prigogine, *Modern thermodynamics: from heat engines to dissipative structures* (John Wiley & Sons, 2014).