Title:

Slow ramping emerges from spontaneous fluctuations in spiking neural networks

Authors: Jake Gavenas[1,2,5]*, Ueli Rutishauser[5,6,7,8], Aaron Schurger[1,3,9,10], Uri Maoz[1,2,3,4,8,11]

[1]Institute for Interdisciplinary Brain and Behavioral Sciences, Chapman University, Orange, CA, USA
[2]Schmid College of Science and Technology, Chapman University, Orange, CA, USA
[3]Crean College of Health and Behavioral Sciences, Chapman University, Orange, CA, USA
[4]Fowler School of Engineering, Chapman University, Orange, CA, USA
[5]Department of Neurosurgery, Cedars-Sinai Medical Center, Los Angeles, CA, USA
[6]Department of Neurology, Cedars-Sinai Medical Center, Los Angeles, CA, USA
[7]Center for Neural Science and Medicine, Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA, USA
[8]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA, USA
[9]INSERM U992, Cognitive Neuroimaging Unit, NeuroSpin Center, Gif sur Yvette 91191, France
[10]Commissariat à l'Energie Atomique, Direction des Sciences du Vivant, I2BM, NeuroSpin Center, Gif sur Yvette 91191, France
[11]Anderson School of Management, University of California, Los Angeles, CA, USA

*J.G. moved from Chapman University to Cedars-Sinai Medical Center during completion of this project.

Keywords:
voluntary action, spontaneous movement, volition, readiness potential, slow-ramping signal, medial frontal cortex, slow synapses, spiking neural network, clustered connectivity, spontaneous neural activity

Highlights:
1. We reveal a mechanism for slow-ramping signals before spontaneous voluntary movements.
2. Slow synapses stabilize spontaneous fluctuations in spiking neural network.
3. We validate model predictions in human frontal cortical single-neuron recordings.
4. The model recreates the readiness potential in an EEG proxy signal.
5. Neurons that ramp together had correlated activity before ramping onset.

Abstract
The capacity to initiate actions endogenously is critical for goal-directed behavior. Spontaneous voluntary actions are typically preceded by slow-ramping activity in medial frontal cortex that begins around two seconds before movement, which may reflect spontaneous fluctuations that influence action timing. However, the mechanisms by which these slow ramping signals emerge from single-neuron and network dynamics remain poorly understood. Here, we developed a spiking neural-network model that produces spontaneous slow ramping activity in single neurons and population activity with onsets ~2 seconds before threshold crossings. A key prediction of our model is that neurons that ramp together have correlated firing patterns before ramping onset. We confirmed this model-derived hypothesis in a dataset of human single neuron recordings from medial frontal cortex. Our results suggest that slow ramping signals reflect bounded spontaneous fluctuations that emerge from quasi-winner-take-all dynamics in clustered networks that are temporally stabilized by slow-acting synapses.

Introduction

Humans and other animals can initiate actions spontaneously, without immediate external triggers. Such spontaneous voluntary actions are critical to purposeful goal-directed behavior, but the neural mechanisms underlying these kinds of actions are poorly understood. Studies investigating the neural precursors of spontaneous voluntary actions have found that movement onset is preceded by neural signals that slowly ramp up or down more than 2 seconds before movement onset. This phenomenon is most evident in human medial frontal cortex and has been captured by scalp electroencephalography (EEG), where it is termed the readiness potential or "RP"[1–3], as well as by fMRI[4–6], and in single-neuron firing rates[7], where it is termed the readiness discharge. Other mammals also exhibit slow-ramping signals before spontaneous actions in analogous brain regions[8–10], and slow ramping has also been reported in invertebrates[11]. This suggests that slow ramping reflects an evolutionarily conserved mechanism related to voluntary action initiation. While widely reported, the origin and significance of slow ramping signals remain debated.

Slow-ramping signals were originally interpreted as reflecting preparatory processes that begin at the onset of ramping[2,12]. However, more recently, it has been proposed that slow ramping signals, like the RP, may instead be the result of autocorrelated fluctuations in neural activity that trigger movement upon crossing a threshold[13–17] or bias the precise time of movement onset[18]. In contrast to classic interpretations, these models (hereafter collectively referred to as "stochastic fluctuation models" or SFMs) ascribe no special meaning to the onset of the slow-ramping signals. Rather, SFMs posit that ramping signals appear only because of back-averaging autocorrelated signals aligned to movement onset. SFMs therefore offer drastically different neural and cognitive implications for action initiation than classic interpretations[19].

A major challenge for SFMs is that the time constant of empirically observed autocorrelations in neural spike trains is much shorter than the observed rate of ramping by about an order of magnitude. For example, slow ramping in the firing rate of human medial frontal cortex neurons evolves on the time scale of seconds[7], but studies have reported that neurons in medial frontal cortex have autocorrelation time constants on the order of hundreds of milliseconds[20–22] (although some have much longer time constants[23]). SFMs predict that more strongly autocorrelated fluctuations will approach the threshold more slowly and thus take longer to ramp up, while less autocorrelated (i.e., less temporally stable) processes are more likely to "jump up" and cross the threshold more abruptly, leading to shorter ramps[13]. It is thus unclear whether neurons that exhibit slow ramping are among the few that have longer autocorrelation time constants[23], or increase their autocorrelation temporarily during slow ramping through some unknown mechanism, or if slow ramping is a network phenomenon that does not rely on the autocorrelation of single units.

Here, we utilized a combination of spiking neural network simulations and analysis of human single-neuron experimental data[7] to investigate (1) the mechanisms by which temporally autocorrelated fluctuations in neural activity emerge in spiking neural networks, and (2) whether such fluctuations can account for the kind of slow ramping activity before spontaneous voluntary actions seen in the empirical data. We show that spiking neural networks can be configured to produce spontaneous fluctuations and slow-ramping activity with properties similar to those recorded[7] from human medial frontal cortex during a spontaneous voluntary movement task

similar to the classic Libet task [2]. Our model is a spiking neural-network with 400 units with
clustered connectivity. Such networks can exhibit spontaneous fluctuations[24,25], but prior studies
found that neurons in these networks exhibit "abrupt switching" activity—i.e. abrupt alternations
between slow and fast firing—which is unlike the gradual changes seen during slow ramping.
We hypothesized that we could stabilize activity by inserting slow synapses, which are critical
for producing temporally stable activity[26,27] and increased autocorrelation in simulated spiking
neurons[28]. We show that the presence of slow synapses led to more gradual spontaneous
fluctuations and facilitated the emergence of slow ramping. The properties of ramping resembled
the ramping signals observed in experimental data both at the single-neuron and population level.
Furthermore, our model predicted that pairs of neurons that both ramped up or down would have
more correlated activity compared to pairs that ramped in different directions even before the
onset of ramping. This prediction was validated in the experimental data. We conclude that slow
ramping signals before spontaneous, voluntary action likely emerge from backward-averaged
spontaneous fluctuations aligned to threshold-crossings, and that clustered connectivity and slow
synapses interact to jointly facilitate temporally smooth fluctuations. These results are
compatible with SFMs of spontaneous voluntary action generation.
111
Results
Slow synaptic transmission temporally stabilizes spontaneous fluctuations in simulated
networks
We simulated networks consisting of N=400 spiking leaky integrate-and-fire neurons (see LIF
Network Simulations in Methods); 80% of the neurons were excitatory, i.e., they provided only
positive inputs to other neurons, while the remainder were inhibitory. Excitatory neurons were
grouped into 4 clusters of 80 neurons each, such that within-cluster connections were more likely
than between-cluster connections. Simulations of a range of network sizes (N=200, 300, 400,
500 neurons) and number of clusters ($N_{clusters}$ = 3, 4, 5, 8) led to the same qualitative results,
described below, suggesting that our main results do not strictly depend on our chosen network
size and number of clusters. The extent of clustering was quantified by the parameter R_{EE}
(clustering degree), which is equal to the ratio of the probability of a within-cluster connection to
the probability of a between-cluster connection (within-cluster connections were also stronger
compared to between-cluster connections, as in[24]; see Connectivity Matrix Construction in
Methods). Inhibitory neurons were not clustered, and their connectivity probability with
excitatory neurons did not depend on R_{EE}[29].
128
R_{EE} describes a continuum between homogenous networks without clustering ($R_{EE} = 1$) and
winner-take-all networks with strong clustering ($R_{EE} \gg 1$) (Fig. 1A). Homogenous networks
with balanced excitation and inhibition exhibit asynchronous spiking activity without slow
spontaneous fluctuations in firing rates[30]. Conversely, in winner-take-all networks, neurons in a
single cluster dominate and suppress neurons in other clusters via lateral inhibition implemented
by inhibitory neurons[31]. In between these two extremes, clustered networks (roughly $2 < R_{EE} < 5$,
depending on network size, number of clusters, and other factors) exhibit slow spontaneous
fluctuations due to a combination of lateral inhibition and lateral excitation between clusters.
Note that in our networks, spontaneous fluctuations were entirely due to internal interactions
because external input to the network was fixed (besides noise).
139

140
Figure 1. Simulation overview and example network activity. A: As the clustering degree (R_{EE})
increases, connectivity transitions from homogenous/balanced, through clustered connectivity, to winner-
take-all. **B:** Schematic of fast vs. slow synapses. **C:** Example 5-second simulations for networks with
100% fast synapses (top row) and 50% slow synapses (bottom row) at various clustering degrees (same
underlying network structure, leading to the same 'winning' cluster, depicted in orange). For each
subpanel, at the top are single-trial raster plots and at the bottom are cluster-averaged firing rates over
time (spikes smoothed with 400 ms Gaussian kernel then averaged across all neurons in each cluster in
different colors). In raster plots, purple are excitatory neurons (4 clusters of 80 neurons), green are
inhibitory neurons. Clustering leads to spontaneous fluctuations, but fluctuation characteristics depend on
synaptic dynamics. Networks with 100% fast synapses (top) switch between high- and low-activity states,
whereas networks with 50% slow synapses (bottom) show smoother, more temporally stable fluctuations.
152
Previous work had only simulated clustered networks with fast synaptic dynamics, in which the
entire postsynaptic potential is delivered within ~5 ms[24,25,32]. We were interested in the role of
slow synapses in modulating spontaneous fluctuations in such networks. We therefore simulated
networks with both fast and slow synapses. We used double-exponential functions to model
synapses[24,33,34], with fast synapses having rising and falling time-constants of 2 and 5 ms,
respectively, and slow synapses having rising and falling time-constants of 20 and 100 ms
respectively[33] (Fig. 1B). Because different types of synapses engage either fast or slow dynamics
(AMPA and GABA-A for fast, NMDA and GABA-B for slow[34]), we modeled mixtures of
synapses as a weighted sum of these two types of synapses, parameterized as the ratio of fast to
slow synaptic signaling (i.e., the percent of post-synaptic potential (PSP) delivered via fast
versus slow synapses; note that this applied to both excitatory and inhibitory connections; see
LIF Network Simulations in Methods).

165
Recreating prior results for clustered networks with 100% fast synapses, we found that
increasing R_{EE} was associated with a transition from asynchronous activity to more winner-take-
all dynamics, with networks in an intermediate range exhibiting slow fluctuations (Fig. 1C top
row). Notably, fluctuations in each cluster manifested as bistable switching activity, with entire
clusters switching between states of low and high activity[24,25]. These fluctuations exhibited
quasi-winner-take-all dynamics: activations in one cluster were accompanied by decreased firing
in all the other clusters. Notably, switches between these states were rapid with no apparent slow
ramping (Fig. 1C, top row; see Fig. 2 for quantification).
174
In contrast, networks with slow synapses exhibited more gradual fluctuations. With 50% slow
synapses, fluctuations in networks with intermediate R_{EE} values were temporally stable, with
rises and falls that occur over a longer timespan when compared to networks with 100% fast
synapses (Fig. 1C bottom row; see below for quantification). Fluctuations in 50% slow-synapse
networks still exhibited quasi-winner-take-all dynamics, with activations in one cluster
suppressing other clusters, but without the all-or-none nature of fluctuations common in 100%
fast-synapse networks. Note that fluctuations are less clear in the inhibitory pool (Fig. 1C green
raster); henceforth we exclusively analyze properties of the excitatory neurons.
183
184 **Network architecture and synaptic dynamics interact to shape network activity**
185
Next, we quantified the effects of clustering degree and slow synaptic transmission on
spontaneous fluctuations. We simulated 5 seconds of activity in 50 different randomly generated
networks (each with 400 neurons, 320 excitatory) for R_{EE} values between 1 and 6 (increment of
0.1), with synaptic ratios ranging from 0% slow (100% fast) to 50% slow (increment of 10%,
overall N=15,300 networks). We then investigated the effects of R_{EE} and synaptic ratio on
measures of the networks' temporal statistics, starting with power spectra. For each network, we
calculated firing rate as a function of time for each neuron by smoothing spike trains with a
gaussian filter (50 ms kernel), and then calculated the power spectra from the resulting smoothed
firing rates (see Simulating & Analyzing Fluctuations in Methods). Next, we averaged power at
matched frequencies across neurons within a network, and then averaged across networks within
bins of increasing R_{EE} values (e.g. between 1.0 and 2.0; Fig. 2A left panel). Spectral power was
generally higher at low frequencies and, on average, dropped monotonically as a function of
frequency. As clustering increased past a critical value, spectral power increased for all
networks. Notably, for low clustering coefficients (R_{EE} approximately between 1 and 3), the ratio
of fast to slow synapses had no or a low impact on power. However, as clustering reached a
critical range (R_{EE} above ~3.0), a larger percentage of slow synapses resulted in higher power at
relatively low (below ~1 Hz) and high (above ~5 Hz) frequencies. At even higher clustering (R_{EE}
above ~5.0), power decreased again, likely signifying the transition from the "fluctuation"
regime to the "winner take all" regime.
205

Figure 2. **Effects of clustering & synaptic dynamics on temporal dynamics of neuronal activity. A:** Effect of clustering and synaptic dynamics power spectra of simulated neurons' firing rates (plotted in log-log scale). **B:** Effect of clustering and synaptic dynamics on spike train autocorrelation at time lags up to 1000 ms. **C-F:** Effect of clustering and synaptic dynamics on four measures of temporal dynamics. **C:** Spectral Slope, the slope of a linear regression of power to frequency in log-log scale (panel A **D:** Autocorrelation timescales, the decay parameter of exponential fit to autocorrelation functions (panel B). **E:** Coefficient of variation (CV), the firing rate standard deviation across time (100 ms non-overlapping bins) divided by trial-average firing rate. **F:** $\hat{S}_T$, a measure of firing rate variability across time like CV. $\hat{S}_T$ was designed to specifically index slow-switching behavior like that seen in 100% fast / 0% slow synapse networks[25].

We next investigated autocorrelation directly, employing the same grouping strategy as above (Fig. 2B; see Simulating & Analyzing Fluctuations in Methods). Unlike power, which did not vary as a function of synaptic dynamics for low clustering values, autocorrelation at short time lags was slightly higher for networks with a higher ratio of slow synapses. Then, at higher clustering values (R_{EE} above ~3.0, roughly commensurate with results from power spectra), higher ratios of slow synapses led to noticeably greater autocorrelation at longer time-lags. As clustering increased further, autocorrelation at longer time-lags decreased and lag-one autocorrelation (corresponding to one 50 ms bin) became negative.

To further quantify the slow fluctuations we observed, we calculated four measures of spiking variability. First, we calculated the spectral slope (equal to the negative 1/f exponent) of neural firing rate power spectra as the slope of a linear fit of frequency to power in log-log space[13,35]

(Fig. 2C.; power spectra calculated as above; see Simulating & Analyzing Fluctuations in
Methods). For low clustering (R_{EE} below ~3.0), spectral slope did not depend on clustering
degree and only little on the ratio of slow synapses, with higher ratios of slow synapses having
slightly lower slope. At higher clustering degree, the dynamics changed, with spectral slope
generally increasing with clustering, with larger increases for networks that had higher ratios of
slow synapses. The higher slopes for networks with greater frequency of slow synapses indicates
that power increases due to greater clustering were relatively larger at higher frequencies for
higher ratios of slow synapses. Next, we calculated the autocorrelation timescale τ[20,21,36] of
neural firing rates as the decay parameter of an exponential fit to the autocorrelation function
(Fig. 2D). Timescales were generally higher for networks with greater ratios of slow synapses for
all clustering values. As clustering increased, autocorrelation timescales increased for all
frequencies of slow synapses (until R_{EE}~3-4) and then dropped off. The drop-off was more
gradual for networks with high percentages of slow synapses, indicating that slow synapses may
impact the transition point from fluctuations to winner-take-all dynamics.
243
Next, we investigated how the coefficient of variation (CV) of individual neurons' firing rates
across time was affected by the clustering (Fig. 2E). The CV equals the standard deviation of
firing rates across time (average firing rates in 100 ms non-overlapping bins; see Simulating &
Analyzing Fluctuations in Methods), normalized by dividing by the trial-average firing rate.
High CV values are expected for networks that exhibit slow fluctuations in firing rates, as those
fluctuations entail higher firing rate standard deviations across time. CV sharply increased after a
critical R_{EE} value (~3-3.5) for all networks, suggesting the emergence of slow fluctuations past
that point. CV then decreased for higher R_{EE} values, indicating the transition to a winner-take-all
regime. Notably, a greater percentage of slow synapses consistently resulted in a higher CV
(though the rate of increase diminished with the proportion of slow synapses), suggesting that
slow synapses increase firing rate variability even without clustering. We next looked at $\hat{S}_T$, a
measure of spiking variability[25] that characterizes slow-switching behavior in clustered
networks, such as those seen in 100% fast synapse networks, Fig 1C top row). We investigated
how $\hat{S}_T$ was affected by clustering and slow synapses (Fig. 2F). $\hat{S}_T$ is the standard deviation of
cluster-averaged firing rates (100 ms non-overlapping bins) across time, averaged across
clusters, and normalized by subtracting the same measure applied to random groupings of
neurons rather than by cluster identity. Higher values of $\hat{S}_T$ therefore increased temporal
variability and slow-switching activity. $\hat{S}_T$ increased steadily as R_{EE} increased, reaching maximal
values for R_{EE} between 3 and 4, and then decreased. Notably, a greater percentage of slow
synapses resulted in lower $\hat{S}_T$ values. These results indicate that networks with slow synapses
exhibit slow fluctuations, but that these fluctuations manifest as gradual fluctuations rather than
slow-switching activity. Notably, however, the changes in temporal statistics we observed were
heterogeneous across networks with the same R_{EE} and synaptic ratio parameters (especially so in
the fluctuation regime; see Fig. S1), and likely depend on the emergence of specific connectivity
patterns as clustering increases. Taken together, clustering is sufficient to lead to spontaneous
slow fluctuations in spiking networks, yet our results suggest that introducing slow synaptic
dynamics can greatly alter the nature of said fluctuations, in this case leading them to be
temporally stable. We next turned to quantifying the speed of ramping in this network
architecture.
273
274

Slow-ramping at the single-neuron level
We next examined the relation between spontaneous fluctuations and slow-ramping signals at the single-neuron level. We simulated activity in 20 randomly connected networks with intermediate clustering degrees that led to spontaneous fluctuations (Min R_{EE} = 2.8; Max R_{EE} = 4; manually tuning for each network so that dynamics were in the fluctuation regime rather than winner-take-all regime; see Network Selection in Methods). For each network, we simulated 100 trials of 10 s each, when synaptic dynamics were either 100% fast or 50% slow. We then aligned data to times when activity in a specific cluster increased their activity above a threshold (termed threshold-crossing; the threshold was defined as 50% of maximum normalized firing rate across the whole trial; see Threshold Alignment in Methods). Neurons that significantly increased or decreased their firing rates in the 400 ms before threshold-crossing relative to a baseline (-3 to -2 s; as in Fried et al.[7]) were categorized as increasing or decreasing, respectively (Wilcoxon rank-sum test, the cutoff was p<0.01, as in Fried et al.[7]). The ratio of neurons categorized as increasing to decreasing was, on average, 50:50 in 100% fast synapse networks and 39:61 in 50% slow synapse networks.

Networks with 50% slow synapses demonstrated slow ramping with increasing and decreasing of firing rates before threshold-crossing, whereas networks with fast synapses showed only a steep jump shortly before threshold-crossing (Fig. 3A). Notably, our networks did not have a specified 'ramping onset,' nor were there any changes to the network's input after trial onset (which was many seconds before the spontaneous ramping onset) beyond those attributable to noise. Therefore, the observed ramping must reflect the spontaneous fluctuations prior to threshold-crossing rather than a buildup driven by a specific intervening event. To further explore the relationship between slow synapses and ramping behavior, we conducted simulations (same as above) for various ratios of fast-to-slow synapses. Those simulations revealed that slow ramping emerged gradually as we increased the proportion of slow synaptic signaling (Fig. 3B). This gradual emergence closely matched the similar gradual emergence of slow ramping with increased autocorrelation (Fig. S2A). Accordingly, we found that increasing the ratio of slow synapses gradually increased activity autocorrelation during the ramping period at both the cluster and single-neuron level, although cluster-level autocorrelation was much higher than single-neuron autocorrelation (Fig. S2B). Together, these simulations support the notion that slow synapses facilitate temporally stable autocorrelated activity and, therefore, slow ramping.

Figure 3: Slow ramping at the single-neuron level in simulated networks and recorded human frontal neurons. A: Firing rates (Δ, relative to baseline; mean ± 95% confidence intervals across networks) of neurons that increased (Inc.; red) or decreased (Dec.; blue) activity before threshold-crossings in networks with 100% fast (dotted lines) or 50% slow synapses (solid lines). **B:** Slow ramping of firing rates, either increasing (left) or decreasing (right), gradually emerges with the growing proportion of slow synapses. **C:** Left: Slow ramping before movement in human MFC neurons (Fried et al.[7]; Δ, relative to baseline; mean ± 95% confidence intervals across neurons). Some neurons show a steplike response profile (solid lines), and others show slow ramping (dashed lines). Right: Goodness-of-fit of simulated data versus experimental slow ramping response profiles (points: individual networks, error bars: 95% confidence intervals). For each network simulated with 0% and 50% slow synapses, we regressed average pre-threshold activity on the average empirical pre-movement activity at equivalent time-points (separately for increasing and decreasing neurons). From those models we extracted the R^2 as a measure of how well each network fit pre-threshold changes in firing rates. 50% slow networks had significantly better fits compared to 100% fast networks for both increasing and decreasing neurons (p < 0.001 both increasing and decreasing). **D:** Slow ramping emerges despite low autocorrelation at the single-neuron level in both simulated (left; mean) and empirical neurons that exhibit slow ramping (right; mean ± 95% confidence interval). **E:** Example raster plots and trial-averaged firing rates from simulated (left) and empirical neurons (right). Notably, slow ramping is not apparent in most single-trial responses.

Comparison of single-neuron slow ramping in simulated and empirical data

We next assessed whether the degree of slow ramping observed in the model is comparable to that typically seen in the motor system preceding the onset of self-initiated actions. One of the key brain areas within which such slow ramping has been observed is in medial frontal cortex (MFC) of humans, including the pre-supplementary motor area, supplementary motor area proper, and anterior cingulate cortex[3,37,38]. We examined whether the properties of empirically measured ramping are comparable to those seen in our model by re-analyzing human MFC

335 single-neuron data recorded by Fried and colleagues[7] that show slow ramping and other firing
336 rate changes before spontaneous voluntary actions.
337
We analyzed n=512 MFC neurons. Of these, 153 (30%) neurons significantly changed their
response relative to movement onset (using the same selection criteria we used for our model
data). Of these 153 neurons, 51 had increasing, and 102 had decreasing firing rates before
movement onset relative to baseline. We next sub-selected neurons that exhibited slow ramping
rather than more abrupt, step-like changes in firing rates. We distinguished between these two
response profiles by fitting a sigmoid to trial-averaged firing rates and extracting an inverse-gain
parameter. Neurons with an inverse-gain >0.4 were categorized as slow ramping, otherwise as
steplike (see Neuron Categorization and Response Profiles in Methods). Thus, 41 MFC neurons
were slow-ramping (27% of MFC neurons that significantly changed their firing rate before
movement; 15 increasing and 26 decreasing) and the remaining 112 were steplike (73%; 36
increasing and 76 decreasing). Notably, slow ramping neurons showed progressive changes to
their firing rates beginning around 2 s before movement onset on average, while steplike neurons
showed a sharper firing rate change around 1 second before movement (Fig. 3C left; see Fig.
S2C for distribution of ramping onset times). Here we focus on slow-ramping neurons.
352
Comparing the response profiles of simulated and recorded MFC neurons revealed that neurons
in the slow ramping group had response profiles that were like those in simulated networks with
slow synapses (compare Fig. 3A vs 3C). To quantify the goodness-of-fit of our networks, we
regressed pre-threshold activity (averaged across neurons; -3 to 0 seconds rel. threshold-
crossing) in simulated networks on the empirical data from matching time points (similarly
averaged across neurons; Fig. 3C right). Networks with 50% slow synapses had higher R^2 values
than those with 100% fast synapses when comparing increasing neurons (mean [95% confidence
interval]: 50% slow: 0.73 [0.71, 0.76] vs 100% fast: 0.24 [0.22, 0.27]; $p < 0.001$ two-sided sign-
rank test) and decreasing neurons (50% slow: 0.66 [0.62, 0.71] vs 100% fast: 0.36 [0.31, 0.41]; p
< 0.001 two-sided sign-rank test). Notably, slow ramping over ~2 s at the single-neuron level
occurred even though autocorrelation in these neurons decayed to zero after ~600 ms, both for
simulated and empirical neurons (Fig. 3D). Therefore, our model reproduces the key apparent
paradox that we sought to resolve. Furthermore, visual inspection of raster plots from neurons
that engaged in slow ramping behaviors showed that smooth, largely monotonic ramping was not
visible on every individual trial—rather, activity was highly heterogeneous across trials (Fig.
3E).
369
Slow ramping at the population-level
We next investigated slow ramping at the population level of all recorded/simulated neurons. We
trained a linear discriminant analysis classifier (LDA) to differentiate activity at different points
during the ramping period from a baseline (-3.0 to -2.6 s). Classification accuracy on statistically
independent data, not used for training (5-fold cross-validation) for simulated networks with
50% slow synapses reached 75% at 641±7 ms before threshold-crossing (mean ± standard error
here and below, averaged across networks), while classification accuracy for simulated networks
with 100% fast synapses only increased above 75% on average 31±30 ms before threshold-
crossing (Fig. 4A shows results from 100 simulations on a single network; further examples are
in Fig. S4A). Classification accuracy in the simulated networks with 50% slow synapses
resembled the gradual rise in accuracy obtained by an analogous analysis of the Fried et al.

(2011) dataset (Fig. 4A right), in which we pooled MFC neurons across participants to create a
pseudo-population for decoding[39]. Decoding accuracy reached 75% at ~1.2 s before movement
onset, and reached ~90% at movement onset, presumably because many empirical neurons show
a sudden 'step like' change in their firing rates at around 0.5s before movement onset (see Fig.
3C). Our model does not contain such neurons. These results suggest that our simulations best
capture the early, gradually increasing accuracy in decoding activity from baseline up to ~0.5 s
before movement onset (Fig. 4A, left vs. right plot).
388

389
Figure 4: Slow ramping at the population level in simulated and empirical data. A: Classification accuracy
versus baseline (linear discriminant analysis; spike count in 400 ms windows, times depicted refer to the leading
edge of that window, versus -3 to -2.6 s baseline window) in simulated (left) and empirical data (right; shaded region
is 95% confidence interval across cross-validations; dashed line is chance level across 100 shuffles of the data).
Accuracy slowly ramps before threshold-crossing / movement, suggesting slow ramping at the population level. **B:**
Performance of LDA classifiers trained at one time bin and tested on another time bin (relative to threshold-crossing
or movement). Left: simulated data. Right: empirical data. Accuracy gradually increased leading up to the time of
threshold-crossing or movement and was higher when the training bin was earlier than the testing bin compared to
vice-versa (above versus below the main diagonal). **C:** Top: EEG proxy (PSPs onto excitatory populations) shows
slow ramping in 50% slow but not 100% fast networks (confidence intervals over networks). Bottom: slow ramping
in EEG proxy signal gradually emerges with an increased percentage of slow synaptic signaling.

401
To test whether gradually increasing accuracy reflected ramping at the population level, we
investigated the temporal generalization of decoders, which can be used to ascertain the temporal
evolution of neural activity[40]. We classified activity at different time points against a baseline (-3
to -2.6 s). If the population as a whole is exhibiting ramping activity, then earlier time points will
be closer to the baseline in neuronal state space, while later time points will be farther from
baseline. Therefore, when training a classifier on activity that is temporally close to baseline
(early), the classifier will find a separatrix that is closer to the baseline in neural state space.
Then, when applied to activity that is temporally further from baseline (later), the same
separatrix will perform well because that activity is farther from baseline activity in state space

(and vice-versa for training on later then generalizing to earlier activity). Thus, we should
observe an asymmetry in across-time generalization accuracy such that classifiers perform better
when generalizing from earlier to later timepoints compared to the reverse.
414
To test this theoretical prediction, we trained an LDA classifier to differentiate between ramping
activity at a given time point relative to threshold crossing and activity during a baseline period
(and tested that classifier's accuracy at different time points (see Decoding Analyses in
Methods). For simulated data, decoding accuracy gradually increased leading up to threshold-
crossing (Fig. 4B left; more examples in Fig. S4B). Notably, as expected for ramping signals,
across-time generalization accuracy was asymmetric—higher when training on earlier and
testing on later timepoints compared to when training on later and testing on earlier timepoints
(Fig. 4B). This can be seen by comparing the average values in the upper vs. lower triangle of
the temporal generalization matrix (one-tailed rank-sum test $p < 0.001$). Similarly, for the MFC
neurons, decoding accuracy showed a similar increase and asymmetry (rank-sum test comparing
upper triangle to lower triangle decoding accuracies, $p < 0.001$, Fig. 4B right; see Fig. S4C for
explicit demonstration of this asymmetry). To verify that this asymmetry reflects ramping, we
generated spike trains that slowly ramped up and down to create a pseudo-population of neurons
that we know a priori exhibit linear ramping (n=20, 50 trials, see Simulated Ramping in
Methods). We conducted the same decoding and temporal generalization procedure on those
spike trains and found the same asymmetry that decoders generalized to future time points better
than to past time (one-tailed rank-sum test $p < 0.001$; Fig. S4D).
432
We also investigated ramping at the population level by calculating an EEG proxy from our
network to compare to the readiness potential, a negative deflection in trial-averaged scalp EEG
that slowly builds up over the last ~1-2 s prior to movement onset[3,19]. Notably, we focus only on
the early (negative) rise of the readiness potential, which is thought to originate in bilateral SMA,
unlike the later parts, closer to movement onset, which originate in lateral motor cortex[3]. The
EEG signal is typically taken to reflect the summation of post-synaptic potentials onto neurons[41].
So, as a proxy signal, we calculated the total signed post-synaptic input (both inhibitory and
excitatory) delivered to each excitatory neuron in the network, then averaged across neurons, and
investigated its trajectory leading up to threshold-crossings (see Threshold Alignment in
Methods). Networks with 50% slow synapses exhibited slow negative ramping in the EEG
proxy. In contrast, networks with 100% fast synapses had a relatively flat signal with an abrupt
jump close to threshold-crossing (Fig. 4C top). Repeating this analysis for multiple ratios of fast-
to-slow synaptic signaling revealed that slow ramping in the EEG proxy signal gradually
emerged with a greater percentage of slow synapses (Fig 3C bottom). The results were similar
for an EEG proxy that uses PSPs onto all simulated neurons rather than just excitatory neurons
(Fig. S4E), accounting for suggestions that the human cortex contains larger proportions of
inhibitory neurons than other species[42].
450
Pre-ramping noise correlations relate to slow-ramping activity
Our model, like other SFMs, predicts that slow ramping is a consequence of averaging
autocorrelated fluctuations aligned to times of threshold-crossings. Those fluctuations do not
begin at what would appear as ramping onset on average but occur throughout the entire trial
(see Fig. 1C bottom). Therefore, a novel prediction of our model is that two neurons that show
ramping behavior in the same direction (both increasing or decreasing) will have more correlated

activity *before ramping onset* in comparison to pairs that show ramping behavior in different
directions (one increasing and one decreasing). Such correlations in trial-by-trial firing rate are
called noise correlations[43,44], and are relevant to behavior[45,46]. Pre-ramping noise correlations that
depend on the later type of ramping should emerge because ramping behavior is related to cluster
membership—two neurons that both increase their firing rates are likely part of the same
'winning' cluster, and two neurons that both decrease their firing rates may be from any of the
three 'losing' clusters, whereas two neurons that show opposite ramping directions almost
certainly belong to different clusters (Fig. 5A).
465

466
Figure 5: Model schematic and correlation structure relates to ramping direction. **A:** Stochastic fluctuation model
(SFM) schematic. In SFMs, ramping reflects autocorrelated fluctuations prior to a threshold-crossing. Because the
fluctuations occur throughout a trial, neurons with the same ramping behavior should exhibit correlated activity even
before ramping onset. **B:** In 50% slow synapse networks, pairs of simulated neurons that exhibited the same ramping
behavior—both increasing (Inc-Inc) or both decreasing (Dec-Dec)—were significantly correlated even before
ramping onset (-3 to -2.5 s). Furthermore, neurons that showed opposite ramping behaviors (Inc-Dec) were
anticorrelated before ramping. **C:** Pairs of empirical neurons that showed the same ramping behavior (increasing or
decreasing) were significantly correlated before ramping onset (-3.5 to -2.5 s; mean & 95% confidence interval of
pairwise correlation calculated via a linear mixed-effects model; steplike responses were categorized as "other" for
this analysis). Pairs of neurons that exhibited the same type of ramping (both increasing or decreasing) were more
correlated in the pre-ramping period compared to pairs that exhibited opposing types of ramping (increasing vs.
decreasing). Although pairs showing opposite ramping behaviors were not reliably anticorrelated, as they were in
the simulations, they were not statistically different from 0 in the positive or negative direction.
480
As predicted, in 50% slow synapse networks, activity in pairs of increasing neurons (i.e., both
belonging to the specific cluster associated with movement) was positively correlated during a
pre-ramping baseline period (-3 to -2.5 s, Fig. 5B, r = 0.277, p < 0.001, LME; see Analysis of
Pairwise Correlations in Methods). In addition, activity in pairs of decreasing neurons (i.e., both
belonging to clusters other than the one whose threshold-crossing determined movement timing)
was also positively correlated, though to a lesser extent than pairs of increasing neurons, because
they may belong to different clusters (r = 0.059, p < 0.001, LME). In contrast, activity in pairs
involving one increasing and one decreasing neuron (i.e., very likely belonging to different
clusters) was negatively correlated (r = -0.115, p < 0.001, LME). Finally, some neurons neither
increased nor decreased their activity before threshold-crossing, likely due to the random nature
of network connectivity leading to them receiving heterogeneous inputs. Activity in pairs of such
neurons had very low but significantly positive correlations before ramping onset (r = 0.003, p <
0.001).
494

We next examined empirical noise correlations between firing rates in the pre-ramping baseline
period (-3.5 to -2.5 s) for MFC neurons recorded in the same session (see Analysis of Pairwise
Correlations in Methods; Fig. 5C). Pairs of neurons that later exhibited the same type of ramping
(i.e., both increasing or both decreasing) were significantly positively correlated before ramping
onset (increasing-increasing N=108 pairs: r = 0.048, 95% CI = [0.022 0.074], p < 0.001, LME;
decreasing-decreasing N=865 pairs: r = 0.038, 95% CI = [0.020 0.057], p < 0.001, LME),
whereas pairs with opposing types of ramping were not significantly correlated at baseline
(N=532 pairs; r =0.011, 95% CI = [-0.008 0.030], p = 0.285, LME; Fig. 5C). Further, same-type
pairs were more correlated than opposite-type pairs (increasing-increasing vs. increasing-
decreasing: p = 0.004; decreasing-decreasing vs. increasing-decreasing: p < 0.001, LME post-
hoc tests, Tukey correction for multiple comparisons) and were also more correlated than pairs
involving neurons that did not increase or decrease or had a steplike response profile (N = 18,384
pairs; increasing-increasing vs. other: p = 0.053, only trending; decreasing-decreasing vs. other:
p < 0.001). It is worth noting that noise correlations in the empirical data were smaller in
magnitude overall compared to model predictions (although their magnitude is in line with prior
studies[45,46]). Furthermore, noise correlations between opposite-type and other pairs were more
positive compared to model predictions (e.g., opposite: $r_{simulated}$ = -0.115, r_{data} = 0.011). Together,
these results are in line with the predictions made by an SFM (Fig. 5A-B), increasing the
plausibility that slow ramping signals reflect correlated fluctuations that are present from trial
onset to threshold-crossing.
515
516 **Discussion**
517
In this study, we demonstrated that spiking neural-networks can be configured to produce
spontaneous fluctuations and slow-ramping activity with properties like those of intracortical
experimental recordings in humans. Specifically, we recreated prior work that showed that
networks with clustered architectures exhibit spontaneous "slow switching" fluctuations[24,25], and
further showed that slow synaptic dynamics temporally stabilized those fluctuations that (Figs. 1-
2) and, in the aggregate, facilitated the emergence of gradual ramping activity before threshold-
crossings (Fig. 3). In networks with 50% slow synapses, simulated neurons slowly ramped their
firing rates, with 39% and 61% of slow ramping neurons increasing and decreasing their firing
rates respectively. Interestingly, individual neurons in networks with 50% slow synapses slowly
ramped their firing rates over 2 seconds before threshold-crossing, despite their autocorrelation
dropping to zero at lags of ~600 ms. By contrast, neurons in networks with 100% fast synapses
did not exhibit changes in their trial-average firing rates until around 300 or 500 ms prior to
movement in increasing and decreasing neurons respectively. Moreover, the accuracy of
classifiers trained to distinguish activity from a baseline period slowly ramped up over time
leading up to threshold-crossings (Fig. 4A left). Further, classifiers' accuracy generalized to later
time points better than they generalized to earlier time points (Fig. 4B left), indicating ramping at
the population-level[40].
535
Importantly, all of these features of simulated activity matched analogous analyses that we
carried out on intracranially recorded human MFC neurons[7] (Figs. 4A right, 4B right). Like
simulated neurons, autocorrelation in real neurons that exhibited slow ramping diminished to
zero within ~600 milliseconds. In addition, an aggregate signal derived from our model, which
could be interpreted as a proxy for EEG, slowly ramped negatively, akin to the readiness

potential[1] (RP) in terms of timing, shape, and ramping direction (Fig. 4C). Critically, all
simulated slow-ramping signals emerged spontaneously from ongoing activity, without
necessitating any external input at ramping onset to drive changes in activity. Finally, our model
(similar to other SFMs) predicted that neurons that show the same ramping behavior (increase or
decrease) after ramping onset would exhibit elevated pairwise correlations in their activity
already before ramping onset. This prediction was borne out in the empirical MFC neurons,
which exhibited a similar correlation structure to our model even before ramping onset (Fig. 5).
This analysis was not a test of our model against classical interpretations of ramping onset
(which suggest ramping onset reflects an event that drives ramping and, eventually,
movement[2,12], and thus make no particular prediction regarding pre-ramping correlations). But
this qualitative confirmation of our model's novel prediction in real data increases our model's
plausibility. Taken together, our study explains how highly autocorrelated spontaneous
fluctuations can emerge in spiking neural networks through a combination of topographical and
synaptic factors and demonstrates that such fluctuations are sufficient to explain several non-
trivial aspects of slow ramping neural activity preceding spontaneous voluntary action.
556
One of our central aims was to investigate the discrepancy between short autocorrelation
(decaying in ~600 ms) and slow ramping (lasting ~2 s—hence the period of slow ramping is
therefore ~3 times longer than the autocorrelation window for simulated and empirical neurons).
However, slow ramping at the single-neuron level emerges only when averaging across trials,
while autocorrelation is calculated from firing rates on individual trials. Inspection of raster plots
from individual neurons that exhibit slow ramping (on average across trials) shows that smooth,
generally monotonic ramping is not apparent on every trial (examples in Fig. 3E; more examples
in Fig. S3; this fact is also noted by Schurger et al.[19]). Rather, activity on each trial is highly
variable, and smoother, slow ramping emerges only in the trial-averaged firing rate. Our model
gives a potential explanation: in our model, movement is triggered after an entire cluster's
average activity crosses a threshold (see Threshold Alignment in Methods). Therefore, individual
neurons can and will show variable activity on individual trials. Still, trial-averaged response
profiles will show smooth ramps because their activity is, across many trials, entrained to the
activity of the cluster to which it belongs, which must increase (due to it later crossing the
threshold) or decrease (due to lateral inhibition) its net firing before threshold-crossings.
572
Our work expands on research into spontaneous voluntary actions by offering what is, to our
knowledge, the first model of slow ramping in spiking neural networks. Previous work had
investigated spontaneous neural activity[24,25,28,30,32] or autocorrelation in spiking neural
networks[27,28,47], but did not explore the connection to slow-ramping signals. Our model provides
an implementation at the level of spiking neurons of previous, more abstract stochastic
fluctuation models (SFMs) of spontaneous voluntary action, demonstrating how the fluctuations
that are central to those models[13–17] may emerge from biologically plausible spiking neural
networks. Also, many brain regions are organized in a modular fashion[48], and individual neural
circuits have clustered connectivity among excitatory neurons[49,50]. Such clustering could
plausibly emerge due to synaptic plasticity, such as Hebbian learning[51–53], and 'soft winner-take-
all' networks are hypothesized to be a canonical cortical circuit[54]. Furthermore, it is known that
synaptic transmission occurs through both fast and slow channels[55] (although, as is typical in
computational modeling, our implementation of slow synaptic transmission is likely an
oversimplification), and that some important cognitive functions rely on slow synapses[26]. Taken

together, clustered connectivity and slow synaptic transmission are a plausible combination that
we demonstrate can lead to spontaneous fluctuations and slow ramping. Additionally, the fact
that our model can parsimoniously explain many features of MFC activity before spontaneous
voluntary actions suggests that stochastic fluctuations in brain activity may underlie slow
ramping signals and play a key role in the genesis of spontaneous voluntary action. Our model
also goes beyond prior abstract computational models such as stochastic accumulation[14] by
describing a process that does not terminate at threshold crossing but is rather continuous in time.
In doing so, our model offers possibilities for investigating strings or cascades of actions rather
than single actions that occur in isolation.
596
Our results have major implications for the use of slow-ramping signals for predicting movement
intentions, for instance in brain-computer interfaces. Because neural activity can stochastically
increase or decrease, movement timing is not completely determined until threshold crossing.
Therefore, slow-ramping signals may be of limited utility in online, real-time prediction systems.
Indeed, predictions based on whole-brain EEG activity could predict movement timing above
chance in real time only ~600 ms before movements[56] (see also[57,58]), whereas the RP and other
slow ramping signals emerge 2 s or more before movement onset. Furthermore, in this respect,
the EEG proxy we derived closely matched the RP. And our proxy is based on post-synaptic
potentials, which exhibits a negative deflection before threshold-crossings due to an overall
increase of inhibitory signaling and decrease of excitatory signaling as part of the quasi-winner-
take-all process inherent to the networks we investigated. Taken together, we suggest that the
apparent onset of the RP and other slow ramping signals several seconds before movement is the
result of aligning spontaneous fluctuations to threshold-crossings rather than the many cognitive
interpretations that have been offered since the RP's discovery[3,19].
611
612 **Future Directions**
613
The model we developed is a "minimal plausible model" for slow ramping activity prior to
spontaneous voluntary action. To our knowledge, it is the first model to describe how such
ramping could arise spontaneously from spiking neural-network dynamics and captures several
nontrivial aspects of premovement slow ramping activity. However, many questions remain
about the neural mechanisms underlying spontaneous voluntary action. One key open question is
regarding the timing of threshold-crossing and what mechanisms underlies its implementation.
Prior models suggested that threshold-crossing leads to movement within about 200
milliseconds[14]. However, many non-ramping neurons in MFC and other brain areas abruptly
change their firing rates 0.5-1s before movement (Fig. 3C), around which time activity also
abruptly changes in parietal cortex[59]. Such activations may reflect the spontaneous ignition of
global activity patterns following threshold-crossings[17,60,61]. Future modeling work could
investigate this by directly implementing a threshold mechanism and by recreating the activity of
not just slow-ramping neurons but also those with a "step-like" response profile.
627
Slow ramping signals in the MFC have received much attention in the spontaneous voluntary
action literature. Supporting this view, Fried and colleagues[7] found that a greater fraction of
neurons in MFC changed their activity prior to movement compared to temporal regions,
suggesting a particular importance of this region. However, slow ramping signals have been
observed in other regions, including medial temporal cortex[7], sensory areas[6], and subcortical

areas such as the locus coeruleus (indexed by pupil dilations)[62]. Also, recordings in rats have
found ramping activity in the dopaminergic substantia nigra[63–65]. Recent large-scale recordings
of neural activity in mice further suggest that spontaneous voluntary movements involve widely
distributed rather than localized activity[66,67], as suggested by Schurger and Uithol[68]. Taken
together, fluctuations influencing movement timing may reflect a distributed, brain-wide process
rather than one localized only in MFC. One—admittedly speculative—possibility is that these
fluctuations are related to fluctuations in arousal[69], which may explain correlations between the
RP and respiration[70] and the presence of pupil dilations before freely timed movement in
humans[62]. If so, our model might thus be implemented not in cortical regions exhibiting slow
ramping activity, such as MFC, but rather in subcortical arousal hubs such as the Locus
Coeruleus. Fluctuations and ramping in cortical regions could then be caused by input from
these areas to the MFC[71]. Future studies should test this hypothesis by investigating the
spatiotemporal relationship between slow ramping activity (including activity prior to the onset
of slow ramping in trial-averaged activity, as we did in Fig. 5) and (1) peripheral signatures of
arousal and activity in arousal related areas, such as the Locus Coeruleus, and (2) activity more
proximal to the time of threshold-crossing or movement.
649
Crucially, because slow-ramping signals before spontaneous voluntary action may reflect
stochastic population-level neural fluctuations, they also offer the opportunity to validate models
that describe how temporally stable fluctuations emerge from neural dynamics. Brain activity
fluctuates over several seconds or tens of seconds[72], and the temporal properties of such
fluctuations correlate with behavior[73] change with age[74], and are disrupted in disease[75]. It is
therefore of interest to understand how these fluctuations emerge more generally. Indeed, prior
work has also suggested that such fluctuations emerge when neural systems are poised at the
edge of a phase transition[24,25,32,76–79] (i.e., at criticality). In the present study, we manipulated the
clustering coefficient of our networks to be just below a critical point, which had previously been
shown to lead to spontaneous fluctuations in network activitys[24,25]. By introducing slow synaptic
transmission, we showed that factors besides the parameter poised near criticality can alter the
spatiotemporal dynamics of emergent fluctuations. Our model suggests that clustering and slow
synapses lead to slow fluctuations emerging spontaneously in spiking networks (for instance,
during resting periods). However, models using different mechanisms could potentially recreate
our results. For instance, in our model we generated fluctuations by grouping neurons into
clusters that had relatively higher (lower) intra- (inter-) cluster probability of connection between
neurons. Schaub and colleagues[25] found that other architectures (e.g. small-world connectivity)
or varying intra-cluster synaptic strength instead of connection probability can also lead to
spontaneous fluctuations. Furthermore, in our model, we temporally smoothed fluctuations via
slow synaptic transmission. However, short-term plasticity or neuro-modulatory factors could
potentially also stabilize fluctuations, and short-term plasticity could potentially alter the
clustering degree, leading to further changes in fluctuation dynamics We elected to use slow
synapses over these other possibilities due to their simplicity—slow synapses do not involve
changes in connectivity, as short-term plasticity would, or factors external to the network, as
neuromodulatory factors would, and thus our results emerge spontaneously. Future modeling
efforts could explore whether combinations of these other factors would lead to results like ours
and compare models to establish which factor, or combination of factors, best explain
fluctuations in neural activity. Future studies could also investigate networks such as these
analytically, as Schaub and colleagues did for clustered networks with fast synapses[25], although

we note that slow synapses might complicate such analysis by making network activity
dependent on firing further in the past due to slower decay of synaptic currents (essentially
making dynamics non-Markovian).
682
Slow ramping signals also emerge in other behavioral contexts, such as delayed response tasks
(the contingent negative variation, or CNV, which is dissociated from the RP[80,81]), motor
preparation[82,83], and time perception[84,85]. Crucially, ramping in these other behavioral contexts
occurs between two well-defined points—usually an external stimulus that leads to a response—
whereas ramping onset in the spontaneous voluntary action context emerges without an external
cue. So, whether slow ramping reflects similar processes across these contexts is unclear. In
other "externally cued" contexts, reaction times are often too fast (hundreds of milliseconds) for
"slow ramping" to occur beforehand, and even without time pressure to respond, pre-movement
ramping, at least in MFC, is sometimes absent when decisions are not made spontaneously (e.g.
according to stimulus value)[15].
693
Interestingly, slow ramping with a seemingly spontaneous onset also occurs before other types of
spontaneous behaviors, including generation of creative ideas[86], spontaneous abstract
decisions[87,88], eureka moments in problem solving[89], free recall of memories[79,90,91], and switches
between bistable percepts during binocular rivalry[92]. Notably, slow ramping signals before
creative ideas have been linked to the temporal properties of spontaneous resting-state
fluctuations[86], similar to the relationship between autocorrelation and slow ramping[13] that we
investigated. Others have noticed these similarities and proposed that threshold-crossings by
slow spontaneous fluctuations underlie various types of spontaneous behaviors. It is thus
tempting to suggest that the neural mechanism we investigated in the present study may apply to
spontaneous cognitive and perceptual processes more generally, offering an intriguing direction
for future empirical studies.
705

Acknowledgements
We thank Gabriel Kreiman for granting us access to the neural recordings we analyzed as part of
this study and for discussion and feedback on an earlier version of this manuscript. We also
thank Bill Newsome and Robert Kim for feedback on an earlier version of this manuscript. J.G.,
A.S., and U.M. were supported by the John Templeton Foundation and the Fetzer Institute
(Consciousness and Free Will: A Joint Neuroscientific-Philosophical Investigation (John
Templeton Foundation #61283; Fetzer Institute, Fetzer Memorial Trust #4189)). U.R. was
supported by the US National Institutes of Health (U01NS117839) and J.G. and U.R. were
supported by the NSF (BCS-2219800).

Code and Data Availability
Code used for simulating data & analyzing simulated data will be made publicly available on
OSF upon acceptance.
719

720   **Author Contributions**
721   J.G., U.R., A.S., and U.M. conceptualized the project. J.G. conducted simulations and analysis of
722   simulated and empirical data. U.R., A.S., and especially U.M. supervised and provided feedback
723   on simulations and data analysis. J.G., U.R., A.S., and U.M. wrote the paper.
724

725 **Declaration of Interests**
726 The authors declare no competing interests.
727
728 **Methods**
729 **LIF Network Simulations**
730 We simulated networks of leaky integrate-and-fire (LIF) neurons where the membrane potential
731 V_m was governed (and updated at a resolution of 1 ms) by the equations:
732

733 $$\tau_m \frac{dV_m}{dt} = (E_L - V_m) + I_{app} + RI + \zeta$$

734
Here τ_m is the synaptic time-constant (10 ms for our simulations). E_L is the reversal potential (-
70 mV). I_{app} is a constant bias current set at the rheobase (300 pA). $\zeta \sim N(0, 0.5\ mV)$ is the
noise term, which changes every 10 ms. In other words, each neuron receives constant noise for
10 ms, which then changes. The membrane potential, V_m, had a threshold of -40 mV, and upon
crossing it was reset to -65 mV and held constant for a refractory period of 2 ms. In every
simulation $V_{m\ initial} \sim U(-70, -50\ mV)$ for each neuron.
741
742 RI is the synaptic current vector. For each neuron it is a sum (weighted according to the
743 connectivity matrix W; details below) of the other neurons' synaptic output. Each such synaptic
744 output is itself a weighted sum of fast and slow synaptic currents. RI follows the equation:
745

746 $$RI = W * \left(0.9 * K_{syn} * r_{slow} + \left(1 - K_{syn}\right) * r_{fast}\right)$$

747
Here, $0 \leq K_{syn} \leq 1$ is the ratio of fast to slow synapses. Hence, if $K_{syn} = 0$ or $K_{syn} = 1$ the
network's activity would entirely be governed by fast or slow synapses, respectively. And
$0 < K_{syn} < 1$, the network's activity is governed by a weighted sum of fast and slow synaptic
currents. We multiplied the slow synaptic currents by 0.9 to afford better comparisons between
networks with similar clustering values (see below). Synaptic currents, r, follow a double-
exponential time course including a rising and falling time-constant which differs between fast
and slow synapses. Thus r follows the equations:

$$\frac{dr}{dt} = -\frac{1}{\tau_{fall}} r + s$$

$$\frac{ds}{dt} = -\frac{1}{\tau_{rise}} s + \frac{1}{\tau_{rise}\tau_{fall}} * Spikes(t)$$

757
Here τ_{rise} and τ_{fall} are the time-constants for the synaptic currents to rise to their peak and then
decay, respectively. Fast synapses had $\tau_{rise} = 2\ ms$ and $\tau_{fall} = 5ms$ and slow synapses had
$\tau_{rise} = 20\ ms$ and $\tau_{fall} = 100\ ms$ [33]. $Spikes(t)$ is a binary (0 or 1) spiking vector reflecting,
for each time-step in the simulation, whether each neuron had spiked on the previous timestep.
These spiking values were normalized by the rising and falling time constants, which ensured
that the same total current was delivered for fast and slow synapses (see Fig. 1B).
764

Connectivity Matrix Construction

Much of the dynamics of neural networks are determined by how neurons are interconnected, encoded in the connectivity matrix W. Previous research has found that *balanced* networks with roughly equal excitatory and inhibitory signaling exhibit stable asynchronous dynamics[30]. Balanced networks are usually implemented with more excitatory than inhibitory neurons (usually 75-80% excitatory), though with relatively weaker synaptic signaling in those excitatory neurons (to compensate for their higher proportion). To introduce slow changes in firing rates, we grouped the excitatory neurons into "clusters" both topographically and synaptically. Topographically, we raised the probability of connection for inter-cluster excitatory neurons in comparison with intra-cluster excitatory neurons. Synaptically, we increased the weights representing connections between excitatory neurons in the same cluster[24] (although similar slow fluctuations in firing rates can be achieved by increasing intra-cluster connectivity strength alone, as well as through other methods[25]). Notably, however, the slow fluctuations in firing rates that emerged due to this clustering were depended significantly on the synaptic dynamics (see LIF Network Simulations section above).

We wanted to maintain sparse connectivity in our network, hence we used a 20% connection probability between any two excitatory neurons (P_{EE}), taking into account the different number of intra- and inter-cluster connections. In a network with more than two clusters, there will be a larger percentage of inter-cluster connection opportunities than intra-cluster ones. So, calculating the precise values of the intra-cluster connection probability (P_{EE}^{In}) and inter-cluster connection probability (P_{EE}^{Out}) is non-trivial. To do so, we first note that the overall probability of connection (P_{EE}) is a sum of intra- and inter-cluster connections divided by the total number of possible connections between excitatory neurons:

$$P_{EE} = \frac{E(\# \, Intra - cluster \, connections) + E(\# \, Inter - cluster \, connections)}{N_E^2} = 0.2$$

Next, we find the expected number of intra-cluster connections:

$$E(\# \, Intra - cluster \, connections) = P_{EE}^{In} * N_{Clusters} * C^2 = P_{EE}^{In} \frac{N_E^2}{N_{Clusters}}$$

Where $N_{Clusters}$ is the number of clusters, C is the number of neurons in each cluster (which equals $\frac{N_E}{N_{Clusters}}$), and N_E is the total number of excitatory neurons. The expected number of inter-cluster connections similarly follows:

$$E(\# Inter - cluster \, connections) = P_{EE}^{Out} * N_{Clusters}(N_{Clusters} - 1) * C^2$$
$$= P_{EE}^{Out} \frac{N_E^2 (N_{Clusters} - 1)}{N_{Clusters}}$$

Then, letting $A = \frac{N_E^2}{N_{Clusters}}$ and $B = \frac{N_E^2(N_{Clusters} - 1)}{N_{Clusters}}$ for simplicity, we obtain:

$$P_{EE} = \frac{P_{EE}^{In} * A + P_{EE}^{Out} * B}{N_E^2}$$

Finally, defining the clustering ratio R_{EE} as

$$R_{EE} = \frac{P_{EE}^{In}}{P_{EE}^{Out}}$$

we can then solve explicitly for P_{EE}^{In} and P_{EE}^{Out}:

$$P_{EE}^{In} = \frac{N_E{}^2 P_{EE} R_{EE}}{B + A R_{EE}}$$

$$P_{EE}^{Out} = \frac{N_E{}^2 P_{EE}}{B + A R_{EE}}$$

Solving for these probabilities explicitly thereby allows for a relatively easy construction of
connectivity matrices corresponding to clustered networks, while maintaining a constant
probability of connection overall (P_{EE}=0.2, or 20% between excitatory neurons). Furthermore,
these equations demonstrate that the precise values of P_{EE}^{In} and P_{EE}^{Out} depend not only on the
clustering coefficient (R_{EE}), but also the number of excitatory neurons (N_E), and the number of
clusters ($N_{Clusters}$) (or, equivalently, on the number of clusters and size of each cluster).
811
Simulating & Analyzing Fluctuations
During simulations we varied two control parameters: the clustering coefficient, R_{EE} (defined
below), and the ratio of slow-to-fast synapses, K_{SYN} (defined above). We simulated 5 s of activity
in 50 unique network architectures for a range of R_{EE} values (1 to 6 at increments of 0.1) and a
range of synapse ratios (0% to 50% slow synapses at increments of 10%). After discarding the
first 500 ms of each simulation, we calculated several measures of temporal variability from
simulated firing rates. First, we calculated the power spectra of firing rates (obtained by
convolving spike trains with a gaussian kernel of width 400 ms) using the fast-Fourier transform
in MATLAB and then taking the absolute value of the resulting complex-valued data. For
calculating the spectral slope of these power spectra, we transformed data into log-log space and
linearly regressed power on frequency, and then took the slope of the resulting linear fit as the
spectral slope. For autocorrelation, we calculated firing rates using 50 ms non-overlapping bins
and then calculated the autocorrelation of the resulting time-series up to lag 20 (corresponding to
1000 ms). For autocorrelation timescales, we fit an exponential function to the autocorrelation at
lag t using the following equation:

$$ACF(t) = a + b * e^{\frac{-t}{\tau}}$$

And took the decay constant (τ) as the autocorrelation timescale. We used the approach of
Cavanagh and colleagues[22] by fitting an exponential function to autocorrelation averaged across
all excitatory neurons in a given network (as opposed to fitting and then averaging). We also
followed prior research[20,22] and excluded neurons with trial-average firing rates less than 2.5 Hz
from further analysis, and omitted τ and autocorrelation of a network from further analyses if any
of the following criteria were met: (1) fitted τ was not between 0 and 1000 ms, (2) fitted
parameter b was below zero, or (3) if fitting of the exponential model did not converge. These
rejection criteria led to omission of 3675 out of 15300 simulated trials (24%), but more than half
of that (1913) came from networks with clustering coefficients above 4, at which point activity
started transitioning into a winner-take-all regime.
838
Next, for CV, we calculated the firing rates of each excitatory neuron i ∈ {1, …, N}, where N is
320 (80% of 400), over time using non-overlapping bins of 100 ms. From these we calculated the
coefficient of variation of firing rates across time (standard deviation divided by mean). For each
neuron, with T time bins and firing rates at each bin we get a sequence Fi = [fi(t1), fi(t2), …,
fi(tT)], over which we calculated the standard deviation σi and mean μi. Then, the network-level
coefficient of variation is as follows:

$$CV_{Net} = \frac{1}{N} \sum_{i=1}^{N} \frac{\sigma_i}{\mu_i}$$

We calculated a single CV_{Net} value for each network, and plot grand means and standard error
across networks in Fig. S1A. In Fig. S1B we plot individual network-average values (for the first
10 of the 50 networks to improve visibility).

In order to quantify spiking variability that was due to slow-switching or bistable activity, we
also calculated the $\hat{S}T$ value used by Schaub and colleagues[25] to investigate slow-switching
behavior. This measure quantifies spiking variability over time by taking the average firing rate
in each cluster $i \in \{1, \ldots, C\}$ in 100 ms non-overlapping bins. Then, for each cluster with firing
rates in T time bins we get a sequence of firing rates $F_i = [f_i(t_1), f_i(t_2), \ldots, f_i(t_T)]$ where t_k
represents the kth time bin, for which we calculate the standard deviation $\sigma_T(i)$, and then sum to
obtain an initial measure of spiking variability.

$$S_T = \frac{1}{C} \sum_{i=1}^{C} \sigma_T(i)$$

860   We then normalize this value by subtracting a bootstrapped value S_{Boot} calculated on shuffled
861   data (shuffling each neuron's cluster membership), averaged over 20 random shuffles.

$$\hat{S}_T = S_T - S_{Boot}$$

865   Like for CV, we calculated a single $\hat{S}_T$ value for each network, and plot grand means and
866   standard error across networks in Fig. S1C. For Fig. S1D we once again plot individual network-
867   average values for the first ten of fifty networks.

Network Selection for Threshold-Aligned Analyses

Prior work had shown that clustered networks exhibit fluctuations for a critical range of
clustering coefficients ($2 < R_{EE} < 5$ typically, although this depends on network specifics;
Litwin-Kumar & Doiron, 2012; Schaub et al., 2015). Because we generated network structures
pseudo-randomly (setting the seed for random generation using MATLAB's rng command), the
critical range was slightly different for each network. We identified an R_{EE} value suitable for
simulation in each network by visually inspecting activity at a range of values. We selected R_{EE}
values for each network such that the network exhibited fluctuations at the given value for both
100% fast and 50% slow synapses, but exhibited winner-take-all behavior in at least one case if
we increased the R_{EE} by 0.3. Prior work had linked the R_{EE} value demarking the transition to the
winner-take-all regime to the leading eigenvalue of the connectivity matrix exceeding a value of
1 [25]. However, our simulations did not exhibit such a relation, likely because that prior work used
non-dimensionalized equations for simulating our activity whereas we used the dimensionalized
versions of those equations.

Threshold Alignment

We hypothesized that spontaneous fluctuations trigger movements upon crossing a threshold,
leading to slow ramping in trial-averaged activity. To investigate this hypothesis, we simulated

10-second trials 100 times for each of the 20 network architectures and then aligned to threshold-
crossings (defined below). We identified the most-active cluster across all simulated trials and
aligned to threshold crossings in that cluster (reflecting the notion that threshold crossings in
only some subpopulation of SMA would lead to a movement). If no threshold crossing was
detected in a trial, the trial was omitted from further analyses.
892
Threshold crossings were defined as the time at which the most-active cluster crossed 50% of its
maximum normalized firing rate across the entire simulation period. This was obtained by
averaging firing rates across neurons in each cluster, subtracting the mean firing rate of the first
500 ms, and then dividing by the maximum firing rate on each trial. The choice to use a different
threshold for each network was motivated by the fact that the magnitude and other features of
fluctuations were heterogeneous across different networks even with similar clustering
coefficients. Our choice of 50% of the maximum normalized firing rate was motivated by Fried
and colleagues'[7] finding that increasing neurons reached about 50% of their maximum
normalized firing rate right before movement (Fig. 4 in Fried et al., 2011). Only trials where
crossing occurred more than 3 s into the trial were retained for analysis of neural data (as is
commonly done in self-initiated action paradigms[7]), as we were looking at changes across this
period.
905
Neurons were categorized as increasing or decreasing by comparing their firing rates in the 400
ms before threshold crossings to their firing rates during a baseline period (-3 to -2 s, rank-sum
test with a cut-off of p=0.01, as in[7]). Firing rates were smoothed with a gaussian kernel (width
400 ms), then averaged across response types (increasing and decreasing) to get a single trace for
each network. Those traces were then averaged across networks for Fig. 2A; the shaded regions
reflect the confidence intervals across networks.
912
Our EEG proxy was obtained by averaging the post-synaptic potentials (PSPs) onto excitatory
neurons[41]. We retained the PSPs onto each neuron (*RI* in the above equations). We then sub-
selected the PSPs to retain only those going onto excitatory neurons, averaging across neurons to
obtain a single EEG proxy per trial. We then averaged across trials to get a single EEG proxy
trace per network. Those traces were again averaged across networks for Fig. 2B, and the shaded
regions reflect the confidence intervals across networks.
919
Autocorrelation at the single-neuron and cluster level was obtained using the autocorr function in
MATLAB on the firing rates (smoothed using a moving average filter with a window of 10 ms to
avoid smoothing artifacts; then averaged across neurons in each cluster for cluster-average firing
rates) for each -3 to 0.5-second trial aligned to threshold-crossings. Those traces were then
averaged across trials to get a single autocorrelation trace for each neuron or cluster. Then, those
traces were averaged across neurons or clusters to get a single trace per network and then
averaged across networks to obtain grand averages for Fig. 2C.
927
Neuron Categorization & Response Profiles
We categorized empirical and simulated neurons as increasing or decreasing following the
process described by Fried et al.[7]. We compared firing rates in the 400 ms before threshold
crossing or movement to firing rates during a baseline period (-3 to -2 s) using a non-parametric
rank-sum test. Neurons that significantly increased (decreased) their activity relative to baseline

were categorized as increasing (decreasing; cut-off was p=0.01). Notably, this process omits
neurons with more complex response profiles (e.g. Fig. 3 in Fried et al., 2011[7]), but these are out
of the scope of this study.
936
MFC neurons had a mixture of response profiles. In particular, some neurons slowly ramped
their firing rates up or down before movement, and others exhibited a steep jump up or down
before movement. Simulated neurons tended to exhibit the same type of response (slow ramp or
steplike), but responses in empirical data were mixed and therefore required a method to
distinguish them. To do so, we followed a process similar to the supplementary analysis
conducted by Fried et al.[7] (supplementary analysis/Fig. S2 in their article). We fit a sigmoid
function to the trial-averaged smoothed firing rates (400 ms Gaussian) using nonlinear least-
squares via the Levenberg-Marquardt algorithm in MATLAB. We fit a different sigmoid
depending on if the neuron was increasing or decreasing:

946
$$\frac{F_W \cdot (-1)^d}{1 + exp(-(x - t_0)/\alpha)} + F_B.$$

Here F_B is the firing rate at baseline, F_W is the firing rate in the 400 ms before movement. t_0 is the
(fitted) sigmoid inflection point. α is the (fitted) inverse-gain parameter (i.e. higher values of α
correspond to a smoother ramp). And d is 0 for increasing neurons and 1 for decreasing ones. We
selected an inverse-gain of 0.4 as the cut-off for steplike versus slow ramp response profiles
based on visualizing averaged firing rates from the Fried et al.[7] (2011) dataset (see Fig. 2C).
Thus, neurons with an α parameter greater than 0.4 were categorized as slow-ramping, and
neurons with an α parameter less than 0.4 were categorized as steplike.
954

Decoding Analyses
We also assessed whether above-chance decoding accuracy in the seconds before movement[7,37]
might emerge from aligning fluctuations like those in our networks to threshold-crossings. For
simulated data, at each time point, we compared spike counts in the time bin [t-0.4, t] s to spike
counts in the baseline of (-3,-2.6) s for all neurons. We first constructed a pseudopopulation of
neurons for which we had more than 40 trials by grouping them into one dataset for purposes of
the decoding analysis. We trained a linear discriminant analysis classifier on 80% of the data and
recorded its performance on a reserved 20% test set (i.e., 5-fold cross-validation). Fig. 3A shows
the average accuracy and standard error across 5 cross-validation splits for individual networks
(the same splits were used for $K_{syn} = 0$ and $K_{syn} = 0.5$), as well as performance on 100 random
splits of the data to assess chance-level performance. For empirical data, we constructed a
pseudo population of all recorded MFC neurons. We aggregated across subjects with at least 50
trials and conducted the same analysis as above, with time relative to movement onset rather than
threshold-crossing.
969

To assess the temporal generalization of decoding (Fig. 3B), we completed a similar analysis as
above. However, rather than splitting the data at a single time point into a training and testing set,
we trained the classifier on the data from a given time point and then tested the trained classifier
on a different time point[40]. Therefore, the main diagonal (train and test timepoint are the same)
corresponds to the classifiers' training accuracy; the off-diagonal results refer to the classifiers'
test accuracy.
976

Simulating Ramping Spike-Trains

We simulated spike trains where we knew that the ground-truth was a gradual ramp in order to
see if the asymmetry we observed in our temporal generalization analysis (where decoders
trained to differentiate activity from a baseline generalize better into the future than into the past)
is due to ramping. We simulated 50 spike trains for a pseudopopulation of 20 neurons. Half of
the neurons increased their firing rates from 5 Hz to 8 Hz, and the other half decreased their
firing rates from 8 Hz to 5 Hz. We simulated 3 s of activity (labeled as -3 to 0 to match other
analyses) and generated spike trains in 50 ms bins, where the instantaneous firing rate was
obtained by linearly interpolating the starting and ending firing rates. Then, we conducted the
same temporal generalization analysis as above using a baseline of [-3,-2.6]. We found a striking
asymmetry in the AUCs such that decoders generalized better into the future than into the past
(one-tailed rank-sum test on upper vs lower triangle of AUC matrix, p < 0.001; Fig S4D).
989
Analysis of Pairwise Correlations at Baseline
Our model suggests that early ramping reflects fluctuations in clustered networks aligned to
threshold-crossings. Crucially, those fluctuations emerge naturally due to the clustered
connectivity, and also because neurons that ramp up tend to be in the same cluster, and those that
ramp down tend to belong to clusters other than the one ramping up. However, in our model,
connectivity is static, as would be expected in biological networks at the time scales in which we
simulate here. Neurons in the same cluster therefore show correlated activity across the entire
trial, not just during ramping. Hence, our model predicts that neurons that ramp in up or down
together before movement or threshold crossings should have correlated activity even before
ramping onset.
1000
We thus investigated pairwise correlations at baseline in empirical neurons in a baseline period
of (-3.5,-2.5) (a baseline of (-3,-2.5) was used for simulated data because we did not save data
earlier than -3 s). We first smoothed (Gaussian kernel, 400 ms wide), then soft-normalized
(divided by the square root of the maximum firing rate[39]) firing rates, and finally down-sampled
to a sample every 50 ms to avoid inflating correlations due to autocorrelations in the data. After
that, we correlated firing rates between every pair of neurons that were recorded in the same
session, keeping track of whether the pair was both increasing, both decreasing, one increasing
and one decreasing, or involved a neuron that didn't change its firing rate (for these analyses we
categorized neurons with a steplike response as "other" in order to focus solely on slow-ramping
behavior). At that point, we used linear mixed-effects models (LME; with the network or subject
as a random intercept) to investigate whether correlation structure at baseline depended on later
ramping behavior. We found that increasing and decreasing pairs show significantly correlated
activity at baseline. In contrast, increasing-decreasing pairs do not; they also show significantly
lower correlations than increasing-increasing pairs and decreasing-decreasing pairs. Notably,
running the same analysis without separating slow ramping and steplike responses led to all
shown statistical tests in Fig. 4C becoming highly significant (both increasing vs increasing-
decreasing p = 0.001; both increasing vs other p = 0.013; both decreasing vs increasing-
decreasing p < 0.001; both decreasing vs other p < 0.001; LME post-hoc tests, Tukey correction),
as well as the difference between increasing-decreasing and other pairs becoming significant (p =
0.040). Additionally, omitting pairwise comparisons involving two neurons on the same channel
(accounting for potential issues due to spike sorting) largely did not change the results for
decreasing-decreasing pairs, which had many examples. However, only few slow ramping
increasing-increasing pairs remained after this correction, reducing statistical power and leading

to insignificant differences between increasing-increasing vs increasing-decreasing pairs (p = 0.439), and increasing-increasing vs other pairs (p = 0.777). However, the estimated correlation was similar to decreasing neurons after that correction, so we suggest that the lack of statistical power is the main cause of these differences.

References

1. Kornhuber, H. H. & Deecke, L. [CHANGES IN THE BRAIN POTENTIAL IN VOLUNTARY MOVEMENTS AND PASSIVE MOVEMENTS IN MAN: READINESS POTENTIAL AND REAFFERENT POTENTIALS]. *Pflugers Arch Gesamte Physiol Menschen Tiere* **284**, 1–17 (1965).
2. Libet, B., Gleason, C. A., Wright, E. W. & Pearl, D. K. Time of conscious intention to act in relation to onset of cerebral activity (readiness-potential). The unconscious initiation of a freely voluntary act. *Brain* **106 (Pt 3)**, 623–642 (1983).
3. Shibasaki, H. & Hallett, M. What is the Bereitschaftspotential? *Clin Neurophysiol* **117**, 2341–2356 (2006).
4. Cunnington, R., Windischberger, C., Deecke, L. & Moser, E. The Preparation and Execution of Self-Initiated and Externally-Triggered Movement: A Study of Event-Related fMRI. *NeuroImage* **15**, 373–385 (2002).
5. Cunnington, R., Windischberger, C. & Moser, E. Premovement activity of the pre-supplementary motor area and the readiness for action: Studies of time-resolved event-related functional MRI. *Human Movement Science* **24**, 644–656 (2005).
6. Sakata, H. *et al.* Slow Accumulations of Neural Activities in Multiple Cortical Regions Precede Self-Initiation of Movement: An Event-Related fMRI Study. *eNeuro* **4**, ENEURO.0183-17.2017 (2017).
7. Fried, I., Mukamel, R. & Kreiman, G. Internally Generated Preactivation of Single Neurons in Human Medial Frontal Cortex Predicts Volition. *Neuron* **69**, 548–562 (2011).
8. Bauer, H. H. & Rebert, C. S. Preliminary study on subcortical slow potentials related to the readiness potential in monkey. *Int J Psychophysiol* **9**, 269–278 (1990).
9. Murakami, M., Vicente, M. I., Costa, G. M. & Mainen, Z. F. Neural antecedents of self-initiated actions in secondary motor cortex. *Nat Neurosci* **17**, 1574–1582 (2014).
10. Mitelut, C. *et al.* Mesoscale cortex-wide neural dynamics predict self-initiated actions in mice several seconds prior to movement. *eLife* **11**, e76506 (2022).
11. Kagaya, K. & Takahata, M. Readiness Discharge for Spontaneous Initiation of Walking in Crayfish. *J. Neurosci.* **30**, 1348–1362 (2010).
12. Libet, B. Unconscious cerebral initiative and the role of conscious will in voluntary action. *THE BEHAVIORAL AND BRAIN SCIENCES* 38 (1985).
13. Schurger, A. Specific Relationship between the Shape of the Readiness Potential, Subjective Decision Time, and Waiting Time Predicted by an Accumulator Model with Temporally Autocorrelated Input Noise. *eNeuro* ENEURO.0302-17.2018 (2018) doi:10.1523/ENEURO.0302-17.2018.
14. Schurger, A., Sitt, J. D. & Dehaene, S. An accumulator model for spontaneous neural activity prior to self-initiated movement. *PNAS* **109**, E2904–E2913 (2012).
15. Maoz, U., Yaffe, G., Koch, C. & Mudrik, L. Neural precursors of decisions that matter—an ERP study of deliberate and arbitrary choice. *eLife* **8**, e39787 (2019).
16. Khalighinejad, N., Schurger, A., Desantis, A., Zmigrod, L. & Haggard, P. Precursor processes of human self-initiated action. *NeuroImage* **165**, 35–47 (2018).
17. Moutard, C., Dehaene, S. & Malach, R. Spontaneous Fluctuations and Non-linear Ignitions: Two Dynamic Faces of Cortical Recurrent Loops. *Neuron* **88**, 194–206 (2015).

18. Schmidt, S., Jo, H.-G., Wittmann, M. & Hinterberger, T. 'Catching the waves' – slow cortical
potentials as moderator of voluntary action. *Neuroscience & Biobehavioral Reviews* **68**,
639–650 (2016).
19. Schurger, A., Hu, P. 'Ben', Pak, J. & Roskies, A. L. What Is the Readiness Potential?
Trends in Cognitive Sciences **25**, 558–570 (2021).
20. Cavanagh, S. E., Wallis, J. D., Kennerley, S. W. & Hunt, L. T. Autocorrelation structure at
rest predicts value correlates of single neurons during reward-guided choice. *eLife* **5**,
e18937 (2016).
21. Murray, J. D. *et al.* A hierarchy of intrinsic timescales across primate cortex. *Nature
Neuroscience* **17**, 1661–1663 (2014).
22. Cavanagh, S. E., Hunt, L. T. & Kennerley, S. W. A Diversity of Intrinsic Timescales Underlie
Neural Computations. *Frontiers in Neural Circuits* **14**, (2020).
23. Bernacchia, A., Seo, H., Lee, D. & Wang, X.-J. A reservoir of time constants for memory
traces in cortical neurons. *Nat Neurosci* **14**, 366–372 (2011).
24. Litwin-Kumar, A. & Doiron, B. Slow dynamics and high variability in balanced cortical
networks with clustered connections. *Nature Neuroscience* **15**, 1498–1505 (2012).
25. Schaub, M. T., Billeh, Y. N., Anastassiou, C. A., Koch, C. & Barahona, M. Emergence of
Slow-Switching Assemblies in Structured Neuronal Networks. *PLOS Computational Biology*
11, e1004196 (2015).
26. Wong, K.-F. & Wang, X.-J. A Recurrent Network Mechanism of Time Integration in
Perceptual Decisions. *J. Neurosci.* **26**, 1314–1328 (2006).
27. Li, Y., Kim, R. & Sejnowski, T. J. Learning the Synaptic and Intrinsic Membrane Dynamics
Underlying Working Memory in Spiking Neural Network Models. *Neural Computation* **33**,
3264–3287 (2021).
28. Klinshov, V. V., Kovalchuk, A. V., Franović, I., Perc, M. & Svetec, M. Rate chaos and
memory lifetime in spiking neural networks. *Chaos, Solitons & Fractals* **158**, 112011 (2022).
29. Fino, E. & Yuste, R. Dense inhibitory connectivity in neocortex. *Neuron* **69**, 1188–1203
(2011).
30. van Vreeswijk, C. & Sompolinsky, H. Chaos in neuronal networks with balanced excitatory
and inhibitory activity. *Science* **274**, 1724–1726 (1996).
31. Oster, M., Douglas, R. & Liu, S.-C. Computation with spikes in a winner-take-all network.
Neural Comput **21**, 2437–2465 (2009).
32. Cramer, B. *et al.* Autocorrelations from emergent bistability in homeostatic spiking neural
networks on neuromorphic hardware. Preprint at https://doi.org/10.48550/arXiv.2208.08329
(2022).
33. Gerstner, W. & Kistler, W. M. *Spiking Neuron Models: Single Neurons, Populations,
Plasticity*. (Cambridge University Press, Cambridge, U.K. ; New York, 2002).
34. Rothman, J. S. & Silver, R. A. Data-Driven Modeling of Synaptic Transmission and
Integration. *Prog Mol Biol Transl Sci* **123**, 305–350 (2014).
35. Kramer, M. A. & Chu, C. J. The 1/f-like behavior of neural field spectra are a natural
consequence of noise driven brain dynamics. *bioRxiv* 2023.03.10.532077 (2023)
doi:10.1101/2023.03.10.532077.
36. Cavanagh, S. E., Towers, J. P., Wallis, J. D., Hunt, L. T. & Kennerley, S. W. Reconciling
persistent and dynamic hypotheses of working memory coding in prefrontal cortex. *Nat
Commun* **9**, 3498 (2018).
37. Soon, C. S., Brass, M., Heinze, H.-J. & Haynes, J.-D. Unconscious determinants of free
decisions in the human brain. *Nature Neuroscience* **11**, 543–545 (2008).
38. Zapparoli, L. *et al.* Dissecting the neurofunctional bases of intentional action. *PNAS* **115**,
7440–7445 (2018).
39. Fu, Z. *et al.* The geometry of domain-general performance monitoring in the human medial
frontal cortex. *Science* **376**, eabm9922 (2022).

40. King, J.-R. & Dehaene, S. Characterizing the dynamics of mental representations: the
temporal generalization method. *Trends Cogn Sci* **18**, 203–210 (2014).
41. Nunez, P. L. & Srinivasan, R. *Electric Fields of the Brain: The Neurophysics of EEG, 2nd*
Edition. (Oxford University Press, Oxford ; New York, 2005).
42. Vanderhaeghen, P. & Polleux, F. Developmental mechanisms underlying the evolution of
human cortical circuits. *Nat Rev Neurosci* **24**, 213–232 (2023).
43. Panzeri, S., Moroni, M., Safaai, H. & Harvey, C. D. The structures and functions of
correlations in neural population codes. *Nat Rev Neurosci* **23**, 551–567 (2022).
44. Cohen, M. R. & Kohn, A. Measuring and interpreting neuronal correlations. *Nat Neurosci* **14**,
811–819 (2011).
45. Daume, J. *et al.* Control of working memory maintenance by theta-gamma phase amplitude
coupling of human hippocampal neurons. *bioRxiv* 2023.04.05.535772 (2023)
doi:10.1101/2023.04.05.535772.
46. Valente, M. *et al.* Correlations enhance the behavioral readout of neural population activity
in association cortex. *Nat Neurosci* **24**, 975–986 (2021).
47. Kim, R. & Sejnowski, T. J. Strong inhibitory signaling underlies stable temporal dynamics
and working memory in spiking neural networks. *Nature Neuroscience* **24**, 129–139 (2021).
48. Sporns, O. & Betzel, R. F. Modular Brain Networks. *Annu Rev Psychol* **67**, 613–640 (2016).
49. Song, S., Sjöström, P. J., Reigl, M., Nelson, S. & Chklovskii, D. B. Highly nonrandom
features of synaptic connectivity in local cortical circuits. *PLoS Biol* **3**, e68 (2005).
50. Perin, R., Berger, T. K. & Markram, H. A synaptic organizing principle for cortical neuronal
groups. *Proceedings of the National Academy of Sciences* **108**, 5419–5424 (2011).
51. Tetzlaff, C., Dasgupta, S., Kulvicius, T. & Wörgötter, F. The Use of Hebbian Cell Assemblies
for Nonlinear Computation. *Sci Rep* **5**, 12866 (2015).
52. Gerstner, W. & Kistler, W. M. Mathematical formulations of Hebbian learning. *Biol Cybern*
87, 404–415 (2002).
53. Nachstedt, T., Wörgötter, F. & Tetzlaff, C. Self-organized cell assembly formation. *BMC*
Neurosci **15**, P30 (2014).
54. Douglas, R. J. & Martin, K. A. C. Neuronal Circuits of the Neocortex. *Annual Review of*
Neuroscience **27**, 419–451 (2004).
55. Greengard, P. The Neurobiology of Slow Synaptic Transmission. *Science* **294**, 1024–1030
(2001).
56. Bai, O. *et al.* Prediction of human voluntary movement before it occurs. *Clinical*
Neurophysiology **122**, 364–372 (2011).
57. Lew, E., Chavarriaga, R., Zhang, H., Seeck, M. & Millán, J. del R. Self-paced movement
intention detection from human brain signals: Invasive and non-invasive EEG. in *2012*
Annual International Conference of the IEEE Engineering in Medicine and Biology Society
3280–3283 (2012). doi:10.1109/EMBC.2012.6346665.
58. Lew, E. Y. L., Chavarriaga, R., Silvoni, S. & Millán, J. del R. Single trial prediction of self-
paced reaching directions from EEG signals. *Frontiers in Neuroscience* **8**, (2014).
59. Aflalo, T. *et al.* Implicit mechanisms of intention. *Current Biology* **32**, 2051-2060.e6 (2022).
60. Baars, B., Franklin, S. & Ramsøy, T. Global Workspace Dynamics: Cortical "Binding and
Propagation" Enables Conscious Contents. *Frontiers in Psychology* **4**, (2013).
61. Baars, B. J., Geld, N. & Kozma, R. Global Workspace Theory (GWT) and Prefrontal Cortex:
Recent Developments. *Frontiers in Psychology* **12**, (2021).
62. Richer, F. & Beatty, J. Pupillary Dilations in Movement Preparation and Execution.
Psychophysiology **22**, 204–207 (1985).
63. Hamilos, A. E. *et al.* Slowly evolving dopaminergic activity modulates the moment-to-
moment probability of reward-related self-timed movements. *eLife* **10**, e62583 (2021).
64. Lee, I. H. & Assad, J. A. Putaminal Activity for Simple Reactions or Self-Timed Movements.
Journal of Neurophysiology **89**, 2528–2537 (2003).

65. Schultz, W. & Romo, R. Role of primate basal ganglia and frontal cortex in the internal generation of movements. I. Preparatory activity in the anterior striatum. *Exp Brain Res* **91**, 363–384 (1992).

66. Stringer, C. *et al.* Spontaneous behaviors drive multidimensional, brainwide activity. *Science* **364**, eaav7893 (2019).

67. Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial neural dynamics are dominated by richly varied movements. *Nat Neurosci* **22**, 1677–1686 (2019).

68. Schurger, A. & Uithol, S. Nowhere and Everywhere: The Causal Origin of Voluntary Action. *Rev.Phil.Psych.* **6**, 761–778 (2015).

69. Raut, R. V. *et al.* Global waves synchronize the brain's functional systems with fluctuating arousal. *Science Advances* **7**, eabf2709 (2021).

70. Park, H.-D. *et al.* Breathing is coupled with voluntary action and the cortical readiness potential. *Nature Communications* **11**, 1–8 (2020).

71. Heilbronner, S. R. & Haber, S. N. Frontal Cortical and Subcortical Projections Provide a Basis for Segmenting the Cingulum Bundle: Implications for Neuroimaging and Psychiatric Disorders. *J Neurosci* **34**, 10041–10054 (2014).

72. Huang, Z., Liu, X., Mashour, G. A. & Hudetz, A. G. Timescales of Intrinsic BOLD Signal Dynamics and Functional Connectivity in Pharmacologic and Neuropathologic States of Unconsciousness. *J. Neurosci.* **38**, 2304–2317 (2018).

73. Palva, J. M. *et al.* Neuronal long-range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws. *Proceedings of the National Academy of Sciences* **110**, 3585–3590 (2013).

74. Hill, A. T., Clark, G. M., Bigelow, F. J., Lum, J. A. G. & Enticott, P. G. Periodic and aperiodic neural activity displays age-dependent changes across early-to-middle childhood. *Dev Cogn Neurosci* **54**, 101076 (2022).

75. Wiest, C. *et al.* The aperiodic exponent of subthalamic field potentials reflects excitation/inhibition balance in Parkinsonism. *eLife* **12**, e82467 (2023).

76. Chaudhuri, R., He, B. J. & Wang, X.-J. Random Recurrent Networks Near Criticality Capture the Broadband Power Distribution of Human ECoG Dynamics. *Cereb Cortex* **28**, 3610–3622 (2018).

77. Poil, S.-S., Hardstone, R., Mansvelder, H. D. & Linkenkaer-Hansen, K. Critical-state dynamics of avalanches and oscillations jointly emerge from balanced excitation/inhibition in neuronal networks. *J. Neurosci.* **32**, 9817–9823 (2012).

78. di Santo, S., Villegas, P., Burioni, R. & Muñoz, M. A. Landau–Ginzburg theory of cortex dynamics: Scale-free avalanches emerge at the edge of synchronization. *Proceedings of the National Academy of Sciences* **115**, E1356–E1365 (2018).

79. Yellin, D., Siegel, N., Malach, R. & Shriki, O. Modulation of proximity to criticality enhances slow activity fluctuations during free recall. 2023.02.24.529043 Preprint at https://doi.org/10.1101/2023.02.24.529043 (2023).

80. Ikeda, A. *et al.* Dissociation between contingent negative variation (CNV) and Bereitschaftspotential (BP) in patients with parkinsonism. *Electroencephalography and Clinical Neurophysiology* **102**, 142–151 (1997).

81. Ikeda, A. *et al.* Dissociation between contingent negative variation and Bereitschaftspotential in a patient with cerebellar efferent lesion. *Electroencephalography and Clinical Neurophysiology* **90**, 359–364 (1994).

82. Inagaki, H. K. *et al.* Neural Algorithms and Circuits for Motor Planning. *Annual Review of Neuroscience* **45**, 249–271 (2022).

83. Inagaki, H. K. *et al.* A midbrain-thalamus-cortex circuit reorganizes cortical dynamics to initiate movement. *Cell* **185**, 1065-1081.e23 (2022).

84. Zhou, S. & Buonomano, D. V. Neural population clocks: Encoding time in dynamic patterns of neural activity. *Behavioral Neuroscience* **136**, 374–382 (2022).

85. Jazayeri, M. & Shadlen, M. N. A Neural Mechanism for Sensing and Reproducing a Time
Interval. *Curr Biol* **25**, 2599–2609 (2015).
86. Broday-Dvir, R. & Malach, R. Resting-State Fluctuations Underlie Free and Creative Verbal
Behaviors in the Human Brain. *Cerebral Cortex* **31**, 213–232 (2021).
87. Soon, C. S., He, A. H., Bode, S. & Haynes, J.-D. Predicting free choices for abstract
intentions. *PNAS* **110**, 6217–6222 (2013).
88. Alexander, P. *et al.* Readiness potentials driven by non-motoric processes. *Consciousness*
and Cognition **39**, 38–47 (2016).
89. Jung-Beeman, M. *et al.* Neural Activity When People Solve Verbal Problems with Insight.
PLOS Biology **2**, e97 (2004).
90. Gelbard-Sagiv, H., Mukamel, R., Harel, M., Malach, R. & Fried, I. Internally generated
reactivation of single neurons in human hippocampus during free recall. *Science* **322**, 96–
101 (2008).
91. Norman, Y., Yeagle, E. M., Harel, M., Mehta, A. D. & Malach, R. Neuronal baseline shifts
underlying boundary setting during free recall. *Nat Commun* **8**, 1301 (2017).
92. Gelbard-Sagiv, H., Mudrik, L., Hill, M. R., Koch, C. & Fried, I. Human single neuron activity
precedes emergence of conscious perception. *Nat Commun* **9**, 2057 (2018).
1242
1243

1244 **Supplementary Material**

1245

1246
1247 **Supplementary Figure 1: Homogeneity of fluctuation statistics across networks.** In all figures, shown are
1248 average statistics (solid line), 95% confidence interval (shaded region) and example values for different networks at
1249 different R_{EE} values. **A:** Spectral slope, as in Fig. 2C. **B:** Autocorrelation timescales, as in Fig. 2D. **C:** Coefficient of
1250 variation, as in Fig. 2E. **D:** $\hat{S}_T$ measure of slow-switching spiking variability, as in Fig. 2F.

1251

1252

1253

1254

1255
Supplementary Figure 2: Relation between autocorrelation and ramping onset & characterization of ramping onset
in simulated & empirical data. **A:** Relation between autocorrelation and slow ramping in pink noise. We simulated
pink noise with 1/f exponents ranging from 0.5 to 2.5 using the *colorednoise* Python package. We simulated 10,000
trials of 20 s of noise and aligned to threshold-crossings (threshold of 0.3, baselined using the period [-3, -2] as in
our other simulations). Left: Two examples of simulated trials with lower and higher autocorrelation (1/f exponent
of 1 and 2, respectively). Right: When aligned to threshold crossings, high-autocorrelated noise shows slow ramping
in the preceding signal, whereas low-autocorrelated noise shows little to no ramping. As autocorrelation (1/f
exponent) increases, slow ramping behavior emerges and, at least to some degree, has earlier onsets. **B:**
Autocorrelation at the cluster-level (left, autocorrelation calculated for each cluster-average firing rate and then
averaged across clusters) and single-neuron level (right, autocorrelation calculated for each neuron's firing rate and
then averaged across neurons). Autocorrelation increases with increased proportion of slow synapses. Note from the
Y-Axis that cluster-level autocorrelation is much larger than the single-neuron level. **C:** Characterization of the
onset of slow ramping simulated (left) and empirical neurons (right) (red = increasing neurons, blue = decreasing
neurons) with respect to movement onset (at t = 0 s). We used the MATLAB function findchangepts with the linear
statistical method to conduct a breakpoint analysis in the timeframe of -3.5 to -1 s to determine the onset of ramping.
(Determining the onset of a change in slope in a noisy signal is a difficult problem, and methods using means are
often biased; hence we used the linear statistical method, which takes into account both mean and slope to find a
breakpoint.) In simulated and empirical data, a range of ramping onsets are apparent, with an average ramping onset
of about -2 s.
1275

1276
1277 **Supplementary Figure 3:** Further example of raster plots & trial-averaged firing rates for empirical and simulated
1278 neurons (increasing in red, decreasing in blue).
1279

1280
Supplementary Figure 4: More examples of slow ramping decoding accuracy in simulated networks. **A:** As in Fig.
3A, decoding AUC over time for 3 other example networks with 100% fast (blue) and 50% slow (black) synapses.
B: As in Fig. 3B, temporal generalization of decoders trained at one time bin and tested on another time bin. **C:**
AUC averaged across train-test bin pairs that are the same distance from diagonal. AUC was higher when the
training bin was earlier than the test bin (i.e., for negative x-axis values) for both simulated (left) and empirical data
(right). That asymmetry is consistent with a largely monotonic drift in population firing rates away from a baseline.
D: Results of temporal generalization analysis conducted on Poisson spike-trains that ramp. In a case where ramping
is a priori known to be present, we see an asymmetry such that decoders that classify activity at a given time from
baseline (-3 to -2.6 s in this case) generalize better to the future than they do to the past, indicated by higher AUC in
the lower-triangle compared to the upper-triangle of this matrix. **E:** Pre-threshold changes in total PSP delivered to
all network neurons (EEG proxy). Lines & shaded region is grand-average mean & 95% confidence intervals across
20 simulated networks. Magnitude is smaller than the EEG proxy used in main text (PSPs onto excitatory neurons
only), but still shows an early negative deflection reminiscent of the EEG readiness potential before self-initiated
actions.