Three separable roles for context in breaking environmental symmetries for place cells

Han Yin Cheng[1], Dorothy W Overington and Kate J. Jeffery*

Institute of Behavioural Neuroscience, Division of Psychology & Language Sciences, University College London, 26 Bedford Way, London WC1H 0AP, United Kingdom

*For correspondence: Kate.Jeffery@glasgow.ac.uk (K.J.J)

Running title: Contextual symmetry-breaking in place cells

Key words: place cells, head direction cells, context, hippocampus, spatial cognition, olfaction

Conflict of interest statement: The authors declare the following competing interests: K.J.J is a non-shareholding director of Axona Ltd.

Acknowledgements

This research was funded in part by the Wellcome Trust [Grant number WT103896AIA]. For the purpose of open access, the author has applied a CC BY public copyright licenses to any Author Accepted Manuscript version arising from this submission. The work was also supported by a grant from the Biotechnology and Biological Sciences Research Council (BB/Joo9792/1) to K.J.J, and an A*STAR (Singapore) National Science Scholarship to HYC.

Author contributions: Han Cheng: Conceptualization, Methodology, Software, Investigation, Formal analysis, Data curation, Visualization, Writing – Original Draft, Review & Editing; Dorothy Overington: Conceptualization, Methodology, Software, Investigation, Formal analysis, Data curation; Kate Jeffery: Conceptualization, Methodology, Data curation, Supervision, Resources, Funding acquisition, Writing – Original Draft, Review & Editing.

The authors are grateful to Joanna Holeniewska for technical support during the project's execution.

Data and materials availability

Summarized datasets are documented in an online repository xxx, and the analysis code can be found on xxx (xxx).

Author current affiliations

[1] University of Vermont, Department of Psychological Science, 2 Colchester Avenue ,

Burlington, Vermont, VT 05401, United States, han-yin.cheng@uvm.edu

[2]The Purple Agency, ENI House, Basing View, Basingstoke RG21 4YY, United Kingdom

[3]School of Psychology & Neuroscience, University of Glasgow, 62 Hillhead Street,

Glasgow G12 8QB, United Kingdom Kate.Jeffery@glasgow.ac.uk

Abstract

We investigated how place cells resolve spatial ambiguities, such as identical compartments

and rotational symmetry, by recording from hippocampal CA1 in rats exploring a visually

ambiguous environment, the '2-box', comprising two identical, connected rectangular

compartments with directionally opposing layouts. The layouts distinguish the two ends of

each box, while the boxes themselves are distinguished by their odor context. We found that

firing field ensembles were usually oriented consistently relative to the odor cues, indicating

use of odor context to break the overall global symmetry. Some place cells also maintained

a consistent context-specific relationship to the layout cues, thus 'remapping' between

boxes. Other place cells ignored both the context and the layout cues but maintained the

same relationship to walls of given orientations, thus repeating their fields in the two boxes.

No cells ignored orientation altogether by rotating independently of the ensemble. And

finally, sometimes the whole ensemble rotated its firing *en bloc*, decoupling from the odor

context cues. To explain these observations we propose that odor establishes a context

signal having three simultaneous but separable roles, to: (i) orient the global directional

signal via an association with layout cues, (ii) gate the spatial inputs onto place cells, and (iii)

organize the configural association of head direction with layout. This configuration can in

turn regenerate the context signal if the original sensory input becomes unavailable,

explaining how spatial firing patterns originally established by the odor context can become

independent of it. We speculate that these interactions occur in retrosplenial cortex.

Significance statement

Can the hippocampal place cells resolve spatial ambiguities such as identical compartments

and rotational symmetry? We found that place cells used context information in conjunction

with the environment structure. The pattern of findings suggests three simultaneous but

separable roles for the context cues: to orient the overall pattern, to position place fields, and

to guide formation of stable associations between viewpoints and head direction. This

network of interactions may be located in a multi-sensory cortical region such as

retrosplenial cortex.

Introduction

Place cells fire in specific locations in an environment, forming the neural basis for the

brain's internal representation of space ('cognitive map'; O'Keefe and Nadel, 1978). The

spatial pattern of place cell firing locations (place fields) is unique to each environment.

Given this, how does the system distinguish environments that have repeated identical
elements, and locate the animal within each one?

We have been exploring this issue using environments possessing an inherent symmetry,
broken only by sensory cues of interest. Symmetry refers to a system's property of
invariance under a transformation. For a spatial environment, symmetries of relevance are
translational, rotational and scale (Jeffery, 2023). If place cell firing patterns break these
symmetries (respond differently in the environments), this indicates that these cues are used
by the network, and suggests how (for orientation, positioning etc). The emerging picture so
far is as follows. First, the cells distinguish identically shaped and oriented environments
using context cues such as odor (Anderson and Jeffery, 2003): essentially breaking a
translational symmetry, where odor is the dimension of translation. By contrast, they do not
readily do this using path integration (movement-tracking; Spiers et al., 2015; Grieves et al.,
2016). They do, however, use path integration to break rotational symmetry, enabling them
to, for example, distinguish one end of a rectangular compartment from the other (Taube and
Burton, 1995; Jeffery et al., 1997). They also readily use local (mainly visual) cues such as
landmarks, to do this (Muller and Kubie, 1987). This rotational symmetry-breaking is thought
to be mediated primarily by head direction (HD) cells (Taube et al., 1990; Harland et al.,
2017).

HD cells are central to the organization of these firing patterns, so we recently asked the
question of how the HD system itself solves this same problem of resolving environmental
symmetries. We found to our surprise that a subset of cells in rat dysgranular retrosplenial
cortex (dRSC), which outwardly resemble classic HD cells, are responsive to local but *not*
global rotational symmetry-breaking cues (Jacob et al., 2017; Zhang et al., 2022). This was
found by recording in the '2-box', which comprises two connected rectangular compartments
in which the visual layouts are 180 degrees reversed from each other. This arrangement
created a twofold global rotational symmetry for the whole apparatus. Additionally, if the door
and cue cards are ignored there is also a translational symmetry and a local onefold
rotational symmetry, arising from the rectangular geometry (Figure 1). We found that as the
animal crossed from one box to another through the doorway in the central dividing wall,
some cells ignored the global symmetry-breaking cues and rotated their directional firing to
remain aligned with the local visual layouts. We proposed that these layout-following
'multidirectional' cells represent a link in the chain between local visual scene-processing
and global head-direction orientation (Page and Jeffery, 2018). These experiments also
revealed that ordinary HD cells can break the symmetry of the two-compartment layout by

using the odors in the two compartments, even though odor is not in and of itself a precise
directional cue.

The question arises as to whether all this information is passed to place cells. Thus, the
present study used this same 2-box environment with its multiple symmetries to investigate
the spatial organization of place fields. Given our finding that the retrosplenial HD signal is
not unitary, but rather has cells with differential sensitivity to the global vs. local directional
cues, we wondered what place cells would do. Would they (see Figure 1):

(i)   Repeat their fields between the two compartments, showing translational
108   invariance, and sensitivity to global but not local rotational symmetry-breaking

109   (ii)   Change their firing (remap), showing sensitivity to either translational or rotational
110   symmetry-breaking cues (or both), or

111   (iii)   Rotate, showing sensitivity to local rotational symmetry-breaking cues (the visual
112   layout) but insensitivity to both the translational *and* the global rotational
113   symmetry-breaking cues.

114 Furthermore, would their responses be global (across the whole population) or partial
115 (affecting some cells only)?

116   ---
117   Figure 1 near here
118   ---

We found here that only (i) and (ii) held true; place cells repeated or remapped their fields
but never rotated them. In terms of complete vs. partial responding, place cells showed
complete (ensemble) responses to global rotational symmetry-breaking cues, but partial
responses to the local cues. Furthermore, sometimes they collectively and spontaneously
broke the global symmetry independently of the odor cues, indicating more than one way to
signal global orientation.

This behavior is not readily explained by current models of place cells orientation and
positioning. We therefore propose a new model, adapted from an existing 'contextual gating'
model, that could explain these findings and map them to known circuit properties. It
proposes that context plays three different roles in this setting, by organizing linkages
between directional, positional and viewpoint signals that enable symmetry-breaking.

Materials and methods

Subjects

Five Lister Hooded rats were used for place cell recordings. All animals except one were naïve to the experimental apparatus prior to any recordings. The remaining animal had been exposed to the experimental apparatus as part of a behavioral experiment before entering the current experiment.

The animals were housed individually and maintained on a 12hr/12hr light/dark cycle. A week after surgery and throughout the whole experimental period, all animals were food restricted and their body weight maintained at 90% of their free-feeding body weight. All procedures were done in accordance with the restrictions and provisions listed in the Animals (Scientific Procedures) Act (United Kingdom, 1986) and European Communities Council Directive of November 24,1986 (86/609/EEC)).

Microdrive implantation

Axona microdrives (Axona Ltd, UK) containing four drivable bundles of tetrodes were prepared for implantation in rats. Each tetrode was made from four 25μm polyimide coated 90% platinum-10% iridium wires (California Fine Wire, Grover Beach, CA) twisted together and threaded through a 21G stainless steel cannula (Cooper's Needle Works Ltd, UK). Gold plating was used to reduce impedance to within the range of 180-300 kΩ (with a target impedance of 200 kΩ).

Standard procedures were used for stereotaxic implantation of microdrives into the cortex overlying the hippocampus (AP: -4.0, ML: ± 2.5). Carpofen (0.1ml of 10% v/v per 100g animal weight; Pfizer Ltd, UK) was given pre-surgically while Metacam was given post-surgically to reduce pain and inflammation.

Experimental apparatus

The animals were screened in a separate room distinct from the recording room. The screening room contained a square open box environment (120cm x 120cm) lifted off the floors by four stools, one in each corner. Strips of polypropylene sheeting were mounted onto the box wall to discourage the animals from climbing on or over the edge. Distal cues were available in the screening room.

The recording room contained the recording environment (2-box) surrounded by black floor-ceiling curtains, thus depriving animals of distal visual cues. The 2-box was constructed from

wood, painted black and lined with polypropylene sheets for the walls, and vinyl sheet for the floor. A white cue card (A4-sized; 21cm x 29.7cm) was mounted on one short end of each compartment, 'North' for one and 'South' for the other. The floors were scented at the start of every trial by wiping the vinyl flooring with a diluted (~50%) solution of either vanilla or lemon odors (Dr Oetker, Germany). Four halogen lamps were mounted onto the ceiling near each corner of the environment to provide symmetrical lighting, and white noise was played on a portable speaker mounted on the ceiling to mask extra-maze directional auditory cues. During recordings, the room lights were switched off and the halogen lights switched on.

Recordings and experimental protocol

A week following surgery, the animals began screening for place cells in the screening room. For screening and recording, the connector on the microdrive was connected to a headstage that in turn was connected via a thin flexible cable to the dacqUSB pre-amplifier and recording system (Axona Ltd, UK). The signals acquired from the tetrodes were digitized (48 kHz), amplified 10000-20000 times and high-pass filtered to detect neuronal spikes. Meanwhile, local field potential signals were amplified 2000-3000 times and low-pass filtered. The positions of the animals were tracked via a single LED on the headstage via a ceiling-mounted camera. If place cells were observed, the animals were transferred to the recording room in a black transport box after which the recording protocol commenced.

Rats were always started in the same compartment. They were not actively disoriented before the first trial, but were mildly disoriented in a gently rotated dark box between trials. Rice pops (Waitrose, UK) were thrown into the box to encourage the animal to explore while neural recordings were conducted. In two animals (R590 and R609), vanilla flavored rice was thrown into the vanilla compartment and non-flavored rice was used in the lemon compartment, to maximize the possibility for context discrimination. A single recording session consisted of five trials in sequence (Figure 2A): (i) baseline trial, (ii) rotation trial, (iii)-(iv) two single compartment trials, and (v) a second baseline trial. The rotation trial (90º or 180º) was to ensure that the cells were using the box cues and not uncontrolled distal cues, and the second baseline trial was to check the stability of the place cells. All two-compartment trials ranged from 15 to 25 min while the single compartment trials lasted for 10 min. In between trials, the animals were removed from the environment, placed back into black transport box and disoriented. The environment was cleaned of any urine or feces, the odors reapplied, and the box rotated to the appropriate configuration. At the end of a recording session, the screw on the microdrive was turned one-eighth of a turn (~25μm) to find new cells for the next recording session.

195   Data were analyzed as follows.

196   *Behavior analysis*

197   To examine for any bias in the animals' preference for either compartment, a preference
198   index was calculated as follows:

199
$$Preference\ Index = \frac{T_V - T_L}{T_V + T_L}$$

where T_V and T_L are time the animals spent in the vanilla and lemon compartments,
respectively. PI values > 0 indicate a preference for the vanilla compartment while a PI < 0
indicate a preference for the lemon compartment. A value of 0 will indicate no preference for
either compartment. The preferences for odors were only examined for two-compartment
trials.

Spike sorting

After recording, neuronal spikes were sorted using graphical cluster-cutting software Tint
(Axona, UK) with the help of clustering algorithm KlustaKwik 3.0 (Kadir et al., 2014).
Recording data obtained across trials in the same session were loaded into Tint together for
clustering. The resulting clusters were manually refined by examining their waveforms and
cross-correlograms, and merging them if they occupied similar spaces in the cluster space
and did not spike within the 2 ms refractory period. The sorted spikes were further analyzed
offline in Matlab (MathWorks).

Place cell identification

Data from potential single units were preprocessed and analyzed in Matlab 2019a/b. First,
positional data and spikes were speed-filtered. Spikes from epochs with an instantaneous
speed < 5 cm/s were removed to exclude non-local spiking (replay events). Positional data
and spikes with instantaneous speed > 100cm/s were considered to be tracking errors and
removed.

To ensure good unit isolation, any units with more than 0.2% of spikes falling within an inter-
spikes interval < 2ms and mean firing rate < 0.1Hz in both baseline trials were excluded.
Units were considered to be putative pyramidal cells if the waveform peak to trough time >
250 μs and their mean firing rate was between 0.1 and 5 Hz in both baseline trials.

After identification of putative pyramidal cells, rate maps were constructed to examine for
spatial firing. Rate maps were generated in the same manner as described in Leutgeb et al.,

(2005), but with slightly different parameters. Positional data were divided into 3x3 cm bins,

and spikes falling within each bin were accumulated accordingly into spike maps, and

position points similarly as dwell maps. Positions and spikes were smoothed separately with

a Gaussian kernel (smoothing factor = 5 cm) centered on the bin. Any bins > 9 cm distant

from the current bin were given a weight of 0. Bins in which the animals spent less than 0.1s

were designated as unvisited. The smoothed rate maps were then generated by dividing the

smoothed spike maps by the smoothed dwell maps.

Spatial firing was then examined using the smoothed rate maps. A place field was defined

as region of at least 16 contiguous pixels with firing rates > 40% of the peak firing rate and

peak firing of > 1Hz. Putative pyramidal cells with fields that had summed pixels > 50% of

the dwell map pixels were considered 'messy' and excluded from further analysis.

To further remove cells with non-spatial firing, we calculated the spatial information content

of each cell (Skaggs et al., 1992) and compared that to a shuffled null distribution. The

shuffled null distribution is generated by circularly time shifting the spikes by a random time

interval between 20s and the duration of the session minus 20s, for each round of the

shuffle. This maintains the temporal relationship of the spikes but dissociates the spikes from

the animal's position. For each round of the shuffle, we generated a rate map based on the

new spike positions and calculated the spatial information content. This process was

repeated 100x for each cell, and the cell was considered to be spatially modulated (i.e.

having a place field) if the cell's spatial information content > 95th percentile of the shuffled

null distribution.

For visualization purposes, spikes were plotted on positional data for each trial to create a

spike plot. Paths of the animals in the lemon compartment were plotted orange, and in the

vanilla compartment in brown. Spikes of neurons from the corresponding time point were

overlaid as red dots.

Null distributions

To determine the chance level for which a place field is in a specific location between two

correlation comparisons, null distributions were generated. To do this, randomly selected

pairs of rate maps from the entire pool were correlated, and this procedure repeated 10,000

times. The 95th percentile of the distribution of the resulting values was used as a threshold

for the probability a place cell could have a place field in the exact same location by chance.

Correlation values above threshold indicate similar rate maps while correlation values at or

below threshold suggest dissimilar rate maps and/or global remapping.

Rate maps correlation

To examine how similar or different spatial firing was across trials or compartments,
Pearson's correlations on the smoothed rate maps were carried out between two trials or
two compartments respectively. Before correlation, all rate maps were rotated to the
appropriate orientation (either same orientation or 180º rotation depending on the question
to be answered). If the rate maps were found to be of different sizes (due to unvisited
regions), the smaller rate map was padded to the size of the larger rate map with NaN in all
dimensions. The equivalent spatial bins between the two rate maps were then correlated on
a bin-by-bin basis. The threshold correlation value for what constituted a similar spatial map
depends on the null distributions generated for each type of comparison. For the entire 2-box
taken as a whole, the null distribution threshold was 0.37, while for each sub-compartment
the threshold was 0.46.

Duplicate cell detection/removal

To avoid over- or under-estimating the proportion of place cells that could display local or
global encoding, place cells that were recorded over days were identified systematically
using information about the waveform and the place fields. This method assumed that place
cells in the two-compartment context box are stable and do not remap over days.

Place cells that displayed similar spatial firing were first identified by correlating the rate
maps of all cells recorded from the same tetrode between consecutive recording sessions.
Cell pairs with spatial correlation > 0.375 were then examined for similarities in waveforms
using Tolias distances as a measure for waveform similarities (Tolias et al., 2007; Duvelle et
al., 2019). Specifically, Tolias distances can be described using two separate metrics –
Tolias distance 1 and Tolias distance 2 respectively. Tolias distance 1 captures the
differences in waveform shapes and can be calculated by scaling one of the two waveforms
to be compared by a factor α. After finding the best α fit for each channel, Tolias distance 1
can then be calculated by looking at the normalized Euclidean distance between the two
waveforms. Tolias distance 2 examines the differences in amplitudes across the four
channels. If the same cell was recorded over multiple days with minimal changes in
waveforms, both Tolias distances will return small values.

We next sought to determine an objective threshold that would indicate if the two waveforms
are similar or different. To do so, we fitted a linear discriminant analysis classifier using the
Tolias distances calculated from experimentally defined *same cells* or *different cells*. *Same*

cells were defined as the Tolias distances obtained by comparing the waveforms of cells
determined to be the same (via spike sorting) across the two different two-compartment trials
(baseline 1, rotation and baseline 2) within a recording session. Meanwhile, *different cells*
were defined as the Tolias distances obtained by comparing the waveforms of cells recorded
in the very first and the very last session of an animals. Separate linear discriminant analysis
classifiers were generated for each animal under the assumption that each animal (and each
microdrive) has its own set of electrical noise during recording.

Tolias distances were then calculated for cells recorded on the same tetrode on consecutive
sessions, and the posterior probability of cell pairs having similar waveform ('*same cell*')
predicted from the trained classifiers. Spatially similar cell pairs were considered to be the
same if the trained classifier predicted a posterior probability > 0.5. Occasionally, two cells
recorded in the first session mapped to a single cell in the next session ('convergence'), or
alternatively a single cell in the first session mapped to two different cells in the subsequent
session ('divergence'). In these cases, If the difference in the posterior probability between
the two cell pairs exceeded 0.1, the cell pair with the higher posterior probability was
considered to be the same cell. Alternatively, if the difference in the posterior probability was
less than 0.1, the cell pair with the higher spatial correlation was considered to be the same
cell.

Using this method, 104 cells were found to have been recorded in multiple sessions (2
sessions n = 70; 3-8 sessions n = 34). We removed these cells from every session except
the first. In addition, we removed place cells that did not exhibit cue control ($R < 0.37$
between rotation trials) or were unstable ($R < 0.37$ across two baseline trials).

Histology

Results

The protocol for the experiment is shown in Figure 2A. All tetrodes were situated in the CA1
area of the hippocampus (Figure 2B). Five rats were recorded for 37 sessions of five trials
yielding 437 unique place cells in the final analysis.

Figure 2 near here

Comparison of behavioral dwell time and place cell activity between compartments

First, in order to rule out potential effects of the odors, which may affect how place cells

represent the environment (Mamad et al., 2017), we examined the preference of the rats for

spending time in one compartment vs. the other and expressed it as a preference index (PI;

see materials and methods, behavior analysis). We found a slight preference across all

animals for the vanilla compartment (Figure 3A; one sample t-test $t(110) = 3.59$, $p < 0.001$).

This declined over trials: a repeated-measures ANOVA found a main effect of trials [$F(2,34)$

$= 6.55$, $p = 0.002$] with a significantly greater preference on the first trial than subsequently

(baseline 1 vs rotation trials: $t(36) = 3.08$, $p = 0.004$; baseline 1 vs baseline 2 trials: $t(36) =$

3.33, $p = 0.002$) while the latter two trials did not differ significantly ($t(36) = 0.50$, $p = 0.62$).

Thus, the vanilla preference slowly diminished over the trials such that by the rotation trial,

the animals displayed no preference and explored both compartments equally.

----
335   Figure 3 near here

We next compared the properties of CA1 place cells between the two boxes in Baseline

trial 1, when this preference was the most apparent. Place cell activity did not differ between

the boxes (Figure 3B) either in the number of active cells ($X^2(1, N = 322) = 0.075$, $p = 0.78$)

or the number of place fields ($X^2(1, N = 670) = 0.59$, $p = 0.44$). There were also no

differences in firing statistics (Figure 3C; Area: *t* test $t(668) = -0.22$, $p = 0.83$; Peak firing rate:

$t(668) = 0.14$, $p = 0.89$; Mean firing rate: $t(668) = 0.037$, $p = 0.97$). Thus, overall there was

little to distinguish the representations of the two boxes other than firing field locations, as

analyzed below.

Some place cells repeated their fields, most remapped and none rotated independently

We then looked at the spatial pattern of place cell firing in the two boxes, correlating the rate

maps on a pixel-by-pixel basis under either translational or rotational transformation, or

between the same box in different trials. The remapping correlation threshold was set to 0.46

based on the 95th percentile of a shuffle (see Methods), and we evaluated the firing patterns

based on the three predictions outlined in Figure 1. Note that sometimes the whole firing

pattern rotated 180 degrees independently of the box (examined in more detail below) and

so we corrected for this before comparing the firing patterns, as described in the Methods.

A small but substantial proportion of place cells (~13.5%) repeated place fields between the

boxes (see Figure 4 for an example), with average translational correlations of 0.66 ± 0.02

and rotational correlations of -0.12 ± 0.02. [comparing correlations for same-box, translational comparison and rotational comparison]. This proportion was significantly greater than chance (X^2(1, N = 437) = 65.53, p < 0.001), and suggests spatial encoding independent of the local visual environment. The number of place cells with repeating place fields did not appear to change as a function of experience, and they could be recorded throughout the recording sessions.

Figures 4 and 5 near here

The majority of place cells (83%) had fields that remapped between the boxes (Figure 5A and B), with a low correlation for both translational and rotational comparisons (translational correlation of 0.041 ± 0.009 and a rotational correlation of -0.004 ± 0.009), but a high correlation for repeated trials in the same box (0.68 ± 0.009), which is greater than the 2-box remapping threshold of 0.37, indicating stable recording and stable encoding.

A small population of cells (~3.2%) had rotated place fields, but this was not significantly different from the chance level of 5% (based on the 95th percentile threshold from the null distribution; X^2(1, 437) = 3.06, p = 0.08). Visual inspection showed these to be spurious correlations from cells having very few spikes in one of the boxes.

We examined whether place cells generated their box-specific firing patterns even in single boxes when the central door was closed. To do so, we aligned the boxes according to the visual scene and then correlated the immediately preceding rotation-trial maps with those from the single-box closed-door trials. If place cells can distinguish the compartments during the single-compartment trials based purely on odors, same-odor maps should have significantly higher correlations than different-odor maps. This was indeed the case (Figure 5 C; Two-tailed Wilcoxon signed rank test, Z = 17.46, p < 0.001). This indicates that place cells can use odor to recognize the single compartment.

Place cells sometimes rotated their fields en bloc independently of odor

As mentioned earlier, when we examined the place cell maps within a recording session, we saw that sometimes place cells rotated their entire 2-box firing patterns 180° relative to the odors from one trial to the next (Figure 6A). We call this phenomenon 'odor-switching'.

Figure 6 near here

To detect odor-switching systematically, we correlated every pair of rate maps for the two
baseline trials, and baseline 1 with the rotation trial, in either the same orientation or with a
180° rotation: we then compared the result to a 95th percentile threshold determined using a
null distribution. When these two sets of correlations were plotted against each other
(Figure 6B) then most data points, as expected, fell within the lower right quadrant, meaning
there was a high correlation for the unrotated maps and low correlation when one of the
maps was rotated 180 degrees. However a smaller but distinct cluster showed the reverse,
occupying the upper left quadrant. Cells that showed this effect co-occurred, and never co-
occurred with those that did not, indicating rotation of the whole ensemble. This effect was
more evident at the beginning of the session and had disappeared by the last trial,
suggesting that the system had consolidated the relationship between odor and direction
(see Discussion).

Discussion (1500)

In order to investigate how place cells resolve environmental ambiguities, we recorded the
cells in a two-compartment environment, the '2-box', which has several symmetries that are
broken with experimenter-controlled cues. Global rotational symmetry is broken with odor,
while local rotational symmetry is broken with the within-box layout (the cue card and the
doorway). The experiment was motivated by the recent finding that head direction (HD) cells
in dysgranular retrosplenial cortex (dRSC) show two types of responding in this environment,
with some cells sensitive to the local and some to the global cues (Jacob et al., 2017). We
wanted to see if place cells, connected to RSC, would show this mixed pattern, expressing
information about the ambiguous cues that could be useful for cognitive mapping.

We looked for three possible response patterns (Figure 1), but found only two of them.
Specifically, we found that some place cells repeated their firing fields between the boxes,
while some altered their firing altogether (remapped). However, we saw virtually no cells that
rotated their firing to follow the rotated layouts. It thus seems that the directional ambiguity
seen in RSC has been resolved by the time information reaches the place cells. Importantly,
an additional observation was that although the orientation of the firing-fields ensemble
generally matched that of the odor cues, sometimes the whole pattern dissociated from them
and rotated *en bloc*. And finally, we observed that some types of symmetry-breaking were
complete (affected all the cells together) and some partial.

A couple of things were surprising about the above observations. First, we had expected to
see partial rotational remapping, whereby some cells might rotate their fields to follow the
rotated environment cues. This is because the directionally ambiguous signal found in dRSC

could potentially have found its way into the place cell system via the RSC-HPC interconnections; and also because a previous experiment dissociating directional and local cues did find that a subset of CA1 cells would follow the local cues (Knierim, 2002). Our observation is, by contrast, more similar to that of Fuhs et al. (2005) who found remapping but no rotation (albeit in a small sample from only two rats). It seems that no trace of the directional ambiguity remains in the place cell map.

The other surprising observation from the present study was that the place field ensemble could dissociate from the odor cues without remapping (Figure 6). The reason this is surprising is that the cells definitely could detect the odor cues, because the odors usually oriented the ensemble firing pattern, including in the single compartment trials. We had thus assumed that the remapping cells were also using the odors to distinguish the compartments and position their fields appropriately, since odors have been previously shown to influence such positioning (Anderson and Jeffery, 2003). However, that the ensemble would sometimes decouple from the odors and rotate *en bloc* suggests that odor-induced remapping was not at play here (or at least not always).

How, then, did the remapping cells distinguish the compartments in order to remap, rather than rotate? The answer might lie with the only other symmetry-breaking information available, which is the relationship of the visual scene, arising from the layout, to global direction. If we suppose that in the lemon compartment the cue card is 'North' and the doorway 'East', then in the vanilla compartment the cue card is 'South' and the doorway 'West'. Cells could thus, in principle, distinguish the compartments if they could associate these scenes with the global directional signal. This notion is unpacked, below.

Boundary vectors and context gating

An association between global direction and local environmental cues was earlier proposed by Burgess and colleagues, to explain why place fields 'follow' environment walls when these are moved relative to each other (O'Keefe and Burgess, 1996; Hartley et al., 2000; Barry et al., 2006). In their 'boundary vector' (BV) model (Figure 7A), each place cell is driven by inputs from some (not all) of the walls located at given allocentric directions. The BV model can explain the behavior of the cells that repeated their fields, because there are two 'East' and two 'West' walls in the 2-box.

Figure 7 near here

The BV model does not explain all the results in the 2-box, however, because it predicts that

every place cell should repeat its fields (except at the doorway) and thus does not account

for the remapping cells. To explain remapping, such as when environment odor or color is

changed, we have previously extended the BV model to include contextual cues (Jeffery et

al., 2004; Hayman and Jeffery, 2008; Figure 7B). In our 'contextual gating' (CG) model, the

subset of BVs that drives a place cell is selected (gated) by the extant context cues. The CG

model can thus explain the remapping cells in the 2-box. The co-occurrence of remapping

and repeating cells can further be explained by assuming that repeating cells receive un-

gated BV inputs while the remapping cells receive context-gated ones.

However, the CG model does not explain the results of our experiment, in which we saw

occasional decoupling of the place field ensemble from the odor context, because when this

happens the BVs driving the remapping place cells have dissociated from their contexts, so

how are these cells able to fire in their usual locations?

We come back to this issue in a moment. First, however, we consider the issue of how head

direction is determined in the 2-box, given that this environment lacks a global polarizing

visual cue.

Environmental determination of the HD signal

The BV model and its CG variant assume the presence of a unitary global directional signal.

It is notable that, as mentioned earlier, we never saw cells rotate to follow the environment

layout cues independently of the ensemble, so we assume (as most models do) that a

unitary HD signal is passed to all the place cells together. In most experiments, directional

information is provided to HD cells by a single polarizing cue such as a cue card on the wall

of the recording chamber, but in our 2-box there are two such cards, in opposing directions.

We broke this twofold structural symmetry by scenting the boxes with different odors: we

know that HD cells are able to use this sometimes, based on direct observation in RSC HD

cells (Jacob et al., 2017) and also inferred here from the ensemble behavior of our place

cells. But how does odor orient the HD signal, given odor's inherent non-directionality? Also:

given this role of odor in orienting the signal, how is it that the signal sometimes dissociates

from the odors?

In fact, there *is* actually directional information provided by the odors, since the lemon and

vanilla compartments are spatially separated and thus could in principle define an axis

across the space. However the place cell patterns persisted even in the single

compartments, so this seems unlikely (though not impossible). An alternative explanation,

which we prefer, is that odor can drive the head direction signal by forming an association
with the visual scene perceived at a given facing direction (Figure 7C). Thus, if (for example)
the 'North' HD cells are active when the rat is in the lemon box and facing the cue card, then
the combination of lemon odor plus cue card scene could associate with this subset of cells
by Hebbian learning, such that in future, that combination of odor and visual scene could
reactivate those HD cells. With the 'South' HD cells, although the visual scene is the same,
the presence of the other (vanilla) odor forms a distinguishable configuration that prevents
confusion.

Another way of thinking about this is to formulate it in the standard ring attractor framework
within which theoretical discussions of HD cells are usually situated. By this view, the two
identical cue card scenes drive the HD ring attractor at two different places180 degrees
apart, North and South; the odors break this ambiguity by selectively associating the cue
card scene inputs to the 'North' or 'South' cells, as appropriate for that box, adding additional
drive to one or other of the ring attractor locations and thus breaking the activation
symmetry. This function for the odor cues is analogous to the gating role proposed for
context in the CG model of place cell positioning.

Three roles for context

By the above proposition, context has two separable roles: one in which it associates with
visual scenes to enable retrieval of heading direction (gating the ring attractor, as just
described), and one in which it interacts with the BV inputs and controls which ones drive the
place cells (gating the BVs, as described earlier). This can explain how it is that the place
field ensemble can act coherently in the directional domain (all cells orient together) but
incoherently in the spatial domain (yielding both repeating and remapping cells). It does not,
however, explain how the whole ensemble can decouple from the odor cues and rotate,
because according to the foregoing, odor is the source of the context signal that organizes
both the HD cells *and* the BVs.

We can explain this decoupling, however, if we propose a third role for context, which is to
organize formation of a linkage between visual scene and HD cells (Figure 7D), again by
Hebbian coactivation. We assume that during learning, simultaneous presence of a visual
scene, an HD cell signal and the context cues causes formation of a three-way connection
such that activation of two of the elements can subsequently retrieve the third, again by a
process of pattern completion. This is shown by the blue arrows in Figure 7E. The
consequence is that after learning, if odor inputs are weak or absent, the co-occurrence of a

visual scene plus previously associated HD activity can regenerate the context signal and
gate the BVs appropriately.

This provides an explanation for why the whole ensemble sometimes rotates independently
of the odor cues and yet the place cells retain their usual relative firing locations, as shown
on the right in Figure 7E. Suppose the odor cue inputs occasionally drop out: in this case the
HD ring attractor is driven in two places, and presumably randomly settles on one or other
orientation by chance. Having done this, the learned association of visual scene with HD
orientation reactivates the context signal, substituting for the odor cues, and then drives the
BVs in the usual way. Thus, all the place fields are correctly positioned relative to each
other.

Anatomical locations of the interactions

We can speculate about the possible location of these processes based on the known
anatomy and physiology of the region. Visual scene information from visual cortex may
possibly pass through a scene-selective region such as parahippocampal cortex. Olfactory
or other context information comes via the peripheral senses to a multimodal sensory area,
which could be retrosplenial cortex or even hippocampus. Head direction information comes
from an interaction of cortical and subcortical brain regions that combine self-motion
information with static environmental information.

Where do these signals converge and interact? We propose that retrosplenial cortex is the
most likely site. The Jacob et al (2017) experiment in the 2-box found cells that respond to
visual layout independently of context (the so-called 'bidirectional cells'), as well as
directional cells that are context-sensitive (the classic HD cells). RSC receives context either
directly from hippocampus or indirectly via the subiculum (see Vann et al., 2009; Mitchell et
al., 2018 for review) and receive scene information from the visual cortex directly (van
Groen and Wyss, 1992; Groen and Wyss, 2003) or from as-yet-unidentified scene-selective
regions of cortex. We propose here that this information can set the HD signal. This differs
from previous formulation of HD cells in which that signal produces a unidirectional influence
on place cells, and suggests that the construction of HD signals is, like many other signals in
the brain, the product of feed-back as well as feed-forward inputs. Recordings from these
areas as the cues are manipulated will be required to confirm the existence and determine
the formation mechanism of these configural signals.

References

Anderson MI, Jeffery KJ (2003) Heterogeneous modulation of place cell firing by changes in

context. JNeurosci 23:8827–8835.

Barry C, Lever C, Hayman R, Hartley T, Burton S, O'Keefe J, Jeffery K, Burgess N (2006)
The boundary vector cell model of place cell firing and spatial memory. Rev Neurosci
17:71–97.

Duvelle É, Grieves RM, Hok V, Poucet B, Arleo A, Jeffery KJ, Save E (2019) Insensitivity of
place cells to the value of spatial goals in a two-choice flexible navigation task. J
Neurosci 39:2522–2541.

Fuhs MC, VanRhoads SR, Casale AE, McNaughton B, Touretzky DS (2005) Influence of
path integration versus environmental orientation on place cell remapping between
visually identical environments. J Neurophysiol 94:2603–2616.

Grieves RM, Jenkins BW, Harland BC, Wood ER, Dudchenko PA (2016) Place field
repetition and spatial learning in a multicompartment environment. Hippocampus
26:118–134.

Groen T Van, Wyss JM (2003) Connections of the retrosplenial granular b cortex in the rat. J
Comp Neurol 463:249–263.

Harland B, Grieves RM, Bett D, Stentiford R, Wood ER, Dudchenko PA (2017) Lesions of
the head direction cell system increase hippocampal place field repetition. Curr Biol
27:2706-2712.e2 Available at: http://dx.doi.org/10.1016/j.cub.2017.07.071.

Hartley T, Burgess N, Lever C, Cacucci F, O'Keefe J (2000) Modeling place fields in terms of
the cortical inputs to the hippocampus. Hippocampus 10:369–379.

Hayman RM, Jeffery KJ (2008) How heterogeneous place cell responding arises from
homogeneous grids - A contextual gating hypothesis. Hippocampus 18.

Jacob P-Y, Casali G, Spieser L, Page H, Overington D, Jeffery K (2017) An independent,
landmark-dominated head-direction signal in dysgranular retrosplenial cortex. Nat
Neurosci 20:173–175.

Jeffery KJ (2023) Symmetries and asymmetries in the neural encoding of 3D space. Philos
Trans R Soc Lond B Biol Sci 378:20210452.

Jeffery KJ, Anderson MI, Hayman R, Chakraborty S (2004) A proposed architecture for the
neural representation of spatial context. Neurosci Biobehav Rev 28.

Jeffery KJ, Donnett JG, Burgess N, O'Keefe JM (1997) Directional control of hippocampal
place fields. Exp Brain Res 117.

Kadir SN, Goodman DFM, Harris KD (2014) High-dimensional cluster analysis with the
masked EM algorithm. Neural Comput 26:2379–2394.

Knierim JJ (2002) Dynamic interactions between local surface cues, distal landmarks, and
intrinsic circuitry in hippocampal place cells. J Neurosci 22:6254–6264.

Leutgeb JK, Leutgeb S, Moser MB, Moser EI (n.d.) Pattern separation in the dentate gyrus
and CA3 of the hippocampus. Science (80-) 315:961–966.

Leutgeb S, Leutgeb JK, Barnes CA, Moser EI, McNaughton BL, Moser MB (2005)
Neuroscience: Independent codes for spatial and episodic memory in hippocampal
neuronal ensembles. Science (80-) 309:619–623.

Mamad O, Stumpp L, McNamara HM, Ramakrishnan C, Deisseroth K, Reilly RB, Tsanov M
(2017) Place field assembly distribution encodes preferred locations.

Mitchell AS, Czajkowski R, Zhang N, Jeffery KJ, Nelson AJD (2018) Retrosplenial cortex and
its role in spatial cognition. Brain Neurosci Adv 1:doi: 10.1177/2398212818757098.

Muller RU, Kubie JL (1987) The effects of changes in the environment on the spatial firing of
hippocampal complex-spike cells. J Neurosci 7:1951–1968.

O'Keefe J, Burgess N (1996) Geometric determinants of the place fields of hippocampal
neurons. Nature 381:425–428.

O'Keefe J, Nadel L (1978) The Hippocampus as a Cognitive Map. Clarendon Press.

Page HJI, Jeffery KJ (2018) Landmark-based updating of the head direction system by
retrosplenial cortex: A computational model. Front Cell Neurosci 12:1–17 Available at:
https://www.frontiersin.org/article/10.3389/fncel.2018.00191/full.

Skaggs WE, McNaughton B, Gothard KM, Markus EJ (1992) information theoretic approach
to deciphering the Hippcampal code. Adv Neural Inf Process Syst 5:1030–1038.

Spiers HJ, Hayman RMA, Jovalekic A, Marozzi E, Jeffery KJ (2015) Place field repetition
and purely local remapping in a multicompartment environment. Cereb Cortex 25:10–
25.

Taube JS, Burton H (1995) Head direction cell activity monitored in a novel environment and
during a cue conflict situation. J Neurophysiol 74:1953–1971.

Taube JS, Muller RU, JB JR (1990) Head-direction cells recorded from the postsubiculum in
freely moving rats. II. Effects of environmental manipulations. J Neurosci 10:436–447.

Tolias AS, Ecker AS, Siapas AG, Hoenselaar A, Keliris GA, Logothetis NK (2007) Recording
chronically from the same neurons in awake, behaving primates. J Neurophysiol
98:3780–3790 Available at: http://www.physiology.org/doi/10.1152/jn.00260.2007.

van Groen T, Wyss JM (1992) Connections of the retrosplenial dysgranular cortex in the rat.
J Comp Neurol 315:200–216.

Van Groen T, Wyss JM (1992) Connections of the retrosplenial dysgranular cortex in the rat.
J Comp Neurol 315:200–216.

Vann SD, Aggleton JP, Maguire EA (2009) What does the retrosplenial cortex do? Nat Rev
Neurosci 10:792–802.

Zhang N, Grieves RM, Jeffery KJ (2022) Environment symmetry drives a multidirectional
code in rat retrosplenial cortex. J Neurosci 42:9227–9241.

Figure legends

Figure 1 Multiple symmetries in the two-compartment context box. Left: The two-compartment context box comprises two visually identical compartments, painted grey and scented with either lemon or vanilla. The local symmetry of each sub-compartment is broken by the asymmetrically placed visual cues (cue card, plus the doorway). The global symmetry is broken by the relative positions of the two odors. Right: Three types of symmetry-breaking that could be detected by place cells (see text). (i) The place fields (ovals) could be identical in both compartments but both located at the same end, which is a translational symmetry. This would mean they do not use local odor cues nor local visual asymmetry, but are oriented by the global odor cues.(and hence follow the global environment orientation when the whole environment is rotated). (ii) The cells might remap between compartments. Here it is not possible to determine whether symmetry is locally broken by the odors (inducing contextual remapping) or by the directional cues, in which the system detects that the cue cards have different relationships to the global directional orientation. (iii) The cells might generate fields in the same local location in the two sub-compartments, ignoring both local odor context and the global odor-referenced directional cues.

Figure 2 recording protocol in the two-compartment context box. (A) The experimental protocol showing the 5-trial sequence (door was closed for trials 3 and 4). (B) Recording locations, showing an example photomicrograph, and a schematic of the sites for the 5 animals.

Figure 3 (A) Preference index of all trials (grey) and also averaged across trials for individual animals (colored bars showing mean, s.e. and inter-quartile range) showing mild preference for the vanilla compartment. (B) Comparison of the number of active cells (left) and place fields (right) in the two boxes showing no preference of the cells for encoding one vs. the other box. (C) Place field firing statistics did not differ between the two compartments.

Figure 4 Examples of a repeating cell. Plots of five trials. Note that the firing fields are located at the same end of the box in a global reference frame even though both the odors and the local visual layout are different.

Figure 5 Global place encoding in the two-compartment context box. *(*A) A cell that distinguished the compartments (remapped) by firing in only one of them, maintaining a constant within-compartment position. (B) A cell that remapped by shifting its firing location depending on the box identity. This cells thus receives two sets of positional inputs, which are selected (gated) by the odor context. (C) Quantification of remapping across the odors vs. repeated trials in the same odor (rotate trial compared with first single-compartment trial).

Bars and lines show mean, s.e. and interquartile range; dots show individual cells. The low

mean correlation for the different-odor comparison shows significant discrimination of the

compartments based on their odor and/or relative location.

Figure 6 Odor switching. (A) Firing pattern of two example cells in which the firing location

within a given box reversed its relationship to the box odors. (B) Correlations between rate

maps in the first trial vs. the second trial, made either with the maps in their canonical

orientations (unrotated maps) or with one of the maps reversed 180 degrees (rotated maps).

For the majority of fields (grey circles) the correlations were high in the unrotated condition

and low in the rotated condition, meaning that the fields maintained the original relationship

to the box configuration. For some fields (colored circles) the relationship reversed 180

degrees in the second trial such that the correlation was low in the unrotated condition and

high in the rotated condition. These fields are color-coded by session; note that the rotated

fields generally belonged to the same session, meaning that the rotation was an ensemble

behavior.

Figure 7 The logic of the configural encoding of sensory inputs driving place cells. (A)

The boundary vector (BV) model of Burgess and colleagues (Barry et al., 2006). Separate

signals about facing direction, boundaries and the distance from the boundaries combine to

form boundary vectors (signals conveying distance to boundaries located at a given

direction). When two or more orthogonal boundary vectors meet, they uniquely specify a

location in the environment. (B) The context gating model of Jeffery et al. (Jeffery et al.,

2004; Hayman and Jeffery, 2008). This extends the BV model by proposing that contextual

cues control which boundary vectors can drive a place cell. (C) The configural combining of

context and viewpoint to recover heading direction. In Context 1 (e.g., the lemon box in the

2-box) then the rat sees Viewpoint 1 when 'North' HD cells are active, and Viewpoint 2 when

'South' are active (S). In Context 2 the reverse holds. This configural signal (viewpoint +

direction) can then input to boundary vectors. (D) The configural encoding of viewpoint and

direction to recover context. A direction signal, which may have come from the process in

(C) or may alternatively have come from spontaneous jumping of the HD network to the

opposing orientation, combines with the viewpoint. If 'North' (N) HD cells are active when the

rat sees Viewpoint 1 then it must be in Context 1, etc. This information can then be used to

gate the boundary vectors driving the place cells, retrieving Context 1 or Context 2 pattern

as appropriate. (E) How the foregoing scheme explains how ensemble rotation of the place

fields decouples the firing patterns from odor. The diagrams show hypothetical changes in

information flows in the 2-box depending on whether or not the odor context cues are driving

HD cells. Left: when they are, they combine configurally with viewpoint, enabling the HD

system to infer facing direction using just the environmental cues, and in turn driving the

boundary vector cells. These odor cues simultaneously also gate the boundary vectors so

that the place fields are appropriately positioned in the two boxes (with the gated cells

remapping). This is the canonical place field alignment that remains consistent with the odor

cues most of the time. A consequence of this process is that the viewpoint and HD signal,

which have been coactive, become configurally combined with the context (dotted lines),

such that they also can gate the boundary vectors. Right: Occasionally the odor cues are

weak. The HD system, lacking odor context inputs, chooses randomly from the two attractor

states linked to that viewpoint, meaning that sometimes it aligns inconsistently with respect

to the odor. However it can combine with viewpoint cues to recover a context signal, as in

(D). This can then gate boundary vectors.

← 120 cm →

Vanilla

Lemon

(i)

(ii)

(iii)

Figure 1

Figure 2

Figure 3

Figure 4

A   Active in one box only

B   Active in both boxes but in different places

C

Figure 5

A

B

Figure 6

Figure 7