Title: The developmental gene *disco* regulates diel-niche evolution in adult moths

Authors: Yash Sondhi1[1,2] (ORCID:0000-0002-7704-3944), Rebeccah L. Messcher[2], Anthony J.
Bellantuano[1], Caroline G. Storer[2], Scott D. Cinel[2], R. Keating Godfrey[2], Deborah Glass[4], Ryan A. St
Laurent[2,6], Chris A. Hamilton[5], Chandra Earl[7], Colin J. Brislawn[2], Ian J. Kitching[4], Seth M. Bybee[3], Jamie
C. Theobald[1], Akito Y. Kawahara[2]

Author Affiliation:

[1]Department of Biology, Florida International University, Miami FL 33174 USA

[2]McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of
Florida, Gainesville, FL 32611 USA

[3]Department of Biology, Monte L. Bean Museum, Brigham Young University, 4102 Life Science
Building, Provo, UT 84602, USA

[4]Natural History Museum, Cromwell Road, London SW7 5BD, UK.

[5]Department of Entomology, Plant Pathology & Nematology, University of Idaho, Moscow, ID 83844,
USA

[6]Department of Entomology, Smithsonian Institution, National Museum of Natural History,
Washington, D.C., U.S.A

[7]Bernice P. Bishop Museum, Honolulu, HI 96817, USA

Corresponding author: Yash Sondhi, yashsondhi@gmail.com

Keywords: circadian; development, diel-niche; gene expression; insects; Lepidoptera; light cycle; RNA-
seq transcriptome, transcription factor

Significance: Insects have been longstanding models for understanding circadian and sensory
biology. The diversity of diel-activity patterns of insects is tremendous, yet our understanding of the
genetic and evolutionary processes leading to this diversity is limited. Light environment is a
powerful circadian entrainer that drives diel-niche and sensory evolution. To investigate the
evolutionary impact of highly shifted light environments on animals with otherwise similar
ecologies, we compared gene expression in closely related day- and night-active wild silkmoths.

Genes linked to eye development, neural plasticity and energy cellular maintenance showed
expression patterns linked to diel activity. Notably, *disco*, a gene encoding a zinc finger transcription
factor involved in fly (*Drosophila melanogaster)* optic lobe and antennal development in the pupa,
shows robust adult circadian mRNA cycling in moth heads, is highly conserved in moths, and has
additional zinc-finger domains with specific nocturnal and diurnal mutations. We hypothesize that
disco may contribute to diversification of adult diel-activity patterns in moths.
38
39

Abstract:
Circadian rhythms drive many biological patterns, such as activity periods. The temporal partitioning
that results can reduce predation, minimize competition, or enable new resource utilization. It can
also drive the evolution of sensory systems, such as the highly specialized antennae with which male
moths find mates, and the visual specializations with which butterflies escape from birds. However,
the mechanisms that underlie such niche-shift adaptations are poorly understood. Many model
organisms in circadian biology are too distantly related from one another to elucidate the genetic
drivers of diel-niche transitions. To address this, we examined gene expression patterns between two
closely related wild silk moth species (Saturniidae), that utilize similar host plants and habitats, but
have undergone temporal niche divergence. We found 200-700 differentially expressed genes,
including day upregulation in eye development and visual processing genes, and night upregulation
of antennal and olfactory brain development genes. We also found clusters of circadian, sensory, and
brain development genes co-expressing with diel-activity. Eight genes showed a clear and
statistically significant correlation between expression and diel activity in both species, and were
involved in vision, olfaction, brain development, neural plasticity, energy utilization, and cellular
maintenance. Disconnected (*disco)*, a zinc-finger transcription factor involved in antennal
development, circadian activity, and optic lobe brain development in flies, was repeatedly
recovered in multiple analyses. While *disco* mutants have circadian arrhythmia, most studies
have attributed this to improper clock neuron development, and not directly to adult circadian
maintenance. Comparing predicted 3D protein structure across moth (*Bombyx mori*) and fly
(*Drosophila melanogaster)* genetic models revealed *disco* has likely retained its developmental
function with a conserved zinc finger domain, but it has also likely gained novel functional zinc

finger domains, highly conserved in moths, but absent in *D. melanogaster.* These regions had
several mutations between nocturnal and diurnal study species that co-occur with higher levels of
predicted phosphorylation sites. Together, robust circadian gene expression, functional nocturnal
and diurnal mutations with high phosphorylation, structural and sequence conservation, leads us to
hypothesize that *disco* may be a master regulator contributing to the diversification of adult diel-
activity patterns in adult moths.

Introduction

Circadian rhythms regulate many biological processes, including day-night activity patterns.
Research to date has explored genes, circuits, and environmental cues mostly in the context of
control within a single organism (1, 2). Diel-niche stems from the concept of temporal partitioning of
activity periods, driven by the variation in resources, temperature, light level, or predation, across the
day (3–6). Periods are often binned into three distinct groups: diurnal (day active), nocturnal (night
active), or crepuscular (active at dusk, dawn, or both), and although this is useful for comparative
analysis, it is often an overgeneralization (7–11), and the evolution of diverse diel-niches across
organisms (7, 11–15) is poorly understood (16–19).
Several studies have documented sensory adaptation accompanying evolutionary diel-niche
transitions in mammals, birds, and insects (8, 18, 20, 21). Specific examples include colour vision
gene expansions in diurnal moths (22, 23), improved olfactory senses in dark-bred flies (24), and the
evolution of hearing organs in nocturnal butterflies (25). Comparisons of circadian genes across
model organisms catalogue variation at long evolutionary timescales and highlight conserved
elements (26, 27), but diel switches can occur over short timescales (28), and distantly related species
provide less insight into mechanisms that enable faster switches. Moths and butterflies (Lepidoptera)
are an ideal group to address this, as their evolutionary history is well known and there are many
diel-niche switches, often between closely related species (7, 29). They are also one of the few insect
groups, outside of *Drosophila*, in which both sensory and circadian genes have been characterized
(23, 30–35).
Wild silk moths (Saturniidae) are important models for understanding chronobiology, with
seminal experiments confirming the brain functions as the primary circadian control (36, 37). While
most are nocturnal, many fly during the day, and temporal niche switches may function as a

mechanism for reproductive isolation in sympatric species (38, 39). Saturniid phylogeny is relatively

well known, and there are several annotated genomes, chromosomal maps. Furthermore, they belong

to the superfamily Bombycoidea, which includes the domesticated silk moth, *Bombyx mori*, a key

biological model organism serving as a reference taxon for comparative genomic studies (32, 40–43).

Saturniids such as *Antheraea* are of major importance to the silk industry, and they have a well-

resolved taxonomy and well-documented life histories (38, 44–46). Despite this, the genetic elements

that control day-night activity and diel-niche evolution, in this group and insects in general, remain

largely unknown (28, 37).

We used RNA-Seq to characterize expression patterns during peaks and troughs (midday and

midnight) across two closely related wild silkmoths: the diurnal *Anisota pellucida*, and nocturnal

Dryocampa rubicunda. They are sister genera, feeding on large deciduous trees in the temperate

forests in Eastern North America, with a recent divergence in temporal niche ~3.8 Mya (47). We

sequenced head tissue at different time periods to identify expression in sensory, circadian, and

neural genes that correlate with diel activity. Expression in eight genes clearly and significantly

correlates with diel activity, with functions in vision, olfaction, brain development, neural plasticity,

energy utilization, and cellular maintenance. Of these, a single gene emerges consistently across

analyses: *disco*, which encodes a zinc-finger transcription factor. In moths, it has likely retained

functions also found in flies (eye and brain development during the pupal stage) through a conserved

zinc finger domain. But it has also gained an extra zinc-finger domain, surrounded by phosphorylated

sites and with mutations in both nocturnal and diurnal species, which may contribute to adult moth

circadian regulation.

Results

We compared gene expression across two wild silk moth species, *Anisota pellucida* and *Dryocampa*

rubicunda, whose males are diurnal and nocturnal, respectively (Fig. 1, Table 1). We generated head

(eyes, brain) transcriptomes from moths collected and flash frozen at midday and midnight, referred

to as 'day' and 'night' hereafter. Using multiple programs to assemble high-quality *de novo*

assemblies (Table 2), we characterized the level of gene (mRNA) expression with quasi-read

mapping, which can be used to correlate protein expression (48).

Day-night gene expression patterns switch between nocturnal and diurnal species

We found 350 & 393 significantly differentially expressed genes (DEGs) when comparing day and night treatments for each species (Table 3, Fig. 2A). *Anisota* had more day-upregulated genes (56%) and *Dryocampa* was slightly more night-upregulated (53%). In order to compare DEG sets between species, we mapped our DEGs to *Bombyx mori* orthologs using the Orthofinder software (49) (Supplementary Data 1). Approximately 60% of DEGs from each species had identifiable orthologs in *B. mori* (Table 3), and only a small number of DEGs (6-8 genes) overlapped between both species (Fig. 2B). We also replicated this analysis using other software to assure that our results were robust, despite different normalization methods (50). With DESeq2, we found 498 & 697 DEGs (Fig. 2C, Table 3, Supplementary Data 2), with similar *Bombyx* annotation rates (61%), although the proportion of day upregulated genes increased considerably in the *Anisota* (79%) compared to being more even split in *Dryocampa* (50%) (Table 3). The total number of overlapping genes increased (19-26, Fig. 2D) when using DESeq2. A comparison of the two methods revealed that 174 and 216 genes were shared between *Anisota* and *Dryocampa*, respectively.

Divergently expressed genes are linked to brain optic lobe, antennal and neural development

To identify important regulators involved in diel-niche evolution, we applied two filtering criteria to our gene expression data. First, we selected genes that exhibited highly significant differential expression in both species. Second, we focused on genes that displayed upregulation patterns consistent with the natural diel-activity of each species. Our rationale was that this subset of genes was more likely to contain key regulators. To compare DEG overlap between the two species, we grouped the transcripts to their matching orthologs from *Bombyx mori*; if two transcripts from different species mapped to the same ortholog, we treated them as being the same. This allowed us to examine overlapping genes between the species to see if any genes switched fold-change sign from positive to negative or vice versa. (Fig. 2, Supplementary Data 3). We found 51 overlapping DEG transcripts that mapped to 28 unique *Bombyx* genes. Nine genes showed flipped patterns of expression between the two species, and eight of them coincided with known diel activity patterns (Table 7). Examining gene ontology (GO) annotations and comparing orthologs from flybase (https://flybase.org/), we found genes linked to optic lobe and antennal development (*disco*),

locomotion and energy use (*SLC2A6, SLC17A5*), brain and neural development (*TUBG1*) and other

essential biological processes like transcription, ribosomal translation, protein processing,

mitochondrial maintenance (*RpS4, PARL, Mrps5*) and wound response (*PRP2*) (Table 7, Fig. 2B, D).

Of these, only *disco* was recovered with both methods.

Gene network analysis identifies diel activity and species-specific co-expressed clusters

Identifying highly expressed genes helps understand which genes are activated during particular

biological processes. However, determining only those that are highly expressed can often overlook

genes with important biological functions (51). We examined co-expressed genes that may be

correlated with diel-niche or RNA collection time. We used WGCNA, a weighted correlation

network analysis tool to cluster genes together based on their normalized counts (52). After

examining co-expression patterns for each species separately, we found two modules in each species

that clustered with day-night treatment (cluster-*grey60* and cluster-*tan*) (Fig. S6, Supplementary data

4). Since we were interested in species-specific differences, we reran analyses and combined reads

from *Anisota* and *Dryocampa,* using only normalized counts for genes that had valid *Bombyx*

annotations for both species. This approach narrowed our focus to 2000 genes. Among these, we

discovered two clusters (cluster-blue and cluster-turquoise) consisting of 50 genes each that exhibited

different expression patterns across species (Fig. 4).

GO enrichment of photoperiodism, circadian control, muscle and neural growth genes

We used a gene enrichment analysis to determine if gene ontology (GO) terms were significantly

overrepresented in the DEG and WGCNA sets compared to the appropriate background of GO terms.

Using TopGo, which allows custom gene sets, we found an overrepresentation of genes involved in

several biological functions (Supplementary Table 1). Those that seemed significant for diel-niche

and vision were 'response to stimulus' and 'smooth muscle control', as well as folic acid serine,

glycine and retinoic acid metabolism. We also used ShinyGo to examine WGCNA clusters

(Supplementary Data 5). We matched orthologous genes in the less duplicated, filtered

transcriptomes to obtain identifications of the most related *Bombyx* genes (Table 4). We examined

the enrichment of both tan and grey60 modules, listing the non-redundant terms using ReviGo (Fig.

S7, Supplementary Table 2, 3). Gene clusters that co-expressed in the same direction together in the

day and night treatments of both species included photoperiodism, circadian control, negative

phototaxis and nervous system development. We next checked for the enrichment of the modules that

showed species-specific patterns (blue and turquoise, Fig. S8, Supplementary Table 2). These

included genes involved in muscle proliferation and nerve growth, neural signaling, glycolysis,

oxidative stress response, and basic cellular functioning such as protein processing and

transcriptional regulation.

Day upregulation of vision genes in the diurnal moth

Since both EdgeR and DESeq2 analyses use different normalization methods and statistical model

assumptions (50, 53), we repeated enrichment analyses by combining datasets and examining genes

that appeared in both analyses. For the *Anisota,* we tested over-enrichment of a smaller subset (FC $\leq$

-5) of diurnally highly upregulated genes (Fig. 3, Table 5). We found gene enrichment for visual

perception, excretion regulation, negative gravitaxis, synaptic plasticity, along with genes associated

with other biological processes, such as RNA interference, endopeptidase activity and endocytosis. A

reduction in stringency (FC $\leq$ - 2) did not alter results considerably (Fig. S4). Night-upregulated

genes (FC $\geq$ 2) included ocellar pigment genes, eye-photoreceptor cell development, snRNA

processing, post embryonic development and neurotransmitter secretion, among a host of other

processes that may be required for growth, development and metabolism (wnt signaling,

tricarboxylic acid cycle, and cellular response to insulin, glucose transport) (Fig. S4).

Night upregulation of antennal and olfactory brain regions mushroom development genes

We repeated the same analyses for *Dryocampa* and tested highly nocturnally upregulated genes (FC $\geq$ 5).

Our results show upregulation in genes known to be associated with mushroom body development,

locomotor rhythm, synaptic growth, energy utilization (Sialin transport) and mitochondrial translation

(Fig. 3B, Table 6). A reduction in stringency (|FC| $\geq$ 2) showed entrainment of the clock cycle, and

antennal development genes. Genes associated with innate immune response, DNA repair, cell division,

histone acetylation, circadian rhythm, retinoid cycle were upregulated during the day, possibly indicating

a period of cellular repair during a time when these moths are inactive (Fig. S5).

Key sensory, circadian, eye development and behavioral genes can drive diel-niche switches

We combined results from the DEG (EdgeR and DESeq2) and gene network analyses (WGCNA) to

create a cumulative list of 1700 transcripts (Supplementary Data 5, 6). Focusing on genes that were

recovered across *Anisota* and *Dryocampa* reduced the set to 274 transcripts (Supplementary Data 7).

Because many transcripts had poorly annotated *Bombyx* hits, we improved annotations using the

program eggNog mapper (54). We tested if these genes had GO terms associated with sensory,

circadian, brain and neural development, or behavioral regulatory genes (Fig. 4, Table 7,

Supplementary Data 8). We found that several genes in each category had associated GO terms, with

a predominance of vision and brain development genes (Supplemental Table 4).

Predicted functional regions and homology patterns identified for genes of interest

We examined protein and gene evolution for a set of genes which we found were of interest based on

results from DGE, WGCNA, GO annotations (Supplemental Table 5). In order to infer functional

homology from structural conservation, we downloaded high quality genomes of Bombycoidea

moths and relatives from Darwin Tree of Life [https://www.darwintreeoflife.org/] (Supplemental

Table 6). We assigned a reference protein sequence for each gene of interest from the *Bombyx mori*

predicted proteome. We used Orth finder to identify orthologs and filtered orthogroups containing

the reference sequence (49, 55). Since we had more than one gene per species, we reduced the

number down to a single gene by choosing the highest identity sequence relative to the *B. mori*

reference sequence. We also modelled the 3D structure of *B. mori* proteins and mapped the

evolutionary conservation onto the 3D predicted structure for proteins above a certain conservation

threshold (Supplementary Data 9). These analyses predict structurally and functional conserved

regions of proteins (Fig S9, Supplementary Data 10). We repeated this analysis with 38 insect

genomes (Supplemental Table 7) and mapped evolutionary conservation onto 3D protein structure

(Supplementary Data 9,10). We include results of evolutionary conservation analyses for two

regulatory candidates (*disco* and *tk*) that showed varying levels of sequence and protein evolution

between insects and moths (Fig. S10).

Modeling predicts additional functional zinc finger domains for disco in Lepidoptera

Disco was recovered across multiple analyses. In *Drosophila melanogaster* it is known for its role in

eye development, important for circadian maintenance, and for leg and antennal appendage

formation (56–60). To determine if *disco* was conserved between moths and *Drosophila*, we

compared the primary sequence and 3D protein structure of *Drosophila* and *Bombyx mori*. In the

moth, the sequence length of *disco* was nearly double that of *Drosophila* (Fig 6A). However, a region

spanning over 100 amino acids was highly conserved, contained the zinc-finger domain important for

its function, and showed strong 3D structural conservation (Whole protein alignment RMSD(align,

super) = 37.833, 2.566, MatchAlign score (align, super) = 540 (2234 atoms), 333.1 (554 atoms) vs.

alignment of conserved region RMSD=2.699,2.566, MatchAlignScore =443(671 atoms), 351 (615)

atoms). This indicates the DNA binding function of *disco* has likely been conserved. However, an

additional ~500 amino acid region absent in *Drosophila* is highly conserved across moths, and

includes several regions predicted to be functional (Fig. 6A, Fig. S10). We hypothesize that *disco* has

a novel role in moths for diel-niche regulation. To further test this, we compared *disco* sequences

across *Anisota* and *Dryocampa* and found 23 mutations between them, three of which mapped to the

predicted functional region (Fig. 6B, Supplementary Data 10). InterProScan predicted four zinc

finger domains, three in this region, although the CATH-Gene3D databases prediction combined the

two separate domains into a single predicted domain (Fig. S11). We also found many sites (53) with

predicted phosphorylation potential, especially around the second zinc finger domain (18). Reducing

the stringency increased the total to 142, which were still enriched around the second domain.

Discussion

We identified the genetic mechanisms of diel-niche switches by examining gene expression during

the day and night in two closely related moth species. We used RNA-Seq to measure gene expression

profiles of head tissue and isolated sensory, circadian and neural processes. We found between 300

and 700 genes that significantly altered day night circadian expression patterns. The diurnal species

had enriched visual perception genes during the day and the nocturnal species had locomotor and

olfactory genes upregulated at night.

Thirty overlapping DEGs were present in both species, with some DGEs showing divergent

patterns of expression matching the species' diel-niche. Examining expression data with a sensitive

clustering analysis yielded over 170 genes that showed day-night or species specific co-expressed

clusters. We also used GO terms to search for genes associated with sensory, neural, or circadian

patterns. We found genes in each category (Supplemental Table 4) and modeled the 3D structure and

predicted functional regions for a subset (Supplemental Table 5). The divergently expressed gene

disconnected (*disco*) was implicated in vision, hearing, locomotion and brain development. Given the

multiple lines of evidence supporting the importance of this gene in regulating diurnal and nocturnal

activity in moths, we explored the evolution of *disco* by modeling fly and moth *disco* structures. This

analysis revealed novel zinc-finger conserved domains in moths, which were lacking in *Drosophila,*

surrounded by phosphorylated sites. Several mutations between *Anisota* and *Dryocampa* also

mapped to these regions, further strengthening evidence for its role in diel-niche shifts.

Visual and olfaction

Visual systems often accompany diel and photic environment shifts (18). For example, nocturnal

carpenter bees have much larger facets in their eyes than their diurnal counterparts (61). Nocturnal

moths have higher photoreceptor light sensitivity and have neurons more suited to pooling than close

diurnal relatives (62). Some of these trends seem to hold across the phylogeny, with diurnal species

evolving more complex color visual systems, often reflected in visual opsin sequence evolution (22,

23). We did not find diel-expression patterns in color vision opsins, a result corroborated by a recent

study (22). However, we discovered a cerebral opsin (ceropsin) upregulated during the day. Ceropsin

has been implicated in photoperiodism and is expressed in the brain (63).

We also found several eye development genes (*TENM2, ANKRD17, EHD4, JAK2*),

phototransduction genes (*PPAP2*, *RDH11*), and retina homeostasis, eye-antennal disc development,

and photoreceptor cell maintenance genes (*disco, glass*) (64). Surprisingly a few visual genes, such

as *garnet* and *rugose* also appeared to have different isoforms present, showing both day and night

upregulation. *Garnet* is an eye color mutant gene in flies (65) and *rugose* is implicated in retinal

pattern formation (66).

One of the classical differences between day flying butterflies and night flying moths is that

butterflies have a clubbed antennae, while moths often have a sensilla-covered antenna; likely a

function of increased investment in olfactory sensing, similarly, there is also evidence for increased

investment in olfactory sensing in the brain, with relatively larger mushroom bodies in nocturnal

species (67, 68). In our diel gene-expression dataset, we found several olfactory genes, including

those involved in odorant binding (*Obp84a,Obp58b*), pheromone response (*tk*), mushroom body

development (*DAAM2 and DST*) and antennal development (*disco*).

Hearing and mechanoreception

Many moths have active hearing organs, and these have evolved repeatedly across Lepidoptera (69), and their use varies depending on the ecological context of the moth. Several drivers for moth hearing include sexual selection and as a defense against insectivorous bats (70). The primary nocturnal group of butterflies, Hedylidae, has reverted to a nocturnal niche and regained hearing organs (25). Thus, although rudimentary vibrational hearing exists in some diurnal moths (71) it is significantly more developed and used at broader frequency ranges at night, when light is less available. While Saturniidae lack hearing organs (72) they may instead still have vibration mechanosensors that can be useful in evading predation similar to those in locusts and crickets (73–76). Indeed, we did find diel co-expression of several mechanosensory and ear development genes, including *PI4KB, KCTD15, unc-22,* and *RhoGAP92B* (77)

Brain and neural rewiring

Finding an upregulation of neural and brain development was expected since we sequenced head transcriptomes. However, we tested gene expression two days after pupal eclosion, where presumably gross structures are already developed, so it is possible these genes we found expressed regulate adult plasticity. Adult neural plasticity has been showed in Lepidoptera and other insects (78–80), and we recovered several genes linked to axon regeneration (*APOD*), central complex development (*Ten-a, TENM2, DST, ALDH3A2, OGT*), central nervous system development (*disco, RpL4*), and neuropeptide hormonal activity (*tk*). Several genes were specific to retinal ganglion cell axon guidance (*TENM2*) and mushroom body development (*DAAM2*), leading us to speculate that plasticity could occur through neural wiring and plasticity are shaping sensory adaptation. Many Lepidoptera have distinct phenotypic plasticity in wing patterns and coloration in different seasons (81), and there is some evidence that they also have seasonal plasticity in their behavior and foraging preferences (82). Research has found Lepidoptera can override their innate preferences and learning preferences for new visual and olfactory cues after eclosion (83). It is possible the diel-niche and circadian rhythms too have plasticity, there are reports of some moth species like *Hyles lineata* showing relatively labile diel-niches possibly driven by temperature and resources (84–86). These

same mechanisms allowing flexibility within a species, might be involved in diel-niche evolution

between species.

Circadian and behavioral regulators

The behavioral state of an animal to engage in any activity, e.g., feeding, flight or mating, is likely a

function of circadian and behavioral regulators that respond to certain stimuli. We found differential

expression of genes involved in locomotion (*unc-22, KCTD15,Tk*) and circadian or rhythmic

behavior (*SREBF1/SREBF2, OGT, disco, JAK2*) in both species. We also found several key circadian

regulators like *per*, *tim* although they were downregulated only in *Dryocampa*. C*lock-like* or *takeout*

another gene under circadian control, (87) was expressed in both species, although without any

significant up or down regulation. *Takeout* was moderately conserved in the moths we examined, but

had diverged considerably among other insects, to the point where orthology searches with

Orthofinder failed to recover orthologs among insects. Even among *D. melanogaster*, its closest

homologs were *Jhbp5* and *takeout* which had 21-25% sequence identity (Supplementary Data 11).

Surprising, , it's 3D structure was highly conserved (RMSD (align, super) = 2.556, 2.755,

MatchAlign score (align, super) =199 , 642.937, Supplementary Data 11, Fig. S12), highlighting an

interesting example of likely functional convergence despite primary sequence divergence.

Energy use and general cellular maintenance

In addition to the genes mentioned above, several key divergently expressed genes were involved in

energy utilization. This finding makes biological sense, as once activity is initiated, energy

mobilization and upregulation of basic cellular processes need to be maintained. Two of these genes

were *SLC175A/MFS10* and *SLC2A6/Tret1-1*, which encode different trehalose sugar transporters.

Trehalose is an important sugar present in insect hemolymph (88). Other genes found were ribosomal

(*RpL4, mRpS5*), golgi (*GCC*), and mitochondrial maintenance (*PARL/rho7*) along with

transcriptional regulation (*Spt5).*

Disco as putative master diel-niche regulator in adult moths

Many genes that we discovered likely play a critical role in maintaining sensory shifts, but in order to

identify genes that might regulate upstream, we looked for genes that were: 1) expressed in both

species, 2) showed coincident expression patterns with respect to diel-niche, and 3) played a role in

sensory and neural functioning and circadian control. Only *disconnected* (*disco*) fit these criteria.

Disco is a key developmental and patterning gene, first discovered for its role in neural migration

from the disconnected optic lobe mutant phenotype (89). This gene is now considered an appendage

development gene, with a well characterized pupal role in antennal and leg distal patterning in

Drosophila melanogaster (60). Mutants also have a disrupted circadian locomotor rhythm due to the

improper formation of neurons that express clock genes (90, 91). As a gene that is involved both in

optic lobe, antennal formation, and in neural wiring and circadian control, it is a very strong

candidate for driving behavioral diel-niche shifts and sensory adaptations (56, 57, 92). *Disco*

expression data from *B. mori* suggests strong adult head, antennal, and nucleus expression,

suggesting that it still acts as a transcription factor, and can regulate other genes (40).

Disco encodes the acid zinc finger transcription factor and is approximately 500 and 1000

amino acids in length in *Drosophila* and *B. mori,* respectively. Our modeling showed that a ~150

amino acid region is conserved structurally at the sequence level, and this region also contains zinc-

finger motifs associated with DNA binding (93). This indicates that *disco* likely has retained its DNA

binding and likely pupal and appendage patterning function in moths. We predicted the functional

regions of *disco* in *Bombyx* based on evolutionary conservation modelling, and found an additional

500 amino acids, absent in *Drosophila*, were predicted to be functional (conserved and exposed).

Domain and family level modeling predicted at least two additional zinc finger domains in this

region (Fig. S10,Fig. S11). We also found many phosphorylated sites surrounding the zinc finger

domains. Examining mutations between *Anisota* and *Dryocampa* revealed three mutations mapped to

these predicted functional regions (Fig. 6B).

Given *disco's* adult diel-specific expression in moths, additional zinc-finger DNA binding

domains, and the high number of phosphorylated sites, we propose it as a candidate transcriptional

regulator for diel-regulation in adult moths.

Study species and their evolution and variation in life histories

Anisota and *Dryocampa* are closely related and ecologically similar saturniids of the subfamily

Ceratocampinae, thought to have diverged ~3.81 Mya (47). They largely occupy the same range and

forest habitat in the Americas, feeding on large trees (*Anisota* on oaks and *Dryocampa* on maples)

where population flares cause host plant defoliation (38). Because they reportedly hybridize (94), and

pre- and postzygotic barriers are potentially ineffective, temporal partitioning may be important in

their evolutionary history.

Conclusion

394   This study provides a framework for how diel-niche evolution in Lepidoptera can occur.

395 Sensory and neural developmental genes appear to be key. We identify the pivotal role of the

396 transcription factor *disco*, uncovering a second functional region with species-specific mutations in

397 moths. Our findings provide useful targets for further genetic manipulation and highlight the insight

398 non-model systems provide to genetics and development.

Supplementary information

401   Supplementary tables and data are available for the reviewers and will be uploaded to repository

402 [xxxxx] on submission.

403   .

Author contributions

Y.S.: conceptualization, data curation, formal analysis, investigation, methodology, project

administration, software, visualization, writing–original draft. R.M: investigation, methodology, project

administration, data curation, writing - review & editing. A.J.: conceptualization, formal analysis, writing

- review & editing C.S.: conceptualization, data curation, methodology, project administration,

supervision. S.C.: formal analysis, software. K.G.: formal analysis, software, visualization, writing-

review and editing, D.G.: investigation, project administration, data curation. R.L.: conceptualization,

data curation, investigation, methodology, writing-original draft, writing -review & editing. C.H.:

conceptualization, data curation, investigation, methodology, project administration, writing - review and

editing I.K.: conceptualization, funding acquisition, methodology, resources, writing–review and editing.

C.E.: formal analysis, software, visualization, data curation, writing- original draft, writing- review &

editing. C.B.: software, validation, writing -review and editing. S.B.: funding acquisition, methodology,

validation, writing–review and editing. J.T: conceptualization, funding acquisition, supervision,

validation, writing–review & editing. A.K: conceptualization, methodology, resources, writing–review

and editing, visualization, supervision, funding acquisition

Acknowledgements

We thank Jesse Breinholt, Martijn Timmermans and Andreas Zwick for help with project
conceptualization. Kelly Dexter, Amanda Markee, and members of the McGuire Center for
Lepidoptera and Biodiversity at the University of Florida assisted with animal care and wet-lab
troubleshooting. We thank Belinda Pinto, Danielle DeLeo, Heather Bracken-Grissom, Jorge Perez-
Moreno, Zhou Lei for discussions, advice and assistance with bioinformatics pipelines and wet-lab
protocols. We thank Nicolas Alexandre, Sachit Daniel, Riddhi Deshmukh, David Plotkin, Chelsea
Skojec and Nitin Ravikanthachari for comments and feedback on the manuscript. Janelle Nunez-
Castilla, Jessica Liberles, Raghavan Venket, Yi-Ming Weng, Yu Fahong and Xiaokang Zhang
assisted with troubleshooting bioinformatic pipelines and provided analytical recommendations. The
authors acknowledge the University of Florida Research Computing for providing computational
resources and support that have contributed to the research results reported in this publication
(http://researchcomputing.ufl.edu). We acknowledge NBAF and UF-ICBR for assistance with
sequencing and generating libraries.
Funding for the project was through NSF DEB-1557007, NSF PRFB-1612862, and NSF IOS-
1750833 and NERC NE/P003915/1. The Florida International University Presidential fellowship
supported YS.

Figure Legends and Figures:

Figure 1: Nocturnal and diurnal moths on a phylogeny with RNA-seq sampling design.

(A): Collapsed phylogeny of Saturniidae, adapted to show where the two study species, diurnal
Anisota pellucida (Pink-striped oakworm moth) and nocturnal *Dryocampa rubicunda* (Rosy Maple
moth) sit. Faded numbers on the tip represent the number of genera in the tree before collapsing.
Phylogeny adapted from Rougerie et al. 2022 (**B**): Sampling design showing the number of replicates
for each species and collection period (day/night) sampled. Collection of heads was done 2-days
post eclosion at midday (sun) and midnight (moon) and tissue was flash frozen for RNA
preservation. Care was taken that the eclosed moths were exposed to a natural light cycle and red
lights were used when collecting the moths at night. Photo credits *Anisota pellucida* © Mike
Chapman; *Dryocampa rubicunda* (CC) Andy Reacgo and Chrissy Mclearan;

Figure 2: Nocturnal and diurnal species show divergent patterns of gene expression across two different analyses software (EdgeR and DESeq2)

Volcano plots and Venn diagrams showing EdgeR (A-B) and DESeq2 (C-D) results across **day vs. night** sampling times for both nocturnal and diurnal species. Venn diagrams represent the number of differentially expressed genes (DEGs) across both species with the number of common DEGs across each pair. **(A/C)**: Volcano diagrams illustrating fold change and adjusted p-values for the significant differentially expressed genes between midday and midnight samples. Circles in the top left represent the number of genes expressed in both species and the colors correspond to FC values for those genes in the volcano plots. Yellow and blue represent genes expressed only in the nocturnal or diurnal species, green/purple indicates DEGs present in both species. **(B/D)**: Only genes expressed in both species are shown and genes that display opposite trends in expression are highlighted. See Supplementary Material for a detailed list of DEGs. Positive fold change indicates night overexpression and negative fold change indicates day overexpression. Genes had FDR or adjusted p-value < 0.05 and FC > |2|. Identification of common genes was done using orthogroup clustering with Orthofinder with *Bombyx mori*. Gene names and annotations were transferred from *B. mori*.

Figure 3: Visual genes are upregulated in the diurnal species during the day, and energy utilization,
brain olfactory region and locomotion in the nocturnal species during the night.
Go enrichment of highly upregulated genes recovered from both DEG analyses coinciding with the
species. highlighting the two modules that showed species specific clustering patterns.
A: Enrichment of **day** upregulated genes in diurnal *Dryocampa*. B Enrichment of **night** upregulated
genes in nocturnal *Anisota.* Go enrichment was done using the custom ShinyGo database v0.75c
using *Bombyx* gene IDs. FDR cutoff <=0.17, and only Biological Process GO terms were selected
with Min. pathway size =1. Input genes had FC>5, padj. <0.05 and recovered both in EdgeR and
DeSeq2 were used with the background being all orthologous *Bombyx mori* genes for either species.
481

482
Figure 4: Modules of clustered co-expressed genes grouped using normalized expression
highlighting the two modules that showed species specific clustering patterns.
A: Cluster dendrogram shows WGCNA clusters. B: Shows how patterns of gene expression correlate

across samples and modules. C: Shows the normalized expression for all

genes across samples. The species show two sets of 50 genes (Blue) and (Turquoise) that have clear

species specific expression patterns. Normalization was done with DESeq2 and reads were mapped

to the more stringently filtered transcriptome. A soft power analysis was done and the picked

power=9 for the WGCNA analysis. AN: *Anisota*, DR: *Dryocampa,* Numbered D and N represent the

different samples collected at day and night time points.

Figure 5: Model for diel-niche shift from and genes of interest obtained from GO searches and
annotating divergently expressed genes. For representative purposes, homologs from *Drosophila*

melanogaster have been used, with colors in the 3d structures representing Alpha fold per-residue confidence metric (pLDDT), the range for each color is shown on the top left. Model moth *(Bombyx mori)* proteins were also modeled for a subset of proteins, see Supplementary data.

Figure 6: *Disco* has 150 amino acid long highly conserved region across *Bombyx* (1054 aa) & *Drosophila* (568 aa) involved in its role as a transcription factor, but it also has other roles in Lepidoptera due to the presence of other highly conserved and predicted functional regions, that are not present in flies, and only in Lepidoptera. 3/14 of the mutations between nocturnal to diurnal species map to these regions (>500 aa).
(A) Top: *Disco* best Uniprot hit in *Drosophila* using default settings (blastp, e-threshold 10, Auto-Blosum62). Bottom: *disco* aligned and superimposed predicted structures for the silk moth and vinegar fly. Peach structure: *Bombyx mori* alpha-fold predicted structure for *disco,* Cyan: *Drosophila melanogaster* alpha-fold predicted structure for *disco* **(B)** Top: Partial views of the Consurf predicted residues for *disco.* The views encompassing the region where 3 mutations of the 14 mutated sites between Nocturnal *Dryocampa* and Diurnal *Anisota* overlap with predicted functional residues. Bottom: Residues that were mutated between the nocturnal and diurnal species are highlighted on an overlay of the Consurf scores mapped onto the predicted alpha fold structure of disco from *Bombyx mori.* Green residues indicate mutated and predicted functional sites, red indicate mutated sites that did not have a predicted residue. **(C)** Overlap of the highly conserved region of *disco* this region includes the zinc-finger domain that is characteristic of disco transcription factor.

Supplemental Methods

Insect rearing:

Moths were reared under natural light-dark cycles at room temperature (25°C) and natural light conditions at the McGuire Center for Lepidoptera and Biodiversity (MGCL). Adults were sampled two days after eclosion at the two time points by C.H. and R.S.L.

Sampling design:

We collected 3-5 replicates (i.e., individual moths) per experimental treatment. We sampled at two time points per species at two midpoints of their circadian cycle at noon ('day') and midnight ('night') (Fig. 1, Table 1). Moths were allowed to eclose naturally and sexed, To ensure vision genes were not artificially stimulated, for collection at night we used red light, which is thought to not stimulate most Lepidoptera insect visual systems (95). We used a sterilized scalpel to remove the head, which was placed in a 1.5 ml Eppendorf tube, which was immediately placed in liquid nitrogen and stored at -80°C. It is worth noting that only males of most *Anisota* species are diurnal, including *A. pellucida*, our focal taxon. Females of all *Anisota* and both sexes of at least *A. stigma*, are nocturnal. In *Dryocampa*, both males and females are nocturnal. We therefore limited our transcriptomic work to males of *A. pellucida* and *D. rubicunda*.

RNA extraction and quality control:

Tissue was removed from liquid nitrogen and placed into a flat-bottomed cryo-tube with Trizol and 2-4 sterile, stainless-steel beads. The tissue was homogenized using a bead beater for 2.5 minutes at 1900 RPM. Beads were removed with a magnet, and chloroform added to separate phases. Isopropanol precipitated the RNA, followed by several ethanol wash steps to pellet the RNA. Quality Control (QC) was undertaken with a high sensitivity Qubit assay. Samples with yields > 20 ng/μL were cleaned using a Thermo Scientific RapidOut DNA Removal Kit (Waltham, Massachusetts, USA). Sample quality was verified using Agilent 2200 Tapestation. Since heads comprised of a relatively small amount of tissue, initial extractions using kit-based methods resulted in low yields. Several rounds of troubleshooting were required before completion of extraction using a modified Trizol based protocol with a DNA clean up step. We measured the quantity of RNA using a Qubit

fluorometer and used Agilent tapestation or Bioanalyzer to characterize extraction quality, weight,

and size distributions. Where possible, a concentration of more than 10 ng/μL was used for library

preparation. Yields below 10 ng/μL were prone to higher loss when purifying with a DNAse agent,

which often caused 20-30 % loss in RNA yield, approaching 5ng/μL is below the Qubit detection

threshold (Table S1).

Library preparation and NGS sequencing:

Libraries were prepared by sequencing cores at NBAF-Liverpool using the NEB polyA selection and

NEBNext Ultra II directional stranded kit, suitable for low input yields. 12 samples were run on one

lane of an Illumina NovaSeq using SP chemistry (Paired-end, 2x150 bp sequencing). Samples were

shipped overnight from MCGL to the NERC-NBAF after dehydrating in a biosafety chamber using

GenTegra RNA tubes.

Read trimming and cleanup:

For *Anisota pellucida* and *Dryocampa rubicunda*, trimming was undertaken by the NERC-NBAF

core and the raw Fastq files were trimmed for the presence of Illumina adapter sequences using

Cutadapt version 1.2 (96). The option -O 3 was used, so the 3' end of any reads that match the

adapter sequence for 3 bp or more were trimmed. Trimming was also done with trimmomatic. The

reads are further trimmed using Sickle version 1.200 with a minimum window quality score of 20.

Reads shorter than 15 bp after trimming were removed.

Transcriptome library sizes, QC, and PCA:

Quality Control (QC) was conducted on the trimmed reads, library size varied for each species (Fig.

S1). We examined expression data and removed genes with TPM < 1. PCA results showed that

Anisota and *Dryocampa* did not separate with diel-niche (Fig. S3).

De novo **assemblies:**

We combined reads from multiple samples and generated several reference de novo transcriptomes

using different assemblers, and then combining them, compared their quality using BUSCO scores

and number of single-copy BUSCO genes across the different versions (see Table 2 for the BUSCO

scores, redundancy and duplication for the various assemblies). We found that unfiltered assemblies

were highly redundant, but had the highest BUSCO scores. These assemblies were too

overrepresented to be used in any downstream analysis so we used Transrate, CD-HIT, MMeqs2, as

well as Transdecoder, to filter the assemblies. We chose v5 and v6 to use in downstream analysis,

listed as the last two assemblies for each species (Table 3) and repeated certain analyses with both

assemblies to see how stringent or a less stringent filter would affect the results.

Multi assembler *de novo* transcriptome assembly:

To our knowledge, there is no publicly available annotated reference genome for the two species

used in this study at the time the analyses were performed; therefore, we chose to build a de novo

transcriptome assembly with the pre-processed reads using the NCGAS pipeline, which combines

multiple assemblers and then uses an evidence-based gene modeler to decide on the final set. For

each species, we combined forward and reverse reads from all samples and normalized them using a

normalization script in Trinity v2.12.0 (97). Normalized reads were used as input to generate

assemblies using Velvet v1.2.10 (98), TransAbyss v2.0.1 (99) , SOAPdenovo v1.03 (100) with

various kmer sizes. These assemblies were combined and filtered using Evigenes v4: 2020, March

(101, 102), transdecoder and filtered with scripts provided in the NCGAS pipeline (103).

Transcriptome QC, annotation and filtering:

Further quality control and filtering was conducted using BUSCO (104), transrate (105), QUAST

(106) (v5.02), CD-HIT(v4.6.8) (107, 108)and cascaded clustering with MMseqs2 (v12) (109).

BUSCO scores were greater than 95%, but many genes had multiple versions from the different

assemblers, and because this would bias the downstream analyses we attempted to reduce the

redundancy at the cost of lowering BUSCO scores. Protein coding regions were predicted using

TransDecoder (https://github.com/TransDecoder/TransDecoder) and the cds file from was run

through Transrate (105), which further pulled contigs based on mapping rate and identified a "good"

collection of transcripts, the resulting protein-coding sequences were filtered with additional CD-HIT

(107, 108)and MMseqs2 (109) filtering. This resulted in lower BUSCO scores 75-85% but also much

lower redundancy in the gene set of about 1-3 fold. We ran the WGCNA (52) and GO enrichment

(110) analyses on the more stringent assemblies (BUSCO 75%, 2% genes with duplication) since

they are more sensitive to redundancy, but RNA-seq was run across the slightly lower stringency

analyses to be more inclusive (BUSCO~85%, 60-70% genes with duplications).

Annotation of reference transcriptomes:

Reference genomes were annotated with eggNOGmapper and Orthofinder (54, 55). We also tried

annotating the reference transcriptomes with Trinotate (111), which run against the swissprot and

pfam databases using hmmscan, blastx, blastp, signalP and tmmhmm, KEGG, and GO and eggNOG

(49, 110, 112–120). ORF predictions for orthofinder were obtained using *Antheraea pernyi* as a

model. Orthofinder was run with well-annotated Bombycidae and Saturniidae moths, *Bombyx mori*,

Antheraea pernyi, and *Antheraea yamamai*, to generate clusters. *Bombyx mori* was a useful reference

since the Orthofinder cluster for each species generated a list of mostly single-copy genes for which

more Lepidoptera annotations were available from SilkDBv3 (40). While this was not as complete as

trinotate, the annotations provided from this procedure were more accurate than Trinotate. Trinotate

annotation had many more human/ vertebrate hits than insect matches than eggNOGmapper and

Orthofinder transferred annotations, so while the files are provided as a reference, they were not used

in downstream analyses.

Differential gene expression (DGE):

Reads were mapped using Salmon (121) and differential gene expression analysis was conducted

using EdgeR (122) and DESeq2 (123). EdgeR(v 3.38.1) was used to normalize and test for

significantly expressed genes. DESeq2(v1.36.0) was also used to normalize and to generate PCAs

and other summary statistics. We used 'Day' as the treatment and 'Night' as the control for all

groups with EdgeR, although the order of fold change switched for DESeq2. The p-values obtained

were adjusted to account for multiple hypothesis testing with FDR for EdgeR and adjusted p-value

from DESeq2. Genes with FDR or adjusted p-values <0.05 and |FC| > 2 were used as a criteria to

identify significantly expressed genes.

Gene enrichment analysis:

We performed gene enrichment analysis using GO terms with the tools TopGO v2.48.0 (124) and

ShinyGo v0.75c (125) (http://bioinformatics.sdstate.edu/goc/). This analysis involved comparing the

selected genes of interest to a gene universe with GO term annotations to determine if there were

overrepresented or underrepresented GO terms. To obtain genes with GO annotations, we utilized

similarity clusters to map annotations from *Bombyx mori*.

In TopGO, we employed a more stringent approach by using filtered transcriptomes to

mitigate bias from gene duplications. We selected genes of interest based on a False Discovery Rate

(FDR) threshold of less than 0.05. We ran two different enrichment algorithms, 'classic' and 'elim',

and used two test statistics, Fisher's exact test and a Kolmogorov-Smirnov-like test. The tests were

performed for both 'Biological Process' (BP) and 'Molecular Function' (MF) categories. While we

generated tables of significantly enriched GO terms, obtaining individual genes within each group

was limited due to the constraints of TopGO with custom annotations.

For ShinyGO, we utilized the custom v0.75c which included the updated *Bombyx mori*

genome stringent. For all analysis, we used the following settings (pathway size: min.=1,

max=2000). The settings for number of pathways to show and FDR cut-offs were chosen to represent

the entire list of top 100 genes in the final datasets, although in some cases a fewer number are

represented in the visualizations. We focused on the 'Biological Process' (BP) category for the

various tests. Tables of GO terms, gene information, and graphs summarizing the significantly

enriched GO terms were generated. We repeated several different analyses **1)** A less stringent

analysis was run where genes with FC<|2| and padj. <0.05. This was useful to visualize the genes up

and downregulated functionally and the background used was all the genes recovered from the DEA

analysis that had orthologs. An FDR cut-off <0.1-0.3. For the background gene set, we used the

species' de-novo transcriptome matching *Bombyx* ortholog set.

2) A more stringent criteria was used to examine highly upregulated genes a FC<|5| and padj.

<0.05 cutoff was used, and filtering only genes that occurred in both EdgeR and DESeq2 analyses.

We selected a single representative when multiple Bombyx orthologs were recovered, using the

ortholog with the most complete annotation. An FDR cut-off <0.17 was used. For the background

gene set, we used the species' de-novo transcriptome matching Bombyx ortholog set.

3) For WGCNA clusters tan and grey60, we used their respective species background

expressed transcript set with a FDR cutoff <= 0.1. For blue and turquoise, the background of all

orthologs found in *Bombyx mori* was used. An FDR cutoff<1 was used, since smaller values

provided insufficient genes given the larger number of genes being tested.

Gene network analysis:

669 Gene network analysis was undertaken with WGCNA (1.71) (52, 126). DeESeq2 was used to

670 normalize reads and data was formatted with tidyverse. WGCNA identifies modules of co-expressed

671 genes. We chose modules that showed clear day-night differences and mapped the GO terms with

672 Revigo and ShinyGo. We also combined counts from both species and tested for clusters. Here we

673 chose modules that showed species specific clusters.

Gene annotation and filtering:

DEGs, enriched genes and or co-expressed transcripts/ genes were annotated by cross referencing

them with the *B. mori* annotations (see above). Since *B. mori* mappings were not always 'one-to-

one', mappings that were 'one-to-many' were dropped, i.e., transcripts that mapped to multiple

Bombyx genes were dropped. We also improved annotation by mapping the overlapping transcripts

with eggnog mapper. We further filtered these genes into sub functional categories, by identifying

genes with GO annotations or descriptions linked to vision, smell, hearing, circadian, behavior and

brain the list of go terms for each group was obtained by querying amigo

(http://amigo.geneontology.org/) (Table 8).

Gene Mining and in-silico evolution:

We mined genes of interest from ~bombycoidea moths and closely related families that had well

annotated genomes on Ensembl and NCBI (Supp Table 6) and across model insects (Supp. Table 7).

These genomes were chosen as all Bombycoidea from Ensembl with well annotated assemblies and

five species Noctuidae and Geometridae each were included as close relatives. We added *Bombyx*

mori as a reference and *Antheraea pernyi* to represent Saturniidae, and their peptide files were taken

from their respective source papers (40–42). For insects, we chose long-read well annotated genomes

from Ensembl (Ensembl Gene build) and five model systems from insect base (127). We ran

Orthofinder (v2.5.2) (49, 55) with 'orthofinder -f bom-rel/ -S diamond -M msa -A mafft -T fasttree -a

1 -X -z -t 16' to recover orthologs. Each orthogroup often contained multiple sequences per species.

We chose the sequence with the highest identity to the reference *Bombyx mori* sequence using

custom python scripts. In the rare instances where there were multiple references, we took the longest

one. To model the 3D structure of each reference *Bombyx mori* sequence, we ran AlphaFold v2.1.2

(128) using Deepmind's run_alphafold.py

(https://github.com/deepmind/alphafold/blob/main/run_alphafold.py) on University of Florida's

HiperGator. Model database files (i.e. BFD, MAGnify, PDB70, mmCIF PDB, UniRef30, UniRef90)

downloaded from Deepmind in February 2022 were used as parameter value inputs. All other

parameters were used with default settings. The relaxed predicted model with highest ranked pLDDT

(per-residue estimate corresponding to model's predicted score on the lDDT-Cα metric) was chosen

as the final model (129).

We calculated a conservation score for each alignment using Alistat. For alignments less than 1500

amino acids and with high conservation score (Proteins larger than 1300 aa were difficult to model

without high memory GPU's), we modeled conservation using Consurf (130–134). The results were

displayed using Jmol first glance viewer (http://firstglance.jmol.org/) and as screenshots from the

default viewer. The output for alpha fold runs and consurf is available in the (Supplementary Data).

PyMol (135) was used to align the 3D structures and Aliview was used to compare alignments. We

used InterProscan (https://www.ebi.ac.uk/interpro/about/interproscan/) and NetPhos3.1 (136, 137)

for predicting the domains and phosphorylated sites.

References

1.   Y. Xiao, Y. Yuan, M. Jimenez, N. Soni, S. Yadlapalli, Clock proteins regulate spatiotemporal
organization of clock genes to control circadian rhythms. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

2.   K. Beer, C. Helfrich-Förster, Model and Non-model Insects in Chronobiology. *Front. Behav.*
Neurosci. **14**, 601676 (2020).

3.   D. Székely, D. Cogălniceanu, P. Székely, M. Denoël, Adult-Juvenile interactions and temporal niche
partitioning between life-stages in a tropical amphibian. *PLoS One* **15**, e0238949 (2020).

4.   D. Wang, G. Yang, W. Chen, Diel and Circadian Patterns of Locomotor Activity in the Adults of
Diamondback Moth (Plutella xylostella). *Insects* **12** (2021).

5.   J. H. Fullard, N. Napoleone, Diel flight periodicity and the evolution of auditory defences in the
Macrolepidoptera. *Anim. Behav.* **62**, 349–368 (2001).

6.   R. Refinetti, The diversity of temporal niches in mammals. *Biol. Rhythm Res.* **39**, 173–192 (2008).

7.   A. Y. Kawahara, *et al.*, Diel behavior in moths and butterflies: a synthesis of data illuminates the
evolution of temporal activity. *Organisms Diversity and Evolution* **18**, 13–27 (2018).

8.   D. T. C. Cox, A. S. Gardner, K. J. Gaston, Diel niche variation in mammals associated with
729   expanded trait space. *Nat. Commun.* **12**, 1753 (2021).

730   9.   Y. Basset, N. D. Springate, Diel activity of arboreal arthropods associated with a rainforest tree. *J.*
731   *Nat. Hist.* **26**, 947–952 (1992).

732   10.   P. J. Devries, G. T. Austin, N. H. Martin, Diel activity and reproductive isolation in a diverse
733   assemblage of Neotropical skippers (Lepidoptera: Hesperiidae). *Biol. J. Linn. Soc. Lond.* **94**, 723–
734   736 (2008).

735   11.   S. R. Anderson, J. J. Wiens, Out of the dark: 350 million years of conservatism and evolution in diel
736   activity patterns in vertebrates. *Evolution* **71**, 1944–1959 (2017).

737   12.   S. Frey, J. T. Fisher, A. C. Burton, J. P. Volpe, Investigating animal activity patterns and temporal
738   niche partitioning using camera-trap data: challenges and opportunities. *Remote Sens. Ecol. Conserv.*
739   **3**, 123–132 (2017).

740   13.   R. Steen, Diel activity, frequency and visit duration of pollinators in focal plants: in situ automatic
741   camera monitoring and data processing. *Methods Ecol. Evol.* **8**, 203–213 (2017).

742   14.   H. K. Gill, G. Goyal, R. McSorley, Diel Activity of Fauna in Different Habitats Sampled at the
743   Autumnal Equinox. *Fla. Entomol.* **95**, 319–325 (2012).

744   15.   B. Murugavel, A. Kelber, H. Somanathan, Light, flight and the night: effect of ambient light and
745   moon phase on flight activity of pteropodid bats. *J. Comp. Physiol. A Neuroethol. Sens. Neural*
746   *Behav. Physiol.* **207**, 59–68 (2021).

747   16.   H. Somanathan, *et al.*, Nocturnal bees feed on diurnal leftovers and pay the price of day – night
748   lifestyle transition. *Front. Ecol. Evol.* **8** (2020).

749   17.   R. M. Borges, H. Somanathan, A. Kelber, Patterns and Processes in Nocturnal and Crepuscular
750   Pollination Services. *Q. Rev. Biol.* **91**, 389–418 (2016).

751   18.   S. M. Tierney, *et al.*, Consequences of evolutionary transitions in changing photic environments.
752   *Austral Entomology* **56**, 23–46 (2017).

753   19.   W. T. Wcislo, *et al.*, The evolution of nocturnal behaviour in sweat bees, Megalopta genalis and M.
754   ecuadoria (Hymenoptera: Halictidae): an escape from competitors and enemies? *Biol. J. Linn. Soc.*
755   *Lond.* **83**, 377–387 (2004).

756   20.   A. L. Stockl, W. A. Ribi, E. J. Warrant, Adaptations for Nocturnal and Diurnal Vision in the
757   Hawkmoth Lamina. *J. Comp. Neurol.* **524**, 160–175 (2016).

758   21.   Y. Wu, *et al.*, Retinal transcriptome sequencing sheds light on the adaptation to nocturnal and
759   diurnal lifestyles in raptors. *Sci. Rep.* **6**, 1–12 (2016).

760   22.   T. Akiyama, H. Uchiyama, S. Yajima, K. Arikawa, Y. Terai, Parallel evolution of opsin visual
761   pigments in hawkmoths by tuning of spectral sensitivities during transition from a nocturnal to a
762   diurnal ecology. *J. Exp. Biol.* **225**, jeb244541 (2022).

763 23. Y. Sondhi, E. Elis, S. Bybee, J. Theobald, A. Kawahara, Light environment drives evolution of color
764   vision genes in butterflies and moths (2021) https:/doi.org/10.5061/dryad.gmsbcc2kr (May 13,
765   2022).

766 24. M. Izutsu, A. Toyoda, A. Fujiyama, K. Agata, N. Fuse, Dynamics of Dark-Fly Genome Under
767   Environmental Selections. *G3* **6**, 365–376 (2015).

768 25. J. E. Yack, J. H. Fullard, Ultrasonic hearing in nocturnal butterflies. *Nature* **403**, 265–266 (2000).

769 26. F. Sandrelli, R. Costa, C. P. Kyriacou, E. Rosato, Comparative analysis of circadian clock genes in
770   insects. *Insect Mol. Biol.* **17**, 447–463 (2008).

771 27. R. Závodská, I. Sauman, F. Sehnal, Distribution of PER protein, pigment-dispersing hormone,
772   prothoracicotropic hormone, and eclosion hormone in the cephalic nervous system of insects. *J. Biol.*
773   *Rhythms* **18**, 106–122 (2003).

774 28. C. Hermann, *et al.*, The circadian clock network in the brain of different Drosophila species. *J.*
775   *Comp. Neurol.* **521**, 367–388 (2013).

776 29. C. A. Hamilton, *et al.*, Phylogenomics resolves major relationships and reveals significant
777   diversification rate shifts in the evolution of silk moths and relatives. *BMC Evol. Biol.* (2019)
778   https:/doi.org/10.1186/s12862-019-1505-1.

779 30. A. Kobelková, R. Závodská, I. Sauman, O. Bazalová, D. Dolezel, Expression of clock genes period
780   and timeless in the central nervous system of the Mediterranean flour moth, Ephestia kuehniella. *J.*
781   *Biol. Rhythms* **30**, 104–116 (2015).

782 31. D. Brady, A. Saviane, S. Cappellozza, F. Sandrelli, The Circadian Clock in Lepidoptera. *Front.*
783   *Physiol.* **12**, 776826 (2021).

784 32. S. Iwai, Y. Fukui, Y. Fujiwara, M. Takeda, Structure and expressions of two circadian clock genes,
785   period and timeless in the commercial silkmoth, Bombyx mori. *J. Insect Physiol.* **52**, 625–637
786   (2006).

787 33. A. Macias-Muñoz, A. G. Rangel Olguin, A. D. Briscoe, Evolution of Phototransduction Genes in
788   Lepidoptera. *Genome Biol. Evol.* **11**, 2107–2124 (2019).

789 34. X.-C. Jiang, *et al.*, Identification of Olfactory Genes From the Greater Wax Moth by Antennal
790   Transcriptome Analysis. *Front. Physiol.* **12**, 663040 (2021).

791 35. A. de Fouchier, *et al.*, Functional evolution of Lepidoptera olfactory receptors revealed by
792   deorphanization of a moth repertoire. *Nat. Commun.* **8**, 15709 (2017).

793 36. J. W. Truman, L. M. Riddiford, Neuroendocrine control of ecdysis in silkmoths. *Science* **167**, 1624–
794   1626 (1970).

795 37. I. Sauman, S. M. Reppert, Circadian clock neurons in the silkmoth Antheraea pernyi: novel
796   mechanisms of Period protein regulation. *Neuron* **17**, 889–900 (1996).

38. P. M. Tuskes, J. P. Tuttle, M. M. Collins, *The Wild Silk Moths of North America: A Natural History*
of the Saturniidae of the United States and Canada (Cornell University Press, 1996).

39. M. M. Collins, P. M. Tuskes, REPRODUCTIVE ISOLATION IN SYMPATRIC SPECIES OF
DAYFLYING MOTHS (HEMILEUCA: SATURNIIDAE). *Evolution* **33**, 728–733 (1979).

40. F. Lu, *et al.*, SilkDB 3.0: visualizing and exploring multiple levels of data for silkworm. *Nucleic*
Acids Res. **48**, D749–D755 (2020).

41. S.-R. Kim, *et al.*, Genome sequence of the Japanese oak silk moth, Antheraea yamamai: the first
draft genome in the family Saturniidae. *Gigascience* **7**, 1–11 (2018).

42. J. Duan, *et al.*, A chromosome-scale genome assembly of Antheraea pernyi (Saturniidae,
Lepidoptera). *Mol. Ecol. Resour.* **20**, 1372–1383 (2020).

43. H. Langer, G. Schmeinck, F. Anton-Erxleben, Identification and localization of visual pigments in
the retina of the moth, Antheraea polyphemus (Insecta, Saturniidae). *Cell Tissue Res.* **245**, 81–89
(1986).

44. I. Kitching, *et al.*, A global checklist of the Bombycoidea (Insecta: Lepidoptera). *Biodiversity Data*
Journal **6**, e22236 (2018).

45. M. J. Scoble, *The Lepidoptera. Form, function and diversity* (Oxford University Press, 1992).

46. Lemaire, Minet, 18. The Bombycoidea and their Relatives. *Volume 1: Evolution, Systematics, and*
(2013).

47. R. Rougerie, *et al.*, Phylogenomics Illuminates the Evolutionary History of Wild Silkmoths in Space
and Time (Lepidoptera: Saturniidae). *bioRxiv*, 2022.03.29.486224 (2022).

48. Z. Wang, M. Gerstein, M. Snyder, RNA-Seq: a revolutionary tool for transcriptomics. *Nat. Rev.*
Genet. **10**, 57–63 (2009).

49. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157 (2015).

50. D. Li, *et al.*, An evaluation of RNA-seq differential analysis methods. *PLoS One* **17**, e0264246
(2022).

51. N. Sánchez-Baizán, L. Ribas, F. Piferrer, Improved biomarker discovery through a plot twist in
transcriptomic data analysis. *BMC Biol.* **20**, 208 (2022).

52. P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network analysis. *BMC*
Bioinformatics **9**, 559 (2008).

53. Lafayette Lafayette Press, *A Survey of Best Practices for RNA-Seq Data Analysis* (CreateSpace
Independent Publishing Platform, 2016).

54. C. P. Cantalapiedra, A. Hernández-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas, eggNOG-mapper v2:

Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale.
Mol. Biol. Evol. **38**, 5825–5829 (2021).

55. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative genomics.
Genome Biol. **20**, 238 (2019).

56. P. E. Hardin, J. C. Hall, M. Rosbash, Behavioral and molecular analyses suggest that circadian
output is disrupted by disconnected mutants in D. melanogaster. *EMBO J.* **11**, 1–6 (1992).

57. M. S. Dushay, M. Rosbash, J. C. Hall, The disconnected visual system mutations in Drosophila
melanogaster drastically disrupt circadian rhythms. *J. Biol. Rhythms* **4**, 1–27 (1989).

58. K. J. Lee, M. Mukhopadhyay, P. Pelka, A. R. Campos, H. Steller, Autoregulation of the Drosophila
disconnected gene in the developing visual system. *Dev. Biol.* **214**, 385–398 (1999).

59. V. Suri, Z. Qian, J. C. Hall, M. Rosbash, Evidence that the TIM light response is relevant to light-
induced phase shifts in Drosophila melanogaster. *Neuron* **21**, 225–234 (1998).

60. B. K. Dey, X.-L. Zhao, E. Popo-Ola, A. R. Campos, Mutual regulation of the Drosophila
disconnected (disco) and Distal-less (Dll) genes contributes to proximal-distal patterning of antenna
and leg. *Cell Tissue Res.* **338**, 227–240 (2009).

61. H. Somanathan, A. Kelber, R. M. Borges, R. Wallén, E. J. Warrant, Visual ecology of Indian
carpenter bees II: adaptations of eyes and ocelli to nocturnal and diurnal lifestyles. *J. Comp. Physiol.
A Neuroethol. Sens. Neural Behav. Physiol.* **195**, 571–583 (2009).

62. A. L. Stöckl, D. O'Carroll, E. J. Warrant, Higher-order neural processing tunes motion neurons to
visual ecology in three species of hawkmoths. *Proceedings of the Royal Society B: Biological
Sciences* **284**, 20170880 (2017).

63. I. Shimizu, Y. Yamakawa, Y. Shimazaki, T. Iwasa, Molecular cloning of Bombyx cerebral opsin
(Boceropsin) and cellular localization of its expression in the silkworm brain. *Biochem. Biophys.
Res. Commun.* **287**, 27–34 (2001).

64. K. Moses, M. C. Ellis, G. M. Rubin, The glass gene encodes a zinc-finger protein required by
Drosophila photoreceptor cells. *Nature* **340**, 531–536 (1989).

65. R. Tearle, Tissue specific effects of ommochrome pathway mutations in Drosophila melanogaster.
Genet. Res. **57**, 257–266 (1991).

66. H. K. Shamloula, *et al.*, rugose (rg), a Drosophila A kinase anchor protein, is required for retinal
pattern formation and interacts genetically with multiple signaling pathways. *Genetics* **161**, 693–710
(2002).

67. A. Stöckl, *et al.*, Differential investment in visual and olfactory brain areas reflects behavioural
choices in hawk moths. *Sci. Rep.* **6** (2016).

68. A. Sourakov, R. W. Chadd, *The Lives of Moths: A Natural History of Our Planet's Moth Life*
(Princeton University Press, 2022).

69.   A. Y. Kawahara, *et al.*, Phylogenomics reveals the evolutionary timing and pattern of butterflies and
866   moths. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 22657–22663 (2019).

867   70.   J. J. Rubin, *et al.*, The evolution of anti-bat sensory illusions in moths. *Science Advances* (2018)
868   https:/doi.org/10.1126/sciadv.aar7428.

869   71.   J. E. Yack, L. D. Otero, J. W. Dawson, A. Surlykke, J. H. Fullard, Sound production and hearing in
870   the blue cracker butterfly Hamadryas feronia (Lepidoptera, nymphalidae) from Venezuela. *J. Exp.*
871   *Biol.* **203**, 3689–3702 (2000).

872   72.   J. R. Barber, *et al.*, Anti-bat ultrasound production in moths is globally and phylogenetically
873   widespread. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2117485119 (2022).

874   73.   G. A. Jacobs, J. P. Miller, Z. Aldworth, Computational mechanisms of mechanosensory processing
875   in the cricket. *J. Exp. Biol.* **211**, 1819–1828 (2008).

876   74.   J. M. Camhi, Locust wind receptors: I. Transducer mechanics and sensory response. *J. Exp. Biol.* **50**,
877   335–348 (1969).

878   75.   J. M. Camhi, Locust wind receptors. 3. Contribution to flight initiation and lift control. *J. Exp. Biol.*
879   **50**, 363–373 (1969).

880   76.   J. C. Tuthill, R. I. Wilson, Mechanosensation and Adaptive Motor Control in Insects. *Curr. Biol.* **26**,
881   R1022–R1038 (2016).

882   77.   J. D. Baker, S. Adhikarakunnathu, M. J. Kernan, Mechanosensory-defective, male-sterile unc
883   mutants identify a novel basal body protein required for ciliogenesis in Drosophila. *Development*
884   **131**, 3411–3422 (2004).

885   78.   D. D. Dell'Aglio, W. O. McMillan, S. H. Montgomery, Shifting balances in the weighting of sensory
886   modalities are predicted by divergence in brain morphology in incipient species of Heliconius
887   butterflies. *Anim. Behav.* **185**, 83–90 (2022).

888   79.   S. H. Montgomery, R. M. Merrill, S. R. Ott, Brain composition in Heliconius butterflies,
889   posteclosion growth and experience-dependent neuropil plasticity. *J. Comp. Neurol.* **524**, 1747–1769
890   (2016).

891   80.   S. Anton, W. Rössler, Plasticity and modulation of olfactory circuits in insects. *Cell Tissue Res.* **383**,
892   149–164 (2021).

893   81.   S. Halali, *et al.*, Predictability of temporal variation in climate and the evolution of seasonal
894   polyphenism in tropical butterfly communities. *J. Evol. Biol.* **34**, 1362–1375 (2021).

895   82.   W. Kuenzinger, *et al.*, Innate colour preferences of a hawkmoth depend on visual context. *Biol. Lett.*
896   **15**, 20180886 (2019).

897   83.   A. Kelber, Pattern discrimination in a hawkmoth: innate preferences, learning performance and
898   ecology. *Proc. Biol. Sci.* **269**, 2573–2577 (2002).

84. G. T. Broadhead, T. Basu, M. von Arx, R. A. Raguso, Diel rhythms and sex differences in the
locomotor activity of hawkmoths. *J. Exp. Biol.* **220**, 1472–1480 (2017).

85. M. Bischoff, R. A. Raguso, A. Jürgens, D. R. Campbell, Context-dependent reproductive isolation
mediated by floral scent and color. *Evolution* **69**, 1–13 (2015).

86. S. Jaeger, C. Girvin, N. Demarest, E. LoPresti, Secondary pollinators contribute to reproductive
success of a pink-flowered sand verbena population. *Ecology* **104**, e3977 (2023).

87. W. V. So, *et al.*, takeout, a novel Drosophila gene under circadian clock transcriptional regulation.
Mol. Cell. Biol. **20**, 6935–6944 (2000).

88. T. Kikawada, *et al.*, Trehalose transporter 1, a facilitated and high-capacity trehalose transporter,
allows exogenous trehalose uptake into cells. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11585–11590
(2007).

89. H. Steller, K. F. Fischbach, G. M. Rubin, Disconnected: a locus required for neuronal pathway
formation in the visual system of Drosophila. *Cell* **50**, 1139–1153 (1987).

90. C. Helfrich-Förster, Robust circadian rhythmicity of Drosophila melanogaster requires the presence
of lateral neurons: a brain-behavioral study of disconnected mutants. *J. Comp. Physiol. A* **182**, 435–
453 (1998).

91. L. R. Sanders, M. Patel, J. W. Mahaffey, The Drosophila gap gene giant has an anterior segment
identity function mediated through disconnected and teashirt. *Genetics* **179**, 441–453 (2008).

92. E. Blanchardon, *et al.*, Defining the role of Drosophila lateral neurons in the control of circadian
rhythms in motor activity and eclosion by targeted genetic ablation and PERIOD protein
overexpression. *Eur. J. Neurosci.* **13**, 871–888 (2001).

93. A. D. Keller, T. Maniatis, Only two of the five zinc fingers of the eukaryotic transcriptional
repressor PRDI-BF1 are required for sequence-specific DNA binding. *Mol. Cell. Biol.* **12**, 1940–
1949 (1992).

94. S. Johnson, N. Boob, A strange pair. *News of the Lepidopterists' Society* **55**, 116 (2013).

95. C. J. van der Kooi, D. G. Stavenga, K. Arikawa, G. Belušič, A. Kelber, Evolution of Insect Color
Vision: From Spectral Sensitivity to Visual Ecology. *Annu. Rev. Entomol.* **66**, 435–461 (2021).

96. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**, 10–12 (2011).

97. M. G. Grabherr, *et al.*, Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

98. D. R. Zerbino, E. Birney, Velvet: algorithms for de novo short read assembly using de Bruijn graphs.
Genome Res. **18**, 821–829 (2008).

99. G. Robertson, *et al.*, De novo assembly and analysis of RNA-seq data. *Nat. Methods* **7**, 909–912

(2010).

100.   R. Luo, *et al.*, SOAPdenovo2: an empirically improved memory-efficient short-read de novo
        assembler. *Gigascience* **1**, 18 (2012).

101.   D. G. Gilbert, Longest protein, longest transcript or most expression, for accurate gene
        reconstruction of transcriptomes? *bioRxiv*, 829184 (2019).

102.   D. Gilbert, Gene-omes built from mRNA seq not genome DNA (2016)
        https:/doi.org/10.7490/f1000research.1112594.1 (June 6, 2022).

103.   , *https://github.com/NCGAS/de-novo-transcriptome-assembly-pipeline* (Github) (June 6, 2022).

104.   M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO Update: Novel and
        Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
        Eukaryotic, Prokaryotic, and Viral Genomes. *Mol. Biol. Evol.* **38**, 4647–4654 (2021).

105.   R. Smith-Unna, C. Boursnell, R. Patro, J. M. Hibberd, S. Kelly, TransRate: reference-free quality
        assessment of de novo transcriptome assemblies. *Genome Res.* **26**, 1134–1144 (2016).

106.   A. Mikheenko, A. Prjibelski, V. Saveliev, D. Antipov, A. Gurevich, Versatile genome assembly
        evaluation with QUAST-LG. *Bioinformatics* **34**, i142–i150 (2018).

107.   W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein or
        nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

108.   L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-generation
        sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

109.   M. Steinegger, J. Söding, MMseqs2 enables sensitive protein sequence searching for the analysis
        of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

110.   Gene Ontology Consortium, The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids
        Res.* **49**, D325–D334 (2021).

111.   D. M. Bryant, *et al.*, A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of
        Limb Regeneration Factors. *Cell Rep.* **18**, 762–776 (2017).

112.   R. D. Finn, J. Clements, S. R. Eddy, HMMER web server: interactive sequence similarity
        searching. *Nucleic Acids Research* **39**, W29–W37 (2011).

113.   M. Punta, *et al.*, The Pfam protein families database. *Nucleic Acids Res.* **40**, D290–301 (2012).

114.   R. D. Finn, *et al.*, The Pfam protein families database: towards a more sustainable future. *Nucleic
        Acids Res.* **44**, D279–85 (2016).

115.   T. N. Petersen, S. Brunak, G. von Heijne, H. Nielsen, SignalP 4.0: discriminating signal peptides
        from transmembrane regions. *Nat. Methods* **8**, 785–786 (2011).

965   116.   A. Krogh, B. Larsson, G. von Heijne, E. L. Sonnhammer, Predicting transmembrane protein
966   topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580
967   (2001).

968   117.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool.
969   *J. Mol. Biol.* **215**, 403–410 (1990).

970   118.   M. Kanehisa, S. Goto, Y. Sato, M. Furumichi, M. Tanabe, KEGG for integration and
971   interpretation of large-scale molecular data sets. *Nucleic Acids Res.* **40**, D109–14 (2012).

972   119.   M. Ashburner, *et al.*, Gene ontology: tool for the unification of biology. The Gene Ontology
973   Consortium. *Nat. Genet.* **25**, 25–29 (2000).

974   120.   S. Powell, *et al.*, eggNOG v3.0: orthologous groups covering 1133 organisms at 41 different
975   taxonomic ranges. *Nucleic Acids Res.* **40**, D284–9 (2012).

976   121.   R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and bias-
977   aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

978   122.   M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential
979   expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

980   123.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
981   seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

982   124.   Alexa, Rahnenfuhrer, topGO: Enrichment analysis for Gene Ontology. R package version 2.28. 0.
983   *Cranio* (2016).

984   125.   S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants.
985   *Bioinformatics* **36**, 2628–2629 (2020).

986   126.   P. Langfelder, S. Horvath, Fast R Functions for Robust Correlations and Hierarchical Clustering.
987   *J. Stat. Softw.* **46** (2012).

988   127.   Y. Mei, *et al.*, InsectBase 2.0: a comprehensive gene resource for insects. *Nucleic Acids Res.* **50**,
989   D1040–D1045 (2022).

990   128.   J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
991   589 (2021).

992   129.   V. Mariani, M. Biasini, A. Barbato, T. Schwede, lDDT: a local superposition-free score for
993   comparing protein structures and models using distance difference tests. *Bioinformatics* **29**, 2722–
994   2728 (2013).

995   130.   B. Yariv, *et al.*, Using evolutionary data to make sense of macromolecules with a "face-lifted"
996   ConSurf. *Protein Sci.* **32**, e4582 (2023).

997   131.   H. Ashkenazy, *et al.*, ConSurf 2016: an improved methodology to estimate and visualize
998   evolutionary conservation in macromolecules. *Nucleic Acids Res.* **44**, W344–50 (2016).

999   132.   G. Celniker, *et al.*, ConSurf: Using evolutionary data to raise testable hypotheses about protein
1000   function. *Isr. J. Chem.* **53**, 199–206 (2013).

1001   133.   H. Ashkenazy, E. Erez, E. Martz, T. Pupko, N. Ben-Tal, ConSurf 2010: calculating evolutionary
1002   conservation in sequence and structure of proteins and nucleic acids. *Nucleic Acids Res.* **38**, W529–
1003   33 (2010).

1004   134.   M. Landau, *et al.*, ConSurf 2005: the projection of evolutionary conservation scores of residues
1005   on protein structures. *Nucleic Acids Res.* **33**, W299–302 (2005).

1006   135.   Schrödinger, LLC, The PyMOL Molecular Graphics System, Version 1.8 (2015).

1007   136.   N. Blom, S. Gammeltoft, S. Brunak, Sequence and structure-based prediction of eukaryotic
1008   protein phosphorylation sites. *J. Mol. Biol.* **294**, 1351–1362 (1999).

1009   137.   N. Blom, T. Sicheritz-Pontén, R. Gupta, S. Gammeltoft, S. Brunak, Prediction of post-
1010   translational glycosylation and phosphorylation of proteins from the amino acid sequence.
1011   *Proteomics* **4**, 1633–1649 (2004).

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031   **Supplemental Figures**

1032
1033   **Figure S1**: Variation in salmon mapped libraries for *Dryocampa rubicunda* and *Anisota pellucida*
1034
1035

1036
1037   **Figure S2**: Variation in normalised libraries for *Dryocampa rubicunda* and *Anisota pellucida*
1038

1039

Dryocampa rubicunda   *Anisota pellucida*

1040
1041   **Figure S3**: PC1 and PC2 of libraries labelled for *Dryocampa rubicunda* and *Anisota pellucida*.
1042   Orange and blue labels refer to midday 'Day' and midnight 'Night' collection time points,
1043   respectively. This PCA was generated from mapped salmon reads after normalising with DESeq2.
1044

FC > 2 , padj <0.05, *Anisota pellucida* (Diurnal) : **Night Upregulation***
ShinyGO (FDR <0.1, Pathway (1-2000), top 100 pathways)

FC < -2 , padj <0.05, *Anisota pellucida* (Diurnal) : **Day Upregulation***
ShinyGO (FDR <0.1, Pathway (1-2000), top 20 pathways)

1045

1046
Figure S4: Enriched lists of GO terms associated with significant positively and negatively
expressed genes recovered from 2 or more analyses (DESeq2, WGCNA, EdgeR) for *Anisota*
pellucida. ShinyGo was used to highlight terms with genes showing representation compared to the
background of detectable genes values representing ShinyGo overrepresentation. Top: Enriched Day
Upregulated genes (FDR>0.1) with gene expression FC>=2, adjusted p-value <0.05. Bottom:
Enriched Night Upregulated genes (FDR>0.1) with gene expression FC<=2, adjusted p-value <0.05
and that showed at least 2 occurrences in the combined dataset. Values are sorted by -log FDR
*Genes that showed up in at least 2 separate analyses are included, (>=2 occurrences in the combined
dataset). The background used was all transcripts that were recovered in the DEG analysis and had
Bombyx orthologs

FC > 2 , padj <0.05, *Dryocampa* (Nocturnal): **Night Upregulation***
ShinyGO (FDR <0.32, Pathway (1-2000), top 40 pathways)

FC < 2 , padj <0.05, *Dryocampa* (Nocturnal) : **Day Upregulation***
ShinyGO (FDR <0.1, Pathway (1-2000), top 100 pathways)

Figure S5 : Enriched lists of GO terms associated with significant positively and negatively
expressed genes recovered from 2 or more analyses (DESeq2,WGCNA, EdgeR) for *Dryocampa*
rubicunda . ShinyGo was used to highlight terms with genes showing representation compared to the
background of detectable genes values representing ShinyGo overrepresentation. Top: Enriched Day
Top 40 upregulated genes (FDR>0.32) with gene expression FC>=2, adjusted p-value <0.05, 77
terms enriched , but not all are displayed, see supplementary material for entire list. Bottom:
Enriched Night Upregulated genes (FDR>0.3) with gene expression FC<=2, adjusted p-value <0.05
and that showed at least 2 occurrences in the combined dataset ***Genes** that showed up in at least 2
separate analyses are included, (>=2 occurrences in the combined dataset). The background used was
all transcripts that were recovered in the DEG analysis and had *Bombyx* orthologs

1074
Figure S6: Modules of clustered genes grouped using normalized expression for *Anisota pellucida*
(top) and *Dryocampa rubicunda* (bottom) Grey60 (38 genes) and Tan (32 genes) modules show
species- time of day linked patterns. Left: Shows how patterns of gene expression correlate across
samples for modules Right: Shows the normalized expression for all. Normalization was done with
DESeq2 and reads were mapped to the more stringently filtered transcriptome. A soft power analysis
was done and the picked power=9 for the WGCNA analysis.

1081

1082

1083

1084 **Figure S7:** Shiny Go enrichment for Modules of clustered genes grouped using normalized

1085 expression for *Anisota pellucida* (top) and *Dryocampa rubicunda* (bottom) Grey60 and Tan.

1086 Background used was all *Bombyx mori* genes and FDR cut-off =0.1 min pathway =1 and GO

1087 Biological processes are displayed.

Figure S8: Shiny Go enrichment for Modules of clustered genes grouped using normalized expression for combined overlapping reads for the two species (Top) Blue and (Bottom) Turquoise. Background used was all *Bombyx mori* genes and FDR cut-off =1 min pathway =1 and GO Biological processes are displayed.

Figure S9: Variation in structural conservation across moths for predicted proteins for genes interest.
Structures were obtained using AlphaFold predictions of *Bombyx mori* homologs. Left: Space filling
models, Right cartoon models, colors represent conservation across the alignment from or
Bombycoidea and related moth species. Conservation scores were calculated and mapped onto the

 alpha fold structures using the Consurf (131) web server (For other proteins, see Supplementary data)

Figure S10: Illustration of conservation across the alignments and structure for **(A)** *disco* and **(B)** *tk* across Bombycoidea and relatives (Left) and insects (Right). Each inset contains the *Consurf* mapped pdb structures on the left. The *alistat* pairwise sequence conservation in the alignment scores on the right and the consensus of the protein alignments were visualized using *Geneious*. For conservation scores for other genes and lists of the species used for this modelling, see Supplementary Tales (Supp. Table: 5-7). Supplementary Dataset 10 has the associated models and output from the analyses.

Figure S11: InterPro and NetPhos predictions for *B. mori disco* protein. **(A)**InterPro predicted
several zinc finger domains, disordered regions and disco and disco-related families. The numbers in
black highlight the regions for the predicted domain and the numbers in red list the NetPhos
predicted phosphorylated sites (threshold >0.9). **(B)** An overall distribution of the predicted
phosphorylated sites with different thresholds (Left, Threshold=0.9, Right Threshold= 0.5)

Figure S12: Lack of sequence level conservation in takeout, but high structural overlap

(A) Right: Takeout UniProt blastp hits. (BLOSUM62, filtered for *D. melanogaster*), alignments =250, e-threshold= 10, database = uniprotkb_refprot+swissprot, list of illustrative hits is shown. Left: Similar blastp results from flybase limiting the species to *D. melanogaster*. **(B)** Consurf predicted scores for closely related moths mapped on to the *B, mori* takeout predicted protein structure. **(C)** 3D structural alignments for fly (*D. melanogaster)* and moth (*B. mori)* takeout 3D models. **(D)** Uniprot protein sequence alignment for fly and moth takeout proteins.

Tables

Species	Diel-niche	Collection Time	Replicates (n)
Anisota pellucida	Diurnal	Day	4
		Night	2
Dryocampa rubicunda	Nocturnal	Day	3
		Night	3

Table 1: Sampling design showing species, collection time and number of replicates in each treatment. All specimens were male.

Species	Assembler	kmer-size	Max- BUSCO groups	Lep BUSCO Score(%)	max-N50
Ani_pel	Velvet	35-85	4703 (k55)	88	3918 (k55)
Ani_pel	Soap	35-85	4682 (k35)	88.5	1661 (k35)
Ani_pel	Transabyss	35-85	4216 (k35)	79	1332 (k35)
Ani_pel	Trinity	25	4948	93	1608
Dry_rub	Velvet	45-85	4768 (k55)	90	3957 (k55)
Dry_rub	Soap	35-85	4753 (k35)	89	1915 (k75)
Dry_rub	Transabyss	55-85	3159 (k55)	59	1058 (k55)
Dry_rub	Trinity	25	4939	93	1595

Table 2: Assembly variation using different assemblers and kmer settings.

Key: Dry_rub: *Dryocampa rubicunda*, Ani_pel: *Anisota pellucida.* All models included day and
night samples.

Species	Method	Total DEGs	DEGs (Up)	DEGs (Down)	Annotated DEGs	Unannotated %
A. pellucida D	EdgeR	350	154	196	211	39.71
D, rubicunda N	EdgeR	393	211	182	236	39.95
A. pellucida D	DESeq2	697	141	556	420	39.7
D. rubicunda N	DESeq2	498	253	245	300	39.7

Table 3: DEG recovery and gene annotation for EdgeR and DESeq2 analyses along with percent
annotation using *Bombyx.* Up=FC>2, Implies night expression is greater than day expression.
Down=FC<2, implies day expression is greater (control=Day, Night = Treatment)

Species	Assembly annotation	BUSCO score	BUSCO duplication	No. of contigs*	Redundancy (/15k)
Dry_rub	Combined	94.3	90.8	281790	18.8x
Dry_rub	Combined_cds_cdhit_95	90.1	80.2	184808	12.3x
Dry_rub	Combined_cds_filtered (v5)	85.2	59.6	41712	2.8x
Dry_rub	Combined_cds_stringent_filter (v6)	75.4	4.9	19877	1.3x
Ani_pel	Combined	94.3	90.8	281790	18.78
Ani_pel	Combined_cds	93.9	90.3	233097	13.3x
Ani_pel	Combined_cds_filtered_cds (v5)	85.2	59.2	41712	2.8x
Ani_pel	Combined_cds_stringent_filter (v6)	79.3	4.7	19087	1.3x

1152

Table 4: BUSCO and contig distribution statistics for different species' *de-novo* assemblies with

and without filtering. Increased filtering reduces redundancy and duplication but comes at the

cost of reducing BUSCO scores. *All statistics are based on contigs of size >= 500 bp.

1156

nGenes	Fold Enrichment	Pathway	GO_ID	Gene Names
2	85.53	**Regulation of excretion**	GO:0044062	NPHS1,SLC9A3R1
1	42.76	Response to herbicide	GO:0009635	Alad
1	42.76	Carbohydrate catabolic process	GO:0016052	BMSK0015853
1	42.76	Blastocyst formation	GO:0001825	Actl6a
1	42.76	Tryptophan catabolic process to kynurenine	GO:0019441	KFase
1	42.76	Microvillus assembly	GO:0030033	SLC9A3R1
1	42.76	Parallel actin filament bundle assembly	GO:0030046	jv
1	42.76	Heparan sulfate proteoglycan metabolic process	GO:0030201	botv
1	42.76	Inactivation of MAPK activity	GO:0000188	Spred1
1	42.76	B cell activation	GO:0042113	AKAP17A
1	42.76	Negative regulation of glycogen biosynthetic process	GO:0045719	Gfpt1
1	42.76	Behavioral response to pain	GO:0048266	pain
1	42.76	Interphase microtubule nucleation by interphase microtubule organizing center	GO:0051415	Grip91
1	28.51	Cellular defense response	GO:0006968	PNLIPRP2
1	28.51	Cartilage condensation	GO:0001502	COL11A1
1	28.51	Short-chain fatty acid import	GO:0015913	SLC5A8
1	28.51	Male genitalia development	GO:0030539	ken
1	28.51	Microtubule organizing center organization	GO:0031023	Gcc2
1	28.51	Telomere maintenance via semi-conservative replication	GO:0032201	RFC3
1	28.51	Glomerular basement membrane development	GO:0032836	NPHS1
1	28.51	Endosomal vesicle fusion	GO:0034058	Ankfy1
1	28.51	High-density lipoprotein particle assembly	GO:0034380	ABCA7

1	28.51	Positive regulation of Rho protein signal transduction	GO:0035025	MCF2L
1	28.51	Xenobiotic transport	GO:0042908	Mdr49
1	28.51	DNA replication-removal of RNA primer	GO:0043137	Rnaseh1
2	28.51	Negative regulation of endocytosis	GO:0045806	ABCA7,Mctp1
2	28.51	**Negative gravitaxis**	GO:0048060	pain,pyx
1	28.51	Imaginal disc-derived wing expansion	GO:0048526	Spn88Ea
1	28.51	**Positive regulation of synapse structural plasticity**	GO:0051835	FRMPD4
1	28.51	Chitin biosynthetic process	GO:0006031	chs-2
2	19.01	Lysosomal transport	GO:0007041	Vps53,g
2	19.01	RNA interference	GO:0016246	Dcr-1,bel
2	15.55	Negative regulation of endopeptidase activity	GO:0010951	Spn88Ea,Serpinb1a
2	13.16	Formation of cytoplasmic translation initiation complex	GO:0001732	eIF3-S5, BMSK0008517
2	10.69	Protein heterotetramerization	GO:0051290	pyx,Agrn
2	10.06	**Positive regulation of synaptic growth at neuromuscular junction**	GO:0045887	Agrn,Sin1
2	8.55	Endoplasmic reticulum organization	GO:0007029	atl,Lman1
2	7.78	**Regulation of synaptic growth at neuromuscular junction**	GO:0008582	atl,hiw
2	7.13	Negative regulation of cell migration	GO:0030336	SLC9A3R1,Mctp1
5	5.94	**Visual perception**	GO:0007601	RDH11,COL11A1, OP1,WDR36,disco

1157

1158 **Table 5:** Top significantly enriched GO terms for *A. pellucida* day upregulation

1159

No. Genes	Fold Enrichment	Pathway	GO_ID	Gene Names
1	41.61	Female meiosis II	GO:0007147	polo
1	41.61	Female germline ring canal formation-actin assembly	GO:0008302	Src64B
1	41.61	Dosage compensation by hyperactivation of X chromosome	GO:0009047	mle
1	41.61	10-formyltetrahydrofolate catabolic process	GO:0009258	aldh1l1
1	41.61	**Sialic acid transport**	**GO:0015739**	**SLC17A5**

1	41.61	Arachidonic acid metabolic process	GO:0019369	FAAH2
1	41.61	Regulation of cellular metabolic process	GO:0031323	melt
1	41.61	Protein exit from endoplasmic reticulum	GO:0032527	SEC16A
1	41.61	Single strand break repair	GO:0000012	gkt
1	41.61	Ring gland development	GO:0035271	gl
1	41.61	Notch receptor processing-ligand-dependent	GO:0035333	Nct
1	41.61	Positive regulation of RNA polymerase II transcriptional preinitiation complex assembly	GO:0045899	PSMC6
1	41.61	Tetrahydrofolate metabolic process	GO:0046653	MTHFS
1	41.61	Neurotrophin TRK receptor signaling pathway	GO:0048011	Bcar1
1	41.61	Glucose catabolic process	GO:0006007	agp
1	41.61	Glutaminyl-tRNAGln biosynthesis via transamidation	GO:0070681	gatA
1	41.61	Error-free translesion synthesis	GO:0070987	POLH
1	41.61	Protein deubiquitination involved in ubiquitin-dependent protein catabolic process	GO:0071947	trbd
1	41.61	Negative regulation of release of cytochrome c from mitochondria	GO:0090201	Parl
1	41.61	Regulation of cardiac conduction	GO:1903779	nkx-2.5
1	41.61	Methionyl-tRNA aminoacylation	GO:0006431	MetRS-m
2	33.29	Tetrahydrofolate interconversion	GO:0035999	Sardh,MTHFS
1	27.74	Tubulin complex assembly	GO:0007021	Tbcd
1	27.74	Double-strand break repair via break-induced replication	GO:0000727	Cdc45
1	27.74	Aminophospholipid transport	GO:0015917	ATP11B
1	27.74	**Synaptic target attraction**	**GO:0016200**	**Ten-a**
1	27.74	N-terminal peptidyl-methionine acetylation	GO:0017196	Naa60
1	27.74	Peptidyl-lysine methylation	GO:0018022	Eef1akmt2
1	27.74	Negative regulation of ossification	GO:0030279	LRP4
1	27.74	Negative regulation of phosphoprotein phosphatase activity	GO:0032515	Bod1
1	27.74	Store-operated calcium entry	GO:0002115	olf186-F
1	27.74	Cytoplasmic sequestering of transcription factor	GO:0042994	Sufu
1	27.74	Melanin biosynthetic process from tyrosine	GO:0006583	BMSK0009475
1	20.80	Cytoplasmic mRNA processing body assembly	GO:0033962	patl1

1	20.80	Positive regulation of transcription from RNA polymerase I promoter	GO:0045943	HEATR1
1	20.80	Negative regulation of insulin secretion	GO:0046676	HADH
1	16.64	Microtubule nucleation	GO:0007020	TUBG1
1	16.64	**Eclosion rhythm**	**GO:0008062**	**disco**
1	16.64	Late endosome to vacuole transport	GO:0045324	chmp3
1	16.64	**Negative regulation of axon extension involved in axon guidance**	**GO:0048843**	**tap**
1	16.64	Glucose transmembrane transport	GO:1904659	SLC2A3
1	16.64	Nucleocytoplasmic transport	GO:0006913	Anp32a
2	13.87	Chromosome segregation	GO:0007059	PPP2R1A,Naa60
1	13.87	Nucleotide catabolic process	GO:0009166	BMSK0009399
1	13.87	Phosphatidylethanolamine biosynthetic process	GO:0006646	PCYT2
2	12.80	Negative regulation of sequence-specific DNA binding transcription factor activity	GO:0043433	melt,Sufu
2	10.40	Protein heterotetramerization	GO:0051290	pyx,LRP4
2	9.25	Phospholipid biosynthetic process	GO:0008654	Mboat1,PCYT2
2	9.25	**Locomotor rhythm**	**GO:0045475**	**Ork1,disco**
2	8.76	Pupal chitin-based cuticle development	GO:0008364	Gld,Gld
3	8.61	Mitochondrial translation	GO:0032543	BMSK0007759,MRPS35,gatA
2	8.32	Sperm storage	GO:0046693	Gld,Gld
2	7.93	Cytoplasmic translation	GO:0002181	RpL4,
2	7.93	Synaptic growth at neuromuscular junction	GO:0051124	LRP4,Ten-a
2	7.56	Negative regulation of canonical Wnt signaling pathway	GO:0090090	PSMC6,LRP4
2	6.93	**Mushroom body development**	**GO:0016319**	**tap,Src64B**
2	6.93	Post-translational protein modification	GO:0043687	TULP4,PSMC6
2	6.93	**Synapse organization**	**GO:0050808**	**Ten-a,LRP4**
2	5.74	**Trehalose transport**	**GO:0015771**	**Tret1-1,Tret1-1**
2	5.20	Positive regulation of canonical Wnt signaling pathway	GO:0090263	PSMC6,trbd
3	5.09	Cell migration	GO:0016477	trbd,Bcar1,SPEF1
2	5.04	Response to endoplasmic reticulum stress	GO:0034976	SEC16A,Pdi

3	4.62	Mitotic cell cycle			GO:0000278		TUBG1,Tbcd,polo
2	4.62	Microtubule cytoskeleton organization			GO:0000226		TUBG1,Tbcd
3	3.37	G-protein coupled receptor signaling pathway			GO:0007186		Bcar1,Atrnl1,TRHR

Table 6: Top significantly enriched GO terms for *D. rubicunda* night upregulation

Gene Symbol	Gene Name	Function	DESeq2	EdgeR	WGCNA	Flybase ID	SILKD ID
BNC1/disco	disconnected	vision	1*	1*		FBgn0000459	BMSK0012241
Spt5	Spt5	transcription	1*	1*		FBgn0040273	BMSK0011143
1q/trbd	trabid	development		1		FBgn0037734	BMSK0015294
RpL4	Ribosomal protein L4	energy use, Ribosomal production	1*			FBgn0003279	BMSK0006059
Prp2/PPO2	Prophenoloxidase 2	Wound melanization	1*			FBgn0033367	BMSK0009475
mRpS5	mitochondrial ribosomal proteins	energy use, mitochondrial maintenance	1*			FBgn0287187	BMSK0014662
SLC2A6/Tret1-1	Solute Carrier Family 2 Member 6	locomotion, energy metabolism	1*			FBgn0050035	BMSK0003817**, BMSK0003818
TUBG1/γTub23C	tubulin gamma 1	brain development in adult	1*			FBgn0260639	BMSK0002451
PARL/rho-7	presenilin associated rhomboid like	mitochondrial maintenance	1*			FBgn0033672	BMSK0008635
SLC17A5/MFS10	Solute carrier family 17 member 5	transmembrane ion transporter	1*			FBgn0030452	BMSK0001219
titin1/sls	sallimus	locomotion	1			FBgn0086906	BMSK0000202
unc-22/unc	uncoordinated	adult locomotion, hearing	1		1	FBgn0003950	BMSK0015609**, BMSK0000066
ARHGAP17/RhoGAP92B	rho GTPase-activating protein 17	auditory, brain development			1	FBgn0038747	BMSK0009977
tk	tachykinin	locomotion and circadian rhythms			1	FBgn0037976	BMSK0003725
BmorOBP1/	Odorant Binding	odor binding			1	FBgn0034768	BMSK0013390

Obp58a	Protein 1						
PPAP2A/wun	Inorganic Pyrophosphatase 2/ wunnen	vision			1	FBgn0016078	BMSK0011305
BmorOBP2/ Obp84a	Odorant Binding Protein 2	odor binding			1	FBgn0011282	BMSK0009610
takt/JHBP	takeout-like	circadian			1	FBgn0038395	BMSK0013046

1165

Table 7: Gene symbol from EggNog or *B. mori* annotation. Gene names, function and closest

flybase ID orthologs used to collect evidence for the various listed functions.* Indicated that the

gene altered direction or trend of expression across a diel pair. ** Indicates this ortholog was

used for structural modeling and conservation analyses. This list is only a subset of the genes

recovered from various analyses. The genes were chosen because they had either 1) multiple

lines of evidence, robust divergent expression pattern, or 2) a gene ontology term for sensory or

circadian function among three datasets. For exhaustive lists, see Supplementary Data

1173

Process	Search Query	Excluding
Vision	"vision" or "eye" or "visual" or " compound eye" or "visual perception" or "light" or "ocellus" or "visual" or "photoreceptor"	- "chain" - "immunoglobulin" - "light harvesting" - "photosynthesis" - "plastid" -"plant" -actin -myosin -photoperiod - touch -smell -"light chain"
Smell	"olfaction" or "smell" or "antennal" or "olfactory" or "odor" or "pheromone"	
Hearing	"audition" or "hearing" or "sound" or "auditory" or "tympanum" or "ear"	

Circadian	"circadian" or "rhythm" or "clock" or "diurnal" or "nocturnal" or "entrainment"	
Behaviour	"locomotion" or "flight" or "behavioral"	- "cell" - "morphogenesis" - "development" or - "pheromone"
Brain	"neural plasticity" or "brain" or "neuropeptide"	

Table 8: GO query terms for functional lookup