Manuscript type: Research Article

Title: The developmental gene *disco* regulates diel-niche evolution in adult moths

Authors: Yash Sondhi[1,2] (ORCID:0000-0002-7704-3944), Rebeccah L. Messcher[2], Anthony J.

Bellantuano[1], Caroline G. Storer[2], Scott D. Cinel[2], R. Keating Godfrey[2], Deborah Glass[4], Ryan A. St

Laurent[2,6], Chris A. Hamilton[5], Chandra Earl[7], Colin J. Brislawn[2], Ian J. Kitching[4], Seth M. Bybee[3], Jamie

C. Theobald[1], Akito Y. Kawahara[2]

Author Affiliation:

[1]Department of Biology, Florida International University, Miami FL 33174 USA

[2]McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of

Florida, Gainesville, FL 32611 USA

[3]Department of Biology, Monte L. Bean Museum, Brigham Young University, 4102 Life Science

Building, Provo, UT 84602, USA

[4]Natural History Museum, Cromwell Road, London SW7 5BD, UK.

[5]Department of Entomology, Plant Pathology & Nematology, University of Idaho, Moscow, ID 83844,

USA

[6]Department of Entomology, Smithsonian Institution, National Museum of Natural History,

Washington, D.C., U.S.A

[7]Bernice P. Bishop Museum, Honolulu, HI 96817, USA

Corresponding author: Yash Sondhi, yashsondhi@gmail.com

Keywords: circadian; development, diel-niche; gene expression; insects; Lepidoptera; light cycle; RNA-

seq transcriptome, transcription factor

Significance: Insect diel-activity patterns are diverse, yet the underlying evolutionary processes are

poorly understood. Light environment powerfully entrains circadian rhythms and drives diel-niche

and sensory evolution. To investigate its impact, we compared gene expression in closely related

day- and night-active wild silk moths, with otherwise similar ecologies. Expression patterns that

varied with diel activity included genes linked to eye development, neural plasticity and cellular

maintenance. Notably, *disco*, which encodes a zinc-finger transcription factor involved in pupal

Drosophila optic lobe and antennal development, shows robust adult circadian mRNA cycling in

moth heads, is highly conserved in moths, and has additional zinc-finger domains with specific

nocturnal and diurnal mutations. We hypothesize that *disco* may contribute to diversification of adult

diel-activity patterns in moths.

Abstract: Animals shift activity periods to reduce predation, minimize competition, or exploit new

resources, and this can drive sensory system evolution. But adaptive mechanisms underlying niche-

shifts are poorly understood, and model organisms are often too distantly related to reveal the genetic

drivers. To address this, we examined expression patterns between two closely related silk moths that

have undergone temporal niche divergence. We found 200-700 differentially expressed genes,

including day upregulation in eye development and visual processing genes, and night upregulation

of antennal and olfactory brain development genes. Further, clusters of circadian, sensory, and brain

development genes co-expressed with diel-activity. In both species, eight genes showed expression

significantly correlated to diel activity, and are involved in vision, olfaction, brain development,

neural plasticity, energy utilization, and cellular maintenance. We repeatedly recovered *disco*, a zinc-

finger transcription factor involved in antennal development, circadian activity, and optic lobe brain

development in flies. While *disco* mutants have circadian arrhythmia, most studies attribute this to

improper clock neuron development, not adult circadian maintenance. Comparing predicted 3D

protein structure across moth and fly genetic models revealed *disco* likely retained developmental

function with a conserved zinc finger domain, but gained functional zinc finger domains absent in *D.*

melanogaster. These regions have several mutations between nocturnal and diurnal species that co-

occur with higher levels of predicted phosphorylation sites. With robust circadian expression,

functional nocturnal and diurnal mutations, and structural and sequence conservation, we hypothesize

that *disco* may be a master regulator contributing to diel-activity diversification in adult moths.

Introduction

Circadian rhythms regulate many biological processes, including day-night activity patterns.

Research to date has explored genes, circuits, and environmental cues mostly in the context of

control within a single organism (1, 2). Diel-niche stems from the concept of temporal partitioning of

activity periods, driven by the variation in resources, temperature, light level, or predation, across the

day (3–6). Periods are often binned into three distinct groups: diurnal (day active), nocturnal (night

active), or crepuscular (active at dusk, dawn, or both), and although this is useful for comparative

analysis, it is often an overgeneralization (7–11), and the evolution of diverse diel-niches across

organisms (7, 11–15) is poorly understood (16–19).

Several studies have documented sensory adaptation accompanying evolutionary diel-niche

transitions in mammals, birds, and insects (8, 18, 20, 21). Specific examples include colour vision

gene expansions in diurnal moths (22, 23), improved olfactory senses in dark-bred flies (24), and the

evolution of hearing organs in nocturnal butterflies (25). Comparisons of circadian genes across

model organisms catalogue variation at long evolutionary timescales and highlight conserved

elements (26, 27), but diel switches can occur over short timescales (28), and distantly related species

provide less insight into mechanisms that enable faster switches. Moths and butterflies (Lepidoptera)

are an ideal group to address this, as their evolutionary history is well known and there are many

diel-niche switches, often between closely related species (7, 29). They are also one of the few insect

groups, outside of *Drosophila*, in which both sensory and circadian genes have been characterized

(23, 30–35).

Wild silk moths (Saturniidae) are important models for understanding chronobiology, with

seminal experiments confirming the brain functions as the primary circadian control (36, 37). While

most are nocturnal, many fly during the day, and temporal niche switches may function as a

mechanism for reproductive isolation in sympatric species (38, 39). Saturniid phylogeny is relatively

well known, and there are several annotated genomes, chromosomal maps. Furthermore, they belong

to the superfamily Bombycoidea, which includes the domesticated silk moth, *Bombyx mori*, a key

biological model organism serving as a reference taxon for comparative genomic studies (32, 40–43).

Saturniids such as *Antheraea* are of major importance to the silk industry, and they have a well-

resolved taxonomy and well-documented life histories (38, 44–46). Despite this, the genetic elements

that control day-night activity and diel-niche evolution, in this group and insects in general, remain

largely unknown (28, 37).

We used RNA-Seq to characterize expression patterns during peaks and troughs (midday and

midnight) across two closely related wild silk moths: the diurnal *Anisota pellucida*, and nocturnal

Dryocampa rubicunda. They are sister genera, feeding on large deciduous trees in the temperate

forests in Eastern North America, with a recent divergence in temporal niche ~3.8 Mya (47). We

sequenced head tissue at different time periods to identify expression in sensory, circadian, and

neural genes that correlate with diel activity. Expression in eight genes clearly and significantly

correlates with diel activity, with functions in vision, olfaction, brain development, neural plasticity,

energy utilization, and cellular maintenance. Of these, a single gene emerges consistently across

analyses: *disco*, which encodes a zinc-finger transcription factor. In moths, it has likely retained

functions also found in flies (eye and brain development during the pupal stage) through a conserved

zinc-finger domain. But it has also gained an extra zinc-finger domain, surrounded by

phosphorylated sites and with mutations in both nocturnal and diurnal species, which may contribute

to adult moth circadian regulation.

Results

We compared gene expression across two wild silk moth species, *Anisota pellucida* and *Dryocampa*

rubicunda, whose males are diurnal and nocturnal, respectively (Fig. 1, Table S1). We generated

head (eyes, brain) transcriptomes from moths collected and flash frozen at midday and midnight,

referred to as 'day' and 'night' hereafter. Using multiple programs to assemble high-quality *de novo*

assemblies (Table S2), we characterized the level of gene (mRNA) expression with quasi-read

mapping, which can be used to correlate protein expression (48).

Day-night gene expression patterns switch between nocturnal and diurnal species

We found 350 & 393 significantly differentially expressed genes (DEGs) when comparing day and

night treatments for each species (Table S3, Fig. 2A, Supplementary Data 1). *Anisota* had more day-

upregulated genes (56%) and *Dryocampa* was slightly more night-upregulated (53%). In order to

compare DEG sets between species, we mapped our DEGs to *Bombyx mori* orthologs using the

Orthofinder software (49) (Supplementary Data 1). Approximately 60% of DEGs from each species

had identifiable orthologs in *B. mori* (Table S3), and only a small number of DEGs (6-8 genes)

overlapped between both species (Fig. 2B). We also replicated this analysis using other software to

assure that our results were robust, despite different normalization methods (50). With DESeq2, we

found 498 & 697 DEGs (Fig. 2C, Table S3, Supplementary Data 2), with similar *Bombyx* annotation

rates (61%), although the proportion of day upregulated genes increased considerably in the *Anisota*

(79%) compared to being more even split in *Dryocampa* (50%) (Table S3). The total number of

overlapping genes increased (19-26, Fig. 2D) when using DESeq2. A comparison of the two methods

revealed that 174 and 216 genes were shared between *Anisota* and *Dryocampa*, respectively.

Divergently expressed genes are linked to brain optic lobe, antennal and neural development

To identify important regulators involved in diel-niche evolution, we applied two filtering criteria to

our gene expression data. First, we selected genes that exhibited highly significant differential

expression in both species. Second, we focused on genes that displayed upregulation patterns

consistent with the natural diel-activity of each species. Our rationale was that this subset of genes

was more likely to contain key regulators. To compare DEG overlap between the two species, we

grouped the transcripts to their matching orthologs from *Bombyx mori*; if two transcripts from

different species mapped to the same ortholog, we treated them as being the same. This allowed us to

examine overlapping genes between the species to see if any genes switched fold-change sign from

positive to negative or vice versa. (Fig. 2, Supplementary Data 3). We found 51 overlapping DEG

transcripts that mapped to 28 unique *Bombyx* genes. Nine genes showed flipped patterns of

expression between the two species, and eight of them coincided with known diel activity patterns

(Table S7). Examining gene ontology (GO) annotations and comparing orthologs from flybase

(https://flybase.org/), we found genes linked to optic lobe and antennal development (*disco*),

locomotion and energy use (*SLC2A6, SLC17A5*), brain and neural development (*TUBG1*) and other

essential biological processes like transcription, ribosomal translation, protein processing,

mitochondrial maintenance (*RpS4, PARL, Mrps5*) and wound response (*PRP2*) (Table S7, Fig. 2B,

D). Of these, only *disco* was recovered with both methods.

Gene network analysis identifies diel activity and species-specific co-expressed clusters

Identifying highly expressed genes helps understand which genes are activated during particular

biological processes. However, determining only those that are highly expressed can often overlook

genes with important biological functions (51). We examined co-expressed genes that may be

correlated with diel-niche or RNA collection time. We used WGCNA, a weighted correlation

network analysis tool to cluster genes together based on their normalized counts (52). After

examining co-expression patterns for each species separately, we found two modules in each species

that clustered with day-night treatment (cluster-*grey60* and cluster-*tan*) (Fig. S6, Supplementary data

4). Since we were interested in species-specific differences, we reran analyses and combined reads

from *Anisota* and *Dryocampa,* using only normalized counts for genes that had valid *Bombyx*

annotations for both species. This approach narrowed our focus to 2000 genes. Among these, we

discovered two clusters (cluster-blue and cluster-turquoise) consisting of 50 genes each that exhibited

different expression patterns across species (Fig. 4).

GO enrichment of photoperiodism, circadian control, muscle and neural growth genes

We used a gene enrichment analysis to determine if gene ontology (GO) terms were significantly

overrepresented in the DEG and WGCNA sets compared to the appropriate background of GO terms.

Using TopGo, which allows custom gene sets, we found an overrepresentation of genes involved in

several biological functions (Supplementary Data 5). Those that seemed significant for diel-niche and

vision were 'response to stimulus' and 'smooth muscle control', as well as folic acid serine, glycine

and retinoic acid metabolism. We also used ShinyGo to examine WGCNA clusters (Supplementary

Data 5). We matched orthologous genes in the less duplicated, filtered transcriptomes to obtain

identifications of the most related *Bombyx* genes (Table S4). We examined the enrichment of both

tan and grey60 modules, listing the non-redundant terms using ReviGo (Fig. S7, Supplementary Data

5). Gene clusters that co-expressed in the same direction together in the day and night treatments of

both species included photoperiodism, circadian control, negative phototaxis and nervous system

development. We next checked for the enrichment of the modules that showed species-specific

patterns (blue and turquoise, Fig. S8, Supplementary Data 5). These included genes involved in

muscle proliferation and nerve growth, neural signaling, glycolysis, oxidative stress response, and

basic cellular functioning such as protein processing and transcriptional regulation.

Day upregulation of vision genes in the diurnal moth

Since both EdgeR and DESeq2 analyses use different normalization methods and statistical model

assumptions (50, 53), we repeated enrichment analyses by combining datasets and examining genes

that appeared in both analyses. For the *Anisota,* we tested over-enrichment of a smaller subset (FC $\leq$

-5) of diurnally highly upregulated genes (Fig. 3, Table S5). We found gene enrichment for visual

perception, excretion regulation, negative gravitaxis, synaptic plasticity, along with genes associated

with other biological processes, such as RNA interference, endopeptidase activity and endocytosis. A

reduction in stringency (FC ≤- 2) did not alter results considerably (Fig. S4). Night-upregulated

genes (FC ≥ 2) included ocellar pigment genes, eye-photoreceptor cell development, snRNA

processing, post embryonic development and neurotransmitter secretion, among a host of other

processes that may be required for growth, development and metabolism (wnt signaling,

tricarboxylic acid cycle, and cellular response to insulin, glucose transport) (Fig. S4).

Night upregulation of antennal and olfactory brain regions mushroom development genes

We repeated the same analyses for *Dryocampa* and tested highly nocturnally upregulated genes (FC ≥ 5).

Our results show upregulation in genes known to be associated with mushroom body development,

locomotor rhythm, synaptic growth, energy utilization (Sialin transport) and mitochondrial translation

(Fig. 3B, Table S6). A reduction in stringency (|FC| ≥ 2) showed entrainment of the clock cycle, and

antennal development genes. Genes associated with innate immune response, DNA repair, cell division,

histone acetylation, circadian rhythm, retinoid cycle were upregulated during the day, possibly indicating

a period of cellular repair during a time when these moths are inactive (Fig. S5).

Key sensory, circadian, eye development and behavioral genes can drive diel-niche switches

We combined results from the DEG (EdgeR and DESeq2) and gene network analyses (WGCNA) to

create a cumulative list of 1700 transcripts (Supplementary Data 6). Focusing on genes that were

recovered across *Anisota* and *Dryocampa* reduced the set to 274 transcripts (Supplementary Data 7).

Because many transcripts had poorly annotated *Bombyx* hits*,* we improved annotations using the

program eggNog mapper (54). We tested if these genes had GO terms associated with sensory,

circadian, brain and neural development, or behavioral regulatory genes (Fig. 4, Table S7,

Supplementary Data 8). We found that several genes in each category had associated GO terms, with

a predominance of vision and brain development genes (Supplementary Data 9).

Predicted functional regions and homology patterns identified for genes of interest

We examined protein and gene evolution for a set of genes which we found were of interest based on

results from DGE, WGCNA, GO annotations (Supplementary Data 10). In order to infer functional

homology from structural conservation, we downloaded high quality genomes of Bombycoidea

moths and relatives from Darwin Tree of Life [https://www.darwintreeoflife.org/] (Supplementary

Data 11). We assigned a reference protein sequence for each gene of interest from the *Bombyx mori*

predicted proteome. We used Orthofinder to identify orthologs and filtered orthogroups containing

the reference sequence (49, 55). Since we had more than one gene per species, we reduced the

number down to a single gene by choosing the highest identity sequence relative to the *B. mori*

reference sequence. We also modelled the 3D structure of *B. mori* proteins and mapped the

evolutionary conservation onto the 3D predicted structure for proteins above a certain conservation

threshold (Supplementary Data 12). These analyses predict structurally and functional conserved

regions of proteins (Fig S9, Supplementary Data 13). We repeated this analysis with 38 insect

genomes (Supplemental Data 11) and mapped evolutionary conservation onto 3D protein structure

(Supplementary Data 12-13). We include results of evolutionary conservation analyses for two

regulatory candidates (*disco* and *tk*) that showed varying levels of sequence and protein evolution

between insects and moths (Fig. S10).

Modeling predicts additional functional zinc-finger domains for disco in Lepidoptera

Disco was recovered across multiple analyses. In *Drosophila melanogaster* it is known for its role in

eye development, important for circadian maintenance, and for leg and antennal appendage

formation (56–60). To determine if *disco* was conserved between moths and *Drosophila*, we

compared the primary sequence and 3D protein structure of *Drosophila* and *Bombyx mori*. In the

moth, the sequence length of *disco* was nearly double that of *Drosophila* (Fig 6A). However, a region

spanning over 100 amino acids was highly conserved, contained the zinc-finger domain important for

its function, and showed strong 3D structural conservation (*Whole protein alignment*: RMSD: align,

super) = 37.833, 2.566, MatchAlign score (align, super) = 540 (2234 atoms), 333.1 (554 atoms) vs.

alignment of conserved region: RMSD=2.699,2.566, MatchAlignScore =443(671 atoms), 351 (615)

atoms). This indicates the DNA binding function of *disco* has likely been conserved. However, an

additional ~500 amino acid region absent in *Drosophila* is highly conserved across moths, and

includes several regions predicted to be functional (Fig. 6A, Fig. S10). We hypothesize that *disco* has

a novel role in moths for diel-niche regulation. To further test this, we compared *disco* sequences

across *Anisota* and *Dryocampa* and found 23 mutations between them, three of which mapped to the

predicted functional region (Fig. 6B, Supplementary Data 14). InterProScan predicted four zinc-

finger domains, three in this region, although the CATH-Gene3D databases prediction combined the

two separate domains into a single predicted domain (Fig. S11). We also found 53 sites with

predicted phosphorylation potential, especially around the second zinc-finger domain (18). Reducing

the stringency increased the total to 142, which were still enriched around the second domain.

Discussion

We identified the genetic mechanisms of diel-niche switches by examining gene expression during

the day and night in two closely related moth species. We used RNA-Seq to measure gene expression

profiles of head tissue and isolated sensory, circadian and neural processes. We found between 300

and 700 genes that significantly altered day night circadian expression patterns. The diurnal species

had enriched visual perception genes during the day and the nocturnal species had locomotor and

olfactory genes upregulated at night.

Thirty overlapping DEGs were present in both species, with some DGEs showing divergent

patterns of expression matching the species' diel-niche. Examining expression data with a sensitive

clustering analysis yielded over 170 genes that showed day-night or species specific co-expressed

clusters. We also used GO terms to search for genes associated with sensory, neural, or circadian

patterns. We found genes in each category and modeled the 3D structure and predicted functional

regions for a subset. The divergently expressed gene *disconnected* (*disco*) was implicated in vision,

hearing, locomotion and brain development. Given the multiple lines of evidence supporting the

importance of this gene in regulating diurnal and nocturnal activity in moths, we explored the

evolution of *disco* by modeling fly and moth *disco* structures. This analysis revealed novel zinc-

finger conserved domains in moths, which were lacking in *Drosophila,* surrounded by

phosphorylated sites. Several mutations between *Anisota* and *Dryocampa* also mapped to these

regions, further strengthening evidence for its role in diel-niche shifts.

Visual and olfaction

Visual systems often accompany diel and photic environment shifts (18). For example, nocturnal

carpenter bees have much larger facets in their eyes than their diurnal counterparts (61). Nocturnal

moths have higher photoreceptor light sensitivity and have neurons more suited to pooling than close

diurnal relatives (62). Some of these trends seem to hold across the phylogeny, with diurnal species

evolving more complex color visual systems, often reflected in visual opsin sequence evolution (22,

23). We did not find diel-expression patterns in color vision opsins, a result corroborated by a recent

study (22). However, we discovered a cerebral opsin (ceropsin) upregulated during the day. Ceropsin

has been implicated in photoperiodism and is expressed in the brain (63).

We also found several eye development genes (*TENM2, ANKRD17, EHD4, JAK2*),

phototransduction genes (*PPAP2*, *RDH11*), and retina homeostasis, eye-antennal disc development,

and photoreceptor cell maintenance genes (*disco, glass*) (64). Surprisingly a few visual genes, such

as *garnet* and *rugose* also appeared to have different isoforms present, showing both day and night

upregulation. *Garnet* is an eye color mutant gene in flies (65) and *rugose* is implicated in retinal

pattern formation (66).

One of the classical differences between day flying butterflies and night flying moths is that

butterflies have a clubbed antennae, while moths often have a sensilla-covered antenna; likely a

function of increased investment in olfactory sensing, similarly, there is also evidence for increased

investment in olfactory sensing in the brain, with relatively larger mushroom bodies in nocturnal

species (67, 68). In our diel gene-expression dataset, we found several olfactory genes, including

those involved in odorant binding (*Obp84a, Obp58b*), pheromone response (*tk*), mushroom body

development (*DAAM2 and DST*) and antennal development (*disco*).

Hearing and mechanoreception

Many moths have active hearing organs, and these have evolved repeatedly across Lepidoptera (69),

and their use varies depending on the ecological context of the moth. Several drivers for moth

hearing include sexual selection and as a defense against insectivorous bats (70). The primary

nocturnal group of butterflies, Hedylidae, has reverted to a nocturnal niche and regained hearing

organs (25). Thus, although rudimentary vibrational hearing exists in some diurnal moths (71) it is

significantly more developed and used at broader frequency ranges at night, when light is less

available. While Saturniidae lack hearing organs (72) they may instead still have vibration

mechanosensors that can be useful in evading predation similar to those in locusts and crickets (73–

76). Indeed, we did find diel co-expression of several mechanosensory and ear development genes,

including *PI4KB, KCTD15, unc-22,* and *RhoGAP92B* (77)

Brain and neural rewiring

Finding an upregulation of neural and brain development was expected since we sequenced head

transcriptomes. However, we tested gene expression two days after pupal eclosion, where

presumably gross structures are already developed, so it is possible these expressed genes we found,

regulate adult plasticity. Adult neural plasticity has been showed in Lepidoptera and other insects

(78–80), and we recovered several genes linked to axon regeneration (*APOD*), central complex

development (*Ten-a, TENM2, DST, ALDH3A2, OGT*), central nervous system development (*disco,*

RpL4), and neuropeptide hormonal activity (*tk*). Several genes were specific to retinal ganglion cell

axon guidance (*TENM2*) and mushroom body development (*DAAM2*), leading us to speculate that

plasticity could occur through neural wiring and plasticity are shaping sensory adaptation. Many

Lepidoptera have distinct phenotypic plasticity in wing patterns and coloration in different seasons

(81), and there is some evidence that they also have seasonal plasticity in their behavior and foraging

preferences (82). Research has found Lepidoptera can override their innate preferences and learning

preferences for new visual and olfactory cues after eclosion (83). It is possible the diel-niche and

circadian rhythms too have plasticity, there are reports of some moth species like *Hyles lineata*

showing relatively labile diel-niches possibly driven by temperature and resources (84–86). These

same mechanisms allowing flexibility within a species, might be involved in diel-niche evolution

between species.

Circadian and behavioral regulators

The behavioral state of an animal to engage in any activity, e.g., feeding, flight or mating, is likely a

function of circadian and behavioral regulators that respond to certain stimuli. We found differential

expression of genes involved in locomotion (*unc-22, KCTD15,Tk*) and circadian or rhythmic

behavior (*SREBF1/SREBF2, OGT, disco, JAK2*) in both species. We also found several key circadian

regulators like *per*, *tim* although they were downregulated only in *Dryocampa*. C*lock-like* or *takeout*

another gene under circadian control, (87) was expressed in both species, although without any

significant up or down regulation. *Takeout* was moderately conserved in the moths we examined, but

had diverged considerably among other insects, to the point where orthology searches with

Orthofinder failed to recover orthologs among insects. Even among *D. melanogaster*, its closest

homologs were *Jhbp5* and *takeout* which had 21-25% sequence identity (Supplementary Data 10,14).

Surprising, it's 3D structure was highly conserved (RMSD (align, super) = 2.556, 2.755,

MatchAlign score (align, super) =199 , 642.937, Supplementary Data 14, Fig. S12), highlighting an

interesting example of likely functional convergence despite primary sequence divergence.

Energy use and general cellular maintenance

In addition to the genes mentioned above, several key divergently expressed genes were involved in

energy utilization. This finding makes biological sense, as once activity is initiated, energy

mobilization and upregulation of basic cellular processes need to be maintained. Two of these genes

were *SLC175A/MFS10* and *SLC2A6/Tret1-1*, which encode different trehalose sugar transporters.

Trehalose is an important sugar present in insect hemolymph (88). Other genes found were ribosomal

(*RpL4, mRpS5*), golgi (*GCC*), and mitochondrial maintenance (*PARL/rho7*) along with

transcriptional regulation (*Spt5*).

Disco as putative master diel-niche regulator in adult moths

Many genes that we discovered likely play a critical role in maintaining sensory shifts, but in order to

identify genes that might regulate upstream, we looked for genes that were: 1) expressed in both

species, 2) showed coincident expression patterns with respect to diel-niche, and 3) played a role in

sensory and neural functioning and circadian control. Only *disconnected* (*disco*) fit these criteria.

Disco is a key developmental and patterning gene, first discovered for its role in neural migration

from the disconnected optic lobe mutant phenotype (89). This gene is now considered an appendage

development gene, with a well characterized pupal role in antennal and leg distal patterning in

Drosophila melanogaster (60). Mutants also have a disrupted circadian locomotor rhythm due to the

improper formation of neurons that express clock genes (90, 91). As a gene that is involved both in

optic lobe, antennal formation, and in neural wiring and circadian control, it is a very strong

candidate for driving behavioral diel-niche shifts and sensory adaptations (56, 57, 92). *Disco*

expression data from *B. mori* suggests strong adult head, antennal, and nucleus expression,

suggesting that it still acts as a transcription factor, and can regulate other genes (40).

Disco encodes the acid zinc-finger transcription factor and is approximately 500 and 1000

amino acids in length in *Drosophila* and *B. mori,* respectively. Our modeling showed that a ~150

amino acid region is conserved structurally at the sequence level, and this region also contains zinc-

finger motifs associated with DNA binding (93). This indicates that *disco* likely has retained its DNA

binding and likely pupal and appendage patterning function in moths. We predicted the functional
regions of *disco* in *Bombyx* based on evolutionary conservation modelling, and found an additional
500 amino acids, absent in *Drosophila*, were predicted to be functional (conserved and exposed).
Domain and family level modeling predicted at least two additional zinc-finger domains in this
region (Fig. S10,Fig. S11). We also found many phosphorylated sites surrounding the zinc-finger
domains. Examining mutations between *Anisota* and *Dryocampa* revealed three mutations mapped to
these predicted functional regions (Fig. 6B).
Given *disco's* adult diel-specific expression in moths, additional zinc-finger DNA binding
domains, and the high number of phosphorylated sites, we propose it as a candidate transcriptional
regulator for diel-regulation in adult moths.
372

Study species and their evolution and variation in life histories
Anisota and *Dryocampa* are closely related and ecologically similar saturniids of the subfamily
Ceratocampinae, thought to have diverged ~3.81 Mya (47). They largely occupy the same range and
forest habitat in the Americas, feeding on large trees (*Anisota* on oaks and *Dryocampa* on maples)
where population flares cause host plant defoliation (38). Because they reportedly hybridize (94), and
pre- and postzygotic barriers are potentially ineffective, temporal partitioning may be important in
their evolutionary history.
380

Conclusion

This study provides a framework for how diel-niche evolution in Lepidoptera can occur.
Sensory and neural developmental genes appear to be key. We identify the pivotal role of the
transcription factor *disco*, uncovering a second functional region with species-specific mutations in
moths. Our findings provide useful targets for further genetic manipulation and highlight the insight
non-model systems provide to genetics and development.
387

388 ## Supplementary information

389   Supplementary tables and data are available for the reviewers and will be uploaded to repository
390 [xxxxx] on submission.
391   .

Author contributions

Y.S.: conceptualization, data curation, formal analysis, investigation, methodology, project administration, software, visualization, writing–original draft. R.M: investigation, methodology, project administration, data curation, writing - review & editing. A.J.: conceptualization, formal analysis, writing - review & editing C.S.: conceptualization, data curation, methodology, project administration, supervision. S.C.: formal analysis, software. K.G.: formal analysis, software, visualization, writing-review and editing, D.G.: investigation, project administration, data curation. R.L.: conceptualization, data curation, investigation, methodology, writing-original draft, writing -review & editing. C.H.: conceptualization, data curation, investigation, methodology, project administration, writing - review and editing I.K.: conceptualization, funding acquisition, methodology, resources, writing–review and editing. C.E.: formal analysis, software, visualization, data curation, writing- original draft, writing- review & editing. C.B.: software, validation, writing -review and editing. S.B.: funding acquisition, methodology, validation, writing–review and editing. J.T: conceptualization, funding acquisition, supervision, validation, writing–review & editing. A.K: conceptualization, methodology, resources, writing–review and editing, visualization, supervision, funding acquisition

Acknowledgements

We thank Jesse Breinholt, Martijn Timmermans and Andreas Zwick for help with project conceptualization. Kelly Dexter, Amanda Markee, and members of the McGuire Center for Lepidoptera and Biodiversity at the University of Florida assisted with animal care and wet-lab troubleshooting. We thank Belinda Pinto, Danielle DeLeo, Heather Bracken-Grissom, Jorge Perez-Moreno, Zhou Lei for discussions, advice and assistance with bioinformatics pipelines and wet-lab protocols. We thank Nicolas Alexandre, Sachit Daniel, Riddhi Deshmukh, David Plotkin, Chelsea Skojec and Nitin Ravikanthachari for comments and feedback on the manuscript. Janelle Nunez-Castilla, Jessica Liberles, Raghavan Venket, Yi-Ming Weng, Yu Fahong and Xiaokang Zhang assisted with troubleshooting bioinformatic pipelines and provided analytical recommendations. The authors acknowledge the University of Florida Research Computing for providing computational resources and support that have contributed to the research results reported in this publication (http://researchcomputing.ufl.edu). We acknowledge NBAF and UF-ICBR for assistance with sequencing and generating libraries.

Funding for the project was through NSF DEB-1557007, NSF PRFB-1612862, and NSF IOS-1750833 and NERC NE/P003915/1. The Florida International University Presidential fellowship supported YS.

Figure Legends and Figures:

Figure 1: Nocturnal and diurnal moths on a phylogeny with RNA-seq sampling design.

(A): Collapsed phylogeny of Saturniidae, adapted to show where the two study species, diurnal *Anisota pellucida* (Pink-striped oakworm moth) and nocturnal *Dryocampa rubicunda* (Rosy Maple moth) sit. Faded numbers on the tip represent the number of genera in the tree before collapsing. Phylogeny adapted from Rougerie et al. 2022 (**B**): Sampling design showing the number of replicates for each species and collection period (day/night) sampled. Collection of heads was done 2-days post eclosion at midday (sun) and midnight (moon) and tissue was flash frozen for RNA preservation. Care was taken that the eclosed moths were exposed to a natural light cycle and red lights were used when collecting the moths at night. Photo credits *Anisota pellucida* © Mike Chapman; *Dryocampa rubicunda* (CC) Andy Reacgo and Chrissy Mclearan;

Figure 2: Nocturnal and diurnal species show divergent patterns of gene expression across two different analyses software (EdgeR and DESeq2)

Volcano plots and Venn diagrams showing EdgeR (A-B) and DESeq2 (C-D) results across **day vs. night** sampling times for both nocturnal and diurnal species. Venn diagrams represent the number of differentially expressed genes (DEGs) across both species with the number of common DEGs across each pair. **(A/C)**: Volcano diagrams illustrating fold change and adjusted p-values for the significant differentially expressed genes between midday and midnight samples. Circles in the top left represent the number of genes expressed in both species and the colors correspond to FC values for those genes in the volcano plots. Yellow and blue represent genes expressed only in the nocturnal or diurnal species, green/purple indicates DEGs present in both species. **(B/D)**: Only genes expressed in both species are shown and genes that display opposite trends in expression are highlighted. See Supplementary Material for a detailed list of DEGs. Positive fold change indicates night overexpression and negative fold change indicates day overexpression. Genes had FDR or adjusted p-value < 0.05 and FC > |2|. Identification of common genes was done using orthogroup clustering with Orthofinder with *Bombyx mori*. Gene names and annotations were transferred from *B. mori*.

470
Figure 4: Modules of clustered co-expressed genes grouped using normalized expression
highlighting the two modules that showed species specific clustering patterns.
A: Cluster dendrogram shows WGCNA clusters. B: Shows how patterns of gene expression correlate

across samples and modules. C: Shows the normalized expression for all
genes across samples. The species show two sets of 50 genes (Blue) and (Turquoise) that have clear
species specific expression patterns. Normalization was done with DESeq2 and reads were mapped
to the more stringently filtered transcriptome. A soft power analysis was done and the picked
power=9 for the WGCNA analysis. AN: *Anisota*, DR: *Dryocampa,* Numbered D and N represent the
different samples collected at day and night time points.

480
Figure 5: Model for diel-niche shift from and genes of interest obtained from GO searches and
annotating divergently expressed genes. For representative purposes, homologs from *Drosophila*

483 *melanogaster* have been used, with colors in the 3d structures representing Alpha fold per-residue
484 confidence metric (pLDDT), the range for each color is shown on the top left. Model moth *(Bombyx*
485 *mori)* proteins were also modeled for a subset of proteins, see Supplementary data.
486

487
488

Figure 6: *Disco* has 150 amino acid long highly conserved region across *Bombyx* (1054 aa) &
Drosophila (568 aa) involved in its role as a transcription factor, but it also has other roles in
Lepidoptera due to the presence of other highly conserved and predicted functional regions, that are
not present in flies, and only in Lepidoptera. 3/14 of the mutations between nocturnal to diurnal
species map to these regions (>500 aa).
(A) Top: *Disco* best Uniprot hit in *Drosophila* using default settings (blastp, e-threshold 10, Auto-
Blosum62). Bottom: *disco* aligned and superimposed predicted structures for the silk moth and
vinegar fly. Peach structure: *Bombyx mori* alpha-fold predicted structure for *disco,* Cyan: *Drosophila*
melanogaster alpha-fold predicted structure for *disco* **(B)** Top: Partial views of the Consurf predicted
residues for *disco.* The views encompassing the region where 3 mutations of the 14 mutated sites
between Nocturnal *Dryocampa* and Diurnal *Anisota* overlap with predicted functional residues.
Bottom: Residues that were mutated between the nocturnal and diurnal species are highlighted on an
overlay of the Consurf scores mapped onto the predicted alpha fold structure of disco from *Bombyx*
mori. Green residues indicate mutated and predicted functional sites, red indicate mutated sites that
did not have a predicted residue. **(C)** Overlap of the highly conserved region of *disco* this region
includes the zinc-finger domain that is characteristic of disco transcription factor.

Methods

Insect rearing:

Moths were reared under natural light-dark cycles at room temperature (25°C) and natural light
conditions at the McGuire Center for Lepidoptera and Biodiversity (MGCL). Adults were sampled
two days after eclosion at the two time points by C.H. and R.S.L.

Sampling design:

We collected 3-5 replicates (i.e., individual moths) per experimental treatment. We sampled at two
time points per species at two midpoints of their circadian cycle at noon ('day') and midnight
('night') (Fig. 1, Table 1). Moths were allowed to eclose naturally and sexed, To ensure vision genes
were not artificially stimulated, for collection at night we used red light, which is thought to not
stimulate most Lepidoptera insect visual systems (95). We used a sterilized scalpel to remove the
head, which was placed in a 1.5 ml Eppendorf tube, which was immediately placed in liquid nitrogen
and stored at -80°C. It is worth noting that only males of most *Anisota* species are diurnal, including
A. pellucida, our focal taxon. Females of all *Anisota* and both sexes of at least *A. stigma*, are
nocturnal. In *Dryocampa*, both males and females are nocturnal. We therefore limited our
transcriptomic work to males of *A. pellucida* and *D. rubicunda*.

RNA extraction and quality control:

Tissue was removed from liquid nitrogen and placed into a flat-bottomed cryo-tube with Trizol and
2-4 sterile, stainless-steel beads. The tissue was homogenized using a bead beater for 2.5 minutes at
1900 RPM. Beads were removed with a magnet, and chloroform added to separate phases.
Isopropanol precipitated the RNA, followed by several ethanol wash steps to pellet the RNA. Quality
Control (QC) was undertaken with a high sensitivity Qubit assay. Samples with yields > 20 ng/μL
were cleaned using a Thermo Scientific RapidOut DNA Removal Kit (Waltham, Massachusetts,
USA). Sample quality was verified using Agilent 2200 Tapestation. Since heads comprised of a
relatively small amount of tissue, initial extractions using kit-based methods resulted in low yields.
Several rounds of troubleshooting were required before completion of extraction using a modified
Trizol based protocol with a DNA clean up step. We measured the quantity of RNA using a Qubit

fluorometer and used Agilent tapestation or Bioanalyzer to characterize extraction quality, weight, and size distributions. Where possible, a concentration of more than 10 ng/µL was used for library preparation. Yields below 10 ng/µL were prone to higher loss when purifying with a DNAse agent, which often caused 20-30 % loss in RNA yield, approaching 5ng/µL is below the Qubit detection threshold.

Library preparation and NGS sequencing:

Libraries were prepared by sequencing cores at NBAF-Liverpool using the NEB polyA selection and NEBNext Ultra II directional stranded kit, suitable for low input yields. 12 samples were run on one lane of an Illumina NovaSeq using SP chemistry (Paired-end, 2x150 bp sequencing). Samples were shipped overnight from MCGL to the NERC-NBAF after dehydrating in a biosafety chamber using GenTegra RNA tubes.

Read trimming and cleanup:

For *Anisota pellucida* and *Dryocampa rubicunda*, trimming was undertaken by the NERC-NBAF core and the raw Fastq files were trimmed for the presence of Illumina adapter sequences using Cutadapt version 1.2 (96). The option -O 3 was used, so the 3' end of any reads that match the adapter sequence for 3 bp or more were trimmed. Trimming was also done with trimmomatic. The reads are further trimmed using Sickle version 1.200 with a minimum window quality score of 20. Reads shorter than 15 bp after trimming were removed.

Transcriptome library sizes, QC, and PCA:

Quality Control (QC) was conducted on the trimmed reads, library size varied for each species (Fig. S1). We examined expression data and removed genes with TPM < 1. PCA results showed that *Anisota* and *Dryocampa* did not separate with diel-niche (Fig. S3).

***De novo* assemblies:**

We combined reads from multiple samples and generated several reference de novo transcriptomes using different assemblers, and then combining them, compared their quality using BUSCO scores and number of single-copy BUSCO genes across the different versions (see Table 2 for the BUSCO

scores, redundancy and duplication for the various assemblies). We found that unfiltered assemblies

were highly redundant, but had the highest BUSCO scores. These assemblies were too

overrepresented to be used in any downstream analysis, so we used Transrate, CD-HIT, MMeqs2, as

well as Transdecoder, to filter the assemblies. We chose v5 and v6 to use in downstream analysis,

listed as the last two assemblies for each species (Table 3) and repeated certain analyses with both

assemblies to see how stringent or a less stringent filter would affect the results.

Multi assembler *de novo* transcriptome assembly:

To our knowledge, there is no publicly available annotated reference genome for the two species

used in this study at the time the analyses were performed; therefore, we chose to build a de novo

transcriptome assembly with the pre-processed reads using the NCGAS pipeline, which combines

multiple assemblers and then uses an evidence-based gene modeler to decide on the final set. For

each species, we combined forward and reverse reads from all samples and normalized them using a

normalization script in Trinity v2.12.0 (97). Normalized reads were used as input to generate

assemblies using Velvet v1.2.10 (98), TransAbyss v2.0.1 (99) , SOAPdenovo v1.03 (100) with

various kmer sizes. These assemblies were combined and filtered using Evigenes v4: 2020, March

(101, 102), transdecoder and filtered with scripts provided in the NCGAS pipeline (103).

Transcriptome QC, annotation and filtering:

Further quality control and filtering was conducted using BUSCO (104), transrate (105), QUAST

(106) (v5.02), CD-HIT(v4.6.8) (107, 108)and cascaded clustering with MMseqs2 (v12) (109).

BUSCO scores were greater than 95%, but many genes had multiple versions from the different

assemblers, and because this would bias the downstream analyses we attempted to reduce the

redundancy at the cost of lowering BUSCO scores. Protein coding regions were predicted using

TransDecoder (https://github.com/TransDecoder/TransDecoder) and the cds file from was run

through Transrate (105), which further pulled contigs based on mapping rate and identified a "good"

collection of transcripts, the resulting protein-coding sequences were filtered with additional CD-HIT

(107, 108)and MMseqs2 (109) filtering. This resulted in lower BUSCO scores 75-85% but also much

lower redundancy in the gene set of about 1-3 fold. We ran the WGCNA (52) and GO enrichment

(110) analyses on the more stringent assemblies (BUSCO 75%, 2% genes with duplication) since

they are more sensitive to redundancy, but RNA-seq was run across the slightly lower stringency

analyses to be more inclusive (BUSCO~85%, 60-70% genes with duplications).

Annotation of reference transcriptomes:

Reference genomes were annotated with eggNOGmapper and Orthofinder (54, 55). We also tried

annotating the reference transcriptomes with Trinotate (111), which run against the swissprot and

pfam databases using hmmscan, blastx, blastp, signalP and tmmhmm, KEGG, and GO and eggNOG

(49, 110, 112–120). ORF predictions for orthofinder were obtained using *Antheraea pernyi* as a

model. Orthofinder was run with well-annotated Bombycidae and Saturniidae moths, *Bombyx mori*,

Antheraea pernyi, and *Antheraea yamamai*, to generate clusters. *Bombyx mori* was a useful reference

since the Orthofinder cluster for each species generated a list of mostly single-copy genes for which

more Lepidoptera annotations were available from SilkDBv3 (40). While this was not as complete as

trinotate, the annotations provided from this procedure were more accurate than Trinotate. Trinotate

annotation had many more human/ vertebrate hits than insect matches than eggNOGmapper and

Orthofinder transferred annotations, so while the files are provided as a reference, they were not used

in downstream analyses.

Differential gene expression (DGE):

Reads were mapped using Salmon (121) and differential gene expression analysis was conducted

using EdgeR (122) and DESeq2 (123). EdgeR(v 3.38.1) was used to normalize and test for

significantly expressed genes. DESeq2(v1.36.0) was also used to normalize and to generate PCAs

and other summary statistics. We used 'Day' as the treatment and 'Night' as the control for all

groups with EdgeR, although the order of fold change switched for DESeq2. The p-values obtained

were adjusted to account for multiple hypothesis testing with FDR for EdgeR and adjusted p-value

from DESeq2. Genes with FDR or adjusted p-values <0.05 and |FC| > 2 were used as a criteria to

identify significantly expressed genes.

Gene enrichment analysis:

We performed gene enrichment analysis using GO terms with the tools TopGO v2.48.0 (124) and

ShinyGo v0.75c (125) (http://bioinformatics.sdstate.edu/goc/). This analysis involved comparing the

selected genes of interest to a gene universe with GO term annotations to determine if there were

overrepresented or underrepresented GO terms. To obtain genes with GO annotations, we utilized

similarity clusters to map annotations from *Bombyx mori*.

In TopGO, we employed a more stringent approach by using filtered transcriptomes to

mitigate bias from gene duplications. We selected genes of interest based on a False Discovery Rate

(FDR) threshold of less than 0.05. We ran two different enrichment algorithms, 'classic' and 'elim',

and used two test statistics, Fisher's exact test and a Kolmogorov-Smirnov-like test. The tests were

performed for both 'Biological Process' (BP) and 'Molecular Function' (MF) categories. While we

generated tables of significantly enriched GO terms, obtaining individual genes within each group

was limited due to the constraints of TopGO with custom annotations.

For ShinyGO, we utilized the custom v0.75c which included the updated *Bombyx mori*

genome stringent. For all analysis, we used the following settings (pathway size: min.=1,

max=2000). The settings for number of pathways to show and FDR cut-offs were chosen to represent

the entire list of top 100 genes in the final datasets, although in some cases a fewer number are

represented in the visualizations. We focused on the 'Biological Process' (BP) category for the

various tests. Tables of GO terms, gene information, and graphs summarizing the significantly

enriched GO terms were generated. We repeated several different analyses **1)** A less stringent

analysis was run where genes with FC<|2| and padj. <0.05. This was useful to visualize the genes up

and downregulated functionally and the background used was all the genes recovered from the DEA

analysis that had orthologs. An FDR cut-off <0.1-0.3. For the background gene set, we used the

species' de-novo transcriptome matching *Bombyx* ortholog set.

2) A more stringent criteria was used to examine highly upregulated genes a FC<|5| and padj.

<0.05 cutoff was used, and filtering only genes that occurred in both EdgeR and DESeq2 analyses.

We selected a single representative when multiple Bombyx orthologs were recovered, using the

ortholog with the most complete annotation. An FDR cut-off <0.17 was used. For the background

gene set, we used the species' de-novo transcriptome matching Bombyx ortholog set.

3) For WGCNA clusters tan and grey60, we used their respective species background

expressed transcript set with a FDR cutoff <= 0.1. For blue and turquoise, the background of all

orthologs found in *Bombyx mori* was used. An FDR cutoff<1 was used, since smaller values

provided insufficient genes given the larger number of genes being tested.

Gene network analysis:

Gene network analysis was undertaken with WGCNA (1.71) (52, 126). DeESeq2 was used to
normalize reads and data was formatted with tidyverse. WGCNA identifies modules of co-expressed
genes. We chose modules that showed clear day-night differences and mapped the GO terms with
Revigo and ShinyGo. We also combined counts from both species and tested for clusters. Here we
chose modules that showed species specific clusters.

Gene annotation and filtering:

DEGs, enriched genes and or co-expressed transcripts/ genes were annotated by cross referencing
them with the *B. mori* annotations (see above). Since *B. mori* mappings were not always 'one-to-
one', mappings that were 'one-to-many' were dropped, i.e., transcripts that mapped to multiple
Bombyx genes were dropped. We also improved annotation by mapping the overlapping transcripts
with eggnog mapper. We further filtered these genes into sub functional categories, by identifying
genes with GO annotations or descriptions linked to vision, smell, hearing, circadian, behavior and
brain the list of go terms for each group was obtained by querying amigo
(http://amigo.geneontology.org/) (Table 8).

Gene Mining and in-silico evolution:

We mined genes of interest from ~bombycoidea moths and closely related families that had well
annotated genomes on Ensembl and NCBI (Supp Table 6) and across model insects (Supp. Table 7).
These genomes were chosen as all Bombycoidea from Ensembl with well annotated assemblies and
five species Noctuidae and Geometridae each were included as close relatives. We added *Bombyx*
mori as a reference and *Antheraea pernyi* to represent Saturniidae, and their peptide files were taken
from their respective source papers (40–42). For insects, we chose long-read well annotated genomes
from Ensembl (Ensembl Gene build) and five model systems from insect base (127). We ran
Orthofinder (v2.5.2) (49, 55) with 'orthofinder -f bom-rel/ -S diamond -M msa -A mafft -T fasttree -a
1 -X -z -t 16' to recover orthologs. Each orthogroup often contained multiple sequences per species.
We chose the sequence with the highest identity to the reference *Bombyx mori* sequence using
custom python scripts. In the rare instances where there were multiple references, we took the longest

one. To model the 3D structure of each reference *Bombyx mori* sequence, we ran AlphaFold v2.1.2

(128) using Deepmind's run_alphafold.py

(https://github.com/deepmind/alphafold/blob/main/run_alphafold.py) on University of Florida's

HiperGator. Model database files (i.e. BFD, MAGnify, PDB70, mmCIF PDB, UniRef30, UniRef90)

downloaded from Deepmind in February 2022 were used as parameter value inputs. All other

parameters were used with default settings. The relaxed predicted model with highest ranked pLDDT

(per-residue estimate corresponding to model's predicted score on the lDDT-Cα metric) was chosen

as the final model (129).

We calculated a conservation score for each alignment using Alistat. For alignments less than 1500

amino acids and with high conservation score (Proteins larger than 1300 aa were difficult to model

without high memory GPU's), we modeled conservation using Consurf (130–134). The results were

displayed using Jmol first glance viewer (http://firstglance.jmol.org/) and as screenshots from the

default viewer. The output for alpha fold runs and consurf is available in the (Supplementary Data).

PyMol (135) was used to align the 3D structures and Aliview was used to compare alignments. We

used InterProscan (https://www.ebi.ac.uk/interpro/about/interproscan/) and NetPhos3.1 (136, 137)

for predicting the domains and phosphorylated sites.

References

1. Y. Xiao, Y. Yuan, M. Jimenez, N. Soni, S. Yadlapalli, Clock proteins regulate spatiotemporal
organization of clock genes to control circadian rhythms. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

2. K. Beer, C. Helfrich-Förster, Model and Non-model Insects in Chronobiology. *Front. Behav.*
Neurosci. **14**, 601676 (2020).

3. D. Székely, D. Cogălniceanu, P. Székely, M. Denoël, Adult-Juvenile interactions and temporal niche
partitioning between life-stages in a tropical amphibian. *PLoS One* **15**, e0238949 (2020).

4. D. Wang, G. Yang, W. Chen, Diel and Circadian Patterns of Locomotor Activity in the Adults of
Diamondback Moth (Plutella xylostella). *Insects* **12** (2021).

5. J. H. Fullard, N. Napoleone, Diel flight periodicity and the evolution of auditory defences in the
Macrolepidoptera. *Anim. Behav.* **62**, 349–368 (2001).

6. R. Refinetti, The diversity of temporal niches in mammals. *Biol. Rhythm Res.* **39**, 173–192 (2008).

7. A. Y. Kawahara, *et al.*, Diel behavior in moths and butterflies: a synthesis of data illuminates the
evolution of temporal activity. *Organisms Diversity and Evolution* **18**, 13–27 (2018).

8.   D. T. C. Cox, A. S. Gardner, K. J. Gaston, Diel niche variation in mammals associated with
expanded trait space. *Nat. Commun.* **12**, 1753 (2021).

9.   Y. Basset, N. D. Springate, Diel activity of arboreal arthropods associated with a rainforest tree. *J.
Nat. Hist.* **26**, 947–952 (1992).

10.   P. J. Devries, G. T. Austin, N. H. Martin, Diel activity and reproductive isolation in a diverse
assemblage of Neotropical skippers (Lepidoptera: Hesperiidae). *Biol. J. Linn. Soc. Lond.* **94**, 723–
736 (2008).

11.   S. R. Anderson, J. J. Wiens, Out of the dark: 350 million years of conservatism and evolution in diel
activity patterns in vertebrates. *Evolution* **71**, 1944–1959 (2017).

12.   S. Frey, J. T. Fisher, A. C. Burton, J. P. Volpe, Investigating animal activity patterns and temporal
niche partitioning using camera-trap data: challenges and opportunities. *Remote Sens. Ecol. Conserv.*
3, 123–132 (2017).

13.   R. Steen, Diel activity, frequency and visit duration of pollinators in focal plants: in situ automatic
camera monitoring and data processing. *Methods Ecol. Evol.* **8**, 203–213 (2017).

14.   H. K. Gill, G. Goyal, R. McSorley, Diel Activity of Fauna in Different Habitats Sampled at the
Autumnal Equinox. *Fla. Entomol.* **95**, 319–325 (2012).

15.   B. Murugavel, A. Kelber, H. Somanathan, Light, flight and the night: effect of ambient light and
moon phase on flight activity of pteropodid bats. *J. Comp. Physiol. A Neuroethol. Sens. Neural*
Behav. Physiol. **207**, 59–68 (2021).

16.   H. Somanathan, *et al.*, Nocturnal bees feed on diurnal leftovers and pay the price of day – night
lifestyle transition. *Front. Ecol. Evol.* **8** (2020).

17.   R. M. Borges, H. Somanathan, A. Kelber, Patterns and Processes in Nocturnal and Crepuscular
Pollination Services. *Q. Rev. Biol.* **91**, 389–418 (2016).

18.   S. M. Tierney, *et al.*, Consequences of evolutionary transitions in changing photic environments.
Austral Entomology **56**, 23–46 (2017).

19.   W. T. Wcislo, *et al.*, The evolution of nocturnal behaviour in sweat bees, Megalopta genalis and M.
ecuadoria (Hymenoptera: Halictidae): an escape from competitors and enemies? *Biol. J. Linn. Soc.*
Lond. **83**, 377–387 (2004).

20.   A. L. Stockl, W. A. Ribi, E. J. Warrant, Adaptations for Nocturnal and Diurnal Vision in the
Hawkmoth Lamina. *J. Comp. Neurol.* **524**, 160–175 (2016).

21.   Y. Wu, *et al.*, Retinal transcriptome sequencing sheds light on the adaptation to nocturnal and
diurnal lifestyles in raptors. *Sci. Rep.* **6**, 1–12 (2016).

22.   T. Akiyama, H. Uchiyama, S. Yajima, K. Arikawa, Y. Terai, Parallel evolution of opsin visual
pigments in hawkmoths by tuning of spectral sensitivities during transition from a nocturnal to a
diurnal ecology. *J. Exp. Biol.* **225**, jeb244541 (2022).

23. Y. Sondhi, E. Elis, S. Bybee, J. Theobald, A. Kawahara, Light environment drives evolution of color
vision genes in butterflies and moths (2021) https:/doi.org/10.5061/dryad.gmsbcc2kr (May 13,
2022).

24. M. Izutsu, A. Toyoda, A. Fujiyama, K. Agata, N. Fuse, Dynamics of Dark-Fly Genome Under
Environmental Selections. *G3* **6**, 365–376 (2015).

25. J. E. Yack, J. H. Fullard, Ultrasonic hearing in nocturnal butterflies. *Nature* **403**, 265–266 (2000).

26. F. Sandrelli, R. Costa, C. P. Kyriacou, E. Rosato, Comparative analysis of circadian clock genes in
insects. *Insect Mol. Biol.* **17**, 447–463 (2008).

27. R. Závodská, I. Sauman, F. Sehnal, Distribution of PER protein, pigment-dispersing hormone,
prothoracicotropic hormone, and eclosion hormone in the cephalic nervous system of insects. *J. Biol.*
Rhythms **18**, 106–122 (2003).

28. C. Hermann, *et al.*, The circadian clock network in the brain of different Drosophila species. *J.*
Comp. Neurol. **521**, 367–388 (2013).

29. C. A. Hamilton, *et al.*, Phylogenomics resolves major relationships and reveals significant
diversification rate shifts in the evolution of silk moths and relatives. *BMC Evol. Biol.* (2019)
https:/doi.org/10.1186/s12862-019-1505-1.

30. A. Kobelková, R. Závodská, I. Sauman, O. Bazalová, D. Dolezel, Expression of clock genes period
and timeless in the central nervous system of the Mediterranean flour moth, Ephestia kuehniella. *J.*
Biol. Rhythms **30**, 104–116 (2015).

31. D. Brady, A. Saviane, S. Cappellozza, F. Sandrelli, The Circadian Clock in Lepidoptera. *Front.*
Physiol. **12**, 776826 (2021).

32. S. Iwai, Y. Fukui, Y. Fujiwara, M. Takeda, Structure and expressions of two circadian clock genes,
period and timeless in the commercial silkmoth, Bombyx mori. *J. Insect Physiol.* **52**, 625–637
(2006).

33. A. Macias-Muñoz, A. G. Rangel Olguin, A. D. Briscoe, Evolution of Phototransduction Genes in
Lepidoptera. *Genome Biol. Evol.* **11**, 2107–2124 (2019).

34. X.-C. Jiang, *et al.*, Identification of Olfactory Genes From the Greater Wax Moth by Antennal
Transcriptome Analysis. *Front. Physiol.* **12**, 663040 (2021).

35. A. de Fouchier, *et al.*, Functional evolution of Lepidoptera olfactory receptors revealed by
deorphanization of a moth repertoire. *Nat. Commun.* **8**, 15709 (2017).

36. J. W. Truman, L. M. Riddiford, Neuroendocrine control of ecdysis in silkmoths. *Science* **167**, 1624–
1626 (1970).

37. I. Sauman, S. M. Reppert, Circadian clock neurons in the silkmoth Antheraea pernyi: novel
mechanisms of Period protein regulation. *Neuron* **17**, 889–900 (1996).

38.   P. M. Tuskes, J. P. Tuttle, M. M. Collins, *The Wild Silk Moths of North America: A Natural History*
of the Saturniidae of the United States and Canada (Cornell University Press, 1996).

39.   M. M. Collins, P. M. Tuskes, REPRODUCTIVE ISOLATION IN SYMPATRIC SPECIES OF
DAYFLYING MOTHS (HEMILEUCA: SATURNIIDAE). *Evolution* **33**, 728–733 (1979).

40.   F. Lu, *et al.*, SilkDB 3.0: visualizing and exploring multiple levels of data for silkworm. *Nucleic*
Acids Res. **48**, D749–D755 (2020).

41.   S.-R. Kim, *et al.*, Genome sequence of the Japanese oak silk moth, Antheraea yamamai: the first
draft genome in the family Saturniidae. *Gigascience* **7**, 1–11 (2018).

42.   J. Duan, *et al.*, A chromosome-scale genome assembly of Antheraea pernyi (Saturniidae,
Lepidoptera). *Mol. Ecol. Resour.* **20**, 1372–1383 (2020).

43.   H. Langer, G. Schmeinck, F. Anton-Erxleben, Identification and localization of visual pigments in
the retina of the moth, Antheraea polyphemus (Insecta, Saturniidae). *Cell Tissue Res.* **245**, 81–89
(1986).

44.   I. Kitching, *et al.*, A global checklist of the Bombycoidea (Insecta: Lepidoptera). *Biodiversity Data*
Journal **6**, e22236 (2018).

45.   M. J. Scoble, *The Lepidoptera. Form, function and diversity* (Oxford University Press, 1992).

46.   Lemaire, Minet, 18. The Bombycoidea and their Relatives. *Volume 1: Evolution, Systematics, and*
(2013).

47.   R. Rougerie, *et al.*, Phylogenomics Illuminates the Evolutionary History of Wild Silkmoths in Space
and Time (Lepidoptera: Saturniidae). *bioRxiv*, 2022.03.29.486224 (2022).

48.   Z. Wang, M. Gerstein, M. Snyder, RNA-Seq: a revolutionary tool for transcriptomics. *Nat. Rev.*
Genet. **10**, 57–63 (2009).

49.   D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157 (2015).

50.   D. Li, *et al.*, An evaluation of RNA-seq differential analysis methods. *PLoS One* **17**, e0264246
(2022).

51.   N. Sánchez-Baizán, L. Ribas, F. Piferrer, Improved biomarker discovery through a plot twist in
transcriptomic data analysis. *BMC Biol.* **20**, 208 (2022).

52.   P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network analysis. *BMC*
Bioinformatics **9**, 559 (2008).

53.   Lafayette Lafayette Press, *A Survey of Best Practices for RNA-Seq Data Analysis* (CreateSpace
Independent Publishing Platform, 2016).

54.   C. P. Cantalapiedra, A. Hernández-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas, eggNOG-mapper v2:

Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

55. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238 (2019).

56. P. E. Hardin, J. C. Hall, M. Rosbash, Behavioral and molecular analyses suggest that circadian output is disrupted by disconnected mutants in D. melanogaster. *EMBO J.* **11**, 1–6 (1992).

57. M. S. Dushay, M. Rosbash, J. C. Hall, The disconnected visual system mutations in Drosophila melanogaster drastically disrupt circadian rhythms. *J. Biol. Rhythms* **4**, 1–27 (1989).

58. K. J. Lee, M. Mukhopadhyay, P. Pelka, A. R. Campos, H. Steller, Autoregulation of the Drosophila disconnected gene in the developing visual system. *Dev. Biol.* **214**, 385–398 (1999).

59. V. Suri, Z. Qian, J. C. Hall, M. Rosbash, Evidence that the TIM light response is relevant to light-induced phase shifts in Drosophila melanogaster. *Neuron* **21**, 225–234 (1998).

60. B. K. Dey, X.-L. Zhao, E. Popo-Ola, A. R. Campos, Mutual regulation of the Drosophila disconnected (disco) and Distal-less (Dll) genes contributes to proximal-distal patterning of antenna and leg. *Cell Tissue Res.* **338**, 227–240 (2009).

61. H. Somanathan, A. Kelber, R. M. Borges, R. Wallén, E. J. Warrant, Visual ecology of Indian carpenter bees II: adaptations of eyes and ocelli to nocturnal and diurnal lifestyles. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **195**, 571–583 (2009).

62. A. L. Stöckl, D. O'Carroll, E. J. Warrant, Higher-order neural processing tunes motion neurons to visual ecology in three species of hawkmoths. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20170880 (2017).

63. I. Shimizu, Y. Yamakawa, Y. Shimazaki, T. Iwasa, Molecular cloning of Bombyx cerebral opsin (Boceropsin) and cellular localization of its expression in the silkworm brain. *Biochem. Biophys. Res. Commun.* **287**, 27–34 (2001).

64. K. Moses, M. C. Ellis, G. M. Rubin, The glass gene encodes a zinc-finger protein required by Drosophila photoreceptor cells. *Nature* **340**, 531–536 (1989).

65. R. Tearle, Tissue specific effects of ommochrome pathway mutations in Drosophila melanogaster. *Genet. Res.* **57**, 257–266 (1991).

66. H. K. Shamloula, *et al.*, rugose (rg), a Drosophila A kinase anchor protein, is required for retinal pattern formation and interacts genetically with multiple signaling pathways. *Genetics* **161**, 693–710 (2002).

67. A. Stöckl, *et al.*, Differential investment in visual and olfactory brain areas reflects behavioural choices in hawk moths. *Sci. Rep.* **6** (2016).

68. A. Sourakov, R. W. Chadd, *The Lives of Moths: A Natural History of Our Planet's Moth Life* (Princeton University Press, 2022).

69.   A. Y. Kawahara, *et al.*, Phylogenomics reveals the evolutionary timing and pattern of butterflies and
moths. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 22657–22663 (2019).

70.   J. J. Rubin, *et al.*, The evolution of anti-bat sensory illusions in moths. *Science Advances* (2018)
https:/doi.org/10.1126/sciadv.aar7428.

71.   J. E. Yack, L. D. Otero, J. W. Dawson, A. Surlykke, J. H. Fullard, Sound production and hearing in
the blue cracker butterfly Hamadryas feronia (Lepidoptera, nymphalidae) from Venezuela. *J. Exp.*
Biol. **203**, 3689–3702 (2000).

72.   J. R. Barber, *et al.*, Anti-bat ultrasound production in moths is globally and phylogenetically
widespread. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2117485119 (2022).

73.   G. A. Jacobs, J. P. Miller, Z. Aldworth, Computational mechanisms of mechanosensory processing
in the cricket. *J. Exp. Biol.* **211**, 1819–1828 (2008).

74.   J. M. Camhi, Locust wind receptors: I. Transducer mechanics and sensory response. *J. Exp. Biol.* **50**,
335–348 (1969).

75.   J. M. Camhi, Locust wind receptors. 3. Contribution to flight initiation and lift control. *J. Exp. Biol.*
50, 363–373 (1969).

76.   J. C. Tuthill, R. I. Wilson, Mechanosensation and Adaptive Motor Control in Insects. *Curr. Biol.* **26**,
R1022–R1038 (2016).

77.   J. D. Baker, S. Adhikarakunnathu, M. J. Kernan, Mechanosensory-defective, male-sterile unc
mutants identify a novel basal body protein required for ciliogenesis in Drosophila. *Development*
131, 3411–3422 (2004).

78.   D. D. Dell'Aglio, W. O. McMillan, S. H. Montgomery, Shifting balances in the weighting of sensory
modalities are predicted by divergence in brain morphology in incipient species of Heliconius
butterflies. *Anim. Behav.* **185**, 83–90 (2022).

79.   S. H. Montgomery, R. M. Merrill, S. R. Ott, Brain composition in Heliconius butterflies,
posteclosion growth and experience-dependent neuropil plasticity. *J. Comp. Neurol.* **524**, 1747–1769
(2016).

80.   S. Anton, W. Rössler, Plasticity and modulation of olfactory circuits in insects. *Cell Tissue Res.* **383**,
149–164 (2021).

81.   S. Halali, *et al.*, Predictability of temporal variation in climate and the evolution of seasonal
polyphenism in tropical butterfly communities. *J. Evol. Biol.* **34**, 1362–1375 (2021).

82.   W. Kuenzinger, *et al.*, Innate colour preferences of a hawkmoth depend on visual context. *Biol. Lett.*
15, 20180886 (2019).

83.   A. Kelber, Pattern discrimination in a hawkmoth: innate preferences, learning performance and
ecology. *Proc. Biol. Sci.* **269**, 2573–2577 (2002).

84.   G. T. Broadhead, T. Basu, M. von Arx, R. A. Raguso, Diel rhythms and sex differences in the
888   locomotor activity of hawkmoths. *J. Exp. Biol.* **220**, 1472–1480 (2017).

889   85.   M. Bischoff, R. A. Raguso, A. Jürgens, D. R. Campbell, Context-dependent reproductive isolation
890   mediated by floral scent and color. *Evolution* **69**, 1–13 (2015).

891   86.   S. Jaeger, C. Girvin, N. Demarest, E. LoPresti, Secondary pollinators contribute to reproductive
892   success of a pink-flowered sand verbena population. *Ecology* **104**, e3977 (2023).

893   87.   W. V. So, *et al.*, takeout, a novel Drosophila gene under circadian clock transcriptional regulation.
894   *Mol. Cell. Biol.* **20**, 6935–6944 (2000).

895   88.   T. Kikawada, *et al.*, Trehalose transporter 1, a facilitated and high-capacity trehalose transporter,
896   allows exogenous trehalose uptake into cells. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11585–11590
897   (2007).

898   89.   H. Steller, K. F. Fischbach, G. M. Rubin, Disconnected: a locus required for neuronal pathway
899   formation in the visual system of Drosophila. *Cell* **50**, 1139–1153 (1987).

900   90.   C. Helfrich-Förster, Robust circadian rhythmicity of Drosophila melanogaster requires the presence
901   of lateral neurons: a brain-behavioral study of disconnected mutants. *J. Comp. Physiol. A* **182**, 435–
902   453 (1998).

903   91.   L. R. Sanders, M. Patel, J. W. Mahaffey, The Drosophila gap gene giant has an anterior segment
904   identity function mediated through disconnected and teashirt. *Genetics* **179**, 441–453 (2008).

905   92.   E. Blanchardon, *et al.*, Defining the role of Drosophila lateral neurons in the control of circadian
906   rhythms in motor activity and eclosion by targeted genetic ablation and PERIOD protein
907   overexpression. *Eur. J. Neurosci.* **13**, 871–888 (2001).

908   93.   A. D. Keller, T. Maniatis, Only two of the five zinc fingers of the eukaryotic transcriptional
909   repressor PRDI-BF1 are required for sequence-specific DNA binding. *Mol. Cell. Biol.* **12**, 1940–
910   1949 (1992).

911   94.   S. Johnson, N. Boob, A strange pair. *News of the Lepidopterists' Society* **55**, 116 (2013).

912   95.   C. J. van der Kooi, D. G. Stavenga, K. Arikawa, G. Belušič, A. Kelber, Evolution of Insect Color
913   Vision: From Spectral Sensitivity to Visual Ecology. *Annu. Rev. Entomol.* **66**, 435–461 (2021).

914   96.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.
915   *EMBnet.journal* **17**, 10–12 (2011).

916   97.   M. G. Grabherr, *et al.*, Full-length transcriptome assembly from RNA-Seq data without a reference
917   genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

918   98.   D. R. Zerbino, E. Birney, Velvet: algorithms for de novo short read assembly using de Bruijn graphs.
919   *Genome Res.* **18**, 821–829 (2008).

920   99.   G. Robertson, *et al.*, De novo assembly and analysis of RNA-seq data. *Nat. Methods* **7**, 909–912

921   (2010).

100.   R. Luo, *et al.*, SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *Gigascience* **1**, 18 (2012).

101.   D. G. Gilbert, Longest protein, longest transcript or most expression, for accurate gene reconstruction of transcriptomes? *bioRxiv*, 829184 (2019).

102.   D. Gilbert, Gene-omes built from mRNA seq not genome DNA (2016) https:/doi.org/10.7490/f1000research.1112594.1 (June 6, 2022).

103.   , *https://github.com/NCGAS/de-novo-transcriptome-assembly-pipeline* (Github) (June 6, 2022).

104.   M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol. Biol. Evol.* **38**, 4647–4654 (2021).

105.   R. Smith-Unna, C. Boursnell, R. Patro, J. M. Hibberd, S. Kelly, TransRate: reference-free quality assessment of de novo transcriptome assemblies. *Genome Res.* **26**, 1134–1144 (2016).

106.   A. Mikheenko, A. Prjibelski, V. Saveliev, D. Antipov, A. Gurevich, Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics* **34**, i142–i150 (2018).

107.   W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

108.   L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

109.   M. Steinegger, J. Söding, MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

110.   Gene Ontology Consortium, The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res.* **49**, D325–D334 (2021).

111.   D. M. Bryant, *et al.*, A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of Limb Regeneration Factors. *Cell Rep.* **18**, 762–776 (2017).

112.   R. D. Finn, J. Clements, S. R. Eddy, HMMER web server: interactive sequence similarity searching. *Nucleic Acids Research* **39**, W29–W37 (2011).

113.   M. Punta, *et al.*, The Pfam protein families database. *Nucleic Acids Res.* **40**, D290–301 (2012).

114.   R. D. Finn, *et al.*, The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res.* **44**, D279–85 (2016).

115.   T. N. Petersen, S. Brunak, G. von Heijne, H. Nielsen, SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods* **8**, 785–786 (2011).

116. A. Krogh, B. Larsson, G. von Heijne, E. L. Sonnhammer, Predicting transmembrane protein
topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580
(2001).

117. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool.
J. Mol. Biol. **215**, 403–410 (1990).

118. M. Kanehisa, S. Goto, Y. Sato, M. Furumichi, M. Tanabe, KEGG for integration and
interpretation of large-scale molecular data sets. *Nucleic Acids Res.* **40**, D109–14 (2012).

119. M. Ashburner, *et al.*, Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nat. Genet.* **25**, 25–29 (2000).

120. S. Powell, *et al.*, eggNOG v3.0: orthologous groups covering 1133 organisms at 41 different
taxonomic ranges. *Nucleic Acids Res.* **40**, D284–9 (2012).

121. R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and bias-
aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

122. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

123. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

124. Alexa, Rahnenfuhrer, topGO: Enrichment analysis for Gene Ontology. R package version 2.28. 0.
Cranio (2016).

125. S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants.
Bioinformatics **36**, 2628–2629 (2020).

126. P. Langfelder, S. Horvath, Fast R Functions for Robust Correlations and Hierarchical Clustering.
J. Stat. Softw. **46** (2012).

127. Y. Mei, *et al.*, InsectBase 2.0: a comprehensive gene resource for insects. *Nucleic Acids Res.* **50**,
D1040–D1045 (2022).

128. J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).

129. V. Mariani, M. Biasini, A. Barbato, T. Schwede, lDDT: a local superposition-free score for
comparing protein structures and models using distance difference tests. *Bioinformatics* **29**, 2722–
2728 (2013).

130. B. Yariv, *et al.*, Using evolutionary data to make sense of macromolecules with a "face-lifted"
ConSurf. *Protein Sci.* **32**, e4582 (2023).

131. H. Ashkenazy, *et al.*, ConSurf 2016: an improved methodology to estimate and visualize
evolutionary conservation in macromolecules. *Nucleic Acids Res.* **44**, W344–50 (2016).

132.   G. Celniker, *et al.*, ConSurf: Using evolutionary data to raise testable hypotheses about protein
function. *Isr. J. Chem.* **53**, 199–206 (2013).

133.   H. Ashkenazy, E. Erez, E. Martz, T. Pupko, N. Ben-Tal, ConSurf 2010: calculating evolutionary
conservation in sequence and structure of proteins and nucleic acids. *Nucleic Acids Res.* **38**, W529–
33 (2010).

134.   M. Landau, *et al.*, ConSurf 2005: the projection of evolutionary conservation scores of residues
on protein structures. *Nucleic Acids Res.* **33**, W299–302 (2005).

135.   Schrödinger, LLC, The PyMOL Molecular Graphics System, Version 1.8 (2015).

136.   N. Blom, S. Gammeltoft, S. Brunak, Sequence and structure-based prediction of eukaryotic
protein phosphorylation sites. *J. Mol. Biol.* **294**, 1351–1362 (1999).

137.   N. Blom, T. Sicheritz-Pontén, R. Gupta, S. Gammeltoft, S. Brunak, Prediction of post-
translational glycosylation and phosphorylation of proteins from the amino acid sequence.
Proteomics **4**, 1633–1649 (2004).

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019