The Genetic Landscape of a Metabolic Interaction

Thuy N. Nguyen[1,2,3,†], Christine Ingle[1,2,3], Samuel Thompson[4, ‡], Kimberly A. Reynolds[1,2,3*]

[1]The Green Center for Systems Biology, [2]Department of Bioinformatics, and [3]Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, USA, 75390

[4]Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, CA 94158

[†]Present address: Form Bio, Dallas, USA, 75226

[‡]Present address: Department of Bioengineering, Stanford University, Stanford, CA 94305

* Correspondence: kimberly.reynolds@utsouthwestern.edu (K.A. Reynolds)

Keywords: dihydrofolate reductase, DHFR, thymidylate synthase, TYMS, deep mutational scan, epistasis, genotype-to-phenotype, genetic interaction

Abstract: Enzyme abundance, catalytic activity, and ultimately sequence are all shaped by the need of growing cells to maintain metabolic flux while minimizing accumulation of deleterious intermediates. While much prior work has explored the constraints on protein sequence and evolution induced by physical protein-protein interactions, the sequence-level constraints emerging from non-binding functional interactions in metabolism remain unclear. To quantify how variation in the activity of one enzyme constrains the biochemical parameters and sequence of another, we focused on dihydrofolate reductase (DHFR) and thymidylate synthase (TYMS), a pair of enzymes catalyzing consecutive reactions in folate metabolism. We used deep mutational scanning to quantify the growth rate effect of 2,696 DHFR single mutations in 3 TYMS backgrounds under conditions selected to emphasize biochemical epistasis. Our data are well-described by a relatively simple enzyme velocity to growth rate model that quantifies how metabolic context tunes enzyme mutational tolerance. Together our results reveal the structural distribution of epistasis in a metabolic enzyme and establish a foundation for the design of multi-enzyme systems.

INTRODUCTION

Enzymes function within biochemical pathways; exchanging substrates and products to generate useful metabolites. This metabolic context places constraints on enzyme velocity — the product of both catalytic activity and enzyme abundance. For example, the relative velocities of some enzymes must be coordinated to avoid accumulation of deleterious metabolic intermediates[1–3]. In other instances, optimal enzyme abundance is set by a tradeoff between the cost of protein synthesis and the benefit of efficient nutrient utilization[4–6]. Considered at the pathway scale, metabolic enzymes are often produced in evolutionarily conserved stoichiometric ratios across species[7], providing further indication that relative — not just absolute — enzyme velocity is under selection. More generally, the relationship between the velocity of a given enzyme, metabolic flux through a pathway, and cellular growth rate is non-linear and shaped by interactions amongst pathway enzymes (**Fig.1a**). Indeed, a key result of metabolic control theory is that the control coefficient of an enzyme — defined as the fractional change in pathway-level flux given a fractional change in enzyme velocity — depends on the starting (native) velocity of the enzyme, but *also* the velocity of all other enzymes in the pathway[8,9]. That is to say, given that enzymes act sequentially to produce metabolites, the effects of mutations on cellular phenotype can be buffered or amplified depending on which enzymatic reactions control metabolic flux. As a consequence, enzyme mutations that are neutral in one context may have profound consequences for metabolic flux and growth rate in the background of variation in another[10–13]. This context-dependence, or epistasis, amongst metabolic enzymes need not be mediated by direct physical binding, but emerges indirectly through shared metabolite pools and a need to maintain flux while avoiding the accumulation of deleterious intermediates[6,11,14].

While much prior work has explored the constraints on protein sequence and evolution induced by physical protein-protein interactions, the sequence-level constraints emerging from these sorts of non-binding functional interactions in metabolism remain unclear. How is this biochemically-mediated epistasis organized in the protein structure and reflected in the

sequence? A quantitative understanding of how pathway context shapes sequence and activity would assist in the interpretation of disease-associated mutations, the design of new enzymes, and directing the laboratory evolution of metabolic pathways. To begin to address this, we examined the residue-level epistatic interactions between a pair of enzymes that catalyze consecutive reactions in folate metabolism: dihdyrofolate reductase (DHFR) and thymidylate synthase (TYMS). The activity of these enzymes is strongly linked to *E. coli* growth rate, they are frequent targets of antibiotics and chemotherapeutics, and our prior work showed that they co-evolve as a module both in the laboratory and across thousands of bacterial genomes[1]. Taking this enzyme pair as a simplified model system in which to examine a biochemically-mediated epistatic interaction, we created a mathematical model relating variation in DHFR and TYMS catalytic parameters to growth rate using a focused set of well-characterized point mutants. Then, to more deeply test this model and comprehensively map the pattern of epistasis between these two enzymes, we measured the effect of nearly all possible DHFR single mutations (2,696 in total) in the context of three TYMS variants selected to span a range of catalytic activities. The model predicted – and the data showed – that TYMS background profoundly changed both the sign (buffering vs. aggravating) and magnitude of DHFR epistasis. Mapping the epistatic effects of mutation to the DHFR tertiary structure revealed that they are organized into distinct clusters based on epistatic sign. Additionally, mutations with the largest magnitude epistatic effect to TYMS centered around the DHFR active site, while more weakly epistatic positions radiated outwards. Finally, we inferred approximate values for DHFR catalytic power (k_{cat}/K_m) across all 2,696 mutations by using the growth rate measurements across TYMS backgrounds to constrain the enzyme velocity to growth rate model. The residues linked to catalysis form a structurally distributed network inside the enzyme and are highly evolutionarily conserved. Taken together, our data demonstrates at single-residue resolution how epistasis mediated through a biochemical interaction reshapes a mutational landscape. Our results indicate that metabolic context can strongly modulate enzyme evolution in both the clinic and the lab by facilitating or frustrating

available mutational paths. More generally, our results invite one to consider new ideas for the

joint design of multi-enzyme systems that take into account shared constraints on relative activity

and sequence.

RESULTS

An enzyme velocity to growth rate model for DHFR and TYMS

We constructed a mathematical model relating changes in DHFR and TYMS catalytic parameters

to growth rate phenotype. Our goals were to (1) formalize our previous empirical findings

describing DHFR/TYMS biochemical coupling[1], (2) quantify the absolute and relative constraints

on DHFR and TYMS catalytic activity, and (3) define the relationship between biochemical activity

and epistasis. DHFR and TYMS play a central role in folate metabolism, a well-conserved

biochemical pathway involved in the synthesis of purine nucleotides, thymidine, glycine, and

methionine[15] (**Fig. 1b**). Consequently, this pathway is strongly linked to cell growth and a frequent

target of antibiotics and chemotherapeutics. DHFR is a 159 amino acid enzyme that catalyzes

the reduction of dihydrofolate (DHF) to tetrahydrofolate (THF) using NADPH as a cofactor. The

reduced THF then serves as a carrier for activated one-carbon units in downstream metabolic

processes. TYMS catalyzes the oxidation of THF back to DHF during deoxythymidine synthesis

and is the sole enzyme responsible for recycling the DHF pool[16,17]. Prior work from ourselves and

others indicates that these two enzymes are strongly functionally coupled to each other and less

coupled to the remainder of the pathway: they co-evolve in terms of synteny and gene presence-

absence across bacterial species[1], inhibition of DHFR with trimethoprim is rescued by suppressor

mutations in TYMS in both the lab and the clinic[1,18], and loss-of-function mutations in DHFR are

rescued by loss-of-function mutations in TYMS[1,19,20]. Metabolomics data indicated that loss of

DHFR function resulted in accumulation of DHF and depletion of reduced folates, and that

compensatory loss of function mutations in TYMS help to restore DHF and THF pools to more

native-like levels[1,21,22]. Thus, DHFR and TYMS are a growth-linked two-enzyme system where

epistasis is driven by a biochemical interaction, with the added simplification that they are

relatively decoupled from surrounding metabolic context.

We first developed our mathematical model using a previously collected set of

metabolomics and growth rate data for five DHFR point mutants in the background of both WT

TYMS and TYMS R166Q (**Table S1, Table S2**). TYMS R166Q is an active site mutation with near

complete loss of catalytic activity[23]. The model comprises two parts: the relationship between

intracellular THF abundance and growth rate, and the relationship between enzymatic activity and

intracellular THF. First, we considered the relationship between intracellular THF abundance and

growth rate as measured across all ten DHFR/TYMS sequence combinations. THF limitation

restricts the production of several growth-linked factors, including thymidine, methionine, glycine,

and the purine precursors inosine and AICAR. Under the experimental conditions of our growth

rate assays — M9 minimal media with 0.4% glucose, 0.2% amicase, and 50 µg/ml thymidine —

thymidine is not growth limiting (TYMS R166Q is rescued to WT-like growth) and amicase

provides a source of free amino acids. These conditions — which remove selection pressure on

TYMS due to thymidine production — emphasize coupling between DHFR and TYMS through

the shared THF pool which must be used to produce purine nucleotides. We previously observed

a hyperbolic dependence of growth rate on reduced folate abundance for many THF species in

these experimental conditions[1]. We selected 10-formyl THF with three glutamates as a

representative growth-linked reduced folate for modeling given it's clear relationship to growth

and proximity to purine biosynthesis. Following a similar approach as Rodrigues et al, we fit a

single four-parameter sigmoidal function relating growth rate to the experimental measurements

of intracellular THF concentration[24] (**Fig. 1c**).

139   Eqn. 1 $g = \dfrac{g_{max} - g_{min}}{1 + (K/[THF])^n} + g_{min}$

Here, g_{max} represents the maximal growth rate, g_{min} is the minimal growth rate, K is a constant

that captures the concentration of THF that yields 50% growth, and n is a Hill coefficient (**Table**

S3).

Next, we sought to connect variation in DHFR and TYMS enzyme velocity to intracellular

THF concentrations. To simplify our model, we reduced the pathway to a cycle in which DHFR

and TYMS catalyze opposing oxidation and reduction reactions (**Fig. 1d**). This abstraction

assumes that DHFR and TYMS dominate turnover of the DHF and THF pools, and that the

reduced folates are considered as a single THF pool. While this simplification clearly omits much

of folate metabolism, it allows us to write a rate equation that isolates the recycling of THF in

terms of a small number of measurable biochemical parameters:

$$\text{Eqn. 2} \quad \frac{d[\text{THF}]}{dt} = \frac{[\text{DHFR}] * k_{cat}^{\text{DH}}}{1 + \text{K}_m^{\text{DH}}/([\text{fol}_{\text{tot}}] - [\text{THF}])} - \frac{[\text{TYMS}] * k_{cat}^{\text{TS}}}{1 + \text{K}_m^{\text{TS}}/([\text{THF}])}$$

152   In this equation, DHFR and TYMS are treated as catalyzing opposing reactions with Michaelis

153   Menten kinetics, providing a relationship between steady state kinetics parameters (k_{cat}^{DH}, K_m^{DH},

154   k_{cat}^{TS}, K_m^{TS}) and intracellular THF abundance. From this equation one can find an analytical solution

155   for the steady state concentration of THF in the form of the Goldbeter-Koshland equation [25,26].

$$\text{Eqn. 3} \quad \frac{[\text{THF}_{ss}]}{[\text{fol}_{\text{tot}}]} = \frac{\frac{V_1}{V_2} * (1 - \hat{K}_{m2}) - \hat{K}_{m1} - 1 + \sqrt{4\hat{K}_{m2}\frac{V_1}{V_2}(\frac{V_1}{V_2} - 1) + (\frac{V_1}{V_2}(\hat{K}_{m2} - 1) + \hat{K}_{m1} + 1)^2}}{2 * (\frac{V_1}{V_2} - 1)}$$

158   Where:

$$V_1 = [\text{DHFR}]k_{cat}^{\text{DH}} \qquad \hat{K}_{m1} = \text{K}_m^{\text{DH}}/\text{fol}_{\text{tot}}$$

$$V_2 = [\text{TYMS}]k_{cat}^{\text{TS}} \qquad \hat{K}_{m2} = \text{K}_m^{\text{TS}}/\text{fol}_{\text{tot}}$$

In our initial model construction, the steady state catalytic parameters (k_{cat}^{DH}, K_m^{DH}, k_{cat}^{TS}, K_m^{TS}) were

experimentally measured *in vitro* using purified samples of all mutants, with the exception of

TYMS R166Q which is near-inactive and assigned an arbitrarily low k_{cat} and high K_m (**Table**

S1,S2). Four fit parameters remain in Equation 3: (1) the concentration of the total folate pool

($[\text{fol}_{tot}]$) (2) the intracellular concentration of DHFR ($[\text{DHFR}]$), which we treated as identical across

all variants (as our model will eventually describe thousands of DHFR mutations, and we wished

to avoid overparameterization), (3) the intracellular concentration of WT TYMS ($[\text{TYMS}_{WT}]$), and

(4) the intracellular concentration of TYMS R166Q ($[\text{TYMS}_{R166Q}]$). This relatively simplified model

showed good correspondence to the data when fit (R^2 = 0.96, **Fig. 1e, Table S3**). Equations 1

and 3 were then combined to estimate growth rate as a function of both DHFR and TYMS activity,

by linking catalytic activity to THF abundance, and then THF abundance to growth rate. The

complete model worked well to predict growth rate on our initial training set (**Fig. 1f**).

The growth rate effect of DHFR mutations changes magnitude and sign depending upon

TYMS background.

To more rigorously test our model and understand its' predictions, we expanded our dataset to

include more DHFR and TYMS variants with experimentally characterized activities. As our initial

model was developed using only two extreme TYMS variants (wild-type and a near complete loss

of function variant, R166Q), we were particularly curious to evaluate model performance for TYMS

mutations with intermediate effects on catalysis and *E. coli* growth. We identified candidate TYMS

mutations by examining an earlier growth complementation study[27]. A handful of these mutants

were then cloned, screened for expression, and when possible, purified and characterized.

Through this mini-screen we selected two mutations that stably expressed, purified robustly, and

yielded intermediate activities: TYMS R127A and Q33S (**Fig. 2a**). The R127A mutation is located

in the TYMS active site and is one of four arginines that coordinate the substrate (dUMP)

phosphate group. The Q33S mutation is located at the TYMS dimer interface, distal to the active

site. We observed that R127A was more deleterious to catalytic function than Q33S, but that both

mutations were more active than R166Q (which shows almost no measurable activity *in vitro*, **Fig.**

2b, Table S2).

We measured growth rates for seven catalytically characterized DHFR variants (a set of

single and double mutants selected to span a range of catalytic activities) in the background of

these four TYMS mutants (WT, R127A, Q33S and R166Q). The point mutants were created in a

plasmid encoding both DHFR and TYMS (see Methods for details), and the plasmids were

transformed into an E. coli selection strain lacking the endogenous DHFR and TYMS genes

(ER2566 *ΔfolA ΔthyA*). Growth rates were measured in triplicate using a plate-reader-based

assay (28 measurements total; **Fig. 2c**, **Fig S1a,c**). We used this focused dataset to re-

parameterize the model equations, this time fitting five total parameters

($[\text{fol}_{\text{tot}}], [\text{DHFR}], [\text{TYMS}_{\text{WT}}], [\text{TYMS}_{\text{Q33S}}], [\text{TYMS}_{\text{R127A}}], [\text{TYMS}_{\text{R166Q}}]$, **Table S3**). This second

round of fitting tested the ability of growth rate data alone to constrain the model — an important

step because metabolomics data are available for only a limited number of DHFR and TYMS

mutants and are inherently far lower throughput to collect than growth rates. This iteration of

parameterization also tested the capacity of the model to capture TYMS mutations with

intermediate effects on activity. The data were again well described by the model (**Fig. 2C, Fig.**

S1b,d). We observed some variation in the fit parameters (relative to the older data in Fig.1); this

difference might be attributed to the fact that our newer experiments used a revised selection

vector backbone. As a control for overfitting, we tested the ability of the model to predict growth

rates for arbitrary catalytic data. We randomly shuffled the catalytic parameters (k_{cat} and K_m)

among mutations for both DHFR and TYMS, refit all free model parameters, and calculated the

RMSD and R^2 values between the best fit model and the shuffled data. Importantly, the model

was generally unable to describe the experimental growth rate data when catalytic parameters

were shuffled across both DHFR and TYMS (**Fig S1e,f**). This indicated that the model provided

a specific description of our experiment and was not trivially overfit. The model was less sensitive

to shuffling TYMS catalytic parameters (presumably because we included fit parameters

describing the abundance of each TYMS mutation that can compensate for this shuffling, **Fig.**

S1h). However, it was strongly sensitive to shuffling DHFR parameters (**Fig. S1g**). Taken

together, this analysis indicated that the model provides a good description of the enzyme-

velocity-to-growth-rate relationship and can be used to predict and interpret how molecular

changes in DHFR and TYMS activity modulate growth rate phenotype.

To examine the model more closely, we considered the relationship between TYMS

catalytic activity and DHFR mutational sensitivity. As in previous work, we observed that loss-of-

function mutations in DHFR can be partly or even entirely rescued by the loss-of-function mutation

TYMS R166Q in the presence of thymidine (**Fig. S1a,c**). TYMS R127A, a less severe loss of

function mutation, showed a similar albeit more modest trend – this mutation was able to partly

rescue growth for some (though not all) DHFR mutations. A central factor behind DHFR and

TYMS biochemical coupling is that loss-of-function mutations in TYMS help to preserve reduced

folate pools, allowing THF to shuttle one-carbon units in downstream biochemical processes like

purine biosynthesis even when DHFR activity is low. Moreover, loss of TYMS activity reduces

accumulation of DHF, which can inhibit upstream reactions[1,22]. Thus, the TYMS R166Q and

R127A variants show positive (buffering) epistasis to low-activity DHFR mutations. In contrast to

our expectation that a more intermediate mutation would also demonstrate intermediate levels of

buffering epistasis, TYMS Q33S shows negative (or amplifying) epistasis to some DHFR

mutations (**Fig. S1c**). This means that these DHFR mutations are more deleterious in the

background of TYMS Q33S than in the native TYMS context. Our model accounted for this

observation by increasing the intracellular concentration of TYMS Q33S by three fold (a fit

parameter, **Table S3**) such that the V_{max} of TYMS Q33S becomes greater than wildtype

$([\text{TYMS}_{\text{Q33S}}]k_{cat}^{\text{TYMS\_Q33S}} > [\text{TYMS}_{\text{WT}}]k_{cat}^{\text{TYMS\_WT}})$. This in turn increased the intracellular

requirement for DHFR activity, resulting in negative epistasis.

To further explore the pattern of epistasis across TYMS backgrounds, we simulated

growth rates over a range of DHFR k_{cat} and K_m values in each TYMS background (**Fig. 2d**). This

provided a comprehensive prediction of the TYMS-induced constraints on DHFR activity. In

particular, we obtained a regime of DHFR k_{cat} and K_m values that is sufficient to support growth

for each TYMS mutation. From these data we computed epistasis. These results indicated that

TYMS Q33S has negative epistasis to DHFR variants spanning a well-defined band of catalytic

parameters. The simulations also indicated that R127A has weak positive epistasis over a regime

of moderately impaired DHFR variants, but is insufficient to rescue growth for the strongest loss

of function variants. Finally, TYMS R166Q was observed to be broadly rescuing; DHFR variants

need only a negligible amount of activity to support growth in this context. Together, our

simulations showed that the sign and magnitude of DHFR epistasis are strongly tuned by TYMS

background, and provided quantitative predictions of the catalytic regimes where epistasis is most

apparent.

The single-mutant landscape of DHFR is strongly modulated by TYMS context

Next we examined the structural pattern of biochemical epistasis at the residue level across

DHFR. This also provided an opportunity to see if the model predictions — negative epistasis for

Q33S and broadly positive epistasis for R166Q — held true across a larger dataset. To

accomplish this, we created a plasmid-based saturation mutagenesis library of DHFR containing

all possible single mutations at every position (3002 total). This library was subcloned into all three

TYMS backgrounds. Sequencing showed that these libraries are well-distributed and approach

full coverage of all single mutations (97.1% - WT TYMS, 94.6% - TYMS Q33S, 99.3% - TYMS

R166Q) (**Fig. S2**). We transformed these libraries into the *E. coli* selection strain (ER2566 *ΔfolA*

ΔthyA). Transformants for each library were then grown as a mixed population in selective media

in a turbidostat to ensure maintenance of exponential growth and constancy of media conditions.

By quantifying the change in the relative frequency of individual mutants over time with next

generation sequencing, we obtained a growth rate difference relative to WT DHFR for nearly all

mutations in the library (**Fig. 3a**, **Table S4**, see methods for details). All relative growth rate

measurements were made in triplicate, with good concordance among replicates (**Fig. S3**).

The entire dataset showed that the DHFR mutational landscape was strongly dependent

on TYMS background (**Fig. 3b-e**). In all three TYMS backgrounds, the distribution of growth rate

effects was bi-modal and reasonably well-described by a double gaussian containing one peak

of near-neutral mutations and another (far smaller) peak of mutations with highly deleterious

growth rate effects. This is the expected result for an enzyme that shows a sigmoidal relationship

between activity and growth. In the native TYMS context, the vast majority of mutations fall into

the near-neutral peak. However, there is a substantial fraction (12%, 343 total) that display growth

rates at or below that of "inactive", where "inactive" was defined as the average growth rate across

nonsense mutations in the first 120 residues of DHFR. Consistent with expectation, mutations at

known positions of functional importance tended to be deleterious in the WT TYMS context (W22,

D27, F31, T35, L54, R57, T113, G121, and D122)[28]. For example, both W22 and D27 are directly

in the active site and serve to coordinate substrate through a hydrogen bonding network[29], G121

and D122 are part of the βf-βg loop and stabilize conformational changes associated to

catalysis[30,31], and F31 contacts the substrate and is associated to the "network of promoting

motions"[32,33]. In the TYMS Q33S context, many of these deleterious mutations had even more

severe effects or were classified as "Null". Null mutations disappeared from our sequencing

counts within the first three time points (8 hours) of the selection experiment, preventing accurate

inference of growth rate. For example, mutations at position 22 are deleterious in the WT TYMS

context, and appear as Null or very deleterious in the Q33S context. The same pattern can be

readily observed for positions 7,14,15, 22, 27, 31, 35, and 121. Again we saw that 12% of

mutations have growth rates at or below that of "inactive" variants. Finally, in the TYMS R166Q

context, there are very few deleterious mutations. Nearly all mutations are contained in the near-

neutral peak, including mutations at highly conserved active site positions like M20, W22, and

L28. Stop codons and mutations at the active site residue D27 continued to be deleterious,

indicating that DHFR activity was still under (very weak) selection in the TYMS R166Q

background. Nonetheless, only 5% of mutations displayed growth rates at or below those of

inactive mutations. Thus TYMS R166Q is broadly buffering to DHFR variation.

To quantify the context dependence of mutational effects, we computed epistasis for all

DHFR mutations with measurable relative growth rates in each of the three TYMS backgrounds

(2,696 in total, see also methods) (**Fig. 4, Fig. S4, Table S5**). We assessed the statistical

significance of epistasis by unequal variance t-test under the null hypothesis that the mutations

have equal mean growth rates in both TYMS backgrounds. These p-values were compared to a

multiple-hypothesis testing adjusted p-value determined by Sequential Goodness of Fit (P = 0.035

for TYMS Q33S and P = 0.029 for TYMS R166Q, **Fig. 4a,b**) [34]. In the TYMS Q33S background,

95 mutations (3%) showed significant negative epistasis and 280 mutations (9%) showed

significant positive epistasis. Many of the DHFR mutations with positive epistasis to Q33S were

near-neutral in the WT context, and displayed small improvements in growth rate that were highly

significant due to the low experimental error for the best-growing mutations (**Fig. 4c**). In contrast,

the mutations with negative epistasis exhibited a range of growth rate effects in the WT context.

For the TYMS R166Q background the overall proportion of significant epistatic mutations was

larger: while only 41 mutations (1%) showed significant negative epistasis, 851 mutations (28%)

showed significant positive epistasis. A smaller number of deleterious mutations in the WT context

were not rescued by R166Q (grey points with relative growth rates near 0.25). One possibility is

that these mutations are deleterious for reasons beyond disrupting metabolic flux; for example

they may result in protein aggregation or off target physical interactions. Regardless, direct

comparison of the relative growth rates of mutations across the WT, Q33S, and R166Q TYMS

backgrounds makes it very obvious that TYMS R166Q was broadly rescuing, while TYMS Q33S

had a more subtle effect that sometimes yielded negative epistasis (**Fig. 4c,d**).

The enzyme velocity to growth-rate model captures the observed fitness landscapes and

allows global estimation of mutational effects on catalysis

Next we sought to further test our enzyme velocity to growth-rate model using the deep mutational

scanning data. We refit the model a third time, in order to include all available experimental

information in parameterizing the model for this larger set of mutants. This included a larger

dataset of 34 DHFR single mutants with previously reported k_{cat} and K_m values. We additionally

characterized k_{cat} and K_m for four new DHFR mutations (I5K, V13H, E17V and M20Q) that

exhibited strong sign epistasis to TYMS to more completely test our ability to predict epistasis.

Together this yielded a set of 114 growth rate measurements with matched k_{cat} and K_m values for

DHFR and TYMS (38 DHFR mutations in 3 TYMS backgrounds, **Table S1**). We used these data

to perform a bootstrap analysis; iteratively subsampling the data and refitting the model 1000

times to obtain standard deviations in our model fit and the eight associated parameters (**Fig. 5a**).

The inferred parameters for this large set of sequencing-based growth rate measurements were

qualitatively similar to those obtained for the smaller set of 28 plate-reader based growth rate

measurements (7 DHFR mutants in 4 TYMS backgrounds, **Fig. 2**), but we observed some

discrepancy in the estimated total folate pool and intracellular concentrations of TYMS (**Table**

S3). In general, the total folate pool and concentrations of TYMS R166Q were more variable

across the bootstrap analysis, indicating that these two parameters are less well constrained by

our data (as indicated by the estimated variances in **Table S3**). Nevertheless, the inferred

parameters in both the plate reader and sequencing-based experiments suggested similar relative

expression levels of DHFR and the three TYMS point mutants, with Q33S being roughly 2-3 times

more abundant than WT DHFR. Overall both the predicted growth rates and pattern of epistasis

showed good agreement to our experimental observations (**Fig. 5a,b**).

Having established model performance on this subset of 114 biochemically characterized

DHFR and TYMS sequences, we next examined consistency of the model with all growth rate

measurements (the total model fit). However the effect of most mutations on catalysis is unknown.

Thus, for each DHFR point mutant we used Monte Carlo sampling to identify a space of k_{cat} and

K_m values consistent with the three growth rate measurements (in the three TYMS backgrounds).

While three growth rate measurements were insufficient to uniquely constrain both k_{cat} and K_m

(the solution space is degenerate), this process did permit estimation of $\log_{10}$ catalytic power

(k_{cat}/K_m) for all 2,696 characterized point mutants. For the subset of biochemically characterized

DHFR mutants we observed reasonable agreement between the *in vitro* measurements and those

inferred from our experimental growth rate data (R^2 = 0.74, **Fig. 5c**). Once catalytic parameters

were estimated across all point mutants, we put them back into the model to assess the

correspondence between the predicted (modeled) growth rates, predicted epistasis, and our

experimental observations, yielding a global picture of model fit quality. Overall, we observed that

the model well-described the data with two exceptions. First, there was a small proportion of

DHFR mutations that were predicted to be rescued by TYMS R166Q but in actuality were not (70

total, 2% of all DHFR mutations, the horizontal stripe of red dots in **Fig. 5d**). It is possible that

these mutations caused a growth rate defect through DHFR mis-folding and aggregation, a factor

not captured by our model. Second, there was a proportion of DHFR mutations predicted to have

negative epistasis to TYMS R166Q but observed to exhibit mild positive epistasis (**Fig. 5e**). These

differences may be related to the fact that DHFR abundance is modeled with a single parameter

across all mutants. These discrepancies could be addressed in future work by including additional

high-throughput assay data on stability or abundance to tease apart mutational effects on

catalysis from stability[35]. Nevertheless, the data indicated that our model can globally describe

growth rate phenotypes given variation in enzyme velocity.

The resulting model and inferred catalytic parameters now permit estimation of DHFR

single mutant fitness in any TYMS background. We computed the fraction of DHFR point mutants

that are neutral (growth rate above 0.9) as a function of variation in TYMS k_{cat} and K_m. These

calculations highlighted that selection on DHFR activity is strongly shaped by TYMS background,

with low-activity TYMS variants increasing the mutational tolerance of DHFR (**Fig. 5f,g**). This

suggests that TYMS inhibition or loss of function could promote the evolvability of DHFR both in

the clinic and laboratory settings.

Epistasis between DHFR and TYMS is organized into structurally localized groups

Next, we examined the structural pattern of DHFR positions with epistasis to TYMS Q33S and

TYMS R166Q. Given that mutations tend to have similar epistatic effects at a particular DHFR

position in our data set (**Fig. S4**), we used k-means clustering to sort positions into four categories

according to their pattern of epistasis: negative, insignificant, positive, and strong positive (**Fig.**

6a, Table S6). The strong positive category solely contained DHFR mutations in the TYMS

R166Q background, while the negative epistasis category was predominantly occupied by DHFR

mutations in the TYMS Q33S background. Mapping these positions to the DHFR structure

showed that epistasis is organized into spatially distinct regions of the tertiary structure (**Fig.**

6b,c). Mutations with negative epistasis to Q33S tended to be proximal to the DHFR active site,

particularly the folate binding pocket. The negative epistasis cluster included several key positions

near or in the Met-20 loop, which is known to undergo conformational fluctuations associated with

catalysis(residues A9, V13, E17 and M20) [28,31]. It also encompassed positions I5, L24, L28, and

F31 which surround the folate substrate. Several of these positions have known roles in catalysis;

mutations at position 31 promoted product release (while slowing hydride transfer), and dynamics

of the M20 loop (which includes V13,E17) are associated with substrate binding and product

release[32,36]. Additionally, specific mutations at positions 5, 20, and 28 result in trimethoprim

resistance by altering trimethoprim affinity[36]. These structural and biochemical observations are

consistent with the finding that mutations with negative epistasis tended to yield moderate to

severe growth rate defects. In contrast, positions with positive epistasis to Q33S often had very

little (or sometimes a beneficial) effect on growth rate, and were distributed around the DHFR

surface (**Fig. 4c, Fig. 6b**). In the context of TYMS R166Q only one position — C85 — was

included in the negative epistasis cluster (**Fig. 6c**). A large fraction of DHFR positions (53%, 84

total) displayed positive epistasis to TYMS R166Q; these positions were distributed throughout

the DHFR structure. The positions in the strong positive epistasis cluster included mutations with

some of the most severe effects on growth rate in the WT TYMS context. A number of these

positions were previously established as important to DHFR catalysis, including residues F31,

L54, G121, D122, and S148[28]. Mutations at these sites can be detrimental to k_{cat}, K_m, or both.

Epistasis and the structural encoding of DHFR catalysis

When the epistatic clusters are viewed together on the structure, one sees that they form

approximate distance-dependent shells around the active site (**Fig. 7a-d**). Considering the pattern

of epistasis to TYMS Q33S, positions with negative epistasis were closest to the active site,

surrounded by positions with insignificant epistasis, and finally positions in the positive epistasis

cluster form an outer shell (**Fig. 7a,b**). For TYMS R166Q, positions in the strong positive epistasis

cluster were closest to the active site, followed by positive epistasis positions, and finally those

with insignificant epistasis (**Fig. 7c,d**). For comparison, we also mapped the model-predicted

catalytic power averaged across all mutations at a position to the structure (**Fig. 7e**). Together,

these structural images paint a picture of the molecular encoding of catalysis and epistasis.

Mutations with predicted intermediate-to-large effects on catalysis were nestled near the active

site and showed negative epistasis to Q33S and strong positive to positive epistasis to R166Q.

Mutations with more mild effects on catalysis showed weaker positive to insignificant epistasis to

R166Q and Q33S. Though catalysis and epistasis showed an approximate distance-dependent

relationship to the DHFR active site, there a number of key positions distal to the active site that

exhibited large growth rate effects, strong positive epistasis to TYMS R166Q, and likely act

allosterically to tune catalytic activity (e.g. L110, G121, D122, W133, S148, and Y151). The

positions with the largest estimated effects on catalysis were highly evolutionarily conserved (P <

10^{-10} by Fisher's exact test, **Table S7**, **Fig. 7f**), indicating that our model and experimental data

are capturing features relevant to the fitness of DHFR. Together, these data show that TYMS

metabolic context strongly modulates the constraints on DHFR activity and catalysis.

DISCUSSION

It is well-appreciated that physical protein interactions place constraints on the individual

interacting monomers. Protein interfaces are organized to bind with appropriate affinity and avoid

non-specific interactions[37,38]. The individual components of physical complexes tend to be

expressed in similar ratios to avoid dosage related toxicity and aggregation[39,40]. However the

extent to which interactions mediated by biochemistry (rather than binding) constrain the function

and sequence of individual monomers has remained less clear. We have explicitly revealed these

interactions at single-residue resolution for one model system and coupled them with a

mathematical model to quantify the intracellular constraints on DHFR and TYMS relative catalytic

activities.

Our mutagenesis data and modeling show that TYMS activity strongly modifies the

constraints on DHFR catalytic parameters; shaping both the range and relative importance of k_{cat}

and K_m in modulating growth. This biochemical interaction results in an approximately shell-like

pattern of mutational sensitivity to TYMS background (epistasis) in the DHFR tertiary structure.

Extreme loss-of-TYMS function buffered variation in some of the most conserved DHFR active

site positions, while moderate loss of function buffered variation at more peripheral surface

exposed sites. Given these data, we expect that inhibition or loss-of-function in TYMS will promote

the evolvability of DHFR in nutrient rich environments, a finding with consequences for both

laboratory and clinical evolution. For example, inhibiting TYMS activity in the clinic may promote

the evolution of drug resistance in DHFR, while activating TYMS may restrict evolutionary

accessible paths. In the laboratory, strains with reduced TYMS activity could provide a less

stringent context for testing designed sequences or evolving new DHFR function.

The existence of an enzyme velocity to growth-rate mapping — by definition — allows us

to relate variation in DHFR and TYMS catalytic parameters to growth rate. It also allows one (in

principle) to do the inverse: infer *in vitro* catalytic parameters from growth rate measurements.

The intuition follows from classic steady-state Michaelis Menten experiments: to quantify steady

state kinetics *in vitro* one measures enzyme initial velocity as a function of substrate

concentration. In our sequencing-based experiments, variation in TYMS background effectively

titrates intracellular concentrations of DHF (substrate) while growth rate provides an estimate of

velocity. Though our current dataset of three TYMS backgrounds is insufficient to uniquely

constrain precise fits for k_{cat} and K_m, we anticipate that the addition of a few additional TYMS

backgrounds and/or the use of more sophisticated fitting approaches will permit more accurate

biochemical parameter inference. Indeed, recent work on the small peptide binding proteins (the

PDZ and SH3 domains) has shown how measuring the growth rate effect of mutations in different

genetic backgrounds and assay conditions can well-constrain biophysical parameters for binding

affinity and protein stability[35,41]. One might follow a conceptually similar strategy to learn

quantitative biochemical parameters from high throughput growth rate data. New microfluidics-

based approaches for high-throughput biochemistry could play a key role in refining and testing

such methodology[42].

Together our findings shape how we think about designing enzymes and metabolic

systems. Typical strategies for designing enzymes do not explicitly consider cellular context[43]. As

a result, a significant fraction of designs could fail simply because they are not properly "matched"

in terms of velocity to the surrounding pathway. The limited ability of homologs to complement

growth in another species has been observed for a number of enzymes[44–48], including DHFR[49,50].

Thus, even a well-designed catalytically active synthetic enzyme could fail to rescue growth if

placed in the wrong cellular context. Just as computational protein design considers entire

physical complexes to create binding interactions with altered affinity and specificity, one might

consider the joint design of biochemically-interacting enzymes to alter metabolic efficiency and

growth. Further study of enzyme rates and abundance across species, as well as

characterizations of enzyme velocity to growth rate mappings, will help shape our understanding

of the system level constraints placed on metabolic enzymes.

MATERIALS AND METHODS

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Escherichia coli expression and selection strains: ER2566 Δ*folA* Δ*thyA E. coli* were used for

all growth rate measurements; this strain was a kind gift from Dr. Steven Benkovic and is the

same used in Reynolds et al., 2011 and Thompson et al., 2020[51,52]. XL1-Blue *E. coli* (genotype:

recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac [F' *proAB lacI*q*ZΔM15* Tn*10*(Tet*r*)]) from

Agilent Technologies were used for cloning, mutagenesis, and plasmid propagation. BL21(DE3)

E. coli (genotype*: fhuA2 [lon] ompT gal (λ DE3) [dcm] ΔhsdS. λ DE3 = λ sBamHIo ΔEcoRI-B*

int::(lacI::PlacUV5::T7 gene1) i21 Δnin5) from New England Biolabs were used for protein

expression.

Selection vector for DHFR constructs: DHFR variants were cloned into a modified version of

the pACYC-Duet 1 vector (Novagen), which we refer to as pTet-Duet. pTet-Duet is a low-copy

number vector containing two multiple cloning sites; the first is under control of the T7 promoter

and the second was modified to be regulated by the tetracycline repressor (TetR). DHFR (*folA*) is

cloned into the first MCS; TYMS (*thyA*) is cloned into the second MCS. During selections we do

not induce expression of either gene but instead rely on leaky expression in ER2566 Δ*folA* Δ*thyA*

E. coli. The vector map for these constructs can be found on Addgene: 81596.

Expression vector for DHFR constructs: *E.coli folA* (the gene encoding DHFR) was cloned into

the pHis8-3 expression vector using restriction sites NcoI and XhoI. DHFR was tagged in-frame

with an N-terminal 8X-Histidine tag separated from the *folA* reading frame by a thrombin cleavage

site. Individual point mutant clones were constructed using the Quikchange II site-directed

mutagenesis kit (Agilent).

Expression vector for TYMS constructs: The *thyA* gene (encoding TYMS) was amplified by

PCR from *E. coli* MG1655 and cloned into the vector pET24A using XbaI/Xho restriction sites.

The point mutants of TYMS (Q33S, R127A, and Q33S) were made using the Agilent QuikChange

II site-directed mutagenesis kit.

METHOD DETAILS

Plate-reader Based Growth Rate Assays

DHFR and TYMS point mutant combinations in the selection vector were transformed into

ER2566 Δ*folA* Δ*thyA* chemically competent cells by heat shock. The cells were recovered for 60

minutes at 37°C with shaking at 220 rpm, spread on agar plates (Luria Broth (LB) containing 30

µg/ml chloramphenicol and 50 µg/ml thymidine), and grown at 37°C overnight. The next day,

liquid overnight cultures were inoculated from a streak over multiple colonies and grown overnight

at 37°C in LB supplemented with 30 µg/ml chloramphenicol and 50 µg/ml thymidine. These

overnight cultures were pelleted and washed with M9 minimal media, then resuspended in pre-

warmed M9 media supplemented with 0.4% glucose, 0.2% amicase, 2 mM $MgSO_4$, 0.1µM $CaCl_2$,

30 µg/ml chloramphenicol (henceforth referred to as M9 selection media). Next, OD_{600} for all

resuspended cultures was measured in a Perkin Elmer Victor X3 plate reader. Cultures were then

diluted to OD_{600}=0.1 in prewarmed M9 selection media with 50 µg/ml thymidine and incubated for

4 hours at 30°C, shaking at 220 rpm. After this period of adaptation and regrowth, cultures were

back-diluted to OD_{600} = 0.1 in 1 ml prewarmed M9 selection media with 50 µg/ml thymidine. These

cells were inoculated into 96-well culture plate at OD_{600} = 0.005 (10 µl cells into 200 µl total well

volume) containing prewarmed M9 selection media with 50 µg/ml thymidine; plates were sealed

with EasySeal permeable covers (Sigma Aldrich). All growth rate measurements were made in

triplicate. Plates were shaken for 10 seconds before reading, and Readings of OD_{600} were taken

every 6 minutes over 24 hours using a BioTek Synergy Neo2 plate reader in a 30°C climate-

controlled room.

DHFR Saturation Mutagenesis Library Construction

The DHFR saturation mutagenesis library was constructed as four sub-libraries in the pTet-Duet

selection vector (see above for selection vector details). Each sub-library combines mutations

within 40 contiguous amino acid positions to ensure that the mutated region can be completely

covered with short read sequencing (a 300 cycle v2 Illumina sequencing kit). The regions spanned

by each sub-library were as follows: amino acid positions 1-40 (sub-library 1, SL1), 41-80 (sub-

library 2, SL2), 81-120 (sub-library 3, SL3), and 121-159 (sub-library 4, SL4). 'Round the Horn' or

inverse PCR (iPCR) with mutagenic NNS primers (N = A/T/G/C, S = G/C) was used to introduce

all 20 amino acid substitutions at a single amino acid position as described in Thompson et al[52].

Library completeness was verified by deep sequencing. In our initial validation sequencing run

we found that mutations at positions W22 and L104 were systematically under-represented; iPCR

was repeated for these positions and they were supplemented into their respective assembled

sub-libraries.

After sub-library assembly, restriction digest and ligation were used to subclone each

sublibrary into pTet-Duet plasmids containing the three different TYMS backgrounds (WT,

R166Q, or Q33S). The entire DHFR coding region containing restriction sites (NotI and EcoNI)

was amplified by PCR. PCR reaction was size-verified with agarose gel electrophoresis with an

expected band size of 627 bp. The library PCR products and target plasmids were double

digested with NotI and EcoNI for 3 hours at 37°C. To prevent re-circularization, the digested

plasmid was treated with Antarctic phosphatase for 1 hour at 37°C. The DHFR insert and treated

plasmid were ligated with T4 DNA ligase overnight at 16°C. The concentrated ligation product

was then transformed into *E. coli* XL1-blue by electroporation, and recovered in SOB for 1 hour

at 37°C. 20 µL of the recovery culture was serially diluted and plated on LB-agar with 50 µg/mL

thymidine and 30 µg/mL chloramphenicol, to permit quantification of transformants and estimate

library coverage following ligation into the alternate TYMS backgrounds. The minimum library

coverage was 1000 colony forming units (CFU) per mutation. The remaining recovery culture was

grown in a flask containing 12 ml LB with 30 µg/mL chloramphenicol and 50 µg/mL thymidine at

37°C, with 220 rpm shaking overnight. 10 ml of the overnight culture was miniprepped with the

Gene-Jet Mini-prep kit (Fisher Scientific, K0503) to obtain the plasmid library.

Growth Rate Measurements in the Turbidostat for all DHFR mutant libraries

All sublibraries were inoculated, grown, and sampled in triplicate. Each plasmid sub-library was

transformed into the *E. coli* double knockout strain ER2566 $\Delta folA$ $\Delta thyA$ by electroporation and

recovered in SOB for one hour at 37°C. At this transformation step we again estimated library

coverage to ensure the complete library was transformed into our selection strain. To estimate

library coverage, 20 µL of the recovery culture was serial diluted with SOB and plated on LB agar

plates containing 30 µg/mL chloramphenicol and 50 µg/mL of thymidine. The remainder of the

recovery culture was inoculated into M9 selection media supplemented with 50 µg/mL thymidine

and grown overnight at 37°C. All selection experiments in this work had an estimated library

coverage of 1000 CFU/mutant or greater. The overnight liquid culture was washed and back-

diluted to OD_{600}=0.1 in M9 selection media supplemented with 50 µg/mL thymidine, and incubated

for four hours at 30°C to allow adaptation to selection temperature and to return the culture to log-

phase growth. Following adaptation, selection was initiated by back-diluting these cultures to an

OD_{600} of 0.1 into 17 mL of pre-warmed M9 selection media supplemented with 50 µg/mL thymidine

in continuous culture vials. These vials were then incubated in a turbidostat with a target OD_{600}

of 0.15 at a temperature of 30°C. The turbidostat maintained a set optical density by adding 2.8

mL dilutions of M9 selection media supplemented with 50 µg/mL thymidine in response OD

detection, and was built according to the design of Toprak et al[53]. Culture samples (1 mL each)

were taken at the beginning of selection (t = 0 hours) and at 4, 8, 12, 20, and 24 hours into

selection. Immediately after each time point, these 1 mL samples were pelleted by centrifugation,

supernatant removed, and stored at -20°C.

Next Generation Sequencing Amplicon Sample Preparation

Each turbidostat selection sample (representing a particular timepoint for a sub-library and

replicate) was prepared for sequencing as a PCR amplicon using Illumina TruSeq-HT i5 and i7

indexing barcodes. To generate these amplicons, each cell pellet from the growth rate assay was

thawed and lysed by resuspending the cells with 100 µL dH$_2$O and incubation at 95°C for 5

minutes. Lysates were then clarified by centrifugation at maximum speed for 10 minutes in a room

temperature bench top microcentrifuge. Supernatants containing plasmids were isolated from the

pellet. 1 µL of each supernatant was used as the template for an initial round of PCR with Q5-Hot

Start Polymerase (NEB) that amplified the DHFR coding region of the sublibrary (10 PCR cycles

total, standard Q5 reaction conditions). From this first PCR reaction, 1 µL was used in a second

round of PCR (22 cycles of denaturation/anneal/elongation) with primers that added Illumina

sequencing adaptors. Together, these two rounds of PCR yielded a final product of size: 315 bp

(SL1), 308 bp (SL2), 298 bp (SL3), 304 bp (SL4). The amplicons were size verified using 1%

agarose gel electrophoresis. In the case where a sample did not produce an amplicon, the first

round PCR was repeated with 2 µL of the supernatant rather than 1 µL, with the remaining

preparation identical.   All amplicons were individually quantified using with Quant-iT™

PicoGreen™ dsDNA Assay Kit (ThermoFischer Scientific) and mixed in equimolar ratio, with a

final target amount greater than or equal to 2000 ng. Errors in pipetting volume were minimized

by ensuring that more than 2 µL was taken from each amplicon. This mixture was gel-purified and

then cleaned and concentrated using the Zymo Research DNA Clean & Concentrator-5 kit. Purity

was assessed by A260/A80 and A260/A230 nm absorbance ratios. The sample library DNA

concentration was measured using a Qubit dsDNA HS Assay in a Qubit 3 Fluorometer (Invitrogen

by Thermo Fischer Scientific). The sample library was diluted to 30 nM in a volume of 50 µL of

TE buffer (1 mM Tris-HCl (pH 8.5), 10 mM EDTA (pH 4)). This mixed and quantified library was

sequenced on an Illumina HiSeq (150 cycle x 2 paired-end) by GeneWiz. The NGS sequencing

run resulted in 252 GB of data, consisting of 337,353,664 reads. The raw data have been

deposited with the NCBI sequencing read archive under project identifier PRJNA791680.

DHFR Expression and Purification

DHFR mutant variants were expressed in BL21(DE3) *E. coli* grown at 30°C in 50 ml Terrific Broth

(TB) with 35 µg/ml Kanamycin (Kan) for selection. Expression was induced at an OD_{600} = 0.6-0.8

with 250 uM IPTG, and cells were grown at 18°C for 16-18 hours. Cultures were pelleted by

centrifugation for 10 minutes at 5000 x g, 4°C and supernatant removed; cell pellets were stored

at -80°C. Thawed cell pellets were lysed by sonication in 10ml lysis buffer (50 mM Tris, 500 mM

NaCl, 10 mM imidazole, pH 8.0 buffer containing 0.1 mM PMSF, 0.001 mg/ml pepstatin, 0.01

mg/ml leupeptin, 20 µg/ml DNAseI and 5 µg/mL lysozyme). The resulting lysate was clarified by

centrifugation and incubated with 0.1ml Ni-NTA agarose (Qiagen) slurry (0.05 ml column volume)

equilibrated in Nickel Binding Buffer (NiBB, 50 mM Tris pH 8.0, 500 mM NaCl, 10 mM imidazole)

for 15 minutes on a tube rocker at 4°C. The slurry was then transferred to a disposable

polypropylene column (BioRad). After washing with 10 column volumes (CV) of NiBB, DHFR was

eluted with 0.5 mL 50mM Tris pH 8.0, 500 mM NaCl, 400 mM imidazole. The eluted protein was

concentrated and buffer-exchanged to 50 mM Tris, pH 8.0 in a 10 kDa Amicon centrifugal

concentrator (Millipore) and centrifuged 15 min at 21,000 x g, 4°C to pellet any precipitates.

Following buffer exchange, the protein was purified by anion exchange chromatography (using a

BioRad HiTrapQ HP column on a BioRad NGC Quest FPLC). A linear gradient was run from 0-1

M NaCl in 50 mM Tris pH 8.0 over 30 ml (30 column volumes, CV) while collecting 0.5 ml fractions.

Fractions containing DHFR were combined, concentrated, flash-frozen in liquid nitrogen, and

stored at -80°C.

TYMS Expression and Purification

Individual TYMS mutants were expressed in BL21(DE3) *E. coli* grown at 37°C in 50 ml Terrific

Broth (TB) with 35 µg/ml Kanamycin (Kan) for selection. Expression was induced with 1mM IPTG

when the cells reached an OD_{600} = 0.6-0.8, and the cells were then grown at 18°C for 16-18 hours

before harvesting pellets for storage at -80°C. TYMS was purified from the frozen pellets following

a protocol adapted from Changchien et al [54]. Cell pellets were thawed and resuspended in TYMS

lysis buffer (20 mM Tris, 10 mM MgCl2, 0.1% deoxycholic acid, pH 7.5 with 5 mM DTT, 0.2 mg/ml

lysozyme, 5 µg/ml DNAse I) and incubated at room temperature while rocking for 15 minutes. The

resulting supernatant was clarified by centrifugation. Next, streptomycin sulfate was added to a

final concentration of 0.75% to separate nucleic acids. The cells were incubated rocking at 4°C

for 10 minutes and the supernatant was retained following centrifugation for 10 minutes at

>10,000 x g. Ammonium sulfate was then added at 50% saturation (0.3 g/ml), mixed for 10

minutes at 4°C, then centrifuged as above, retaining supernatant. Additional ammonium sulfate

was then added to the supernatant at 80% saturation (an additional 0.2g/ml), mixed for 10 minutes

at 4°C, and centrifuged as above, retaining the pellet. The pellet was dissolved in 25mM

potassium phosphate pH 6.5 and dialyzed overnight at 4°C against 1L 25 mM potassium

phosphate pH 6.5. Following dialysis the protein was purified by anion exchange (HiTrap Q HP

column, Cytiva) with a 25 CV linear gradient from 0M NaCl to 1M NaCl in 25mM potassium

phosphate pH 6.5. FPLC fractions containing TYMS were combined and concentrated using a 10

kDa Amicon concentrator (Millipore) and stored at 4°C for up to a week.

DHFR Steady-state Michaelis Menten Kinetics

DHFR k_{cat} and K_m were determined under Michaelis-Menten conditions with saturating

concentrations of NADPH as in prior work[55,56]. Briefly, DHFR protein concentration was

determined by measuring A_{280} (extinction coefficient = 33500 $M^{-1}cm^{-1}$). DHF (Sigma Aldrich) was

prepared in MTEN buffer (50 mM MES, 25 mM Tris base, 25 mM Ethanolamine, 100 mM NaCl,

pH 7.0) containing 5 mM DTT (Sigma Aldrich). 100 nM DHFR protein and 100 µM NADPH (Sigma

Aldrich) were combined in MTEN buffer with 5 mM DTT and pre-incubated for 1 hour at 25°C prior

to measurement. To initiate the reaction, the protein-NADPH solution was mixed with DHF in a

quartz cuvette (sampling DHF over a range of concentrations, tuned to the Km of the mutant).

The initial velocity of DHFR was measured spectrophotometrically by monitoring the consumption

of NADPH and DHF (decrease in absorbance at 340 nm, $\Delta\varepsilon_{340}$=13.2 mM^{-1} cm^{-1}). All

measurements were made in triplicate; analysis was performed using the Michaelis-Menten

nonlinear regression function of Graph Pad Prism.

Preparation of TYMS substrate for assaying enzyme activity and steady state kinetics

(6R)-methylenetetrahydrofolic acid (MTHF, CH_2H_4fol) was purchased from Merck & Cie

(Switzerland) and dissolved to 100 mM in nitrogen-sparged citrate-ascorbate buffer (10 mM

ascorbic acid, 8.5 mM sodium citrate, pH 8.0). 30 µL aliquots were made in light-safe

microcentrifuge tubes, flash-frozen in liquid nitrogen, and stored at -80C. Before use, the stock

was thawed and diluted to 10 mM in TYMS kinetics reaction buffer (100 mM Tris base, 5 mM

Formaldehyde, 1 mM EDTA, pH 7.5) and quantified in an enzymatic assay: 50 µM MTHF, 200

µM dUMP and 1µM TYMS protein were combined and A_{340} measured until steady-state reached.

Actual concentration was then calculated from the difference in A_{340} before and after the reaction

using Beer's Law (MTHF extinction coefficient: 6.4 $mM^{-1}cm^{-1}$).

TYMS Steady-state Michaelis Menten Kinetics

TYMS k_{cat} and K_m were determined for both dUMP and MTHF under Michaelis-Menten conditions

by varying one substrate and holding the other saturating as in prior work[57,58]. Briefly, TYMS

protein concentration was determined by measuring A_{280} (extinction coefficient = 53400 $M^{-1}cm^{-1}$).

TYMS protein was prepared in TYMS assay buffer (100 mM Tris base, 5 mM Formaldehyde, 1

mM EDTA, pH 7.5) containing 50 mM DTT (Sigma Aldrich). 50 nM TYMS protein and either 100

µM dUMP (Sigma Aldrich) or 150 µM MTHF (Merck & Cie) were combined with varying

concentrations of the other substrate to initiate the reaction. The production of DHF was monitored

spectrophotometrically (increase in absorbance at 340nm, $\Delta\varepsilon_{340}$=6.4 $mM^{-1}cm^{-1}$) for 2 minutes per

reaction. All measurements were made in triplicate; analysis was performed using the Michaelis-

Menten nonlinear regression function of Graph Pad Prism.

QUANTIFICATION AND STATISTICAL ANALYSIS

Enzyme Velocity to Growth Rate Model Construction and Parameterization

For the purposes of modeling, we approximated DHFR and TYMS as a two-enzyme cycle in which

DHFR produces THF and consumes DHF, and TYMS produces DHF and consumes THF. This

abstraction ignores the different carbon-carrying THF species, instead collapsing them into a

single "reduced folate" pool. This simplification allows us to construct an analytically solvable

model for steady state THF concentration that we can then relate to growth (Equation 3).

First, we fit the free parameters in the Goldbeter-Koshland equation ($[\text{DHFR}]$, $[\text{TYMS}_{\text{WT}}]$,

$[\text{TYMS}_{\text{R166Q}}]$, fol_{tot}) using a set of ten metabolomics measurements for the relative abundance of

3-glutamate form of formyl THF as obtained in prior work[1]. These measurements were made for

DHFR mutations G121V, F31Y/L54I, M42F/G121V, F31Y/G121V and the WT in the background

of WT TYMS and TYMS R166Q. So why this particular folate species? We noticed that the relative

abundance of many of the reduced THF species in our data set was correlated, and chose formyl

THF to model because the experimental data were less variable and showed a strong, monotonic

relationship with cell growth. Then, we fit the free parameters in equation two (g_{max}, g_{min}, K, n)

using a set of ten growth rate measurements for the same DHFR/TYMS mutation pairs. This fitting

process gave rise to the fits shown in figure 1. When assessing model performance against the

larger set of TYMS variants (as in figure 2) we refit all parameters (g_{max}, g_{min}, K, n, [DHFR],

[$TYMS_{WT}$], [$TYMS_{R166Q}$], [$TYMS_{Q33S}$], [$TYMS_{R127A}$], fol_{tot}) to the growth rate data only since we did

not have metabolomics data for this larger set. All parameter fits were made in python using the

least_squares fitting function of the scipy.optimize module[59]; the complete fitting process is

documented in Jupyter notebook 1_KGmodel.ipynb in the associated github repository.

705   We assessed the model sensitivity to shuffling the data (Figure 2 – supplement 2) by

randomly shuffling all catalytic parameters (k_{cat}, K_m) 50 times across DHFR and TYMS and

computing an R^2 value. We also assessed model sensitivity to subsampling the data; error bars

in Figures 1, 2, and Figure 2 – supplement 2 correspond to SEM across "jackknife" re-samplings

of the data wherein one DHFR/TYMS combination was left out for each re-sampling. Finally, to

assess the global model fit to the data (as in Figure 6 and Figure 6 – supplement 1) we first fit the

9 model parameters (g_{max}, g_{min}, K, n, [DHFR], [$TYMS_{WT}$], [$TYMS_{R166Q}$], [$TYMS_{Q33S}$], fol_{tot}) using the

growth rate measurements of 16 DHFR mutations for which experimental k_{cat} and K_m were known

(48 total observations given the three TYMS backgrounds). Then, fixing these parameters, we fit

k_{cat} and K_m values for all 2696 mutations with growth rate measurements in all three TYMS

backgrounds to the complete data set of 8,088 sequencing-based growth rate observations. This

process is documented in Jupyter notebook 4_ModelAndDMSData.ipynb in the associated github

repository.

Next Generation Sequencing Data Processing and Read Counting

All PCR amplicons (corresponding to individual replicates, timepoints and sublibraries) were

sequenced on an Illumina HiSeq using 2 x 150 paired end reads. The resulting fastq files were

processed and filtered prior to read counting. Briefly, the forward and reverse reads were merged

using USEARCH. Each read was quality score filtered (Q-Score ≥ 20) and identified as a WT or

mutant of DHFR using a custom python script. This python script filtered for full length reads and

base call quality scores greater than 20 (error rate ≤ 1:100). The reads passing these quality

control criteria were compared against the wild-type reference sequence to determine mutation

identify. Reads that contained multiple point mutations or mutations outside the sublibrary of

interest were removed from analysis. This process resulted in counts for the WT and each mutant

at each time point and replicate. These counts were further corrected given the expected error in

the data (q-score) and Hamming distance from the WT codon to account for potential hopping of

WT reads to mutations; a process that was detailed in McCormick et al [56].

Relative Growth Rate Calculations

We calculated relative growth rates for individual mutations and the WT over time from the

sequencing-based counts (N_t^{mut}, N_t^{WT}). Mutants with fewer than 10 counts were considered

absent from the data set and were set to zero to reduce noise. From these thresholded counts,

we calculated a log normalized relative frequency of each mutation over time:

$$log_2(f(t)) \ = \ log_2(N_t^{mut}/N_t^{WT}) \ - \ log_2(N_{t=0}^{mut}/N_{t=0}^{WT})$$

We then calculated relative growth rate (m_{DHmut}^{TS}) as the slope of the log relative frequency over

time by linear regression. Linear regression was performed using scikit Learn, and individual

points were weighted by the number of counts (in order to down weight less-sampled mutants at

later time points). Relative growth rate or a mutant was only calculated if the mutant was present

over at least the first three time points, otherwise it was classified as a "Null" mutant. Finally, all

relative growth rates were normalized such that WT has a relative growth rate of 1. Growth rates

were additionally normalized by the bulk culture growth rate (estimated from the turbidostat, in

units of generations per hour) to account for small vial-to-vial variations culture doublings across

the experiment. The standard error in growth rate was computed across triplicate measurements.

All calculations are shown in Jupyter notebook 2_DMSGrowthRates.ipynb in the associated

github repository.

Epistasis Analysis

Epistasis was calculated according to an additive model:

$$\varepsilon_{DHmut,TSmut} = m_{DHmut}^{TS\_mut} - m_{DHmut}^{TS\_WT}$$

In our experiments TYMS R127A, Q33S and R166Q have no growth rate effect in the WT DHFR

context due to thymidine supplementation. Under this formalism, mutations that show improved

growth in the mutated TYMS background have positive epistasis, while mutations with reduced

growth in the mutated TYMS background have negative epistasis. We assessed the statistical

significance of epistasis by unequal variance t-test under the null hypothesis that the mutations

have equal mean growth rates in both TYMS backgrounds (across three replicate

measurements). These p-values were compared to a multiple-hypothesis testing adjusted p-value

determined by Sequential Goodness of Fit (P = 0.035 for TYMS Q33S and P = 0.029 for TYMS

R166Q) [34]. K-means clustering of epistatic positions was performed using a custom script based

on that described in Thompson et al [52]. All epistasis calculations are shown in Jupyter notebook

3_Epistasis.ipynb in the associated github repository.

ACKNOWLEDGEMENTS

The authors thank Olivier Rivoire for early conversations regarding our data, and Elliott Ross for

critical feedback on both the modeling and manuscript. We also thank the Reynolds lab for

feedback on experimental design, data analysis, and the manuscript.

FUNDING

Research reported in this publication was supported by the National Institute of General Medical

Sciences of the National Institutes of Health under Award Number R01GM136842. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health. This work was also partly supported in its early stages by the

Gordon and Betty Moore Foundation's Data Driven Discovery Initiative through grant GBMF4557

to KAR.

AUTHOR CONTRIBUTIONS

TNN and KAR conceptualized the work and designed experiments. KAR created the

mathematical model. TNN collected all deep mutational scanning data. TNN and KAR analyzed

the data. CI collected plate-reader based growth rate measurements (used in model

development) and performed all Michaelis-Menten enzyme kinetics assays. SMT constructed the

deep mutational scanning library and contributed to data interpretation. KAR wrote the original

draft. TNN, CI, SMT and KAR revised the paper. KAR provided supervision and funding

acquisition.

COMPETING INTERESTS

The authors have no competing interests to declare.

DATA AVAILABILITY

The raw sequencing data is available in FASTQ format through the NCBI sequencing read

archive, under **BioProject ID PRJNA791680**.

CODE AVAILABILITY

Code for the enzyme velocity to growth rate model, and analysis of all deep mutational scanning

data is available on github: https://github.com/reynoldsk/dhfr-tyms-epistasis

MATERIALS AND CORRESPONDENCE

The DHFR deep mutational scanning libraries (in all three TYMS backgrounds) have been

deposited at addgene under **deposit number 81596**.

Requests for any other materials or data should be made by email to Dr. Kimberly Reynolds,

with Christine Ingle in carbon copy (cc).

 FIGURES AND LEGENDS

806
Figure 1. Constructing a biochemistry-to-growth model for DHFR and TYMS.
a) Schematic describing the relationship between metabolic pathway flux and enzyme velocity.
Many enzymes show a hyperbolic relationship between velocity and flux; the enzyme control
coefficient describes the fractional change in flux given a fractional change in velocity. Control
coefficients vary with the starting enzyme velocity (purple and dark blue arrows, background
"A") and can change with genetic background (violet and light blue arrows, background "B").
Consequently mutations can have a strong effect on flux and growth in one context but not in
another (compare violet and purple arrows).
b) The role of DHFR and TYMS in folate metabolism. Metabolites are labeled in grey or black
italic text. Dotted lines indicate multiple intermediate reactions that are summarized with a
single line.
c) The relationship between the experimentally measured relative abundance of [10-formyl-THF]
and *E. coli* growth rate. Red points indicate five DHFR variants in the background of TYMS
R166Q (a near catalytically inactive variant) and black indicates the same DHFR variants in
the context of WT TYMS. Error bars indicate the standard deviation across N=3 replicates for
both growth rate (y-axis) and 10-formyl-THF abundance (x-axis). The blue dotted line
indicates the best fit for a hyperbolic model (Equation 1) relating THF abundance to growth.
d) A simplified, abstracted version of the DHFR and TYMS system. Again dotted lines indicate
multiple intermediate reactions that are summarized with a single line.

e)   The correlation between experimentally measured $\log_{10}$[10-formyl-THF] relative abundance
and the model prediction (as computed with Equation 3). The grey dotted line indicates x=y.
Color coding is identical to **c**. Error bars in x indicate the standard deviation across N=3
replicate experimental measurements, error bars in y describe the standard deviation across
ten fits obtained by jackknife (leave-one-out) sub-sampling the data and refitting the model.
f)   The correlation between experimentally measured and predicted growth rates for five DHFR
point mutations in two different TYMS backgrounds (same mutants as in **c,e**). The grey dotted
line indicates x=y. Color coding is identical to **c**. Error bars in x indicate the standard deviation
across N=3 replicate experimental measurements, error bars in y describe the standard
deviation across ten fits obtained by jackknife (leave-one-out) sub-sampling the data and
refitting the model.

Figure 2. **TYMS catalytic activity modulates the epistatic constraints on DHFR sequence and catalytic activity**.

a) Location of the TYMS point mutations (PDBID: 1BID[60]). TYMS functions as an obligate domain-swapped homodimer; active sites include residues from both monomers (white and grey cartoon). Positions mutated in this study are in colored spheres, and indicated with arrows (Q33 – cyan, R127 – navy, R166 – red). The dUMP substrate is in sticks, labeled and colored green.

b) Michaelis Menten enzyme kinetics for WT TYMS (black), TYMS Q33S (cyan), and TYMS R127A (navy). Experimental replicates (3 total) are plotted individually. Points indicate experimental data and lines the best fit steady state model.

c) Correlation between experimentally measured and model-predicted relative growth rates for seven DHFR variants in four TYMS backgrounds. Each point represents one DHFR/TYMS combination. Error bars in the x direction are SEM across triplicate growth rate measurements, error bars in y are the SEM estimated from jackknife (leave-one-out) sub-sampling the data and refitting the model.

d) Heatmaps of simulated growth rates (top row) and epistasis (bottom row) as computed over a range of DHFR kinetic parameters in four TYMS backgrounds. In the left-most column of heatmaps a red star marks the highest activity enzyme (low K_m, high k_{cat}), while a yellow star marks the lowest activity enzyme. Growth rates are indicated with a blue-white-red color map, where a relative growth rate of one (white) is equivalent to WT. Epistasis values are indicated with a green-white-pink color map, where zero epistasis is shown in white.

GAVLMIFYWKRHDESTCNQP*

Figure 3. The effects of DHFR mutation on growth rate in three TYMS backgrounds.

a) Sequencing-based growth rate measurements for DHFR F31V in three TYMS backgrounds: R166Q (red), Q33S (cyan), and WT (black). Each point represents one triplicate experimental measurement. Dotted lines indicate linear regression fits to each replicate, the slope of each line is the inferred growth rate (relative to WT) for that DHFR/TYMS mutant combination.

b) Heatmaps of the growth rate effect for all DHFR single mutations. DHFR positions are along the horizontal axis; amino acid residues (along the vertical axis) are organized by physiochemical similarity. The displayed relative growth rate is an average across three replicates, and is normalized such that the WT DHFR is equal to one. Red indicates mutations that increase growth rate, white indicates mutations with wild-type like growth, and blue indicates mutations that decrease growth rate. Null mutations (black squares) were not

observed by sequencing after the first two time points, and thus there was insufficient data for
growth rate inference. Small dots mark the WT residue identity in each column.
c) The distribution of DHFR mutational effects in the WT TYMS background. The red line
indicates a best-fit double gaussian, grey bars are the data. The red, dashed "inactive" line
marks the average relative growth rate for nonsense mutations (stop codons) in the first 120
positions of DHFR. The WT DHFR growth rate is equal to one.
d) The distribution of DHFR mutational effects in the TYMS Q33S background, color coding
identical to (**c**)
e) The distribution of DHFR mutational effects in the TYMS R166Q background, color coding
identical to (**c**). Note that the y-axis for (**e**) is distinct from (**c**) and (**d**).
879
880

Figure 4. Epistatic coupling of DHFR to two TYMS backgrounds.

a) Volcano plot examining the statistical significance of epistasis across all DHFR point mutations in the Q33S background. P-values were calculated by unequal variance t-test under the null hypothesis that the mutations have equal mean growth rates in both TYMS backgrounds (across triplicate measurements). The red horizontal dashed line marks the standard significance cutoff of P=0.05, the black horizontal dashed line indicates a multiple-hypothesis testing adjusted p-value (P=0.035). The grey vertical dashed lines indicate an empirical threshold for epistasis. Pink and green indicate statistically significant positive and negative epistasis respectively.

b) Volcano plot examining the statistical significance of epistasis across all DHFR point mutations in the R166Q background. P-values were calculated as in (**a**); the multiple-hypothesis testing adjusted p-value for the R166Q background was (P=0.029). Color coding is identical to (**a**).

c) Comparison of the relative growth rate effects for DHFR single mutants in the WT and TYMS Q33S backgrounds. The marginal distribution of growth rate effects is shown along each axis.

Mutations with statistically significant positive and negative epistasis are indicated in pink and
green respectively. The WT relative growth rate equals one, and is indicated with a dashed
grey line across each axis. The dashed red line marks x=y. Error bars (in x and y) indicate
standard deviation across three experimental measurements.

d) Comparison of the relative growth rate effects for DHFR single mutants in the WT and TYMS
R166Q backgrounds. Plot layout and color coding is identical to (**c**). Error bars (in x and y)
indicate standard deviation across three experimental measurements.

904
905

Figure 5. Global comparison of the biochemistry-to-growth model and deep mutational scanning data set.

a) Correlation between the experimentally measured and predicted growth rates of 114 DHFR/TYMS mutant combinations (circles colored according to TYMS background). Horizontal error bars indicate standard deviation in experimentally measured growth rates across three replicate measurements, vertical error bars are the standard deviation in the predicted growth rates estimated by performing 1000 bootstrap re-samplings (and model fits) of the data. The dashed grey line indicates y=x.

b) Correlation between the experimentally measured and model-predicted epistasis, as computed from the growth rate data in (**a**). Again, color coding indicates TYMS background (identical to **a**). The dashed grey line indicates y=x. Horizontal error bars indicate standard deviation propagated from the experimentally measured growth rates across three replicate measurements, vertical error bars are the standard deviation in the predicted epistasis values estimated by performing 1000 bootstrap re-samplings (and model fits) of the data.

c) Correlation between the experimentally measured and computationally inferred $\log_{10}(k_{cat}/K_m)$ values for 38 mutants of DHFR. Horizontal error bars describe the standard deviation across triplicate experimental measurements, vertical error bars indicate the standard deviation across 50 iterations of stochastic (Monte-Carlo based) model inference.

d) Correlation between experimentally measured and predicted growth rates across the entire deep mutational scanning dataset. The marginal distribution of growth rate effects is shown along each axis.

e) Correlation between experimentally measured and predicted epistasis across the entire deep
mutational scanning dataset. The marginal distribution of epistatic effects is shown along
each axis.
f) Mutational tolerance of DHFR as a function of TYMS background. The heatmap shows the
fraction of DHFR mutations with growth rates of 0.9 or better as TYMS k_{cat} and K_m are
discretely varied. Both axes are natural log spaced, TYMS k_{cat} was sampled at 50 points
between ln(-2) and ln(2), while TYMS K_m was sampled at 50 points between ln(-1) and ln(3).
The values for TYMS R166Q, Q33S and WT are marked with red, cyan and black circles
respectively.
g) A zoomed-in version of (**f**), focusing on the mutational tolerance of DHFR for TYMS
backgrounds similar in velocity to WT and Q33S TYMS.
938
939

Figure 6. DHFR positions clustered by epistatic mutational effect.

a) Clusters of DHFR positions organized by predominant epistasis type. In each heat map DHFR positions are ordered along the vertical axis; amino acid residues are organized by physiochemical similarity along the horizontal axis. As in earlier plots, green indicates negative epistasis, and pink indicates positive epistasis. Grey pixels mark mutations with statistically insignificant epistasis.

b) Structural location of epistatic clusters for DHFR to TYMS Q33S. The DHFR backbone is in grey cartoon (PDBID: 1RX2[31]). Folate, the DHFR substrate is indicated with yellow sticks. The NADP+ cofactor is in green sticks.

c) Structural location of epistatic clusters for DHFR to TYMS R166Q. Color coding is identical to panel (**b**).

954
Figure 7. The structural organization of epistasis in DHFR.
a) Epistasis of individual DHFR positions to TYMS Q33S. The DHFR structure is shown in space
filling spheres (PDBID: 1RX2), with the NADP co-factor in green sticks, and folate in yellow
sticks. A slice through the structure shows the interior arrangement of epistasis. Positions in
the negative epistasis cluster are colored green, positions in the positive epistasis cluster are
colored pink. Grey spheres indicate positions in the insignificant epistasis cluster.
b) Cumulative distribution of positions in each epistatic cluster by distance to the DHFR active
site for the TYMS Q33S background. In this case, active site was defined as the C6 atom of
the folate substrate. Color coding follows from (**a**)
c) Epistasis of individual DHFR positions to TYMS R166Q. The DHFR structure is shown in
space filling spheres (PDBID: 1RX2), with the NADP co-factor in green sticks, and folate in
yellow sticks. A slice through the structure shows the interior arrangement of epistasis.
Positions with strong positive epistasis are colored magenta, positions in the positive epistasis
cluster are colored pink. Grey spheres indicate positions in the insignificant epistasis cluster.
d) Cumulative distribution of positions in each epistatic cluster by distance to the DHFR active
site for the TYMS R166Q background. In this case, active site was defined as the C6 atom of
the folate substrate. Color coding follows from (**c**)
e) The position-averaged $\log_{10}$ catalytic power across measured mutations. All residues are
indicated in space filling and color coded by the average mutational effect. Blue indicates
positions where mutations have a deleterious effect on catalytic power (on average), while

white indicates mutations that have little to no effect on catalytic power. Again, the NADP co-
factor is shown in green sticks, and folate in yellow sticks. A slice through the structure shows
the interior distribution of mutational effects on catalysis.
f) The structural overlap between positions associated to catalysis and evolutionary
conservation. The DHFR backbone is shown in grey cartoon, the NADP co-factor in green
sticks, and folate in yellow sticks. Positions where mutations have (on average) a deleterious
effect on catalysis at least half a standard deviation below the mean are shown in blue space
filling (color coding identical to **c**). Evolutionarily conserved positions (as computed by the
Kullback-Leibler relative entropy in a large alignment of DHFR sequences) are outlined in red
mesh.
985
986

REFERENCES

1. Schober, A. F. *et al.* A two-enzyme adaptive unit within bacterial folate metabolism. *Cell Reports* **27**, 3359-3370.e7 (2019).

2. Kemble, H. E. *et al. Flux, toxicity and protein expression costs shape genetic interaction in a metabolic pathway*. http://biorxiv.org/lookup/doi/10.1101/362327 (2018) doi:10.1101/362327.

3. Lilja, E. E. & Johnson, D. R. Metabolite toxicity determines the pace of molecular evolution within microbial populations. *BMC Evol Biol* **17**, 52 (2017).

4. Noor, E. *et al.* The Protein Cost of Metabolic Fluxes: Prediction from Enzymatic Rate Laws and Cost Minimization. *PLOS Comput. Biol.* **12**, e1005167 (2016).

5. Dekel, E. & Alon, U. Optimality and evolutionary tuning of the expression level of a protein. *Nature* **436**, 588–592 (2005).

6. Bar-Even, A. *et al.* The Moderately Efficient Enzyme: Evolutionary and Physicochemical Trends Shaping Enzyme Parameters. *Biochemistry* **50**, 4402–4410 (2011).

7. Lalanne, J.-B. *et al.* Evolutionary Convergence of Pathway-Specific Enzyme Expression Stoichiometry. *Cell* **173**, 749-761.e38 (2018).

8. Kacser, H. & Burns, J. A. The control of flux. *Biochem Soc Trans* **23**, 341–366 (1995).

9. Dykhuizen, D. E., Dean, A. M. & Hartl, D. L. Metabolic Flux and Fitness. *Genetics* **115**, 25–31 (1987).

10. Kacser, H. & Burns, J. A. The Molecular Basis of Dominance. *Genetics* **97**, 639–666 (1981).

11. Domingo, J., Baeza-Centurion, P. & Lehner, B. The Causes and Consequences of Genetic Interactions (Epistasis). *Annu. Rev. Genom. Hum. Genet.* **20**, 433–460 (2019).

12. Pokusaeva, V. O. *et al.* An experimental assay of the interactions of amino acids from orthologous sequences shaping a complex fitness landscape. *PLoS Genet.* **15**, e1008079 (2019).

13. Kondrashov, A. S., Sunyaev, S. & Kondrashov, F. A. Dobzhansky-Muller incompatibilities in

protein evolution. *Proc Natl Acad Sci U S A* **99**, 14878–14883 (2002).

14. Noda-Garcia, L., Liebermeister, W. & Tawfik, D. S. Metabolite-Enzyme Coevolution: From

Single Enzymes to Metabolic Pathways and Networks. *Annu Rev Biochem* **87**, 187–216

(2018).

15. Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell

Metabolism* **25**, 27–42 (2017).

16. Nijhout, H. F., Reed, M. C., Budu, P. & Ulrich, C. M. A Mathematical Model of the Folate

Cycle: NEW INSIGHTS INTO FOLATE HOMEOSTASIS. *J. Biol. Chem.* **279**, 55008–55016

(2004).

17. Leduc, D. *et al.* Flavin-Dependent Thymidylate Synthase ThyX Activity: Implications for the

Folate Cycle in Bacteria. *Journal of Bacteriology* **189**, 8537–8545 (2007).

18. King, C. H., Shlaes, D. M. & Dul, M. J. Infection caused by thymidine-requiring,

trimethoprim-resistant bacteria. *Journal of Clinical Microbiology* **18**, 79–83 (1983).

19. Howell, E. E., Foster, P. G. & Foster, L. M. Construction of a dihydrofolate reductase-

deficient mutant of Escherichia coli by gene replacement. *J Bacteriol* **170**, 3040–3045

(1988).

20. Rodrigues, J. V. & Shakhnovich, E. I. Adaptation to mutational inactivation of an essential

gene converges to an accessible suboptimal fitness peak. *Elife* **8**, e50509 (2019).

21. Kwon, Y. K., Higgins, M. B. & Rabinowitz, J. D. Antifolate-Induced Depletion of Intracellular

Glycine and Purines Inhibits Thymineless Death in *E. coli*. *ACS Chem. Biol.* **5**, 787–795

(2010).

22. Kwon, Y. K. *et al.* A domino effect in antifolate drug action in Escherichia coli. *Nat Chem Biol*

4, 602–608 (2008).

23. Sotelo-Mundo, R. R., Changchien, L., Maley, F. & Montfort, W. R. Crystal structures of

thymidylate synthase mutant R166Q: Structural basis for the nearly complete loss of

catalytic activity. *Journal of Biochemical and Molecular Toxicology* **20**, 88–92 (2006).

24. Rodrigues, J. V. *et al.* Biophysical principles predict fitness landscapes of drug resistance.

PNAS **113**, E1470–E1478 (2016).

25. Ferrell, J. E. & Ha, S. H. Ultrasensitivity part I: Michaelian responses and zero-order

ultrasensitivity. *Trends in Biochemical Sciences* **39**, 496–503 (2014).

26. Goldbeter, A. & Koshland, D. E. An amplified sensitivity arising from covalent modification in

biological systems. *Proceedings of the National Academy of Sciences* **78**, 6840–6844

(1981).

27. Michaels, M. L., Kim, C. W., Matthews, D. A. & Miller, J. H. Escherichia coli thymidylate

synthase: amino acid substitutions by suppression of amber nonsense mutations. *PNAS* **87**,

3957–3961 (1990).

28. Schnell, J. R., Dyson, H. J. & Wright, P. E. Structure, Dynamics, and Catalytic Function of

Dihydrofolate Reductase. *Annual Review of Biophysics and Biomolecular Structure* **33**,

119–140 (2004).

29. Shrimpton, P. & Allemann, R. K. Role of water in the catalytic cycle of E. coli dihydrofolate

reductase. *Protein Sci* **11**, 1442–1451 (2002).

30. Gekko, K., Kunori, Y., Takeuchi, H., Ichihara, S. & Kodama, M. Point Mutations at Glycine-

121 of Escherichia coli Dihydrofolate Reductase: Important Roles of a Flexible Loop in the

Stability and Function. *J Biochem* **116**, 34–41 (1994).

31. Sawaya, M. R. & Kraut, J. Loop and subdomain movements in the mechanism of

Escherichia coli dihydrofolate reductase: crystallographic evidence. *Biochemistry* **36**, 586–

603 (1997).

32. Chen, J. T., Taira, K., Tu, C. P. D. & Benkovic, S. J. Probing the functional role of

phenylalanine-31 of Escherichia coli dihydrofolate reductase by site-directed mutagenesis.

Biochemistry **26**, 4093–4100 (1987).

33. Watney, J. B., Agarwal, P. K. & Hammes-Schiffer, S. Effect of Mutation on Enzyme Motion

in Dihydrofolate Reductase. *J. Am. Chem. Soc.* **125**, 3745–3750 (2003).

34. Carvajal-Rodríguez, A., de Uña-Alvarez, J. & Rolán-Alvarez, E. A new multitest correction

(SGoF) that increases its statistical power when increasing the number of tests. *BMC*

Bioinformatics **10**, 209 (2009).

35. Faure, A. J. *et al.* Mapping the energetic and allosteric landscapes of protein binding

domains. *Nature* **604**, 175–183 (2022).

36. Tamer, Y. T. *et al.* High-Order Epistasis in Catalytic Power of Dihydrofolate Reductase

Gives Rise to a Rugged Fitness Landscape in the Presence of Trimethoprim Selection.

Molecular Biology and Evolution **36**, 1533–1550 (2019).

37. Karanicolas, J. & Kuhlman, B. Computational design of affinity and specificity at protein–

protein interfaces. *Current Opinion in Structural Biology* **19**, 458–463 (2009).

38. McClune, C. J., Alvarez-Buylla, A., Voigt, C. A. & Laub, M. T. Engineering orthogonal

signalling pathways reveals the sparse occupancy of sequence space. *Nature* **574**, 702–

706 (2019).

39. Li, G.-W., Burkhardt, D., Gross, C. & Weissman, J. S. Quantifying Absolute Protein

Synthesis Rates Reveals Principles Underlying Allocation of Cellular Resources. *Cell* **157**,

624–635 (2014).

40. Papp, B., Pál, C. & Hurst, L. D. Dosage sensitivity and the evolution of gene families in

yeast. *Nature* **424**, 194–197 (2003).

41. Otwinowski, J. Biophysical Inference of Epistasis and the Effects of Mutations on Protein

Stability and Function. *Molecular Biology and Evolution* **35**, 2345–2354 (2018).

42. Markin, C. J. *et al.* Revealing enzyme functional architecture via high-throughput microfluidic

enzyme kinetics. *Science* **373**, eabf8761 (2021).

43. Lovelock, S. L. *et al.* The road to fully programmable protein catalysis. *Nature* **606**, 49–58

(2022).

44. Michener, J. K., Camargo Neves, A. A., Vuilleumier, S., Bringel, F. & Marx, C. J. Effective

use of a horizontally-transferred pathway for dichloromethane catabolism requires post–

transfer refinement. *eLife* **3**, e04279 (2014).

45. Michener, J. K., Vuilleumier, S., Bringel, F. & Marx, C. J. Phylogeny poorly predicts the utility

of a challenging horizontally transferred gene in Methylobacterium strains. *J Bacteriol* **196**,

2101–2107 (2014).

46. Kim, J. & Copley, S. D. Inhibitory cross-talk upon introduction of a new metabolic pathway

into an existing metabolic network. *Proc Natl Acad Sci U S A* **109**, E2856-2864 (2012).

47. Kachroo, A. H. *et al.* Systematic humanization of yeast genes reveals conserved functions

and genetic modularity. *Science* **348**, 921–925 (2015).

48. Russ, W. P. *et al.* An evolution-based model for designing chorismate mutase enzymes.

Science **369**, 440–445 (2020).

49. Bhattacharyya, S. *et al.* Transient protein-protein interactions perturb E. coli metabolome

and cause gene dosage toxicity. *Elife* **5**, e20309 (2016).

50. Bershtein, S. *et al.* Protein homeostasis imposes a barrier on functional integration of

horizontally transferred genes in bacteria. *PLoS Genet* **11**, e1005612 (2015).

51. Reynolds, K. A., McLaughlin, R. N. & Ranganathan, R. Hot Spots for Allosteric Regulation

on Protein Surfaces. *Cell* **147**, 1564–1575 (2011).

52. Thompson, S., Zhang, Y., Ingle, C., Reynolds, K. A. & Kortemme, T. Altered expression of a

quality control protease in E. coli reshapes the in vivo mutational landscape of a model

enzyme. *eLife* **9**, e53476 (2020).

53. Toprak, E. *et al.* Building a morbidostat: an automated continuous-culture device for

studying bacterial drug resistance under dynamically sustained drug inhibition. *Nature*

Protocols **8**, 555–567 (2013).

54. Changchien, L.-M. *et al.* High-Level Expression of Escherichia coli and Bacillus subtilis

Thymidylate Synthases. *Protein Expression and Purification* **19**, 265–270 (2000).

55. Reynolds, K. A., McLaughlin, R. N. & Ranganathan, R. Hot Spots for Allosteric Regulation

on Protein Surfaces. *Cell* **147**, 1564–1575 (2011).

56. McCormick, J. W., Russo, M. A., Thompson, S., Blevins, A. & Reynolds, K. A. Structurally

distributed surface sites tune allosteric regulation. *Elife* **10**, e68346 (2021).

57. Agrawal, N., Hong, B., Mihai, C. & Kohen, A. Vibrationally enhanced hydrogen tunneling in

the Escherichia coli thymidylate synthase catalyzed reaction. *Biochemistry* **43**, 1998–2006

(2004).

58. Wang, Z. *et al.* Mg2+ Binds to the Surface of Thymidylate Synthase and Affects Hydride

Transfer at the Interior Active Site. *J. Am. Chem. Soc.* **135**, 7583–7592 (2013).

59. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat*

Methods **17**, 261–272 (2020).

60. Stout, T. J., Sage, C. R. & Stroud, R. M. The additivity of substrate fragments in enzyme–

ligand binding. *Structure* **6**, 839–848 (1998).

1128